1

# Sex differences in post-stroke aphasia rates are caused by age. A meta-analysis and database query

*Mikkel Wallentin* [1,2,3,*]

1) Department of Linguistics, Cognitive Science and Semiotics, Aarhus University, Jens Chr.

Skous Vej 2, 8000 Aarhus C, Denmark;

2) Center of Functionally Integrative Neuroscience, Aarhus University Hospital, Denmark;

3) Interacting Minds Centre, Aarhus University, Denmark

*) Corresponding author

Email: mikkel@cc.au.dk

Word count (including title page, abstract, references, tables and captions): 7785

# Abstract

*Background:* Studies have suggested that aphasia rates are different in men and women following stroke. One hypothesis says that men have more lateralized language function than women. Given unilateral stroke, this would lead to a prediction of men having higher aphasia rates than women. Another line of observations suggest that women are more severely affected by stroke, which could lead to a higher aphasia rate. An additional potential confounding variable could be age at stroke, given that women are typically older at the time of stroke.

*Methods & Procedures:* This study consists of two parts. First, a meta-analysis of the available reports of aphasia rates in the two sexes was conducted. A comprehensive literature search yielded 25 studies with sufficient aphasia and gender information. These studies included a total of 48,362 stroke patients for which aphasia rates were calculated. Second, data were analyzed from an American health database with 1,655,295 stroke patients, in order to include age and stroke severity into a regression analysis of sex differences in aphasia rates.

*Outcomes & Results:* Both analyses revealed significantly larger aphasia rates in women than in men. This speaks against the idea that men should be more lateralized in their language function. Furthermore, when age and stroke severity were included as covariates, sex failed to explain any aphasia rate sex difference above and beyond that which is explained by age differences at time of stroke.

# Introduction

A stroke is a medical condition in which blood flow to the brain is restricted, due to occlusion (ischemic stroke) or haemorrhage (haemorrhagic stroke), resulting in cell death (WHO). In the US alone, approximately 800,000 people experience a stroke each year, according to the American Heart Association [1]. Stroke is the leading cause of motor and cognitive disability in western countries and aphasia, the inability to comprehend and formulate language because of brain damage, is one of the most common deficits after stroke. A large variability in the reported frequency of aphasia can be found in the literature [e.g. 2, 3-5], ranging from 15 to 68 percent of acute patients. A recent meta-analysis, however, concluded that aphasia is present in approximately 30 % of acute patients and 34 % in rehabilitation settings [6]. Variability of measured prevalence of aphasia has many causes. The method for aphasia identification differs between hospitals and countries and some sub-scores for aphasia in stroke scales have been found to be limited in their accuracy and reliability [7, 8]. Another potential source of variance may be sex. The above-mentioned meta-analysis did not take potential sex differences into account.

Stroke has been noted to affect the sexes differently. Stroke is more common among men [9]. The symptoms of stroke have also been found to differ somewhat between men and women. Women are often more severely affected overall, more often experience paralysis, impaired consciousness and altered mental status together with a generalized weakness, while men more often experience dysarthria, diplopia, sensory loss, ataxia and balancing problems [10]. An association between pre-stroke dementia, which is more prevalent in women, and stroke severity has also been noted [11]. Lastly, aphasia following stroke has been reported to affect women to a larger degree than men [see 10 for a review], although evidence has been conflicting [e.g. 12, 13, 14].

Potential sex differences in aphasia may shed light on overall sex differences in language and

cognition. Sex differences in certain linguistic domains are known to exist within the normal

population, with differences in first language acquisition speed [15] and reading and writing

abilities [16] being the most consistent, favoring girls/women over boys/men. Differences in

word use have also been documented [17]. The underlying causes for these differences are

probably complex and research trying to tie them to brain structure and function has yielded

inconsistent results [18]. Some studies, however, have argued for the hypothesis that language

is more bilaterally organized in the brains of women than in men [e.g. 19, 20-22], although

this is highly controversial [18, 23, 24]. A sex difference in language lateralization would

ultimately lead to a sex difference in aphasia following unilateral stroke. If men's language is

more lateralized in the brain than the language of women, we would expect them to be more

prone to aphasia following unilateral stroke and vice versa, if women have greater language

lateralization than men, we would expect women's language function to be more vulnerable

to stroke.

In this paper I conduct a meta-analysis on aphasia prevalence given stroke across published

peer-reviewed papers and test if the frequency for the two sexes differ. I then compare the

results of the meta-analysis to data from a large American patient database (Healthcare Cost

and Utilization Project (HCUP) under Agency for Healthcare Research and Quality, U.S.

Department of Health & Human Services: https://hcupnet.ahrq.gov). Given that both analyses

are based on fully anonymized and publicly available data, the study poses no ethical

concerns.

# Meta-analysis methods

This report was prepared according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA, http://www.prisma-statement.org). PRISMA is an evidence-based minimum set of items for reporting in systematic reviews and meta-analyses [25]. The supporting PRISMA checklist for this meta-analysis is available as supporting information; see S1 PRISMA Checklist.

The main outcome measures for the analysis was aphasia rate (percentage of stroke patients with aphasia diagnosis) and sex ratio of aphasia rate. A pub-med search including the terms "stroke" AND "aphasia" AND "gender" (which automatically includes the term "sex") generated 211 citations in June 2018 (see Fig 1 for a flow-chart of the data sampling procedure). No time or language constraints were put on the sampled reports. References in review articles on stroke and aphasia were also investigated. A total of 419 titles were considered. 90 papers were selected for further inspection on the basis of their title and abstract. The analysis set out to study post-stroke aphasia in the broadest sense. Speech pathology test still lack standardization and diagnostic data for identifying aphasia in stroke populations [26]. Different diagnostic approaches to aphasia were therefore not distinguished. All studies including some overall aphasia diagnosis were deemed relevant. Inclusion criteria were the following: 1) Primary peer reviewed studies reporting aphasia frequency among stroke patients; 2) Studies reporting overall number of aphasia patients and stroke patients from a unselected stroke cohort (thus limiting bias), i.e. studies including only sub-types of aphasia were excluded and studies of group comparisons between aphasia patients and matched control groups were also not considered; 3) Reports reporting aphasia counts for both male and female aphasia patients as well as for stroke patients or where these could be extracted from reported percentages; 4) First/Primary report of data: The same data could

only be included once. If more than one paper investigated the same patient group, the

earliest/most comprehensive publication was chosen.

For each study, the number of stroke patients for each sex, the number of aphasia patients for

each sex and the percentage of aphasia patients were extracted. Studies with more patients

were considered less at risk of within-study bias and thus analyses used study size weights.

Study year, and place was also noted, as well as potential additional information about the

patients' age and time/type of study relative to disease onset. All data were analyzed using R.

Heterogeneity between studies was assessed using the metaphor package [27] and a funnel

plot was used to inspect potential bias. Year of study was added as a covariate to a subsequent

analysis in order to investigate if sex differences in aphasia diagnoses have been changing

over time.

124
**Fig 1. Data gathering flow-chart.** Flow chart depicting the different phases of data

gathering.

Twenty-five studies were included in the final dataset [4, 5, 11-14, 28-46]. One study [12]

contained information about aphasia frequency for different age bands. These groups were

included separately (see Table 1 and Fig 2).

**Table 1. Studies included in the meta-analysis**

| Study | Place | Note | N(stroke) | Aphasia Rate % (Female) | Aphasia Rate % (Male) | Ratio | 95% CI lower | 95% CI upper |
|---|---|---|---|---|---|---|---|---|
| Brust et al. 1976 | New York, USA | acute stroke | 850 | 18.4 | 23.8 | 0.774 | 0.580 | 1.040 |
| Siirtola et al. 1977 | Turku, Finland | acute stroke | 338 | 28.7 | 28.2 | 1.016 | 0.670 | 1.540 |
| Miceli et al. 1981 | Rome, Italy | 223 CVA, 128 tumours, 29 traumas etc | 390 | 60.8 | 62.3 | 0.975 | 0.750 | 1.280 |
| Scarpa et al. 1987 | Modena, Italy | left hemisphere stroke, post 14 days | 196 | 62.5 | 50.0 | 1.250 | 0.860 | 1.820 |
| Hier et al. 1994 | four sites, USA | acute stroke | 1805 | 22.6 | 19.4 | 1.103 | 0.920 | 1.330 |
| Pedersen et al. 1995 | Copenhagen, Denmark | acute stroke | 881 | 39.7 | 34.8 | 1.140 | 0.920 | 1.420 |

| Study | Place | Note | N(stroke) | Aphasia Rate % (Female) | Aphasia Rate % (Male) | Ratio | 95% CI lower | 95% CI upper |
|---|---|---|---|---|---|---|---|---|
| Laska et al. 2001 | Danderyd, Sweden | acute stroke | 106 | 33.3 | 34.7 | 0.961 | 0.500 | 1.850 |
| Godefroy et al. 2002 | Lille, France | Acute stroke, Aphasia | 308 | 68.5 | 66.1 | 1.037 | 0.790 | 1.360 |
| Di Carlo et al. 2003 | 7 European countries | acute stroke | 4499 | 34.8 | 30.3 | 1.337 | 1.060 | 1.690 |
| Kelly-Hayes et al. 2003 | Framingham MA, USA | at 6 months post stroke | 108 | 23.8 | 11.6 | 2.143 | 0.780 | 5.900 |
| Roquer et al. 2003 | Barcelona, Spain | acute stroke | 1581 | 28.9 | 21.6 | 1.335 | 1.100 | 1.630 |
| Engelter et al. 2006 | Basel, Switzerland | acute ischemic stroke | 269 | 34.0 | 23.9 | 1.422 | 0.890 | 2.260 |
| Kyrozis et al. 2009 | Acadia, Greece | 28 days post-stroke | 555 | 27.6 | 18.8 | 1.473 | 1.040 | 2.090 |
| Tsouli et al. 2009 | Athens, Greece | acute stroke | 2297 | 41.3 | 31.5 | 1.313 | 1.140 | 1.510 |
| Brkic et al. 2009 | Tuzla, Bosnia and Herzegovina | acute stroke | 993 | 23.0 | 17.6 | 1.309 | 0.990 | 1.730 |
| Bersano et al. 2009a, <64 years | seven regions, Italy | acute stroke, <64 years | 1751 | 21.0 | 20.0 | 1.044 | 0.840 | 1.300 |
| Bersano et al. 2009b, 64-74 years | seven regions, Italy | acute stroke, 64-74 years | 2663 | 26.0 | 24.0 | 1.081 | 0.930 | 1.260 |
| Bersano et al. 2009c, 75-84 years | seven regions, Italy | acute stroke, 75-84 years | 2853 | 31.0 | 27.0 | 1.160 | 1.010 | 1.330 |
| Bersano et al. 2009d, >84 years | seven regions, Italy | acute stroke, 84< years | 1581 | 43.0 | 35.0 | 1.223 | 1.030 | 1.450 |
| Dickey et al. 2010 | Ontario, Canada | at discharge | 15327 | 33.4 | 31.0 | 1.078 | 1.020 | 1.140 |
| Gall et al. 2010 | Melbourne, Australia | acute stroke, Dysphasia | 843 | 46.1 | 35.6 | 1.294 | 1.040 | 1.600 |
| Gialanella et al. 2011 | Lumezzane, Italy | acute stroke | 262 | 55.9 | 44.4 | 1.258 | 0.890 | 1.770 |
| Croquelois & Bogousslavsky 2011 | Lausanne, Switzerland | acute stroke | 5880 | 28.1 | 24.9 | 1.128 | 1.020 | 1.250 |
| Jerath et al. 2011 | Rochester, USA | acute stroke | 449 | 45.8 | 37.4 | 1.222 | 0.910 | 1.640 |
| Kadojic et al. 2012 | Osijek, Croatia | acute ischemic stroke | 177 | 48.2 | 37.2 | 1.294 | 0.820 | 2.040 |
| Flowers et al. 2013 | Toronto, Canada | acute ischemic stroke | 221 | 35.7 | 26.0 | 1.373 | 0.850 | 2.220 |
| Wasserman et al. 2015 | Ottawa, Canada | isolated aphasia as only deficit of stroke | 1155 | 4.1 | 3.0 | 1.347 | 0.720 | 2.510 |
| Chang et al. 2015 | Colombo, Sri Lanka | data from questionnaire | 24 | 62.5 | 56.2 | 1.111 | 0.370 | 3.320 |
| **ALL** | | | **48362** | **29.6** | **26.0** | **1.139** | **1.100** | **1.180** |

Overview of the studies included in the meta-analysis on sex differences in post-stroke

aphasia rate.

**Fig 2. Meta-analysis forest plot.** Forest plot of aphasia rate ratios between males and

females for the 25 studies included in the meta-analysis (total n=48,362), showing that across

studies a small but significant effect of sex exists, indicating that women are more likely to

get aphasia from stroke. This effect, however, does not take age or stroke severity into

account.

Below, I report the excluded full-text papers (see Fig 1). This is done in order to give the

reader as complete an overview of the selection process as possible. As mentioned, studies

were included on the basis of the overall number of stroke patients being available for

calculation of aphasia prevalence. I excluded reviews [6, 9, 10, 47-54], studies with

insufficient information about sex [2, 55-87], studies with insufficient aphasia information

[88-93], studies with insufficiently specified sampling procedure for aphasia patients from

stroke cohorts [94-107], and studies with repetition of data use [39].

## Results of meta-analysis

The 25 studies included a total of 48,362 stroke patients (23,085 women, 25,297 men). Of

these 13,398 (6,828 women, 6,570 men) were diagnosed with aphasia (27.7%). 29.6 % of

female stroke patients were diagnosed, while 26 % of males were diagnosed with aphasia (see

Table 1 and Fig 2). This difference was found to be statistically significant using a paired and

weighted t-test on the aphasia rates across studies, weighted to add emphasis on studies with

larger patient samples, $t(27)=6.76$, $p<0.001$, forcing a rejection of the null-hypothesis that

there is no difference in aphasia rate between women and men. The overall sex aphasia rate

ratio was found to be 1.14 (1.10-1.18 95% CI) with a Cohen's $d$ of 0.37 which is usually

considered a small effect [108]. Low to medium heterogeneity between studies was observed:

$I^2$= 41 % (CI: 4%-77%) (see Fig 1 for a forest plot) and a funnel plot did not suggest any

outspoken bias in the reports (see Fig 3). Including publication year as a covariate in a

regression analysis did not alter the result and was not in itself a significant predictor of sex

differences (p>0.05).

**Fig 3. Funnel plot.** A Funnel plot did not indicate any outspoken bias in the meta-analysis data.

## Interim discussion

A higher aphasia rate after stroke for women than for men was found across studies in the meta-analysis. The average aphasia rate for women was 29.6% and for men 26% (see Table 1). The effect-size was in the small range.

The aphasia rate across studies and sexes (27.7%) was comparable to that reported in a recent meta-analysis (30%) [6]. The slightly lower estimate may in part be related to the inclusion of a study of cases with isolated aphasia [43] which had a much smaller aphasia rate than studies with a regular aphasia diagnosis (see Table 1). The aphasia rate ratio for the sexes reported on isolated aphasia by [43], however, was comparable to that of most other studies in the sample. Study year did not explain any variance in sex differences, suggesting that e.g. evolving diagnostic procedures for aphasia [26] has not had an observable impact on sex differences in aphasia rates. The forest plot in Fig 2 is ordered according to publication year on the y-axis, and given that there had been changes as a function of time, this should be visible as either a leftward or rightward trend in the data points. This seems not to be the case.

The fact that a higher aphasia rate after stroke for women than for men is found across studies in the meta-analysis contradicts the notion that language in men is more lateralized than in women (see introduction). If men have more lateralized language, one would expect their language to be more vulnerable to unilateral stroke than women's language, which, as shown, is not the case. But the findings are also at odds with previous critical suggestions that there are no sex differences in language lateralization between women and men [e.g. see 18, 109].

At face value, the findings would suggest that women in fact have more lateralized language

than men. There are, however, reasons to hesitate before arriving at such a conclusion, based

on the present analysis. As mentioned in the introduction, stroke is known to affect men and

women differently on a number of accounts, including general severity. The sexes also differ

on general health levels, resulting in women being older, on average, when they get a stroke

[9]. Age has previously been found to be a predictor of acquiring aphasia [53]. In order to

investigate if the effects of sex found in the meta-analysis are specific to language or may

relate to more general differences that are unlikely to be caused by a sex difference in core

language function, an investigation of aphasia rates that include additional explanatory

variables is needed. Unfortunately, very few studies in the current cohort make detailed,

reports of age effects on aphasia stratified for sex. One exception is Bersano et al. [12] who

report aphasia rates for 4 different age groups. Here an interaction between age and sex

differences seemingly can be observed. The sex difference is almost non-existing in the

youngest age group (under 64), but gradually grows larger and larger in older age groups (see

Table 1). It thus seems that taking age into account is important when trying to understand the

sex difference in aphasia rates.

Another possible explanation for the higher aphasia rate in women is that they may be

affected more severely by stroke in a non-discriminant manner [10]. If aphasia rates can be

explained by severity alone, it would again suggest that the sex difference is not restricted or

related to language in any meaningful way. But again, this type of information is not reported

in the papers covered by the present meta-analysis.

I have therefore added a 2$^{nd}$ dataset from an American healthcare database (see below) that

will allow me to investigate aphasia rates while taking age and stroke severity into account.

## Methods for database analysis

Data from the Healthcare Cost and Utilization Project (HCUP) from community hospitals in the United States were used for the analysis. The database (https://hcupnet.ahrq.gov/) contains inpatient diagnostic information using the International Classification of Diseases and Health Related Problems (ICD-9) codes from 35 US American states from the years 2011-2014. The ICD system is used by US hospitals for reimbursement purposes and subsequent research, e.g. to study patterns and outcome of disease [110]. Patient numbers from each year for each state, stratified by sex, was used in the analysis. Data from this database has previously been used to study post-stroke aphasia rates [2], but here we add sex as an explanatory variable and incorporate all available states for all the years in which the ICD-9 diagnoses were used (i.e. 10 times more patients).

To identify number of patients with stroke, the combined number of diagnoses from the database related to stroke was obtained. Cases with the following IDC-9 codes were included: "434.00 Crbl Thrmbs Wo Infrct", "434.01 Crbl Thrmbs W Infrct", "434.10 Crbl Emblsm Wo Infrct", "434.11 Crbl Emblsm W Infrct", "434.90 Crbl Art Oc Nos Wo Infrc", "434.91 Crbl Art Ocl Nos W Infrc", "436 Cva". To identify the number of patients with aphasia, the IDC-9 code: "784.3 Aphasia" was used. As a proxy for stroke severity, the number of hemiparesis/hemiplegia diagnoses were included (using the IDC-9 code: "342.90 Unsp Hemiplga Unspf Side", which is the most commonly used ICD-9 code for hemiplegia/paresis in the data). Hemiparesis, hemiplegia and aphasia are comorbid deficits. Around 90% of aphasia patients have hemiparesis [65], but if a sex difference in number of aphasias is accompanied by a similar sex difference in hemiparesis/hemiplegia diagnoses, then the difference is likely to be explained by stroke severity rather than being a specific language related phenomenon.

The database allows for two different ways to draw data. Either one can draw "Principal"

diagnoses or "all-listed" diagnoses. As aphasia is often unlikely to be the principal diagnosis

in a hospital visit, "all-listed" diagnoses were used. However, age information is only

available with "principal" diagnoses, and age information was therefore drawn from this data.

The assumption is that age differences in principal diagnosis will be representative for age

differences in the "all-listed" diagnoses as well. It is important to note that the database does

not offer information about age for any individual patient. These data are all summery data for

the participating states for a particular year (2011-2014). Yet, similarly to a meta-analysis of

data from different studies, summary data from different states, each representing a large

group of patients may nevertheless be representative of generalizable differences between

groups (males and females) in the data.

To evaluate statistical significance of these findings, a linear mixed-effects regression analysis

was conducted, fit by REML, using the *lmertest* package in R [111]. P-values were estimated

using Satterthwaite's method. The model incorporated aphasia rate as the dependent variable

and sex as the main fixed dependent variable. Age and rate of hemiplegia diagnoses (proxy

for stroke severity) were z-score scaled and added as additional covariates. The model also

included all possible interactions between the three variables. US state and year for each data-

point were included as random effects. The regression was weighted by number of stroke

cases in a particular state/year, to put more weight on data-points from larger states.

## Results of database analysis

A total of 1,655,295 stroke patients were found in the database (863,612 women, 791,683

men) in the period from 2011 to 2014. Aphasia was diagnosed in 623,942 cases (336,604

women, 287,338 men) or 37.7%. Using this method, 38.9 % of female stroke patients were

diagnosed, while 36.2 % of males were diagnosed with aphasia (see Fig 4).

**Fig 4**. **US data forest plot.** Aphasia rate ratios (uncorrected for age) for each US state in the

HCUP database from 2011-2014. This analysis replicates the findings from the meta-analysis

and provides unequivocal evidence for a higher aphasia rate among women compared to men

given stroke (see Fig 1, but note the scale difference between plots). However, as Fig 5

shows, this effect can be explained completely by the sex difference in age at stroke.

The overall female/male aphasia rate ratio was found to be 1.073 (1.068-1.079 95% CI) with a

Cohen's $d$ effect size across states of 0.63 which is usually considered a medium effect size

[108]. A paired t-test again yielded support to the existence of a sex difference, $t(143)=-13.74$,

$p<0.001$.

281
When including age and stroke severity in a regression analysis, however, no significant

effect of sex over and above that explained by age and severity could be observed,

$t(272.33)=-0.826$, $p=0.4$. A significant effect of age, $t(275.23)=2.21$, $p(uncorrected)<0.05$,

and a significant effect of stroke severity, $t(268.66)=5.34$, $p<0.001$ was observed. Fig 5

displays how sex is completely confounded by age of stroke and does not add any explanatory

power to the analysis. A significant interaction between age and stroke severity was also

observed, $t(275.9)=-2.30$, $p(uncorrected)<0.05$. No other interactions were significant.

**Fig 5. Aphasia rate as a function of age at stroke.** A scatterplot of stroke average age

against aphasia rate for each US state and year (2011-2014) in the HCUP database. The plot

illustrates the large age difference between men and women at time of stroke. It also shows a

positive correlation between average age and aphasia rate, suggesting that older stroke

patients more often get aphasia. When this relationship is taken into account, sex effects are

no longer significant in the aphasia rates.

## Discussion

On this very large cohort of patients, we replicate the findings from the meta-analysis. Based

on raw aphasia rates, women are more likely to get an aphasia diagnosis following stroke than

men. The sizes of the effects vary somewhat across the two analyses. The aphasia ratio is

smaller for the database study, but the effect size is found to be larger. This is likely due to the

fact that the database data is more homogeneous than the data included in the meta-analysis,

where the time of diagnosis post-stroke differs and also to some extent the definition of

aphasia (e.g. one study includes dysphasia as well as aphasia and one study only looks at

isolated aphasia cases – see Table 1 for details).

At the same time, our $2^{nd}$ analysis completely revokes the conclusions that one could have

drawn from the meta-analysis. In the database, we find no evidence of any sex difference in

aphasia rates over and above that which can be explained by the age differences between

females and males at time of stroke. This replicates previous findings that age is a predictor of

aphasia following stroke [53], and given that age is a more fundamental causal variable than

language (i.e. your language cannot change your age, but the opposite may be true), it is likely

that most if not all of the sex differences in aphasia rates is caused by the age difference in

stroke between women and men.

We also find an independent effect of stroke severity on aphasia rates as measured by

diagnoses of hemiplegia. Aphasia and hemiparesis/hemiplegia are known to be highly co-

morbid. In this study we find that severity effects on aphasia are independent of the sex

effects. The sex differences thus do not seem to be related to stroke severity per se. It has to

be said, however, that this analysis uses a somewhat crude proxy for stroke severity. Other

measures, such as general stroke scale scores [112, 113] might interact more with sex.

Bersano et al. [12] found indications of increasing sex differences in aphasia rates with age.

Contrary to this, the database analysis shows no indication of an interaction between sex and

age. Bersano and co-workers did not report inferential statistics documenting an actual

interaction, but looking at their data, the increasing discrepancy in aphasia between males and

females with age is striking. For patients below 64 years the aphasia rate gender ratio is 1.04

and grows to 1.08 in 64-74 year old patients, 1.16 in 75-84 and 1.22 in patients above 84

years of age (see Table 1). How does this fit with the current data not showing any interaction

between age and sex? One possible explanation is that there is an inherent bias in the way that

the Bersano and co-workers' age data is distributed. When lumping the data into 10 year age

bins, one needs to take into account that women and men may not be equally distributed

within each bin. The data from the Bersano et al. study were collected in 2001 in Italy. If one

looks at the gender and age distribution of the Italian population in January 2002 using

population statistics (http://demo.istat.it/pop2002/index_e.html), one finds that because

women live longer than men, the average age of women within the different age bins from

middle age and onwards is higher than that of men and that this difference gets larger for the

older groups. For the 54 to 64 year old Italians, the mean age difference between men and

women is 0.07 years, but for the age group above 84 years, it has grown to 0.52 years. There

is a very strong linear correlation between these mean age differences in the Italian population

in these age bins and the reported differences in aphasia rate ($r = 0.96$, data available from

author on request), which suggests that at least some of the interaction between sex and age

reported by Bersano and co-workers is based on unequal sampling of the different ages. This

is not to say that there could not be sex and age interactions that are not picked up by the

current analysis. The data from the database is distributed on a state by year basis and each

data-point for age is the result of averaging across many individual patients. Underneath this

gross data reduction may be hidden lots of interesting variability. Further studies are needed

in order to rule out a potential interaction between sex, age and aphasia.

The present analyses are also limited in that they say nothing about the different types of

aphasia symptoms that patients may suffer from and the potential interactions that might be

found with gender if one looks more carefully at aphasia subtypes.

Taken together, the results are in line with a critical stance towards any large-scale brain base

for sex differences in language [18]. This, of course, does not mean that the observed sex

differences in language related behavior (see introduction) do not have brain correlates, just

that these differences will be dynamic, complex and to a large extent dependent on gender

differences in experience and context rather than being tied to genetic sex.

## Conclusion

358    Women were found to be diagnosed with aphasia following stroke more often than men. This

359    is in direct opposition to the hypothesis that women have less lateralized language function

360    than men. The sex difference is most likely caused by age differences in the two groups at the

361    time of stroke.

## Acknowledgements

363    The author has no conflict of interests to report.

# References

1.  Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, et al. Heart Disease and Stroke Statistics-2017 Update: A Report From the American Heart Association. Circulation. 2017;135(10):e146-e603.

2.  Ellis C, Hardy RY, Lindrooth RC, Peach RK. Rate of aphasia among stroke patients discharged from hospitals in the United States. Aphasiology. 2018;32(9):1075-86.

3.  Inatomi Y, Yonehara T, Omiya S, Hashimoto Y, Hirano T, Uchino M. Aphasia during the acute phase in ischemic stroke. Cerebrovasc Dis. 2008;25(4):316-23.

4.  Tsouli S, Kyritsis AP, Tsagalis G, Virvidaki E, Vemmos KN. Significance of Aphasia after First-Ever Acute Stroke: Impact on Early and Late Outcomes. Neuroepidemiology. 2009;33(2):96-102.

5.  Godefroy O, Dubois C, Debachy B, Leclerc M, Kreisler A. Vascular Aphasias. Main Characteristics of Patients Hospitalized in Acute Stroke Units. Stroke. 2002;33(3):702.

6.  Flowers HL, Skoretz SA, Silver FL, Rochon E, Fang J, Flamand-Roze C, et al. Poststroke Aphasia Frequency, Recovery, and Outcomes: A Systematic Review and Meta-Analysis. Archives of Physical Medicine and Rehabilitation. 2016;97(12):2188-201.e8.

7.  Thommessen B, Thoresen GE, Bautz-Holter E, Laake K. Validity of the Aphasia Item from the Scandinavian Stroke Scale. Cerebrovascular Diseases. 2002;13(3):184-6.

8.  Meyer BC, Lyden PD. The modified National Institutes of Health Stroke Scale: Its Time Has Come. International Journal of Stroke. 2009;4(4):267-73.

9.  Appelros P, Stegmayr B, Terént A. Sex differences in stroke epidemiology: a systematic review. Stroke. 2009;40(4):1082-90.

391    10.    Berglund A, Schenck-Gustafsson K, von Euler M. Sex differences in the presentation

392           of stroke. Maturitas. 2017;99:47-50.

393    11.    Gall SL, Donnan G, Dewey HM, Macdonell R, Sturm J, Gilligan A, et al. Sex

394           differences in presentation, severity, and management of stroke in a population-based

395           study. Neurology. 2010;74(12):975.

396    12.    Bersano A, Burgio F, Gattinoni M, Candelise L. Aphasia Burden to Hospitalised

397           Acute Stroke Patients: Need for an Early Rehabilitation Programme. International

398           Journal of Stroke. 2009;4(6):443-7.

399    13.    Pedersen PM, Jørgensen HS, Nakayama H, Raaschou HO, Olsen TS. Aphasia in acute

400           stroke: incidence, determinants, and recovery. Annals of Neurology. 1995;38(4):659-

401           66.

402    14.    Miceli G, Caltagirone C, Gainotti G, Masullo C, Silveri Maria C, Villa G. Influence of

403           age, sex, literacy and pathologic lesion on incidence, severity and type of aphasia.

404           Acta Neurologica Scandinavica. 1981;64(5):370-82.

405    15.    Bleses D, Vach W, Slott M, Wehberg S, Thomsen P, Madsen T, et al. The Danish

406           Communicative Developmental Inventories: validity and main developmental trends. J

407           Child Lang. 2008;35(03):1-19.

408    16.    Reilly D, Neumann DL, Andrews G. Gender differences in reading and writing

409           achievement: Evidence from the National Assessment of Educational Progress

410           (NAEP). American Psychologist. in press.

411    17.    Schwartz HA, Eichstaedt JC, Kern ML, Dziurzynski L, Ramones SM, Agrawal M, et

412           al. Personality, gender, and age in the language of social media: the open-vocabulary

413           approach. PLoS ONE. 2013;8(9):e73791.

414    18.    Wallentin M. Putative sex differences in verbal abilities and language cortex: a critical

415           review. Brain and Language. 2009;108(3):175-83.

416    19.    Kansaku K, Kitazawa S. Imaging studies on sex differences in the lateralization of

417            language. Neurosci Res. 2001;41(4):333-7.

418    20.    Shaywitz BA, Shaywitz SE, Pugh KR, Constable RT, Skudlarski P, Fulbright RK, et

419            al. Sex differences in the functional organization of the brain for language. Nature.

420            1995;373(6515):607-9.

421    21.    Baron-Cohen S, Knickmeyer RC, Belmonte MK. Sex Differences in the Brain:

422            Implications for Explaining Autism. Science. 2005;310(5749):819-23.

423    22.    Hausmann M. Why sex hormones matter for neuroscience: A very short review on

424            sex, sex hormones, and functional brain asymmetries. Journal of Neuroscience

425            Research. 2016;95(1-2):40-9.

426    23.    Sommer IEC, Aleman A, Bouma A, Kahn RS. Do women really have more bilateral

427            language representation than men? A meta-analysis of functional imaging studies.

428            Brain. 2004;127(8):1845-52.

429    24.    Hirnstein M, Hugdahl K, Hausmann M. Cognitive sex differences and hemispheric

430            asymmetry: A critical review of 40 years of research. Laterality: Asymmetries of

431            Body, Brain and Cognition. 2018:1-49.

432    25.    Moher D, Liberati A, Tetzlaff J, Altman DG, The PG. Preferred Reporting Items for

433            Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLOS Medicine.

434            2009;6(7):e1000097.

435    26.    Rohde A, Worrall L, Godecke E, O'Halloran R, Farrell A, Massey M. Diagnosis of

436            aphasia in stroke populations: A systematic review of language tests. PLOS ONE.

437            2018;13(3):e0194143.

438    27.    Viechtbauer W. Conducting meta-analyses in R with the metafor package. Journal of

439            Statistical Software. 2010;36(3):1-48.

440    28.    Croquelois A, Bogousslavsky J. Stroke Aphasia: 1,500 Consecutive Cases.

441            Cerebrovascular Diseases. 2011;31(4):392-9.

442    29.    Kyrozis A, Potagas C, Ghika A, Tsimpouris PK, Virvidaki ES, Vemmos KN.

443            Incidence and predictors of post-stroke aphasia: The Arcadia Stroke Registry.

444            European Journal of Neurology. 2009;16(6):733-9.

445    30.    Laska AC, Hellblom A, Murray V, Kahan T, Von Arbin M. Aphasia in acute stroke

446            and relation to outcome. Journal of Internal Medicine. 2001;249(5):413-22.

447    31.    Roquer J, Campello AR, Gomis M. Sex Differences in First-Ever Acute Stroke.

448            Stroke. 2003;34(7):1581.

449    32.    Brkić E, Sinanović O, Vidović M, Smajlović D. Incidence and clinical

450            phenomenology of aphasic disorders after stroke [Ucestalost i klinicka fenomenologija

451            afazickih poremećaja nakon mozdanog udara.]. Medicinski arhiv. 2009;63(4):197-9.

452    33.    Brust JC, Shafer SQ, Richter RW, Bruun B. Aphasia in acute stroke. Stroke.

453            1976;7(2):167.

454    34.    Hier DB, Yoon WB, Mohr JP, Price TR, Wolf PA. Gender and Aphasia in the Stroke

455            Data Bank. Brain and Language. 1994;47(1):155-67.

456    35.    Engelter ST, Gostynski M, Papa S, Frei M, Born C, Ajdacic-Gross V, et al.

457            Epidemiology of Aphasia Attributable to First Ischemic Stroke. Stroke.

458            2006;37(6):1379.

459    36.    Flowers HL, Silver FL, Fang J, Rochon E, Martino R. The incidence, co-occurrence,

460            and predictors of dysphagia, dysarthria, and aphasia after first-ever acute ischemic

461            stroke. Journal of Communication Disorders. 2013;46(3):238-48.

462    37.    Dickey L, Kagan A, Lindsay MP, Fang J, Rowland A, Black S. Incidence and Profile

463            of Inpatient Stroke-Induced Aphasia in Ontario, Canada. Archives of Physical

464            Medicine and Rehabilitation. 2010;91(2):196-202.

38.    Kadojić D, Rostohar Bijelić B, Radanović R, Porobić M, Rimac J, Dikanović M.

Aphasia in Patients with Ischemic Stroke. Acta clinica Croatica. 2012;51(2):221-4.

39.    Gialanella B, Bertolinelli M, Lissi M, Prometti P. Predicting outcome after stroke: the

role of aphasia. Disability and Rehabilitation. 2011;33(2):122-9.

40.    Di Carlo A, Lamassa M, Baldereschi M, Pracucci G, Basile AM, Wolfe CDA, et al.

Sex Differences in the Clinical Presentation, Resource Use, and 3-Month Outcome of

Acute Stroke in Europe. Stroke. 2003;34(5):1114.

41.    Kelly-Hayes M, Beiser A, Kase CS, Scaramucci A, D'Agostino RB, Wolf PA. The

influence of gender and age on disability following ischemic stroke: the Framingham

study. Journal of Stroke and Cerebrovascular Diseases. 2003;12(3):119-26.

42.    Jerath NU, Reddy C, Freeman WD, Jerath AU, Brown RD. Gender Differences in

Presenting Signs and Symptoms of Acute Ischemic Stroke: A Population-Based

Study. Gender Medicine. 2011;8(5):312-9.

43.    Wasserman JK, Perry JJ, Dowlatshahi D, Stotts G, Sivilotti MLA, Worster A, et al.

Isolated transient aphasia at emergency presentation is associated with a high rate of

cardioembolic embolism. CJEM. 2015;17(6):624-30.

44.    Chang T, Gajasinghe S, Arambepola C. Prevalence of Stroke and Its Risk Factors in

Urban Sri Lanka. Stroke. 2015;46(10):2965.

45.    Scarpa M, Colombo A, Sorgato P, De Renzi E. The Incidence of Aphasia and Global

Aphasia in Left Brain-Damaged Patients. Cortex. 1987;23(2):331-6.

46.    Siirtola M, Narva EV, Siirtola T. On the occurrence and prognosis of aphasia in

patients with cerebral infarction. Scand J Soc Med Suppl. 1977;14:128-33.

47.    Crinion JT, Leff AP. Recovery and treatment of aphasia after stroke: functional

imaging studies. Current Opinion in Neurology. 2007;20(6):667-73.

48.    Watila MM, Balarabe SA. Factors predicting post-stroke aphasia recovery. Journal of

490            the Neurological Sciences. 2015;352(1):12-8.

491    49.    Plowman E, Hentz B, Ellis C. Post-stroke aphasia prognosis: a review of patient-

492            related and stroke-related factors. Journal of Evaluation in Clinical Practice.

493            2011;18(3):689-94.

494    50.    Jongbloed L. Prediction of function after stroke: a critical review. Stroke.

495            1986;17(4):765.

496    51.    Ferro JM, Mariano G, Madureira S. Recovery from Aphasia and Neglect.

497            Cerebrovascular Diseases. 1999;9(suppl 5)(Suppl. 5):6-22.

498    52.    Vuković V, Galinović I, Lovrenčić-Huzjan A, Budišić M, Demarin V. Women and

499            Stroke: How Much do Women and Men Differ? A Review – Diagnostics, Clinical

500            Differences, Therapy and Outcome. Collegium antropologicum. 2009;33(3):977-84.

501    53.    Ellis C, Urban S. Age and aphasia: a review of presence, type, recovery and clinical

502            outcomes. Topics in Stroke Rehabilitation. 2016;23(6):430-9.

503    54.    Lazar RM, Boehme AK. Aphasia As a Predictor of Stroke Outcome. Current

504            Neurology and Neuroscience Reports. 2017;17(11):83.

505    55.    Hoffmann M, Chen R. The Spectrum of Aphasia Subtypes and Etiology in Subacute

506            Stroke. Journal of Stroke and Cerebrovascular Diseases. 2013;22(8):1385-92.

507    56.    Leśniak M, Bak T, Czepiel W, Seniów J, Członkowska A. Frequency and Prognostic

508            Value of Cognitive Disorders in Stroke Patients. Dementia and Geriatric Cognitive

509            Disorders. 2008;26(4):356-63.

510    57.    Lubart E, Leibovitz A, Baumoehl Y, Klein C, Gil I, Abramovitz I, et al. Progressing

511            stroke with neurological deterioration in a group of Israeli elderly. Archives of

512            Gerontology and Geriatrics. 2005;41(1):95-100.

58.     Paolucci S, Antonucci G, Gialloreti LE, Traballesi M, Lubich S, Pratesi L, et al.

Predicting Stroke Inpatient Rehabilitation Outcome: The Prominent Role of

Neuropsychological Disorders. European Neurology. 1996;36(6):385-90.

59.     Stegmayr B, Asplund K, Wester PO. Trends in incidence, case-fatality rate, and

severity of stroke in northern Sweden, 1985-1991. Stroke. 1994;25(9):1738.

60.     Stürmer T, Schlindwein G, Kleiser B, Roempp A, Brenner H. Clinical Diagnosis of

Ischemic versus Hemorrhagic Stroke: Applicability of Existing Scores in the

Emergency Situation and Proposal of a New Score. Neuroepidemiology.

2002;21(1):8-17.

61.     Tilling K, Sterne JAC, Rudd AG, Glass TA, Wityk RJ, Wolfe CDA. A New Method

for Predicting Recovery After Stroke. Stroke. 2001;32(12):2867.

62.     Troisi E, Paolucci S, Silvestrini M, Matteis M, Vernieri F, Grasso MG, et al.

Prognostic factors in stroke rehabilitation: the possible role of pharmacological

treatment. Acta Neurologica Scandinavica. 2002;105(2):100-6.

63.     Vidović M, Sinanović O, Šabaškić L, Hatičić A, Brkić E. Incidence and Types of

Speech Disorders in Stroke Patients. Acta clinica Croatica. 2011;50(4):491-3.

64.     Lazar RM, Speizer AE, Festa JR, Krakauer JW, Marshall RS. Variability in language

recovery after first-time stroke. Journal of Neurology, Neurosurgery &

Psychiatry. 2008;79(5):530.

65.     Wade DT, Hewer RL, David RM, Enderby PM. Aphasia after stroke: natural history

and associated deficits. Journal of Neurology, Neurosurgery & Psychiatry.

1986;49(1):11.

66.     Candelise L, Gattinoni M, Bersano A, Micieli G, Sterzi R, Morabito A. Stroke-unit

care for acute stroke patients: an observational follow-up study. The Lancet.

2007;369(9558):299-305.

67.  Hillis AE, Heidler J. Mechanisms of early aphasia recovery. Aphasiology.

2002;16(9):885-95.

68.  Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, et al.

Heart Disease and Stroke Statistics—2010 Update. Circulation. 2010;121(7):e46.

69.  Maas MB, Lev MH, Ay H, Singhal AB, Greer DM, Smith WS, et al. The Prognosis

for Aphasia in Stroke. Journal of Stroke and Cerebrovascular Diseases.

2012;21(5):350-7.

70.  Kremer C, Perren F, Kappelin J, Selariu E, Abul-Kasim K. Prognosis of aphasia in

stroke patients early after iv thrombolysis. Clinical Neurology and Neurosurgery.

2013;115(3):289-92.

71.  Pedersen PM, Vinter K, Olsen TS. Aphasia after Stroke: Type, Severity and

Prognosis. Cerebrovascular Diseases. 2004;17(1):35-43.

72.  Hachioui HE, van de Sandt-Koenderman MWME, Dippel DWJ, Koudstaal PJ, Visch-

Brink EG. The Screeling: Occurrence of Linguistic Deficits in Acute Aphasia Post-

Stroke. Journal of Rehabilitation Medicine. 2012;44(5):429-35.

73.  Fennis TFM, Compter A, van den Broek MWC, Koudstaal PJ, Algra A, Koehler PJ. Is

Isolated Aphasia a Typical Presentation of Presumed Cardioembolic Transient

Ischemic Attack or Stroke? Cerebrovascular Diseases. 2013;35(4):337-40.

74.  Gialanella B, Santoro R, Ferlucci C. Predicting outcome after stroke: the role of basic

activities of daily living predicting outcome after stroke. European Journal of Physical

and Rehabilitation Medicine. 2013;49(3):629-37.

75.  González-Fernández M, Davis C, Molitoris JJ, Newhart M, Leigh R, Hillis AE.

Formal Education, Socioeconomic Status, and the Severity of Aphasia After Stroke.

Archives of Physical Medicine and Rehabilitation. 2011;92(11):1809-13.

76.    Kemper C, Koller D, Glaeske G, van den Bussche H. Mortality and Nursing Care

563           Dependency One Year After First Ischemic Stroke: An Analysis of German Statutory

564           Health Insurance Data. Topics in Stroke Rehabilitation. 2011;18(2):172-8.

565    77.    Miller N, Gray WK, Howitt SC, Jusabani A, Swai M, Mugusi F, et al. Aphasia and

566           Swallowing Problems in Subjects With Incident Stroke in Rural Northern Tanzania: A

567           Case-Control Study. Topics in Stroke Rehabilitation. 2014;21(1):52-62.

568    78.    Ali M, Bath PM, Lyden PD, Bernhardt J, Brady M. Representation of People with

569           Aphasia in Randomized Controlled Trials of Acute Stroke Interventions. International

570           Journal of Stroke. 2014;9(2):174-82.

571    79.    Ojala-Oksala J, Jokinen H, Kopsi V, Lehtonen K, Luukkonen L, Paukkunen A, et al.

572           Educational History Is an Independent Predictor of Cognitive Deficits and Long-Term

573           Survival in Postacute Patients With Mild to Moderate Ischemic Stroke. Stroke.

574           2012;43(11):2931.

575    80.    Yuan S-M, Humuruola G. Stroke of a cardiac myxoma origin. Brazilian Journal of

576           Cardiovascular Surgery. 2015;30:225-34.

577    81.    Giaquinto S, Buzzelli S, Francesco L, Lottarini A, Montenero P, In PT, et al. On the

578           prognosis of outcome after stroke. Acta Neurologica Scandinavica. 1999;100(3):202-

579           8.

580    82.    Worrall LE, Hudson K, Khan A, Ryan B, Simmons-Mackie N. Determinants of

581           Living Well With Aphasia in the First Year Poststroke: A Prospective Cohort Study.

582           Archives of Physical Medicine and Rehabilitation. 2017;98(2):235-40.

583    83.    Kauhanen ML, Korpelainen JT, Hiltunen P, Määttä R, Mononen H, Brusin E, et al.

584           Aphasia, Depression, and Non-Verbal Cognitive Impairment in Ischaemic Stroke.

585           Cerebrovascular Diseases. 2000;10(6):455-61.

586    84.    Bogousslavsky J, Van Melle G, Regli F. The Lausanne Stroke Registry: analysis of

587           1,000 consecutive patients with first stroke. Stroke (1970). 1988;19(9):1083-92.

588    85.    Haselbach D, Renggli A, Carda S, Croquelois A. Determinants of Neurological

589           Functional Recovery Potential after Stroke in Young Adults. Cerebrovascular

590           Diseases Extra. 2014;4(1):77-83.

591    86.    Maehlum S, Roaldsen K, Kolsrud M, Dahl M. Rehabilitering etter hjerneslag

592           [Rehabilitation after a stroke]. Tidsskrift for Den norske legeforening. 1990;110:2657-

593           9.

594    87.    Wade DT, Hewer RL, Wood VA. Stroke: influence of patient's sex and side of

595           weakness on outcome. Archives of physical medicine and rehabilitation.

596           1984;65(9):513-6.

597    88.    Kelly PJ, Furie KL, Shafqat S, Rallis N, Chang Y, Stein J. Functional recovery

598           following rehabilitation after hemorrhagic and ischemic stroke11No commercial party

599           having a direct financial interest in the results of the research supporting this article

600           has or will confer a benefit on the authors or on any organization with which the

601           authors are associated. Archives of Physical Medicine and Rehabilitation.

602           2003;84(7):968-72.

603    89.    Ween JE, Mernoff ST, Alexander MP. Recovery Rates after Stroke and Their Impact

604           on Outcome Prediction. Neurorehabilitation and Neural Repair. 2000;14(3):229-35.

605    90.    Fuh JL, Wang SJ, Larson EB, Liu HC. Prevalence of Stroke in Kinmen. Stroke.

606           1996;27(8):1338.

607    91.    Matsumoto N, Whisnant JP, Kurland LT, Okazaki H. Natural History of Stroke in

608           Rochester, Minnesota, 1955 Through 1969: An Extension of a Previous Study, 1945

609           Through 1954. Stroke. 1973;4(1):20.

92.   Safaz I, Kesikburun S, Adigüzel E, Yilmaz B. Determinants of disease-specific health-
related quality of life in Turkish stroke survivors. International Journal of
Rehabilitation Research. 2016;39(2).

93.   Kunitz SC, Gross CR, Heyman A, Kase CS, Mohr JP, Price TR, et al. The pilot Stroke
Data Bank: definition, design, and data. Stroke. 1984;15(4):740-6.

94.   Sarno MT, Buonaguro A, Levita E. Gender and recovery from aphasia after stroke. J
Nerv Ment Dis. 1985;173(10):605-9.

95.   Pickersgill MJ, Lincoln NB. Prognostic indicators and the pattern of recovery of
communication in aphasic stroke patients. Journal of Neurology, Neurosurgery
& Psychiatry. 1983;46(2):130.

96.   Law J, Rush R, Pringle AM, Irving AM, Huby G, Smith M, et al. The incidence of
cases of aphasia following first stroke referred to speech and language therapy
services in Scotland. Aphasiology. 2009;23(10):1266-75.

97.   Jodzio K, Drumm DA, Nyka WM, Lass P, Gąsecki D. The contribution of the left and
right hemispheres to early recovery from aphasia: A SPECT prospective study.
Neuropsychological Rehabilitation. 2005;15(5):588-604.

98.   De Renzi E, Faglioni P, Ferrari P. The Influence of Sex and Age on the Incidence and
Type of Aphasia. Cortex. 1980;16(4):627-30.

99.   Basso A, Capitani E, Moraschini S. Sex Differences in Recovery from Aphasia.
Cortex. 1982;18(3):469-75.

100.  Oliveira FF, Damasceno BP. Short-term prognosis for speech and language in first
stroke patients. Arquivos de Neuro-Psiquiatria. 2009;67:849-55.

101.  Sundet K. Sex differences in severity and type of aphasia1. Scandinavian Journal of
Psychology. 1988;29(3-4):168-79.

102.    Bhatnagar SC, Jain SK, Bihari M, Bansal NK, Pauranik A, Jain DC, et al. Aphasia

635            type and aging in Hindi-speaking stroke patients. Brain and Language.

636            2002;83(2):353-61.

637    103.    Eslinger PJ, Damasio AR. Age and type of aphasia in patients with stroke. Journal of

638            Neurology, Neurosurgery & Psychiatry. 1981;44(5):377.

639    104.    Kertesz A, Benke T. Sex equality in intrahemispheric language organization. Brain

640            and Language. 1989;37(3):401-8.

641    105.    Pizzamiglio L, Mammucari A, Razzano C. Evidence for sex differences in brain

642            organization in recovery in aphasia. Brain and Language. 1985;25(2):213-23.

643    106.    Kertesz A, Sheppard ANN. The Epidemiology of Aphasic and Cognitive impairment

644            in Stroke: Age, Sex, Aphasia Type and Laterality Differences. Brain.

645            1981;104(1):117-28.

646    107.    McGlone J, Kertesz A. Sex Differences in Cerebral Processing of Visuospatial Tasks.

647            Cortex. 1973;9(3):313-20.

648    108.    Cohen J. A power primer. Psychological bulletin. 1992;112(1):155-9.

649    109.    Wallentin M, Michaelsen JLD, Rynne I, Nielsen RH. Lateralized task shift effects in

650            Broca's and Wernicke's regions and in visual word form area are selective for

651            conceptual content and reflect trial history. NeuroImage. 2014;101:276-88.

652    110.    O'Malley KJ, Cook KF, Price MD, Wildes KR, Hurdle JF, Ashton CM. Measuring

653            Diagnoses: ICD Code Accuracy. Health Services Research. 2005;40(5p2):1620-39.

654    111.    Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear

655            Mixed Effects Models. Journal of Statistical Software. 2017;82(13).

656    112.    Lindenstrøm E, Boysen G, Waage Christiansen L, à Rogvi Hansen B, Würtzen

657            Nielsen P. Reliability of Scandinavian Neurological Stroke Scale. Cerebrovascular

658            Diseases. 1991;1(2):103-7.

659    113.    Hantson L, De Weerdt W, De Keyser J, Diener HC, Franke C, Palm R, et al. The

660            European Stroke Scale. Stroke. 1994;25(11):2215-9.

## Supporting information

**S1 File. PRISMA Checklist.** Checklist for meta-analyses according to http://www.prisma-
statement.org

Female/Male Aphasia Rate Ratio + 95% CI