Ancestry informative alleles captured with reduced representation

library sequencing in *Theobroma cacao*

Jaime A. Osorio-Guarín[1], Corey R. Quackenbush[2,3], Omar E. Cornejo[2]

[1]Centro de Investigación Tibaitatá, Corporación Colombiana de Investigación Agropecuaria

– Agrosavia, Mosquera, Cundinamarca, Colombia.

[2]School of Biological Sciences, Washington State University, Pullman, WA, United States.

[3]Division of Molecular and Translational Sciences, U. S. Army Medical Research Institute

of Infectious Diseases (USAMRIID), Fort Detrick, Md, United States.

14   * Corresponding author: Omar E. Cornejo

15   E-mail: omar.cornejo@wsu.edu (OC)

Abstract

As the source of chocolate, cacao has become one of the most important crops in the world. The identification of molecular markers to understand the demographic history, genetic diversity and population structure plays a pivotal role in cacao breeding programs. Here, we report the use of a modified genotyping-by-sequencing (GBS) approach for large-scale single nucleotide polymorphism (SNP) discovery and allele ancestry mapping. We identified 12,357 bi-allelic SNPs after filtering, of which, 7,009 variants were ancestry informative. The GBS approach proved to be rapid, cost-effective, and highly informative for ancestry assignment in this species.

Introduction

Theobroma cacao L, the tree from which chocolate is confectioned, originated in the Amazonian basin [1] and has become a globally cultivated crop in recent times [2]. There are more than 20 species within the *Theobroma* genus, which belongs to the Malvaceae family [3]. *T. cacao* is the only species that is cultivated extensively because it can be manufactured into chocolate, liquors, confections, cosmetics, and animal feed [4,5]. Chemical and archaeological evidence shows that human settlements have been consuming cacao since 1,400 BC [6,7]. Since 900 BC *T. cacao* has been widely cultivated in Mesoamerica, which is considered the center of domestication. Recent genetic work has supported these findings suggesting that domestication of cacao occurred 3.6 Kya [8]. The domestication of cacao has resulted in the development of an important crop that sustains the livelihood of communities throughout Central and South America, Asia, and Africa [1,9,10], and is the principal income

for about six million smallholder farmers globally that produce up to 90% of the world's

cocoa [11].

Cacao is a diploid species with a chromosome complement of 2n=20 [12]. Recently, genomic

resources have been developed to facilitate the study of the species evolutionary history, the

inference of the genetic basis of phenotypes of interest, and the development of marker

assisted breeding programs. There are two reference genomes that have been sequences from

different cultivars, the B97-61/B2 genome (Criollo), with an estimated size of 409 million

base pairs (Mbp) [13] and the Matina 1-6 genome (Amelonado), with an estimated genome

size of 445 Mbp [14]. In addition, Cornejo et al. 2017 [8], performed whole genome

sequencing on 200 cultivars in order to understand the domestication process in this species.

Recently, approaches to infer genetic diversity in *T. cacao* have been completed using a small

panel of single nucleotide polymorphisms (SNPs, ranging from 87 to 6,000) developed with

a Fluidigm array system [15–17]. Also, using the information generated from a subset of the

200 cultivars project, an Ilumina Infinity II genotyping array of 13,530 markers was recently

developed [18]. However, it is well known that genotyping strategies based on arrays are

biased towards the capture of known specific loci in reference populations which introduces

ascertainment bias in the genotyping of samples from different origins [19]. There is a

growing interest in developing strategies to rapidly genotype accessions at a reduced cost in

order to perform extensive association studies and better identify accessions with specific

markers that will accelerate assisted breeding programs.

The generation of genomic resources in cacao has benefited from advances in sequencing

chemistry, cost reductions, and increased computational capacity to analyze data [20]. Yet,

despite the general reduction of costs, the sequencing of complete cacao genomes remains a

costly approach for breeders and researchers. Due to the high cost of whole genome

sequencing, genotyping arrays are typically used. However, genotyping arrays suffer from

the aforementioned issues of ascertainment bias, which hinders the identification of novel

untapped genetic variation in the species. It has been shown that genotyping-by-sequencing

(GBS) is a cost effective and efficient methodology for the generation of a large number of

SNPs, while avoiding ascertainment biases [21,22]. The genetic information generated by

GBS has been shown to facilitate robust studies of genetic diversity, population structure,

phylogeny, and association in a large number of plant and animal species [23–29] but have

never been applied to the study of *T. cacao*. The original protocol proposed for the digestion

and generation of reduced representation genomic libraries used the *ApeKI* enzyme [21].

However, Cooke et al. (2016) [30], demonstrated that the use of enzymes that cut far from

the recognition site, such as *BsaXI*, increases the diversity of the library and therefore the

number and quality of reads sequenced. In the original protocol the restriction enzymes

would produce libraries with nearly identical initial sequence causing saturation of the image

capture during sequencing. To avoid this issue, the original protocol incorporated the barcode

in one of the adapter sequences upstream of the cut-site to introduce variation and minimize

the problems associated with saturation [31]. In the proposed protocol (taken from Cooke et

al. 2016 [30]), the chosen enzyme generates libraries in which the initial sequence is random

due to the fact that the enzyme cuts 14 base pairs (bp) away from its recognition site.

In this work, we adapt and modify an existing GBS method [30] to generate a large number

of ancestry informative SNPs in cacao. We further show that the data generated produced

genetic diversity estimates consistent with what has been shown in previous work. We also

provide an estimate of the proportion of novel versus common SNPs (based on the relative

frequency of SNPs) that can be generated using the proposed methodology. Due to the lower

cost of generating this type of data we propose that this strategy could be widely used to

perform genetic diversity analysis on thousands of samples, facilitating association mapping

studies aimed at identifying the genetic basis of important agronomic traits in cacao such as

disease resistance and bean quality.

Methods

Plant materials

A total of 30 accessions from the germplasm bank from Cocoa Research Centre at the

University of West Indies in Trinidad **(S1 Table)** were included in the analysis. Young leaves

from one individual per accession were sampled in Trinidad and sent to the Cornejo Lab at

Washington State University. Small strips (100 mg) were cut with sterile scissors (washed

with 1% hydrochloric acid and ethanol prior to and in between sample processing), placed in

1.5 ml Eppendorf tubes and stored at -80 ºC for DNA extraction. Five samples (GU 175/P,

ICS 1, IMC 67, M 8[SUR], SCA 6) from the 200 genomes project were genotyped in this

study in order to compare the concordance between SNPs identified using GBS to those

identified using full genome sequencing [8].

DNA extraction and GBS library preparation

Total DNA was isolated from 100 mg of young leaves collected from each accession using

a DNeasy Plant Mini Kit (QIAGEN, Germany) according to manufacturer's instructions. The

final elution volume was adjusted to 60 μl with 1X TE solution buffer. Integrity of the total

DNA was checked on a 0.7% agarose gel. DNA concentration was measured using a Qubit

Fluorometer (Life Technologies, Thermo Fisher Scientific Inc., Waltham, MA, USA) with a

dsDNA HS Assay Kit (Life Technologies). All samples were normalized to the same volume

(50 μl) containing 500 ng of DNA. For each sample, genomic DNA was digested with 2.4

units of *BsaXI* (New England Biolabs, NEB, R0609S) at 37 ºC overnight in a 100 μl reaction

buffered with Cut Smart NEBuffer (NEB, B7204S). After digestion, we performed a 0.7X

size selection using AMPure XP beads to remove any large undigested fragments. The DNA

was then prepared for sequencing using an NEBNext Ultra DNA Library Prep Kit for

Illumina (NEB E7370S; version 1.2) with the following modifications. Adapter ligation was

followed with a bead clean up without size selection. Library enrichment was accomplished

using 10 cycles of PCR amplification. Finally, we evaluated the distribution of fragment

lengths for each library using an Agilent 2100 BioAnalyzer. Samples were sequenced on an

Illumina HiSeq 2500 (Illumina Inc. San Diego, CA), with paired-end 100 bp reads.

SNP discovery and data processing

Raw sequence reads were demultiplexed and converted to Fastq files by the Spokane

Sequencing Core at Washington State University using Casava v1.8.2 (Illumina Inc.). SNP

quality control and trimming were conducted using FastQC [32] and Trim Galore v0.5.0 [33],

respectively. Reads were trimmed to remove 3' bases with a quality score less than 25. Reads

in which complete or partial adapter sequences were identified were trimmed accordingly.

Reads less than 60 bp in length after trimming were removed. The trimmed sequences were

aligned to the Matina reference genome using BWA v0.17.0 [34] with the default parameters

(except for mismatch penalty -B 6). We performed quality assessment of the BAM files,

validation read groups, removal of PCR duplicates, and correction of mapping quality

assignment with Picard v2.18.9 [35]. The genome analysis toolkit (GATK) v3.8.1 [36] was

used to perform local realignment and base quality recalibration. We performed SNP calling

using the unified genotyper module in GATK to facilitate comparison with previous work.

Hard filters were applied to the data (quality by depth higher than 2, strand bias FS 60, Root

mean square of mapping quality 40.0) and only SNPs that passed the filters were retained.

For population structure analysis we used VCFtools v0.1.13 [37] to performed additional

variant filtering in which only bi-allelic variants with a minimum minor allele frequency

(MAF) greater than 0.05 were retained. Coverage statistics were obtained by analyzing these

loci with Plink-Seq v0.10 [38]. To analyze the types of mutations we used the program

SnpEff with the genome built in database [39]. The R scripts are provided in the GitHub of

the project.

In silico digest

In silico digestion of the cacao chromosomes were performed with the *T. cacao* cv. Matina

reference genome [14]. To identify all restriction cut site positions for *BsaXI*, the program

RESTRICT from the Emboss package v6.5.7.0 was used [40]. Digested fragments were

ordered per chromosome and then were summed using two size ranges (200-700 bp and 200-

1000 bp). The desired range for the libraries to be sequenced on an Illumina HiSeq is between

200-700 bp, however fragments of up to 1000 bp can be sequenced [41]. We report the

distribution of the fragments as a histogram and the location of the *in silico* digested

fragments along the genome.

To estimate the number of fragments produced from the *BsaXI* GBS experiment and to

compare them with the *in silico* predictions, the alignment files were analyzed using

INTERSECT included in BEDtools v2.27.0 [42]. Using the output files from this command

we estimated the number of *BsaXI* fragments for all chromosomes and compared the results

with those from the *in silico* digestion.

Genetic diversity

Observed heterozygosity (Ho), expected heterozygosity (He), and polymorphism

information content (PIC) were calculated with Power Marker v1.25 [43]. Tajima's D was

computed with VCFtools v0.1.13. We also estimated allele frequencies using VCFtools for

each of the sites that met the following criteria: i) bi-allelic and ii) no missing data. For each

site we used folded site frequency spectrum (SFS) for the cacao GBS dataset, which describes

the genetic variation as a distribution of allele frequencies across the genome. The site

frequency spectrum was generated in R.

Admixture and population genetic structure analysis

The population analysis was carried out on the filtered set of SNPs. We analyzed the ancestry

and admixture patterns for the 30 unrelated accessions using cacao samples recently

sequenced [8]. The intersected dataset of MAF filtered SNPs between our cacao GBS dataset

(test set) and the 79 individuals from the whole sequence dataset (reference set) consisted of

5,225 SNPs. We ran ADMIXTURE v1.3 [44] in supervised mode assuming ten populations,

following Motamayor et al. 2008 [45] and Cornejo et al [8]. Principal component analysis

(PCA) was performed with the algorithm implemented in EIGENSOFT [46]. All figures

were generated using R version v3.5.1 [47]. The precision of the ancestry analysis was

evaluated by obtaining bias and standard error for each ancestry out of a bootstrap with 2000

pseudo-replicates in ADMIXTURE v1.3 [44].

Results

In silico restriction enzyme digestion

To determine i) what the expected number of fragments to be sequenced was; and ii) to infer

how many of the sequenced fragments matched locations in the genome that were expected

from the digestion, we evaluated the distribution of fragments throughout the genome (**S1**

Fig). Those sequenced regions that did not match the locations identified from the *in silico*

digest were considered off target. The *in silico* digestion was done considering two sets: 1)

lengths ranging from 200 to 700 bp generated a total of 5,479,477 bp **(Fig 1A)** and 2) lengths

ranging from 200 to 1000 bp generated a total of 10,217,990 bp **(Fig 1B)**. These represent 1

and 3% of the genome, which suggests that *BsaXI* is an appropriate cutter. We used the larger

range to evaluate overlap of sequenced target and predicted target.

Fig 1. *In silico* analysis of restriction enzyme sites in the *T. cacao* genome. (A) Number

of fragments computed in the size range between 200 and 700 bp. **(B)** Number of fragments

computed in the size range between 200 and 1000 bp.

The comparison between the predicted fragment-size distribution with the fragments

produced through our assay **(Fig 2)** suggests high variability in the accuracy of sequenced

targets across samples. Despite the high variability, 73% of the samples had an overlap of

over 57% with the *in silico* digest, and 53% of samples had an overlap over 76% with the

expectation

Fig 2. Assessment of sequencing on target per sample. The orange bar corresponds to the

total number of fragments that are predicted to be digested *in silico* within the length

distribution of 200 to 700 bp. Blue bars correspond to the number of sequenced fragments

that meet our callability criteria overlapping with those predicted *in silico*.

SNP Discovery

We obtained a total of 366 million paired-end 100 bp reads. The number of raw sequence

reads per individual ranged from 0.4 million to 52.2 million. After removal of low-quality

sequences and adapter trimming, 341 million reads remained. Of these reads, 322 million

aligned to the Matina reference genome, and were used for SNP calling. The mean read depth

ranged from 8.9 to 160.1. The observed differences in read depth across samples is most

likely due to differences in quality and/or variations in barcode efficiencies. Individual data

information is shown in **Table 1**.

224   **Table 1.** Summary statistics of sequenced data per individual.

Accession Number	Reads raw-data (Millions)	Reads after trimming (Millions)	Mapping reads (and percent)	Depth
CL 10/11	0.53	0.50	0.49 (97.6%)	15.2

CL 10/3	4.8	4.7	4.5 (96.8%)	67.1
CLM 100	14.9	14.6	14.2 (97.4%)	131.2
GS 6	14.5	14.1	13.8 (97.8%)	122.9
GS 61	10.2	9.9	9.5 (96.4%)	104.2
GU 175/P_1	5.3	5.1	5.0 (96.7%)	73.6
GU 175/P_2	1.1	1.04	1.01 (96.6%)	9.0
ICS 1_1	3.2	3.1	3.0 (96.9%)	52.6
ICS 1_2	4.9	4.7	4.5 (95.6%)	86.7
ICS 28	1.9	1.88	1.82 (96.8%)	26.6
ICS 87	3.1	1.67	1.62 (96.9%)	21.6
IMC 6	7.9	5.3	5.1 (95.7%)	71.6
IMC 67_1	1.91	1.90	1.83 (96.4%)	26.6
IMC 67_2	0.4	0.36	0.35 (97.6%)	8.9
LCT EEN23	10.4	8.3	8.2 (98.2%)	126.6
LCT EEN246	8.2	7.8	7.6 (97.1%)	100.4
LCT EEN31	7.8	6.2	6.0 (96.3%)	74.8
LP 3/29[POU]	7.4	5.2	5.0 (96 %)	62.3
LZ 33	8,0	6.2	6.0 (97.4%)	66.3
M8 [SUR]_1	4.6	3.4	3.3 (96.6%)	52.1
M8 [SUR]_2	15.2	14.8	14.4 (97.4%)	125.9
NA 189	7.2	5.2	5.1 (96.9%)	72.4
NA 26	12,0	11.7	11.2 (95.9%)	128.5
NA 68	6.6	6.4	6.2 (96.7%)	86.6
RIM 113[MEX]	10.8	10.5	10.1 (96.1%)	39.9
SCA 6_1	2.6	2.5	2.4 (96.4%)	39.9
SCA 6_2	8.4	8.1	7.9 (96.6%)	101.2
SLC 18	13.9	13.1	12.7 (97.1%)	123.9
SLC 8	20.6	19.5	18.8 (96.6%)	141.9
UF 38	52.2	50.9	49.05 (96.2%)	160.1

A total of 12,357 SNPs were identified in the collection. The SNPs were classified into

transitions (Ti) and transversions (Tv) based on nucleotide substitution. The number of

A/G transitions was very similar to the number of C/T transitions. The numbers of A/C

and A/T transversions were relatively higher than the C/G and similar to G/T

transversions. Transitions are the most common type of nucleotide substitutions [48]. In

this dataset 62.7% of the base changes were transitions and 37.3% were transversions

(Table 2).

Table 2. Summary of polymorphisms identified in cacao dataset.

Total number of polymorphisms		
SNPs	12,357	100%
Transition		
A/G	3,856	31.2%
C/T	3,894	31.5%
Transversion		
A/C	1,243	10.1%
A/T	1,403	11.3%
C/G	758	6.2%
G/T	1,203	9.7%
Ti/Tv ratio	1.79	

Genetic diversity analysis

The dataset was reduced to 7,009 SNPs after excluding markers showing: 1) more than 20%

missing data, 2) a MAF $\leq$ 0.05, 3) multi-allelic sites, 4) insertions and deletions. Expected

heterozygosity (He) (mean 0.272), Ho (mean 0.200) and PIC (mean 0.226) values estimated

per position are listed in **S2 Table**. He, Ho and PIC ranged from 0.117 to 0.500, from 0 to

0.437, and from 0.374 to 0.375, respectively. Tajima's D was 0.458, indicating an excess of

intermediate frequency alleles that could result from demographic processes such as

population bottlenecks, population subdivision or migration. The SFS showed a significantly

high proportion of rare alleles (45.3% of identified alleles). These results quantitatively show

the proportion of rare variants that would be missed if genotyping arrays were employed for

detecting and characterizing genetic variation **(Fig 3)**. The SFS shows a variable pattern of

frequencies of alleles at intermediate minor allele counts, which is consistent with

unaccounted for population structure.

Fig 3. The site frequency spectrum of *T. cacao* samples used in the study. The orange

bars correspond to alleles found in one, two or three chromosomes in the sample, considered

to be low frequency in this study. The green bars correspond to alleles with a minor allele

count of 4 or more and are considered common variants for the purpose of this experiment.

Conservative estimates suggest that nearly 50% of the SNPs are rare and would be missed if

an array were used to genotype these accessions.

Admixture and population genetic structure analysis

The filtered set of SNPs was used for admixture and population structure analysis. The 30

accessions were compared with ten reference populations that are related to the cacao groups

previously defined [45]. Fifty percent of analyzed individuals were assigned to unique

ancestry in the Curaray, Nanay, Contamana, Amelonado, Nacional, Guianna and Iquitos

populations. The other half of the individuals presented a variety of mixed ancestries **(Fig 4)**.

The analysis of bias and error supports a high level of precision in the assignment of ancestry

for the majority of the samples **(S2 Fig A)**. For the most part, bias in the assignment was

lower than 5%. However, LCTEEN23 presented a relatively higher bias (10%) in the

assignment to Nacional ancestry **(S2 Fig B)**. Standard error (SE) is relatively low overall

with the majority being below 0.05. Sample LCTEEN23 presented a relatively higher SE in

the assignment of Nacional ancestry **(S2 Fig C)**.

Fig 4. Supervised ADMIXTURE using ten reference populations. Each bar corresponds

to an individual and the height to the proportion of ancestry explained by one of the 10

previously described populations. The first uniformly colored 79 columns correspond to

individuals in the reference set. The remaining 30 columns show the proportions of ancestry

assignment for each one of the newly sequenced cacao samples with our GBS approach.

In addition, pairwise allele (SNP) sharing distances were computed between all samples and

reference populations and PCA was used to illustrate the resulting pairwise distances **(Fig 5)**.

Plotting the two first axes showed that they capture the previously described genetic groups

with distinct clusters reflecting the different origin in cacao today. The first principal

component (13.74% of the variation) showed a clear relationship between the samples and

the Criollo and Amelonado ancestral populations whereas the remaining populations

(Curaray, Nanay, Contamana, Maranon, Nacional, Gianna, Iquitos, and Purus) separated

along the second principal component (11.77% of the variation).

Fig 5. Principal component analysis of reference and cacao GBS samples. PCA plots of

the same cacao collection according to subgroups, as identified by ADMIXTURE software.

Although reference samples are closer together, the newly sequenced individuals are placed

close to other individuals showing similar ancestry. Admixed individuals in the new cacao

GBS set are more variable and overdispersed when compared to the reference set.

Discussion

Marker assisted breeding in cacao currently faces a great challenge in the development of

alternative fast, cheap, and accurate genotyping strategies for a large number of accessions.

The need for inexpensive and accurate genotyping strategies is especially true when

considering that most cacao producers are located in developing countries with limited

resources. Some of the technologies developed for array genotyping, which have been

advanced by us and others, work well for the identification of known variants but are limited

in the ability to discover new and potentially informative variants. Genotyping-by-

sequencing is a high-throughput, low-cost technology, which is useful for variant

identification in diverse species and populations and allows for the discovery of previously

unidentified variants [21,49]. In this work, we have adapted and modified a GBS genotyping

assay for cacao. The most important changes to the library preparation are: 1) the use of the

restriction enzyme *BsaXI* that cut 14 bp away from the recognition motif [30], 2) the use of

the Illumina Y-adapters allowing for PCR incorporation of dual-indexed barcodes, which

facilitates large-scale, inexpensive multiplexing, and 3) the efficient cleaning of non-ligated

adapter excess and PCR duplicates using cleaning beads, which is a widely used technique

already employed in previous work [8].

There was variation across samples in the number of mapped reads that overlapped with the

fragments predicted from the *in silico* digestion. The performance of our genotyping in on-

target regions is similar to other studies [21], while we produced a large number of reads per

sample (0.5 M – 14.3 M reads), only a fraction overlapped with predicted regions for a

number of samples. The lack of overlap for some samples suggests that a large number of

fragments sequenced were off target. For example, GU 175.2 had a low percent of overlap

(5%), whereas UF 38 had 95% overlap with the *in silico* digest (**Fig 2**). The number of

sequenced fragments overlapping with *in silico* predicted regions is likely explained by

variations in DNA quality that result from fragmentation during DNA extraction, prior to the

digestion.

We identified a large number of variants, 12,357 SNPs, that were retained after filtering.

Estimates of genetic variability are comparable with those estimated in previous studies of

cacao [18]. Enhanced SNP discovery, SNP quality, production steps, and optimization of

parameters improved the SNP detection when compared to those obtained from 96 SNPs

from small Fluidigm arrays or the 6K SNP array previously reported [15–17]; or even from

the 15K SNP arrays considering the fact that a large proportion of SNPs cannot be identified

with array systems. According to Romay et al. 2013 [50], one important advantage of GBS

over the SNP array systems, is the MAF distribution. We are interested in the identification

of novel genetic variation that might prove to be useful in the identification of traits of interest

and much of this genetic variation may not be captured with genetic arrays. The allele

frequency distribution of variants identified using GBS followed an expected pattern with an

excess of rare variants **(Fig 3)**. More than 45% of bi-allelic variants were rare and the

distribution of intermediate variants revealed signatures of population structure.

The mean expected heterozygosity value was higher than the observed heterozygosity,

indicating a deficit of heterozygotes. Compared with recent published genetic diversity

studies in cacao [16,17,51], our results present slightly lower values of He, Ho and PIC,

probably due to the high presence of homozygous genotypes of the Criollo and Amelonado

populations and the impact of a well-known effect of allelic dropout in GBS datasets [30,52].

The pattern of population structure was further validated when ancestry and structure were

analyzed together with the reference panel set. To study the ancestry of the newly sequenced

samples 5,225 bi-allelic SNPs were successfully intersected with a reference panel, which

consisted of 79 fully sequenced genomes. The samples used for the reference set were

recently sequenced [8] and represent the 10 different genetic groups from the Upper Amazon,

Lower Amazon, Orinoco, and Guyana [45]. Our ancestry characterization of the newly

sequenced 30 samples was consistent with the known passport data for the accessions **(S1**

Table). More specifically, accessions NA 26 and NA 189 were successfully assigned to the

Nanay group and both SCA 6 accessions were correctly assigned to Contamana, showing

that the procedure allows us to correctly identify their population of provenance. In the case

of admixed individuals, we could correctly assign ancestry to individuals with known

admixture patterns. For example, for ICS or UF individuals which are known to be a mix

between Criollo and Amelonado. These results make us confident that the generated data and

procedure could be used to disentangle more complex patterns of genetic admixture like those

obtained for previously uncharacterized individuals such as LCT EEN 246. The accession

LCT EEN 246 is comprised of Criollo, Marañon, Iquitos, and Contamana ancestry, which

could open new possibilities for studying traits of interest.

Overall, the observed patterns of ancestry are consistent with a known history of the

hybridization process in cacao breeding programs and natural hybridization [8]. Studies with

a limited set of markers have already hinted at the possibility of further substructure within

the ten main populations described for cacao [45,53]. Our results show the usefulness of GBS

to address these questions.

Conclusions

We have shown the increased potential for the identification of ancestry informative and novel variants, which are useful for addressing questions about population demographic history and genomic variation in *Theobroma cacao*. We were able to identify 12,357 SNPs markers from only 30 samples, out of which 7,009 could be used for ancestry determination. There is a need to increase our knowledge about cacao accessions throughout the world to produce high yielding and disease resistance genotypes. Our results have practical implications for the development of strategies for genotyping large numbers of accessions with the aim of performing association mapping studies. In our future work, the GBS technique will be refined to include a two digestion enzymes protocol to genotype larger cacao collections and increase the number of common targets/SNPs identified with the method.

Acknowledgments

We thank Path Umaharan, Lambert Motilal, and Annelle Holder-John for providing the germplasm material and associated information used in this study. The authors thank Joanna Kelley for assistance in revising the final version of the manuscript. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Jaime Osorio was supported through funds assigned from Agrosavia-Colombia to the Cornejo Lab to work on the library preparation of the samples.

Data availability statement

383   Raw sequence reads (data to SRA) is available at NCBI Bioproject ID No. PRJNA486150.

384   The scripts are available at GitHub repository (https://github.com/jaog224/cacao_gbs).

Supporting information

S1 Table. Summary of the accessions of the *Theobroma cacao* collection used in this

study.

S2 Table. Summary statistics of genetic diversity of the 7,009 SNPs.

S1 Fig. Distribution of *BsaXI* enzyme cut sites across the *T. cacao* cv. Matina genome.

S2 Fig. Precision of the ancestry analysis using 200 bootstrap pseudoreplicates. (A)

ADMIXTURE bar plot, **(B)** bias bar plot and **(C)** standard error for each ancestry.

References

1.   Motamayor J, Risterucci A, Lopez P, Ortiz C, Moreno A, Lanaud C. Cocoa

domestication I: the origin of the cocoa cultivated by the Mayas. Heredity (Edinb).

2002;89: 380–386.

2.   Bartley BGD. The genetic diversity of cacao and its utilization [Internet]. Bartley

BGD, editor. Genetic Diversity of Cacao and Its Utilization. Wallingford: CABI;

2005. doi:10.1079/9780851996196.0000

3.   Wood GAR, Lass R. Cocoa. 4th ed. Longman, London; 1985.

4.   Donald PF. Biodiversity Impacts of Some Agricultural Commodity Production

Systems. Conserv Biol. 2004;18: 17–38. doi:10.1111/j.1523-1739.2004.01803.x

5.   Othman A, Ismail A, Abdul Ghani N, Adenan I. Antioxidant capacity and phenolic

content   of   cocoa   beans.   Food   Chem.   2007;100:   1523–1530.

doi:https://doi.org/10.1016/j.foodchem.2005.12.021

6.   Henderson JS, Joyce RA, Hall GR, Hurst WJ, McGovern PE. Chemical and

archaeological evidence for the earliest cacao beverages. Proc Natl Acad Sci. National

Academy of Sciences; 2007;104: 18937–18940. doi:10.1073/pnas.0708815104

7.   Ogata N, Gómez-pompa A, Taube KA. The domestication and distribution of

Theobroma cacao L. in the Neotropics. In: McNeil C, editor. Chocolate in

Mesoamerica. Gainesville, USA: University Press of Florida; 2009. pp. 69–89.

doi:10.5744/florida/9780813029535.003.0003

8.   Cornejo OE, Yee M-C, Dominguez V, Andrews M, Sockell A, Strandberg E, et al.

Genomic insights into the domestication of the chocolate tree, *Theobroma cacao* L.

bioRxiv. Cold Spring Harbor Laboratory; 2017; doi:10.1101/223438

9.   Cuatrecasas J. Cacao and its allies: a taxonomic revision of the genus *Theobroma*.

Contrib US Herb. 1964;35: 379–614.

10.   Hunter JR. The status of cacao (*Theobroma cacao*, sterculiaceae) in the western

hemisphere. Econ Bot. 1990;44: 425–439. doi:10.1007/BF02859775

11.   World Cocoa Foundation. Cocoa market update [Internet]. 2014 [cited 9 Jul 2018].

Available: http://www.worldcocoafoundation.org/wp-content/uploads/Cocoa-

Market-Update-as-of-4-1-2014.pdf.

12.   Dantas LG, Guerra M. Chromatin differentiation between *Theobroma cacao* L. and *T.*

grandiflorum Schum. Genet Mol Biol. scielo; 2010;33: 94–98. Available:

http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1415-

47572010000100017&nrm=iso

13. Argout X, Salse J, Aury J-M, Guiltinan MJ, Droc G, Gouzy J, et al. The genome of

Theobroma cacao. Nat Genet. The Author(s); 2010;43: 101. doi:10.1038/ng.736

14. Motamayor JC, Mockaitis K, Schmutz J, Haiminen N, Livingstone D, Cornejo O, et

al. The genome sequence of the most widely cultivated cacao type and its use to

identify candidate genes regulating pod color. Genome Biol. BioMed Central Ltd;

2013;14: r53. doi:10.1186/gb-2013-14-6-r53

15. Livingstone D, Royaert S, Stack C, Mockaitis K, May G, Farmer A, et al. Making a

chocolate chip: development and evaluation of a 6K SNP array for *Theobroma cacao*.

DNA Res An Int J Rapid Publ Reports Genes Genomes. Oxford University Press;

2015;22: 279–291. doi:10.1093/dnares/dsv009

16. Cosme S., Cuevas HE, Zhang D, Oleksyk TK, Irish BM. Genetic diversity of

naturalized cacao (*Theobroma cacao* L.) in Puerto Rico. Tree Genet Genomes. Tree

Genetics & Genomes; 2016;12: 88. doi:10.1007/s11295-016-1045-4

17. Osorio-Guarín JA, Berdugo-Cely J, Coronado RA, Zapata YP, Quintero C, Gallego-

Sánchez G, et al. Colombia a source of cacao genetic diversity as revealed by the

population structure analysis of germplasm bank of *Theobroma cacao* L. Front Plant

Sci. 2017;8: 1994. doi:10.3389/fpls.2017.01994

18. Livingstone D, Stack C, Mustiga GM, Rodezno DC, Suarez C, Amores F, et al. A

larger chocolate chip—development of a 15k *Theobroma cacao* L. SNP array to create

high-density   linkage   maps.   Front   Plant   Sci.   2017;8:   2008.

doi:10.3389/fpls.2017.02008

19.   Ganal MW, Polley A, Graner E-M, Plieske J, Wieseke R, Luerssen H, et al. Large

453   SNP arrays for genotyping in crop plants. J Biosci. 2012;37: 821–828.

454   doi:10.1007/s12038-012-9225-3

455   20.   Loman NJ, Misra R V, Dallman TJ, Constantinidou C, Gharbia SE, Wain J, et al.

456   Performance comparison of benchtop high-throughput sequencing platforms. Nat

457   Biotechnol. Nature Publishing Group, a division of Macmillan Publishers Limited. All

458   Rights Reserved.; 2012;30: 434. Available: http://dx.doi.org/10.1038/nbt.2198

459   21.   Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, et al. A robust,

460   simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS

461   One. Public Library of Science; 2011;6: 1–10. doi:10.1371/journal.pone.0019379

462   22.   Poland JA, Rife TW. Genotyping-by-Sequencing for plant breeding and genetics.

463   Plant Genome J. 2012;5: 92–102. doi:10.3835/plantgenome2012.05.0005

464   23.   Johnson JL, Wittgenstein H, Mitchell SE, Hyma KE, Temnykh S V, Kharlamova A

465   V, et al. Genotyping-By-Sequencing (GBS) detects genetic structure and confirms

466   behavioral qtl in tame and aggressive foxes (*Vulpes vulpes*). PLoS One. Public Library

467   of Science; 2015;10: 1–22. doi:10.1371/journal.pone.0127013

468   24.   Lu F, Romay MC, Glaubitz JC, Bradbury PJ, Elshire RJ, Wang T, et al. High-

469   resolution genetic mapping of maize pan-genome sequence anchors. Nat Commun.

470   The Author(s); 2015;6: 6914. Available: http://dx.doi.org/10.1038/ncomms7914

471   25.   Osorio-Guarín JA, Enciso-Rodríguez FE, González C, Fernández-Pozo N, Mueller

472   LA, Barrero LS. Association analysis for disease resistance to *Fusarium oxysporum*

473   in cape gooseberry (*Physalis peruviana* L). BMC Genomics. BMC Genomics;

474   2016;17: 248. doi:10.1186/s12864-016-2568-7

26.   Torkamaneh D, Laroche J, Belzile F. Genome-wide snp calling from genotyping by

sequencing (GBS) data: a comparison of seven pipelines and two sequencing

technologies. PLoS One. Public Library of Science; 2016;11: 1–14.

doi:10.1371/journal.pone.0161333

27.   Alipour H, Bihamta MR, Mohammadi V, Peyghambari SA, Bai G, Zhang G.

Genotyping-by-Sequencing (GBS) revealed molecular genetic diversity of iranian

wheat landraces and cultivars. Front Plant Sci. 2017;8: 1293.

doi:10.3389/fpls.2017.01293

28.   Bastien M, Boudhrioua C, Fortin G, Belzile F. Exploring the potential and limitations

of genotyping-by-sequencing for SNP discovery and genotyping in tetraploid potato.

Genome. 2018;61: 449–456. doi:10.1139/gen-2017-0236

29.   Gurgul A, Miksza-Cybulska A, Szmatoła T, Jasielczuk I, Piestrzyńska-Kajtoch A,

Fornal A, et al. Genotyping-by-sequencing performance in selected livestock species.

Genomics. 2018; doi:https://doi.org/10.1016/j.ygeno.2018.02.002

30.   Cooke TF, Yee M-C, Muzzio M, Sockell A, Bell R, Cornejo OE, et al. GBStools: a

statistical method for estimating allelic dropout in reduced representation sequencing

data. PLOS Genet. Public Library of Science; 2016;12: 1–18.

doi:10.1371/journal.pgen.1005631

31.   He J, Zhao X, Laroche A, Lu Z-X, Liu H, Li Z. Genotyping-by-sequencing (GBS), an

ultimate marker-assisted selection (MAS) tool to accelerate plant breeding. Front Plant

Sci. 2014;5: 484. doi:10.3389/fpls.2014.00484

32.   Andrews S. FastQC: a quality control tool for high throughput sequence data.

[Internet]. 2018 [cited 9 Jul 2018]. Available:

http://www.bioinformatics.babraham.ac.uk/projects/fastqc

33.   Krueger F. Trim Galore [Internet]. 2018 [cited 9 Jul 2018]. Available:

http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/.

34. Li H, Durbin R. Fast and accurate short read alignment with burrows–wheeler

transform. Bioinformatics. Oxford University Press; 2009;25: 1754–1760.

doi:10.1093/bioinformatics/btp324

35. Picard toolkit [Internet]. Broad Institute; 2018. Available:

http://broadinstitute.github.io/picard/

36. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The

Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation

DNA sequencing data. Genome Res. Cold Spring Harbor Laboratory Press; 2010;20:

1297–1303. doi:10.1101/gr.107524.110

37. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The

variant call format and VCFtools. Bioinformatics. Oxford University Press; 2011;27:

2156–2158. doi:10.1093/bioinformatics/btr330

38. PLINK/SEQ: A library for the analysis of genetic variation data [Internet]. 2018.

Available: https://atgu.mgh.harvard.edu/plinkseq/

39. Cingolani P, Platts A, Wang LLL, Coon M, Nguyen T, Wang LLL, et al. A program

for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff:

SNPs in the genome of Drosophila melanogaster strain w(1118); iso-2; iso-3. Fly

(Austin). Landes Bioscience; 2012;6: 80–92. doi:10.4161/fly.19695

40. Rice P, Longden I, Bleasby A. EMBOSS: the european molecular biology open

software suite. Trends Genet. 2000;16: 276–277. doi:https://doi.org/10.1016/S0168-

9525(00)02024-2

41. Bronner IF, Quail MA, Turner DJ, Swerdlow H. Improved protocols for illumina

sequencing. Curr Protoc Hum Genet. 2009;0 18: 10.1002/0471142905.hg1802s62.

doi:10.1002/0471142905.hg1802s62

42.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. Oxford University Press; 2010;26: 841–842.

doi:10.1093/bioinformatics/btq033

43.   Liu K, Muse S V. PowerMarker: an integrated analysis environment for genetic

marker analysis. Bioinformatics. 2005;21: 2128–9. doi:10.1093/bioinformatics/bti282

44.   Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in

unrelated individuals. Genome Res. Cold Spring Harbor Laboratory Press; 2009;19:

1655–1664. doi:10.1101/gr.094052.109

45.   Motamayor JC, Lachenaud P, Wallace J, Loor R, Kuhn DN, Brown S, et al.

Geographic and genetic population differentiation of the amazonian chocolate tree

(*Theobroma cacao* L.). PLoS One. Public Library of Science; 2008;3: 1–8.

doi:10.1371/journal.pone.0003311

46.   Patterson N, Price AL, Reich D. Population Structure and Eigenanalysis. PLOS Genet.

Public Library of Science; 2006;2: e190. Available:

https://doi.org/10.1371/journal.pgen.0020190

47.   R development core team. R: a language and environment for statistical computing.

[Internet]. Vienna: R Foundation for Statistical Computing; 2008. Available:

http://www.r-project.org

48.   Duchêne S, Ho SYW, Holmes EC. Declining transition/transversion ratios through

time reveal limitations to the accuracy of nucleotide substitution models. BMC Evol

Biol. London: BioMed Central; 2015;15: 36. doi:10.1186/s12862-015-0312-6

49.   Chung YS, Choi SC, Jun TH, Kim C. Genotyping-by-sequencing: a promising tool

for plant genetics research and breeding. Hortic Environ Biotechnol. 2017;58: 425–

431. doi:10.1007/s13580-017-0297-8

50.   Romay MC, Millard MJ, Glaubitz JC, Peiffer J a, Swarts KL, Casstevens TM, et al.

Comprehensive genotyping of the USA national maize inbred seed bank. Genome

Biol. BioMed Central Ltd; 2013;14: R55. doi:10.1186/gb-2013-14-6-r55

51.   Ji K, Zhang D, Motilal LA, Boccara M, Lachenaud P, Meinhardt LW. Genetic

diversity and parentage in farmer varieties of cacao (*Theobroma cacao* L.) from

Honduras and Nicaragua as revealed by single nucleotide polymorphism (SNP)

markers. Genet Resour Crop Evol. 2013;60: 441–453. doi:10.1007/s10722-012-9847-

1

52.   Gautier M, Gharbi K, Cezard T, Foucaud J, Kerdelhué C, Pudlo P, et al. The effect of

RAD allele dropout on the estimation of genetic variation within and between

populations. Mol Ecol. Wiley/Blackwell (10.1111); 2012;22: 3165–3178.

doi:10.1111/mec.12089

53.   Zhang D, Martínez WJ, Johnson ES, Somarriba E, Phillips-Mora W, Astorga C, et al.

Genetic diversity and spatial structure in a new distinct *Theobroma cacao* L.

population in Bolivia. Genet Resour Crop Evol. 2012;59: 239–252.

doi:10.1007/s10722-011-9680-y