A Chromosome-level Sequence Assembly Reveals the Structure of the *Arabidopsis*

***thaliana* Nd-1 Genome and its Gene Set**

Boas Pucker[1], Daniela Holtgräwe[1], Kai Bernd Stadermann[1], Katharina Frey[1], Bruno

Huettel[2], Richard Reinhardt[2] and Bernd Weisshaar[1*]

[1] Bielefeld University, Faculty of Biology & Center for Biotechnology, Bielefeld, Germany

[2] Max Planck Genome Centre Cologne, Max Planck Institute for Plant Breeding Research,

Cologne, Germany

[*] Corresponding author (BW)

Email addresses:
BP: bpucker@cebitec.uni-bielefeld.de
DH: dholtgra@cebitec.uni-bielefeld.de
KS: kstaderm@cebitec.uni-bielefeld.de
KF: katharina.frey@uni-bielefeld.de.
BH: huettel@mpipz.mpg.de
RR: reinhardt@mpipz.mpg.de
BW: bernd.weisshaar@uni-bielefeld.de
20
21 ORCIDs
22   BP: https://orcid.org/0000-0002-3321-7471
23   DH: https://orcid.org/0000-0002-1062-4576
24   KS: https://orcid.org/0000-0002-8036-1492
25   BW: https://orcid.org/0000-0002-7635-3473
26
27

Abstract

Background

In addition to the BAC-based reference sequence of the accession Columbia-0 from the year 2000, several short read assemblies of THE plant model organism *Arabidopsis thaliana* were published during the last years. Also, a SMRT-based assembly of Landsberg *erecta* has been generated that identified translocation and inversion polymorphisms between two genotypes of the species.

Results

Here we provide a chromosome-arm level assembly of the *A. thaliana* accession Niederzenz-1 (AthNd-1_v2c) based on SMRT sequencing data. The best assembly comprises 69 nucleome sequences and displays a contig length of up to 16 Mbp. Compared to an earlier Illumina short read-based NGS assembly (AthNd-1_v1), a 75 fold increase in contiguity was observed for AthNd-1_v2c. To assign contig locations independent from the Col-0 gold standard reference sequence, we used genetic anchoring to generate a *de novo* assembly. In addition, we assembled the chondrome and plastome sequences.

Conclusions

Detailed analyses of AthNd-1_v2c allowed reliable identification of large genomic rearrangements between *A. thaliana* accessions contributing to differences in the gene sets that distinguish the genotypes. One of the differences detected identified a gene that is lacking from the Col-0 gold standard sequence. This *de novo* assembly extends the known proportion of the *A. thaliana* pan-genome.

Background

Introduction

Arabidopsis thaliana became the most important model for plant biology within decades due to properties valuable for basic research like short generation time, small footprint and a small genome [1]. Shortcomings of the BAC-by-BAC assembled 120 Mbp long Col-0 gold standard sequence [2] are some missing sequences and gaps in almost inaccessible regions like repeats in the centromeres [3, 4], at the telomeres and throughout NORs as well as few mis-assemblies [5, Kawakatsu, 2016 #4823]. Information about genomic differences between *A. thaliana* accessions were mostly derived from short read data [6, 7, The 1001 Genomes Consortium, 2016 #4933]. Only selected accessions were sequenced deep enough and with sufficient read length to reach almost reference-size assemblies [6, 8-13]. While the identification of SNPs can be based on short read mappings, the identification of structural variants had an upper limit of 40 bp for most of the investigated accessions [14]. Larger insertions and deletions, which will often result in presence/absence variations of entire genes, are often missed in short read data sets but are easily recovered by long read sequencing [12, 13, 15]. *De novo* assemblies based on long sequencing reads are currently emphasized to resolve structural variants without an upper limit and to facilitate *A. thaliana* pan-genomics. Even a fully complete Col-0 genome sequence would not reveal the entire diversity of this species, as this accession is assumed to have a relatively small genome compared to other *A. thaliana* accessions.

The strong increase in the length of sequencing reads that was technically realized during the last years is enabling new assembly approaches [16, 17]. Despite the high error rate of 'Single Molecule, Real Time' (SMRT) sequencing, the long reads significantly improve contiguity of *de novo* assemblies due to an efficient correction of the almost unbiased errors [18-20], provided that sufficient read coverage is available. SMRT sequencing offered by PacBio results routinely in average read lengths above 10 kbp [8, 21, 22]. These long reads were incorporated into high quality hybrid assemblies involving Illumina short read data [12, 22], but increasing sequencing output supports the potential for so called 'PacBio only

assemblies' [8-10, 13, 19]. Oxford Nanopore Technology's (ONT) sequencing provides even

longer reads with recent reports of longest reads over 2Mbp [23]. However, the error rate of

5-15% [24] is still an issue in plant genome assembly although it has been shown that a high

contiguity assembly is possible for *A. thaliana* [13].

Here we provide a SMRT sequencing-based *de novo* genome assembly of Nd-1 comprising

contigs of chromosome-arm size anchored to chromosomes and oriented within

pseudochromosome sequences based on genetic linkage information. The application of

long sequencing reads abolished limitations of short read mapping and short read

assemblies for genome sequence comparison. Based on this genome sequence assembly,

we identified genomic rearrangements between Col-0 and Nd-1 ranging from a few kbp up to

one Mbp. Gene duplications between both accessions as well as lineage specific genes in

Nd-1 and Col-0 were revealed by this high quality sequence. The current assembly version

outperforms the Illumina-based version (AthNd-1_v1) about 75 fold with respect to assembly

contiguity calculated with respect to number of contigs [11] and is in the same quality range

as the recently released L*er* and KBS-Mac-74 genome sequence assembly [12, 13].

Methods

Plant material

Niederzenz-1 (Nd-1) seeds were obtained from the European Arabidopsis Stock Centre

(NASC; stock number N22619). The DNA source was the same as described earlier [11].

DNA extraction

The DNA isolation procedure was based on previously published protocols [10, 25] and

started with 5 g of frozen leaves which were homogenized by grinding. Samples were mixed

in a 1:10 ratio with extraction buffer (300mM Tris pH8.0, 25mM EDTA, 2M NaCl, 2%

polyvinylpyrrolidone (PVP), 2% hexadecyltrimethylammonium bromide (CTAB)) and

incubated at 65°C for 30 minutes with six inversions to mix the samples again. After five

minute spinning at 5,000xg, the supernatant was transferred and mixed with one volume of

chlorophorm/isoamylalcohol (24:1). Again, the upper phase was transferred after repetition of

the centrifugation step for ten minutes. RNA was removed by adding 30µL RNase A

(10mg/ml) and incubation for 30 minutes at 37°C. Addition of chlorophorm/isoamylalcohol,

centrifugation and transfer of supernatant were repeated. One volume of isopropanol and 0.1

volumes of 3M NaOAc (pH 5.2) were added and mixed. DNA was precipitated by incubating

at -80°C for 30 minutes and spinning for 45 minutes at 5,000g and a final ethanol wash step

was performed. Finally, 500µl of 10mM Tris/HCl (pH 8.0) were added and samples were

incubated over night at 4°C for resuspension.

Library preparation and sequencing

Sequencing was performed using PacBio RS II (Menlo Park, CA, USA). Five microgram high

molecular weight DNA without further fragmentation was used to prepare a SMRTbell library

with PacBio SMRTbell Template Prep Kit 1 (Pacific Biosciences, Menlo Park, CA, USA)

according to the manufacturer's recommendations. The resulting library was size-selected

using a BluePippin system (Sage Science, Inc. Beverly, MA, USA) to enrich for molecules

larger than 11 kbp. The recovered library was again damage repaired and then sequenced

on a total of 25 SMRT cells with P6-C4v2 chemistry and by MagBead loading on the PacBio

RSII system (Pacific Biosciences, Menlo Park, CA, USA) with 360 min movie length.

Assembly parameters

A total of 1,972,766 subreads with an N50 read length of 15,244 bp and containing

information about 16,798,450,532 bases were generated. Assuming a genome size of 150

Mbp, the data cover the genome at 112 fold.

Read sequences derived from the plastome [GenBank: AP000423.1] or chondrome

[GenBank: Y08501.2] were extracted from the raw data set by mapping to the respective

sequence of Col-0 as previously described [26]. Canu v1.4 [27] was used for the assembly of

the organell genome sequences. Beside default parameters, a genome size of 0.37 Mbp for

the chondrome and 0.167 Mbp for the plastome were assumed in the assembly process.

Scaffolding of initial contigs was performed with SSPACE-LongRead v1.1 [28] (parameters: -

a 0 -i 70 -k 3 -r 0.3 -o 10). The quality of both assemblies was checked by mapping of NGS

reads from Nd-1 [11] and Col-0 [29]. Manual inspection and polishing with Quiver v.2.3.0

(parameters: minSubReadLength=500, readScore>80, minLength=500, maxHits=10,

maxDivergence=30, minAnchorSize=12, --maxHits=1, --minAccuracy=0.75, --minLength=50,

--algorithmOptions="-useQuality", --seed=1, --minAccuracy=0.75, --minLength=50, --

concordant --algorithmOptions="-useQuality", minConfidence=40, minCoverage=5,

diploidMode=False) [10] let to the final sequences. The start of the Nd-1 plastome and

chondrome sequences was set according to the corresponding Col-0 sequences to ease

comparisons. Finally, small assembly errors were corrected via CLC basic variant detection

(ploidy=2; coverage<100,000; minimalCoverage=5; minimalCount=5;

minimalFrequency=0.2) based on mapped Illumina paired-end reads (SRX1683594, [11])

and PacBio reads. Sequence properties like GC content and GC skew were determined and

visualized by CGView [30].

A total of 166,600 seed reads spanning 4,500,092,354 nt (N50 = 26,295 nt) and covering the

expected 150 Mbp genome sequence were used for the assembly thus leading to a

coverage of 30 fold (see AdditionalFile1 for details). Assemblies were performed with Canu

[27], FALCON [10], Flye [31] and miniasm [32]. These assemblies are named Ath-Nd1_v2

with an assembler specific suffix (c=Canu, f=FALCON, y=Flye, m=miniasm).

All available SMRT sequencing reads were subjected to Canu v.1.7.1 [27] for read

correction, trimming, overlap detection, and *de novo* assembly (see AdditionalFile2 for

parameters). Contigs with a length below 50 kb were discarded to avoid artifacts. The

remaining contigs were checked for contaminations with bacterial sequences and organell

genome sequences as previously described [11]. BWA-MEM v0.7.13 [33] was used with

default parameters and -m to map Nd-1 Illumina reads [11] to this assembly. The coverage

depth was extracted from the resulting BAM file as previously described [34] and used to

support the identification of plastome and chondrome contigs. Pilon v.1.22 [35] was applied

twice with default parameters for polishing of the assembly.

Release version 1.7.5 of the FALCON assembler

https://github.com/PacificBiosciences/FALCON/ [10] was used for a *de novo* assembly (see

AdditionalFile3 for parameters) of the nuclear genome sequence. Resulting contigs were

checked for contaminations with bacterial sequences and organell genome sequences as

previously described [11]. Small fragments with low coverage were removed prior to

polishing and error correction with Quiver [10].

SMRT sequencing reads were corrected via Canu and subjected to miniasm v0.3-r179 [32]

for *de novo* assembly. Illumina reads were mapped with BWA-MEM for two rounds of

polishing via Pilon as described above. The assembly was reduced to nucleome contigs as

described above for the Canu assembly.

Flye v2.3.1 [31] was deployed on the subreads with an estimated genome size of 150 Mbp.

The resulting assembly was polished twice via Pilon and reduced to nucleome contigs as

described above for the Canu assembly.

Construction of pseudochromosomes based on genetic information

All 26 contigs of Ath-Nd1_v2f were sorted and orientated based on genetic linkage

information derived from 63 genetic markers (AdditionalFile4, AdditionalFile5,

AdditionalFile6), which were analyzed in about 1,000 F2 plants, progeny of a reciprocal

cross of Nd-1xCol-0 and Col-0xNd-1. Genetic markers belong to three different types: (1)

fragment length polymorphisms, which can be distinguished by agarose gel electrophoresis,

(2) small nucleotide polymorphisms which can be distinguished by Sanger sequencing and

(3) small nucleotide polymorphisms, which were identified by high resolution melt analysis.

Design of oligonucleotides was performed manually and using Primer3Plus [36]. DNA for

genotyping experiments was extracted from *A. thaliana* leaf tissue using a

cetyltrimethylammonium bromide (CTAB) based method [37]. PCRs were carried out using

GoTaq® G2 DNA Polymerase (Promega) generally based on the suppliers' protocol. The

total reaction volume was reduced to 15 μl and only 0.2u of the polymerase were used per

reaction. Sizes of amplicons generated were checked on agarose gels. If required, samples

were purified for sequencing by ExoSAP-IT (78201.1.ML ThermoFisher Scientific) treatment

as previously described [38]. Sanger sequencing on ABI3730XL was applied to identify

allele-specific SNPs for the genotyping. Manual inspection of gel pictures and

electropherograms lead to genotype calling. High resolution melt analysis was performed on

a CFX96 Touch Real-Time PCR Detection System (BioRad) using the Precision Melt

Supermix according to suppliers instructions (BioRad). All data were combined, processed

by customized Python scripts to calculate recombination frequencies between genetic

markers. Linkage of genetic markers provided information about relationships of assembled

sequences. The north-south orientation of the chromosomes was transferred from the

reference sequence based on RBH support. Contigs were joined to pseudochromosome

sequences for Ath-Nd1_v2f. Subsequently, positioning was transferred to the 69 contigs of

the Canu assembly to create pseudochromosome sequences for AthNd-1_v2c

(AdditionalFile7).

Since the assemblies except Ath-Nd1_v2f contain smaller contigs in the range of 50-100 kb,

anchoring via genetic linkage information was not feasible. Therefore, these short contigs

were placed based on best BLASTn matches to the TAIR9 sequence (i.e. the Col-0 gold

standard reference) and integrated into pseudochromosomes as previously described [11].

We used the TAIR9 sequence, because this is the sequence basis of the structural and

functional annotation provided in TAIR9, TAIR10 and Araport11 [39]. Since the total length of

the TAIR9 sequence exceeds that of the Ath-Nd1_v2f assembly, TAIR9 was selected for

anchoring of small contigs.

All data generated is stored in the PGP Repository [40] and is accessible via DOI

(https://doi.org/10.5447/IPK/2019/4).

Genome structure investigation

Characteristic elements of the Nd-1 genome sequence were annotated by mapping of known

sequences as previously described [11]. Fragments and one complete 45S rDNA unit were

discovered based on gi|16131:848-4222 and gi|16506:88-1891. AF198222.1 was subjected

to a BLASTn for the identification of 5S rDNA sequences. Telomeric repeats were used to

validate the assembly completeness at the pseudochromosome end as well as centromere

positions as previously described [11].

BUSCO analysis

BUSCO v3 [41] was run with default parameters on the Nd-1 pseudochromosomes and on

the TAIR9 reference sequence to produce a gold standard for Arabidopsis. The

'embryophyta_odb9' was used as reference gene set.

Genome sequence alignment

Nd-1 pseudochromosome sequences were aligned to the Col-0 gold standard sequence [2]

via nucmer [42] and NucDiff [43] based on default parameters of NucDiff. Customized Python

scripts were deployed to process the results and to assess the genome-wide distribution of

differences. Spearman correlation coefficient was calculated using the implementation in the

Python module scipy to validate the indication of increased numbers of SV around the

centromeres.

Gene prediction and RBH analysis

AUGUSTUS v.3.3 [44] was applied to all four Nd-1 assemblies with previously optimized

parameters [38]. Afterwards, the identification of RBHs at the protein sequence level

between Nd-1 and Col-0 (Araport11, representative peptide sequences) was carried out with

a custom Python script as previously described [11]. Additionally, gene prediction was run on

the Col-0 gold standard sequence [2] as well as on the L*er* chromosome sequences [12].

Parameters were set as described before to generate two control data sets. Previously

trimmed and filtered ESTs [11, 45] were matched via BLASTn [46] to the predicted mRNAs.

Pairwise global alignments were constructed via MAFFT v.7.299b [47] to validate the

annotation quality. After processing GCA_000001735.1 (Col-0), Ath-Nd-1_v2c, and the most

recent L*er* assembly [12] with RepeatMasker v4 [48] all three were subjected to cactus [49]

for alignment. CAT [50] was run on this alignment with the ENSEMBL v42 annotation of Col-

0.

INFERNAL [51] with default parameters and based on Rfam13 [52] was applied to detect

various non-coding RNA genes. In addition, tRNAscan-SE [53] with –G option was deployed

to identify tRNA and rRNA genes. Overlaps between both methods were analysed.

Transposable element annotation

All annotated transposable element (TE) sequences of Araport11 (derived from TAIR) [39]

were mapped via BLASTn to the Nd-1 assembly AthNd-1_v2c and against the Col-0 gold

standard sequence. The top BLAST score for each element in the mapping against the

TAIR9 reference sequence was identified. All hits against Nd-1 with at least 90% of this top

score were considered for further analysis. Overlapping hits were removed to annotate a final

TE set. All predicted Nd-1 genes which overlapped TEs with more than 80% of their gene

space were flagged as putative protein encoding TE genes. In addition, RepeatModeler

v.1.0.11 [48] was deployed to identify novel TE sequences in the Ath-Nd-1_v2c assembly.

Identification of gene space differences

Genes in insertions in Ath-Nd1_v2c were searched in the Col-0 gold standard sequence and

vice versa. The BLAST results on DNA and peptide level indicated the absence of any truly

novel genes thus indicating that presence/absence variants are the results of gene

duplications. Apparent gene space differences could be caused by regions missing in the

assembly. Therefore, Nd-1 (SRR2919279, SRR3340908, SRR3340909) [11] and Col-0

(SRR1810832, SRR1945757)[14, 29] Illumina sequencing reads were mapped to Ath-

Nd1_v2c via BWA MEM [33] using the -m flag to discard spurious hits. The sequencing read

coverage depth was calculated via bedtools by a customized Python script [34]. Average

coverage per accession was calculated as the median of all coverage values. Per gene

coverage was calculated as the median of all coverage values at positions in the respective

gene and normalized to the average coverage of the respective accession. To correct for

accession specific mapability differences caused e.g. by sequence divergence from Nd-1,

the resulting values were additionally normalized to the median of all per gene ratios. Genes

were considered as duplicated in Col-0 if their relative coverage in Nd-1 was below 50 % of

the Col-0 value. Genes with Nd-1 values above 150 % of the Col-0 value were considered

duplicated in Nd-1. These cutoff values were validated based on experimentally validated

gene differences. Corresponding AGIs were identified via BLASTp to transfer the functional

annotation of Araport11 if possible. Following this initial identification, putative TE genes

were removed based on the annotation or the overlap with annotated TE sequences

(AdditionalFile8), respectively.

Sequencing data of 1,135 *A. thaliana* accessions [14] were retrieved from the Sequence

Read Archive, mapped to Ath-Nd1_v2c, and processed as described above. Accessions

displaying an average coverage below 10x were excluded from the following identification of

presence/absence variations. GeneSet_Nd-1_v2.0 genes were classified as dispensable if

their relative coverage was below 0.1 in more than 100 accessions. Remaining genes were

classified as core or TE genes depending on their overlap with TE features (see TE

annotation).

Validation of rearrangements and duplications

LongAmpTaq (NEB) was used for the generation of large genomic amplicons up to 18 kbp

based on the suppliers' protocol. Sanger sequencing was applied for additional confirmation

of generated amplicons. The amplification of small fragments and the following procedures

were carried out with standard polymerases as previously described [11].

Investigation of collapsed region

The region around At4g22214 was amplified in five overlapping parts using the Q5 High

Fidelity polymerase (NEB) with genomic DNA from Col-0. Amplicons were checked on

agarose gels and finally cloned into pCR2.1 (Invitrogen) or pMiniT 2.0 (NEB), respectively,

based on the suppliers' recommendations. Cloned amplicons were sequenced on an

ABI3730XL by primer walking. Sequencing reads were assembled using CLC

GenomicsWorkbench (v. 9.5 CLC bio). In addition, 2x250 nt paired-end Illumina reads of

Col-0 [29] were mapped to correct small variants in the assembled contigs and to close a

small gap between cloned amplicons.

Analysis of gaps in the Col-0 reference sequence

Flanking sequences of gaps in the Col-0 gold standard sequence were submitted to a

BLASTn against the Ath-Nd-1_v2c genome sequence. Nd-1 sequences enclosed by hits of

pairs of 30 kbp long flanking sequences from Col-0 were extracted. Homotetramer

frequencies were calculated for all sequences and compared against the frequencies in

randomly picked sequences. A Mann-Whitney U test was applied to analyze the difference

between both groups.

Results

Nd-1 genome

Assemblies generated via Canu (Ath-Nd-1_v2c), FALCON (Ath-Nd-1_v2f), Flye (Ath-Nd-

1_v2y) and miniasm (Ath-Nd-1_v2m) were compared based on numerous assembly

statistics (Table 1). AthNd-1_v2f exceeds the previously reported assembly version AthNd-

1_v1 by 2.5 Mbp, while reducing the number of contigs by a factor of about 200 to 26. AthNd-

1_v2c is adding 6.9 Mbp to the previous assembly version (AthNd-1_v1), but at the expense

of a higher number of contigs than AthNd-1_v2f. We selected AthNd-1_v2c that contains 69

contigs as the representative assembly which should be used for further comparison.

Pseudochromosomes of AthNd-1_v2f were constructed truly *de novo* from 3-7 contigs based

on genetic linkage information. All 26 contigs were anchored based on 63 genetic markers,

but precise positioning around the centromeres of chromosome 4 and 5 was ambigious.

Pseudochromosomes reach similar lengths as the corresponding chromosome sequences in

the Col-0 gold standard sequence. Pseudochromosomes for AthNd-1_v2c were generated

by transferring orientation and position of large contigs from AthNd-1_v2f. The Nd-1 genome

sequence AthNd-1_v2c contains complete 45S rDNA units on pseudochromosome 3 as well

as several fragments of additional 45S rDNA units on all other pseudochromosomes (Fig. 1).

Centromeric and telomeric repeat sequences as well as 5S rDNA sequences were detected

at centromere positions. Completeness of most assembled sequences was confirmed by the

occurrence of telomeric repeat sequences (Fig. 1). The high assembly quality and

completeness of AthNd-1_v2c is supported by the detection of 98.2% of all embryophyta

BUSCO genes – even one more than detected in the Col-0 gold standard sequence

(AdditionalFile9).

The plastome and chondrome sequences comprise 154,443 bp and 368,216 bp, respectively

(https://doi.org/10.5447/IPK/2019/4). A total of 148 small variants were identified from a

global alignment between the Nd-1 and Col-0 plastome sequences. General sequence

properties like GC content and GC skew (AdditionalFile10, AdditionalFile11) are almost

identical to the plastome and chondrome of Col-0. Nevertheless, there are some

rearrangements between the chondrome sequences of Nd-1 and Col-0.

Genome structure differences

Sequence comparison between AthNd-1_v2c and the Col-0 gold standard sequence

revealed a large inversion on chromosome 4 involving about 1 Mbp (Fig. 2). The left break

point is at 1,637,889 bp and the right break point at 2,708,850 bp on chromosome 4

(NdChr4). The inverted sequence is 120 kbp shorter than the corresponding Col-0 sequence.

PCR amplification of both inversion borders (AdditionalFile12) and Sanger sequencing of the

generated amplicons was used to validate this rearrangement.

The recombination frequency in this region was analyzed using the marker pair M84/M74.

Only a single recombination was observed between these markers while investigating 60

plants. Moreover, only 8 recombination events in 108 plants were observed between another

pair of markers, spanning a larger region across the inversion (AdditonalFile5). In contrast,

the average recombination frequency per Mbp at the corresponding position on other

chromosomes was between 12%, observed for M31/M32, and 18%, observed for M13/M14.

Statistical analysis revealed a significant difference in the recombination frequencies

between the corresponding positions on different chromosomes ($p < 0.001$, prop.test() in R)

supporting a reduced recombination rate across the inversion on chromosome 4.

Comparison of a region on Chr2, which is probably of mitochondrial origin (mtDNA), in the

Col-0 gold standard sequence with the Nd-1 genome sequence revealed a 300 kbp highly

divergent region (Fig. 3). Sequences between position 3.20 Mbp and 3.29 Mbp on NdChr2 of

AthNd-1_v2c display low similarity to the Col-0 gold standard sequence, while there is almost

no similarity between 3.29 Mbp and 3.48 Mbp. However, the length of both regions is roughly

the same. Comparison against the L*er* genome sequence assembly revealed the absence of

the entire region between 3.29 Mbp and 3.48 Mbp on chromosome 2. The Nd-1 sequence

from this region lacks continuous similarity to any other region in the Col-0 or Nd-1 genome

sequence. However, the 28 genes encoded in this region in Nd-1 show weak similarity to

other Arabidopsis genes. Comparison of gene space sequences from this region against the

entire Nd-1 assembly revealed some similarity on chromosome 3, 4, and 5

(AdditionalFile13).

An inversion on chromosome 3 of 170 kbp which was described between Col-0 and L*er* [12]

is not present in Nd-1. The sequence similarity between Col-0 and Nd-1 is high in this region.

In total, 2206 structural variants larger than 1 kbp were identified between Col-0 and Nd-1.

The genome-wide distribution of these variants indicated a clustering around the

centromeres (AdditionalFile14). A Spearman correlation coefficient of -0.79 ($p = 1.1*10^{-27}$) was

calculated for the correlation of the number of SVs in a given interval and the distance of this interval to the centromere (AdditionalFile15). Therefore, these large structural variants are significantly more frequent in the centromeric and pericentromic regions. A total, of 148 new regions larger than 1 kbp were identified in Ath-Nd-1_v2c. These regions are also more frequent in proximity of the centromere ($r=-0.43$, $p=3.7*10^{-7}$).

Hint-based gene prediction

Hint-based gene prediction using AUGUSTUS with the *A. thaliana* species parameter set on the Nd-1 pseudochromosomes resulted in 30,126 nuclear protein coding genes (GeneSet_Nd-1_v2.0) with an average predicted transcript length of 1,798 bp, an average predicted CDS length of 1,391 bp and an average exon number per predicted transcript of 5.98. The number of predicted genes is reduced compared to the GeneSet_Nd-1_v1.1 [38] by 708 genes. In total, 28,042 (93%) representative peptide sequences were matched to Araport11 sequences and functionally annotated based on Araport11 information (AdditionalFile16). As controls we run the gene prediction with same parameters on Col-0 and L*er* pseudochromosome sequences resulting in 30,352 genes and 29,302 genes, respectively. There were only minor differences concerning the average transcript and CDS length as well as the number of exons per gene.

Based on 35,636 TEs detected in Nd-1 (AdditionalFile8) 2,879 predicted Nd-1 genes were flagged as putative TE genes (AdditionalFile17, AdditionalFile18). This number matches well with the difference between the predicted genes in Nd-1 and the annotated protein coding genes in Araport11, which is supposed to be free of TE genes. The predicted mRNAs were supported by ESTs, which matched almost perfectly with an average similarity of 98.7% (AdditionalFile19). Additionally, the assembly was screened for TEs resulting in 613 consensus sequences of TE families (https://doi.org/10.5447/IPK/2019/4).

The comparative gene prediction with CAT resulted in 26,717 genes in Nd-1 and 26,681 genes in L*er*, respectively. Average CDS lengths were 1,292 bp and 1,291 bp, respectively. Since TE associated genes were not identified in this gene prediction process, the number of

gene models predicted by CAT should be smaller than the number predicted by

AUGUSTUS. The difference of 3,409 for Nd-1 is slightly exceeding the number of 2,879 TE

associated genes that were predicted and flagged in the GeneSet_Nd-1_v2.0.

Besides protein encoding genes, 557 tRNA genes and 963 rRNA genes were predicted by

INFERNAL. Comparison of these predicted tRNA genes to the result of tRNAscan-SE

revealed an overlap of 552 (96.5%).

Detection of gene space differences between Nd-1 and Col-0

A BLASTp-based comparison of all predicted Nd-1 peptide sequences and Col-0 Araport11

representative peptide sequences in both directions revealed 24,453 reciprocal best hits

(RBHs) (AdditionalFile20). In total, 89.1% of all 27,445 nuclear Col-0 genes are represented

in this RBH set. Analysis of the colinearity of the genomic location of all 24,453 RBHs (see

AdditionalFile21 for a list) between Nd-1 and Col-0 showed overall synteny of both genomes

as well as an inversion on chromosome 4 (AdditionalFile22). While most RBHs are properly

flanked by their syntenic homologs and thus lead to a diagonal positioning of points in the

scatter plot, there are 345 outliers (see AdditionalFile23 for a list). In general, outliers occur

frequently in regions around the centromeres. Positional analysis revealed an involvement of

many outliers in the large inversion on chromosome 4. An NGS read mapping at the

positions of randomly selected outliers was manually inspected and indicated

rearrangements between Nd-1 and Col-0. Structural variants, which affect at least three

different genes in a row of RBH pairs, were identified from the RBH analysis. Examples

beside the previously mentioned 1.2 Mbp inversion on chromosome 4 (*At4g03820-*

At4g05497) are a translocation on chromosome 3 (*At3g60975-At3g61035*) as well as an

inversion on chromosome 3 around *At3g30845*.

As a control we identified 25,556 (91.1%) RBHs between our gene prediction on Col-0 and

the manually curated reference annotation Araport11. In addition, 24,329 (88.6 %) RBHs

were identified between our gene prediction on the L*er* assembly and the Col-0 annotation

Araport11.

In total, 947 protein encoding genes in Nd-1 (AdditionalFile24) were detected to be copies of

only 421 genes annotated in Araport11. *SEC10* (*At5g12370*) [5] was previously described as

an example for a tandem gene duplication collapsed in the Col-0 gold standard sequence.

However, this region was already properly represented in AthNd-1_v1 [11]. Gene

duplications of *At2g06555* (unknown protein), *At3g05530* (*RPT5A*) and *At4g11510*

(*RALFL28*) in Nd-1 were confirmed by PCR amplification and Sanger sequencing of the

sequences enclosed by both copies as well as through amplification of the entire event locus.

On the other hand, there are 383 predicted genes in Nd-1 (AdditionalFile25) which appeared

at least duplicated in Col-0.

Pan-genomic analyses

Presence/absence variations of genes were inspected across a panel of 964 additional

accessions (https://doi.org/10.5447/IPK/2019/4). In total, 25,809 genes were present in

almost all accessions, 1,438 genes were considered dispensable, and the remaining 2,879

genes were flagged as TE or at least TE-associated genes (AdditionalFile26). Dispensable

genes and TE genes are frequently located in proximity of the centromeres while core genes

are less frequent in these regions (AdditionalFile27).

Hidden locus in Col-0

At4g22214 was identified as a gene duplicated in Nd-1 in our analysis. During experimental

validation, we did not detect the expected difference between Col-0 and Nd-1 DNA

concerning the locus around *At4g22214*. However, the PCR results from Col-0 matched the

expectation based on the Nd-1 genome sequence thus suggesting a collapsed gene

sequence in the Col-0 gold standard sequence. This hypothesis was supported by PCR

results with outwards facing primers (Fig. 4). Cloning of the At4g22214 region of Col-0 in five

overlapping fragments was done to enable Sanger sequencing. The combination of Sanger

and paired-end Illumina sequencing reads revealed a tandem duplication with modification of

the original gene (Fig. 4). The copies were designated At4g22214a and At4g22214b based

on their position in the genome (GenBank: MG720229). While At4g22214b almost perfectly

matches the Araport11 annotation of At4g22214, a significant part of the CDS of At4g22214a

is missing. Therefore, the gene product of this copy is probably functionless. At4g22214 is

annotated as defensin-like family protein [54]. Since the family of defensin proteins contains

about 300 members in *A. thaliana* [54], the functional implications of this duplication are

probably low.

Gaps in the Col-0 reference sequence

Despite its very high quality, the Col-0 gold standard sequence contains 92 N stretches of

various sizes representing regions of unknown sequence like the NOR clusters or

centromeres. Ath-Nd1_v2c enabled the investigation of some of these sequences based on

homology assumptions. A total of 13 Col-0 gold standard sequence gaps were spanned with

high confidence by Ath-Nd1_v2c and therefore selected for homopolymer frequency

analysis. The corresponding regions in Nd-1 are significantly enriched with homopolymers in

comparison to randomly picked control sequences (p=0.000048, Mann-Whitney U test)

(AdditionalFile28).

Discussion

Genome structure of the *A. thaliana* accession Nd-1

In order to further investigate large variations in the range of several kbp up to several Mbp

between *A. thaliana* accessions, we performed a *de novo* genome assembly for the Nd-1

accession using long sequencing reads and cutting-edge assembly software. Based on

SMRT sequencing reads the assembly contiguity was improved by over 75 fold considering

the number of contigs in the previously released NGS-based assembly [11]. Assembly

statistics were comparable to other projects using similar data [9, 12, 13, 55, 56]. Despite the

very high contiguity, regions like NORs still pose a major challenge. These regions are not

just randomly clustered repeats, but highly controlled and ordered repetitions of sequences

[57]. Therefore, the identification of accession-specific sequence differences could explain

phenotypic differences. NOR repeat unit sequences in Ath-Nd-1_v2c are located at on the

distal short arm of NdChr2 and NdChr4. This NOR position matches the situation in Col-0

where the NOR2 is located distal on the short arm of Chr2 [58] and NOR4 on Chr4,

respectively. In addition to NORs, the assembly of chromosome ends remains still

challenging, since the absence of some telomeric sequences in a high quality assembly was

observed before [12]. Despite the absence of challenging repeats, regions close to the

telomeres including the genes At3g01060 (BUSCO ID: EOG09360D4T) and At5g01010

(BUSCO ID: EOG09360DFK) were not assembled by FALCON (Ath-Nd1_v2f) although

sequence reads covering these regions were present in the input data. Also, this region is

represented in the Canu assembly (Ath-Nd1_v2c). In our hands, Canu performed best for the

assembly of the Nd-1 genome sequence based on SMRT sequencing reads.

Nuclear genome sequence differences

The increased contiguity of this long read assembly was necessary to discover an

approximately 1 Mbp inversion through sequence comparison as well as RBH analysis. An

earlier Illumina short read based assembly [11] lacked sufficient contiguity in the region of

interest to reveal both breakpoints of this variant between Col-0 and Nd-1. The large

inversion at the north of NdChr4 relative to the Col-0 gold standard sequence turned out as a

modification of the allele originally detected in L*er* [12, 59]. However, the Nd-1 allele is

different from the L*er* allele. This could explain previous observations in several hundred

A. thaliana accessions, which share the left inversion border with L*er*, but show a different

right inversion border [12].

Despite the very much improved contiguity and selection of Canu as the currently best

assembler for our dataset, there are only very small parts of pericentromeric sequences

represented in the Ath-Nd1_v2c assembly. Centromeric regions with an estimated size of 5

Mbp each [11] were not assembled. The absence of these highly repetitive regions is in

agreement with previous findings that the error rate of long reads is still too high to resolve

NORs and centromeres [13]. The pericentromeric and to a very limited extent also

centromeric regions are represented in the assemblies by small contigs of 50-100 kbp.

However, absence of telomeric sequences from some pseudochromsome ends was

observed before even for a very high quality assembly [12, 13]. The detected telomeric

repeats at the centromere positions support previously reported hypothesis about the

evolution of centromers out of telomere sequences [60], and telomeric or centromeric

sequences, respectively, at the end of at least some pseudochromosomes indicated the

completeness of the Ath-Nd1_v2c assembly at these points. However, almost 20 years after

the release of the first chromosome sequences of *A. thaliana*, we are still not able to

assemble complete centromere sequences continuously.

Sequence differences observed on the short arm of chromosome 2 between Col-0 and Nd-1

could be due to the integration of mtDNA into the Chr2 of Col-0 [2]. This region was reported

to be collapsed in the Col-0 reference genome sequence, harboring in real Col-0 DNA about

600 kbp of mtDNA instead of the 270 kbp represented in the reference genome sequence

[61]. However, Ath-Nd1_v2c did not reveal such a duplication of the mtDNA located in

NdChr2. Filtering of plastome and chondrome sequences during the polishing process could

be responsible for this if copies would have been fragmented into multiple contigs. In this

region, we detected in the Nd-1 assembly sequences unrelated to the corresponding Col-0

reference sequence. Since Nd-1 genes of this region show similarity to gene clusters on

other chromosomes, they could be relics of a whole genome duplication as reported before

for several regions of the Col-0 gold standard sequence [62]. This difference on Chr2 at

about 3.2 to 3.5 Mbp is only one example for a large variant region between Col-0 and Nd-1.

Similar though shorter such differences detected around centromeres could be explained by

TEs and pseudogenes which were previously reported as causes for intra-species variants in

these regions [61, 63].

Plastome and chondrome

Size and structure of the Nd-1 plastome is very similar to Col-0 [2] or L*er* [26]. In accordance

with the overall genome similarities, the observed number of small differences between the

plastome sequences of Col-0 and Nd-1 is slightly higher than the value reported before for

the Col-0 comparison to L*er* [26].

The size of the Nd-1 chondrome matches previously reported values for the large chondrome

configuration of other *A. thaliana* accessions [64]. Large structural differences between the

Col-0 chondrome [64] and the Nd-1 chondrome could be due to the previously described

high diversity of this subgenome including the generation of substoichiometric DNA

molecules [65, 66]. In addition, the mtDNA level was reported to differ between cell types or

cells of different ages within the same plant [67, 68]. The almost equal read coverage of the

assembled Nd-1 chondrome could be explained by the young age of the plants at the point of

DNA isolation, as the amount of all chondrome parts should be the same in young leafs [68].

Nd-1 gene space

Many diploid plant genomes contain on average around or even slightly below 30,000 protein

encoding genes [34, 69] with the *A. thaliana* genome harboring 27,445 nuclear protein-

coding genes according to the most recent Araport11 annotation [39]. Since there are only

few other chromosome-level assembly sequences of *A. thaliana* available at the moment, we

do not know the precise variation range of gene numbers between different accessions. The

number of 27,247 predicted non-TE genes in Nd-1 is further supported by the identification of

24,453 RBHs with the Araport11 [39] annotation of the Col-0 gold standard sequence. This

number exceeds the matches between Col-0 and Ler-0 [12]. Our chromosome-level

assembly further enhances the gene prediction quality since at least 89.1% of all Col-0

genes were recovered. This reinforces previous studies that also reported annotation

improvements through improved assembly quality [70]. The number of 35,636 TEs annotated

for Ath-Nd-1_v2c exceeds the number of 33,892 such elements identified in the previous

assembly version Ath-Nd-1_v1 [11] by 1,744. Such an increase in the number of resolved

TEs as well as an improved assembly of TE-rich regions in assemblies based on long reads

was reported before [71].

Due to the high proportion of genes within the *A. thaliana* genome assigned to paralogous

groups with high sequence similarity [72, 73], we speculated that the identification of

orthologous pairs via RBH analysis might be almost saturated. Gene prediction with the

same parameters on the Col-0 gold standard genome sequence prior to RBH analysis

supported this hypothesis. Since there are even some RBHs at non-syntenic positions

between the control Col-0 annotation and the Araport11 annotation, our Nd-1 annotation is

already of very high accuracy. The precise annotation of non-canonical splice sites via hints

as described before [38] contributed to the new GeneSet_Nd-1_v2.0. Gene duplication and

deletion numbers, or accession specific presence/absence variations of genes, in Nd-1 and

Col-0 are in the same range as previously reported values of up to a few hundred [12, 74].

Since we were searching genome-wide for copies of genes without requiring an annotated

feature in each genome sequence, both numbers might include some pseudogenes due to

the frequent occurrence of these elements within plant genomes [75, 76]. Since all

comparisons rely on the constructed sequences we cannot absolutely exclude that a small

number of other genes were detected as amplified due to collapsed sequences similar to

SEC10 (At5g12370) [5]. Removing TE genes based on sequence similarity to annotated

features reduced the proportion of putative pseudogenes. However, it is impossible to

unequivocally distinguish between real genes and pseudogenes in all cases, because even

genes with a premature stop codon or a frameshift mutation could function as a truncated

versions or give rise to regulatory RNAs [73, 76-78]. In addition, the impact of copy number

variations involving protein encoding genes in *A. thaliana* might be higher than previously

assumed thus supporting the existence of multiple gene copies [79]. Gene expression

analysis could support the discrimination of pseudogenes, because low gene expression in

A. thaliana was reported to be associated with pseudogenization [80]. Despite the unclear

status of the gene product, the mere presence of these sequences revealed fascinating

insights into genome evolution and contributed to the pan-genome [81, 82].

Conclusions

We report a high quality long read *de novo* assembly (AthNd-1_v2c) of the *A. thaliana*

accession Nd-1, which improved significantly on the previously released NGS assembly

sequence AthNd-1_v1.0 [11]. Comparison of the GeneSet_Nd-1_v2.0 with the Araport11

nuclear protein coding genes revealed 24,453 RBHs supporting an overall synteny between

both *A. thaliana* accessions except for an approximately 1 Mbp inversion at the north of

chromosome 4. Moreover, large structural variants were identified in the pericentromeric

regions. Comparisons with the Col-0 gold standard sequence also revealed a collapsed

locus around At4g22214 in Col-0. Therefore, this work contributes to the increasing

A. thaliana pan-genome with significantly extended details about genomic rearrangements.

List of abbreviations

NGS   next generation sequencing

NOR   nucleolus organizing region

RBH   reciprocal best hit

SMRT single molecule real time

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The data sets supporting the results of this article are included within the article and its additional files. The Ath-Nd-1_v2 assemblies (Table 1) and derived files are available as external downloads from http://doi.org/10.5447/IPK/2019/4. Sequencing reads were submitted to the SRA (SRP066294). Python scripts developed and applied for this study are available on github: https://github.com/bpucker/Nd1_PacBio (http://doi.org/10.5281/zenodo.2590750).

Competing interest

The authors declare that they have no competing interest.

Funding

We acknowledge the financial support of the German Research Foundation (DFG) and the Open Access Publication Fund of Bielefeld University for the article processing charge. The funding body did not influence the design of the study, the data collection, the analysis, the interpretation of data, or the writing of the manuscript.

Author's contributions

BP, DH and BW conceived and designed research. BP, KS, KF, BH and RR conducted experiments. BP, DH and BW interpreted the data. BP and BW wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Willy Keller for isolating high molecular DNA, Katharina Kemmet for extensive genotyping of plants for the genetic map, Helene Schellenberg, Ann-Christin Polikeit and Prisca Viehöver for Sanger sequencing, and Melanie Kuhlmann as well as Andrea Voigt for taking excellent care of the plants. We are very grateful for support from the Bioinformatics Resource Facility and Stefan Albaum.

Additional Files

AdditionalFile1. Sequencing Statistics.

Statistical information about the generated SMRT sequencing data for the *A. thaliana* Nd-1 genome assembly are listed in this table. The expected genome size is based on several analyses reporting values around 150 Mbp [83, 84].

AdditionalFile2. Canu assembly parameters.

Listing of all parameters that were adjusted for the Canu assembly of the Nd-1 nucleome. While most default parameters were kept, some were specifically adjusted for this plant genome assembly.

AdditionalFile3. FALCON assembly parameters.

All parameters that were adjusted for the FALCON assembly of the Nd-1 nucleome are listed in this table. While most default parameters were kept, some were specifically adjusted for this plant genome assembly.

AdditionalFile4. Molecular markers for genetic linkage analysis.

All markers require the amplification of a genomic region using the listed oligonucleotides under the specified conditions (annealing temperature, elongation time). Depending on the fragment size differences, the resulting PCR products can allow the separation of both alleles by agarose gel electrophoresis (length polymorphism) or might require Sanger sequencing to investigate single SNPs.

AdditionalFile5. Distribution of genetic markers over physical map.

The positions of all genetic markers on the pseudochromosome sequences are illustrated. Assembled sequences were positioned based on the genetic linkage information. Some

genetic marker combinations allowed the investigation of recombination frequencies within

continuous sequences.

AdditionalFile6. Oligonucleotide sequences for genetic linkage analysis.

Sequences, names and recommended annealing temperatures of all oligonucleotides used

in this work are listed in this table. Usage remarks for the oligonucleotides are provided as

well.

AdditionalFile7. Alignment of Ath-Nd1_v2c and Ath-Nd1_v2f.

Assemblies generated by Canu and FALCON, respectively, were compared via BLASTn

search of 10 kb sequence chunks. Color and position of dots in the figure indicate the

position of the best hit on the respective sequence.

AdditionalFile8. TE positions in the Nd-1 genome sequence.

TE genes, TEs, and TE fragments in the Nd-1 genome sequence were identified based on

sequence similarity to annotated TEs from the Col-0 gold standard sequence (Araport11)

[39].

AdditionalFile9. BUSCO analysis of the Col-0 and Nd-1 genome sequences.

BUSCO v2.0 was run on the genomic sequences of Col-0 and Nd-1 using AUGUSTUS 3.3

with default parameters for the gene prediction process.

AdditionalFile10. Nd-1 plastome map.

The GC content (black) and GC skew (green for positive GC skew, purple for negative GC

skew) of the plastome sequence were analyzed by CGView [30]. The sequence and its

properties are very similar to the Col-0 plastome sequence.

AdditionalFile11. Nd-1 chondrome map.

The GC content (black) and GC skew (green for positive GC skew, purple for negative GC

skew) of the chondrome sequence were analyzed by CGView [30]. The sequence and its

properties are very similar to the Col-0 chondrome sequence.

AdditionalFile12. Experimental validation of 1 Mbp inversion on chromosome 4.

The identified inversion between Nd-1 and Col-0 on chromosome 4 is different from the

inversion described before between Col-0 and L*er* [12]. However, the left breakpoint is the

same for both alleles enabling the use of previously published oligonucleotide sequences

[12]. The right breakpoint was identified by manual investigation of sequence alignments.

Both breakpoints were validated via PCR using the oligonucleotides (for sequences see

AdditionalFile6) as illustrated in (a). The results support the expected inversion borders (b).

AdditionalFile13. Genome-wide distribution of genes inserted on chromosome 2 in Nd-

1.

AthNd-1_v2c and the Col-0 gold standard sequence display a highly diverged region at the

north of chromosome 2, which is about 300 kbp long. BLASTn of the complete Nd-1 gene

sequences from this region revealed several regions on other Nd-1 chromosomes with

copies of these genes.

AdditionalFile14. Genome-wide distribution of large structural variants.

The distribution of structural variants (SVs) >10 kbp (red dots) between Col-0 and Nd-1 over

all five pseudochromosome sequences (black lines) is illustrated. Additionally, the assumed

centromere (CEN) positions are indicated (blue dots). Most SVs are clustered in the (peri-

)centromeric region.

AdditionalFile15. Clustering of SVs around centromeres.

The correlation between the number of SVs in a given part of the genome sequence (1 Mbp)

and the distance of this region to the centromere position is illustrated. SVs are clustered

around the centromeres (Spearman correlation coefficient = -0.66, p-value = $1.7*10^{-16}$).

AdditionalFile16. Functional annotation of GeneSet_Nd-1_v2.0.

Functional annotations were transferred from Araport11 to corresponding RBHs in

GeneSet_Nd-1_v2.0. In addition, genes were annotated based on the best BLAST hit

if annotation via RBH was not possible.

AdditionalFile17. TE overlap with GeneSet_Nd-1_v2.0.

The overlap between annotated TEs (AdditionalFile8) and predicted protein coding genes

was analyzed to identify TE genes. This figure illustrates the fraction of a gene that is

covered by a TE. Since TEs might occur within the intron of a gene, only genes with at least

80% TE coverage were flagged as TE genes (AdditionalFile18).

AdditionalFile18. TE genes in GeneSet_Nd-1_v2.0.

These genes were predicted by AUGUSTUS as protein coding genes. Due to their positional

overlap with TEs (AdditionalFile8), they were flagged as TE genes and excluded from further

gene set analysis.

AdditionalFile19. EST mapping.

Percentage of nucleotides in ESTs matching predicted transcripts are displayed.

AdditionalFile20. Gene set overlap between Araport11, GeneSet_Nd-1_v1.1, and

GeneSet_Nd-1_v2.0.

RBHs were identified pairwise between gene sets. The overlap was identified by mapping all

genes onto Araport11 identifiers. Venn diagram construction was performed at

http://bioinformatics.psb.ugent.be/webtools/Venn/.

AdditionalFile21. Reciprocal best hits (RBH) pairs between Col-0 and Nd-1.

Reciprocal best hits between predicted peptide sequences of Nd-1 and the representative

peptide sequences of Col-0 (Araport11).

AdditionalFile22. Reciprocal best hits (RHB) indicates inversion between Nd-1 and

Col-0.

Genes in RBH pairs were sorted based on their position on the five pseudochromosomes of

the two genome sequences to form the x (Col-0) and y (Nd-1) axes of this diagram. Plotting

the positions of each RBH pair leads to a bisecting line of black dots representing genes at

perfectly syntenic positions. Red and green dots indicate RBH gene pair positions deviating

from the syntenic position. Red dots symbolize a unique match to another gene, while green

dots indicate multiple very similar matches. Positions of the centromere (CEN4) on the

chromosomes of both accessions are indicated by purple lines. An inversion involving 131

genes in RBH pairs just north of CEN4 distinguishes Nd-1 and Col-0.

AdditionalFile23. RBH outliers in GeneSet_Nd-1_v2.0.

RBHs between Araport11 and GeneSet_Nd-1_v2.0 were identified based on encoded

representative peptide sequences. All 242 RBHs at positions deviating from the syntenic

diagonal line were collected. The functional annotation of these genes was derived from

Araport11.

AdditionalFile24. Duplicated genes in Nd-1.

The listed 385 Col-0 genes (Araport11 [39]) have at least two copies in Nd-1. Exons of these

genes showed an increased copy number in Ath-Nd-1_v2c compared to the Col-0 gold

standard sequence. The annotation was derived from Araport11.

AdditionalFile25. Duplicated genes in Col-0.

The listed 394 Nd-1 genes have at least two copies in Col-0. Exons of these genes showed

an increased copy number in the Col-0 gold standard sequence compared to Ath-Nd-1_v2c.

AdditionalFile26. Classification of genes as core, dispensable, or TE.

GeneSet_Nd-1_v2.0 genes were classified as core, dispensable, or TE genes based on coverage in a read mapping involving 1,137 Illumina read data sets.

AdditionalFile27. Genome-wide distribution of core, dispensable, and TE genes.

Visualisation of the position of core genes, dispensable genes and TEs along the chromosomes.

AdditionalFile28. Critical regions in the Col-0 gold standard sequence.

The high contiguity of the Ath-Nd-1_v2c assembly enabled the investigation of 13 sequences corresponding to gaps in the Col-0 gold standard sequence. This figure illustrates the homotetranucleotide occurrence in these sequences (red dots) in comparison to some randomly selected reference sequences (green dots). While there is a clear enrichment of homotetranucleotides in the gap-homolog sequences, there was no clear correlation between the length of a gap and the composition of the corresponding sequence observed.

References

1. Koornneef M, Meinke D. The development of Arabidopsis as a model plant. Plant J. 2010;61(6):909-21.
2. The Arabidopsis Genome Initiative. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. Nature. 2000;408(6814):796-815.
3. Kumekawa N, Hosouchi T, Tsuruoka H, Kotani H. The size and sequence organization of the centromeric region of arabidopsis thaliana chromosome 5. DNA Res. 2000;7(6):315-21.
4. Long Q, Rabanal FA, Meng D, Huber CD, Farlow A, Platzer A, et al. Massive genomic variation and strong selection in Arabidopsis thaliana lines from Sweden. Nat Genet. 2013;45(8):884-90. Epub 2013 Jun 23.
5. Vukašinović N, Cvrčková F, Eliáš M, Cole R, Fowler JE, Žárský V, et al. Dissecting a hidden gene duplication: the Arabidopsis thaliana SEC10 locus. PLoS ONE. 2014;9(4):e94077.
6. Schneeberger K, Ossowski S, Ott F, Klein JD, Wang X, Lanz C, et al. Reference-guided assembly of four diverse Arabidopsis thaliana genomes. Proceedings of the National Academie of Sciences of the United States of America. 2011;108(25):10249-54. Epub 2011 Jun 6.
7. Li YH, Zhou G, Ma J, Jiang W, Jin LG, Zhang Z, et al. De novo assembly of soybean wild relatives for pan-genome analysis of diversity and agronomic traits. Nat Biotechnol. 2014;32(10):1045-54. Epub 2014 Sep 14.
8. Kim KE, Peluso P, Babayan P, Yeadon PJ, Yu C, Fisher WW, et al. Long-read, whole-genome shotgun sequence data for five model organisms. Scientific Data. 2014;1:140045.
9. Berlin K, Koren S, Chin CS, Drake JP, Landolin JM, Phillippy AM. Assembling large genomes with single-molecule sequencing and locality-sensitive hashing. Nat Biotechnol. 2015;33(6):623-30. Epub 2015 May 25.
10. Chin CS, Peluso P, Sedlazeck FJ, Nattestad M, Concepcion GT, Clum A, et al. Phased diploid genome assembly with single-molecule real-time sequencing. Nature Methods. 2016;13(12):1050-4. Epub 2016 Oct 17.
11. Pucker B, Holtgräwe D, Rosleff Sörensen T, Stracke R, Viehöver P, Weisshaar B. A De Novo Genome Sequence Assembly of the Arabidopsis thaliana Accession Niederzenz-1 Displays Presence/Absence Variation and Strong Synteny. PLoS ONE. 2016;11(10):e0164321.
12. Zapata L, Ding J, Willing EM, Hartwig B, Bezdan D, Jiao WB, et al. Chromosome-level assembly of Arabidopsis thaliana Ler reveals the extent of translocation and inversion polymorphisms. Proc Natl Acad Sci USA. 2016.
13. Michael TP, Jupe F, Bemm F, Motley ST, Sandoval JP, Lanz C, et al. High contiguity Arabidopsis thaliana genome assembly with a single nanopore flow cell. Nature Communications. 2018;9(1):541.
14. Consortium TG. 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. Cell. 2016;166(2):481-91. Epub 2016 Jun 9.
15. Huddleston J, Chaisson MJP, Steinberg KM, Warren W, Hoekzema K, Gordon D, et al. Discovery and genotyping of structural variation from long-read haploid genome sequence data. Genome Res. 2017;27(5):677-85. Epub 2016 Nov 28.
16. Simpson JT, Pop M. The Theory and Practice of Genome Sequence Assembly. Annual Review of Genomics and Human Genetics. 2015;16:153-72. Epub 2015 Apr 22.
17. Rhoads A, Au KF. PacBio Sequencing and Its Applications. Genomics Proteomics Bioinformatics. 2015;13(5):278-89. Epub 2015 Nov 2.
18. Lam KK, Khalak A, D. T. Near-optimal assembly for shotgun sequencing with noisy reads. BMC Bioinf. 2014;15:S4. Epub 2014 Sep 10. doi: 10.1186/1471-2105-15-S9-S4.

19.   Koren S, Phillippy AM. One chromosome, one contig: complete microbial genomes
from long-read sequencing and assembly. Current Opinion in Microbiology. 2015;23:110-20.
Epub 2014 Dec 1.
20.   Shoromony I, Courtade T, Tse D. Do Read Errors Matter for Genome Assembly?
IEEE International Symposium on Information Theory (ISIT); Hong Kong2015. p. 919-23.
21.   Koren S, Harhay GP, Smith TP, Bono JL, Harhay DM, Mcvey SD, et al. Reducing
assembly complexity of microbial genomes with single-molecule sequencing. Genome Biol.
2013;14(9):R101.
22.   Pendleton M, Sebra R, Pang AW, Ummat A, Franzen O, Rausch T, et al. Assembly
and diploid architecture of an individual human genome via single-molecule technologies.
Nature Methods. 2015;12(8):780-6. Epub 2015 Jun 29.
23.   Payne A, Holmes N, Rakyan V, Loose M. BulkVis: a graphical viewer for Oxford
nanopore bulk FAST5 files. Bioinformatics. 2018.
24.   Istace B, Friedrich A, d'Agata L, Faye S, Payen E, Beluche O, et al. de novo
assembly and population genomic survey of natural yeast isolates with the Oxford Nanopore
MinION sequencer. Gigascience. 2017;6(2):1-13.
25.   Healey A, Furtado A, Cooper T, Henry RJ. Protocol: a simple method for extracting
next-generation sequencing quality genomic DNA from recalcitrant plant species. Plant
Methods. 2014;10(21). doi: 10.1186/1746-4811-10-21.
26.   Stadermann KB, Holtgräwe D, Weisshaar B. Chloroplast Genome Sequence of
Arabidopsis thaliana Accession Landsberg erecta, Assembled from Single-Molecule, Real-
Time Sequencing Data. Genome Announcements. 2016;4(5):e00975-16.
27.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable
and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res. 2017;27(5):722-36. Epub 2017 Mar 15.
28.   Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. Scaffolding pre-assembled
contigs using SSPACE. Bioinformatics. 2011;27(4):578-9. Epub 2010 Dec 12.
29.   Kleinboelting N, Huep G, Appelhagen I, Viehoever P, Li Y, Weisshaar B. The
Structural Features of Thousands of T-DNA Insertion Sites Are Consistent with a Double-
Strand Break Repair-Based Insertion Mechanism. Molecular Plant. 2015;8(11):1651-64.
Epub 2015 Sep 5.
30.   Stothard P, Wishart DS. Circular genome visualization and exploration using CGView.
Bioinformatics. 2005;21(4):537-9. Epub 2004 Oct 12.
31.   Kolmogorov M, Yuan J, Lin YR, Pevzner PA. Assembly of Loing Error-Prone Reads
Using Repeat Graphs. 2018.
32.   Li H. Minimap and miniasm: fast mapping and de novo assembly for noisy long
sequences. Bioinformatics. 2016;32(14):2103-10. Epub 2016 Mar 19.
33.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. Oxford University Press. 2013:1-3.
34.   Pucker B, Brockington SF. Genome-wide analyses supported by RNA-Seq reveal
non-canonical splice sites in plant genomes. BMC Genomics. 2018;19(1):980.
35.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an
integrated tool for comprehensive microbial variant detection and genome assembly
improvement. PLoS ONE. 2014;9(11):e112963.
36.   Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JA. Primer3Plus,
an enhanced web interface to Primer3. Nucleic Acids Res. 2007;35:W71-4. Epub 2007 May
7.
37.   Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B. An *Arabidopsis*
thaliana T-DNA mutagenised population (GABI-Kat) for flanking sequence tag based reverse
genetics. Plant Mol Biol. 2003;53(1):247-59.
38.   Pucker B, Holtgräwe D, Weisshaar B. Consideration of non-canonical splice sites
improves gene prediction on the Arabidopsis thaliana Niederzenz-1 genome sequence. BMC
Res Notes. 2017;10(1):667.
39.   Cheng CY, Krishnakumar V, Chan A, Thibaud-Nissen F, Schobel S, Town CD.
Araport11: a complete reannotation of the Arabidopsis thaliana reference genome. Plant J.
2017;(89):789-804.

40.   Arend D, Junker A, Scholz U, Schüler D, Wylie J, Lange M. PGP repository: a plant
phenomics and genomics data publication infrastructure. Database. 2016.
41.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 2015;31(19):3210-2. Epub 2015 Jun 9.
42.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al. Versatile
and open software for comparing large genomes. Genome Biol. 2004;5(2):R12. Epub 2004
Jan 30.
43.   Khelik K, Lagesen K, Sandve GK, Rognes T, Nederbragt AJ. NucDiff: in-depth
characterization and annotation of differences between two sets of DNA sequences. BMC
Bioinf. 2017;18(1):338.
44.   Keller O, Kollmar M, Stanke M, Waack S. A novel hybrid gene prediction method
employing protein multiple sequence alignments. Bioinformatics. 2011;27(6):757-63. Epub
2011 Jan 6.
45.   Schmid KJ, Sorensen TR, Stracke R, Torjek O, Altmann T, Mitchell-Olds T, et al.
Large-scale identification and analysis of genome-wide single-nucleotide polymorphisms for
mapping in *Arabidopsis thaliana*. Genome Res. 2003;13(6A):1250-7. Epub 2003/06/12. doi:
10.1101/gr.728603. PubMed PMID: 12799357.
46.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
tool. J Mol Biol. 1990;215(3):403-10.
47.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80. Epub 2013
Jan 16.
48.   Smit AFA, Hubley R, Green P. RepeatMasker Open-4.0 2013-2015. Available from:
http://www.repeatmasker.org.
49.   Paten B, Earl D, Nguyen N, Diekhans M, Zerbino D, Haussler D. Cactus: Algorithms
for genome multiple sequence alignment. Genome Res. 2011;21(9):1512-28. Epub 2011 Jun
10.
50.   Fiddes IT, Armstrong J, Diekhans M, Nachtweide S, Kronenberg ZN, Underwood JG,
et al. Comparative Annotation Toolkit (CAT)-simultaneous clade and personal genome
annotation. Genome Res. 2018;28(7):1029-38. Epub 2018 Jun 8.
51.   Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics. 2013;29(22):2933-5. Epub 2013 Sep 4.
52.   Kalvari I, Argasinska J, Quinones-Olvera N, Nawrocki EP, Rivas E, Eddy SR, et al.
Rfam 13.0: shifting to a genome-centric resource for non-coding RNA families. Nucleic Acids
Res. 2018;46(D1):D335-D42.
53.   Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA
genes in genomic sequence. Nucleic Acids Res. 1997;25(5):955-64.
54.   Silverstein KA, Graham MA, Paape TD, VandenBosch KA. Genome organization of
more than 300 defensin-like genes in Arabidopsis. Plant Physiol. 2005;138(2):600-10.
55.   Cho YS, Kim H, Kim HM, Jho S, Jun J, Lee YJ, et al. An ethnically relevant
consensus Korean reference genome is a step towards personal reference genomes. Nature
Communications7. 2016:13637.
56.   Seo JS, Rhie A, Kim J, Lee S, Sohn MH, Kim CU, et al. De novo assembly and
phasing of a Korean human genome. Nature. 2016;538(7624):243-7. Epub 2016 Oct 5.
57.   Chandrasekhara C, Mohannath G, Blevins T, Pontvianne F, Pikaard CS.
Chromosome-specific NOR inactivation explains selective rRNA gene silencing and dosage
control in Arabidopsis. Genes Dev. 2016;30(2):177-90. Epub 2016 Jan 7.
58.   Copenhaver GP, Pikaard CS. RFLP and physical mapping with an rDNA-specific
endonuclease reveals that nucleolus organizer regions of Arabidopsis thaliana adjoin the
telomeres on chromosomes 2 and 4. Plant J. 1996;9(2):259-72.
59.   Hu TT, Pattyn P, Bakker EG, Cao J, Cheng JF, Clark RM, et al. The Arabidopsis
lyrata genome sequence and the basis of rapid genome size change. Nat Genet.
2011;43(5):476-81. Epub 2011 Apr 10.

60. Villasante A, Abad JP, Méndez-Lago M. Centromeres were derived from telomeres
during the evolution of the eukaryotic chromosome. Proceedings of the National Academy of
Sciences of the United Stated of America. 2007;104(25):10542-7. Epub 2007 Jun 8.
61. Stupar RM, Lilly JW, Town CD, Cheng Z, Kaul S, Buell CR, et al. Complex mtDNA
constitutes an approximate 620-kb insertion on Arabidopsis thaliana chromosome 2:
implication of potential sequencing errors caused by large-unit repeats. Proceedings of the
National Academy of Sciences of the United Stated of America. 2001;98(9):5099-103. Epub
2001 Apr 17.
62. Kowalski SP, Lan TH, Feldmann KA, Paterson AH. Comparative mapping of
Arabidopsis thaliana and Brassica oleracea chromosomes reveals islands of conserved
organization. Genetics. 1994;138(2):499-510.
63. Fransz PF, Armstrong S, de Jong JH, Parnell LD, van Drunen C, Dean C, et al.
Integrated cytogenetic map of chromosome arm 4S of A. thaliana: structural organization of
heterochromatic knob and centromere region. Cell. 2000;100(3):367-76.
64. Unseld M, Marienfeld JR, Brandt P, Brennicke A. The mitochondrial genome of
Arabidopsis thaliana contains 57 genes in 366,924 nucleotides. Nat Genet. 1997;15(1):57-
61.
65. Martínez-Zapater JM, Gil P, Capel J, Somerville CR. Mutations at the Arabidopsis
CHM locus promote rearrangements of the mitochondrial genome. Plant Cell. 1992;4(8):889-
99.
66. Christensen AC. Plant mitochondrial genome evolution can be explained by DNA
repair mechanisms. Genome Biology and Evolution. 2013;5(6):1079-86.
67. Preuten T, Cincu E, Fuchs J, Zoschke R, Liere K, Börner T. Fewer genes than
organelles: extremely low and variable gene copy numbers in mitochondria of somatic plant
cells. Plant J. 2010;64(6):948-59. Epub 2010 Nov 4.
68. Woloszynska M, Gola EM, Piechota J. Changes in accumulation of heteroplasmic
mitochondrial DNA and frequency of recombination via short repeats during plant lifetime in
Phaseolus vulgaris. Acta Biochim Pol. 2012;59(4):703-9. Epub 2012 Dec 6.
69. Wendel JF, Jackson SA, Meyers BC, Wing RA. Evolution of plant genome
architecture. Genome Biol. 2016;17(37):s13059-016-0908-1.
70. Denton JF, Lugo-Martinez J, Tucker AE, Schrider DR, Warren WC, Hahn MW.
Extensive error in the number of genes inferred from draft genome assemblies. PLoS
Computational Biology. 2014;10(12):e1003998.
71. Belser C, Istace B, Denis E, Dubarry M, Baurens FC, Falentin C, et al. Chromosome-
scale assemblies of plant genomes using nanopore long reads and optical maps. Natural
Plants. 2018;4(11):879-87. Epub 2018 Nov 2.
72. Paquette SM, Bak S, Feyereisen R. Intron-exon organization and phylogeny in a
large superfamily, the paralogous cytochrome P450 genes of Arabidopsis thaliana. DNA Cell
Biol. 2000;19(5):307-17.
73. Panchy N, Lehti-Shiu M, Shiu SH. Evolution of Gene Duplication in Plants. Plant
Physiol. 2016;171(4):2294-316. Epub 2016 Jun 10.
74. Tan S, Zhong Y, Hou H, Yang S, Tian D. Variation of presence/absence genes
among Arabidopsis populations. BMC Evolutionary Biology. 2012;12(86):1471-2148/12/86.
75. Benovoy D, Drouin G. Processed pseudogenes, processed genes, and spontaneous
mutations in the Arabidopsis genome. J Mol Evol. 2006;62(5):511-22. Epub 2006 Apr 11.
76. Zou C, Lehti-Shiu MD, Thibaud-Nissen F, Prakash T, Buell CR, Shiu SH. Evolutionary
and expression signatures of pseudogenes in Arabidopsis and rice. Plant Physiol.
2009;151(1):3-15. Epub 2009 Jul 29.
77. Yamada K, Lim J, Dale JM, Chen H, Shinn P, Palm CJ, et al. Empirical analysis of
transcriptional activity in the Arabidopsis genome. Science. 2003;302(5646):842-6.
78. Siena LA, Ortiz JP, Calderini O, Paolocci F, Cáceres ME, Kaushal P, et al. An
apomixis-linked ORC3-like pseudogene is associated with silencing of its functional homolog
in apomictic Paspalum simplex. J Exp Bot. 2016;67(6):1965-78. Epub 2016 Feb 2.
79. Zmienko A, Samelak-Czajka A, Kozlowski P, Szymanska M, Figlerowicz M.
Arabidopsis thaliana population analysis reveals high plasticity of the genomic region

spanning MSH2, AT3G18530 and AT3G18535 genes and provides evidence for NAHR-driven recurrent CNV events occurring in this location. BMC Genetics. 2016;17(1):893.

80.   Yang L, Takuno S, Waters ER, Gaut BS. Lowly expressed genes in Arabidopsis thaliana bear the signature of possible pseudogenization by promoter degradation. Mol Biol Evol. 2011;28(3):1193-203. Epub 2010 Nov 8.

81.   Marroni F, Pinosio S, Morgante M. Structural variation and genome complexity: is dispensable really dispensable? Curr Opin Plant Biol. 2014;18:31-6. Epub 2014 Feb 16.

82.   Golicz AA, Batley J, Edwards D. Towards plant pangenomics. Plant Biotechnol J. 2016;14(4):1099-105. Epub 2015 Nov 23.

83.   Arumuganathan K, Earle ED. Nuclear DNA Content of Some Important Plant Species. Plant Mol Biol Reptr. 1991;9(3):208-18.

84.   Höfte H, Desprez T, Amselm L, Chiapello H, Caboche M, Moisan A, et al. An inventory of 1152 expressed sequence tags obtained by partial sequencing of cDNAs from *Arabidopsis thaliana.* Plant J. 1993;4(6):1051-61.

 Figure Legends

 Figure 1: Nd-1 genome structure.

 Schematic pseudochromosomes are represented by black lines with positions of genomics

 features highlighted with colored icons as indicated in the insert.

 Figure 2: Inversion on chromosome 4.

 The dotplot heatmaps show the similarity between small fragments of two sequences. Each

 dot indicates a match of 1 kbp between both sequences, while the color is indicating the

 similarity of the matching sequences. A red line highlights an inversion between Nd-1 and

 Col-0 or L*er* and Col-0, respectively. A red arrow points at the position where the inversion

 alleles differ between Nd-1 and L*er*. (a) Comparison of the Nd-1 genome sequence against

 the Col-0 gold standard sequence reveals a 1 Mbp inversion. (b) The L*er* genome sequence

 displays another inversion allele [12].

 Figure 3: Highly divergent region on chromosome 2.

 There is a very low similarity region (light blue) between the sequences in region A and

 almost no similarity between the sequences in region B (white). The complete region

 between 3.29 Mbp and 3.48 Mbp on NdChr2 is missing in the L*er* genome.

 Figure 4: Hidden locus in the Col-0 reference sequence.

 Differences between the Nd-1 and Col-0 genome sequences lead to the discovery of a

 collapsed region in the Col-0 gold standard sequence. There are two copies of At4g22214

 (blue) present in the Col-0 genome, while only one copy is represented in the Col-0 gold

 standard sequence. This gene duplication was initially validated through PCR with outwards

facing oligonucleotides N258 and N259 (purple) which lead to the formation of the expected

PCR product (black). Parts of this region were cloned into plasmids (grey) for sequencing.

Sanger and paired-end Illumina sequencing reads revealed one complete gene

(At4g22214b) and a degenerated copy (At4g22214a). Moreover, the region downstream of

the complete gene copy in Nd-1 indicates the presence of at least one additional

degenerated copy.

1087

1088

1089 **Table 1: Nd-1 *de novo* assembly statistics.**

1090 Metrics of assemblies of the Nd-1 nucleome sequence generated by Canu, FALCON,

1091 miniasm, and Flye. All described assemblies are the final version after polishing.

parameter	Ath-Nd1_v2c	Ath-Nd-1_v2f	Ath-Nd-1_v2m	Ath-Nd-1_v2y
Assembler	Canu	FALCON	Miniasm	Flye
number of contigs	69	26	72	44
total number of bases	123,513,866	119,540,544	120,159,079	116,964,092
average contig length	1,790,056 bp	4,597,713 bp	1,668,876 bp	2,658,274 bp
minimal contig length	50,345 bp	86,055 bp	50,142 bp	53,207 bp
maximal contig length	15,898,009 bp	15,877,978 bp	14,338,505 bp	14,857,908 bp
GC content	36.14 %	36.04%	36.07%	36.01%
N25	14,369,729 bp	14,534,675 bp	11,880,610 bp	12,510,540 bp
N50	13,422,481 bp	9,302,209 bp	8,595,164 bp	10,607,548 bp
N75	8,555,326 bp	6,666,836 bp	3,513,050 bp	6,001,858 bp
N90	2,928,047 bp	2,829,734 bp	1,430,525 bp	2,524876 bp

1092
1093