A Humanized Monoclonal Antibody against the Enzymatic Subunit of Ricin Toxin Rescues Rhesus macaques from the Lethality of Aerosolized Ricin

Chad J. Roy[1,*], Dylan J. Ehrbar[2], Natasha Bohorova[3], Ognian Bohorov[3], Do Kim[3], Michael Pauly[3], Kevin Whaley[3], Yinghui Rong[2], Fernando J Torres-Velez[2], Ellen S Vitetta[4], Peter J. Didier[1], Lara Doyle-Meyers[1], Larry Zeitlin[3,*], and Nicholas J. Mantis[2,*]

[1]Tulane National Primate Research Center, Covington, Louisiana USA; [2]Division of Infectious Disease, Wadsworth Center, New York State Department of Health, Albany, NY 12208; [3]Mapp Biopharmaceutical, Inc. San Diego, CA 92121; [4]Departments of Immunology and Microbiology, University of Texas Southwestern Medical Center, Dallas, TX 75390

Running title: Rescue from ricin aerosol exposure

Keywords: MAb, ricin, toxin, biodefense, therapeutic

*Corresponding authors: Chad J. Roy, Tulane National Primate Research Center, Microbiology Division, 18703 Three Rivers Road, Covington, LA 70433; Tel 985-871-6417, email: croy@tulane.edu; *Larry Zeitlin, Mapp Biopharmaceutical, Inc. San Diego, CA 92121; Phone: 858-625-0335; email: larry.zeitlin@mappbio.com;*Nicholas Mantis, Division of Infectious Disease, Wadsworth Center, 120 New Scotland Avenue, Albany, NY 12208. Phone: 518-473-7487. email: nicholas.mantis@health.ny.gov

Abstract

Ricin toxin (RT) ranks at the top of the list of potential bioweapons of concern to civilian and military personnel alike due to its high potential for morbidity and mortality after inhalation. In non-human primates, aerosolized ricin triggers a severe acute respiratory distress characterized by perivascular and alveolar edema, neutrophilic infiltration, and severe necrotizing bronchiolitis and alveolitis. There are currently no approved countermeasures for ricin intoxication. In this report, we demonstrate the therapeutic potential of huPB10, a toxin-neutralizing humanized monoclonal antibody (MAb) against an immunodominant epitope on ricin's enzymatic A chain (RTA). Five rhesus macaques that received intravenous huPB10 (10 mg/kg) four hours after lethal dose ricin aerosol exposure all survived the toxin challenge, as compared to control animals, which succumbed to ricin intoxication within 30 h. Antibody treatment at 12 h after ricin exposure resulted in the survival of only one of five monkeys, indicating that, in the majority of animals, ricin intoxication and local tissue damage had progressed beyond the point where huPB10 intervention was beneficial. Change in pro-inflammatory cytokine/chemokines levels in bronchial alveolar lavage fluids before and after toxin challenge successfully clustered monkeys based on survival, as well as treatment group. IL-6 was the most apparent marker of ricin intoxication. This study represents the first demonstration in nonhuman primates that the lethal effects of inhalational ricin exposure can be negated by a drug candidate and opens up a path forward for product development.

53   .
54
55   **Introduction**
56   Ricin toxin (RT) is considered a high priority biothreat agent by the Centers for
Disease Control and Prevention (CDC), the US Department of Defense (DOD), and
NATO due to it's accessibility, stability, and high toxicity especially by the aerosol route.
(1, 2). In nonhuman primates (NHPs), inhalation of RT elicits the clinical equivalent of
acute respiratory distress syndrome (ARDS), characterized by widespread apoptosis of
alveolar macrophages, intra-alveolar edema, neutrophilic infiltration, accumulation of
pro-inflammatory cytokines, and fibrinous exudate (3, 4). Ricin also damages the lung
mucosa and triggers vascular leak due to direct damage to endothelial cells (5). Similar
effects are observed in mice, rats and swine (6-10). RT is derived from castor beans
(*Ricinus communis*) and is a 65 kDa heterodimeric glycoprotein consisting of two
subunits, RTA and RTB, joined via a single disulfide bond. RTB binds to glycoproteins
and glycolipids on mammalian cells and facilitates the retrograde transport of RT to the
endoplasmic reticulum (ER). In the ER, RTA is liberated from RTB and retrotranslocated
into the cytoplasm *via* the Sec61 complex (11). RTA is an RNA N-glycosidase that
catalyzes the hydrolysis of a conserved adenine residue within the sarcin/ricin loop of
28S rRNA, resulting in the inhibition of protein synthesis (12, 13) and the activation of
apoptosis (14). Alveolar macrophages are particularly sensitive to the cytotoxic effects
and secrete an array of pro-inflammatory cytokines before undergoing apoptosis (8, 9,
15).
75   In this report we investigated the potential of a humanized monoclonal antibody
(MAb) huPB10 to serve as a therapeutic in an established Rhesus macaques model of
RT inhalation (3, 16). PB10 was first described as a murine MAb with potent-toxin
neutralizing activity *in vitro* and *in vivo* (17). PB10 recognizes an immunodominant
epitope situated at the apex of RTA, relative to RTB (18, 19). Chimeric (cPB10) and fully
humanized (huPB10) versions of PB10 retain *in vitro* toxin-neutralizing activity and have
been shown to passively protect mice against lethal dose of RT administered by
injection or inhalation (20, 21). It was also demonstrated that huPB10 can rescue mice
from intoxication if administered 4-6 h after exposure (21). Based on these preliminary
findings, assessed the therapeutic potential of huPB10 in a well-established nonhuman

primate (NHP) model of ricin aerosol challenge (16). The NHP is believed to be the

model most representative of aerosolized exposure.

Results

A total of 12 rhesus macaques (~7 kg; range 3.8-10.2 kg) bred at Tulane National

Primate Research Center (TNPRC) were randomly assigned to three experimental

groups and then challenged with RT by small particle aerosol at a target dose of 18

µg/kg (**Table 1; Figure 1**). Animal studies were conducted in strict compliance with

protocols approved by TNPRC's Institutional Animal Care and Use Committee (IACUC).

Group 1 (n=2) received intravenous administration of saline 4 h post challenge. Animals

in groups 2 and 3 received huPB10 at 4 h or 12 h post challenge, respectively (**Table**

1). huPB10 was administered intravenously at a final dose of 10 mg/kg (**Table S1**). The

macaques were subjected to whole body plethysmography and radiotelemetry over the

course of the study. Animals surviving on day 14 post challenge were euthanized and

subjected to complete necropsy and histopathological analysis. Serum and bronchial

alveolar lavage (BAL) fluids were collected from the animals before and 24 h after

exposure to RT.

Control animals (n=2) succumbed to RT toxicosis within 36 hours of exposure.

(**Table 1; Figure 1**). The clinical progression of RT-induced intoxication of the control

animals was the same as in previous studies (3). Approximately 12-16 hours after

exposure, animals displayed reduced activity and fever (**Figure 2**). Clinical examination

conducted 24 h post exposure revealed respiratory complications, including bilateral

congestion and crackles, with dyspnea and tachypnea. Arterial oxygen ranged from 75-

85%. The clinical state of the animals continued to decline over the subsequent several

hours, manifested by a marked drop in normal activity and cyanotic mucous membrane.

In contrast, all five of the Rhesus macaques that received huPB10 at 4 h post-

ricin challenge time point survived (*P<0.01* compared to controls) and remained

otherwise normal for the duration of the three-week post-exposure observation period

(**Table 1: Figure 1**). Compared to the control animals, the five macaques in Group 2

presented with only minimal signs of distress: mild dyspnea, minimal tachypnea and

increased lung sounds upon physical examination at 24 h post-exposure. The animals
displayed no such symptoms upon physical examination on day 7.

Finally, only one of the five animals in Group 3 that received huPB10 at 12 h
survived RT challenge (**Table 1; Figure 1;** *P<0.01* and *P<0.05* compared to controls
and +4 h group, respectively). The remaining four animals succumbed to RT intoxication
between 44 and 72 h post challenge and followed a clinical course similar to the control
animals. HuPB10 was detected in the sera and BAL fluids collected at 24 h post RT
challenge, indicating that biodistribution of huPB10 was similar between groups 2 and 3.

Gross examination of the lungs from sham-treated control animals showed
coalescing hemorrhage with frothy exudate marked by fibrin in lung parenchyma
(**Figure S1**). The lungs of control animals were grossly described as fibrinosuppurative
bronchointerstitial pneumonia with pulmonary edema and bronchial epithelial necrosis,
with severe fibrinosuppurative lymphadenitis in the bronchial lymph nodes. The wet
weights of control animal lungs were >150 grams in contrast to a normal wet lung
weight of ~30-40 grams in naïve animals of approximately the same body weight.
Histologically, the lungs of sham-treated animals showed hallmark inflammation
consistent with RT-induced injury, characterized by marked edema, corresponding
hemorrhage, and numerous infiltrates. The pathological outcome of the four animals in
Animals in Group 3 that succumbed to RT intoxication resembled the control animals.
There was extensive pulmonary congestion, edema, inflammation with infiltrates, and
punctate hemorrhage evident. Gross lung weights were similar to RT-challenged control
animals (>140 g) with clear signs of hemorrhage.

Pathological analysis of the lung tissues collected at the time of euthanasia (day 21
post challenge) from the five survivors in Group 2 that had been treated with huPB10 at
4 h and the single animal in Group 3 (DR61) revealed evidence of chronic inflammation
and a distinctive fibrosis proximal to the respiratory bronicholes, reminiscent of past
studies in which animals had received sub-lethal exposures to RT (22).

Sera and BAL fluids collected before (day -7) and 24 h post RT challenge were
subjected to analysis with a 29-plex cytokine/chemokine/growth factor Luminex array as
a means to assess the impact of huPB10 on local and systemic inflammatory
responses. In the sera of the two control animals, there were 12 cytokines/chemokines

that were significantly different in serum post- versus pre-challenge; 11 cytokines were increased, while one (IL-8) was decreased (**Figure 3A; S2**). Most notable were IL-6 (~500-fold increase) and IL-1RA (~256-fold increase). VEGF was also elevated, possibly reflecting response to pulmonary insult. Analysis of sera from animals that received huPB10 at 4 h post challenge indicated that only four cytokine/chemokines were significantly different from pre-challenge levels (3 up;1 down), including a ~4-fold increase in IL-6. IL-1RA and VEGF levels were unchanged. In animals that received huPB10 at 12 h after RT challenge, a total of six cytokines/chemokines changed relative to pre-challenge levels, although the magnitude of these changes was lower than that if the control animals, possibly reflecting a dampening of the inflammatory response as a consequence of huPB10 intervention. Principle component analysis (PCA) of fold-change in cytokine/chemokine levels from serum samples from all 12 monkeys did not reveal any clustering by experimental group or survival (**Figure S3A-C**), signifying that serum inflammatory responses are not indicative of experimental outcome.

The impact of huPB10 intervention was much more apparent in BAL fluids than in sera (**Figure 3B**). Within group 2, a total of 21 cytokines/chemokines were significantly elevated following RT challenge (as compared to pre-challenge levels), with IL-6 being the most pronounced. In group 3, there were 25 cytokines/chemokines that were significantly elevated compared to pre-challenge levels. Seven cytokines/chemokines differed between the groups 2 and 3 with the most notable being IL-6, which was 32-fold elevated in group 2 and 181-fold elevated in group 3. PCA successfully clustered animals by both group and survival status, demonstrating that the localized cytokine responses in the BAL are more closely related to survival than the systemic responses (**Figure S3D-E**). The relative contributions of the different cytokines/chemokines responsible for segregating the animals into clusters are shown in **Figure S3F**.

Discussion

The results of this study constitute a significant advance in longstanding efforts to develop effective medical countermeasures against RT inhalation (2, 23). Foremost, it is the first demonstration in NHPs that a MAb can rescue animals from the lethal effects of aerosolized ricin toxin exposure. Studies in primates are important because anti-RT

products for humans must adhere to the Food and Drug Administration's (FDA) Animal

Rule; human challenge studies with RT are obviously unethical. The well-established

model of aerosolized RT challenge in Rhesus macaques was a prerequisite for the

therapeutic study conducted herein (3, 16, 24). Comparative models in mice (6, 8, 25)

and swine (10) may also be important for therapeutic development under Animal Rule

guidelines. The fact that all five animals in group 2 (+4 h huPB10 intervention), and one

animal in group 3 (+12 h huPB10 intervention), survived exposure to RT indicates that a

significant proportion of RT remains accessible to huPB0 in the alveolar space and/or

interstitial fluids for hours after inhalational exposure. This finding is consistent with

reports in mice with huPB10 and other anti-RT MAbs (6, 7, 21, 26), but surprising all

the same considering the extraordinary sensitivity of the lung mucosa to the effects of

toxin exposure (27).

We postulate that high dose intravenous delivery of huPB10 results in the

accumulation of huPB10 within the lung mucosa and alveolar space where it can

engage free (soluble) or receptor-bound RT. Indeed, IV administration of recombinant

anti-viral IgG1 MAbs results a corresponding linear distribution of antibodies lung (28,

29). Whether transudation or active transport of serum antibodies into the lung mucosa

is triggered by RT exposure has not been evaluated. Nonetheless, once in the lung

mucosa, huPB10 presumably limits toxin uptake into target cells (e.g., macrophages

and airway epithelial cells) and/or interferes with ricin intracellular transport in the event

that endocytosis of RT-antibody complexes should occur (30, 31). Work by Magun and

colleagues made it clear more than a decade ago that protecting alveolar macrophages

is paramount in limiting toxin-induced lung damage (9, 32).

It is worth underscoring the value of the 29-plex monkey cytokine/chemokine/growth

factor array in not only identifying local inflammatory markers like IL-6 that arise

following RT exposure but also enabling through PCA the unbiased clustering of

specific animals based on survival or experimental group. Identifying an inflammatory

"fingerprint" associated with RT exposure has obvious applications in diagnostics,

especially in the context of biodefense where early clinical symptoms following

exposure to different toxins and pathogens may in fact be indistinguishable (33). As

noted above, IL-6 levels were particularly elevated following RT exposure, which is

consistent with what has been observed in mice (34). IL-6 is a particularly potent driver

of pulmonary inflammation and could very well be a major contributor to RT-induced

pathology in conjunction with RT's other properties, including agglutinin activity and the

capacity to induce vascular leak syndrome (5, 35). We recently reported that human

lung epithelial cell lines preferentially secrete IL-6 following RT exposure, especially in

the presence of pro-apoptotic factors like TRAIL (36). It has been suggested that anti-

inflammatory agents may extend the therapeutic window of anti-RT MAbs by

suppressing bystander tissue damage (26). Whether such directed immunotherapies

would have utility in the context of potent toxin-neutralizing antibody like huPb10

remains to be seen. At least early intervention with huPB10 was sufficient to render RT

effectively inert within the context of the lung and it seems unlikely that supplementing

treatment with anti-inflammatory agents would afford much additional benefit.

Methods

Ricin toxin and huPB10. Purified ricin toxin derived from castor beans (*Ricinus*

communis) was produced as previously described (37). HuPB10 was expressed using a

Nicotiana benthamiana-based manufacturing platform. The properties of huPB10 used

in this study are shown in **Table S1.**

Animal care and use. Rhesus macaques were born and housed at the Tulane

National Primate Research Center (Covington, LA), which is US Department of

Agriculture-licensed and fully accredited by the Association for Assessment and

Accreditation of Laboratory Animal Care (AAALAC). Aerosolization, dosing and delivery

of RT were performed as described (16). The LD_{50} of ricin is 5.8 µg/kg body weight and

the target dose for this experiment was set at the equivalent of three LD_{50}s ($\approx$18 µg/kg).

The mean inhaled dose of ricin across all animals was 4.4 ± 1.4 LD_{50}s. At 4 h or 12

hours post-exposure, designated animal groups received a single intravenous

administration of huPB10 by slow infusion at an individualized unit dose of 10 mg/kg.

Sham-treated animals received saline at the 4h time point. Treated animals were

observed for signs of adverse reactions to the MAb during administration and

throughout the anesthesia recovery period. Animals were bled just before and 24 h

following aerosol challenge. Blood was also collected when the animals either

succumbed to intoxication or 21 d after challenge, when the experiment was terminated.

Animals determined to be in respiratory distress and those that survived for 21 d after

exposure to ricin were euthanized by an overdose of sodium pentobarbital, consistent

with the recommendation of the American Veterinary Medical Association's Panel on

Euthanasia, and submitted for necropsy. All methods were approved by the Tulane

University's IACUC. After gross necropsy, tissues were collected in neutral buffered

zinc-formalin solution (Z-Fix Concentrate, Anatach). Tissues were processed,

sectioned, and stained as previously described (3).

Statistics. Statistical analysis was carried out with GraphPad Prism 6 (GraphPad

Software, La Jolla California USA). The difference in outcomes between groups was

determined by Fisher's exact test (two-tailed) and the mean survival times after

exposure to RT were compared by log-rank analysis of Kaplan–Meier survival curves.

The statistical significance of the effects of ricin challenge and huPB10 intervention on

cytokine levels were analyzed with two-way repeated measures ANOVAs in both serum

and BAL, with the repeated measures being pre- and post-exposure status, and

treatment group as the independent measure. Resulting p-values were corrected with

the Benjamini, Krieger, and Yekutieli method to control false discovery rate. All

analyses were performed on $\log^2$ transformed raw fluorescent intensity values to avoid

the need to censor values. PCA analysis was performed using singular value

decomposition with the R package FactoMineR (38). Heatmap construction using raw

fold change values was done using GraphPad Prism 6. Hierarchical clustering and

scaled heatmap construction was completed with the R package pheatmap (39).

Author contributions. NB, OB, DK, MP, KW and ESV generated reagents; PJD, LDM,

YR, and CJR conducted animal studies and animal tissue/sample analysis; DJE

performed statistical analysis; CJR, NJM, and LZ are responsible for experimental

design and CJR, NM and ESV prepared the manuscript.

Conflicts of interest. NB, OB, DK, MP are Mapp Biopharmaceutical employees and

shareholders. KW and LZ are employees, shareholders, and co-owners of Mapp

Biopharmaceutical. The remaining authors declare no conflicts of interest.

Acknowledgements. This work was supported by R01AI098774 to LZ and Contract

No. HHSN272201400021C to NJM from the National Institutes of Allergy and Infectious

Diseases, National Institutes of Health (NIH), as well as the National Center for

Research Resources and the Office of Research Infrastructure Programs (ORIP) of the

NIH through grant OD011104 at the Tulane National Primate Research Center to CJR.

The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health. The funders had no role

in study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

References

1. Cieslak TJ, Kortepeter MG, Wojtyk RJ, Jansen HJ, Reyes RA, Smith JO, et al.

Beyond the Dirty Dozen: A Proposed Methodology for Assessing Future Bioweapon

Threats. *Mil Med.* 2018;183(1-2):e59-e65.

2. Wolfe DN, Florence W, and Bryant P. Current biodefense vaccine programs and

challenges. *Human vaccines & immunotherapeutics.* 2013;9(7):1591-7.

3. Pincus SH, Bhaskaran M, Brey RN, 3rd, Didier PJ, Doyle-Meyers LA, and Roy CJ.

Clinical and Pathological Findings Associated with Aerosol Exposure of Macaques to

Ricin Toxin. *Toxins (Basel).* 2015;7(6):2121-33.

4. Wilhelmsen CL, and Pitt ML. Lesions of acute inhaled lethal ricin intoxication in

rhesus monkeys. *Vet Pathol.* 1996;33(3):296-302.

5. Soler-Rodriguez AM, Ghetie MA, Oppenheimer-Marks N, Uhr JW, and Vitetta ES.

Ricin A-chain and ricin A-chain immunotoxins rapidly damage human endothelial

cells: implications for vascular leak syndrome. *Exp Cell Res.* 1993;206(2):227-34.

6. Poli MA, Rivera VR, Pitt ML, and Vogel P. Aerosolized specific antibody protects

mice from lung injury associated with aerosolized ricin exposure. *Toxicon.*

1996;34(9):1037-44.

7. Pratt TS, Pincus SH, Hale ML, Moreira AL, Roy CJ, and Tchou-Wong KM.

Oropharyngeal aspiration of ricin as a lung challenge model for evaluation of the

therapeutic index of antibodies against ricin A-chain for post-exposure treatment. *Exp*

Lung Res. 2007;33(8-9):459-81.

8. Wong J, Korcheva V, Jacoby DB, and Magun B. Intrapulmonary delivery of ricin at

high dosage triggers a systemic inflammatory response and glomerular damage. *Am*

J Pathol. 2007;170(5):1497-510.

9. Lindauer ML, Wong J, Iwakura Y, and Magun BE. Pulmonary inflammation triggered

by ricin toxin requires macrophages and IL-1 signaling. *J Immunol.*

2009;183(2):1419-26.

10.   Katalan S, Falach R, Rosner A, Goldvaser M, Brosh-Nissimov T, Dvir A, et al. A

novel swine model of ricin-induced acute respiratory distress syndrome. *Dis Model*

Mech. 2017;10(2):173-83.

11.   Spooner RA, and Lord JM. How Ricin and Shiga Toxin Reach the Cytosol of

Target Cells: Retrotranslocation from the Endoplasmic Reticulum. *Curr Top Microbiol*

Immunol. 2012;357:19-40.

12.   Endo Y, Mitsui K, Motizuki M, and Tsurugi K. The mechanism of action of ricin

and related toxic lectins on eukaryotic ribosomes. The site and the characteristics of

the modification in 28 S ribosomal RNA caused by the toxins. *J Biol Chem.*

1987;262(12):5908-12.

13.   Endo Y, and Tsurugi K. RNA N-glycosidase activity of ricin A-chain. Mechanism

of action of the toxic lectin ricin on eukaryotic ribosomes. *J Biol Chem.*

1987;262(17):8128-30.

14.   Tesh VL. The induction of apoptosis by Shiga toxins and ricin. *Curr Top Microbiol*

Immunol. 2012;357:137-78.

15.   Brown RF, and White DE. Ultrastructure of rat lung following inhalation of ricin

aerosol. *Int J Exp Pathol.* 1997;78(4):267-76.

16.   Roy CJ, Brey RN, Mantis NJ, Mapes K, Pop IV, Pop LM, et al. Thermostable ricin

vaccine protects rhesus macaques against aerosolized ricin: Epitope-specific

neutralizing antibodies correlate with protection. *Proc Natl Acad Sci U S A.*

2015;112(12):3782-7.

331 17. O'Hara JM, Neal LM, McCarthy EA, Kasten-Jolly JA, Brey RN, 3rd, and Mantis
332   NJ. Folding domains within the ricin toxin A subunit as targets of protective
333   antibodies. *Vaccine.* 2010;28:7035-46.

334 18. Toth Rt, Angalakurthi SK, Van Slyke G, Vance DJ, Hickey JM, Joshi SB, et al.
335   High-Definition Mapping of Four Spatially Distinct Neutralizing Epitope Clusters on
336   RiVax, a Candidate Ricin Toxin Subunit Vaccine. *Clin Vaccine Immunol.* 2017.

337 19. Vance DJ, and Mantis NJ. Resolution of two overlapping neutralizing B cell
338   epitopes within a solvent exposed, immunodominant alpha-helix in ricin toxin's
339   enzymatic subunit. *Toxicon.* 2012;60(5):874-7.

340 20. Sully EK, Whaley K, Bohorova N, Bohorov O, Goodman C, Kim D, et al. A
341   tripartite cocktail of chimeric monoclonal antibodies passively protects mice against
342   ricin, staphylococcal enterotoxin B and Clostridium perfringens epsilon toxin. *Toxicon.*
343   2014;92:36-41.

344 21. Van Slyke G, Sully EK, Bohorova N, Bohorov O, Kim D, Pauly MH, et al.
345   Humanized Monoclonal Antibody That Passively Protects Mice against Systemic and
346   Intranasal Ricin Toxin Challenge. *Clin Vaccine Immunol.* 2016;23(9):795-9.

347 22. Bhaskaran M, Didier PJ, Sivasubramani SK, Doyle LA, Holley J, and Roy CJ.
348   Pathology of lethal and sublethal doses of aerosolized ricin in rhesus macaques.
349   *Toxicol Pathol.* 2014;42(3):573-81.

350 23. Reisler RB, and Smith LA. The need for continued development of ricin
351   countermeasures. *Adv Prev Med.* 2012;2012:149737.

352 24. Roy CJ, Song K, Sivasubramani SK, Gardner DJ, and Pincus SH. Animal Models
353   of Ricin Toxicosis. *Curr Top Microbiol Immunol.* 2012.

354 25. Wong J, Korcheva V, Jacoby DB, and Magun BE. Proinflammatory responses of
355   human airway cells to ricin involve stress-activated protein kinases and NF-kappaB.
356   *Am J Physiol Lung Cell Mol Physiol.* 2007;293(6):L1385-94.

357 26. Gal Y, Mazor O, Falach R, Sapoznikov A, Kronman C, and Sabo T. Treatments
358   for Pulmonary Ricin Intoxication: Current Aspects and Future Prospects. *Toxins
359   (Basel).* 2017;9(10).

27.   van Deurs B, Hansen SH, Petersen OW, Melby EL, and Sandvig K. Endocytosis,
intracellular transport and transcytosis of the toxic protein ricin by a polarized
epithelium. *EurJCell Biol.* 1990;51(1):96-109.
28.   Dall'Acqua WF, Kiener PA, and Wu H. Properties of human IgG1s engineered for
enhanced binding to the neonatal Fc receptor (FcRn). *J Biol Chem.*
2006;281(33):23514-24.
29.   Wu H, Pfarr DS, Johnson S, Brewah YA, Woods RM, Patel NK, et al.
Development of motavizumab, an ultra-potent antibody for the prevention of
respiratory syncytial virus infection in the upper and lower respiratory tract. *J Mol Biol.*
2007;368(3):652-65.
30.   O'Hara JM, and Mantis NJ. Neutralizing monoclonal antibodies against ricin's
enzymatic subunit interfere with protein disulfide isomerase-mediated reduction of
ricin holotoxin in vitro. *J Immunol Methods.* 2013;395(1-2):71-8.
31.   Yermakova A, Klokk TI, O'Hara JM, Cole R, Sandvig K, and Mantis NJ.
Neutralizing Monoclonal Antibodies against Disparate Epitopes on Ricin Toxin's
Enzymatic Subunit Interfere with Intracellular Toxin Transport. *Sci Rep.*
2016;6:22721.
32.   Wong J, Magun BE, and Wood LJ. Lung inflammation caused by inhaled
toxicants: a review. *Int J Chron Obstruct Pulmon Dis.* 2016;11:1391-401.
33.   Lipscomb MF, Hutt J, Lovchik J, Wu T, and Lyons CR. The pathogenesis of
acute pulmonary viral and bacterial infections: investigations in animal models. *Annu*
Rev Pathol. 2010;5:223-52.
34.   Sabo T, Gal Y, Elhanany E, Sapoznikov A, Falach R, Mazor O, et al. Antibody
treatment against pulmonary exposure to abrin confers significantly higher levels of
protection than treatment against ricin intoxication. *Toxicol Lett.* 2015;237(2):72-8.
35.   Rubini A. Interleukin-6 and lung inflammation: evidence for a causative role in
inducing respiratory system resistance increments. *Inflamm Allergy Drug Targets.*
2013;12(5):315-21.
36.   Rong Y, Westfall J, Ehrbar D, LaRocca T, and Mantis N. TRAIL (CD253)
Sensitizes Human Airway Epithelial Cells to Toxin-Induced Cell Death. *bioRxiv.* 2018.

37.   Smallshaw JE, Firan A, Fulmer JR, Ruback SL, Ghetie V, and Vitetta ES. A novel
recombinant vaccine which protects mice against ricin intoxication. *Vaccine.*
2002;20(27-28):3422-7.
38.   Lê S, Josse J, and Husson F. FactoMineR: An R Package for Multivariate
Analysis. *Journal of Statistical Software.* 2008;25(1):1-18.
39.   Kolde R. pheatmap: Pretty Heatmaps. https://cran.r-
project.org/web/packages/pheatmap/index.html. Updated 2018-05-19.
397
398

Figure Legends

Figure 1. Effect of huPB10 on the survival of animals exposed to aerosolized RT A. Animals were exposed to aerosolized RT aerosol and the individual RT doses , expressed in µg/kg inhaled, and clustered by treatment group, show minimal variation in dosing, bars represent group mean ± standard deviation; the aerosol target dose of 18 µg/kg represented by segmented line. B. Macaque survival is represented by survival curve. The two control animals succumbed to intoxication by day 2; treatment with huPB10 at 4 h post ricin challenge resulted in 5/5 survival, while treatment with huPB10 at 12 h resulted in only 1/5 survivors.

Figure 2. Physiological response to RT. A. Radiotelemetry of core temperature of rhesus macaques treated with huPB10 either at 4 h or 12 h post-exposure to aerosolized RT. Continuous monitoring showed significant differences ($P<0.005$) between the initiation and tempo of pyrexia that was dependent on the time to treat. B. Fever intensity showed differences between the relative change (increase) based upon time to treat. Respiratory function measured by whole-body plethysmography. Respiratory function was measured *via* head-out conductance plethysmography prior to and at timed intervals post challenge. C. Significant changes ($P<0.05$) in post-exposure group tidal volume was observed among animals rescued from ricin intoxication, (D) although compensatory changes in frequency resulted in minimal observable changes in minute volume, with some animals producing increased minute volumes +24 h post-exposure to ricin.

Figure 3. Cytokine profiles of serum and BAL in RT challenged Rhesus macaques. Heatmaps present log_2-fold change of cytokines in (A) sera and (B) BALs, comparing samples collected 24 hours post-challenge to those collected 7 days pre-challenge. Animals are separated into groups for controls, 4-hour, and 12-hour huPB10 interventions. Increases in cytokine levels are shown in red, while decreases are shown in blue. Specific values for the fold changes are shown within each cell and based on single 29-plex analysis. A subset of cytokine changes were confirmed using BD™ CBA Human Inflammatory Cytokines Kit.

430 **Tables**
431

Table 1. Experimental groups and outcome of ricin challenge

					huPB10 (μg/ml)[c]			
Group	**ID**	**kg**	**M/F**[a]	**RT**[b]	**serum**	**BAL**	**TTD**[d]	**#/total**
Control	DH27	9.7	M	14	-	-	28 h	
	JP56	8.4	M	26	-	-	30 h	0/2
+4 h huPB10	LJ80	4.4	M	31	157	2.8	+	
	LB98	4.9	M	29	177	2.7	+	
	EM61	9.5	M	31	207	2.5	+	
	IC70	5.9	F	26	225	3.6	+	
	IL40	6.1	F	33	147	3.2	+	5/5
+12 h huPB10	JM39	9.4	M	22	205	42.8	72 h	
	DR61	10.2	F	24	265	3.9	+	
	KP48	5.8	M	30	203	25.7	48 h	
	KR49	5.7	M	29	151	3.1	49 h	
	LB26	3.8	M	29	191	11.3	44 h	1/5

[a], sex of animal; [b], ricin toxin (RT) dose (μg/kg) received per animal; [c], huPB10 in serum and BAL from samples collected at 24 h post challenge. The average in serum was 182.6 μg/ml and 203 μg/ml for the +4 and +12 h groups, respectively. In the BAL, the was 3.04 μg/ml and 17.3 μg/ml, respectively. Times of huPB10 administration are approximate (~ 30 minutes of point estimate); the logistics of performing slow intravenous infusion of huPB10 in monkeys initiated and was times at 4 hours from each of the individual ricin aerosol challenge events and was performed in sequence; [d], time to death (TTD); +, indicates survival for 21 day duration of study.

432

433

Supplemental Material

<u>Supplemental Methods</u>

Animal Husbandry and Telemetry. Rhesus macaques were born and housed at the Tulane National Primate Research Center (Covington, LA), which is US Department of Agriculture-licensed and fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care. Subcutaneous radiotelemetry transmitters combined with sensors capable of detecting biopotential signals of an electrocardiogram as well as thermistor-type sensors capable of detecting temperature signals (T34G-8; Konigsberg Instruments) were surgically implanted under aseptic conditions in 8 of the 12 vaccinated macaques and both control macaques before the start of the study. Animals determined to be in respiratory distress and those that survived for 21 d after exposure to ricin were euthanized by an overdose of sodium pentobarbital, consistent with the recommendation of the American Veterinary Medical Association's Panel on Euthanasia, and submitted for necropsy. All methods were approved by the Tulane Institutional Animal Care and Use Committee (IACUC).

Treatment of Rhesus macaques. At 4 h or 12 hours postexposure, designated animal groups received a single intravenous administration of huPB10 by slow infusion at an individualized unit dose of 10 mg/kg. Sham-treated animals were administered saline for injection at the four-hour time point. Treated animals were observed for signs of adverse reactions to the antibody during administration and throughout anesthesia recovery period. Animals were bled just before and 24 h following aerosol challenge. Blood was also collected when the animals either succumbed to intoxication or 21 d after challenge, when the experiment was terminated.

RT aerosolization, dosing, and calculation. Aerosolization, dosing and delivery of ricin were performed as described (16). Inductive plethysmography that measures volume of air breathed by each individual animal per minute was performed just before the ricin exposure. Ricin was dissolved in 10 mL sterile phosphate buffer saline to the

desired concentration for each animal based on plethysmography data obtained 2 d

before the exposure. Aerosols were generated directly into a head-only chamber using

a Collision three jet-nebulizer (BGI) with fully automated management control system

(Biaera Technologies, Hagerstown, MD) all within a Class III biological safety cabinet

housed within the TNPRC high-containment (BSL-3) laboratories. The nebulizer

operated at 18 lb/inch^2 equating to a flow of 7.5 L/min and produced 3.0E+04 particles

per cc with a mass median aerodynamic diameter of ~1.4 μm. Each discrete aerosol

exposure lasted 10 minutes (per animal). Air samples were continuously obtained

during the exposure and the protein concentrations of these samples were determined

using a micro-BSA protein assay kit (Thermo Scientific). The aerosol concentrations

were determined and the inhaled dose of RT for each animal was calculated by

multiplying the empirically determined aerosol exposure concentration (μg/liter of air) in

the chamber by volume of air estimated to have been breathed by the animal (via

results of plethysmography just before exposure). The LD_{50} of ricin is 5.8 μg/kg body

weight and the target dose for this experiment was set at the equivalent of three LD_{50}s

(≈18 μg/kg). The mean inhaled dose of ricin across all animals was 4.4 ± 1.4 LD_{50}s.

Statistics. Statistical analysis was carried out with GraphPad Prism 6 (GraphPad

Software, La Jolla California USA). The difference in outcomes between groups was

determined by Fisher's exact test (two-tailed) and the mean survival times after

exposure to ricin were compared by log-rank analysis of Kaplan–Meier survival curves.

The statistical significance of the effects of ricin challenge and huPB10 intervention on

cytokine levels were analyzed with two-way repeated measures ANOVAs in both serum

and BAL, with the repeated measures being pre- and post-exposure status, and

treatment group as the independent measure. Resulting p-values were corrected with

the Benjamini, Krieger, and Yekutieli method to control false discovery rate. All

analyses were performed on log^2 transformed raw fluorescent intensity values to avoid

the need to censor values. PCA analysis was performed using singular value

decomposition with the R package FactoMineR (38). Heatmap construction using raw

fold change values was done using GraphPad Prism 6. Hierarchical clustering and

scaled heatmap construction was completed with the R package pheatmap (39).

Tissue Collection, Histological Analysis, and Special Stains. After gross necropsy,

tissues were collected in neutral buffered zinc-formalin solution (Z-Fix Concentrate,

Anatach). Tissues were processed, sectioned, and stained as previously described (3).

Supplemental Table

Table S1. Characteristics of huPB10 used in study		
Test Parameters	**Test Method**	**Result**
Concentration	UV Absorbance	19.7 mg/mL
Appearance	Visible	clear, liquid
Physical/chemical	pH Determination	5.6
Purity	SDS-PAGE	> 99%
Osmolality	Micro-Osmometer	412 mOS/kg
Bioburden	Bioburden Testing	0 CFU/mL
Aggregates	Size Exclusion HPLC	1.90%
Potency	ELISA	20.0 mg/mL
Safety	Endotoxin	0.2 EU/mg

Figure S1. Gross pathology associated with RT exposure and huPB10 intervention.   The lungs of sham-treated animals (Figure 3a) show hallmark inflammation, marked edema, corresponding hemorrhage, and numerous infiltrates; grossly, wet weight at 3x normal with coalescing hemorrhage. Animals treated with huPB10 at 4 h (Figure 3b & 3c) show remarkably little pulmonary damage, with occasional infiltrates; lung wet weights were essentially normal. Animals treated with huPB10 at 12 hours (Figure 3d) demonstrated mild to moderate inflammation with infiltrates, edema, and punctate hemorrhage evident; gross lung weight were approximately 2x with clear signs of hemorrhage. Original magnification at 10x

Fig. S2 Scaled heatmap and cluster analysis of log2-fold changes in cytokine levels in individual macaques following ricin challenge. Heatmaps visualizing log2-fold changes in cytokines in serum (A) and BAL (B). Both individual animals and cytokines are arranged by hierarchical clustering, represented by dendrograms on top and to the right of the heatmaps, respectively. Surviving animals are marked in white in the top bar above the heatmap, while dead animals are in black. In the bar below this, the 4-hour group of animals are marked in red, the 12-hour group in blue, and the control group in green. Fold change values are centered and scaled for each cytokine by first subtracting the mean fold change from each value and then dividing by the standard deviation of that cytokine.

Fig. S3 Principal component analysis of cytokine fold changes following RT challenge. Scatter plots of first 2 principal components of log2-transformed cytokine fold changes in A-C) BAL and (D-F) serum from all animals included in the study, calculated used singular value decomposition. Each dot represents an individual monkey; red for those in the 4-hour group, black for the 12-hour group, and green for the control group. 95% confidence ellipses of the group means are represented in the color of the group they correspond to. Eigenvectors for (C) BAL and (F) and analyses are colored to show the percent contribution of each variable to the principal components.

A.

B.

Figure 2

Figure 3

A. Serum

	Control		4 hr intervention					12 hr intervention				
	DH27	JP56	LB98	IC70	IL40	LJ80	EM61	KP48	KR49	LB26	DR61	JM39
EGF	1.53	0.83	-0.17	0.73	0.28	-0.83	0.64	0.27	0.64	-0.67	0.79	1.42
Eotaxin	0.62	0.91	-0.01	0.48	0.31	0.28	0.40	-0.29	0.01	-0.05	-0.17	0.61
FGF-basic	0.38	1.49	0	0.33	-0.27	-0.32	0.40	0.25	0.28	-0.35	0.29	0.73
G-CSF	2.58	4.92	-0.07	0.05	-0.93	-0.22	-0.33	-0.14	-0.21	-0.33	-0.10	0.21
GM-CSF	1.58	2.16	-0.21	-0.05	0.08	-0.07	0.07	-0.03	0.04	-0.18	-0.07	0.26
HGF	0.33	0.68	-0.24	-0.10	-1.07	-0.22	0.04	-0.17	-0.16	-0.04	-0.15	0.23
IFN-g	0.53	0.74	-0.20	-0.15	-1.15	-0.22	0	-0.13	-0.21	0	-0.24	0.57
IL-1b	0.99	1.56	-0.15	-0.01	-0.33	-0.06	0.15	-0.06	-0.03	-0.07	-0.38	0.44
IL-1RA	7.17	8.47	0.09	0.04	0.12	-0.47	1.73	1.68	0.84	1.55	1.35	3.74
IL-2	1.04	1.64	-0.08	-0.10	0.10	0	0.07	-0.07	0.07	-0.03	-0.21	0.13
IL-4	0.23	0.35	-0.15	-0.08	-0.58	-0.12	-0.11	0	-0.07	0.08	-0.14	-0.07
IL-5	1.04	1.95	-0.21	0.26	-0.39	-0.97	0.18	-0.06	0.15	-0.20	-0.38	0.06
IL-6	8.65	9.64	1.41	1.94	2.23	2.14	2.61	2.30	0.64	2.49	2.24	4.18
IL-8	-0.85	-1.52	-1.74	-1.87	-1.83	-2.01	-0.84	-2.28	-2.80	-1.78	-3.56	-1.90
IL-10	0.60	0.85	-0.06	-0.03	-0.77	-0.03	0.06	-0.01	0.10	0.05	0	0.12
IL-12	-0.44	0.05	-0.13	-0.27	-0.82	-0.43	0.07	-0.32	-0.25	-0.10	-0.22	-0.55
IL-15	0.57	0.96	-0.18	0.05	-0.61	-0.16	0.27	-0.08	0.02	-0.45	-0.11	0.21
IL-17	0.05	-0.08	-0.10	-0.12	-0.23	-0.40	0.04	-0.21	-0.18	-0.42	-0.18	0.14
I-TAC	-0.32	-0.17	-0.57	-0.74	0.50	-0.56	0.50	-0.19	-0.81	-0.71	-0.36	0.18
MCP-1	3.05	4.02	-0.16	0.73	0.72	0.28	1.68	-0.22	0.30	0.26	-0.31	1.41
MDC	-0.25	0.19	-0.16	-0.28	-0.04	0.17	0.33	0	0.23	0.08	-0.42	-0.36
MIF	2.00	1.74	0.27	1.09	0.22	-0.67	0.78	0.62	0.67	-0.48	1.39	0.86
MIG	0.31	0.38	-0.26	-0.18	-0.94	-0.07	-0.04	0.10	-0.18	-0.17	-0.20	0.25
MIP-1a	0.71	0.77	-0.16	0	-1.04	0.13	0.10	-0.15	0.17	0	-0.18	0.24
MIP-1b	0.44	0.82	-0.17	-0.04	-0.77	-0.20	0.08	-0.10	-0.10	0.14	-0.06	0.42
RANTES	0.04	0.08	-0.02	-0.02	-0.02	0.01	0	-0.04	-0.01	-0.06	-0.07	-0.01
TNF-a	0.30	0.58	-0.19	-0.02	-1.34	-0.11	0	-0.32	0.11	-0.21	-0.12	0.15
VEGF	3.18	3.71	0.32	0.31	1.16	0.14	0.67	0.73	0.20	0.67	0.44	1.57
IP-10	0.33	0.71	-0.10	-0.23	-1.12	-0.07	-0.10	-0.15	0	-0.17	-0.03	0.13

B. BAL

	Control		4 hr intervention				12 hr intervention				
	DH27	LB98	IC70	IL40	LJ80	EM61	KP48	KR49	LB26	DR61	JM39
EGF	0.10	0.07	0.12	0.26	-0.10	0.05	-0.02	0.19	0.22	0.22	0.29
Eotaxin	2.25	0.98	2.84	1.97	1.73	1.09	3.78	2.00	3.69	2.74	4.23
FGF-basic	0.62	0.41	0.13	0.14	-0.08	0.03	0.62	0.49	0.45	0.56	0.62
G-CSF	1.21	0.17	0.20	0.29	0.21	0.20	0.19	0.25	1.21	0.38	3.74
GM-CSF	1.02	0.26	0.50	0.64	0.80	0.57	1.00	1.23	1.29	1.11	1.65
HGF	0.21	0.09	0	0.29	0.13	0.05	0.42	0.37	0.52	0.23	1.02
IFN-g	-0.17	0.09	0.13	0	0.11	0	0.47	0.28	0.35	0.28	0.47
IL-1b	1.76	0.84	0.57	2.16	0.82	1.29	1.04	1.32	1.52	2.76	2.36
IL-1RA	2.28	0.10	0.98	2.09	1.10	1.52	2.25	2.81	1.75	3.72	3.41
IL-2	0.92	0.03	0.21	0.40	0.16	0.16	0.90	0.90	1.04	0.88	1.39
IL-4	0.22	0.15	0.08	0.03	0.12	0	0.35	0.14	0.22	0.29	0.51
IL-5	1.07	0.25	0.58	0.63	0.28	0.29	1.12	1.56	1.46	1.43	1.90
IL-6	6.25	3.79	5.20	5.40	4.16	4.53	6.79	5.27	7.74	7.96	7.79
IL-8	0.85	0.62	0.58	0.77	1.01	0.50	0.55	1.03	1.04	1.08	1.03
IL-10	0.08	0.04	0.08	0.06	0.03	0.01	0.27	0.73	0.34	0.23	0.53
IL-12	0.04	0.04	0.11	0.15	0.17	0.12	0.28	0.24	0.26	0.25	0.55
IL-15	-0.21	-0.02	0.09	0.16	-0.02	0.03	0.54	0.11	0.26	0.20	0.32
IL-17	0.20	0.07	0.17	0.58	0.16	0.03	0.74	0.69	1.52	0.22	0.69
I-TAC	-0.81	1.84	0.93	1.65	0.64	1.63	0.33	1.04	0.75	1.78	1.42
MCP-1	-0.89	2.69	2.02	3.02	3.78	3.51	3.83	3.71	5.19	1.97	4.65
MDC	-0.06	1.73	1.62	2.01	1.34	2.70	1.30	1.51	0.56	2.72	3.62
MIF	1.47	1.10	0.73	0.51	0.73	0.46	-0.48	1.11	0.07	-0.48	-0.86
MIG	0	0	-0.13	0.06	0.06	0.13	0.14	-0.21	0.16	0.29	0.20
MIP-1a	0.51	0.47	0.45	1.78	1.09	0.45	0.42	0.48	2.45	0.78	0.91
MIP-1b	0.64	0.06	0.13	0.27	0.13	0.07	0.72	0.49	1.42	0.46	0.81
RANTES	0.37	-0.22	-0.31	-0.39	-0.18	-0.09	-0.76	-0.08	-0.63	-0.58	-0.40
TNF-a	0.49	0.28	0.36	0.77	0.51	0.47	0.65	0.75	0.74	0.77	1.37
VEGF	-1.36	-0.24	0.50	-0.29	0.27	0.14	0.62	-0.30	1.34	2.02	0.39
IP-10	0.10	0.08	0.14	0.17	0.08	0.15	0.74	0.24	0.68	0.43	0.76

Supplemental Figures

Figure S1

A.

B.

C.

D.

Serum

BAL