The *Aquilegia* genome reveals a hybrid origin of core eudicots

Gökçe Aköz[1,2] and Magnus Nordborg[1,*]

[1]Gregor Mendel Institute, Austrian Academy of Sciences, Vienna Biocenter, Vienna, Austria

[2]Vienna Graduate School of Population Genetics, Vienna, Austria

[*]magnus.nordborg@gmi.oeaw.ac.at

Abstract

Background: Whole-genome duplications (WGD) have dominated the evolutionary history of plants. One consequence of WGD is a dramatic restructuring of the genome as it undergoes diploidization, a process under which deletions and rearrangements of various sizes scramble the genetic material, leading to a repacking of the genome and eventual return to diploidy. Here, we investigate the history of WGD in the columbine genus *Aquilegia*, a basal eudicot, and use it to illuminate the origins of the core eudicots.

Results: Within-genome synteny confirms that columbines are ancient tetraploids, and comparison with the grape genome reveals that this tetraploidy is shared with the core eudicots. Thus, the ancient *gamma* hexaploidy found in all core eudicots must have involved a two-step process: first tetraploidy in the ancestry of all eudicots, then hexaploidy in the ancestry of core eudicots. Furthermore, the precise pattern of synteny sharing suggests that the latter involved allopolyploidization, and that core eudicots thus have a hybrid origin.

Conclusions: Novel analyses of synteny sharing together with the well-preserved structure of the columbine genome reveal that the *gamma* hexaploidy at the root of core eudicots is likely a result of hybridization between a tetraploid and a diploid species.

Background

Whole-genome duplication (WGD) is common in the evolutionary history of plants [reviewed in 1,2]. All flowering plants are descended from a polyploid ancestor, which in turn shows evidence of an even older WGD shared by all seed plants [3]. These repeated cycles of polyploidy dramatically restructure plant genomes. Presumably driven by the "diploidization" process, whereby genomes are returned to an effectively diploid state, chromosomes are scrambled via fusions and fissions, lose both repetitive and genic sequences, or are lost entirely [4–11]. Intriguingly, gene loss after WGD is non-random: not only is there a bias against the retention of certain genes [12,13], but also against the retention of one of the WGD-derived paralog chromosomes [6,9,14–16].

We investigated the history of WGDs in the columbine genus *Aquilegia* for two reasons. The first is related to its phylogenetic position: as a basal eudicot, columbines are members of the very earliest diverging branch of the eudicots [17,18]. This matters

because our understanding of eudicot karyotype evolution is limited to the heavily sampled core eudicots. Using the recently published *Aquilegia coerulea* genome [19], we are able to address key questions about the history of polyploidization in all eudicots. Second, we traced the origins of the columbine chromosomes with a particular focus on chromosome 4, which, compared to the rest of the genome: harbors more genetic polymorphism, has a higher transposable element density, has a lower gene density and reduced gene expression, shows less population structure worldwide, appears more permeable to gene flow, and carries the rDNA clusters [19].

Results

Within genome synteny confirms columbine paleotetraploidy

Ancient WGDs have been commonly inferred from the distribution of divergences between gene duplicates. The simultaneous generation of gene duplicates via WGD is expected to produce a peak in the age distribution relative to the background age distribution of single gene duplicates [20–22]. Such a spike of ancient gene birth was the first evidence of paleotetraploidy in columbines [23], and was later supported by gene count-based modelling [24].

Given an assembled genome, a more direct method to infer ancient polyploidy is to look for regions with conserved gene order [25,26]. Such conservation (a.k.a., synteny) decreases over time due to gene loss and rearrangements, but will still be detectable unless the extent of change is extreme. We detected a total of 121 synteny block pairs harboring at least five paralogous gene pairs within the columbine genome. The distribution of these blocks across the seven columbine chromosomes indicates pairwise synteny between large genomic regions (Fig. 1). This 1:1 relationship suggests a single round of WGD in columbine, and is further supported by similar levels of divergence between synteny pairs (Figs. S1 and S2).

Fig. 1: Intragenomic synteny blocks in the columbine genome. Pairs of synteny blocks are denoted as uniquely colored small rectangles. Larger rectangles of the same color outline large regions of synteny. Arrows under the synteny blocks show the orientation of the alignment between collinear genes. Grey dots highlight BLAST hits of a 329 bp centromeric repeat monomer [19,27].

Columbines share ancient tetraploidy with core eudicots

All sequenced core eudicots appear to share a triplicate genome structure due to paleohexaploidy postdating the separation of monocots and eudicots [9,28–32, and Supplementary Note 5 in 33]. The tetraploidy in columbines, a basal eudicot, might be independent of this ancient "*gamma*" hexaploidy (Scenarios 1 and 2 in Fig. 2) or might be a remnant of a WGD at the base of all eudicots, which formed the first step of the *gamma* hexaploidy in core eudicots (Scenario 3 in Fig.2).

83
84
Fig. 2: Three scenarios for the relationship between columbine tetraploidy
and core eudicot "*gamma*" hexaploidy. The *gamma* hexaploidy is a two-step
process: a single round of WGD creates tetraploids (4n) whose unreduced gametes then
fuse with diploid gametes (+2n). **Scenario 1**: *gamma* hexaploidy precedes the split
between columbine and core eudicots, with the former undergoing an additional
tetraploidy. **Scenario 2**: Both *gamma* hexaploidy and columbine tetraploidy occur
after the split between columbines and core eudicots. **Scenario 3**: Columbine
tetraploidy is derived from the ancient tetraploidy that was the first step of the process
leading to *gamma* hexaploidy.

94
We used the grape (*Vitis vinifera*) genome as a representative of the core eudicots to
distinguish between the three scenarios in Fig. 2. Grape has experienced the least
number of chromosomal rearrangements post-*gamma* and thus strongly resembles the
ancestral pre-hexaploid genome [34]. Given the ploidy level of columbine under each
scenario, we can predict the synteny relationship between the homologous
chromosomes of grape and columbine, which is simply the ratio of haploid chromosome
set in grape to that of in columbine. If tetraploidy in columbines is lineage-specific and
superimposed on the *gamma* hexaploidy (Scenario 1), we would expect to find a 3:6
ratio of grape and columbine synteny blocks. Instead, we observe a 3:2 relationship
(Figs. 3 and S3) as expected under Scenarios 2 or 3. A similar 3:2 pattern is found in
comparisons between grape and sacred lotus [35]. This strongly suggests that basal

eudicots do not share the triplicate genome structure of core eudicots, ruling out
Scenario 1.
108

109
110
111 **Fig. 3: Synteny between the homologous regions of columbine and grape.**
112 The results are shown here only for the columbine chromosomes 1,2 and the grape
113 chromosomes 6,8 and 13 but reflect the overall synteny relationship of 3:2 between
114 grape:columbine chromosomes (see Fig. S3 for the genome-wide synteny). This pattern
115 argues against Scenario 1, but is consistent with either Scenario 2 or Scenario 3.
116
To distinguish between the two remaining scenarios, we compared the divergence at
synonymous sites (Ks) between columbine paralogs, grape paralogs and columbine-
grape homologs. In agreement with the analysis of Jiao et al. [36], the Ks distribution
for grape paralogs shows two major peaks, as expected under the two-step model for
gamma hexaploidy (Fig. 4). However, columbine paralogs and columbine-grape
homologs each show a single peak of divergence — and the peaks overlap each other and
the "older" divergence peak of grape paralogs. This suggests that columbine tetraploidy
is derived from the tetraploidy that eventually led to *gamma* hexaploidy in core eudicots
(Scenario 3).
126

Fig. 4: The distribution of the median Ks across syntenic regions. Synteny blocks are identified within columbine (col_paralogs), between columbine and grape (col_grape) and within grape (grape_paralogs). Note that only the putative WGD-derived blocks (median Ks=1-2) are kept in columbine (Fig. S2).

To further explore the hypothesis of a shared WGD by all eudicots, we focused on the gene order similarity between the homologous regions of columbine and grape. If columbine and grape have descended from a common tetraploid ancestor, they should have inherited diploidization-driven differential gene order on the paralogous chromosomes of the ancestor (Fig. S5). As a result, we expect to see the alternative paralogous gene orders to be uniquely shared between two different pairs of columbine and grape chromosomes. To detect this, we first searched for at least three consecutive genes aligning between a pair of columbine and grape chromosomes and then looked at the distribution of these genes on all the columbine and grape chromosomes. This way of reconstructing chromosomes clearly shows that each of the paralogous chromosome pairs in columbine has a match to a single grape chromosome, with respect to its gene order (Fig. S6). This result was corroborated by a second approach where we quantify

the similarity between columbine and grape chromosomes. When we performed a
pairwise alignment between each sliding window of genes on a columbine chromosome
and all the genes on a grape homolog, we again see that each member of columbine
paralogs gets the best hit to a single grape chromosome (Figs. 5 and S7-9). Reshuffling
genes on grape chromosomes further indicates that this pattern of clustering is highly
unlikely to be produced by chance alone (p=0-0.05).

153
154
Fig. 5: Examples of gene order similarity between the homologous regions
of columbine and grape. For successive windows of genes within a given columbine
chromosomal region, the best alignment score with respect to each of the three grape
chromosomes harboring homologous regions, is given. For example, columbine
chromosomes 1 and 2 share a paralogous region homologous to grape chromosomes 6,
8, and 13 (Figs. 7 and S3). The chromosome 1 region **(top panel)** appears to be most
closely related to grape chromosome 6, whereas its paralogous counterpart on
chromosome 2 **(bottom panel)** appears to be most closely related to grape

chromosome 13. See Fig. S6 for the correspondence between gene orders and scores, which peak towards the end of each columbine region. Note that the results presented here are shown for a window size of 12 genes but remain significant for all the window sizes tested (p=0-0.05).

An eudicot-wide WGD is further supported by the observation that a chromosomal fusion, presumably experienced by the common tetraploid ancestor, is still detectable in the genomes of columbine and grape despite their separation of around 125 million years [37]. The first hint comes from the composition of the chromosomes: columbine chromosome 5 and grape chromosome 7 share the two chromosomal origins (Fig. 7). If these fused chromosomes were created by a single fusion event in the common tetraploid ancestor of eudicots, they should match each other with respect to gene order on each of the two homologous portions ("orange" and "green" portions in Fig. 6). This is what we see: columbine chromosome 5 and grape chromosome 7 cluster together with respect to their gene order on the "orange" portion (Fig. S7). For the "green" portion, columbine chromosome 5 matches grape chromosome 4 (Fig. S8), which used to be fused to grape chromosome 7 [38]. Additional support for shared ancestral fusion comes from the cacao (*Theobroma cacao*) genome [39]. The first chromosome of cacao does not only show a similar pattern of chromosomal ancestry [38,39], but also shares the gene order exclusively with the grape chromosomes 4 and 7 on the corresponding homologous portions (Fig. S10). In summary, the columbine fusion clusters with that of grape, which, in turn, clusters with that of cacao, strongly favoring a common origin of the fusion between "orange" and "green" ancestral chromosomes (Fig. 7).

Fig. 6: Schematic of predicted synteny patterns in the case of shared ancestral fusion. Two ancestral chromosomes (orange and green rectangles, with genes depicted as numbers) undergo WGD. Paralogous chromosome pairs diverge as a part of the diploidization process. A fusion joins one version of the "orange" chromosome ('1, 3, 5') with one version of the "green" chromosome ('7, 10, 8'). If this took place in the common tetraploid ancestor of eudicots, the fused chromosomes in columbine and grape should also carry these versions on their "orange" and "green" portions. In the hypothetical example here, diploidization precedes the fusion event but may well happen afterwards with no effect on the predicted synteny patterns.

Fig. 7: Tracing the genome reshuffling in columbine following tetraploidy.
Grape chromosomes **(bottom right)** are colored by within-genome synteny. Seven distinct colors represent the haploid set of seven ancestral chromosomes before the eudicot-wide WGD. Each color is shared by three grape chromosomes reflecting the triplicate genome structure of core eudicots. The only exception is the "green" chromosome which is shared by four grape chromosomes due to a fission event [38]. Columbine chromosomes **(bottom left)** are colored by their synteny to grape chromosomes. Each color is generally shared by two chromosomes, reflecting columbine paleotetraploidy. As few as 7 fusions and a single fission are enough to explain the current structure of the columbine genome. Of these 7 fusions, 5 are between different chromosomes while 2 are between WGD-derived paralogous chromosomes. Columbine chromosomes 3 and 7 are examples of the latter (Figs. 1 and S4). Note that chromosome 5 of columbine and chromosome 7 of grape (*) both have the colors "orange" and "green" (cf. Fig. 6).

The core eudicots have a hybrid origin

Our inference of shared tetraploidy between basal and core eudicots makes use of the signals presumably generated by diplodization (Figs. 6 and S5). However, hybridization of unreduced gametes from two divergent diploid genomes, "allotetraploidy", would also lead to gene order-based clustering between two different pairs of grape and columbine chromosomes (Figs. S11-12). In this case, the alternative paralogous gene orders of the tetraploid ancestor reflect the gene orders on progenitor chromosomes. Thus, the clustering pattern does not depend on whether the eudicot tetraploid genome evolved via "auto-" or "allopolyploidy". The same is not true for the second of the process leading to hexaploidy. Only allohexaploidy would lead to one of the three paralogous grape chromosomes being an "outlier" to the two grape-columbine pairing (Figs. S11-12) — which is what we see in our data (light blue lines in Figs. 5 and S7-9).

If our interpretation is correct and all core eudicots have a hybrid origin, the pattern of gene order-based clustering should be conserved. That is, we should be able to identify the hexaploidy-derived "outlier" chromosomes in other core eudicot genomes as well. To check this expectation, we again used the cacao genome, one of the most conserved genomes after grape [9,39]. Pairwise alignment between the homologous regions of columbine and cacao confirms our expectation: each member of columbine paralogs pairs up with a single cacao chromosome, leaving one of the cacao paralogs as an outlier (Figs. S13-14). Furthermore, as shown in Fig. 8 (see also Fig. S10), the cacao regions putatively derived from tetraploidy and hexaploidy, respectively, show a very clear one-to-one match to those in grape (detected in the grape-columbine comparison). As expected, the putatively orthologous pairs of cacao and grape regions show similar levels of synteny conservation with their paralogous counterparts, with the "outlier" regions being the most divergent [38]. Thus, the cacao genome provides an independent line of evidence for a hybrid origin, and highlights the key role of the columbine genome in unravelling the history of the eudicot genome.

Fig. 8: The shared history of chromosomes in columbine, grape and cacao.
Gene order-based clustering results **(left panel)** are summarized here for the chromosomes harboring the "orange" and "green" homologous portions. The former corresponds to 5, 7, 14 in grape and 1, 4, 5 in cacao. The latter corresponds to 3, 4+7 (products of a fission), 18 in grape and 1, 2, 8 in cacao. In columbine, the "orange" portions are on chromosomes 2 and 5 while the "green" portions are on chromosomes 6 and 5, each pair of which being denoted as *colum A* and *colum B*, respectively. Both grape- and cacao-columbine pairing distinguish tetraploidy-derived regions (blue and purple rectangles) from hybridization-derived ones (light blue rectangles), defining the orthologous sets of regions across the three eudicot genomes **(right panel)**. The conservation of gene order exclusively between the putatively orthologous regions of grape and cacao (black arrows, Fig. S10) further strengthens our columbine-based inference of orthology.

Current columbine chromosomes have mostly been generated via fusions

It is widely accepted that genome shuffling post-WGD has shaped the present-day karyotypes of all plant genomes [34]. Nevertheless, the extent of genome shuffling as a part of the "re-diploidization" process seems to vary widely: only 3 chromosomal rearrangements post-*gamma* are enough to explain the current structure of the grape genome (Fig. 7) while almost 150 chromosomal rearrangements were necessary for the sunflower genome to reach its current karyotype after several rounds of WGD [11]. To check where columbine falls in this spectrum, we identified chromosomal rearrangements likely to have happened after the tetraploidy shared by all eudicots: if the pre-WGD ancestral eudicot karyotype had a haploid number of 7 chromosomes [28], only seven columbine-specific fusions and a single fission are enough to explain the

reduction in columbine chromosome number from n=13 to n=7 after the ancestral
fusion event (Fig. 7). These rearrangements involve all the chromosomes in columbine
apart from chromosomes 4 and 6, the former of which paradoxically shows the greatest
erosion of synteny with grape chromosomes (Figs. 7 and S3). Given all the evidence
suggesting a "decaying" nature of columbine chromosome 4 [19], we repeated the
analysis of grape-columbine synteny detection with relaxed parameter settings. We did
this by decreasing the minimum number of aligned gene pairs within a block (from 5 to
3) and increasing the maximum genic distance between matches (from 20 to 30). This
allowed us to extend the synteny blocks towards more proximal regions (Fig. S15).
Further zooming into the synteny relationship between grape chromosomes that are
homologous to columbine chromosome 4 confirmed that there is no evidence of a fusion
event (Fig. 9).
282

283
284   **Fig. 9: Synteny between columbine chromosome 4 and grape chromosomes**
285   **12 and 19.** Much smaller grape chromosomes look like the compact versions of
286   columbine chromosome 4. Note that this result is generated with the most relaxed
287   parameter combination in Fig. S15, but holds true for a less relaxed combination of
288   parameters as well (Fig. S16).

289
The lack of a fusion event on columbine chromosome 6 might explain the fact that it is
the smallest chromosome of columbine (Fig. 7). However, chromosome 4 is comparable
in size to the remaining chromosomes, all of which are products of ancient fusion
events. The observations that chromosome 4 has a higher proportion of genes in tandem
duplicates (0.37 versus genome-wide mean of 0.22) and a greater extent of intra-
chromosomal synteny (indicative of segmental duplications) (Fig. S17) suggest that
chromosome 4 has reached a comparable size partly due to numerous tandem and
segmental duplications and partly due to an expansion of repetitive DNA [19]. These
results reinforce the idea that chromosome 4 has followed a distinct evolutionary path
from the rest of genome.

Fusion-dominated genome shuffling [34] is not the only facet of diploidization [40].
Following WGDs, gene duplicates get lost and this happens in a non-random manner.
Genes involved in connected molecular functions like kinases, transcription factors and
ribosomal proteins are retained in pairs [41–45] potentially due to dosage-related
constraints [46]: losing or duplicating some, but not all of these dosage-sensitive genes
might upset the stoichiometric relationship between their protein products [47–49].
Consistent with this dosage balance hypothesis, columbine genes potentially retained
post-WGD (1302 genes across 76 syntenic regions; Supplementary Data 1) are enriched
for the GO categories "structural constituent of ribosome", "transcription factor
activity", "translation" (p<0.001) and "protein tyrosine kinase activity" (p<0.01).
Tandemly duplicated genes (n=6972), on the other hand, are depleted for the GO
categories "structural constituent of ribosome", and "translation" (p=10^{-17}), reflecting
the role of dosage-related purifying selection.
313

Discussion

The evolutionary history of plants is characterized by multiple WGDs and columbine is
no exception. The alignment between chromosomal regions in a 1:1 ratio (Fig. 1)
confirms a single round of WGD in columbine [23,24,50]. Furthermore, we demonstrate
that this tetraploidy is shared with all eudicots, refuting a lineage-specific
polyploidization in columbine [23,50], in favor of an eudicot-wide WGD [26,35,36,51].
Unlike previous attempts based on genetic distance, our approach simply relies on gene
order conservation. It also takes advantage of the well-preserved genome structure of
columbines: free from recent WGDs, the columbine genome carries only the traces of
the ancient tetraploidy.
324

This approach also helps us shed light on the nature of the *gamma* hexaploidy found in
all core eudicots [9,28–32, and Supplementary Note 5 in 33]. WGDs have often been
discussed as if they were "events", ignoring the process by which they originated. We
show here that core eudicot hexaploidy is the result of two processes: an ancient
tetraploidization shared by all eudicots, followed by allopolyploidization leading to core
eudicots. In other words, all core eudicots have a hybrid origin. An allohexaploid origin
has indeed been previously suggested by Murat et al. [9], who identified the three
subgenomes of grape using differential patterns of gene loss on "dominant" versus
"sensitive" subgenomes. Their classification assumes that the most recently added set of
paralogous chromosomes will be "dominant", because they have spent a shorter amount
of time in the polyploid genome and thus experienced fewer gene losses. Contrary to
this, our results suggest that the most recently added grape chromosomes
(chromosomes 3, 8, 9 and 14) largely corresponds to the "sensitive" grape chromosomes
identified by Murat et al. [9]. Instead, we argue that the extensive gene loss in the
"youngest" subgenome reflects its divergence from the other two subgenomes at the
time of hexaploid formation, perhaps similar to the situation in the allotetraploid

Arabidopsis suecica, which is a hybrid between the more ancestral (n=8) genome of *A.*
arenosa, and the heavily reduced (n=5) genome of *A. thaliana* [52]. Another example is
hexaploid wheat, which is a hybrid between tetraploid emmer wheat and a wild diploid
grass, *Aegilops tauschii* [53 and references therein].
345

Conclusions

Our findings help us understand the hybrid structure of core eudicot genomes and will
hopefully encourage larger scale analyses to understand what hybridization has meant
for core eudicots — a group which comprises more than 70% of all living flowering
plants [54]. What are the hybridization-coupled changes that has led to the current
patterns of gene expression, methylation, transposable element density/distribution?
All these questions call for additional genomes from basal eudicots which — as this
study illustrates — have great values as outgroup to the core eudicots.
354

Materials and Methods

Synteny detection

We performed all genes (CDS)-against-all genes (CDS) BLAST for the latest version of
Aquilegia coerulea reference genome (v3.1) using SynMap tool [29] in the online CoGe
portal [55]. We also looked at the synteny within *Vitis vinifera* (v12) and between *V.*
vinifera and *A. coerulea* using both default and more relaxed parameter combinations
in DAGChainer. We filtered the raw output files for both within grape and grape-to-
columbine synteny. For the former, we only kept the blocks that are syntenic between
the polyploidy-derived paralogous chromosomes of grape as identified by Jaillon et al.
[28]. For the latter, we required that a given columbine chromosome is overall syntenic
to all the three paralogous chromosomes of grape. So, for a given pair of columbine and
grape chromosomes, we only kept the blocks if the columbine chromosome also matches
to the other members of paralogous grape chromosomes.
368

The raw output files can be regenerated at the CoGe portal [55] using the id numbers
provided below for each species (Availability of data and material) and changing the
default parameter combination in DAGChainer (D:A=20:5) when needed. D and A
specify the maximum genic distance between two matches and the minimum number of
aligned gene pairs, respectively, to form a collinear syntenic block.

Estimating the divergence between synteny block pairs

We used Ks (the number of synonymous substitutions per synonymous site) values
provided for each duplicate gene pair by the CoGe portal [55]. We estimated the median

Ks of all gene duplicates in a syntenic block after filtering duplicates with Ks>10 due to
saturation effect [56].

Quantifying gene order similarity

We first detected three consecutive genes aligning between a pair of columbine and
grape chromosomes harboring homologous regions (D:A=0:3). We particularly chose
three genes since it is the most stringent value we could use to detect homologous
synteny blocks; we detected almost nothing when we required 4 consecutive genes
(D:A=0:4). We then looked at the distribution of these genes on a given pair of
columbine and grape chromosomes and also on their paralogous counterparts
(D:A=0:1). Once we had the gene order for each chromosome, we assigned a unique
word to each synteny block and the genes forming the block to be able to use the text
alignment provided by the R package *align_local* [57]. Having each chromosome
represented by a sentence, we quantified the gene ("word") similarity as such: for an
initial N number of words on a columbine chromosome (N=window size), we did a
pairwise alignment between these N words and a grape chromosome (match=4, gap=-
1). We repeated the same analysis with the inverted order of N words and picked the
maximum alignment score. We repeated these steps by sliding the window by one word
and keeping the N constant to get a distribution of scores as in Fig. 5. We used different
N values ranging from 4 to 15. Note that we excluded columbine chromosomes 3 and 4
from this analysis since both have a complex history of lineage-specific chromosomal
reshuffling events: fusions and a fission gave rise to columbine chromosome 3 (Figs. 1
and 7) while duplications have shaped the current structure of columbine chromosome
4 (Fig. S17).

We applied the same stringent criteria (D:A=0:3) to detect the homologous regions
between grape and cacao (*Theobroma cacao*, v1). The same criteria led to very few
homologous regions between columbine and cacao. So, we relaxed the parameters for
the synteny detection between these two genomes (D:A=0:2) and quantified the gene
order similarity with greater window sizes (N=20, 30, 35, 40 and 50). Note that we
focused on the triplicated regions distributed across 3 different cacao chromosomes
(Figs. 8, S13-14), which are rather unaffected by lineage-specific shuffling [38].

Statistical testing of gene order similarity

Given the gene order similarity between the two different pairs of columbine and grape
chromosomes harboring homologous regions, we performed permutation tests to
estimate the probability of observing such a clustering just by chance. To do so, we first
combined all the grape genes and sampled the same number of genes ("words") as we
observe to reconstruct each of the paralogous grape chromosome. We repeated the
quantification step as above to get a permuted distribution of alignment score between a

pair of columbine and grape chromosomes. We used Wilcoxon rank sum test (W-statistic) to quantify the shift in the distribution of alignment scores between one of the members of columbine paralogous chromosomes and its best grape hit when combined with the alignment scores between the same columbine chromosome and other grape chromosomes. We repeated the same analysis for the other member of columbine paralogous chromosomes as well. Having these *observed* W-statistics, we counted the number of cases (out of 100) where the permuted distributions generate W-statistics as high as or higher than the observed ones. Note that for columbine chromosome 7, whose structure has been greatly shaped by the fusion of WGD-derived paralog chromosomes (Fig. S4), we created two paralogous chromosomes using the observed distribution of alignment scores (Fig. S9). Columbine chromosome 7 matches best to grape chromosome 11 for the first 14 "words" and to grape chromosome 4 for most of the remaining "words", which define the putative boundaries of columbine paralogous chromosomes before the fusion event. We ran permutation tests for the columbine-cacao pairing as well (Figs. S13-14).

GO enrichment analysis

We used gene annotations provided by JGI [19] to test the null hypothesis that the property for a gene to be retained post-WGD and to belong to a given GO category are independent. We created a 2x2 contingency table as shown below and applied Fisher's exact test for each GO category independently. We repeated the same analysis for tandem gene duplicates as identified by SynMap [29,55]; this time testing the null hypothesis that the property for a gene to be tandemly duplicated and to belong to a given GO category are independent. We excluded genes on scaffolds and reported enriched/depleted categories if they remain significant (p <0.05) after multiple test correction (fdr).

Table 1: 2x2 contingency table obtained by classifying genes into 2 categorical variables. The letters denote the number of genes for a given category (e.g. "a" denotes the number of retained genes annotated with the tested GO category).

	GO	not-GO	SUM
retained	a	b	a+b*
not-retained	c	d	c+d
SUM	a+c	b+d	N=total number of genes =29550 (across 7 chromosomes)

---

*equal to 1302 and 6972 for candidate WGD-derived paralogs and tandem gene duplicates, respectively.

List of abbreviations

WGD: whole genome duplication; Ks: the number of synonymous substitutions per synonymous site; GO: Gene Ontology.

Declarations

Availability of data and material

The columbine, grape and cacao genomes are available at the CoGE portal for the synteny analyses with the id numbers 28620, 19990 and 25287, respectively [55].

Competing interests

The authors declare no competing interests.

Funding

G.A. was supported by the Vienna Graduate School of Population Genetics (Austrian Science Fund, FWF: DK W1225-B20).

Authors' contributions

G.A. performed all analyses. G.A. and M.N. wrote the manuscript.

Acknowledgements

We thank Robin Burns and Claus Vogl for their comments on the manuscript; Daniel Gómez Sánchez and Benjamin Jaegle for the fruitful discussions.

References

1. Wendel JF. The wondrous cycles of polyploidy in plants. Am J Bot. 2015;102:1753–6.

2. Van de Peer Y, Mizrachi E, Marchal K. The evolutionary significance of polyploidy. Nat Rev Genet. 2017;18:411–24.

3. Jiao Y, Wickett NJ, Ayyampalayam S, Chanderbali AS, Landherr L, Ralph PE, et al. Ancestral polyploidy in seed plants and angiosperms. Nature. 2011;473:97–100.

4. Leitch IJ, Bennett MD. Genome downsizing in polyploid plants. Biol J Linn Soc Lond. Oxford University Press; 2004;82:651–63.

5. Comai L. The advantages and disadvantages of being polyploid. Nat Rev Genet. 2005;6:836–46.

6. Thomas BC, Pedersen B, Freeling M. Following tetraploidy in an Arabidopsis
ancestor, genes were removed preferentially from one homeolog leaving clusters
enriched in dose-sensitive genes. Genome Res. 2006;16:934–46.

7. Otto SP. The evolutionary consequences of polyploidy. Cell. 2007;131:452–62.

8. Renny-Byfield S, Chester M, Kovařík A, Le Comber SC, Grandbastien M-A, Deloger
M, et al. Next generation sequencing reveals genome downsizing in allotetraploid
Nicotiana tabacum, predominantly through the elimination of paternally derived
repetitive DNAs. Mol Biol Evol. 2011;28:2843–54.

9. Murat F, Zhang R, Guizard S, Gavranović H, Flores R, Steinbach D, et al. Karyotype
and gene order evolution from reconstructed extinct ancestors highlight contrasts in
genome plasticity of modern rosid crops. Genome Biol Evol. 2015;7:735–49.

10. Murat F, Armero A, Pont C, Klopp C, Salse J. Reconstructing the genome of the most
recent common ancestor of flowering plants. Nat Genet. 2017;49:490–6.

11. Badouin H, Gouzy J, Grassa CJ, Murat F, Staton SE, Cottret L, et al. The sunflower
genome provides insights into oil metabolism, flowering and Asterid evolution. Nature.
2017;546:148–52.

12. Blanc G, Wolfe KH. Functional divergence of duplicated genes formed by polyploidy
during Arabidopsis evolution. Plant Cell. 2004;16:1679–91.

13. De Smet R, Adams KL, Vandepoele K, Van Montagu MCE, Maere S, Van de Peer Y.
Convergent gene loss following gene and genome duplications creates single-copy
families in flowering plants. Proc Natl Acad Sci U S A. 2013;110:2898–903.

14. Woodhouse MR, Schnable JC, Pedersen BS, Lyons E, Lisch D, Subramaniam S, et al.
Following Tetraploidy in Maize, a Short Deletion Mechanism Removed Genes
Preferentially from One of the Two Homeologs. PLoS Biol. Public Library of Science;
2010;8:e1000409.

15. Schnable JC, Springer NM, Freeling M. Differentiation of the maize subgenomes by
genome dominance and both ancient and ongoing gene loss. Proc Natl Acad Sci U S A.
2011;108:4069–74.

16. Tang H, Woodhouse MR, Cheng F, Schnable JC, Pedersen BS, Conant G, et al.
Altered patterns of fractionation and exon deletions in Brassica rapa support a two-step
model of paleohexaploidy. Genetics. 2012;190:1563–74.

17. Hodges SA, Kramer EM. Columbines. Curr Biol. 2007;17:R992–4.

18. Soltis DE, Smith SA, Cellinese N, Wurdack KJ, Tank DC, Brockington SF, et al.
Angiosperm phylogeny: 17 genes, 640 taxa. Am J Bot. 2011;98:704–30.

19. Filiault D, Ballerini E, Mandakova T, Akoz G, Derieg N. The Aquilegia genome:
adaptive radiation and an extraordinarily polymorphic chromosome with a unique

history. bioRxiv [Internet]. biorxiv.org; 2018; Available from:
https://www.biorxiv.org/content/early/2018/02/12/264101.abstract

20. Lynch M, Conery JS. The evolutionary fate and consequences of duplicate genes.
Science. 2000;290:1151–5.

21. Blanc G, Wolfe KH. Widespread paleopolyploidy in model plant species inferred
from age distributions of duplicate genes. Plant Cell. 2004;16:1667–78.

22. Cui L, Wall PK, Leebens-Mack JH, Lindsay BG, Soltis DE, Doyle JJ, et al.
Widespread genome duplications throughout the history of flowering plants. Genome
Res. 2006;16:738–49.

23. Vanneste K, Baele G, Maere S, Van de Peer Y. Analysis of 41 plant genomes supports
a wave of successful genome duplications in association with the Cretaceous–Paleogene
boundary. Genome Res. 2014;24:1334–47.

24. Tiley GP, Ané C, Burleigh JG. Evaluating and Characterizing Ancient Whole-
Genome Duplications in Plants with Gene Count Data. Genome Biol Evol. 2016;8:1023–
37.

25. Doyle JJ, Egan AN. Dating the origins of polyploidy events. New Phytol.
2010;186:73–85.

26. Jiao Y, Paterson AH. Polyploidy-associated genome modifications during land plant
evolution. Philos Trans R Soc Lond B Biol Sci [Internet]. 2014;369. Available from:
http://dx.doi.org/10.1098/rstb.2013.0355

27. Melters DP, Bradnam KR, Young HA, Telis N, May MR, Ruby JG, et al. Comparative
analysis of tandem repeats from hundreds of species reveals unique insights into
centromere evolution. Genome Biol. 2013;14:R10.

28. Jaillon O, Aury J-M, Noel B, Policriti A, Clepet C, Casagrande A, et al. The grapevine
genome sequence suggests ancestral hexaploidization in major angiosperm phyla.
Nature. 2007;449:463–7.

29. Lyons E, Pedersen B, Kane J, Freeling M. The Value of Nonmodel Genomes and an
Example Using SynMap Within CoGe to Dissect the Hexaploidy that Predates the
Rosids. Trop Plant Biol. 2008;1:181–90.

30. Potato Genome Sequencing Consortium, Xu X, Pan S, Cheng S, Zhang B, Mu D, et
al. Genome sequence and analysis of the tuber crop potato. Nature. 2011;475:189–95.

31. Truco MJ, Ashrafi H, Kozik A, van Leeuwen H, Bowers J, Wo SRC, et al. An Ultra-
High-Density, Transcript-Based, Genetic Map of Lettuce. G3 . 2013;3:617–31.

32. Denoeud F, Carretero-Paulet L, Dereeper A, Droc G, Guyot R, Pietrella M, et al. The
coffee genome provides insight into the convergent evolution of caffeine biosynthesis.
Science. 2014;345:1181–4.

33. Bombarely A, Moser M, Amrad A, Bao M, Bapaume L, Barry CS, et al. Insight into
the evolution of the Solanaceae from the parental genomes of Petunia hybrida. Nat
Plants. 2016;2:16074.

34. Salse J. Ancestors of modern plant crops. Curr Opin Plant Biol. 2016;30:134–42.

35. Ming R, VanBuren R, Liu Y, Yang M, Han Y, Li L-T, et al. Genome of the long-living
sacred lotus (Nelumbo nucifera Gaertn.). Genome Biol. 2013;14:R41.

36. Jiao Y, Leebens-Mack J, Ayyampalayam S, Bowers JE, McKain MR, McNeal J, et al.
A genome triplication associated with early diversification of the core eudicots. Genome
Biol. 2012;13:R3.

37. Zeng L, Zhang Q, Sun R, Kong H, Zhang N, Ma H. Resolution of deep angiosperm
phylogeny using conserved nuclear genes and estimates of early divergence times. Nat
Commun. 2014;5:4956.

38. Zheng C, Chen E, Albert VA, Lyons E, Sankoff D. Ancient eudicot hexaploidy meets
ancestral eurosid gene order. BMC Genomics. 2013;14 Suppl 7:S3.

39. Argout X, Salse J, Aury J-M, Guiltinan MJ, Droc G, Gouzy J, et al. The genome of
Theobroma cacao. Nat Genet. 2011;43:101–8.

40. Soltis DE, Visger CJ, Marchant DB, Soltis PS. Polyploidy: Pitfalls and paths to a
paradigm. Am J Bot. 2016;103:1146–66.

41. Seoighe C, Wolfe KH. Yeast genome evolution in the post-genome era. Curr Opin
Microbiol. 1999;2:548–54.

42. Tian C-G, Xiong Y-Q, Liu T-Y, Sun S-H, Chen L-B, Chen M-S. Evidence for an
ancient whole-genome duplication event in rice and other cereals. Yi Chuan Xue Bao.
2005;32:519–27.

43. Aury J-M, Jaillon O, Duret L, Noel B, Jubin C, Porcel BM, et al. Global trends of
whole-genome duplications revealed by the ciliate Paramecium tetraurelia. Nature.
2006;444:171–8.

44. Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, et al. The B73 maize
genome: complexity, diversity, and dynamics. Science. 2009;326:1112–5.

45. Rodgers-Melnick E, Mane SP, Dharmawardhana P, Slavov GT, Crasta OR, Strauss
SH, et al. Contrasting patterns of evolution following whole genome versus tandem
duplication events in Populus. Genome Res. 2012;22:95–105.

46. Birchler JA, Bhadra U, Bhadra MP, Auger DL. Dosage-dependent gene regulation in
multicellular eukaryotes: implications for dosage compensation, aneuploid syndromes,
and quantitative traits. Dev Biol. 2001;234:275–88.

47. Papp B, Pál C, Hurst LD. Dosage sensitivity and the evolution of gene families in

yeast. Nature. 2003;424:194–7.

48. Freeling M, Thomas BC. Gene-balanced duplications, like tetraploidy, provide
predictable drive to increase morphological complexity. Genome Res. 2006;16:805–14.

49. Birchler JA, Veitia RA. Gene balance hypothesis: connecting issues of dosage
sensitivity across biological disciplines. Proc Natl Acad Sci USA. 2012;109:14746–53.

50. Guo L, Winzer T, Yang X, Li Y, Ning Z, He Z, et al. The opium poppy genome and
morphinan production. Science [Internet]. 2018; Available from:
http://dx.doi.org/10.1126/science.aat4096

51. Malacarne G, Perazzolli M, Cestaro A, Sterck L, Fontana P, Van de Peer Y, et al.
Deconstruction of the (paleo)polyploid grapevine genome based on the analysis of
transposition events involving NBS resistance genes. PLoS One. 2012;7:e29762.

52. O'Kane SL, Schaal BA, Al-Shehbaz IA. The Origins of Arabidopsis suecica
(Brassicaceae) as Indicated by Nuclear rDNA Sequences. Syst Bot. American Society of
Plant Taxonomists; 1996;21:559–66.

53. Matsuoka Y. Evolution of polyploid triticum wheats under cultivation: the role of
domestication, natural hybridization and allopolyploid speciation in their
diversification. Plant Cell Physiol. 2011;52:750–64.

54. Friis EM, Pedersen KR, Crane PR. The emergence of core eudicots: new floral
evidence from the earliest Late Cretaceous. Proc Biol Sci [Internet]. 2016;283. Available
from: http://dx.doi.org/10.1098/rspb.2016.1325

55. Lyons E, Freeling M. How to usefully compare homologous plant genes and
chromosomes as DNA sequences. Plant J. 2008;53:661–73.

56. Blanc G, Hokamp K, Wolfe KH. A recent polyploidy superimposed on older large-
scale duplications in the Arabidopsis genome. Genome Res. 2003;13:137–44.

57. Smith DA, Cordell R, Dillon EM. Infectious texts: Modeling text reuse in nineteenth-
century newspapers. 2013 IEEE International Conference on Big Data. 2013. p. 86–94.

616