A systematic review of the relation between species traits and extinction risk

Filipe Chichorro [1*], Aino Juslén[2] and Pedro Cardoso[1]

[1]LiBRE – Laboratory for Integrative Biodiversity Research, Finnish Museum of Natural History, University

of Helsinki, PO Box 17 (Pohjoinen Rautatiekatu 13), 00014 Helsinki, Finland

[2]Finnish Museum of Natural History, University of Helsinki, PO Box 17 (Pohjoinen Rautatiekatu 13), 00014

Helsinki, Finland

*Address for correspondence (Tel: +358 469 508 202; E-mail: filipe.chichorrodecarvalho@helsinki.fi)

ABSTRACT

Biodiversity is shrinking rapidly, and despite our efforts only a small part of it has been assessed for

extinction risk. Identifying the traits that make species vulnerable might help us to predict the outcome for

those less known. We gathered information on the relations of traits to extinction risk from 173 publications,

across all taxa, spatial scales and biogeographical regions, in what we think is the most comprehensive

compilation to date. The Palaearctic is the most represented biogeographical realm in extinction risk studies,

but the other realms are also well sampled for all vertebrate groups. Other taxa are less well represented and

there are gaps for some biogeographical realms. Many traits have been suggested to be good predictors of

extinction risk. Among these, our meta-analyses were successful in identifying two as potentially useful in

assessing risk for the lesser-known species: regardless of the taxon, species with small range and habitat

breadth – two of the three classic dimensions of rarity – are more vulnerable to extinction. On the other hand,

body size (the most studied trait) did not present a consistently positive or negative response. Large species

can be more, or less, threatened than their smaller counterparts, depending on the taxon and spatial setting. In

line with recent research, we hypothesize that the relationship between body size and extinction risk is

shaped by different aspects, namely body size is a proxy for different phenomena depending on the

taxonomic group, and there might be relevant interactions between body size and the specific threat that a

species faces. With implications across all traits, a particular focus in future studies should be given to the

interaction between traits and threats. Furthermore, intraspecific variability of traits should be part of future

studies whenever data is available. We also propose two complementary paths to better understand extinction

risk dynamics: (i) the simulation of virtual species and settings for theoretical insights, and (ii) the use of

comprehensive and comparable empirical data representative of all taxa and regions under a unified study.

Keywords: biological traits, functional diversity, IUCN, machine learning, meta-analysis, threat status.

Contents

I. Introduction

The last annual checklist of the Catalogue of Life reports 1.7 million (eukaryote) species described on this planet (Roskov *et al.*, 2017). Describing the remaining estimated 7 million might still take 1,200 years if the rates of description, average number of new species described per taxonomist's career, and average cost to describe species remain constant (Mora *et al.*, 2011). This latency in describing species is put into a typical crisis context if, as predicted, 3,000 species fall into extinction every year (Wilson, 2003; González-Oreja, 2008), many of them undescribed.

The International Union for Conservation of Nature (IUCN) compiles and keeps updated a database with assessments of risk of extinction for species. As of August 2018, 26 197 (28%) of all 93 577 species in this list were either Critically Endangered, Endangered, or Vulnerable to extinction and 14 616 (16%) were Data Deficient (IUCN, 2018). Yet, species in the IUCN database mostly comprise well-known taxa (e.g. 67% of vertebrates have been assessed versus 0.8% of insects (IUCN, 2018)), and it will probably take decades until a reasonable proportion of many taxa, such as most invertebrates, are assessed (Cardoso *et al.*, 2011a, b, 2012). Increasing the number of species in the database to the point where we have an unbiased picture of extinction risk across all organisms during the next years seems highly unlikely, as is the Barometer of Life goal of assessing 160 000 species by 2020 (Stuart *et al.*, 2010). Moreover, extinction is taxonomically selective (e.g. 63% of cycads are assessed as threatened versus 'only' about 13% of bird species (IUCN, 2018)). The current proportions of endangered species might not represent the greater picture of species diversity. Therefore, alternative ways of predicting the risk of extinction of species are urgently needed.

Understanding which biological/ecological traits of species make them more vulnerable could help us predict their extinction risk and make species protection and conservation planning more efficient. This approach is not new. Some comparative studies can be traced back to the 19th century (see McKinney, 1997, for a thorough historical perspective), and since the beginning of the new millenium many new comparative studies have arisen on the topic, as well as discussions over their usefulness (Fisher & Owens, 2004; Cardillo & Meijaard, 2012; Murray *et al.*, 2014; Verde Arregoitia, 2016). Many traits have been tested across

hundreds of publications. Body size, for example, was found to be positively correlated with extinction risk

across multiple taxa (Seibold *et al.*, 2015; Terzopoulou *et al.*, 2015; Verde Arregoitia, 2016), either through

direct effects (e.g. larger species require more resources) or as a proxy for other traits (e.g. larger species

have slower life cycles and therefore respond more slowly to change). Range size and population density,

even after taking into account that they are often used to quantify extinction risk, have also been extensively

tested and found to be relevant, at least for mammals (Purvis et al., 2000a; González-Suárez, Gómez &

Revilla, 2013; Bland *et al.*, 2015; Verde Arregoitia, 2016). Traits related to exposure to human pressures

have also been relevant in predicting threats to species (Cardillo, 2003), and recently Murray *et al.* (2014)

have called for more studies explicitly incorporating threats and the interplay between traits and threats into

the analyses. The inclusion of threat information in predicting extinction risk has indeed proved to increase

the explanatory power of models (Murray *et al.*, 2014), and in some cases the same trait can bolster

extinction risk or prevent it, depending on the threat (González-Suárez *et al.*, 2013).

Most of the studies to date have focused on mammals (e.g. Purvis *et al.*, 2000a; Cardillo *et al.*, 2008;

González-Suárez *et al.*, 2013) and other vertebrates (e.g. Owens & Bennett, 2000; Luiz *et al.*, 2016), with

relatively few on plants (e.g. Sodhi *et al.*, 2008; Powney *et al.*, 2014; Stefanaki *et al.*, 2015) and invertebrates

(e.g. Sullivan *et al.*, 2000; Koh, Sodhi & Brook, 2004; Arbetman *et al.*, 2017). Each study was made on

different spatial settings and scales, testing different traits (often according to availability of data), and

employed different methods and response variables. While this is necessary and valuable information,

making sense of the plethora of contrasting results is difficult, and perceiving general trends and trying to

cover current gaps and bias are urgent. In this work we attempt to answer the following questions through a

comprehensive bibliography search, data exploration and meta-analysis:

● Which traits have been studied more often?

● Which traits have been suggested as predictors of extinction risk?

● How generalizable are the past results?

● What directions should research take in the future to overcome the shortage and bias of data?

II. Material and Methods

In this review we undertook two sequential analyses of studies that examined the relation between traits of species and their estimated extinction risk: first, an exploratory analysis of the literature, to know which traits have been more studied and which were found to be most relevant, and second, multiple meta-analyses, in which we tried to use comparable data from a subset of the studies to understand and quantify trends across studies and taxa from all published data and to see whether any general conclusions can be made from existing literature.

(1) Bibliography selection

We were aiming to retrieve a list of all publications that have explicitly performed comparative studies of biological/ecological traits and extinction risk/decline of species and to identify which traits, extrinsic factors and taxa were used in each analysis and at which spatial scale. We searched in the Web of Science using the keywords 'trait*' and 'extinct*' in October 2016 and obtained a list of 2,616 manuscripts. To consider a given paper as relevant to our study, all of the following conditions had to be met:

- more than five species were involved in the study;
- for each species there was information on at least one biological trait;
- for each species there was some kind of quantification of its extinction risk;
- there was a statistical model linking the species traits (explanatory variables) to the extinction risk (response variables), assigning scores to each trait involved (not necessarily significance).

We considered as extinction risk measurements any of the following:

- recent (anthropogenic) extinctions versus extant species;
- any variable (continuous, ordinal, categorical or binary) directly indicating relative extinction risk, whether it was based on the IUCN Red List categories or not;
- population trend data, or a proxy of population trend data, in time;
- any other variables that indicated decline of species over time and/or risk of extinction.

On a first reading of a small part of the studies, we realized that most were not relevant to the study, as either

traits or extinction were marginal to our questions (some were not even related to biology) or some of the

minimum requirements were missing. Analysing 2,616 manuscripts would have been too time consuming

when a relatively small fraction would be relevant. For this reason, we used an automated process to sort out

relevant publications.

We used a random forest algorithm trained to identify relevant from non-relevant publications. We first

created two sets of 35 abstracts+titles, one with relevant and one with non-relevant papers (70 in total). As

words with common roots (e.g. extinction and extinct) should be quantified together, we used the R software

(R Core Team, 2018) and package SnowballC (Bouchet-Valat, 2014) to stem these words (e.g. all words in

the 'extinct' family fall into the same stem). We also excluded hyphenation, punctuation and stopwords

(Rajamaran & Ullman, 2011) using R packages SnowballC and tm (Feinerer *& Hornik,* 2008). Then, all the

words and corresponding word counts of each abstract were used as variables in training the random forest

model.

Decision trees were built with packages rpart (Therneau & Atkinson, 2015) and rpart.plot (Milborrow,

2018). The training of the random forest algorithm was run with 50 000 trees and 45 variables tried at each

step (the suggested number of variables is the square root of the number of words, Liaw & Wiener, 2002)

with R package randomForest (Liaw & Wiener, 2002). To validate the model, we assembled another set of

abstracts (n = 76, test set) and calculated the sensitivity, specificity and the true skill statistic (TSS) of the

model based on predictions on the test set. After training and validating the algorithm, we used it to classify

the remaining 2,000+ abstracts. In order to further test for false negatives, i.e. papers incorrectly labelled as

non-relevant leading to the loss of important studies, we randomly picked 100 abstracts labelled as non-

relevant and checked them.

To update the full set of relevant papers, we performed another search using the same keywords in June

2018, retrieving new publications and obtaining 682 further studies from the Web of Knowledge. The trained

algorithm was again used to filter potentially relevant publications from these. We finally added to the

relevant set all papers of which we had prior knowledge by other means.

167   (2) Statistical analysis

168   (a) Data collection

169   We assembled information on each comparative statistical test employed in each article. For each of these

170   tests, we extracted information on the following (see also Appendix S1, S2):

- **Taxonomic group**: mammals, birds, reptiles, amphibians, fishes, insects, molluscs, other invertebrates, plants and fungi.

- **Geographical realm**: Afrotropics, Antarctic, Australasia, Indo-Malaya, Nearctic, Neotropics, Oceania and Palaearctic (Olson *et al.*, 2001).

- **Traits**: continuous, ordinal, categorical or binary – units, the number of observations (usually species), and whether there was a significant response to extinction risk for that test. We grouped traits with similar biological meaning into the same unified trait (e.g. body length and body mass into 'body size'). Henceforth, trait refers to these unified traits. Finally, to identify meaningful groups of traits and facilitate analysis, we clustered these traits into meaningful classes (see next section).

- **Statistical methodology**: generalized linear models and linear models (including parametric tests of the generalized linear models' and linear models' families, Pearson correlations, G-tests, analysis of variance and analysis of covariance, chi-square statistics), generalized linear mixed models (GLMMs) and linear mixed models (parametric tests of the generalized mixed linear models' and linear mixed models' families), decision tree based (including random forests, boosted or non-boosted classification and regression trees), generalized estimating equations, phylogenetic comparative methods (PGCMs; which include phylogenetic independent contrasts and phylogenetic generalized least squares), non-parametric (Kruskal–Wallis, Spearman rank correlations, etc.), and other (including discriminant analysis, path analysis, randomization tests, skewness tests, structural equation modelling).

● **Phylogenetic control**: we distinguished absolute no control of phylogeny from at least some control

of phylogeny (via using phylogenetic trees or a taxonomical higher group as a controlling or

covariable in the analyses).

● **Proxy of extinction used**: IUCN based (where the response variable – ordinal or binary – is based

upon IUCN assessments of species), other red lists (based on non-IUCN red lists), temporal trends

(in which the trend can be the difference between population numbers, or geographical range size,

between two time periods, or the slope of the variation in the population numbers over time), and

everything else as others (recently extinct species versus extant species, extirpated versus non-

extirpated, etc.).

(b) Trait classes

We named nine natural groups (or categories) of traits (Table 1). All these categories describe a property of a

focal species, but they differ in the general type of such property:

● **Individual**: traits measurable at an individual level: morphological traits, traits related to

reproductive output and life-cycle speed.

● **Population**: traits that describe a property of a whole population. Such properties can relate to how

individuals of the populations of the focal species aggregate, whether they are social, or other traits

only measurable at the population level, like population density or abundance.

● **Community**: in this category, traits describe the interactions between the focal species and other

species in the community, whether this interaction is trophic or other, such as symbiosis,

commensalism or pollination.

● **Evolution**: including traits related to how successful a species might be in adapting to change

through evolution, as well as any other traits related to the evolutionary history of the species.

● **Space**: traits describing the current or past geographical distribution in space of the species or its

ability to disperse.

● **Time**: traits that describe how species use time at different scales, such as daily or seasonal cycles.

● **Environmental niche**: traits mainly describing species interactions with their biotic or abiotic niche,

such as habitat affinities and preferential climatic ranges.

● **Human influence**: traits that describe the interplay between species and humans, such as habitat loss

due to anthropogenic causes and hunting or other uses of the species.

● **Taxonomy**: taxonomic affinity or hierarchy of the species.

● **Other**: traits that do not fit into any of the previous groups.

(c) Exploratory analysis

In the exploratory analysis we first compared the number of studies across taxa and combinations of taxa

across biogeographical realms, proxy of extinction used, statistical methodology, and phylogeny. Next, we

compared the number of studies in which each trait was used, and calculated the percentage of significant

measurements of each trait. Statistical tests which did not assign significance levels to traits had to be

excluded from this step (e.g. most decision tree methodologies).

(d) Meta-analysis

To understand whether traits were positively or negatively related to extinction risk across the multiple

studies, we performed meta-analyses for each continuous trait. Meta-analyses are useful because they allow

the comparison of outcomes from different studies by converting the outcomes to effect sizes. The use of

Fisher's Z as the effect size has the advantage of allowing very diverse statistical methodologies into the

same effect size measurement. Effect sizes were obtained by transforming the statistics reported in the

manuscripts (F, z, X^2, t or r^2) into Pearson's product-moment correlation coefficients (r) by applying

equations (1) to (5) (Rosenthal, 1991) and then transforming r into Fisher's Z using equation (6) using R

package metafor (Viechtbauer, 2010):

$r = \dfrac{z}{\sqrt{N}}$ (1)

$r = \sqrt{\dfrac{t^2}{t^2 + df}}$ (2)

$r = \sqrt{\dfrac{F_{1,df}}{F_{1,df} + df}}$ (3)

$r = \sqrt{\dfrac{X_1^2}{N}}$ (4)

$r = \sqrt{r^2}$ (5)

$$Z = \frac{1}{2 \ln\frac{1+r}{1-r}} \qquad (6)$$

To ensure that the outcomes would be comparable, we restricted the analyses to univariate tests. We

followed two different methodologies to detect the overall effect size for each trait. We first performed a

traditional random-effects meta-analytic model for each trait. Random-effect meta-analytic models assume

that in addition to the sampling error within measurements, there is also another source of variation between

measurements due to different methodologies being used. However, traditional meta-analytic methods lack a

tool to control for the existence of multiple measurements within studies. Therefore, we also used linear

mixed models, since these allow for more flexibility by controlling multiple measurements within studies

(and hence non-independence of observations) through the use of random effects (see Prugh, 2009; Chaplin-

Kramer *et al.*, 2011). Fisher's Z was the response variable and was weighted by the inverse of the sample

sizes. Additionally, in the linear mixed models the response variable was tested against the intercept term

only, with random effects being taxonomic group and study.

255 III. Results

(1) Bibliography selection

The random forests classified 488 abstracts as relevant in the first literature search. The sensitivity,

specificity and TSS on the test data were respectively 0.895, 0.868 and 0.763. A number of word roots were

found to be of considerable importance in the abstract filtering, namely 'extinct', 'risk', 'speci' and 'vulner',

all with a decrease in the Gini index above 0.7 (Appendix S3, Fig. S1), even if the most parsimonious tree

used the words 'extinct', 'habitat' and 'caus' (Appendix S3, Fig. S2). The algorithm classified an additional

135 abstracts as relevant from the second literature search. After human filtering and the inclusion of a single

false negative, the number of relevant publications decreased from 623 to 136. To these we added 37 papers

found a priori to be relevant and not identified by the algorithm, for a total of 173 manuscripts fulfilling all

criteria and included in this study (Appendix S1).

(2) Exploratory analysis

(a) Studies

The number of publications relating traits to extinction risk has increased steadily (Fig. 1). Mammals and

birds have received the most attention over the years, followed by fishes, insects and plants. Most studies

were conducted in the Palaearctic region (Fig. 2), particularly for insects and plants. Of particular note,

amphibians, reptiles and mammals have been included in many studies focusing on the Australasian realm.

Oceania and especially the Antarctic were the least represented biogeographical realms, with intermediate

values in all the other regions.

Regarding the response variable, IUCN categories and temporal trends (including range size or population

size contractions over time) were the most used (Table 2). IUCN categories were used mainly in studies

assessing mammals and amphibians but much less so in insect studies. Temporal trends and other red-list

types were mostly used in bird and insect studies.

Most studies used methodologies that take into consideration at least some degree of non-independence of

species, namely GLMMs (by using taxonomic group as random effect) and PGCMs (Table 2). Among these,

PGCMs require phylogenetic trees, often not available for many taxa, and hence they were mostly used in

mammal studies for which a global tree was easy accessible (e.g. Bininda-Emonds *et al.*, 2007).

(b) Traits

The most commonly used trait categories were Individual, Space, and Environmental Niche (Fig. 3 and

Table 1). Similarly, the Individual, Space, and Environmental Niche categories had the highest number of

measurements, along with Human influence (Fig. 3).

Body size was by far the most studied trait both in the Individual category, as well as overall (Fig. 4),

followed by geographical range size. After these, several traits appeared in a similar number of studies:

habitat type, habitat breadth and fecundity (Fig. 4). The taxonomic group was used in some studies to control

for phylogeny and often to guarantee the comparability of results across taxa. The compound metric trait was

in most cases a variable resulting from the first axis of principal component analysis of traits and their

relationships, and therefore it contained a mix of traits from multiple groups.

Geographical range size was the trait with the greatest proportion of studies with not non-significant tests

(almost three quarters) (Fig. 4). Among the traits that were present in at least 10% of the studies, several

were not non-significant in at least 40% of the tests, half of them belonging to the Environmental Niche

category (temperature, (micro)habitat type, habitat breadth) but also to the Individual (body size), Space

(location) and Community (diet breadth) categories.

Even when used in at least 10% of studies, not all of these traits were studied across all taxa. Body size and

geographical range size were the only traits that were studied for all taxa (except for fungi, since the only

study focusing on fungi did not attribute significance levels to traits and thus this group was not included

here) and were significant in at least one test for each taxon. The other 10 traits occurred in at least 7 out of 9

taxa. They belong to the Environmental Niche (altitude, temperature, temperature range, habitat breadth,

habitat type, microhabitat type), Community (diet type), Space (location, motility), and Individual (offspring

number per reproductive event) categories.

Despite occurring in less than 10% of the studies, many traits have been found to be good predictors of

extinction risk for some taxa. A number of traits (see Appendix S4 for the significances of all tested traits

within taxa) were tested in at least three studies and were significant at least once, even if for single taxa

(torpor/hibernation and weaning age in mammals; duration of flight period in birds; temperature for breeding

in fishes; overwintering stage in insects; pollen vector, reproduction type, dispersal agent, and seed size in

plants).

(3) Meta-analysis

Geographical range size, habitat breadth, and body size were the only traits from which we could determine

effect sizes and sample sizes from at least 10 studies including univariate tests – the minimum number that

we considered reasonable in order to have confidence in the results of the meta-analyses. Effect sizes of

geographical range size and body size mostly originated from mammal and bird studies but also from studies

on reptiles, amphibians, fishes, insects, other invertebrates and plants (Appendix S5, Figs S14, S15). Effect

sizes of habitat breadth also originated mostly from mammal and bird studies, yet reptile, amphibian, other

invertebrates and plant studies were included (Appendix S5, Fig. S16).

The random-effect meta-analytic model identified significant negative effect sizes for geographical range

size and habitat breadth, and positive effect sizes for body size. However, these analyses failed to take into

consideration the disproportionate number of effect sizes within studies (some studies with more tests drove

the results) and the number of effect sizes per taxon (some taxa drove the results) in all three analyses

(Appendix S5, Figs S14–S16). Due to the much better control of such bias allowed by the linear mixed

model analyses, we henceforth focus on this.

For geographical range size and habitat breadth, the overall effect size was consistently and significantly

negative across taxa and studies (Table 3, Appendix S5, Figs S14–15). Contrasting with the classic meta-

analysis, the linear mixed model revealed an overall effect size not different from zero for body size

(Table 3). Effect sizes of body size were either positive or negative (Appendix S5, Fig. S16), and while there

was some tendency in mammals and birds for the effect sizes to be positive, in plants these were consistently

negative.

IV. Discussion

Our review clearly reveals the increasing importance of the study of species traits on the understanding and

prediction of extinction risk. The interest in the subject, even if relatively recent, is increasing exponentially

and shows no signs of slowing down. Yet, we also found that past studies were biased in scope in terms of

taxa, with vertebrates having the largest share, and spatial setting, with the Palaearctic dominating across

taxa, although Australasia is much studied for mammals, reptiles and amphibians. Such biases should be

mostly due to a large body of accumulated knowledge of these taxa and regions, to which a predominance of

researchers in these areas continue to contribute. The special interest in the Australasian mammals may be

due to an ongoing debate on the role of body size in extinction risk in this particular region (Verde

Arregoitia, 2016).

Despite being, to our knowledge, the largest review of the relation between traits and extinction risk to date,

we are aware that this contribution might not include all relevant studies in this field. Using less restrictive

rules for inclusion of studies would help to retrieve more publications (like the ones used in Verde

Arregoitia, 2016, and Murray *et al.*, 2014) but would probably lead to a higher bias towards some better

studied taxa (vertebrates). The proportion of non-vertebrate studies included in our study is larger than that

of Murray *et al.* (2014), even though the bias is inevitable in any comprehensive study on this subject.

Publication bias may also occur due to studies with non-significant terms being often less reported in the

literature (Gurevitch *et al.*, 2018), which might cause changes in the magnitude of effect sizes. We are,

however, confident that this review is thorough and as unbiased as possible with current data.

Herein we suggest a new way of grouping traits that works across taxa and is useful to better visualize trends

when many traits are studied. Previous attempts of grouping traits by Verde Arregoitia (2016) followed

functional groups: traits related to reproduction, morphology, ecology, behaviour, among others. Yet, this

categorization was hard to do across taxa, as the same trait might reveal different phenomena of interest

according to each study and in particular each taxon. That is the case, for example, with body size, which in

mammals and invertebrates is often associated with reproductive speed (Purvis *et al.*, 2000a; Cardillo, 2003;

Kotze & O'Hara, 2003; Bland, 2017) but with competitive ability and fitness in plants (Falster & Westoby,

2003; Walker & Preston, 2006; Laanisto *et al.*, 2015). Some traits in the Spatial category would fit equally

well under Individual traits, but in those publications from where they were extracted they were used as

proxy of dispersal ability (e.g. wing length and wing aspect ratio for bats, Safi & Kerth, 2004). Our trait

classes, thus, are not functional classes in the typical sense, but rather follow a more complex set of rules

related to how traits are measured and what they represent for the chance of survival of the species.

(1) Relation between traits and extinction risk

Geographical range size was the best predictor of extinction risk overall. This is not surprising, since small

geographical range is one of the values used in IUCN assessments (criteria B, D2), and these assessments are

the measure of extinction risk in many studies, which might lead to circular reasoning. However, even when

excluding from the analyses all species considered threatened due to small range, range was still strongly

associated with extinction risk (e.g. Purvis *et al.*, 2000a; Wang *et al.*, 2018). The mechanism behind this

relationship is not entirely understood (Purvis *et al.*, 2000a), but geographical range size captures ecological

and dispersal attributes of species that would require harder to obtain variables, such as overall abundance of

species, which are important in understanding extinction risk (Polaina, Revilla & González-Suárez, 2016).

The abundance–occupancy relationship is a well-known and thoroughly studied pattern, and many

mechanisms relate abundance to extinction risk (Gaston *et al.*, 2002). Likewise, range size is related to the

dispersal ability of species, determining the capacity of a species to occupy new areas to escape multiple

pressures, and with habitat breadth, revealing the ability of a species to cope with habitat change or loss.

Among the studies we included in our analysis, species with greater habitat breadth (habitat generalists) were

less prone to becoming extinct. Specialists have long been regarded as losers, and generalists as winners in

the current extinction crisis (McKinney & Lockwood, 1999), and ecological niche theory predicts the

worldwide decline in specialist species (Clavel, Julliard & Devictor, 2011). Whether this trend is due to the

intrinsic specificity of the species or to geographical range size is, however, not trivial to discern. In the

studies included in this review, most habitat breadth measures were derived from maps. Consequently, less

widespread species have less sampling points and therefore might show smaller habitat breadth due to

sampling bias alone (Burgman, 1989), when in reality we lack knowledge of whether they could thrive under

different habitats. Despite this, one of the studies we analysed controlled for range size by using the residuals

of a regression of range size by habitat breadth (Tingley, Hitchmough & Chapple, 2013) thus excluding the

effect of range size from the variable of interest, and they still found a negative relationship of this trait with

extinction risk in New Zealand lizard species. Furthermore, in a review on the topic, Slatyer, Hirst & Sexton

(2013) showed that even after taking into consideration sampling bias, the relationship between habitat

breadth and geographical range size remains significant across taxa. Irrespective of the putative causes or

relations to other variables, species with larger habitat breadth do have more chances to escape from multiple

pressure types and are consistently less threatened across taxa and spatial settings.

Although almost half of all measurements of body size were significant, the meta-analyses revealed that the

relationship between body size and extinction risk is not unidirectional. The interplay between body size and

threat type is one of the reasons for this phenomenon. Owens and Bennett (2000) showed that while larger

bird species were threatened by overexploitation, habitat loss or degradation selected for smaller species. The

same trend seems to apply at least to marine fishes (Olden, Hogan & Zanden, 2007) and mammals

(González-Suárez *et al.*, 2013), taxa that are often targeted directly and selectively by man. Independently of

threats, relationships may not even be linear. Threatened freshwater fishes are found both at the smaller and

larger spectrum of body sizes (Olden *et al.*, 2007), and the same bimodal relationship is found when pooling

all vertebrates together (Ripple *et al.*, 2017). In general, this bimodality seems to be derived from threat type,

with different threats leading to increasing extinction risk of different body size classes.

Other traits for which we could not perform a quantitative analysis have also shown to be useful in predicting

extinction risk under certain circumstances. Among the most studied traits in the Individual category are

traits related to speed of life cycle and reproductive output: fecundity, egg/neonatal size, and generation

length. These traits usually correlate with each other and with body size and longevity: bigger, longer-lived

species often have lower fecundity, bigger egg/neonatal sizes and longer generation lengths. Threat status

has been positively related to species with decreased fecundity (Cardillo, 2003; González-Suárez & Revilla,

2013; Böhm *et al.*, 2016; Ribeiro *et al.*, 2016; but see Pinsky & Byler, 2015; Sreekar *et al.*, 2015), larger

egg/neonatal sizes (Cardillo *et al.*, 2005; Jones, Fielding & Sullivan, 2006; González-Suárez & Revilla,

2013; Pinsky & Byler, 2015) and longer generation lengths (Anderson *et al.*, 2011; Hanna & Cardillo, 2013;

Jeppsson & Forslund, 2014; Comeros-Raynal *et al.*, 2016; but see Chessman, 2013). These traits reduce the

capability of species to compensate for high mortality rates (Pimm, Jones & Diamond, 1988; Purvis *et al.*,

2000a; González-Suárez *et al.*, 2013), even if their longer longevities should make them more apt to resist at

lower densities as they survive longer and might be able to overcome short-lived threats (Pimm *et al.*, 1988).

When species are directly persecuted by man, they are often bigger, with larger fecundity and egg/neonatal

sizes (Owens & Bennett, 2000; González-Suárez *et al.*, 2013), and longer longevity alone is not sufficient to

compensate for the high mortality. But when the threat is habitat loss, which indirectly increases mortality

and/or reduces natality rates, the trend is non-existent or even reversed (Owens & Bennett, 2000; González-

Suárez *et al.*, 2013), this being possibly due to the advantages of longer longevity alone.

Some traits related to Environmental Niche (besides habitat breadth) are commonly used across taxa and

much data are available about them. Among those, temperature (optimal temperature or temperature of the

species across its geographical range) and temperature range (range of temperatures tolerated by the species

or range of temperatures found across its geographical range) were often important predictors in the studies

that used them. Species with lower average temperatures within their range or narrower temperature ranges

are especially at risk due to an increasingly warmer climate (Jiguet *et al.*, 2010; Grenouillet & Comte, 2014;

Flousek *et al.*, 2015). In contrast, thriving under broad temperature ranges grants species the necessary

flexibility to deal with environmental or climatic change and hence lower their extinction risk (Chessman,

2013; Lootvoet, Philippon & Bessa-Gomes, 2015). When exceptions were found, these were due to the

correlation of temperature with the true causes of change in extinction risk. For example, tropical species of

amphibians living in narrower temperature ranges were found to be less at risk, as the higher tropical

temperatures possibly lead to higher fecundity and hence adaptability of the species (Cooper *et al.*, 2008).

Although the generality of the pattern could not be confirmed across studies, species depending on habitats

more affected by human influence are often more threatened (Stefanaki *et al.*, 2015; Powney *et al.*, 2014). In

Greece, flowering plants occurring in coastal or ruderal habitats, under pressure from urbanization and

tourism, were more at risk than flowering plants occurring on cliffs or high-mountain vegetation, the latter

habitats being under lower human pressure (Stefanaki *et al.*, 2015). British plant species with lower affinity

to nitrogen-rich soils are declining due to the intensification of agriculture, which has led to increased inputs

of nitrogen in otherwise nitrogen-poor soils (Powney *et al.*, 2014). Likewise, microhabitat type was a good

predictor of extinction risk in some studies due to some microhabitats becoming more rare with increased

human pressure (Parent & Schriml, 1995; Seibold *et al.*, 2015). A striking example is the decline of

saproxylic beetles that use dead wood of large diameter in Germany, as forest management options often

lead to the scarcity of such microhabitat (Seibold *et al.*, 2015). These observations give support to recent

claims that predicting extinction risk requires taking into account the threat type and using different variables

related to human use of species and habitats (Murray *et al.*, 2014).

Among all variables measured at the community level, both diet breadth and type were significant predictors

across several studies. The diet of a species can be important in leading to and predicting extinction in two

ways. Species restricted to fewer dietary options have shown to be more threatened (Basset *et al.*, 2015;

Jeppsson & Forslund, 2014; González-Suárez *et al.*, 2013; Matsuzaki *et al.*, 2011; Mattila *et al.*, 2008),

probably due to lower flexibility in switching to other options when the availability of their preferred food

source decreases (Purvis, Jones, & Mace, 2000). On the other hand, diet type, namely the trophic position of

a species, may be as important. Species at higher trophic levels tend to be more threatened (Chessman, 2013;

Bender *et al.*, 2013; Cardillo *et al.*, 2004; Purvis *et al.*, 2000a) and often provide early warnings of extinction

across the entire food chain (Cardoso *et al.*, 2010). The greater dependence on the densities and larger

foraging areas of prey species may lead to such a pattern (Carbone & Gittleman, 2002), with synergistic

effects between resource abundance and other factors contributing to the decline of, for example, predators.

With the density of wildlife dwindling everywhere (e.g. Hallmann *et al.*, 2017), and everything else being

equal, top predators are expected to be more at risk.

Within the Space category, location and migration distance were often tested and found to be important

predictors. Location as a trait is very idiosyncratic, and it just informs that the geographical location where

the species is found is a significant predictor of threat. As extinctions are not spatially random, and might

even interact with some traits (see, for example, Fritz, Bininda-Emonds & Purvis, 2009), this is to be

expected and relates more to the threat than to the species itself. Most studies on migration distance are of

birds. Long distance migrants tend to be more at risk, which could be either due to phenological mismatch

due to climate change (Amano & Yamaura, 2007; Jiguet *et al.*, 2010; Thaxter *et al.*, 2010; Flousek *et al.*,

2015), dependence on the good quality of at least two habitats or sites (Jiguet *et al.*, 2010; Flousek *et al.*,

2015), or to increased competition with resident species that, in temperate regions, survive through

increasingly less severe winters (Jiguet *et al.*, 2010; Amano & Yamaura, 2007).

Finally, there are also traits that were found to be significant but only studied for one or two taxa. These

include a wide array of morphological traits that are taxon specific. Some plant growth forms (e.g.

herbaceous, bush or tree) are more threatened than others. Perennial growth forms can sustain populations

through harsh times (Stefanaki *et al.*, 2015) but might be more affected by forest loss (Leão *et al.*, 2014).

Mammals going through a hibernating or torpor phase are less prone to becoming extinct, due to a greater

capacity to avoid harsher seasonal conditions (Liow *et al.*, 2009). The life stage in which an insect

overwinters (egg, larva, pupa or adult) influences vulnerability (e.g. Powney *et al.*, 2015; Jeppsson &

Forslund, 2014; Mattila *et al.*, 2009). At least for some studies with particular applied relevance, Cardillo &

Meijaard (2012) claim that 'researchers should adopt a somewhat "smaller picture" view by restricting the

geographical and taxonomic scope of comparative analyses, and aiming for clearer, more focused outcomes

on particular hypotheses'. We corroborate that restricting the studies in these two dimensions might prove

useful when the goal goes beyond understanding the general pattern and requires true predictive power for

species extinctions.

(2) Generalization

Given the inherent bias of past studies, any generalizations require critical consideration. Geographical range

and habitat breadth seem to be very well supported across taxa and regions, even if most past studies using

such traits were on vertebrates. Both are consistently negatively related to extinction risk and might be seen

as representing a single phenomenon: the range or rarity of a species in two different dimensions (area and

habitat). Species with larger ranges, be these spatial or biotic, have more chance of surviving in case of

diminishing availability of resources, and the risk of their populations or the entire species vanishing is

smaller. These traits can therefore be confidently used as predictors of extinction risk across taxa. Area and

habitat are in fact two of the three dimensions of rarity preconized by Rabinowitz (1981): geographical range

size, habitat breadth, and local abundance. The latter was seldom used probably due to the scarcity of

abundance data for most taxa (the Prestonian shortfall, Cardoso *et al.*, 2011b) but is certainly crucial to fully

understand the extinction phenomenon.

Body size, on the other hand, seems to be at least taxon dependent, probably because, as previously

mentioned, it represents different ways in which species interact with their environment and therefore how

they affect their risk of extinction (González-Suárez *et al.*, 2013; Ripple *et al.*, 2017). This trait is often

studied as a proxy for traits that may be very hard to measure or are very abstract. This implies that the

mechanisms behind the relation between body size and extinction risk are very much dependent on the traits

that body size proxies. Therefore, what it really represents for the way the species copes with different

threats must be treated with caution. If for animals it usually is related to resource availability, as larger

animals require more, often scarce, resources, being these space, food or other, for plants it represents

competitive ability, with larger plants being able to better exploit, for example, the sun, by growing taller and

overshadowing smaller species, or water and mineral resources found deeper underground. Many other traits

were either seldom studied or rarely found to be relevant. We must emphasize, however, that we still lack a

complete and unbiased picture of the relation between traits and extinction risk and that future studies could

and should provide insights much beyond what is possible now.

(3) Next steps

Given the scarcity and/or lack of comparability of data, we were not able to test the importance of the vast

majority of traits used in the past. Many of them might turn out to be crucial to the understanding of

extinction mechanisms and their prediction across multiple taxa. Also important to emphasize, Murray *et al.*

(2014) have already called for more papers that include human pressures as covariables in comparative

studies of extinction risk. Here we further reinforce this need. Furthermore, and somewhat surprisingly given

its perceived and demonstrated importance, not many studies included intraspecific variability of any trait.

Higher intraspecific variability has shown to predispose species to adapt to changing conditions (González-

Suárez & Revilla, 2013); however, morphological, physiological, genetic or behavioural variability have

seldom been studied, probably due to a lack of data. Here we also emphasize the importance of including

intraspecific variability in its different dimensions in future studies whenever data is available.

We propose two complementary paths for the future that will allow research in this area to be advanced

further and eventually allow prediction of the extinction risk of species based on their traits: (1) simulations

of virtual settings, threats and species using agent-based modelling, and (2) exhaustive sampling of unbiased

data across taxa and regions.

Computer simulations, in the form of agent-based models (ABMs, Grimm *et al.*, 2010), offer a framework in

which species, maps and interactions can be designed from scratch and which therefore can help us untangle

the often complicated relationships between traits, threats and the extinction risk of species. Out of simple

rules between interacting agents often emerge patterns not implicitly incorporated into the system (Grimm *et*

al., 2010). This is one of the main advantages of ABMs. By defining simple rules on how agents are created,

get energy, reproduce and die, we can test how changing different aspects related to a functional trait, in

interaction with a landscape, where threats are also modelled, will influence the outcome. Two approaches,

with different objectives, are possible. In one approach, the objective is to understand the mechanisms

behind extinction risk. In this approach, we simplify the system and simulate only the minimum required

complexity to understand the underlying mechanisms under certain conditions. Such a procedure is not very

common in biological sciences, but theoretical results are an important part of science and have proven

crucial to moving science forward in some fields, such as physics (Goldstein, 2018). In a second approach,

the objective is to use the models for prediction and therefore model complexity must be as much as is

needed for robust and realistic predictions. These two approaches are complementary and much needed to

complement the set of tools available to conservation biologists.

A strong predictor of extinction risk is important in a model when the amount of variation it explains is high

even after introducing other traits into the analysis. However, because we analysed very diverse studies,

focusing on multiple taxa and spatial settings, it was impossible to compare the performance of each trait

while taking into consideration other traits, because no two studies compared exactly the same traits in a

multivariable setting. The main idea behind this approach is to perform a single comparative study including

many different taxa, in which species selection, trait data collection, and statistical analysis are performed in

a congruent way thus allowing full comparability of taxa and traits.

V. Conclusions

(1) Most studies investigating drivers of extinction risk have focused on vertebrates, particularly mammals and birds, due to larger amounts of knowledge accumulated for these taxa. Likewise, many studies were restricted to certain regions, namely the Palaearctic, with other regions remaining understudied.

(2) Species with smaller ranges are more at risk of extinction irrespective of the taxon or geographical setting. The underlying mechanism is not entirely understood, yet species that occur in smaller ranges may have a particular set of traits, not captured in other commonly studied traits, that predispose them to elevated extinction risk such as reduced numbers and less dispersal ability.

(3) Species with narrow habitat breadth (habitat specialists) are also consistently more vulnerable to extinction across taxa and regions. When estimating the habitat breadth of a species, one may, however, find a spurious relationship between habitat breadth and extinction risk, due to the former being often correlated with geographical range size. Nonetheless, habitat breadth has been correlated with extinction risk, independently of geographical range. Like geographical range size, habitat breadth is a good predictor for less known species.

(4) Body size, the most studied trait, is an important predictor of risk. However, bigger species may be more, or less, threatened than smaller species. The underlying mechanism is rather more complicated than for the two previous traits, which is explained both by body size being (i) a proxy for very contrasting, often abstract, and hard to quantify traits (e.g. reproductive output in animals, competitive ability in plants), and (ii) very dependent on the threats to each species.

(5) The relationship between the vast majority of species traits, threats and extinction risk remains elusive. Many traits were found to be important across studies but have seldom been studied or are relevant for only some taxa.

(6) Human pressure and other threats should be increasingly used as covariates in comparative studies of extinction risk. Furthermore, intraspecific variability of traits should be part of future studies whenever data is available.

570   (7) We propose two contrasting approaches for future work: simulations with strong theoretical

571   background to understand the mechanisms behind the relation of traits to extinction risk, and

572   performing a single comparative study, across taxa, in order to ensure comparability of results.

VI. Acknowledgements

We would like to thank Cathryn Primrose-Matiasen for correcting the English language, and Sini Seppälä, Jon Rikberg and Fernando Urbano-Tenorio for their suggestions in defining and naming trait classes. We would also like to thank Cathryn Primrose-Mathisen who provided professional English language assistance during the preparation of this article. She was not responsible for reviewing the final version. F.C. was funded by Kone Foundation.

VII. References

References marked with asterisk have been used in the analyses

*ABELSON, E.S. (2016) Brain size is correlated with endangerment status in mammals. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20152772.

*AMANO, T. & YAMAURA, Y. (2007) Ecological and life-history traits related to range contractions among breeding birds in Japan. *Biological Conservation* **137**, 271–282.

*ANDERSON, S.C., FARMER, R.G., FERRETTI, F., HOUDE, A.L.S. & HUTCHINGS, J.A. (2011) Correlates of Vertebrate Extinction Risk in Canada. *BioScience* **61**, 538–549.

*ANGERMEIER, P.L. (1995) Ecological Attributes of Extinction-Prone Species: Loss of Freshwater Fishes of Virginia. *Conservation Biology* **9**, 143–158.

*ARBETMAN, M.P., GLEISER, G., MORALES, C.L., WILLIAMS, P. & AIZEN, M.A. (2017) Global decline of bumblebees is phylogenetically structured and inversely related to species range size and pathogen incidence. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20170204.

*ARBUCKLE, K. (2016) Chemical antipredator defence is linked to higher extinction risk. *Royal Society Open Science* **3**, 160681.

*BARSHEP, Y., ERNI, B., UNDERHILL, L.G. & ALTWEGG, R. (2017) Identifying ecological and life-history drivers of population dynamics of wetland birds in South Africa. *Global Ecology and Conservation* **12**, 96–107.

*BARTOMEUS, I., ASCHER, J.S., GIBBS, J., DANFORTH, B.N., WAGNER, D.L., HEDTKE, S.M. & WINFREE, R. (2013) Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proceedings of the National Academy of Sciences* **110**, 4656–4660.

*BARTONOVA, A., BENES, J. & KONVICKA, M. (2014) Generalist-specialist continuum and life

history traits of Central European butterflies (Lepidoptera) - are we missing a part of the

picture? *European Journal of Entomology*.

*BASSET, Y., BARRIOS, H., SEGAR, S., SRYGLEY, R.B., AIELLO, A., WARREN, A.D., DELGADO, F.,

CORONADO, J., LEZCANO, J., ARIZALA, S., RIVERA, M., PEREZ, F., BOBADILLA, R., LOPEZ, Y. &

RAMIREZ, J.A. (2015) The Butterflies of Barro Colorado Island, Panama: Local Extinction

since the 1930s. *PLoS ONE* **10**, e0136623.

*BENDER, M. G., FLOETER, S. R., MAYER, F. P., VILA-NOVA, D. A., LONGO, G. O., HANAZAKI, N.,

CARVALHO-FILHO, A. & FERREIRA, C. E. L. (2013) Biological attributes and major threats as

predictors of the vulnerability of species: a case study with Brazilian reef fishes. *Oryx* **47**, 259–

265.

*BENNETT, P.M. & OWENS, I.P. (1997) Variation in extinction risk among birds: chance or

evolutionary predisposition? *Proceedings of the Royal Society of London B: Biological*

Sciences **264**, 401–408.

*BENSCOTER, A.M., REECE, J.S., NOSS, R.F., BRANDT, L.A., MAZZOTTI, F.J., ROMAÑACH, S.S. &

WATLING, J.I. (2013) Threatened and Endangered Subspecies with Vulnerable Ecological

Traits Also Have High Susceptibility to Sea Level Rise and Habitat Fragmentation. *PLoS ONE*

8, e70647.

*BETZHOLTZ, P.-E., FRANZÉN, M. & FORSMAN, A. (2017) Colour pattern variation can inform about

extinction risk in moths. *Animal Conservation* **20**, 72–79.

BININDA-EMONDS, O.R.P., CARDILLO, M., JONES, K.E., MACPHEE, R.D.E., BECK, R.M.D.,

GRENYER, R., PRICE, S.A., VOS, R.A., GITTLEMAN, J.L. & PURVIS, A. (2007) The delayed rise

of present-day mammals. *Nature* **446**, 507–512.

*BLAND, L.M. (2017) Global correlates of extinction risk in freshwater crayfish. *Animal*

Conservation **20**, 532–542.

BLAND, L.M., COLLEN, B., ORME, C.D.L. & BIELBY, J. (2015) Predicting the conservation status of

data-deficient species. *Conservation Biology* **29**, 250–259.

*BÖHM, M., WILLIAMS, R., BRAMHALL, H.R., MCMILLAN, K.M., DAVIDSON, A.D., GARCIA, A.,

BLAND, L.M., BIELBY, J. & COLLEN, B. (2016) Correlates of extinction risk in squamate

reptiles: the relative importance of biology, geography, threat and range size. *Global Ecology*

and Biogeography **25**, 391–405.

*BONELLI, S., CERRATO, C., LOGLISCI, N. & BALLETTO, E. (2011) Population extinctions in the

Italian diurnal lepidoptera: an analysis of possible causes. *Journal of Insect Conservation* **15**,

879–890.

*BOTTS, E.A., ERASMUS, B.F.N. & ALEXANDER, G.J. (2013) Small range size and narrow niche

breadth predict range contractions in South African frogs. *Global Ecology and Biogeography*

22, 567–576.

BOUCHET-VALAT, M. (2014) *SnowballC: Snowball stemmers based on the C libstemmer UTF-8*

library. R package version 0.5.1. https://CRAN.R-project.org/package=SnowballC

*BOYLES, J.G. & STORM, J.J. (2007) The Perils of Picky Eating: Dietary Breadth Is Related to

Extinction Risk in Insectivorous Bats. *PLoS ONE* **2**, e672.

*BRADSHAW, C.J.A., GIAM, X., TAN, H.T.W., BROOK, B.W. & SODHI, N.S. (2008) Threat or

invasive status in legumes is related to opposite extremes of the same ecological and life-

history attributes: Twin fates of legume species. *Journal of Ecology* **96**, 869–883.

*Bʀᴏ-Jørgensen, J. (2014) Will their armaments be their downfall? Large horn size increases

extinction risk in bovids: Large horn size increases extinction risk in bovids. *Animal*

Conservation **17**, 80–87.

*Bᴜʙᴏᴠá, T., Kᴜʟᴍᴀ, M., Vʀᴀʙᴇᴄ, V. & Nᴏᴡɪᴄᴋɪ, P. (2016) Adult longevity and its relationship

with conservation status in European butterflies. *Journal of Insect Conservation* **20**, 1021–

1032.

Bᴜʀɢᴍᴀɴ, M.A. (1989) The Habitat Volumes of Scarce and Ubiquitous Plants: A Test of the

Model of Environmental Control. *The American Naturalist* **133**, 228–239.

Cᴀʀʙᴏɴᴇ, C. & Gɪᴛᴛʟᴇᴍᴀɴ, J.L. (2002) A Common Rule for the Scaling of Carnivore Density.

Science **295**, 2273–2276.

*Cᴀʀᴅɪʟʟᴏ, M. & Bʀᴏᴍʜᴀᴍ, L. (2001) Body Size and Risk of Extinction in Australian Mammals.

Conservation Biology **15**, 1435–1440.

Cᴀʀᴅɪʟʟᴏ, M. & Mᴇɪᴊᴀᴀʀᴅ, E. (2012) Are comparative studies of extinction risk useful for

conservation? *Trends in Ecology & Evolution* **27**, 167–171.

*Cᴀʀᴅɪʟʟᴏ, M. (2003) Biological determinants of extinction risk: why are smaller species less

vulnerable? *Animal Conservation* **6**, 63–69.

*Cᴀʀᴅɪʟʟᴏ, M., Mᴀᴄᴇ, G.M., Gɪᴛᴛʟᴇᴍᴀɴ, J.L., Jᴏɴᴇs, K.E., Bɪᴇʟʙʏ, J. & Pᴜʀᴠɪs, A. (2008) The

predictability of extinction: biological and external correlates of decline in mammals.

Proceedings of the Royal Society of London B: Biological Sciences **275**, 1441–1448.

*Cᴀʀᴅɪʟʟᴏ, M., Mᴀᴄᴇ, G.M., Jᴏɴᴇs, K.E., Bɪᴇʟʙʏ, J., Bɪɴɪɴᴅᴀ-Eᴍᴏɴᴅs, O.R.P., Sᴇᴄʜʀᴇsᴛ, W.,

Oʀᴍᴇ, C.D.L. & Pᴜʀᴠɪs, A. (2005) Multiple Causes of High Extinction Risk in Large

Mammal Species. *Science* **309**, 1239–1241.

*Cardillo, M., Purvis, A., Sechrest, W., Gittleman, J.L., Bielby, J. & Mace, G.M. (2004)

Human Population Density and Extinction Risk in the World's Carnivores. *PLoS Biol* **2**, e197.

Cardoso, P., Arnedo, M.A., Triantis, K.A. & Borges, P.A.V. (2010) Drivers of diversity in

Macaronesian spiders and the role of species extinctions. *Journal of Biogeography* **37**, 1034–

1046.

Cardoso, P., Borges, P.A.V., Triantis, K.A., Ferrández, M.A. & Martín, J.L. (2011a)

Adapting the IUCN Red List criteria for invertebrates. *Biological Conservation* **144**, 2432–

2440.

Cardoso, P., Borges, P.A.V., Triantis, K.A., Ferrández, M.A. & Martín, J.L. (2012) The

underrepresentation and misrepresentation of invertebrates in the IUCN Red List. *Biological*

Conservation **149**, 147–148.

Cardoso, P., Erwin, T.L., Borges, P.A.V. & New, T.R. (2011b) The seven impediments in

invertebrate conservation and how to overcome them. *Biological Conservation* **144**, 2647–

2655.

*Cassey, P. (2001) Determining variation in the success of New Zealand land birds. *Global*

Ecology & Biogeography, 161–172.

Chaplin-Kramer, R., O'Rourke, M.E., Blitzer, E.J. & Kremen, C. (2011) A meta-analysis of

crop pest and natural enemy response to landscape complexity. *Ecology Letters* **14**, 922–932.

*Chessman, B.C. (2013) Identifying species at risk from climate change: Traits predict the drought

vulnerability of freshwater fishes. *Biological Conservation* **160**, 40–49.

*CHIBA, S. & ROY, K. (2011) Selectivity of terrestrial gastropod extinctions on an oceanic

archipelago and insights into the anthropogenic extinction process. *Proceedings of the*

National Academy of Sciences **108**, 9496–9501.

*CHISHOLM, R.A. & TAYLOR, R. (2010) Body size and extinction risk in Australian mammals: An

information-theoretic approach: INFORMATION THEORY AND EXTINCTION RISK.

Austral Ecology **35**, 616–623.

CLAVEL, J., JULLIARD, R. & DEVICTOR, V. (2011) Worldwide decline of specialist species: toward a

global functional homogenization? *Frontiers in Ecology and the Environment* **9**, 222–228.

*COLLEN, B., BYKOVA, E., LING, S., MILNER-GULLAND, E.J. & PURVIS, A. (2006) Extinction Risk:

A Comparative Analysis of Central Asian Vertebrates. *Biodiversity & Conservation* **15**, 1859–

1871.

*COLLEN, B., MCRAE, L., DEINET, S., DE PALMA, A., CARRANZA, T., COOPER, N., LOH, J. &

BAILLIE, J.E.M. (2011) Predicting how populations decline to extinction. *Philosophical*

Transactions of the Royal Society B: Biological Sciences **366**, 2577–2586.

*COMEROS-RAYNAL, M.T., POLIDORO, B.A., BROATCH, J., MANN, B.Q., GORMAN, C., BUXTON,

C.D., GOODPASTER, A.M., IWATSUKI, Y., MACDONALD, T.C., POLLARD, D., RUSSELL, B. &

CARPENTER, K.E. (2016) Key predictors of extinction risk in sea breams and porgies (Family:

Sparidae). *Biological Conservation* **202**, 88–98.

*COOPER, N., BIELBY, J., THOMAS, G.H. & PURVIS, A. (2008) Macroecology and extinction risk

correlates of frogs. *Global Ecology and Biogeography* **17**, 211–221.

*CRIMMINS, S.M., BOMA, P. & THOGMARTIN, W.E. (2015) Projected Risk of Population Declines

for Native Fish Species in the Upper Mississippi River. *River Research and Applications* **31**,

135–142.

*DARRAH, S.E., BLAND, L.M., BACHMAN, S.P., CLUBBE, C.P. & TRIAS-BLASI, A. (2017) Using

coarse-scale species distribution data to predict extinction risk in plants. *Diversity and*

Distributions **23**, 435–447.

*DAVIDSON, A.D., BOYER, A.G., KIM, H., POMPA-MANSILLA, S., HAMILTON, M.J., COSTA, D.P.,

CEBALLOS, G. & BROWN, J.H. (2012) Drivers and hotspots of extinction risk in marine

mammals. *Proceedings of the National Academy of Sciences* **109**, 3395–3400.

*DAVIDSON, A.D., HAMILTON, M.J., BOYER, A.G., BROWN, J.H. & CEBALLOS, G. (2009) Multiple

ecological pathways to extinction in mammals. *Proceedings of the National Academy of*

Sciences **106**, 10702–10705.

*DAVIDSON, A.D., SHOEMAKER, K.T., WEINSTEIN, B., COSTA, G.C., BROOKS, T.M., CEBALLOS, G.,

RADELOFF, V.C., RONDININI, C. & GRAHAM, C.H. (2017) Geography of current and future

global mammal extinction risk. *PLoS ONE* **12**, e0186934.

*DAVIES, T.J., SMITH, G.F., BELLSTEDT, D.U., BOATWRIGHT, J.S., BYTEBIER, B., COWLING, R.M.,

FOREST, F., HARMON, L.J., MUASYA, A.M., SCHRIRE, B.D., STEENKAMP, Y., VAN DER BANK,

M. & SAVOLAINEN, V. (2011) Extinction Risk and Diversification Are Linked in a Plant

Biodiversity Hotspot. *PLoS Biology* **9**, e1000620.

*DESENDER, K., DEKONINCK, W., DUFRÊNE, M. & MAES, D. (2010) Changes in the distribution of

carabid beetles in Belgium revisited: Have we halted the diversity loss? *Biological*

Conservation **143**, 1549–1557.

*DI MARCO, M., BUCHANAN, G.M., SZANTOI, Z., HOLMGREN, M., GROTTOLO MARASINI, G.,

GROSS, D., TRANQUILLI, S., BOITANI, L. & RONDININI, C. (2014) Drivers of extinction risk in

African mammals: the interplay of distribution state, human pressure, conservation response

and species biology. *Philosophical Transactions of the Royal Society B: Biological Sciences*

369, 20130198.

*DUNCAN, J.R. & LOCKWOOD, J.L. (2001) Extinction in a field of bullets: a search for causes in the

decline of the world's freshwater fishes. *Biological Conservation* **102**,: 97–105.

*DUNCAN, R.P. & BLACKBURN, T.M. (2004) Extinction and endemism in the New Zealand

avifauna. *Global Ecology and Biogeography* **13**, 509–517.

*ESSENS, T., VAN LANGEVELDE, F., VOS, R.A., VAN SWAAY, C.A.M. & WALLISDEVRIES, M.F.

(2017) Ecological determinants of butterfly vulnerability across the European continent.

Journal of Insect Conservation **21**, 439–450.

FALSTER, D.S. & WESTOBY, M. (2003) Plant height and evolutionary games. *Trends in Ecology &*

Evolution **18**, 337–343.

*FATTORINI, S. (2013) Species ecological preferences predict extinction risk in urban tenebrionid

beetle guilds. *Animal Biology* **63**, 93–106.

FEINERER, I. & HORNIK, K. (2008) Text Mining Infrastructure in R. *Journal of Statistical Software*

25, 1–54.

*FIELD, I.C., MEEKAN, M.G., BUCKWORTH, R.C. & BRADSHAW, C.J.A. (2009) Chapter 4

Susceptibility of Sharks, Rays and Chimaeras to Global Extinction. In *Advances in Marine*

Biology pp. 275–363. Elsevier.

FISHER, D.O. & OWENS, I.P.F. (2004) The comparative method in conservation biology. *Trends in*

Ecology & Evolution **19**, 391–398.

*FISHER, D.O., BLOMBERG, S.P. & OWENS, I.P. (2003) Extrinsic versus intrinsic factors in the

decline and extinction of Australian marsupials. *Proceedings of the Royal Society of London B:*

Biological Sciences **270**, 1801–1808.

*FLOUSEK, J., TELENSKÝ, T., HANZELKA, J. & REIF, J. (2015) Population Trends of Central

European Montane Birds Provide Evidence for Adverse Impacts of Climate Change on High-

Altitude Species. *PLoS ONE* **10**, e0139465.

*FORERO-MEDINA, G., VIEIRA, M.V., GRELLE, C.E. DE V. & ALMEIDA, P.J. (2009) Body size and

extinction risk in Brazilian carnivores. *Biota Neotropica* **9**, 45–49.

*FORISTER, M.L., JAHNER, J.P., CASNER, K.L., WILSON, J.S. & SHAPIRO, A.M. (2011) The race is

not to the swift: Long-term data reveal pervasive declines in California's low-elevation

butterfly fauna. *Ecology* **92**, 2222–2235.

*FRITZ, S.A., BININDA-EMONDS, O.R.P. & PURVIS, A. (2009) Geographical variation in predictors

of mammalian extinction risk: big is bad, but only in the tropics. *Ecology Letters* **12**, 538–549.

*GAGE, G.S., BROOKE, M. DE L., SYMONDS, M.R.E. & WEGE, D. (2004) Ecological correlates of the

threat of extinction in Neotropical bird species. *Animal Conservation* **7**, 161–168.

*GARCIA, V.B., LUCIFORA, L.O. & MYERS, R.A. (2008) The importance of habitat and life history

to extinction risk in sharks, skates, rays and chimaeras. *Proceedings of the Royal Society B:*

Biological Sciences **275**, 83–89.

*GARNETT, S.T. & BROOK, B.W. (2007) Modelling to forestall extinction of Australian tropical

birds. *Journal of Ornithology* **148**, 311–320.

*GASTON, K.J. & BLACKBURN, T.M. (1995) Birds, body size and the threat of extinction. *Phil.*

Trans. R. Soc. Lond. B **347**, 205–212.

*GASTON, K.J. (1997) Evolutionary age and risk of extinction in the global avifauna. *Evolutionary*

Ecology **11**, 557–565.

GASTON, K.J., BLACKBURN, T.M., GREENWOOD, J.J.D., GREGORY, R.D., QUINN, R.M. & LAWTON,

J.H. (2002) Abundance–occupancy relationships. *Journal of Applied Ecology* **37**, 39–59.

*GIAM, X., NG, T.H., LOK, A.F.S.L. & NG, H.H. (2011) Local geographic range predicts freshwater

fish extinctions in Singapore: Extinction correlates of tropical freshwater fish. *Journal of*

Applied Ecology **48**, 356–363.

*GIBB, H., HJÄLTÉN, J., BALL, J.P., PETTERSSON, R.B., LANDIN, J., ALVINI, O. & DANELL, K. (2006)

Wing loading and habitat selection in forest beetles: Are red-listed species poorer dispersers or

more habitat-specific than common congenerics? *Biological Conservation* **132**, 250–260.

*GLASS, W.R., CORKUM, L.D. & MANDRAK, N.E. (2017) Living on the edge: Traits of freshwater

fish species at risk in Canada. *Aquatic Conservation: Marine and Freshwater Ecosystems* **27**,

938–945.

*GODEFROID, S. (2014) Do plant reproductive traits influence species susceptibility to decline?

Plant Ecology and Evolution **147**, 154–164.

GOLDSTEIN, R.E. (2018) Point of View: Are theoretical results 'Results'? *eLife* **7**, e40018.

GONZÁLEZ-OREJA, J.A. (2008) The Encyclopedia of Life vs. the Brochure of Life: Exploring the

relationships between the extinction of species and the inventory of life on Earth. *Zootaxa*

1965, 61–68.

*GONZÁLEZ-SUÁREZ, M. & REVILLA, E. (2013) Variability in life-history and ecological traits is a

buffer against extinction in mammals. *Ecology Letters* **16**, 242–251.

*GONZÁLEZ-SUÁREZ, M., GÓMEZ, A. & REVILLA, E. (2013) Which intrinsic traits predict

vulnerability to extinction depends on the actual threatening processes. *Ecosphere* **4**, 1–16.

*GONZALEZ-VOYER, A., GONZÁLEZ-SUÁREZ, M., VILÀ, C. & REVILLA, E. (2016) Larger brain size

indirectly increases vulnerability to extinction in mammals: BRAIN SIZE AND

VULNERABILITY TO EXTINCTION. *Evolution* **70**, 1364–1375.

*GRENOUILLET, G. & COMTE, L. (2014) Illuminating geographical patterns in species' range shifts.

Global Change Biology **20**, 3080–3091.

GRIMM, V., BERGER, U., DEANGELIS, D.L., POLHILL, J.G., GISKE, J. & RAILSBACK, S.F. (2010) The

ODD protocol: A review and first update. *Ecological Modelling* **221**, 2760–2768.

GUREVITCH, J., KORICHEVA, J., NAKAGAWA, S. & STEWART, G. (2018) Meta-analysis and the

science of research synthesis. *Nature* **555**, 175–182.

HALLMANN, C.A., SORG, M., JONGEJANS, E., SIEPEL, H., HOFLAND, N., SCHWAN, H., STENMANS,

W., MÜLLER, A., SUMSER, H., HÖRREN, T., GOULSON, D. & KROON, H. DE (2017) More than

75 percent decline over 27 years in total flying insect biomass in protected areas. *PLoS ONE*

12, e0185809.

*HAMMERSON, G.A., KLING, M., HARKNESS, M., ORMES, M. & YOUNG, B.E. (2017) Strong

geographic and temporal patterns in conservation status of North American bats. *Biological*

Conservation **212**, 144–152.

*HANNA, E. & CARDILLO, M. (2013) A comparison of current and reconstructed historic geographic

range sizes as predictors of extinction risk in Australian mammals. *Biological Conservation*

158, 196–204.

*HANNA, E. & CARDILLO, M. (2014) Clarifying the relationship between torpor and anthropogenic

extinction risk in mammals: Torpor and extinction risk in mammals. *Journal of Zoology* **293**,

211–217.

*HERO, J.-M., WILLIAMS, S.E. & MAGNUSSON, W.E. (2005) Ecological traits of declining

amphibians in upland areas of eastern Australia. *Journal of Zoology* **267**, 221–232.

*HOF, A.R., RODRÍGUEZ-CASTAÑEDA, G., ALLEN, A.M., JANSSON, R. & NILSSON, C. (2017)

Vulnerability of Subarctic and Arctic breeding birds. *Ecological Applications* **27**, 219–234.

*HOWARD, S.D. & BICKFORD, D.P. (2014) Amphibians over the edge: silent extinction risk of Data

Deficient species. *Diversity and Distributions* **20**, 837–846.

IUCN (2018) The IUCN Red List of Threatened Species. Version 2018-1.

http://www.iucnredlist.org [accessed 13 July 2018].

*JAGER, H.I., ROSE, K.A. & VILA-GISPERT, A. (2008) Life history correlates and extinction risk of

capital-breeding fishes. *Hydrobiologia* **602**, 15–25.

*JENNINGS, S., GREENSTREET, S.P.R. & REYNOLDS, J.D. (1999) Structural change in an exploited

fish community: a consequence of differential fishing effects on species with contrasting life

histories. *Journal of Animal Ecology* **68**, 617–627.

*JENNINGS, S., REYNOLDS, J.D. & MILLS, S.C. (1998) Life history correlates of responses to

fisheries exploitation. *Proceedings of the Royal Society of London B: Biological Sciences* **265**,

333–339.

*JEPPSSON, T. & FORSLUND, P. (2014) Species' traits explain differences in Red list status and long-

term population trends in longhorn beetles: Traits and extinction risk in longhorn beetles.

Animal Conservation **17**, 332–341.

*JIGUET, F., GADOT, A.-S., JULLIARD, R., NEWSON, S.E. & COUVET, D. (2007) Climate envelope,

life history traits and the resilience of birds facing global change. *Global Change Biology* **13**,

1672–1684.

*JIGUET, F., GREGORY, R.D., DEVICTOR, V., GREEN, R.E., VOŘÍŠEK, P., VAN STRIEN, A. & COUVET,

D. (2010) Population trends of European common birds are predicted by characteristics of their

climatic niche. *Global Change Biology* **16**, 497–505.

*JOHNSON, C.N. & ISAAC, J.L. (2009) Body mass and extinction risk in Australian marsupials: The

'Critical Weight Range' revisited. *Austral Ecology* **34**, 35–40.

*JOHNSON, C.N., DELEAN, S. & BALMFORD, A. (2002) Phylogeny and the selectivity of extinction

in Australian marsupials. *Animal Conservation* **5**, 135–142.

*JONES, K.E., PURVIS, A. & GITTLEMAN, J.L. (2003) Biological correlates of extinction risk in bats.

The American Naturalist **161**, 601–614.

*JONES, M.J., FIELDING, A. & SULLIVAN, M. (2006) Analysing Extinction Risk in Parrots using

Decision Trees. *Biodiversity & Conservation* **15**, 1993–2007.

*JUAN-JORDÁ, M.J., MOSQUEIRA, I., FREIRE, J. & DULVY, N.K. (2015) Population declines of tuna

and relatives depend on their speed of life. *Proceedings of the Royal Society B: Biological

Sciences* **282**, 20150322.

*KAMILAR, J.M. & PACIULLI, L.M. (2008) Examining the extinction risk of specialized folivores: a

comparative study of Colobine monkeys. *American Journal of Primatology* **70**, 816–827.

*KEANE, A., BROOKE, M. D. L. & MCGOWAN, P.J.K. (2005) Correlates of extinction risk and

hunting pressure in gamebirds (Galliformes). *Biological Conservation* **126**, 216–233.

*KOH, L.P., SODHI, N.S. & BROOK, B.W. (2004) Ecological Correlates of Extinction Proneness in

Tropical Butterflies: Extinction Correlates of Tropical Butterflies. *Conservation Biology* **18**,

1571–1578.

*KOLEČEK, J., ALBRECHT, T. & REIF, J. (2014) Predictors of extinction risk of passerine birds in a

Central European country: Extinction risk of passerines. *Animal Conservation* **17**, 498–506.

*KOPF, R.K., SHAW, C. & HUMPHRIES, P. (2017) Trait-based prediction of extinction risk of small-

bodied freshwater fishes: Predicting Extinction Risk. *Conservation Biology* **31**, 581–591.

*KOTZE, D.J. & O'HARA, R.B. (2003) Species decline - but why? Explanations of carabid beetle

(Coleoptera, Carabidae) declines in Europe. *Oecologia* **135**, 138–148.

*KRÜGER, O. & RADFORD, A.N. (2008) Doomed to die? Predicting extinction risk in the true hawks

Accipitridae. *Animal Conservation* **11**, 83–91.

*LAANISTO, L., SAMMUL, M., KULL, T., MACEK, P. & HUTCHINGS, M.J. (2015) Trait-based analysis

of decline in plant species ranges during the 20th century: a regional comparison between the

UK and Estonia. *Global Change Biology* **21**, 2726–2738.

*LAUTERBACH, D., RÖMERMANN, C., JELTSCH, F. & RISTOW, M. (2013) Factors driving plant rarity

in dry grasslands on different spatial scales: a functional trait approach. *Biodiversity and*

Conservation **22**, 2337–2352.

*LAVERGNE, S., EVANS, M.E.K., BURFIELD, I.J., JIGUET, F. & THUILLER, W. (2012) Are species'

responses to global change predicted by past niche evolution? *Philosophical Transactions of*

the Royal Society B: Biological Sciences **368**, 20120091–20120091.

*LAVERGNE, S., MOLINA, J. & DEBUSSCHE, M. (2006) Fingerprints of environmental change on the

rare mediterranean flora: a 115-year study. *Global Change Biology* **12**, 1466–1478.

*LAWES, M.J., FISHER, D.O., JOHNSON, C.N., BLOMBERG, S.P., FRANK, A.S., FRITZ, S.A.,

MCCALLUM, H., VANDERWAL, J., ABBOTT, B.N. & LEGGE, S. (2015) Correlates of recent

declines of rodents in northern and southern Australia: habitat structure is critical. *PLoS One*

10, e0130626.

*LEACH, K., KELLY, R., CAMERON, A., MONTGOMERY, W.I. & REID, N. (2015) Expertly Validated

Models and Phylogenetically-Controlled Analysis Suggests Responses to Climate Change Are

Related to Species Traits in the Order Lagomorpha. *PLoS ONE* **10**, e0122267.

*LEÃO, T.C.C., FONSECA, C.R., PERES, C.A. & TABARELLI, M. (2014) Predicting Extinction Risk of

Brazilian Atlantic Forest Angiosperms: Neotropical Plant Extinction Risk. *Conservation*

Biology **28**, 1349–1359.

*LEE, T.M. & JETZ, W. (2010) Unravelling the structure of species extinction risk for predictive

conservation science. *Proceedings of the Royal Society of London B: Biological Sciences*,

rspb20101877.

LIAW, A. & WIENER, M. (2002) Classification and Regression by randomForest. *R News* **2**, 18–22.

*LIOW, L.H., FORTELIUS, M., LINTULAAKSO, K., MANNILA, H. & STENSETH, N.C. (2009) Lower

Extinction Risk in Sleep‐or‐Hide Mammals. *The American Naturalist* **173**, 264–272.

*LIPS, K.R., REEVE, J.D. & WITTERS, L.R. (2003) Ecological Traits Predicting Amphibian

Population Declines in Central America. *Conservation Biology* **17**, 1078–1088.

*LIU, C., COMTE, L. & OLDEN, J.D. (2017) Heads you win, tails you lose: Life-history traits predict

invasion and extinction risk of the world's freshwater fishes: TRAITS PREDICT GLOBAL

FRESHWATER FISH INVASION AND EXTINCTION. *Aquatic Conservation: Marine and*

Freshwater Ecosystems **27**, 773–779.

*LONG, P.R., SZÉKELY, T., KERSHAW, M. & O'CONNELL, M. (2007) Ecological factors and human

threats both drive wildfowl population declines. *Animal Conservation* **10**, 183–191.

*LOOTVOET, A.C., PHILIPPON, J. & BESSA-GOMES, C. (2015) Behavioral Correlates of Primates

Conservation Status: Intrinsic Vulnerability to Anthropogenic Threats. *PLoS ONE* **10**,

e0135585.

*LUIZ, O.J., WOODS, R.M., MADIN, E.M.P. & MADIN, J.S. (2016) Predicting IUCN Extinction Risk

Categories for the World's Data Deficient Groupers (Teleostei: Epinephelidae). *Conservation*

Letters **9**, 342–350.

*MACE, G.M., COLLEN, B., FULLER, R.A. & BOAKES, E.H. (2010) Population and geographic range

dynamics: implications for conservation planning. *Philosophical Transactions of the Royal*

Society B: Biological Sciences **365**, 3743–3751.

*MACHADO, N. & LOYOLA, R.D. (2013) A Comprehensive Quantitative Assessment of Bird

Extinction Risk in Brazil. *PLoS ONE* **8**, e72283.

*MANKGA, L.T. & YESSOUFOU, K. (2017) Factors driving the global decline of cycad diversity.

AoB Plants **9**, plx022.

*MATSUZAKI, S.-I.S., TAKAMURA, N., ARAYAMA, K., TOMINAGA, A., IWASAKI, J. & WASHITANI, I.

(2011) Potential impacts of non-native channel catfish on commercially important species in a

Japanese lake, as inferred from long-term monitoring data. *Aquatic Conservation: Marine and*

Freshwater Ecosystems **21**, 348–357.

*MATTHEWS, L.J., ARNOLD, C., MACHANDA, Z. & NUNN, C.L. (2011) Primate extinction risk and

historical patterns of speciation and extinction in relation to body mass. *Proceedings of the*

Royal Society B: Biological Sciences **278**, 1256–1263.

*MATTILA, N., KAITALA, V., KOMONEN, A., PÄIVINEN, J. & KOTIAHO, J.S. (2011) Ecological

correlates of distribution change and range shift in butterflies: Distribution decline in

butterflies. *Insect Conservation and Diversity* **4**, 239–246.

*MATTILA, N., KOTIAHO, J.S., KAITALA, V. & KOMONEN, A. (2008) The use of ecological traits in

extinction risk assessments: A case study on geometrid moths. *Biological Conservation* **141**,

2322–2328.

*MATTILA, N., KOTIAHO, J.S., KAITALA, V., KOMONEN, A. & PÄIVINEN, J. (2009) Interactions

between Ecological Traits and Host Plant Type Explain Distribution Change in Noctuid

Moths. *Conservation Biology* **23**, 703–709.

MCKINNEY, M.L. & LOCKWOOD, J.L. (1999) Biotic homogenization: a few winners replacing many

losers in the next mass extinction. *Trends in Ecology & Evolution* **14**, 450–453.

MCKINNEY, M.L. (1997) Extinction Vulnerability and Selectivity: Combining Ecological and

Paleontological Views. *Annual Review of Ecology and Systematics* **28**, 495–516.

MILBORROW, S. (2018) *rpart.plot: Plot 'rpart' Models: An Enhanced Version of 'plot.rpart'*. R

package version 2.2.0. https://CRAN.R-project.org/package=rpart.plot

*MONKS, A. & BURROWS, L. (2014) Are threatened plant species specialists, or just more

vulnerable to disturbance? *Journal of Applied Ecology* **51**, 1228–1235.

MORA, C., TITTENSOR, D.P., ADL, S., SIMPSON, A.G.B. & WORM, B. (2011) How Many Species Are

There on Earth and in the Ocean? *PLOS Biology* **9**, e1001127.

*MORROW, E.H. & FRICKE, C. (2004) Sexual selection and the risk of extinction in mammals.

Proceedings of the Royal Society B: Biological Sciences **271**, 2395–2401.

*MORROW, E.H. & PITCHER, T.E. (2003) Sexual selection and the risk of extinction in birds.

Proceedings of the Royal Society of London B: Biological Sciences **270**, 1793–1799.

*MURPHY, B.P. & DAVIES, H.F. (2014) There is a critical weight range for Australia's declining

tropical mammals: Critical weight range for Australia's declining tropical mammals. *Global*

Ecology and Biogeography **23**, 1058–1061.

*MURRAY, B.R. & HOSE, G.C. (2005) Life-history and ecological correlates of decline and

extinction in the endemic Australian frog fauna. *Austral Ecology* **30**, 564–571.

*MURRAY, B.R., THRALL, P.H. & LEPSCHI, B.J. (2002) Relating species rarity to life history in

plants of eastern Australia, 14.

*MURRAY, K.A., ROSAUER, D., MCCALLUM, H. & SKERRATT, L.F. (2011) Integrating species traits

with extrinsic threats: closing the gap between predicting and preventing species declines.

Proceedings of the Royal Society of London B: Biological Sciences, rspb20101872.

MURRAY, K.A., VERDE ARREGOITIA, L.D., DAVIDSON, A., DI MARCO, M. & DI FONZO, M.M.I.

(2014) Threat to the point: improving the value of comparative extinction risk analysis for

conservation action. *Global Change Biology* **20**, 483–494.

*MUSTERS, C.J.M., KALKMAN, V. & VAN STRIEN, A. (2013) Predicting rarity and decline in

animals, plants, and mushrooms based on species attributes and indicator groups. *Ecology and*

Evolution, 3401–3414.

*NÁJERA, A. & SIMONETTI, J.A. (2012) Threatened birds of Guatemala: a random subset of the

avifauna? *Bird Conservation International* **22**, 348–353.

*NORRIS, K. & HARPER, N. (2004) Extinction processes in hot spots of avian biodiversity and the

targeting of pre–emptive conservation action. *Proceedings of the Royal Society of London B:*

Biological Sciences **271**, 123–130.

*NYLIN, S. & BERGSTRÖM, A. (2009) Threat status in butterflies and its ecological correlates: how

far can we generalize? *Biodiversity and Conservation* **18**, 3243–3267.

*OLDEN, J.D., HOGAN, Z.S. & ZANDEN, M.J.V. (2007) Small fish, big fish, red fish, blue fish: size-

biased extinction risk of the world's freshwater and marine fishes. *Global Ecology and*

Biogeography **16**, 694–701.

*OLDEN, J.D., POFF, N.L. & BESTGEN, K.R. (2008) Trait synergisms and the rarity, extirpation, and

extinction risk of desert fishes. *Ecology* **89**, 847–856.

OLSON, D.M., DINERSTEIN, E., WIKRAMANAYAKE, E.D., BURGESS, N.D., POWELL, G.V.N.,

UNDERWOOD, E.C., D'AMICO, J.A., ITOUA, I., STRAND, H.E., MORRISON, J.C., LOUCKS, C.J.,

ALLNUTT, T.F., RICKETTS, T.H., KURA, Y., LAMOREUX, J.F., ET AL. (2001) Terrestrial

Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial

ecoregions provides an innovative tool for conserving biodiversity. *BioScience* **51**, 933–938.

*OWENS, I.P.F. & BENNETT, P.M. (2000) Ecological basis of extinction risk in birds: Habitat loss

versus human persecution and introduced predators. *Proceedings of the National Academy of*

Sciences **97**, 12144–12148.

*PARENT, S. & SCHRIML, L.M. (1995) A model for the determination of fish species at risk based

upon life-history traits and ecological data. *Canadian Journal of Fisheries and Aquatic*

Sciences **52**, 1768–1781.

*PARLATO, E.H., ARMSTRONG, D.P. & INNES, J.G. (2015) Traits influencing range contraction in

New Zealand's endemic forest birds. *Oecologia* **179**, 319–328.

*PAYNE, J.L., BUSH, A.M., HEIM, N.A., KNOPE, M.L. & MCCAULEY, D.J. (2016) Ecological

selectivity of the emerging mass extinction in the oceans. *Science*, aaf2416.

*PEÑARANDA, D.A. & SIMONETTI, J.A. (2015) Predicting and setting conservation priorities for

Bolivian mammals based on biological correlates of the risk of decline: Biological Correlates

of Decline Risk. *Conservation Biology* **29**, 834–843.

PIMM, S.L., JONES, H.L. & DIAMOND, J. (1988) On the Risk of Extinction. *The American Naturalist*

132, 757–785.

*PINSKY, M.L. & BYLER, D. (2015) Fishing, fast growth and climate variability increase the risk of

collapse. *Proceedings of the Royal Society B: Biological Sciences* **282**, 20151053.

*PINSKY, M.L., JENSEN, O.P., RICARD, D. & PALUMBI, S.R. (2011) Unexpected patterns of fisheries

collapse in the world's oceans. *Proceedings of the National Academy of Sciences* **108**, 8317–

8322.

*PLEGUEZUELOS, J.M., BRITO, J.C., FAHD, S., FERICHE, M., MATEO, J.A., MORENO-RUEDA, G.,

REQUES, R. & SANTOS, X. (2010) Setting conservation priorities for the Moroccan

herpetofauna: the utility of regional red lists. *Oryx* **44**, 501–508.

*POCOCK, M.J.O. (2011) Can traits predict species' vulnerability? A test with farmland passerines

in two continents. *Proceedings of the Royal Society B: Biological Sciences* **278**, 1532–1538.

*POLAINA, E., REVILLA, E. & GONZÁLEZ-SUÁREZ, M. (2016) Putting susceptibility on the map to

improve conservation planning, an example with terrestrial mammals. *Diversity and*

Distributions **22**, 881–892.

*POLISHCHUK, L.V., POPADIN, K.Y., BARANOVA, M.A. & KONDRASHOV, A.S. (2015) A genetic

component of extinction risk in mammals. *Oikos* **124**, 983–993.

*POWNEY, G.D., CHAM, S.S.A., SMALLSHIRE, D. & ISAAC, N.J.B. (2015) Trait correlates of

distribution trends in the Odonata of Britain and Ireland. *PeerJ* **3**, e1410.

*POWNEY, G.D., RAPACCIUOLO, G., PRESTON, C.D., PURVIS, A. & ROY, D.B. (2014) A

phylogenetically-informed trait-based analysis of range change in the vascular plant flora of

Britain. *Biodiversity and Conservation* **23**, 171–185.

*PRICE, S.A. & GITTLEMAN, J.L. (2007) Hunting to extinction: biology and regional economy

influence extinction risk and the impact of hunting in artiodactyls. *Proceedings of the Royal*

Society B: Biological Sciences **274**, 1845–1851.

PRUGH, L.R. (2009) An evaluation of patch connectivity measures. *Ecological Applications* **19**,

1300–1310.

*PURVIS, A., GITTLEMAN, J.L., COWLISHAW, G. & MACE, G.M. (2000a) Predicting extinction risk in

declining species. *Proceedings of the Royal Society of London B: Biological Sciences* **267**,

1947–1952.

PURVIS, A., JONES, K.E. & MACE, G.M. (2000b) Extinction. *BioEssays* **22**, 1123–1133.

R Core Team (2018) *R: A language and environment for statistical computing.* R Foundation for

Statistical Computing, Vienna, Austria.

RABINOWITZ, D. (1981) Seven forms of Rarity. In *Biological aspects of rare plant conservation* pp.

205–217. John Wiley & Sons Ltd., New York.

RAJAMARAN, A. & ULLMAN, J.D. (2011) *Mining of Massive Datasets*. Cambridge University Press,

Cambridge.

*REED, R.N. & SHINE, R. (2002) Lying in Wait for Extinction: Ecological Correlates of

Conservation Status among Australian Elapid Snakes. *Conservation Biology* **16**, 451–461.

*REYNOLDS, J.D., WEBB, T.J. & HAWKINS, L.A. (2005) Life history and ecological correlates of

extinction risk in European freshwater fishes. *Canadian Journal of Fisheries and Aquatic*

Sciences **62**, 854–862.

*RIBEIRO, J., COLLI, G.R., CALDWELL, J.P. & SOARES, A.M.V.M. (2016) An integrated trait-based

framework to predict extinction risk and guide conservation planning in biodiversity hotspots.

Biological Conservation **195**, 214–223.

*RIPPLE, W.J., WOLF, C., NEWSOME, T.M., HOFFMANN, M., WIRSING, A.J. & MCCAULEY, D.J.

(2017) Extinction risk is most acute for the world's largest and smallest vertebrates.

Proceedings of the National Academy of Sciences, 201702078.

*ROLLAND, J. & SALAMIN, N. (2016) Niche width impacts vertebrate diversification. *Global*

Ecology and Biogeography **25**, 1252–1263.

ROSENTHAL, R. (1991) *Meta-Analytic Procedures for Social Research*. SAGE Publications,

London.

ROSKOV, Y., ABUCAY, L., ORREL, T., NICOLSON, D., BAILLY, N., KIRK, P., BOURGOIN, T., DEWALT,

R., DECOCK, W., DE WEVER, A., VAN NIEUKERKEN, E., ZARUCCHI, J. & PENEV, L. (2017)

Species 2000 & ITIS Catalogue of Life, 2017 Annual Checklist. Species 2000: Naturalis,

Leiden, The Netherlands.

*RULAND, F. & JESCHKE, J.M. (2017) Threat-dependent traits of endangered frogs. *Biological*

Conservation **206**, 310–313.

*SAAR, L., TAKKIS, K., PÄRTEL, M. & HELM, A. (2012) Which plant traits predict species loss in

calcareous grasslands with extinction debt? *Diversity and Distributions* **18**, 808–817.

*SAFI, K. & KERTH, G. (2004) A Comparative Analysis of Specialization and Extinction Risk in

Temperate-Zone Bats. *Conservation Biology* **18**, 1293–1303.

*SAGOT, M. & CHAVERRI, G. (2015) Effects of roost specialization on extinction risk in bats.

Conservation Biology **29**, 1666–1673.

*SAKAI, A.K., WAGNER, W.L. & MEHRHOFF, L.A. (2002) Patterns of Endangerment in the

Hawaiian Flora. *Systematic Biology* **51**, 276–302.

*SCHMIDT, J.P., STEPHENS, P.R. & DRAKE, J.M. (2012) Two sides of the same coin? Rare and pest

plants native to the United States and Canada. *Ecological Applications* **22**, 1512–1525.

*SEIBOLD, S., BRANDL, R., BUSE, J., HOTHORN, T., SCHMIDL, J., THORN, S. & MÜLLER, J. (2015)

Association of extinction risk of saproxylic beetles with ecological degradation of forests in

Europe: Beetle Extinction and Forest Degradation. *Conservation Biology* **29**, 382–390.

*SEOANE, J. & CARRASCAL, L.M. (2007) Interspecific differences in population trends of Spanish

birds are related to habitat and climatic preferences. *Global Ecology and Biogeography* **17**,

111–121.

*SHULTZ, S., B. BRADBURY, R., L. EVANS, K., D. GREGORY, R. & M. BLACKBURN, T. (2005) Brain

size and resource specialization predict long-term population trends in British birds.

Proceedings of the Royal Society B: Biological Sciences **272**, 2305–2311.

*SILICEO, I. & DÍAZ, J.A. (2010) A comparative study of clutch size, range size, and the

conservation status of island vs. mainland lacertid lizards. *Biological Conservation* **143**, 2601–

2608.

SLATYER, R.A., HIRST, M. & SEXTON, J.P. (2013) Niche breadth predicts geographical range size: a

general ecological pattern. *Ecology Letters* **16**, 1104–1114.

*SODHI, N.S., KOH, L.P., PEH, K.S.-H., TAN, H.T.W., CHAZDON, R.L., CORLETT, R.T., LEE, T.M.,

COLWELL, R.K., BROOK, B.W., SEKERCIOGLU, C.H. & BRADSHAW, C.J.A. (2008) Correlates of

extinction proneness in tropical angiosperms. *Diversity and Distributions* **14**, 1–10.

*SODHI, N.S., LEE, T.M., KOH, L.P. & PRAWIRADILAGA, D.M. (2006) Long-Term Avifaunal

Impoverishment in an Isolated Tropical Woodlot. *Conservation Biology* **20**, 772–779.

*SREEKAR, R., HUANG, G., ZHAO, J.-B., PASION, B.O., YASUDA, M., ZHANG, K., PEABOTUWAGE, I.,

WANG, X., QUAN, R.-C., FERRY SLIK, J.W., CORLETT, R.T., GOODALE, E. & HARRISON, R.D.

(2015) The use of species-area relationships to partition the effects of hunting and

deforestation on bird extirpations in a fragmented landscape. *Diversity and Distributions* **21**,

441–450.

*STEFANAKI, A., KANTSA, A., TSCHEULIN, T., CHARITONIDOU, M. & PETANIDOU, T. (2015) Lessons

from Red Data Books: Plant Vulnerability Increases with Floral Complexity. *PLoS ONE* **10**,

e0138414.

STUART, S.N., WILSON, E.O., MCNEELY, J.A., MITTERMEIER, R.A. & RODRÍGUEZ, J.P. (2010) The

Barometer of Life. *Science* **328**, 177–177.

*SULLIVAN, M.S., GILBERT, F., ROTHERAY, G., CROASDALE, S. & JONES, M. (2000) Comparative

analyses of correlates of Red data book status: a case study using European hoverflies

(Diptera: Syrphidae). *Animal Conservation* **3**, 91–95.

*TERZOPOULOU, S., RIGAL, F., WHITTAKER, R.J., BORGES, P.A.V. & TRIANTIS, K.A. (2015) Drivers

of extinction: the case of Azorean beetles. *Biology Letters* **11**, 20150273.

*THAXTER, C.B., JOYS, A.C., GREGORY, R.D., BAILLIE, S.R. & NOBLE, D.G. (2010) Hypotheses to

explain patterns of population change among breeding bird species in England. *Biological*

Conservation **143**, 2006–2019.

THERNEAU, T. & ATKINSON, B. (2015) *rpart: Recursive Partitioning and Regression Trees*. R

package version 4.1–13. https://CRAN.R-project.org/package=rpart

*TINGLEY, R., HITCHMOUGH, R.A. & CHAPPLE, D.G. (2013) Life-history traits and extrinsic threats

determine extinction risk in New Zealand lizards. *Biological Conservation* **165**, 62–68.

*TINGLEY, R., MAHONEY, P.J., DURSO, A.M., TALLIAN, A.G., MORÁN-ORDÓÑEZ, A. & BEARD,

K.H. (2016) Threatened and invasive reptiles are not two sides of the same coin: Extinction

and invasion risk in reptiles. *Global Ecology and Biogeography* **25**, 1050–1060.

*TRINDER-SMITH, H., COWLING, R.M. & LINDER, H.P. (1996) Profiling a besieged flora: endemic

and threatened plants of the Cape Peninsula, South Africa. *Biodiversity and Conservation* **5**,

575–589.

*VAMOSI, J.C. & VAMOSI, S.M. (2005) Present day risk of extinction may exacerbate the lower

species richness of dioecious clades: Extinction risk of dioecious plants. *Diversity and*

Distributions **11**, 25–32.

*VAN CALSTER, H., VANDENBERGHE, R., RUYSEN, M., VERHEYEN, K., HERMY, M. & DECOCQ, G.

(2008) Unexpectedly high 20th century floristic losses in a rural landscape in northern France:

Floristic changes in rural landscapes. *Journal of Ecology* **96**, 927–936.

*VAN TURNHOUT, C.A.M., FOPPEN, R.P.B., LEUVEN, R.S.E.W., VAN STRIEN, A. & SIEPEL, H.

(2010) Life-history and ecological correlates of population change in Dutch breeding birds.

Biological Conservation **143**, 173–181.

VERDE ARREGOITIA, L.D. (2016) Biases, gaps, and opportunities in mammalian extinction risk

research. *Mammal Review* **46**, 17–29.

*VERDE ARREGOITIA, L.D., LEACH, K., REID, N. & FISHER, D.O. (2015) Diversity, extinction, and

threat status in Lagomorphs. *Ecography* **38**, 1155–1165.

VIECHTBAUER, W. (2010) Conducting Meta-Analyses in R with the metafor Package. *Journal of*

Statistical Software **36**, 1–48.

*VILCHIS, L.I., JOHNSON, C.K., EVENSON, J.R., PEARSON, S.F., BARRY, K.L., DAVIDSON, P.,

RAPHAEL, M.G. & GAYDOS, J.K. (2015) Assessing ecological correlates of marine bird

declines to inform marine conservation: Ecological Correlates of Seabird Declines.

Conservation Biology **29**, 154–163.

*WALKER, K.J. & PRESTON, C.D. (2006) Ecological Predictors of Extinction Risk in the Flora of

Lowland England, UK. *Biodiversity & Conservation* **15**, 1913–1942.

*WALKER, K.J., PRESTON, C.D. & BOON, C.R. (2009) Fifty years of change in an area of intensive

agriculture: plant trait responses to habitat modification and conservation, Bedfordshire,

England. *Biodiversity and Conservation* **18**, 3597–3613.

*WANG, Y., SI, X., BENNETT, P.M., CHEN, C., ZENG, D., ZHAO, Y., WU, Y. & DING, P. (2018)

Ecological correlates of extinction risk in Chinese birds. *Ecography* **41**, 782–794.

WILSON, E.O. (2003) On global biodiversity estimates. *Paleobiology* **29**, 14–14.

1139

VIII. Supporting Information

Appendix S1 – List of all studies and traits included in the analyses

Appendix S2 – Data-frames with all information on studies and traits collected.

Appendix S3 – Important words in the automated filtering of publications

Appendix S4 – Proportion of significant measurements by trait and taxon

Appendix S5 – Forest plots of the meta-analyses

Figures
1147

1148

1149   Figure 1: Cumulative number of publications per taxon relating traits to extinction risk.

1150

1151

Figure 2: Proportion of manuscripts focusing on the different biogeographical regions by taxonomic group.

1153

1154

1155

1156 Figure 3: Number of studies and measurements per class of traits, as well as the number of unique traits in

1157 each major trait class (numbers before the dotted lines). **n_studies:** Number of studies in which the variable

1158 appears. **n_measurements:** total number of measurements for that variable.

1159

1160

Figure 4: Summary information on variable use among all studies, depicting only variables included in at least 17 (10%) studies. The numbers before the dotted lines indicate the percentage of measurements in which the variable was significant. **n_studies**: number of studies in which the variable appears. **n_measurements**: total number of measurements for that variable.

1161
1162
1163
1164

1165

1166   Tables

 Table 1: Traits and categories identified across the 173 studies analysed. The number of studies for each trait

 is shown in parentheses.

	Traits
Individual	Body size (141), fecundity (73), generation length (41), longevity (24), egg/neonatal size (21), nest microhabitat (20), gestation/incubation length (19), development type (13), plant growth form (13), interbirth interval (11), reproduction type (sexual or asexual) (11), parental care (10), sexual dimorphism (10), weaning age (10), pollen vector (9), fertilization mode (8), monogamy (7), brain size (6), population growth rate (6), gonad size (5), competitor richness (4), fledgling duration (4), age at eye opening (3), investment in reproduction (3), mass-specific metabolic rate (3), nest guarding (3), body shape (2), body size at weaning (2), colour detectability category (2), CSR ruderality (2), fecundity variability (2), nest vulnerability (2), vegetative resprouting capacity (2), body armour (1), body size variability (1), chela shape (1), chela size (1), chemical defence (1), clutch volume (1), colour pattern variability (1), CSR competitiveness (1), CSR strategy (1), CSR stress tolerance (1), egg-laying behaviour (1), egg adhesion (1), egg buoyancy (1), egg/neonatal size variability (1), endotherm (1), forearm length (1), forearm length variability (1), generation length variability (1), gestation/incubation length variability (1), incubating sex (1), interbirth interval variability (1), internal or external fertilization (1), larval body size (1), leaf length (1), leaf mass per unit area (1), leaf water content (1), male combat (1), male horn length (1), mouth brooding (1), niche uniqueness (1), number of reproductive years (1), number of teats (1), number of teats variability (1), plant coverage (1), ploidy (1), ploidy variability (1), plumage polymorphism (1), population juvenile ratio (1), population level sex ratio (1), sex change (1), specific leaf area (1), spire index (1), tail length (1), tongue length (1), weaning age variability (1)
Population	Population density (18), social group size (12), sociality (10), population abundance (6), spawning aggregation (2), survivability (2), population density variability (1), social group size variability (1)
Community	Diet type (49), diet breadth (35), floral colour (2), ambush predation (1), amount of floral reward (1), corolla segmentation (1), depending on symbioses (1), floral display (1), floral shape (1), floral size (1), functional reproductive unit (1), length of corolla tube (1), predator type (1)
Space	Geographical range size (81), location (39), migration distance (24), dispersal ability (21), home range size (17), insularity (13), wing size (13), dispersal agent (8), seed size (5), endemism (4), historical geographical range size (4), geographical range size change (3), wing shape (3), diaspore type (2), fruit dehiscence (2), geographic origin (2), location change (2), active dispersal (1), age at dispersal (1), air-breathing capability (1), altitudinal range shift (1), diaspore mass (1), geographical range size of host (1), home range size variability (1), reproductive area (1)
Time	Circadian period (17), overwintering stage (11), duration of reproductive season (10), timing of reproductive season (9), duration of flight period (8), torpor/hibernation (3), seasonality of reproduction (2), circadian range (1), flight period to adult life span (1), timing of migration (1), winter leaf carrying (1)
Evolution	Taxon age (5), evolutionary plasticity (2), rate of diversification (2), taxon richness (2), rate of climatic niche evolution (1), rate of habitat niche evolution (1), rate of trophic niche evolution (1)
Environmental niche	Habitat type (56), habitat breadth (54), microhabitat type (36), temperature (27), altitude (16), precipitation (16), temperature range (12), altitude range (7), evapotranspiration (6), precipitation range (6), productivity within range (6), nest habitat (5), temperature for breeding (3), climate range (2), annual productivity range (1), humidity (1), microhabitat breadth (1), shelter use (1), water balance within range (1), wingtip shape (1), wingtip size (1)
Human influence	Human population density (14), direct exploitation (13), land-use proportion (11), threat range (9), human footprint (7), threat type (7), land-use change (4), change in human population density (3), seed bank longevity (3), threat breadth (3), accessibility (2), gross domestic product (2), human resource intake (2), proportion of range protected (2), considered a pest (1), controlled by man (1), country responsible for management (1), economic value (1), human night-time lights (1), pollution tolerance (1), protection status (1)
Other	Compound metrics (6)
Taxonomy	Taxonomic group (16)

1169

1170　Table 2: Number of publications within each taxonomic group, response variables, controlling phylogeny or

1171　not, and statistical approach. Percentages are of the number of publications within each category divided by

1172　the total number of publications in the given taxonomic group. GEE = generalized estimating equation; GLM

1173　= generalized linear model; GLMM = generalized linear mixed model; LM = linear model; LMM = linear

1174　mixed model; PGCM = phylogenetic comparative method.

		All	Mammals	Birds	Reptiles	Amphibians	Fishes	Insects	Plants
No. of studies		173	50	42	10	14	29	24	23
Response variable	IUCNcateg	76 (44%)	35 (70%)	14 (33%)	4 (40%)	10 (71%)	12 (41%)	3 (12%)	7 (30%)
	Temporal Trend	54 (31%)	9 (18%)	18 (43%)	0 (0%)	3 (21%)	6 (21%)	11 (46%)	8 (35%)
	Other Redlists	33 (19%)	7 (14%)	8 (19%)	6 (60%)	1 (7%)	6 (21%)	8 (33%)	6 (26%)
	Other	19 (11%)	1 (2%)	3 (7%)	0 (0%)	0 (0%)	7 (24%)	4 (17%)	3 (13%)
Approach	DT based	20 (12%)	7 (14%)	3 (7%)	1 (10%)	4 (29%)	6 (21%)	1 (4%)	4 (17%)
	GEE	5 (3%)	1 (2%)	2 (5%)	0 (0%)	0 (0%)	1 (3%)	0 (0%)	1 (4%)
	GLM&LM	82 (47%)	16 (32%)	19 (45%)	7 (70%)	5 (36%)	14 (48%)	13 (54%)	14 (61%)
	GLMM&LMM	21 (12%)	6 (12%)	5 (12%)	1 (10%)	2 (14%)	5 (17%)	3 (12%)	4 (17%)
	Non-parametric	18 (10%)	2 (4%)	4 (10%)	1 (10%)	1 (7%)	5 (17%)	3 (12%)	2 (9%)
	Other	9 (5%)	4 (8%)	2 (5%)	1 (10%)	2 (14%)	3 (10%)	1 (4%)	0 (0%)
	PGCMs	64 (37%)	30 (60%)	17 (40%)	4 (40%)	4 (29%)	3 (10%)	6 (25%)	3 (13%)
Control of phylogeny	yes	116 (67%)	41 (82%)	30 (71%)	8 (80%)	9 (64%)	16 (55%)	15 (62%)	13 (57%)
	no	94 (54%)	23 (46%)	22 (52%)	7 (70%)	7 (50%)	16 (55%)	15 (62%)	15 (65%)

1175

1176 Table 3: Results of the linear mixed-effect models relating extinction risk with body size, geographical range

1177 size and habitat breadth. *df:* Degrees of freedom, *P:* p value.

Model	Number of studies / measurements / different taxa	Taxa (number of studies/measurements)	Intercept estimate (standard error)	*df*	t value	*P*
Effect sizes of Geographical range size ~ (Intercept) + Random(Study) + Random(Taxon)	22/44/9	Mammals (6/10), Birds (6/15), Reptiles (3/3), Amphibians (3/4), Fishes (2/4), Vertebrates (1/2), Insects (2/4), Other Invertebrates (1/1), Plants (1/1)	−0.463 (0.112)	23.424	−4.137	0.0004
Effect sizes of body size ~ (Intercept) + Random(Study) + Random(Taxon)	30/84/9	Mammals (9/33), Birds (10/25), Reptiles (4/9), Amphibians (3/3), Fishes (3/7), Vertebrates (1/1), Insects (1/1), Other invertebrates (1/1), Plants (3/4)	0.126 (0.104)	11.503	1.205	0.2520
Effect sizes of Habitat breadth ~ (Intercept) + Random(Study) + Random(Taxon)	14/25/6	Mammals (2/6), Birds (6/11), Reptiles (2/3), Amphibians (2/3), Vertebrates (1/1), Other invertebrates (1/1), Plants (1/1)	−0.208 (0.040)	11.622	−5.162	0.0003

1178