1   **A Type I Restriction-Modification System Associated with *Enterococcus faecium***

2   **Subspecies Separation**

4   Wenwen Huo[a], Hannah M. Adams[a], Cristian Trejo[a], Rohit Badia[a] and Kelli L. Palmer[a]*

5   [a]Department of Biological Sciences, University of Texas at Dallas, Richardson, Texas 75080

7   *Contact information for corresponding author:

8   Kelli Palmer: kelli.palmer@utdallas.edu

ABSTRACT

The gastrointestinal colonizer *Enterococcus faecium* is a leading cause of hospital-acquired infections. Multidrug-resistant (MDR) *E. faecium* are particularly concerning for infection treatment. Previous comparative genomic studies revealed that subspecies referred to as Clade A and Clade B exist within *E. faecium*. MDR *E. faecium* belong to Clade A, while Clade B consists of drug-susceptible fecal commensal *E. faecium*. Isolates from Clade A are further grouped into two sub-clades, A1 and A2. In general, Clade A1 isolates are hospital epidemic isolates whereas Clade A2 isolates are isolates from animals and sporadic human infections. Such phylogenetic separation indicates that reduced gene exchange occurs between the clades. We hypothesize that endogenous barriers to gene exchange exist between *E. faecium* clades. Restriction-modification (R-M) systems are such barriers in other microbes. We utilized bioinformatics analysis coupled with second generation and third generation deep sequencing platforms to characterize the methylome of two representative *E. faecium* strains, one from Clade A1 and one from Clade B. We identified a Type I R-M system that is Clade A1-specific, is active for DNA methylation, and significantly reduces transformability of Clade A1 *E. faecium*. Based on our results, we conclude that R-M systems act as barriers to horizontal gene exchange in *E. faecium* and propose that R-M systems contribute to *E. faecium* subspecies separation.

IMPORTANCE

Enterococcus faecium is a leading cause of hospital-acquired infections around the world. Rising antibiotic resistance in certain *E. faecium* lineages leaves fewer treatment options. The overarching aim of the attached work was to determine whether restriction-modification (R-M) systems contribute to the structure of the *E. faecium* species, wherein hospital-epidemic and non-hospital-epidemic isolates have distinct evolutionary histories and highly resolved clade

structures. R-M provides bacteria with a type of innate immunity to horizontal gene transfer

(HGT). We identified a Type I R-M system that is enriched in the hospital-epidemic clade and

determined that it is active for DNA modification activity and significantly impacts HGT. Overall,

this work is important because it provides a mechanism for the observed clade structure of *E.*

faecium as well as a mechanism for facilitated gene exchange among hospital-epidemic *E.*

faecium.

INTRODUCTION

Enterococcus faecium is a Gram-positive opportunistic pathogen that normally resides in the gastrointestinal tracts of humans and other animals (1, 2). *E. faecium* can cause life-threatening infections such as endocarditis and is among the leading causes of catheter-associated bloodstream and urinary tract infections in clinical settings (3).

Previous comparative genomic studies revealed that subspecies exist within *E. faecium* (4-7). Different names have been used by different groups to describe these clades; in this study, we use the Clade A/B nomenclature. Generally speaking, MDR *E. faecium* belong to Clade A, while Clade B consists of drug-susceptible fecal commensal *E. faecium* (8). Clade A is further split into two subclades, A1 and A2, with hospital-endemic strains generally clustering in Clade A1 and sporadic infection isolates and animal isolates generally clustering in Clade A2 (8). Specific phenotypes and genomic features are enriched in Clade A1 isolates relative to Clade A2 and B isolates (8). Specifically, Clade A1 isolates have significantly higher mutation rates, larger overall genome sizes including a larger core genome, and possess more mobile elements. On the other hand, Clade A2 possesses a larger pan-genome than Clade A1 and B, possibly reflective of the broader host origins of these strains. Given that Clade A and Clade B strains would be expected to co-mingle in certain environments (for example, in hospital and municipal sewage), the phylogenetic separation among the *E. faecium* clades suggests that they are not sharing genetic information freely because of endogenous barriers to genetic exchange.

Horizontal gene transfer (HGT) is the exchange of genetic material between cells rather than the vertical inheritance of genetic material from a parental cell. Bacteria can encode genome defense mechanisms that can act in opposition to HGT. Two examples of these mechanisms are clustered regularly interspaced short palindromic repeats (CRISPR) and associated proteins (CRISPR-Cas) systems and restriction-modification (R-M) systems. CRISPR-Cas is a dynamic

immune system that utilizes sequence complementarity between self (CRISPR RNAs) and

foreign nucleic acid to carry out its restrictive function, whereas R-M discriminates self from

foreign DNA by DNA methylation patterns. If the *E. faecium* clades encode different defense

mechanisms, they may not exchange genetic information freely, thereby facilitating and

maintaining phylogenetic separation. However, little is known about CRISPR-Cas and R-M in *E.*

faecium. Genomic analysis suggests that these systems could contribute to the observed clade

structure of *E. faecium*. For example, CRISPR-Cas systems have been identified exclusively in

Clade B *E. faecium* and in sporadic Clade A-Clade B recombinant strains (8). For R-M, a

predicted methyl-directed restriction endonuclease (REase) is enriched in Clade A2 and B *E.*

faecium genomes relative to Clade A1 genomes (8).

Here, we focused on R-M systems and their roles in regulating gene exchange in *E. faecium*

because little is known about R-M defense in this species. Moreover, there is precedent in the

literature for R-M systems contributing to bacterial clade structure, as has been observed in

Burkholderia (9) and *Neisseria* (10). Our overarching hypothesis is that the *E. faecium* clades

encode different R-M systems, thereby inhibiting genetic exchange between them. In general,

R-M systems are composed of cognate methyltransferase (MTase) and REase activities and

are classified into different types based on the specific number and types of enzymes in the

system, as well as characteristics such as methylation type and pattern, cofactor requirement,

and restriction activity (11). A MTase recognizes specific sequences in the bacterial genome

and transfers a methyl group to either an adenine or a cytosine, resulting in 6-methyladenine

(m6A), 4-methylcytosine (m4C), or 5-methylcytosine (m5C). A REase may recognize the same

sequence as a MTase and cleave that region if the sequence is unmethylated (or in some

cases, if methylated). With the activities of MTases and REases, bacteria can use R-M to

impede entry of non-self DNA.

In this study, we used single-molecule real-time (SMRT) sequencing and whole genome

bisulfite sequencing to characterize the methylomes of representative *E. faecium* strains from

Clade A1 (*E. faecium* 1,231,502; or Efm502) and Clade B (*E. faecium* 1,141,733; or Efm733).

Two unique m6A methylation patterns were identified, one in each strain. These patterns were

asymmetric and bipartite, which is characteristic of Type I R-M methylation motifs (12).

Bioinformatic analyses were performed to identify candidate genes responsible for the

methylation. A unique Type I R-M system is encoded by each strain. We have named these

systems Efa502I (for Efm502) and Efa733I (for Efm733). Expression of these candidate

systems in *E. faecalis* heterologous hosts followed by SMRT sequencing confirmed that they

are responsible for the methylation patterns observed in Efm502 and Efm733. A functional

analysis was performed in order to assess the abilities of these systems to reduce *E. faecium*

HGT by transformation. In a comparative analysis among 73 *E. faecium* genomes, we found

that Efa502I is significantly enriched among Clade A1 isolates, while the Type I R-M system of

Efm733 appears to be strain-specific. Overall, this study is a first step towards understanding

the role of R-M in regulating HGT in *E. faecium* and the potential for R-M as one mechanism for

the clade structure of *E. faecium*.

METHODS

Bacterial strains and growth conditions. The strains used in this study are shown in Table 1.

All enterococci were grown in Brain Heart Infusion (BHI) broth or agar at 37°C, unless otherwise

stated. *Escherichia coli* strains were grown in Luria Broth (LB) at 37°C and with shaking at 225

rpm unless otherwise stated. Antibiotic concentrations for enterococcal strains were as follows:

rifampin, 50 µg/mL; fusidic acid, 25 µg/mL; spectinomycin, 500 µg/mL; streptomycin, 500 µg/mL;

chloramphenicol, 15 µg/mL. Antibiotic concentrations for *E. coli* strains were as follows:

chloramphenicol, 15 µg/mL; ampicillin, 100 µg/mL. All REases were purchased from New

England Biolabs (NEB) and used per the manufacturer's instructions. PCR was performed using

Taq polymerase (NEB) or Phusion (Fisher). Sanger sequencing to validate all genetic

constructs was performed at the Massachusetts General Hospital DNA Core facility (Boston,

MA).

Isolation of genomic DNA. Enterococcal strains were cultured overnight in BHI broth prior to

genomic DNA (gDNA) extraction. The extraction was performed using a Qiagen Blood and

Tissue DNeasy Kit using a previously published protocol (13). To isolate *E. coli* gDNA, bacteria

were grown overnight in LB broth prior to extraction using either the Blood and Tissue DNeasy

kit (Qiagen) or the UltraClean Microbial DNA Isolation Kit (Qiagen) per the manufacturer's

instructions.

SMRT sequencing and methylome detection. SMRT sequencing was performed by the

Johns Hopkins Medical Institute Deep Sequencing and Microarray Core. After sequencing, the

reads were assembled into contigs and analyzed using the RS modification and motif detection

protocol in SMRT portal v1.3.3. An *in silico* control was used as a methylation baseline. SMRT

sequencing in *E. faecalis* expressing Efa502I or Efa733I was performed by the University of

Michigan sequencing core facility. Reads were mapped to the *E. faecalis* OG1RF reference

sequence (GenBank accession number NC_017316), and the methylation motifs were detected

using the RS modification and motif detection protocol in SMRT portal v.2.3.2.

Bioinformatic analysis of R-M systems in eight *E. faecium* genomes. The entire protein

complement for eight previously sequenced *E. faecium* isolates (14) was analyzed. To identify

potential MTases, the REBASE Gold Standard list (15) was used as a reference. This list is

comprised of biochemically verified MTases and REases. Each protein sequence from *E.*

faecium genomes was analyzed using BlastP against the REBASE Gold Standard list. The

protein sequences with significant (e-value <1e^{-3}) homology to REBASE Gold Standard proteins

were further filtered by protein size. If an *E. faecium* query protein length was less than half of

its subject's length, the match was removed from the prediction list. Due to the sequence

diversity of REases which complicates their bioinformatic identification (15), guilt-by-association

was used to identify full R-M systems as we have previously described (16). The proteins

encoded near candidate DNA MTases were analyzed using BLAST and Pfam for conserved

domains consistent with REase activities and/or sequence identity to confirmed REases. The

amino acid sequence of each R-M candidate was then pairwise compared among all the eight

strains to identify putative orthologs. If two protein sequences shared an amino acid identity

≥90% with query coverage ≥90%, they were considered to be orthologous.

Expression of R-M systems in *E. faecalis* heterologous hosts. Genes encoding the

specificity and methylation subunits of Efa733I (EFSG_05028-EFSG_05027) were PCR-

amplified in their entirety, including the upstream region to retain the native promoter, using

primers 733_T1A_SM_F and 733_T1A_SM_R (see Table S1 for primer sequences). The PCR

product was digested with *Not*I and ligated into *Not*I-digested pWH03 (16) using T4 DNA Ligase

(NEB), generating pHA102. pWH03 is a pLT06 derivative for expression of genes from a

previously validated neutral genomic insertion site (EF2238-EF2239) for expression (GISE) (16,

17). pHA102 constructs were then introduced into *E. coli* DH5α via heat shock for propagation

and sequence confirmation. pHA102 was electroporated into *E. faecalis* OG1SSp using a

previously described method (18). An *E. faecalis* OG1SSp derivative with a chromosomal

integration of Efa733I, referred to as OG1SSp::*efa733I*, was generated by temperature shifts

and *p*-chlorophenylalanine counterselection, as previously described (19).

Genes encoding the specificity and methylation subunits of Efa502I (EFQG_01131-

EFQG_01132) were PCR-amplified using primers 502_T1A_SM_F and 502_T1A_SM_R. The

PCR product was then TA-cloned into the pGEM-T Easy Vector (Promega) and introduced into

DH5α via heat shock to generate pGEM-SMA1. pGEM-SMA1 was then digested with *Not*I, and

the digestion reaction was used as insert for ligation into *Not*I-digested pWH03. The ligation

reaction was then introduced into DH5α, and colonies were screened for chloramphenicol

resistance and ampicillin susceptibility to ensure the pGEM backbone was not ligated into

pWH03. Once the construct, referred to as pHA103, was confirmed via Sanger sequencing, it

was introduced into electrocompetent *E. faecalis* OG1RF. An *E. faecalis* OG1RF derivative with

a chromosomal integration of Efa502I, referred to as OG1RF::*efa502I*, was generated by

temperature shifts and *p*-chlorophenylalanine counterselection. All plasmids and strains for

heterologous expression were validated by PCR and Sanger sequencing.

Generation of *E. faecium* R-M deletion mutants. Regions up- and downstream of Efa502I

and Efa733I were PCR-amplified using primers listed in Table S1, ligated into pLT06, and

transformed into *E. coli* EC1000 (20), generating pWH16 and pWH17 (Table 1). Insert

sequences were confirmed using Sanger sequencing. *E. faecium* strains were made

electrocompetent using previously a published protocol (21). 2 µg of sequence-confirmed

plasmids were electroporated into electrocompetent Efm733 and Efm502. The generation of

deletion mutants was accomplished using temperature shifts and *p*-chlorophenylalanine

counterselection, as previously described (19). The successful deletion mutants were

sequence-confirmed by PCR and Sanger sequencing.

Transformation efficiency test. Efm733, Efm502, and their respective R-M deletion mutants

were made electrocompetent using a modified version of the previously published protocol (21).

Briefly, overnight cultures were diluted 10-fold in BHI and cultured to $OD_{600nm} \sim 0.6$. The bacteria

were then pelleted and treated with filter-sterilized lysozyme buffer (10 mM Tris-HCl pH 8.0, 10

mM EDTA pH 8.0, 50 mM NaCl) supplemented with 83 µL of 2.5 KU/mL mutanolysin stock for

30 min at 37°C. The cells were then pelleted and washed three times with ice-cold filter-

sterilized electroporation buffer (0.5 M sucrose and 10% glycerol). Finally, the cells were

pelleted and resuspended in electroporation buffer and aliquoted for storage at -80°C and future

use. 1 μg pAT28 (22) was electroporated into the electrocompetent *E. faecium* cells. The counts

of total viable cells and spectinomycin-resistant cells were determined by serial dilution and

plating. The transformation efficiency was expressed as percent of transformed (spectinomycin-

resistant) cells per total viable cells. Three independent experiments were performed and the

statistical significance was assessed using the unpaired one-tailed Student's t-test.

Distribution analysis of putative R-M systems and orphan MTases. The amino acid

sequences for select R-M system and orphan MTase candidates were queried against a

collection of 73 *E. faecium* isolates previously analyzed by Lebreton *et al* (8) using BLASTP.

Any proteins which shared >90% query coverage and amino acid identity were considered

orthologs. The Fisher's exact test was used to determine if an orphan MTase or R-M system

was significantly over- or under-represented in a particular clade.

REase protection assays. To identify m5C methylation, gDNA was treated with the

methylation-sensitive REases McrBC, FspEI, and MspJI (NEB). 500 ng gDNA was incubated

with each REase at 37°C for 3 h (McrBC) or 6 h (FspEI and MspJI) followed by analysis by

electrophoresis on a 1% agarose gel with ethidium bromide.

Bisulfite sequencing. Whole-genome bisulfite sequencing libraries were constructed using the

Illumina TruSeq LT PCR FREE kit and the Qiagen EpiTect Bisulfite kit. Native DNA was isolated

as described above. Whole-genome-amplified (WGA) control DNA was generated by

amplification of native gDNA using the Qiagen REPLI-g® kit, per the manufacturer's

instructions. For bisulfite sequencing, briefly, 2 μg each of native and WGA control DNA were

fragmented using NEB fragmentase. DNA fragments ranging from 200 bp to 700 bp were gel

extracted and end-repaired. After A-tailing of DNA fragments, Illumina TruSeq adapters were

ligated. Then, the bisulfite conversion was performed using the Qiagen EpiTect Bisulfite kit, per

the manufacturer's instruction. An 8-cycle PCR enrichment with Illumina primer mix was

performed, followed by size selection and gel purification. The libraries were sequenced using

Illumina MiSeq with 2x75 bp paired-end chemistry.

Whole genome bisulfite sequencing analysis. The sequencing reads were analyzed using

Bismark (23) with additional quality control and filtering as described previously (16). Briefly, the

Illumina reads were mapped to the *in silico* bisulfite-converted references (23). Then, we

quantified the conversion rate of each mapped read by calculating the percentage of converted

C (which will result in T) to the total number of C in the reference within the mapped region. The

mapped reads with ≤80% conversion rate were filtered out from analysis (16). Next, the

coverage depth and methylation ratio were calculated for each C site. The methylation ratio was

calculated by dividing the total number of C by the coverage depth at each C site. A fully

methylated C, thus protected from bisulfite conversion, will have a methylation ratio near 1. An

unmethylated C will have a methylation ratio near 0. To identify consensus methylation motifs, C

sites with ≥0.35 methylation ratio and ≥10 coverage depth, along with the sequences of 5 bp

upstream and 5 bp downstream, were extracted. The extracted sequences were subjected for

MEME motif search (24).

Confirmation of m5C MTase activity. Primers EFSG_00659_F and EFSG_00659_R (Table

S1) were used to amplify the entire Efm733 EFSG_00659 coding region and its upstream

predicted promoter. The PCR product was then cloned into the pGEM-T Easy Vector (Promega)

per the manufacturer's instructions and transformed into *E. coli* STBL4 (Fisher) to generate

pRB01. REase digestion assays with methylation-sensitive enzymes were performed on purified

E. coli and *E. faecium* gDNA as described above.

Accession numbers. DNA sequence data generated in this study have been deposited in the

Sequence Read Archive under accession numbers PRJNA397049 (for SMRT sequencing data)

and PRJNA488088 (for Illumina bisulfite sequencing data).

RESULTS

Identification of Clade A1-specific putative Type I R-M system in *E. faecium*. We previously

reported that a Type II R-M system significantly reduces HGT via conjugation (18) and

transformation (16) in *E. faecalis*. Here, we hypothesize that the *E. faecium* clades encode

distinct R-M systems that reduce the exchange of genetic information between them. We

utilized an approach we previously developed for *E. faecalis* R-M analysis (18) to predict

potential R-M systems in eight previously sequenced *E. faecium* genomes. The 8 genomes

included 3 genomes from Clade A1, 3 genomes from Clade B, one genome from Clade A2, and

one recombinant Clade A1/B hybrid (5, 8). Because REases are difficult to identify with

bioinformatics, and MTase prediction is comparatively straightforward, as has been previously

reported by NEB (15), we first identified predicted DNA MTases in *E. faecium* genomes, and

then analyzed surrounding genes for predicted R-M-related activities. The complete list of

candidates for the eight strains is shown in Table 2.

Interestingly, we predicted at least one putative Type I R-M system for seven of the eight *E.*

faecium strains (Table 2). Type I R-M systems are multisubunit complexes comprised of a

specificity subunit (S), a methylation subunit (M), and a restriction subunit (R) (11, 25-27). The S

subunit is responsible for the specific DNA recognition motif and associates with the DNA to

bring the M and R subunits into contact. The system has two conformations: M_2S_1, which is

capable of methylating DNA based on the recognition sequence, and $R_2M_2S_1$, which is capable

of restricting DNA (27, 28). One predicted *E. faecium* Type I R-M system is comprised of highly

conserved (>90% amino acid sequence identity) M and R subunits in six of eight genomes

across both Clade A1 and Clade B (Table 2 and Dataset S1). The specificity subunit from this

system, however, is highly conserved in Clade A1 genomes but not in Clade B (Table 2 and Fig.

1a-b). S subunits possess two target recognition domains (TRDs) that determine the nucleotide

sequence the subunit binds to (29, 30). The variation in amino acid sequence between the S

subunits occurs within these TRDs (Fig 1a), suggesting that these S subunits recognize

different DNA sequences. Notably, the S subunits from Clade A1 strains are identical to each

other, indicating that they utilize the same recognition sequence.

To examine the distribution of this putative Clade A1-specific system in a larger collection of *E.*

faecium strains, we analyzed 73 *E. faecium* genomes of mostly draft status that were reported

previously (8). This list includes 15 clade B isolates, 21 clade A1 isolates, 35 clade A2 isolates,

and 2 hybrid isolates (Table S2). We selected Efm502 (Clade A1) as our representative Clade

A1 strain for this analysis and used Type I R-M sequences from this genome as references for

analysis against the broader collection of *E. faecium* strains. The M and R subunits of the

putative Clade A1-specific Type I system were detected in 51 and 52, respectively, of 73 *E.*

faecium genomes, including both Clade A and Clade B strains (Fig S1a). However, the

distribution of the S subunits varied (Fig S1a-b). The S subunit present in Efm502,

EFQG_01131, was significantly enriched within Clade A1 isolates (14/21; p-value <0.0001 using

Fisher exact test; Fig S1a) and absent from all other clades with the exception of strain

EnGen002, which is classified as a Clade A1/B hybrid strain. Interestingly, the S subunits

present in most other *E. faecium* strains are strain-specific by the strict thresholds applied here.

Given that the Efm502 S subunit is enriched in Clade A1, we hypothesize that many Clade A1

strains exchange genetic information freely with each other while exchange with other *E.*

faecium strains is restricted.

SMRT sequencing for *E. faecium* methylome analysis. We analyzed the Efm502 and

Efm733 genomes by SMRT sequencing. SMRT sequencing measures the kinetics of DNA

polymerase as it synthesizes DNA in order to identify bases that have been modified (31-33). It

has been extensively utilized for bacterial methylome analysis (34-42). With SMRT sequencing,

6-methyladenine (m6A) and 4-methylcytosine (m4C) can be easily detected with modest

sequence coverage (~25x per strand), while 5-methylcytosine (m5C) detections requires high

coverage (~250x per strand) (41, 43). Using SMRT sequencing, we identified two unique m6A

methylation motifs in Efm502 and Efm733. Efm502 possessed m6A methylation at the

underlined position of the motif 5'-RAYCNNNNNNTTRG-3' and Efm733 possessed m6A

methylation at the underlined position of the motif 5'-AGAWNNNNATTA-3' (Table 3). These

sequences are asymmetric and bipartite, which is characteristic of Type I R-M methylation (12).

Due to the coverage of our SMRT sequencing, m5C modification could not be accurately

detected. The two unique m6A methylation patterns indicate that DNA from one strain would be

recognized as foreign should it cross the strain barrier.

**Expression in heterologous hosts links methylation activity to genes in Efm502 and

Efm733.** According to our predictions (Table 2), there is only one complete Type I R-M system

encoded by each of Efm502 and Efm733. To determine if these systems are responsible for the

methylation patterns identified by SMRT sequencing, we expressed the respective S and M

subunits (EFQG_01131-01132 for Efm502 and EFSG_05028-05027 for Efm733) in the

heterologous host *E. faecalis* OG1RF or its spectinomycin/streptomycin-resistant relative

OG1SSp. Previous work in our lab had characterized the methylome of OG1RF using SMRT

and bisulfite sequencing (16). This allowed us to attribute any new methylation patterns

observed during SMRT sequencing to the *E. faecium* genes that were expressed in the OG1RF

background. SMRT sequencing of these strains detected the same methylation patterns

originally identified in Efm502 and Efm733 (Table 3). These data demonstrate that

EFQG_01131-01132 is responsible for the 5'-RAYCNNNNNNTTRG-3' methylation in Efm502

and that EFSG_05028-05027 is responsible for the 5'-AGAWNNNNATTA-3' methylation in

Efm733. Because we have confirmed the function of these genes, we have named them

Efa502I and Efa733I, which is consistent with the R-M system nomenclature convention

established by New England Biolabs (12).

Efa502I and Efa733I reduce transformation efficiency in *E. faecium*. To determine whether

the Type I R-M systems in Efm733 and Efm502 actively defend against exogenous DNA, we

constructed null strains (Efm733ΔRM and Efm502ΔRM; Table 1) and evaluated their

transformation efficiencies relative to their wild-type parent strains. Here, we utilized the broad

host range plasmid pAT28 (Table 1) (22). pAT28 sequence has motifs recognized by the Type I

R-M systems in both wild-type Efm733 (1 occurrence) and Efm502 (1 occurrence). The

transformation of pAT28 into Efm733 and Efm502 served as a baseline for the experiment. If

Efa733I and Efa502I are active, we expect to see higher pAT28 transformation efficiencies into

Efm733ΔRM and Efm502ΔRM, respectively. Indeed, we observed significantly higher

transformation efficiencies into Type I R-M system null strains (Fig 2; p-value<0.05 using one-

tailed Student's t-test). These data demonstrate that the Type I R-M systems in Efm733 and

Efm502 actively function as mechanisms of genome defense.

m5C methylation occurs in Efm733. As described previously, our SMRT sequencing had

insufficient coverage depth for m5C methylome characterization. Hence, we used REase

protection assays with commercially available methylation-sensitive REases to query the

presence of m5C methylation in our eight *E. faecium* strains. Table 4 summarizes the

recognition sequences and modifications of the enzymes used in this study. Only Efm733

showed evidence of cytosine modification, as it was digested by MspJI (Table 4; Fig S2).

To determine the exact cytosine methylation motif present in Efm733, gDNA was subjected to

whole-genome bisulfite sequencing. Whole genome amplified (WGA) DNA was used as

negative control since WGA removes all modifications. During bisulfite treatment, cytosine

bases are converted to thymine unless they are protected by either m4C or m5C methylation.

Additionally, our lab has previously published a method of distinguishing between m4C and

m5C methylation using thymine conversion ratios of sequencing reads after bisulfite treatment

(44). m5C methylation is sufficient to protect the cytosine residue completely from bisulfite

conversion, so that most sequencing reads at that position contain the original cytosine base.

However, m4C methylation provides only partial protection from bisulfite conversion, so a

thymine conversion rate of 0.5 at a particular position within the sequencing reads suggests the

presence of m4C methylation. Bisulfite conversion and subsequent sequencing revealed that

Efm733 possesses m5C modification at the motif 5'-R^{m}CCGGY-3' (Table 5, Fig 3, and Fig S3;

the methylation occurs at the underlined position), which overlaps the MspJI recognition site (5'-

CNNR-3') and hence supports the evidence of methylation obtained from the MspJI digestion

assays. Based on the cytosine conversion ratio of close to 1.0, m5C modification is supported,

which is consistent with why it was not detected by our SMRT sequencing.

EFSG_00659 is responsible for m5C methylation in Efm733. Based on the bioinformatics

analyses, we hypothesized that EFSG_00659 was responsible for the m5C methylation found in

Efm733 (Table 2). EFSG_00659 possessed no homologs in the other seven strains analyzed,

making it a good candidate for the unique methylation found in Efm733. We queried the

EFSG_00659 protein sequence against the REBASE gold standard list and identified that it has

high sequence similarity to M.AvaIX, M.VchO395I and M.VchAI (e-value ≤ 3e^{-125}; recognition

sites are 5'-RCCGGY-3'). Interestingly, BLASTP identified no significant hits when

EFSG_00659 was queried against the larger collection of 73 *E. faecium* genomes, indicating its

unique presence in Efm733.

In order to link EFSG_00659 with the m5C methylation identified during bisulfite sequencing, we

expressed it in the heterologous host *E. coli* STBL4 and performed an REase protection assay.

The REase AgeI recognizes the motif 5'-A<u>C</u>CGGT-3', and its enzymatic activity is blocked if

m5C methylation is present at the underlined position. This motif overlaps the m5C methylation

motif in Efm733 identified by bisulfite sequencing. If the motif is methylated, DNA will be

protected against digestion. We cloned EFSG_00659 into the vector pGEM-T and transformed it

into *E. coli* STBL4, generating strain *E. coli* STBL4(pRB01). Genomic DNA from Efm733,

STBL4(pGEM-T), and STBL4(pRB01) was treated with AgeI per the manufacturer's instructions.

EcoRI was used as a positive control for digestion. Figure 4 shows representative results of the

digestions on a 1% agarose gel. As expected, Efm733 was digested by EcoRI and protected

against digestion from AgeI. STBL4(pGEM-T) was digested by both EcoRI and AgeI, indicating

that the original host and empty vector pGEM-T did not possess the appropriate m5C

methylation. STBL4(pRB01) was protected against digestion by AgeI, demonstrating that

EFSG_00659 is responsible for the 5'-R^{m}CCGGY-3' methylation found in Efm733.

DISCUSSION

In this study, we used a combination of genomic and genetic approaches to identify a functional

Type I R-M system that is enriched in Clade A1 *E. faecium* and that significantly alters

transformability of a model Clade A1 strain. We propose that this R-M system impacts HGT

rates among *E. faecium* mixed-clade communities, thereby helping to maintain the observed

phylogenetic structure of *E. faecium* and facilitating HGT specifically among Clade A1 strains.

Mixed communities of *E. faecium* clades are expected to occur in environments where healthy

and ill human hosts, human and animal hosts, and/or the feces of any of these hosts co-mingle

(i.e. in sewage). In future studies, we plan to assess the impact of R-M on conjugative plasmid

transfer, which is a major mode of HGT in enterococci and was not assessed in our current

study.

An interesting observation from our study is the sequence diversity of Type I S subunits

encoded within *E. faecium* Type R-M systems having nearly identical R and M subunits (Fig

S1a-b). After further investigation into those alignments, we found that those S subunits sharing

50-70% overall amino acid sequence identities possess sequence diversity within one TRD

domain, where the other TRD domain and the central conserved domain are conserved. This

suggests that these systems share partial recognition sequences. Previous research has

reported that the diversification of Type I R-M recognition sequences is driven by TRD

exchanges, permutation of the dimerization domain, and circular permutation of TRDs (45). Our

observation suggests that TRD recombination and reorganization events occur for *E. faecium*

Type I R-M systems outside Clade A1. Future studies will use genomics to further explore the

relationship between S subunit sequence diversity and its impact on *E. faecium* methylomes

and inter-strain and inter-clade HGT.

Acknowledgments

This work was supported by Public Health Service grants K22AI099088 and R01AI116610 to

K.L.P.

References

1.   Noble CJ. 1978. Carriage of group D streptococci in the human bowel. J Clin Pathol 31:1182-6.

2.   Chenoweth C, Schaberg D. 1990. The epidemiology of enterococci. Eur J Clin Microbiol Infect Dis
9:80-9.

3.   Agudelo Higuita NI, Huycke MM. 2014. Enterococcal Disease, Epidemiology, and Implications for
Treatment. *In* Gilmore MS, Clewell DB, Ike Y, Shankar N (ed), Enterococci: From Commensals to
Leading Causes of Drug Resistant Infection, Boston.

4.   Louis E, Galloway-Peña J, Roh JH, Latorre M, Qin X, Murray BE. 2012. Genomic and SNP Analyses
Demonstrate a Distant Separation of the Hospital and Community-Associated Clades of
Enterococcus faecium. PLoS ONE 7:e30187.

5.   Palmer KL, Godfrey P, Griggs A, Kos VN, Zucker J, Desjardins C, Cerqueira G, Gevers D, Walker S,
Wortman J, Feldgarden M, Haas B, Birren B, Gilmore MS. 2012. Comparative genomics of
enterococci: variation in Enterococcus faecalis, clade structure in E. faecium, and defining
characteristics of E. gallinarum and E. casseliflavus. MBio 3:e00318-11.

6.   Willems RJL, Top J, van Schaik W, Leavis H, Bonten M, Siren J, Hanage WP, Corander J. 2012.
Restricted Gene Flow among Hospital Subpopulations of Enterococcus faecium. mBio 3.

7.   van Schaik W, Top J, Riley DR, Boekhorst J, Vrijenhoek JE, Schapendonk CM, Hendrickx AP,
Nijman IJ, Bonten MJ, Tettelin H, Willems RJ. 2010. Pyrosequencing-based comparative genome
analysis of the nosocomial pathogen Enterococcus faecium and identification of a large
transferable pathogenicity island. BMC Genomics 11:239.

8.   Lebreton F, van Schaik W, McGuire AM, Godfrey P, Griggs A, Mazumdar V, Corander J, Cheng L,
Saif S, Young S, Zeng Q, Wortman J, Birren B, Willems RJ, Earl AM, Gilmore MS. 2013. Emergence
of epidemic multidrug-resistant Enterococcus faecium from animal and commensal strains.
MBio 4.

9.   Nandi T, Holden MTG, Didelot X, Mehershahi K, Boddey JA, Beacham I, Peak I, Harting J,
Baybayan P, Guo Y, Wang S, How LC, Sim B, Essex-Lopresti A, Sarkar-Tyson M, Nelson M, Smither
S, Ong C, Aw LT, Hoon CH, Michell S, Studholme DJ, Titball R, Chen SL, Parkhill J, Tan P. 2015.
Burkholderia pseudomalleisequencing identifies genomic clades with distinct recombination,
accessory, and epigenetic profiles. Genome Research 25:129-141.

10.   Budroni S, Siena E, Hotopp JCD, Seib KL, Serruto D, Nofroni C, Comanducci M, Riley DR,
Daugherty SC, Angiuoli SV, Covacci A, Pizza M, Rappuoli R, Moxon ER, Tettelin H, Medini D. 2011.
Neisseria meningitidis is structured in clades associated with restriction modification systems
that modulate homologous recombination. Proceedings of the National Academy of Sciences
108:4494-4499.

11.   Tock MR, Dryden DT. 2005. The biology of restriction and anti-restriction. Curr Opin Microbiol
8:466-72.

12.   Roberts RJ, Belfort M, Bestor T, Bhagwat AS, Bickle TA, Bitinaite J, Blumenthal RM, Degtyarev S,
Dryden DT, Dybvig K, Firman K, Gromova ES, Gumport RI, Halford SE, Hattman S, Heitman J,
Hornby DP, Janulaitis A, Jeltsch A, Josephsen J, Kiss A, Klaenhammer TR, Kobayashi I, Kong H,
Kruger DH, Lacks S, Marinus MG, Miyahara M, Morgan RD, Murray NE, Nagaraja V, Piekarowicz
A, Pingoud A, Raleigh E, Rao DN, Reich N, Repin VE, Selker EU, Shaw PC, Stein DC, Stoddard BL,
Szybalski W, Trautner TA, Van Etten JL, Vitor JM, Wilson GG, Xu SY. 2003. A nomenclature for
restriction enzymes, DNA methyltransferases, homing endonucleases and their genes. Nucleic
Acids Res 31:1805-12.

13.   Adams HM, Li X, Mascio C, Chesnel L, Palmer KL. 2015. Mutations associated with reduced
surotomycin susceptibility in Clostridium difficile and Enterococcus species. Antimicrob Agents
Chemother 59:4139-47.

14.   Palmer KL, Carniol K, Manson JM, Heiman D, Shea T, Young S, Zeng Q, Gevers D, Feldgarden M,
Birren B, Gilmore MS. 2010. High-quality draft genome sequences of 28 Enterococcus sp.
isolates. J Bacteriol 192:2469-70.

15.   Roberts RJ, Vincze T, Posfai J, Macelis D. 2015. REBASE—a database for DNA restriction and
modification: enzymes, genes and genomes. Nucleic Acids Research 43:D298-D299.

16.   Huo W, Adams HM, Zhang MQ, Palmer KL. 2015. Genome Modification in Enterococcus faecalis
OG1RF Assessed by Bisulfite Sequencing and Single-Molecule Real-Time Sequencing. J Bacteriol
197:1939-51.

17.   DebRoy S, van der Hoeven R, Singh KV, Gao P, Harvey BR, Murray BE, Garsin DA. 2012.
Development of a genomic site for gene integration and expression in Enterococcus faecalis.
Journal of Microbiological Methods 90:1-8.

18.   Price VJ, Huo W, Sharifi A, Palmer KL. 2016. CRISPR-Cas and Restriction-Modification Act
Additively against Conjugative Antibiotic Resistance Plasmid Transfer in Enterococcus faecalis.
mSphere 1.

19.   Thurlow LR, Thomas VC, Hancock LE. 2009. Capsular polysaccharide production in Enterococcus
faecalis and contribution of CpsF to capsule serospecificity. J Bacteriol 191:6203-10.

20.   Leenhouts K, Buist G, Bolhuis A, ten Berge A, Kiel J, Mierau I, Dabrowska M, Venema G, Kok J.
1996. A general system for generating unlabelled gene replacements in bacterial chromosomes.
Mol Gen Genet 253:217-24.

21.   Bhardwaj P, Ziegler E, Palmer KL. 2016. Chlorhexidine Induces VanA-Type Vancomycin
Resistance Genes in Enterococci. Antimicrob Agents Chemother 60:2209-21.

22.   Trieu-Cuot P, Carlier C, Poyart-Salmeron C, Courvalin P. 1990. A pair of mobilizable shuttle
vectors conferring resistance to spectinomycin for molecular cloning inEscherichia coliand in
Gram-positive bacteria. Nucleic Acids Research 18:4296-4296.

23.   Krueger F, Andrews SR. 2011. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. Bioinformatics 27:1571-2.

24. Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble WS. 2009.
MEME SUITE: tools for motif discovery and searching. Nucleic Acids Res 37:W202-8.

25. Luria SE, Human ML. 1952. A nonhereditary, host-induced variation of bacterial viruses. J
Bacteriol 64:557-69.

26. Bertani G, Weigle JJ. 1953. Host controlled variation in bacterial viruses. J Bacteriol 65:113-21.

27. Murray NE. 2000. Type I restriction systems: sophisticated molecular machines (a legacy of
Bertani and Weigle). Microbiol Mol Biol Rev 64:412-34.

28. Taylor I, Patel J, Firman K, Kneale G. 1992. Purification and biochemical characterisation of the
EcoR124 type I modification methylase. Nucleic Acids Res 20:179-86.

29. Gough JA, Murray NE. 1983. Sequence diversity among related genes for recognition of specific
targets in DNA molecules. J Mol Biol 166:1-19.

30. Gann AA, Campbell AJ, Collins JF, Coulson AF, Murray NE. 1987. Reassortment of DNA
recognition domains and the evolution of new specificities. Mol Microbiol 1:13-22.

31. Clarke J, Wu H-C, Jayasinghe L, Patel A, Reid S, Bayley H. 2009. Continuous base identification for
single-molecule nanopore DNA sequencing. Nature Nanotechnology 4:265-270.

32. Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A,
Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dewinter A, Dixon J, Foquet M,
Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S,
Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R,
Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Vieceli J, Wegener J, Wu D, Yang A,
Zaccarin D, et al. 2009. Real-time DNA sequencing from single polymerase molecules. Science
323:133-8.

33. Korlach J, Bjornson KP, Chaudhuri BP, Cicero RL, Flusberg BA, Gray JJ, Holden D, Saxena R,
Wegener J, Turner SW. 2010. Real-time DNA sequencing from single polymerase molecules.
Methods Enzymol 472:431-55.

34. Clark TA, Murray IA, Morgan RD, Kislyuk AO, Spittle KE, Boitano M, Fomenkov A, Roberts RJ,
Korlach J. 2012. Characterization of DNA methyltransferase specificities using single-molecule,
real-time DNA sequencing. Nucleic Acids Research 40:e29-e29.

35. Fang G, Munera D, Friedman DI, Mandlik A, Chao MC, Banerjee O, Feng Z, Losic B, Mahajan MC,
Jabado OJ, Deikus G, Clark TA, Luong K, Murray IA, Davis BM, Keren-Paz A, Chess A, Roberts RJ,
Korlach J, Turner SW, Kumar V, Waldor MK, Schadt EE. 2012. Genome-wide mapping of
methylated adenine residues in pathogenic Escherichia coli using single-molecule real-time
sequencing. Nature Biotechnology 30:1232-1239.

36. Murray IA, Clark TA, Morgan RD, Boitano M, Anton BP, Luong K, Fomenkov A, Turner SW,
Korlach J, Roberts RJ. 2012. The methylomes of six bacteria. Nucleic Acids Res 40:11450-62.

37. Bendall ML, Luong K, Wetmore KM, Blow M, Korlach J, Deutschbauer A, Malmstrom RR. 2013.
Exploring the Roles of DNA Methylation in the Metal-Reducing Bacterium Shewanella oneidensis
MR-1. Journal of Bacteriology 195:4966-4974.

38. Davis BM, Chao MC, Waldor MK. 2013. Entering the era of bacterial epigenomics with single
molecule real time DNA sequencing. Current Opinion in Microbiology 16:192-198.

39. Kozdon JB, Melfi MD, Luong K, Clark TA, Boitano M, Wang S, Zhou B, Gonzalez D, Collier J, Turner
SW, Korlach J, Shapiro L, McAdams HH. 2013. Global methylation state at base-pair resolution of
the Caulobacter genome throughout the cell cycle. Proceedings of the National Academy of
Sciences 110:E4658-E4667.

40. Richardson PM, Lluch-Senar M, Luong K, Lloréns-Rico V, Delgado J, Fang G, Spittle K, Clark TA,
Schadt E, Turner SW, Korlach J, Serrano L. 2013. Comprehensive Methylome Characterization of
Mycoplasma genitalium and Mycoplasma pneumoniae at Single-Base Resolution. PLoS Genetics
9:e1003191.

41. Roberts RJ, Carneiro MO, Schatz MC. 2013. The advantages of SMRT sequencing. Genome
Biology 14.

42. Krebes J, Morgan RD, Bunk B, Spröer C, Luong K, Parusel R, Anton BP, König C, Josenhans C,
Overmann J, Roberts RJ, Korlach J, Suerbaum S. 2014. The complex methylome of the human
gastric pathogen Helicobacter pylori. Nucleic Acids Research 42:2415-2432.

43. Flusberg BA, Webster DR, Lee JH, Travers KJ, Olivares EC, Clark TA, Korlach J, Turner SW. 2010.
Direct detection of DNA methylation during single-molecule, real-time sequencing. Nat Methods
7:461-5.

44. Huo W. 2017. Enterococcus faecalis Genome Defense Systems and Their Impact on Conjugative
Antibiotic Resistance Plasmid Transfer. 10970358. The University of Texas at Dallas, Ann Arbor.

45. Loenen WAM, Dryden DTF, Raleigh EA, Wilson GG. 2013. Type I restriction enzymes and their
relatives. Nucleic Acids Research 42:20-44.

46. Gold OG, Jordan HV, van Houte J. 1975. The prevalence of enterococci in the human mouth and
their pathogenicity in animal models. Arch Oral Biol 20:473-7.

Table 1. Strains used in this study.

Strain	Description	Reference
Enterococcus faecium		
1,231,502	Clade A1 isolate; also referred to as Efm502	(14)
1,230,933	Clade A1 isolate	(14)
1,231,410	Clade A1 isolate	(14)
1,231,408	Hybrid Clade A1/B isolate	(14)
1,231,501	Clade A2 isolate	(14)
1,141,733	Clade B isolate; also referred to as Efm733	(14)
Com12	Clade B isolate	(14)
Com15	Clade B isolate	(14)
Efm733ΔRM	Efm733 with deletion of Efa733I (EFSG_05027-29)	This study
Efm502ΔRM	Efm502 with deletion of Efa502I (EFQG_01130-32)	This study
Enterococcus faecalis		
OG1RF	Rifampicin- and fusidic acid-resistant derivative of *E. faecalis* OG1	(46)
OG1SSp	Streptomycin- and spectinomycin-resistant derivative of *E. faecalis* OG1	(46)
OG1RF::*efa502I*	OG1RF with Efa502I inserted at GISE site	This study
OG1SSp::*efa733I*	OG1SSp with Efa733I inserted at GISE site	This study
Escherichia coli		
EC1000	Plasmid propagation host	(20)
DH5α	Plasmid propagation host	
STBL4	Plasmid propagation host	Fisher
STBL4(pGEM-T)	STBL4 with pGEM-T Easy	This study
STBL4(pRB01)	STBL4 with pGEM-T Easy vector containing EFSG_00659	This study
Plasmids		
pGEM-T Easy	Commercial plasmid for gene propagation	Promega
pLT06	Temperature-sensitive plasmid	(19)
pAT28	Shuttle vector for *E. faecalis*	(22)
pWH03	Used for gene insertion in the enterococci	(16)
pRB01	pGEM-T Easy Vector with EFSG_00659	This study
pHA102	pWH03 containing NotI-digested fragments of Efa733I M and S loci (EFSG_05028-9)	This study
pHA103	pWH03 containing NotI-digested fragments of Efa502I M and S loci (EFQG_01131-2)	This study
pWH16	pLT06 containing fragments from upstream and downstream of Efa502I	This study
pWH17	pLT06 containing fragments from upstream and downstream of Efa733I	This study

562 **Table 2. Distribution of predicted DNA MTases and R-M Systems*.**

Strain name		1,230,933	1,231,410	1,231,502	1,231,501	1,231,408	1,141,733	Com12	Com15
Clade		A1	A1	A1	A2	Hybrid A1/B	B	B	B
Type I Restriction Modification System	Specificity	EFPG_01270	EFTG_00783	EFQG_01131		EFUG_01527	EFSG_05028		EFWG_02518
	Modification	EFPG_01269	EFTG_00782	EFQG_01132		EFUG_01526	EFSG_05029		EFWG_02519
	Restriction	EFPG_01271	EFTG_00784	EFQG_01130		EFUG_01528	EFSG_05027		EFWG_02517
Type I Restriction Modification System	Specificity	EFPG_05435 and EFPG_05434	EFTG_02031						
	Modification	EFPG_05433	EFTG_02032						
	Restriction	EFPG_05436	EFTG_02030						
Type I Restriction Modification System	Specificity				EFRG_00106				
	Modification				EFRG_00107				
	Restriction				EFRG_00105				
Type II Methyltransferase		EFPG_01821	EFTG_02364; EFTG_02653	EFQG_01609, EFQG_02270			EFSG_00659		
Unspecified Methyltransferases			EFTG_02333		EFRG_02239	EFUG_01512		EFVG_00502	

563 *Loci that are in black are strain specific. Loci that are the same color are conserved in their protein sequences based on a >90% sequence
564 identity threshold. An empty cell indicates that the system was not detected.

565   **Table 3. SMRT Sequencing results.**

Strain	Motif (5' -> 3')	Modified Position	Type	% Motif Detected	# of Motifs Detected	# of Motifs in Genome	Mean Motif Coverage
Efm502	RAYCNNNNNNTTRG	2	m6A	98.8	905	916	53.7
	CYAANNNNNNGRTY	4	m6A	97.9	897	916	54.5
Efm733	AGAWNNNNATTA	1	m6A	78.5	278	354	22.3
	TAATNNNNWTCT	3	m6A	73.2	259	354	23.6
OG1RF:: *efa502I*	RAYCNNNNNNTTRG	2	m6A	99.9	795	796	162.8
	CYAANNNNNNGRTY	4	m6A	99.0	788	796	163.2
OG1SSp:: *efa733I*	AGAWNNNNATTA	1	m6A	99.0	323	326	180.4
	TAATNNNNWTCT	3	m6A	98.5	321	326	180.8

567 **Table 4. Methylation-sensitive REase digestion reaction results[1].**

REase	Recognition sequence	Recognition methylation	Strains Digested
McrBC	5'-R^{m}C(N$_{40}$-N$_{3000}$)R^{m}C-3'	m5C, m4C	0/8
FspEI	5'-C^{m}C-3'	m5C	0/8
MspJI	5'-^{m}CNNR-3'	m5C	Efm733

568 [1]Recognition sequences and methylation patterns were retrieved from NEB.

569 **Table 5. Bisulfite sequencing results.**

methylation motif	methylation position	methylation type	# motifs detected[a]	# motifs in genome	# motif pass cvg filter	average methylation ratio[b]	mean motif coverage[c]
733 5'-RCCGGY-3'	2	m5C	748	780	750	96.5% (6.2%)	61.5X

570 [a]# motifs detected is defined as motifs with coverage depth more than 10X and methylation ratio >=0.35.
571 [b]Average methylation ratio is defined as the mean of the methylation ratio from all motifs in the genome.

a).

b).

Figure 1. Conservation and variability of S subunits. The protein sequences of

predicted S subunits from 6 (out of 8) representative *E. faecium* genomes were pairwise

aligned using MacVector. The alignment is shown in (a). Central and distal conserved domain was identified based on sequence homology and labeled in red (30). The target recognition domains were interpreted based on conserved domains as labeled in green. The percent identity of each pair is shown in (b). White to red: low to high percent identities. Each number represents percent identity of one protein sequence (row name) to another (column name).

Figure 2. Type I R-M systems reduce transformability of Efm733 and Efm502.
Three independent transformation experiments were performed. There is a statistical difference between the transformation efficiency of pAT28 into wild type and R-M null strains of Efm733 and Efm502. EOT: Efficiency of transfer. *: $p < 0.05$.

Figure 3. Bisulfite sequencing results for Efm733. a) Efm733 gDNA and its whole genome amplified (WGA) control were bisulfite-converted, and the methylation ratio for each cytosine site was calculated. Efm733 gDNA has cytosines with methylation ratios near 100%, indicating the presence of m5C methylation. b) All sites with ≥0.35 methylation ratio from native gDNA samples were extracted, together with 5 bp upstream and 5 bp downstream. The sequences are subjected for consensus motif analysis using MEME. The consensus motif identified by this analysis is shown.

Figure 4. EFSG_00659 confers protection against AgeI digestion. AgeI digestion reactions were analyzed by agarose gel electrophoresis with ethidium bromide staining. Bacterial gDNA was used as substrate for REase reactions. Expression of the Efm733 gene EFSG_00659 in *E. coli* STBL4 protects *E. coli* gDNA from AgeI digestion. EcoRI is a positive control for digestion. -, no enzyme added.