The melanoma microenvironment dissected through high throughput multiplexing: functional meaning of brisk/non brisk patterns and regression.

<u>Running title</u>: TILs activation in brisk/non-brisk patterns

Francesca Maria Bosisio* (1)(2), Asier Antoranz* (3)(4), Maddalena Maria Bolognesi (2), Lukas Marcelis (1), Clizia Chinello (2), Jasper Wouters (1), Fulvio Magni (2), Leonidas Alexopoulos (3)(4), Giorgio Cattoretti (2), Joost van den Oord (1)

 (1) Laboratory of Translational Cell and Tissue Research, KUL, Leuven, Belgium

(2) Pathology, Department of Medicine & Surgery, Università degli studi di Milano-Bicocca, Milan, Italy

(3) ProtATonce Ltd, Athens, Greece

(4) National Technical University of Athens, Greece

* These authors contributed equally to this work

Francesca Maria Bosisio, MD, PhD

Laboratory of Translational Cell and Tissue Research, KUL, O&N4, Herestraat 49, 3000, Leuven,

Belgium.

+393420342028

f.bosisio1@gmail.com

Abstract

In melanoma, the lymphocytic infiltrate is a prognostic parameter classified morphologically in "brisk", "non-brisk" and "absent". The meaning of "brisk" is "active, energetic", implying a functional connotation which has never been proved. Recently, it has been shown that not all brisk infiltrates have a good prognosis, that lymphocytic populations can be very heterogeneous, and that anti-PD-1 immunotherapy supports activated T cells. Here, we characterize the immune landscape in primary melanoma by high-dimensional single cell multiplex analysis in tissue sections (MILAN technique) followed by image analysis, RT-PCR and shotgun proteomics. We observed that the brisk and non-brisk patterns of tumor infiltrating lymphocytes (TILs) in primary melanoma are heterogeneous functional categories that can be further sub-classified according to the functional status of their lymphocytes as active, transitional or exhausted. We found that only activation of TILs correlates with spontaneous regression, and that the good prognosis of melanomas with a brisk pattern could be due to the fact that in these case the Tcy that are in contact with the melanoma cells are still active, and consequently the melanoma is still under immune control at the moment of surgical excision. The main inflammatory cell subpopulations that are present in the microenvironment associated with activation and exhaustion and their spatial relationships are described using neighbourhood analysis. Finally, we show a bioinformatic pipeline that, starting from common immunofluorescence stainings, can transform the tissue into a digitalized image, which represents the starting point for multiple deeper levels of analysis.

Immune-based therapies have achieved long term survival in metastatic melanoma, but checkpoint inhibitors as monotherapy yield response rates between 15-40%[1,2,3]. Low response rates, possible severe adverse effects and high cost of these drugs have prompted the search for predictive biomarkers. Recent studies have found that durable response is associated with molecular signatures enriched in immune-related genes and heavy CD8+ T cell infiltration of the tumour at the baseline[4,5,6,7,8,9,10,11]. The lymphocytic infiltrate in melanoma is a prognostic parameter reported by the pathologist as patterns of tumour-infiltrating lymphocytes (TILs). The "brisk" pattern (diffuse or complete peripheral TILs infiltration) has a better prognosis compared to the "non-brisk" (tumour areas with TILs alternate with areas without TILs) or to the "absent" pattern (no TILs or no contact with melanoma cells)[12,13,14]. There are multiple pitfalls in the purely morphological evaluation of TILs[15], but the most important one is that morphology alone cannot determine their activation status. The meaning of the word "brisk" according to the dictionary is "active, energetic", a definition implying a functional connotation starting from a morphological evaluation. Surprisingly, this functional connotation has never been proved. The contact between cytotoxic lymphocytes (Tcy) and melanoma does not always lead to tumour eradication but, due to immune modulation, can also result in Tcy inactivation. These "exhausted" Tcy would still be morphologically present, indistinguishable from active lymphocytes. Moreover, the morphological[16] and functional[17] side of the tumour microenvironment has been separately investigated, but integration of both types of data is still lacking.

In the latest years, several methods for single cell-analysis have been implemented in order to obtain a high-resolution landscape of the tumour microenvironment[18]. According to a recent review[19], high resolution means to characterize not only the immune infiltrate but also to define the spatial distribution of each component within it, which allows to make inferences about cell-cell interactions. Nevertheless, most of these methods rely on dissociation of the cells from fresh material, an impractical option in primary melanomas, nowadays diagnosed at early stage, with very limited material, and resulting in loss of the knowledge of the spatial distribution of each component within it. Moreover, several studies for predictive biomarkers relies on peripheral blood[20]. Though, the functional status of circulating inflammatory cells can completely change entering the tumour site[21]. Intuitively, it is the behaviour of the inflammatory cells in the surroundings of the tumour that will make the difference in terms of prognosis and response to therapy.

Here, we characterize the immune landscape at single cell-level in primary melanoma based on a panel of 39 immune markers applied on one single tissue section through a high-dimensional multiplexing method[22], an RT-PCR expression evaluation and a shotgun proteomic analysis. This approach allowed us a) to further categorize the brisk and non-brisk morphological patterns of TILs into three functional categories; b) to define the correlation between T-cell activation and spontaneous melanoma regression; c) to investigate the most important inflammatory subpopulations involved in TILs exhaustion (Figure 1).

Material and methods

<u>Sample Description</u>

Twenty-nine invasive primary cutaneous melanomas from the Department of Pathology of the University Hospitals Leuven (KU Leuven), Belgium, were classified based on the H&E staining according to the pattern of the inflammatory infiltrate into brisk (6 cases) and non-brisk (23 cases). According to their subtype, 24 superficial spreading melanomas, 3 nodular melanomas, and 2 lentigo maligna melanoma were included.

TMA construction

Tissue Micro Arrays (TMAs) were constructed with the GALILEO CK4500 (Isenet Srl, Milan, Italy). For each patient, one to five representative regions of interest were sampled according to the size of the specimen and the morphological heterogeneity both of the melanoma and of the distribution of the infiltrate. According to the morphological TILs pattern, we sampled brisk areas (i.e. 6 brisk areas in melanomas with brisk TILs pattern, termed "brisk in brisk" and 10 brisk areas in melanomas with non-brisk TILs pattern, termed "brisk in non-brisk") and non-brisk areas (i.e. 15 non-brisk areas in non-brisk melanomas, termed "non-brisk in non-brisk"); in addition, areas showing "early regression" (7), " late regression" (5) and "no regression" (17) according to current morphological criteria[33] were sampled. After processing, cutting and staining of the TMA blocks, a total of 60 cores were available for analysis.

Multiplex-Stripping Immunofluorescence

The multiplex-stripping method was performed according to the MILAN technique, as published in Cattoretti G, et al[22]. The antibodies used for the multiplex staining are listed in the Extended Data Table 2. Slides were scanned, and images acquired with the Hamamatsu Nanozoomer S60 scanner (Nikon, Italia) equipped with a Nikon 20X/0.75 lambda objective. Complete removal of previous layers was monitored by: a) checking for unexpected staining consistent with the subcellular and tissue distribution of a previously detected marker; and b) spurious co-clustering of unrelated molecules in the hierarchical clustering images (see below).

Image Pre-processing

Fiji/ImageJ (version 1.51u) were used to pre-process the images (File format: from. ndpi to.tif 8/16 bit, grayscale). Registration was done through the Turboreg and MultiStackReg plugins, by aligning the DAPI channels of different rounds of staining, saving the coordinates of the registration as Landmarks and applying the landmarks of the transformation to the other channels. Registration was followed by autofluorescence subtraction (Image process → subtract), previously acquired in a dedicated channel, for FITC, TRITc and Pacific Orange. A macro was written in Fiji/ImageJ and used for the TMA segmentation into single images. Cell segmentation, mask creation, and single cell measurements were done with a custom pipeline using CellProfiler (version 2014-07-23T17:45:00 6c2d896). Quality Control (QC) over the Mean Fluorescence Intensity (MFI) values was performed using feature and sample selection. In short, those cells that did not have expression in at least three markers, and those markers that were not expressed in at least 1% of the samples were removed. MFIs were further normalized to Z-scores as recommended in Caicedo JC, et al[23]. Z-scores were trimmed between -5 and +5 to avoid a strong influence of any possible outliers in the downstream analysis. The correlation between the different markers was calculated using Pearson's correlation coefficient.

Functional analysis of TILs

We selected two activation (CD69, OX40) and two exhaustion (TIM3, LAG3) markers after literature review and preliminary testing of the antibody performance on control FFPE under the conditions of the multiplex protocol. The expression levels of these markers were measured selectively on CD8+ lymphocytes using a first mask focused only on CD8+ cells. Principal Component Analysis (PCA) was applied over the expression values to evaluate the functional structure of the data and to assign an activation value in the [-1, 1] range to each cell (Supplementary Information 1, Extended Data Figure 5). Briefly, principal components (PCs) 2 and 3 were used as the rotation matrix revealed that PC1 contained all the markers in the same direction

(same sign). The point of maximum activation (Activation = 1), was defined where the projected value of CD69 (marker of activation) over PCs 2 and 3 was at the maximum while the point of maximum exhaustion (Activation = -1) where the projected value of TIM3 (marker of exhaustion) over PCs 2 and 3 was at the maximum (Figure 2). The gradient of transition was defined between the previously defined points and the centroid of the projected dataset. Pairwise t-tests with pooled standard deviation (sd) were used to find significant differences in the level of activation of the images regarding multiple histopathologic parameters (brisk/non-brisk infiltrate, regression, number of lymphocytes, ulceration, Breslow thickness, mitoses, subtype of melanoma). For this purpose, the activation level of each image was represented by the mean of its cells while the intrinsic degree of heterogeneity was captured by the sd. Continuous values of the histopathological parameters were fitted using linear models instead. P-values were adjusted for multiple comparisons using the holm method. A cut-off of 0.05 was used as significance threshold for the adjusted p-values. Images were further classified into: 'Active', 'Transition', and 'Exhausted' (from now on, indicating the core status) using one-tailed t-tests comparing the distribution of the activation values in a specific image versus the background distribution (combination of all images). P-values were adjusted using the False Discovery Rate (fdr) method. A cut-off value of 0.001 over the adjusted p-values was used as classification threshold.

Identification of cell subpopulations

To evaluate the cell subpopulations, a second mask based on the DAPI nuclear staining contour expanded by 5 pixels was created. A two-tier approach was followed for the identification of cell subpopulations: phenotypic and functional. The phenotypic identification was conducted by applying three different clustering methods: PhenoGraph, ClusterX, and K-means, over the phenotypic markers: CD3, CD20, CD4, HLA-DR, Bcl6, CD16, CD68, CD56, CD141, CD1a, CD1c, Blimp1, Langerin, Lysozyme, Podoplanin, FOXP3, S100AB, IRF4, IRF8, CD1a, CXCL13, CD8, CD138, CD123, PD-1, and MelanA. PhenoGraph and ClusterX were implemented using the cytofkit package from R[24]. Clusters were represented by a vector containing the mean of each marker and were used to further associate them to a cell subpopulation using prior knowledge. For a phenotype to be assigned to a cell, at least two clustering methods should agree on their predicted phenotype. Prior to functional identification, PCA was repeated over the Tcy cells (CD8+) using CD69, OX40, LAG3, and TIM3 markers in order to confirm that with the new mask, the same dataset with the same structure as with the CD8+ mask could be retrieved (Extended Data Figure 1). The functional identification was conducted by applying PhenoGraph over the functional markers: CD69, Ki-67, TAP2, GBP1, MYC, p16, MX1, OX40, c-Maf, PD-L1, LAG3, TIM3, and Phospho-Stat1 (with the exception of Tcy cells for which we used a personalized panel consisting of: CD8, CD69, OX40, LAG3, TIM3, PD-1, and Ki-67). Clusters were represented, associated to cell subpopulations, and evaluated for stability as described for the phenotypic identification. Significant differences in the cellular composition of the cores based on activation status were identified using Kruskal-Wallis rank sum test. The same approach was repeated for the brisk infiltrate histopathologic parameter using Mann-Whitney test.

An unbiased quantitative analysis of cell-cell interactions was performed using an adaptation of the algorithm described in Schapiro D, et al[25] for neighbourhood analysis to systematically identify social networks of cells and to better understand the tissue microenvironment. Our adaptation also uses a kernel-based approach (radius = 30 px) to define the neighbourhood of a cell and a permutation test (N = 1000) to compare the number of neighbouring cells of each phenotype in a given image to the randomized case. This allows the assignment of a significance value to a cell-cell interaction representative of the spatial organization of the cells. Significance values were

further classified into: avoidance (-1), non-significant (0), and proximity (1) using a significance threshold of 0.001 (more significant that all the random cases). Interactions across images were integrated according to equation 1:

$$P_{i,j} = \frac{\sum_{k=1}^{M}(c_{i,j,k}\cdot\sqrt{N_{i,j,k}})}{\sum_{k=1}^{M}(\sqrt{N_{i,j,k}})}$$ [1]

where $C_{i,j,k}$ is the significance value (-1, 0, or 1) of the interaction between cell types i and j for image k, and $N_{i,j,k}$ is the geometric average of the number of cells of type i and j for image k. Cell-cell interactions were considered strong if they were significant in at least 75% of the N-adjusted cases (abs(P) > 0.75), moderate if 50%, (0.5 < abs(P) <= 0.75), weak if 25% (0.25 < abs(P) <= 0.5), and non-significant otherwise (abs(P) <= 0.25). A comparative analysis of the above described method was performed for the different core statuses as well as for the different brisk infiltrate cases (Figure 4).

Even though neighbourhood analysis allows evaluation of cell-cell interactions, the mathematical model applied is limited in cases where there are dominant cell types that grow in nests, with cells packed next to each other. Therefore, for melanoma cells, we evaluated the closest neighbours by counting the number of cells of each subpopulation (apart from melanoma cells) that were in their neighbourhood and divided the amount by the geometric average of the number of melanoma cells and the number of cells of the specific population across all the cores in which the specific population appears. This analysis was repeated for the different brisk and activation cases.

Laser Microdissection

18 fresh frozen melanoma metastases with different types of TILs patterns (9 brisk, 7 non brisk, 1 absent and 1 tumoral melanosis) were collected in the Department of Pathology of the University Hospitals Leuven (KU Leuven), Belgium. 10 micrometre-thick sections were cut from each fresh frozen block and put on a film slide (Zeiss, Oberkochen, Germany). Sections were stained with crystal violet. Areas with dense TILs infiltrate were microdissected with the Leica DM6000 B laser microdissection device (Leica, Wetzlar, Germany). A calculation was made in order to microdissect the same surface in all the samples in order to minimize differences between the samples (around 10.000 lymphocytes/sample).

qPCR of laser microdissected samples

RNA extraction was done with RNeasy® Plus Micro Kit (Qiagen) according to the protocol. cDNA retrotranscription followed by an amplification step was done with Ovation® Pico SL WTA System V2 (Nugen) according to protocol. Primers for Interferon gamma (INFg, forward 'TGTTACTGCCAGGACCCA' and reverse 'TTCTGTCACTCTCCTCTTTCCA'), TIM3 (forward 'CTACTACTTACAAGGTCCTCAGAA' and reverse 'TCCTGAGCACCACGTTG'), LAG3 (forward 'CACCTCCTGCTGTTTCTCA' and reverse 'TTGGTCGCCACTGTCTTC'), CD40-L (forward 'GAAGGTTGGACAAGATAGAAGATG' and reverse 'GGATAAGGATCTTTCTCCTGTGTT'), CD45 (forward 'GCTACTGGAAACCTGAAGTGA' and reverse 'CACAGATTTCCTGGTCTCCAT'), Beta2microglobulin (forward 'ACAGCCCAAGATAGTTAAGTG' and reverse 'ATCTTCAAACCTCCATGATGC'), HPRT (forward 'ATAAGCCAGACTTTGTTGGA' and reverse 'CTCAACTTGAACTCTCATCTTAGG') were designed with Perl Primer® and tested in our laboratory. 96-wells plates were loaded with Fast SYBR® Green Master Mix, the primers and the samples in the recommended proportions, and analysed with the 7900 HT Fast Real-Time PCR system (Applied Biosystems). The log fold change (logFC) of the expression values towards the

expression value of CD45 were calculated. If the log(IFNg/CD45) was positive, the sample was classified as positive. On the other hand, exhaustion was defined by expression of LAG3 and/or TIM3 with lack of IFNg and CD40L expression.

Shotgun Proteomics of laser microdissected samples

The materials used for the shotgun proteomics analysis were: Trifluoroacetic acid, MS grade porcine trypsin, DTT (dithiothreitol), IAA (Iodoacetamide), ABC (Ammonium Bicarbonate), HPLC grade water, acetonitrile (ACN), were from Sigma-Aldrich (Sigma-Aldrich Chemie GmbH, Buchs, Switzerland). All solutions for Mass Spectrometry (MS) analysis were prepared using HPLC-grade. LCM collected material corresponding to about 10^4 cells for each sample group was re-suspended in 90 µl of bidistilled water and immediately stored at -80°C. For the bottom-up MS analysis, all the samples were processed and trypsinized. Briefly, thawed cells were submitted to a second lysis adding 60 µl of 0.25% w/v RapiGest surfactant (RG, Waters Corporation) in 125mM ammonium bicarbonate (ABC) and sonicated for 10 min. Samples were then centrifuged at $14\,000 \times g$ for 10 min. About 140 µl of supernatants were collected, transferred in a new tube and quantified using bicinchoninic acid assay (Pierce -Thermo Fisher Scientific). After 5min denaturation (95°C), proteins were reduced with 50mM DTT in 50mM ABC at room temperature and alkylated with 100mM IAA in 50mM ABC (30 min incubation in dark). Digestion of samples was performed overnight at 37°C using 2µg of MS grade trypsin. RG surfactant were removed using an acid precipitation with TFA at a final concentration of 0.5% v/v. Samples were then spun down for 10 min at 14000 x g and supernatants collected for MS analysis. Peptide mixtures were desalted and concentrated using Ziptip[TM] µ-C8 pipette tips (Millipore Corp, Bedford, MA). An equal volume of eluted digests was injected at least three times for each sample into Ultimate[TM] 3000 RSLCnano (ThermoScientific, Sunnyvale, CA) coupled online with Impact HD[TM] UHR-QqToF (Bruker Daltonics, Germany). In details, samples were concentrated onto a pre-column (Dionex, Acclaim PepMap 100 C18 cartridge, 300 µm) and then separated at 40°C with a flow rate of 300 nL/min through a 50 cm nano-column (Dionex, ID 0.075mm, Acclaim PepMap100, C18). A multi-step gradient of 4 hours ranging from 4 to 66% of 0.1% formic acid in 80% ACN in 200min was applied[26]. NanoBoosterCaptiveSpray™ ESI source (Bruker Daltonics) was directly connected to column out. Mass spectrometer was operated in data-dependent acquisition mode, using CID fragmentation assisted by N2 as collision gas setting acquisition parameters as already reported[27]. Mass accuracy was assessed using a specific lock mass (1221.9906 m/z) and a calibration segment (10 mM sodium formate cluster solution) for each single run. Raw data from nLC ESI-MS/MS were elaborated through DataAnalysis[TM] v.4.1 Sp4 (Bruker Daltonics, Germany) and converted into peaklists. Resulting files were interrogated for protein identification through in-house Mascot search engine (version: 2.4.1), as described[27]. Identity was accepted for proteins recognized by at least one unique and significant (p-value < 0.05) peptide.

Pathways Analysis

Gene-set enrichment analysis was performed with DAVID 6.8[28,29]. Pathways were visualized and partially analysed with STRING v10[30].

Results

Functional analysis of TILs

We classified each CD8+ cell in the TMA cores as part of a spectrum ("cell status") ranging from "active" (CD69high and/or OX40high) to "exhausted" (TIM3highCD69lowOX40low) (Figure 2). Two intermediate states between activation and exhaustion were identified, one passing through a gradual decrease of CD69 with parallel increase of TIM3 expression, and the other in which OX40 was co-expressed with LAG3. These latter two markers had also the highest Pearson correlation coefficient among all the activation and exhaustion markers (R = 0.35). When all the cells were included in the analysis (using the DAPI mask instead of the CD8 mask), the same computational approach repeated on the subgroup identified as Tcy (CD8+) gave a similar result, confirming the solidity of this approach (Extended Data Figure 1).

Comparing the activation level of the cells in each core to the activation level of the cells across all the cores, we assigned a label to each core in the TMA ("core status"): active (17/60 cores), in transition (23/60 cores), or exhausted (20/60 cores). At this point, with the exception of 8 patients in which the size of the melanoma allowed to sample only a single core, we could assess the heterogeneity of the immune response in different areas of the melanoma from the same patient. 10/21 showed homogeneous core statuses (4 active, 5 in transition, and 1 exhausted) and 11/21 showed heterogeneity. In summary, we observed a spatially homogeneous TILs infiltrate in 6 active cases (20.8%) and in 5 exhausted ones (17.2%), while in 18 cases (62%) the presence of areas of transition suggested an ongoing dynamic transition toward exhaustion (Figure 2.D).

We then checked whether the cores status was significantly associated with spontaneous regression of the tumour, regarded as the result of a successful Tcy immune response, and with other histopathological parameters (histological subtype, ulceration, Breslow thickness, mitoses). Late regression areas indeed showed significant differences in the mean level of activation of the cores as compared to early regression (p = 0.022) and no regression (p = 0.031). No significant differences were instead found between early regression and no regression. Higher levels of activation were found in Lentigo Maligna Melanoma (p = 0.02). However, since only three LMM cores from two patients were included in our data set, no definite conclusions can be drawn from this data. The other histopathological prognostic parameters did not show significance (Extended Data Table 1).

Clustering and neighbourhood analysis

The inflammatory subpopulations were identified using three different unsupervised clustering methods (KMeans, PhenoGraph, and ClusterX) followed by manual annotation by the pathologist of the clusters and assessment of their solidity as described in the methods (Extended Data Figure 2). On 19 clusters identified, 17 could be associated to a determinate cell lineage, while the remaining two were discarded. Ultimately, we could solidly classify 179304 out of 242224 cells across all of our cores (74.02%). A further subclustering was done on the so-identified clusters using a set of functional markers with PhenoGraph, obtaining a total number of 47 subclusters (Figure 3A). The most abundant cell population consisted of melanoma cells (41.76%). Most of the melanoma cells expressed both melanocytic markers Melan A and S-100, while two minor groups of melanoma cells had loss of expression of one of the two markers. When Melan A was lost, melanocytes were distinguished by dendritic cells thanks to the absence of expression of the other dendritic cell markers included in the panel. Expression of Melan A in melanoma cells was associated with a lower amount of proliferating melanoma cells and occasional expression of p16. The second most abundant cell type were the macrophages ("Macroph", CD68+CD163+Lysozyme+HLA-DR+, 11.48%), one quarter of them expressing the immunosuppressive marker TIM3. Epithelial cells represented 8.4% of the population of our data

set. The lymphoid compartment accounted for multiple subpopulations. Tcy (CD3+CD8+) and T helpers ("Th", CD3+CD4+FOXP3-) were the most abundant subtypes, accounting respectively for 11.19% and 10.10% of all the cells, while regulatory T cells ("Treg", CD3+CD4+FOXP3+) represented 2,82% of the cells. We interpreted the last T cell cluster as T follicular helpers ("Tfh", CXCL13+PD1+, 2.57%) even though this cluster did not express the full Tfh phenotype. Within Tcy, we could identify the active (CD69+OX40+/-, 16,1% of all the lymphocytes), transition (balanced expression of both exhaustion and expression markers, 29,6%) and exhausted (high expression of LAG3 and/or TIM3, low/absent CD69 and OX40, 12,8%) functional subgroups in the Tcy cluster, plus a clonally expanding subgroup (6%), and one with low or absent expression of all functional markers, that we defined as "anergic" (34,9%) but that could represent also naïve T cells. Th could also be subdivided according to their expression of activation and exhaustion markers (28,5% ThCD69+, considered active, 18,46% ThCD69+TIM3+, considered transitional, and 13,08% ThTIM3+, considered immunosuppressive). NK cells, as expected in melanoma, were extremely infrequent (0.51%), but even more infrequent were the B cells ("BC", CD20+), that represented 0.12% of the total number of cells in the cores. The CD20 negative cells characterized by high expression of IRF4 and Blimp1 ("IRF4_Blimp1", 1,19%) were interpreted as plasma cells[23]. Finally, we could identify among the dendritic cell group the classical dendritic cells type 1 ("cDC1", CD141+CD4+IRF8+, 4.13%), classical dendritic cells type 2 ("cDC2", CD1c+CD4+HLA-DR+, 2.41%), Langerhans cells ("Lang", CD1a+Langerin+, 2.13%), and plasmacytoid dendritic cells ("pDC", CD123+, 0.33%). In both the two classical dendritic cell subgroups an immunosuppressive TIM3+ subpopulation was identifiable, while in the pDC group a small subpopulation was found to express PD-L1. No immunosuppressive subpopulation was identified among the Langerhans cells. Some subpopulations were statistically significantly different (p-val < 0.05) comparing brisk vs non-brisk and active vs transition vs exhausted. Brisk cases were significantly enriched in BC (p-val = 0.041), TIM3+cDC1 (p-val = 0.024), macrophages (p-val = 0.043) (including the proliferating subgroup (p-val = 0.034)), NK (p-val = 0.011), anergic Tcy (p-val = 0.039) and proliferating Tcy (p-val = 0.006) (Figure 3B). Active cores had, together with more active Tcy (p-val < 0.001), higher percentages of Tcy in transition (p-val = 0.030), transition cores more Th (p-val = 0.057), while exhausted cases had more TIM3+Tregs (p-val = 0.018) and more proliferating melanoma cells (p-val = 0.045) (Figure 3C).

After having identified the clusters, we applied neighbourhood analysis in order to systematically identify social networks of cells to draw conclusions on actual cell-cell interactions. To understand what is the immune context that determines activation and exhaustion, we compared the results of neighbourhood analysis between the three functional subgroups (Figure 4A). The detection of the interaction of Langerhans cells with the epithelium, identifiable in all the three groups, was considered as positive control and proof of concept for the method. Some other interactions were present in all three activation statuses: TIM3+ macrophages and exhausted/translational Tcy, CD69+TIM3+Th and translational Tcy, active Th and anergic Tcy, TIM3+cDC1 and CD69+TIM3+Th. Only in active cases it was possible to observe neighbourhood between active Th and active Tcy, TIM3+macrophages and CD69+TIM3+Th, NK and Tfh, Tfh and exhausted Tcy, pDC and Langerhans. Peculiar to transition cases were the spatial interactions between CD69+TIM3+Th and exhausted Tcy, CD69+TIM3+Th and proliferating Treg, Treg and Transition Tcy, active Th and transition Tcy, Tfh and PD-L1+pDC. Exclusively in exhausted case we had in close proximity CD69+TIM3+Th and anergic Tcy, anergic Tcy and active Th, anergic Tcy and BC expressing INFgamma molecules, CD69+TIM3+Th and TIM3+cDC2, TIM3+macrophages and transition Tcy, NK and PD-L1+pDC, and the different subtypes of Tfh and cDC1.

To investigate the significance of the neighbourhood composition between the inflammatory subtypes we used a permutation test. Interpreting the results of permeation analysis needs to be done with caution when looking at packed cells: melanoma cells showed avoidance from the other cells of the inflammatory infiltrate, an incoherent result due to the fact that they generally grow in nests. In presence of extensive areas of cells packed close to each other, there are very low chances that other interactions may result stronger than the ones in between them. Therefore, we applied another type of neighbourhood analysis for melanoma cells, as described in the methods, and we observed that macrophages and epithelial cells were in general most often located in strict proximity to the melanoma cells, without differences among the functional and morphologic categories, and brisk cases had more Tcy in close proximity with melanoma cells than non-brisk cases, as expected (Extended Data Figure 3). Interestingly, brisk cases had a higher prevalence specifically of transition and active Tcy in contact with melanoma cells compared with non-brisk cases (Active/Exhausted ratio: Brisk = 2.108762, Non-Brisk = 1.331195), that instead had relatively more exhausted and anergic cells in contact with melanoma cells (Active/Anergic ratio: Brisk = 1.743081, NonBrisk = 0.7704947).

qPCR and shotgun proteomics

To confirm the existence of the same functional subcategories in metastatic melanoma samples, that represents the type of lesions that receive the immunotherapeutic treatment we perform qPCR followed by proteomics on microdissected TILs, comparing brisk and non-brisk cases using an "absent" case and a "tumoral melanosis" (i.e. complete melanoma regression with persistence of melanin-loaded macrophages) case as controls. Furthermore, we could measure directly the levels of expression of IFNg, the best indicator of CD8+ activation, for which no suitable antibody exists for FFPE material. 4/8 melanomas with a brisk TILs pattern, and 3/7 with non-brisk TILs pattern proved active, confirming that we could subclassify from a functional point of view also metastatic lesions (Extended Data Figure 4).

To correlate the gene expression measured by qPCR with the actual protein expression we performed a proteomic analysis on the microdissected material in 3 representative cases, i.e. one with high IFNg expression ("IFNg-high"), one with the high LAG3 and no IFNg expression ("LAG3-high"), and one with none of the four markers strongly expressed ("none"). The results showed a higher number of proteins identified in the "IFNg-high" sample (324) compared to "LAG3-high" (93) and "none" (134), with almost two thirds of them (210/324) not shared with the other two samples and enriched in proteins involved in different inflammatory pathways (innate immunity, TNFR1 signalling pathway, FAS signalling pathway, T cell receptor and Fc-epsilon receptor signalling pathway), including the interferon gamma-mediated signalling pathway. As we minimized the difference in the number of microdissected cells in each sample, these observations could only be explained by a higher production of pro-inflammatory proteins in the "IFNg-high" sample.

Discussion

The fact that not all brisk infiltrates have a good prognosis[16,32], that TILs populations can be very heterogeneous[33,34] and that anti-PD-1 immunotherapy was found to support functionally activated T cells[17] urged a thoughtful investigation of the functional status of the TILs. Since the molecules that mediate exhaustion are expressed upon activation in order to prevent the hyperactivity of the immune system[35], to investigate the activation status of lymphocytes the simultaneous expression of several molecules must be evaluated at a single cell level and at once, as a panel[36]. Moreover, the assessment of the interactions between cells requires preservation of the tissue architecture. Our

study is the first to assess more than 40 markers on tissue sections at single cell resolution without losing their spatial distribution. We obtained by high-dimensional in-situ immunotyping a snapshot of the co-expression patterns of activation and inhibition markers in tissue sections. OX40 was strongly correlated with Ki-67 and PD-1, confirming its role in sustaining Tcy clonal expansion[37]. LAG3 is expressed very early and only transiently during T cell activation, but persistent T cell activation with sustained expression of LAG3 together with other exhaustion markers (e.g. PD-L1) results in T cell dysfunction[38]. LAG3+Tregs have been shown to suppress DC maturation[39].

We could observe that both brisk and non-brisk cases can harbour predominantly exhausted TILs or predominantly active TILs; therefore, with the morphological classification (brisk – non brisk – absent), a complementary functional classification (active – transitional - exhausted) co-exists. The importance of going beyond the morphologic classification of TILs was previously raised by others, who claimed that a functional analysis could help in the application of a better immunoscore for therapeutic prediction[34]. Only a minority (15%) of patients presented with all active cores, and 30% of patients presented with at least one active core, whereas the great majority of the patients presented with only exhausted or transition areas at the moment in which the melanoma was removed. Since the percentage of patients that obtains a durable response with single agent checkpoint therapy[1,2,3] lies between 15 and 30%, and since Krieg et al[17] reported that anti-PD-1 immunotherapy supports functionally activated T cells, it is tempting to speculate that the "mostly active" TILs cases could correspond to the responders to immunotherapy, while the "mostly exhausted" cases could benefit instead from combination approaches with immunotherapy and other type of therapies in order to rescue the exhausted T cells. Therefore, adding the functional evaluation could definitely improve the predictive value of the morphological TILs patterns in melanoma. Since spontaneous melanoma regression, present in only 10-35% of the melanomas, is considered to be the end-results of the melanoma-eliminating capacities of active TILs, we also studied the association of activation of TILs with early and late regression[40]. Our data showed a clear-cut association of activation of TILs with late regression areas, indirectly proving the functional meaning of an active infiltrate.

We obtained a functional picture of the inflammatory landscape in brisk versus non-brisk cases and in active/transition/exhausted TILs microenvironments. A previous study already casted some doubts on the significance of the brisk pattern, as they found no evidence of clonally expanded TILs in some cases with a brisk infiltrate[34], and hypothesized that clonally expanded T cells might represent not only cytotoxic cells but also regulatory cells. We indeed found not only immune stimulating cells such as proliferating and anergic Tcy, cDC1, and NK significantly increased in brisk cases, but also immune suppressive cells such as BC and macrophages. Active and exhausted cases were instead functionally coherent categories. An active microenvironment was enriched for active Tcy, while an exhausted microenvironment was enriched not only with Treg, possible origin of the exhaustion, but also with proliferating melanoma cells, possible effect of the lack of immune control over the neoplasia.

A slight increase in Treg between active and exhausted cases may not be able alone to justify the shift from an active to an exhausted microenvironment. To this end, adding the spatial information and making inferences about the interactions between the cells in the tissue using neighbourhood analysis could represent a better way to investigate this dynamic. In this way, we could identify the most important differences that could explain the transition from an active to an exhausted environment. In active cases, it was possible to identify a stronger spatial association between active Th and active Tcy, while the same association was weaker in transition cases and disappeared in exhausted ones. In general, there was a decrease of interactions between the Th compartment and the active Tcy from active to exhausted cases. This could be explained observing the peculiar

interactions with the other cell types in each of the functional states. In active cases, TIM3+ macrophages interacted with CD69+TIM3+ Th, as well as cDC1 expressing TIM3 interacted with TIM3+ Treg, possibly limiting their suppressive effect on the Tcy. In transition cases instead TIM3+ cDC1 are interacting only with CD69+TIM3+ Th and transitional Tcy, their block on Treg is removed and the Treg population is globally inhibiting all the Th subpopulation, probably reducing the strength of activation of the Tcy compartment. In fact, in active and exhausted cases the active Th are seen more often in contact with transitional Tcy, that are also the direct target of the Treg, while CD69+TIM3+ Th are in direct connection with exhausted Tcy. Starting from the transitional status we see the appearance of interactions between BC expressing INFg-related molecules and CD69+TIM3+ Th and anergic Tcy, and between TIM3+ cDC2 and CD69+TIM3+ Th, confirming the immune suppressive role of this cell types. The presence of a shift of interaction of the active Th towards anergic Tcy observed in the exhausted cases may be a rescue mechanism, meant to induce new active Tcy starting from anergic/naïve Tcy but possibly contrasted by the effects of the surrounding immunosuppressive cells. Finally, in active cases there is a strong interaction between TIM3+ macrophages and exhausted/transitional Tcy along all the statuses, confirming their prominent role in keeping the exhaustion in the tumoral microenvironment. Considering the brisk and non-brisk classification, instead, we can see that both categories are a mixture of immune stimulating and suppressive interactions, once again confirming that the morphological categories are functionally very heterogeneous (Figure 4B). Paradoxically, it looks like non- brisk cases have more activating interactions than brisk cases. Nevertheless, the real determinant that may justify the good prognosis of the brisk cases may be that in brisk case melanoma cells are more in contact specifically with active/transition Tcy than non-brisk cases, that instead had more exhausted and anergic cells in contact with melanoma cells.

In conclusion, in this paper we have shown that the activation status of the TILs does not necessarily parallel the morphological categories, and that within a single melanoma, the inflammatory response may vary considerably. We have shown a bioinformatic pipeline that, starting from common immunofluorescence stainings, can transform the tissue into a digitalized image, which represents the starting point for multiple deeper levels of analysis (Figure 5). We hypothesize that the general good prognosis of melanomas with a brisk pattern of TILs could be based on the fact that the Tcy that are in contact with the melanoma cells at the moment that the melanoma is excised are still active, and consequently the melanoma is still under immune control. In this study we have also described the main interactions between the inflammatory subpopulations in an active, transition and exhausted environment, interactions that should be taken in consideration when assessing the response to immunotherapy and that will ultimately lead to the identification of functional inflammatory microenvironments that may benefit from personalized combined therapy protocols.

Acknowledgements

Francesca Maria Bosisio is funded by the MEL-PLEX research training programme ('Exploiting MELanoma disease comPLEXity to address European research training needs in translational cancer systems biology and cancer systems medicine', Grant agreement no: 642295, MSCA-ITN-2014-ETN, Project Horizon 2020, in the framework of the MARIE SKŁODOWSKA-CURIE ACTIONS).

Asier Antoranz is funded by the SyMBioSys research training programme ('Systematic Modeling of Biological Systems'), grant agreement no: 675585, MSCA-ITN-2015-ETN, Project Horizon 2020, in the framework of the MARIE SKŁODOWSKA-CURIE ACTIONS.

Maddalena Maria Bolognesi is supported by a clinical research project BEL114054 (HGS1006-C1121) of the University of Milano-Bicocca and GlaxoSmithKline, UK.

Disclosures

No conflict of interest to declare.

References

1. Ribas A, Flaherty KT. Gauging the Long-Term Benefits of Ipilimumab in Melanoma. J Clin Oncol. 2015;33[17]:1865-6.
2. Ribas A, Hamid O, Daud A, Hodi FS, Wolchok JD, Kefford R, et al. Association of Pembrolizumab With Tumor Response and Survival Among Patients With Advanced Melanoma. JAMA. 2016;315[15]:1600-9.
3. Robert C, Long GV, Brady B, Dutriaux C, Maio M, Mortier L, et al. Nivolumab in previously untreated melanoma without BRAF mutation. N Engl J Med. 2015;372[4]:320-30.
4. Seremet T, Koch A, Jansen Y, Schreuer M, Wilgenhof S, Del Marmol V, et al. Molecular and epigenetic features of melanomas and tumor immune microenvironment linked to durable remission to ipilimumab-based immunotherapy in metastatic patients. J Transl Med. 2016;14(1):232.
5. Hugo W, Zaretsky JM, Sun L, Song C, Moreno BH, Hu-Lieskovan S, et al. Genomic and Transcriptomic Features of Response to Anti-PD-1 Therapy in Metastatic Melanoma. Cell. 2016;165(1):35-44.
6. Gibney GT, Weiner LM, Atkins MB. Predictive biomarkers for checkpoint inhibitor-based immunotherapy Lancet Oncol. 2016;17(12):e542-e551.
7. Teixido C, Gonzalez-Cao M, Karachaliou N, Rosell R: Predictive factors for immunotherapy in melanoma. Ann Transl Med 2015;3(15): 208-218
8. Vasaturo A, Halilovic A, Bol KF, Verweij DI, Blokx WA, Punt CJ, et al. T cell landscape in a primary melanoma predicts the survival of patients with metastatic disease after their treatment with dendritic cell vaccines. Cancer Res. 2016;76(12):3496-506..
9. Loo K, Daud A. Emerging biomarkers as predictors to anti-PD1/PD-L1 therapies in advanced melanoma. Immunotherapy. 2016;8[7]:775-84.
10. Koguchi Y, Hoen HM, Bambina SA, Rynning MD, Fuerstenberg RK, Curti BD, et al. Serum Immunoregulatory Proteins as Predictors of Overall Survival of Metastatic Melanoma Patients Treated with Ipilimumab. Cancer Res. 2015;75[23]:5084-92.
11. Teng MW, Ngiow SF, Ribas A, Smyth MJ. Classifying Cancers Based on T-cell Infiltration and PD-L1. Cancer Res. 2015;75[11]:2139-45.
12. Clark WH Jr, Elder DE, Guerry D 4th, Braitman LE, Trock BJ, Schultz D, et al. Model predicting survival in stage I melanoma based on tumor progression. J Natl Cancer Inst. 1989;81(24):1893-904.
13. Clemente CG, Mihm MC Jr, Bufalino R, Zurrida S, Collini P, Cascinelli N. Prognostic value of tumor infiltrating lymphocytes in the vertical growth phase of primary cutaneous melanoma. Cancer. 1996;77[7]:1303-10.
14. Mihm MC Jr, Clemente CG, Cascinelli N. Tumor infiltrating lymphocytes in lymph node melanoma metastases: a histopathologic prognostic indicator and an expression of local immune response. Lab Invest. 1996;74[1]:43-7.
15. Bosisio FM, van den Oord JJ. Immunoplasticity in cutaneous melanoma: beyond pure morphology. Virchows Arch. 2017;470(4):357-369.
16. Saltz J, Gupta R, Hou L, Kurc T, Singh P, Nguyen V, et al. Spatial Organization and Molecular Correlation of Tumor-Infiltrating Lymphocytes Using Deep Learning on Pathology Images. Cell Rep. 2018;23(1):181–7.
17. Krieg C, Nowicka M, Guglietta S, Schindler S, Hartmann FJ, Weber LM, et al. High-dimensional single-cell analysis predicts response to anti-PD-1 immunotherapy. Nat Med. 2018;24(2):144-153.

18. Biology of single cells shines a light on collaboration. Nature. 2017;547(7661):5.
19. Binnewies M, Roberts EW, Kersten K, Chan V, Fearon DF, Merad M, et al. Understanding the tumor immune microenvironment (TIME) for effective therapy. Nat Med. 2018;24(5):541-550.
20. Ascierto PA, Agarwala SS, Ciliberto G, Demaria S, Dummer R, Duong CPM, et al. Future perspectives in melanoma research "Melanoma Bridge", Napoli, November 30th-3rd December 2016. J Transl Med. 2017;15(1):236.

21. Buggert M, Nguyen S, Salgado-Montes de Oca G, Bengsch B, Darko S, Ransier A, Roberts ER, Del Alcazar D, et al. Identification and characterization of HIV-specific resident memory CD8+ T cells in human lymphoid tissue. Sci Immunol. 2018;3(24).
22. Cattoretti, G., Bosisio, F.M., Marcelis, L. & Bolognesi, M.M. Multiple Iterative Labeling by Antibody Neodeposition (MILAN). Protocol Exchange doi:10.1038/protex.2018.106
23. Caicedo JC, Cooper S, Heigwer F, Warchal S, Qiu P, Molnar C, et al. Data-analysis strategies for image-based cell profiling. Nat Methods. 2017;14(9):849-863.
24. Chen H, Lau MC, Wong MT, Newell EW, Poidinger M, Chen J. Cytofkit: A Bioconductor Package for an Integrated Mass Cytometry Data Analysis Pipeline. Schneidman D, ed. PLoS Computational Biology. 2016;12(9):e1005112. doi:10.1371/journal.pcbi.1005112.
25. Schapiro D, Jackson HW, Raghuraman S, Fischer JR, Zanotelli VRT, Schulz D, Giesen C, Catena R, Varga Z, Bodenmiller B. miCAT: A toolbox for analysis of cell phenotypes and interactions in multiplex image cytometry data. Nature methods. 2017;14(9):873-876. doi:10.1038/nmeth.4391.
26. Chinello C, Stella M, Piga I, Smith AJ, Bovo G, Varallo M, Ivanova M, Denti V, Grasso M, Grasso A, Del Puppo M, Zaravinos A, Magni F. Proteomics of liquid biopsies: Depicting RCC infiltration into the renal vein by MS analysis of urine and plasma. J Proteomics. 2019 Jan 16;191:29-37.
27. Liu X, Chinello C, Musante L, Cazzaniga M, Tataruch D, Calzaferri G, et al. Intraluminal proteome and peptidome of human urinary extracellular vesicles. Proteomics Clin Appl. 2015;9(5-6):568-73.
28. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. Nat. Protoc.2008; 4,44.

29. Dennis G Jr, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC, et al. DAVID: Database for Annotation, Visualization, and Integrated Discovery. Genome Biol. 2003;4(5):P3.
30. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al. STRING v10: protein-protein interaction networks, integrated over the tree of life. Nucleic Acids Res. 2015;43(Database issue):D447-52.
31. Alinikula J, Nera KP, Junttila S, Lassila O. Alternate pathways for Bcl6-mediated regulation of B cell to plasma cell differentiation. Eur J Immunol. 2011;41(8):2404-13.
32. Thorsson V, Gibbs DL, Brown SD, Wolf D, Bortone DS, Ou Yang TH, et al. The Immune Landscape of Cancer. Immunity. 2018;48(4):812-830.e14.
33. Bernsen MR, Diepstra JH, van Mil P, Punt CJ, Figdor CG, van Muijen GN, et al. Presence and localization of T-cell subsets in relation to melanocyte differentiation antigen expression and tumour regression as assessed by immunohistochemistry and molecular analysis of microdissected T cells. J Pathol. 2004;202(1):70-9.
34. Pao W, Ooi CH, Birzele F, Ruefli-Brasse A, Cannarile MA, Reis B, et al. Tissue-Specific Immunoregulation: A Call for Better Understanding of the "Immunostat" in the Context of Cancer.Cancer Discov. 2018;8(4):395-402.

35. Wherry EJ. T cell exhaustion. Nat Immunol. 2011;12(6):492-9

36. Weixler B, Cremonesi E, Sorge R, Muraro MG, Delko T, Nebiker CA, et al. OX40 expression enhances the prognostic significance of CD8 positive lymphocyte infiltration in colorectal cancer. Oncotarget. 2015;6(35):37588-99

37. Ruea-Yea Huang, Cheryl Eppolito, Shashikant Lele, Protul Shrikant, Junko Matsuzaki, Kunle Odunsi. LAG3 and PD1 co-inhibitory molecules collaborate to limit CD8+ T cell signaling and dampen antitumor immunity in a murine ovarian cancer model. Oncotarget. 2015; 6(29): 27359–27377.

38. Andrews LP, Marciscano AE, Drake CG, Vignali DA. LAG3 (CD223) as a cancer immunotherapy target. Immunol Rev. 2017;276(1):80-96.

39. Spranger S, Luke JJ, Bao R, Zha Y, Hernandez KM, Li Y, et al. Density of immunogenic antigens does not explain the presence or absence of the T-cell–inflamed tumor microenvironment in melanoma. Proc Natl Acad Sci U S A. 201;113(48):E7759-E7768.

40. Botella-Estrada R, Traves V, Requena C, Guillen-Barona C, Nagore E. Correlation of histologic regression in primary melanoma with sentinel node status.JAMA Dermatol. 2014;150(8):828-35

Figure 1. Single cell analysis scheme. A) 60 cores were obtained from 29 patients, assembled in a Tissue Micro Array, and analysed using the MILAN technique[22]; B) CD8+ cells were analysed using image analysis for functionality using an activation parameter derived from multiple activation and exhaustion markers evaluated at single cell resolution; C) All the cell populations in the cores were phenotypically identified using consensus between three clustering methods and manual annotation from the pathologists; D) The social network of the cells was analysed using a permutation test for neighbourhood analysis in order to make inferences on cell-cell interactions. All steps are exemplified using three representative images (35, 40, and 46).

Figure 2. Definition of activation. A biplot showing the projection of the cells as well as the rotation vectors of the markers over PC2 and PC3 (A) was the first step to define a gradient of activation going from the maximum projected value of CD69 (maximum activation) to the maximum projected value of TIM3 (maximum exhaustion) (B). C) Z-scores of the original markers over PC2 and PC3. D) Visual representation of the inter- and intra-patient heterogeneity. Each core is assigned an activation status ('Active', 'Transition', or 'Exhausted'). The cores are grouped for each patient, giving an at-a-glance representation of the heterogeneity of the activation status in different areas of the melanoma in the same patient.

Figure 3. Distribution of the immune cells' subgroups in the morphological and functional TILs categories. The percentage of cells for each inflammatory subpopulation across all the cores is showed in (A). The 17 phenotypic clusters are on the upper side of the graph, while at the bottom each of them is subdivided in the respective functional subclusters. BC = B cells; cDC1 = classical dendritic cells type 1; cDC2 = classical dendritic cells type 2; Epith = epithelial cells; IRF4BLIMP1 = plasma cells; Lang = Langerhans cells; LV = lymph vessels; Macroph = macrophages; pDC = plasmocytoid dendritic cells; S-M+ = S100+MelanA- melanoma cells; S+M- = S100-MelanA+ = melanoma cells; S+M+ = S100+MelanA+ melanoma cells; Tcy = cytotoxic T cells; Tfh = T follicular helpers; Th = T helpers; Treg = regulatory T cells; suffix: "prolif" = proliferating, IFNg = interferon gamma. B) The percentage of the different immune cells subpopulations compared in brisk and non-brisk cases and C) in active, transition and exhausted microenvironments. The populations with p-value < 0.05 are indicated by the *, whereas p-values < 0.1 by (*).

Figure 4. Neighbourhood analysis in the morphological and functional TILs categories. The result of the neighbourhood analysis for active, transition and exhausted environments (A) and for brisk and non-brisk cases (B) is represented as heatmaps on the left and as conceptual maps on the right. In the heatmaps, different intensities of green are used to express different strengths of interaction between the cell subtypes, so as different intensities of red are used to express different strengths of avoidance. In the conceptual maps, the same concept is expressed by the thickness of the line connecting two populations. The color of the line indicates interactions leading to immune suppression = red, to immune stimulation = green, and to a probably sub-optimal/impaired immune stimulation = orange. In the heatmaps are represented all the findings, while in the heatmaps only the relevant interactions were selected, after filtering the interactions internal to the same immune cell population (these may be due to clonal expansion/phenotype shifting instead that a real cross-talking).

Figure 5. Complete digitalization of a core and practical example of a possible downstream analysis. Anti-clockwise: (1) images for the markers stained with the MILAN multiplexing

technique (for this work 39 markers, but we reached in our laboratory an output of around 100 markers per single section) are acquired and composite images with some selected markers can be prepared. (2) All the markers are used to phenotypically identify all the cell subtypes present in the tissue and the tissue is digitally reconstructed. (3) Studies of the functional status of these cells can be done, for example we could localize exhausted and active cells in the tissue. (4) With neighbourhood analysis it is possible to identify the cells that are in proximity with each other more than chance can suggest, inferring possible interactions between these cell types (in the image, two cells identified as "neighbours" are encircled in red).

Active

Transition

Exhausted

Interaction

- Strong Interaction
- Moderate Interaction
- Weak Interaction
- No Interaction
- Weak Avoidance
- Moderate Avoidance
- Strong Avoidance

B