An improved high-quality genome assembly and annotation of Qingke, Tibetan hulless barley

Xingquan Zeng[1,2,3,] †, Xiangfeng Li[5,]†, Lijun Bai[5,]†, Yulin Wang[1,2,3,]†, Tong Xu[5], Hongjun Yuan[1,2,3] , Qijun Xu[1,2,3], Sang Zha[1,2,3], Zexiu Wei[1,3,4], Wangmu Qimei[1,2,3], Yuzhen Basang[1,2,3], Jiabu Dunzhu[1,2,3],Mingzai Yu[1,2,3] & Tashi Nyima[1,3,]*.

Affiliations

1 State Key Laboratory of Barley and Yak Germplasm Resources and Genetic Improvement, Lhasa 850002, China.

2 Research Institute of Agriculture, Tibet Academy of Agricultural and Animal Husbandry Sciences Lhasa, Tibet 850002, China.

3 Tibet Academy of Agricultural and Animal Husbandry Sciences, Lhasa, Tibet 850002, China.

4 Agricultural Resources and Environment Research Institute, Tibet Academy of Agricultural and Animal Husbandry Sciences, Lhasa, Tibet 850002, China.

5. Chengdu Life Baseline Technology Co., LTD, Chengdu 610041, China

† These authors contributed equally to this work.

* Correspondence and requests for materials should be addressed to T.N. (email: nima_zhaxi@sina.com)

Abstract

Background

The Tibetan hulless barley (Hordeum vulgare L. var. nudum), also called "Qingke" in Chinese and "Ne" in Tibetan, is the staple food for Tibetans and an important livestock feed in the Tibetan Plateau. The Qingke in China has about 3500 years of cultivation history, mainly produced in Tibet, Qinghai, Sichuan, Yunnan and other areas. In addition, Qingke has rich nutritional value and outstanding health effects, including the beta glucan, dietary fiber, amylopectin, the contents of trace elements , which are higher than any other cereal crops.

Findings

Here, we reported an improved high-quality assembly of Qingke genome with 4.0 Gb in size. We employed the falcon assembly package ,scaffolding and error correction tools to finish improvement using PacBio long reads sequencing technology, with contig and scaffold N50 lengths of 1.563Mb and 4.006Mb, respectively, representing more continuous than the original Qingke genome nearly two orders of magnitude. We also re-annotated the new assembly, and reported 61,303 stringent confident putative protein-coding genes, of which 40,457 is HC genes. We have developed a new Qingke genome database (QGD) to download and use friendly, as well as to better manage the information of the Qingke genetic resources.

Conclusions

The availability of new Qingke genome and annotations will take the genetics of Qingke to a new level and will greatly simplify the breeders effort. It will also enrich the granary of the Tibetan people.

Keyword: Qingke; Hulless barley; Genome assembly; full-length cDNA; Qingke Genome Database

Introduction

Qingke (Hordeum vulgare L. var. nudum) is a cereal crop, a variant of barley, alias called hulless barley, yuan wheat, rice barley(**Fig.1**). Qingke has a long history of cultivation in the Qinghai-Tibet Plateau. The adaptation to extreme environmental conditions in high altitudes made it the staple food for Tibetans beginning at least 3,500–4,000 years ago [1]. It continues to be the predominant crop in Tibet, occupying ∼70% of crop lands.

Genome sequences provide a substantial basis for understanding the biological essences of crops associated with biologically and economically essential traits[2]. Therefore, it is necessary and crucial to construct the genome of a species. In recent years, with the development of NGS sequencing technology, many crops have published the genome draft or near-finished genome sequences, especially the larger genome of the wheat family[3-13]. At the same time, two studies have published two versions of the Qingke genome in 2015[2] and 2017[14], respectively. And the former assembly strain is Lasa Goumang , the latter is Zangqing320, all excellent cultivars of Qingke. The estimated genome size of the Qingke is 4.48Gb[2], which is much smaller than the 5.1Gb[4, 10] of barley. Although the former is a subspecies of barley, considering that the barley genome is rich in repeats(about 80.8%-84%[2, 4, 10]), the difference in genome size may come from the numbers and length of transposons.

Although the publication of the Qingke genome has attracted a new upsurge in Qingke genetics research[15, 16], due to the current poor genome assembly, such as the particularly short contig, the very low contig N50, and the poor genome annotation, the functional genome studying of Qingke has caused some vital problems. In addition, with the development of single-molecule sequencing platforms, such as those by Pacific Biosciences and Oxford Nanopore Technologies, have become available to researchers and are currently being increasingly applied in *de novo* genome assembly and genome update , and other scaffolding techniques(such as BioNano optical mapping, Chicago, Hi-C and 10X genomics)[17-19], it provides a new opportunity for improving Qingke genome assembly. Therefore, we *de novo* assembled a high-quality genome of Qingke using high-coverage Pacbio long reads, combining published Illumina paired-ends and mate-pairs

76 reads. And then, we applied PacBio Iso-seq to get high confidence full-length transcripts
77 and re-annotation Qingke protein-coding genes. This new assembly, annotations and newly
78 online Qingke genomics database discussed below will serve as valuable resources for
79 investigating the economic and genetic of Qingke and will also help the molecular breeders
80 to continually improve Qingke or barley germplasm resources.

81 DNA and RNA sequencing

DNA isolation, libraries construction and sequencing

The seedling of Qingke (NCBI BioSample ID **SAMN09914874**) were collected from
Tibet, China. After collection, tissues were immediately transferred into liquid nitrogen
and stored until DNA extraction. DNA was extracted using the Cetyltrimethyl Ammonium
Bromide (CTAB) method according to the protocal " Preparing Arabidopsis Genomic
DNA for Size-Selected ~20 kb SMRTbell^{TM} Libraries "(https://www.pacb.com/wp-
content/uploads/2015/09/Shared-Protocol-Preparing-Arabidopsis-DNA-for-20-kb-
SMRTbell-Libraries.pdf). The quality of the extracted genomic DNA was checked using
1% agarose gel electrophoresis, and the concentration was quantified using a Qubit
fluorimeter (Invitrogen, Carlsbad, CA, USA).

Single-molecule Real-time long reads sequencing was performed at NextOmics
Technology Corporation (Wuhan, China) with a PacBio Sequel Sequencer (Pacific
Biosciences, Menlo Park, CA,USA). The SMRT Bell library was prepared using a DNA
Template Prep Kit 1.0, and Six 20-kb SMRT Bell libraries were constructed(**Fig. S1**).
Genomic DNA（10ug）was mechanically sheared using a Covaris g-Tube with a goal of
DNA fragments of approximately 20kb. A Bioanalyzer 2100 12K DNA Chip assay was
used to assess the fragment size distribution. Sheared genomics DNA(5ug) was DNA-
damage repaired and end-repaired using polishing enzymes. A blunt-end ligation reaction
followed by exonuclease treatment was conducted to generate the SMRT Bell template. A
Blue Pippin device(Sage Science) was used to size select the SMRT Bell template and
enrich large fragments(>10kb).The size-selected library was quality inspected and
quantified on an Agilent Bioanalyzer 12Kb DNA Chip (Agilent) and a Qubit fluorimeter
(Life Technologies). A ready-to-sequence SMRT Bell Polymerase Complex was created

using a Binding Kit 2.0 (PacBio p/n 100-862-200), according to the manufacturer's
instructions. The PacBio Sequel instrument was programmed to load and sequenced the
sample on PacBio SMRT cells v2.0 (Pacific Biosciences), acquiring one movie of 360 min
per SMRT cell. The MagBead loading (Pacific Biosciences) method was used to improve
the enrichment of the larger fragments. A total of 64 SMRT cells were processed yielding
300.24G subread sequences, average 4.69Gb per cell, and 9,358bp in length(**Table S1**).

For Illumina sequencing data, we applied previously published paired-ends and mate-
pair library data[2](SRA no. SRA201388 , totally 819.37Gb clean data) for PacBio
assembly error correction and scaffolding, respectively.

RNA isolation and sequencing

Fresh roots, and leaves of Qingke were sampled in Tibet, China. Totally, 12
samples ,different N condition (Prepared for unpublished paper) were used for Iso-seq
sequencing. Total RNA was prepared by grinding tissue in TRIzol reagent (Invitrogen
15596026) on dry ice and processed following the protocol provided by the manufacturer.
To remove DNA, an aliquot of total RNA was treated with RQ1 DNase (Promega M6101),
followed by phenol/chloroform/isoamyl alcohol extraction, chloroform/isoamyl alcohol
extraction using Phase Lock Gel Light tubes (5 PRIME 2302800) and ethanol precipitation.
Precipitated RNA was stored at -20℃.

One microgram of total RNA per reaction per tissue was reverse transcribed using the
Clontech SMARTer cDNA synthesis kit. Specific barcoded oligo dT in separate PCR tubes,
to generate barcoded FL cDNA. Large-scale PCR products were purified with AMPure PB
beads and quality control (QC) was performed on a 2100 BioAnalyzer (Agilent).
Equimolar ratios of six cDNA libraries were pooled together. A total of 3.8 mg cDNA was
subjected to size fractionation using the SageELF system. Size fractions eluted from the
run were subjected QC and pooled in equimolar ratios for subsequent re-amplification to
yield four libraries (1–2, 2–3, 3–6 and 5–10Kb). The pooled PCR products were purified
using AMPure PB beads. One to five micrograms of purified amplicons were subjected to
Iso-Seq SMRT Bell library preparation (https://pacbio.secure.force.com/SamplePrep) .A
total of 17 SMRT cells (unpublished data) were sequenced on the PacBio RS II platform
using P6-C4 chemistry with 3–4 h movies.

Genome assembly

The Sequel raw bam files were converted into subreads in fasta format using the
standard Pacbio software package BAM2fastx. Totally, 300.24Gb PacBio data, covered
genome ~67X(estimated genome size 4.48Gb),was used to initially assemble.
Falcon assembly
we used falcon software package（https://github.com/PacificBiosciences/falcon ）
to construct primary assembly. Firstly, corrected the error reads by the overlap strategy
(parameter length_cutoff control is used to correct the wrong seed reads length);Secondly,
pre-assembled the corrected reads; Thirdly, Constructed contig with long reads obtained
from pre-assembly (parameter length_cutoff_pr controls the pre-assembled reads length
used to build contig); After many attempts, we determined that the following two reference
settings can get better assembly results: length_cutoff = 5000;length_cutoff_pr = 10000.
And, we get the initial assembly, 1.229Mb in length of contig N50.
Arrow polish
There are often errors in the preliminary assembly results of the genome, such as
SNP or InDel, which can be fixed by certain algorithms. Arrow is a genomic polishing tool
based on hidden Markov models. We use the following steps to polish the genome:
1) Align the PacBio sequencing reads to the genome assembly with blasr (pbalign
--algorithm=blasr, the rest of the parameters default);
2) Use Arrow to polish based on the alignment result (variantCaller --
algorithm=arrow, the rest of the parameters default).pbalign and variantCaller are from the
SMRT Link 4 toolkit (https://www.pacb.com/support/software-downloads/)
Scaffolding
The Scaffolding tool can group, sort, and orient contigs to join contigs into longer
sequence fragments, called scaffold. The sequence between some contigs may be unknown
but the distance between them can be roughly known, which is also helpful for subsequent
related work.

SSPACE[20] is a widely used tool for building scaffold. Firstly aligned the reads

to contigs using alignment tools (such as BWA, bowtie), and then built scaffold based on

the connection information between the contigs. This project uses Illumina data previously

published and PacBio data to build scaffold.

Firstly，using Illumina mate-pair library (20kb and 40Kb fragment long)

sequencing data to scaffolding contigs. Aligned the paired-end reads to contigs with

bowtie[21]; then, constructed scaffold (parameter: -p 1 -g 2) with SSPACE-STANDARD-

3.0.

Secondly, using PacBio data to continuously improving the scaffolds. Aligned

PacBio reads to the scaffold version of Illumina data using blasr; and then, built scaffold

with SSPACE-LongRead[22].

The final assembly produced 1,856 scaffolds, 1,092 of which greater than or equal

to 1Mb, with a contig N50 length of 1.563Mb and scaffold N50 of 4.006Mb. Compared to

the preliminary draft of the published two Qingke genomes[2, 14], the contiguity was

improved more than ~80 and 243 times, respectively(**Table 1**;**Table S2-S3**).

Assembly evaluation

First, the 682.57Gb Illumina paired-ends reads(library fragment 250,500 and

800,respectively) were mapped to new assembly, and 99.87-99.89% PE reads were mapped

concordantly (**Table S4**).

Using these short reads, the estimated quality value (QV) of Qingke genome was

calculated according to a previously described method[23]. The erroneous bases in the

genome assembly were identified using the variant calling software Genome Analysis

Toolkit (GATK) v4[24] with default parameters. The QV of the Qingke genome was

estimated to be 52(**Table S5**), which means that the accuracy of the assembly in base level

is very accurate after base correction.

We randomly downloaded 4 *Hordeum vulgare subsp. vulgare* (domesticated

barley) BAC clone sequencing data (NCBI ID: AC248476.1,AC250041.1,AC253057.1

and AC253422.1) to evaluate assembly quality. Using nucmer[25] mapping BAC data to

new assembly, the results manifested very high consistency.(**Fig. S2**).

In addition, we also used BUSCO[26] to perform an assessment of genome
completeness . BUSCO used a total of 1440 genes to evaluate the genome; the genome has
1312 complete BUSCO genes, accounting for 91.1%, of which 1288 have single copies,
accounting for 89.4%, 24 have multiple copies, accounting for 1.7%; 34 gene have only a
few fragments of the gene; 94 genes are missing from the genome.

Together, the results indicated that our dataset represented a genome assembly with
a high level of coverage, quality and completeness.

Repetitive sequences annotation

Repetitive DNA accounts for much of the remarkable variation in plant genome
sizes[27], especially some Gramineae crop plants, above 80% [10, 12, 28] of genome for
repetitive elements . Repetitive sequences are divided into transposons and simple repeats.
According to Tandem repeats finder program[29]，we marked simple repeat elements,
155,813,415 in length，3.89% of genome. For transposons, we performed homology-
based and de novo prediction, respectively. The software packages involved are
RepeatMasker[30],   RepeatProteinMask[30]   ,   repeatmodeler
(http://www.repeatmasker.org/RepeatModeler/) and ltr_finder[31]. The repetitive
database used here is Repbase 21.01[32](**Fig. S3**). Long terminal repeat(LTR)
retroelements, belong to class I elements, is accounted for 70.22% of genome. All repeats
together accounted for 87.48% of the genome(**Table 2**;**Table S6**).

Gene prediction

For the annotation of protein-coding gene, we used a combination of multiple data
sources, followed by a high confidence (HC) and low confidence (LC) classification of the
annotated genes(**Fig. S3-4**).

Combination of multiple annotations

According to Iso-seq methods, we obtained 39,442 high-quality full-length
transcripts. Theses high-quality transcripts were aligned to the genome using GMAP
(version 2017-05-08)[33], with a total of 14,099 perfect alignments; TransDecoder

v4.0.1(https://github.com/TransDecoder) was used to predict the coding region of 14,099

transcripts, of which 13,936 contained open reading frame (ORF), these ORF contained

transcripts are ultimately assigned to 9,360 genes, called perfect genes. We employed

Augustus v3.3[34],which used perfect genes as input train sets, to get 128,400 putative

genes. At the same time ,we downloaded representative proteins of 7 related species

(*Triticum aestivum A*[12]*, Triticum aestivum D*[9], *Brachypodium distachyon*[35],

Hordeum vulgare[10], *Oryza sativa*[36], *Sorghum bicolor*[37] and *Zea mays*[38]) from

public database(**Table S7**). Aligning the protein sequences of each species to the Qingke

genome using tblastn v2.6.0+[39]; and then, using Genewise v2.4.1[40] to predict gene

structure and obtain homologous gene sequences from various species in the Qingke

genome. In addition, using 16 transcriptome data (**Table S8**) from different tissues and

different conditions, we mapped the reads to Qingke genome by hisat2 v2.1.0[41] and

reference-based assembled the 47,490 transcripts by stringtie v1.2.4[42, 43]. The gene

information extracted from PacBio Iso-seq, Augustus, Homolog and RNA-seq was

integrated using EVidenceModeler[44], according to different weight parameters (**Table

S9**), to obtain 129,269 putative genes. The same RNA-seq reads were de novo assembled

using Trinity v2.4.0[45] to obtain 722,803 transcripts. Structural optimization of EVM

integration results was performed using PASA v2.1.0[46] and Trinity assembly results.

129,269 transcripts were annotated with NR, Uniprot, and Pfam database. And filter

the putative genes according to the function assignment result: (1) NR, Uniprot comparison

identity>=50%, coverage>=50%; (2) Not satisfying (1), but protein length >= 50, and

containing the Pfam domain. The tandem duplicated genes were identified using MCScanX.

Only the longest protein of each subgroup was kept to represent the corresponding gene.

The combine Qingke genome annotation included 65,833 genes with names derived from

BLAST analysis and genes with domain names derived from Pfam analysis.

High-confidence Genes

Stringent confidence classification was applied to all predicted genes to

discriminate between loci representing high-confidence (HC) protein-coding genes and

less reliable low-confidence (LC) genes, which potentially consisted of gene fragments,

putative pseudogenes and non-(protein)-coding transcripts. We assigned confidence values

to a gene model in a two-step procedure by using the similar criteria and methods described

previously[10](**Fig. S4**). Finally, we obtained 61,303 stringent confident putative genes, of
which 40,457 is HC genes(**Table 3**; **Fig. S4**).

Functional annotation of all 61,303 protein sequences was performed using NR,
KEGG, SwissProt, Trembl, GO, PFAM, and InterPro databases. 96.57% of all protein
sequences were assigned to at least one database(**Table S10**).

Non-coding Gene prediction

In the process of annotation of non-coding RNAs, tRNAscan-SE[47] software is
used to search for tRNA sequences in the genome according to the structural characteristics
of tRNA; since rRNA is highly conserved, the rRNA sequence of related species can be
selected as a reference sequence, through BLASTN Alignment to find rRNA in the genome;
in addition, using the Rfam family's covariance model, Rfam's own INFERNAL
software[48] can predict miRNA and snRNA sequence information on the genome(**Table**
4).

Annotation evaluation

The BUSCO[26] assessment showed that 91% of the 1,440 genes were completed
annotated. And RNA-seq reads rates mapping to gene set also have a significant
improvement, from 60% previously annotation to exceed 70%(**Table S8**).

In summary, we also acquired a high-quality genome annotation, and it will greatly
benefit future function genomics research.

Genome database

A new Qingke Genome Database(QGD) has been developed to facilitate studies
on Qinghai-Tibet Plateau crop plant genomics. The database currently provides access to
Qingke genome assembly V1,de novo assembly using NGS sequencing and V2,de novo
assembly improvement using combined TGS and NGS. Meanwhile, it also provides access
to genome annotation, including repetitive element annotation, ncRNA prediction and
protein-coding genes prediction(**Fig. 2**). The QGD embedded JBrowse[49] visual genomic
display module to display genome and gene annotation information. One can click on the
gene name on the screen to display specific information about the gene. One also can use

online tools Blast v2.4.0+ to enter the query sequence in fasta format, align and return the

visual aligned results. In the future, we will focus on optimizing the interactive operation

and visualization of the QGD.

You can access to the QGD by visiting http://show.genebang.com/project/index

Discussion

Qingke is a local species of barley and is of great significance to the life and culture

of the Tibetan. Its genome and annotation updates help future research on functional

genomics, and also help breeders to simplify their work and enrich the genomics resources

of barley. The generation of Qingke high-quality reference genomes also demonstrates the

superiority of PacBio long reads sequencing technology for larger genome assembly[50].

Although the genome has a long contig, the scaffold is still a bit short at the state-of-art of

the current genome assembly, mainly because of the complexity and data volume of the

scaffolding mate-pair library is relatively few. In the future, we will use BioNano optical

mapping, Hi-C, and akin weapon to complete the genomic construction at the chromosome

level[17]. Having an excellent Qingke genomics resources, it will enable geneticists to

focus more on functional genomics itself and breeding research.

Availability of supporting data

299   Raw genomic sequence reads are available in the NCBI Sequence Read Archive

300 under project number **SRS3725794**. Genome assembly and annotation file are available in

301 Qingke Genome Database (QGD) or contact the correspondence author. Supplementary

302 files can be downloaded from the GigaScience Website.

Abbreviations

BLAST: Basic Local Alignment Search Tool;

BUSCO: Benchmarking Universal Single-Copy Orthologs;

QV:quality value;

PacBio: Pacifc Biosciences;

RNA-seq: RNA sequencing;

NGS: Next generation sequencing

TGS: Third generation sequencing

QGD: Qingke Genome Database

Conflicts of Interest

313	The authors declare no conflict of interest.

Acknowledgments

This research was supported by the following funding sources: the Tibet
Autonomous Region Financial Special Fund 2017CZZX001, 2017CZZX002, XZNKY-
2018-C-021.

We acknowledge (1) Zuoyi Jian, genome assembly executor; (2) Hao Tang,
genome annotation pipeline writer; (3) Zhen Zeng ,repetitive annotation executor and (4)
NextOmics and BGI-Shenzhen for sequencing supports.

Shan Li, Zhenhua Zhuang and Jie Li also contributed to this Paper.

Author Contributions

Project coordination: X.Z.,X.L.,Y.W.,L.B. and N.T.(Leader).

Sampling and experimentation: X.Z.,Y.W.,H.Y.,Q.X.,S.Z.,Z.W.,W.Q.,Y.B.,J.D.,M.Y..

Genome assembly and Genome annotation guidance and performance：
X.L.(Co-leader) and his team.

Data analysis: T.X., X.Z.,Y.W.,H.Y.,Q.X.,S.Z.,Z.W.,W.Q.,Y.B.,J.D.,M.Y..

Genome database construction: T.X. and his team

Data submission: T.X.

Writing: X.L.(Co-leader).

All authors read and commented on the manuscript.

Reference

334   1.   Fu DX, Ruan RW, Dai XM and Liu YM. Astudy of ancient barley, wheat and millet
335   discovered at Changguo in Tibet. 2000.
336   2.   Zeng X, Long H, Wang Z, Zhao S, Tang Y, Huang Z, et al. The draft genome of
337   Tibetan hulless barley reveals adaptive patterns to the high stressful Tibetan
338   Plateau. Proceedings of the National Academy of Sciences of the United States
339   of America. 2015;112 4:1095-100. doi:10.1073/pnas.1423628112.
340   3.   Brenchley R, Spannagl M, Pfeifer M, Barker GL, D'Amore R, Allen AM, et al.
341   Analysis of the bread wheat genome using whole-genome shotgun sequencing.
342   Nature. 2012;491 7426:705-10. doi:10.1038/nature11650.
343   4.   International Barley Genome Sequencing C, Mayer KF, Waugh R, Brown JW,
344   Schulman A, Langridge P, et al. A physical, genetic and functional sequence
345   assembly of the barley genome. Nature. 2012;491 7426:711-6.
346   doi:10.1038/nature11543.
347   5.   Jia J, Zhao S, Kong X, Li Y, Zhao G, He W, et al. Aegilops tauschii draft genome
348   sequence reveals a gene repertoire for wheat adaptation. Nature. 2013;496
349   7443:91-5. doi:10.1038/nature12028.
350   6.   Ling HQ, Zhao S, Liu D, Wang J, Sun H, Zhang C, et al. Draft genome of the wheat
351   A-genome progenitor Triticum urartu. Nature. 2013;496 7443:87-90.
352   doi:10.1038/nature11997.
353   7.   International Wheat Genome Sequencing C. A chromosome-based draft
354   sequence of the hexaploid bread wheat (Triticum aestivum) genome. Science.
355   2014;345 6194:1251788. doi:10.1126/science.1251788.
356   8.   Avni R, Nave M, Barad O, Baruch K, Twardziok SO, Gundlach H, et al. Wild
357   emmer genome architecture and diversity elucidate wheat evolution and
358   domestication. Science. 2017;357 6346:93-7. doi:10.1126/science.aan0032.
359   9.   Luo MC, Gu YQ, Puiu D, Wang H, Twardziok SO, Deal KR, et al. Genome
360   sequence of the progenitor of the wheat D genome Aegilops tauschii. Nature.
361   2017;551 7681:498-502. doi:10.1038/nature24486.
362   10.   Mascher M, Gundlach H, Himmelbach A, Beier S, Twardziok SO, Wicker T, et al.
363   A chromosome conformation capture ordered sequence of the barley genome.
364   Nature. 2017;544 7651:427-33. doi:10.1038/nature22043.
365   11.   Zhao G, Zou C, Li K, Wang K, Li T, Gao L, et al. The Aegilops tauschii genome
366   reveals multiple impacts of transposons. Nat Plants. 2017;3 12:946-55.
367   doi:10.1038/s41477-017-0067-8.
368   12.   Ling HQ, Ma B, Shi X, Liu H, Dong L, Sun H, et al. Genome sequence of the
369   progenitor of wheat A subgenome Triticum urartu. Nature. 2018;557
370   7705:424-8. doi:10.1038/s41586-018-0108-0.

13. Appels R, Eversole K, Feuillet C, Keller B, Rogers J, Stein N, et al. Shifting the
limits in wheat research and breeding using a fully annotated reference
genome. Science. 2018;361 6403 doi:10.1126/science.aar7191.
14. Dai F, Wang X, Zhang XQ, Chen Z, Nevo E, Jin G, et al. Assembly and analysis of
a qingke reference genome demonstrate its close genetic relation to modern
cultivated barley. Plant Biotechnol J. 2018;16 3:760-70.
doi:10.1111/pbi.12826.
15. Wei Z, Zeng X, Qin C, Wang Y, Bai L, Xu Q, et al. Comparative transcriptome
analysis revealed genes commonly responsive to varied nitrate stress in leaves
of Tibetan hulless barley. Frontiers in plant science. 2016;7:1067.
16. Zeng X, Bai L, Wei Z, Yuan H, Wang Y, Xu Q, et al. Transcriptome analysis
revealed the drought-responsive genes in Tibetan hulless barley. BMC
genomics. 2016;17 1:386.
17. Sedlazeck FJ, Lee H, Darby CA and Schatz MC. Piercing the dark matter:
bioinformatics of long-range sequencing and mapping. Nat Rev Genet. 2018;19
6:329-46. doi:10.1038/s41576-018-0003-4.
18. Yuan Y, Bayer PE, Batley J and Edwards D. Improvements in Genomic
Technologies: Application to Crop Genomics. Trends Biotechnol. 2017;35
6:547-58. doi:10.1016/j.tibtech.2017.02.009.
19. Paajanen P, Kettleborough G, Lopez-Girona E, Giolai M, Heavens D, Baker D, et
al. A critical comparison of technologies for a plant genome sequencing project.
bioRxiv. 2017; doi:10.1101/201830.
20. Boetzer M, Henkel CV, Jansen HJ, Butler D and Pirovano W. Scaffolding pre-
assembled contigs using SSPACE. Bioinformatics. 2011;27 4:578-9.
doi:10.1093/bioinformatics/btq683.
21. Langmead B. Aligning short sequencing reads with Bowtie. Curr Protoc
Bioinformatics. 2010;Chapter 11:Unit 11.7.
doi:10.1002/0471250953.bi1107s32.
22. Boetzer M and Pirovano W. SSPACE-LongRead: scaffolding bacterial draft
genomes using long read sequence information. BMC Bioinformatics. 2014;15
1:211. doi:10.1186/1471-2105-15-211.
23. Xia M, Han X, He H, Yu R, Zhen G, Jia X, et al. Improved de novo genome
assembly and analysis of the Chinese cucurbit Siraitia grosvenorii, also known
as monk fruit or luo-han-guo. Gigascience. 2018;7 6
doi:10.1093/gigascience/giy067.
24. Heldenbrand JR, Baheti S, Bockol MA, Drucker TM, Hart SN, Hudson ME, et al.
Performance benchmarking of GATK3.8 and GATK4. bioRxiv. 2018;
doi:10.1101/348565.
25. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al.
Versatile and open software for comparing large genomes. Genome Biology.
2004;5 2:R12. doi:10.1186/gb-2004-5-2-r12.
26. Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV and Zdobnov EM.
BUSCO: assessing genome assembly and annotation completeness with single-
copy orthologs. Bioinformatics. 2015;31 19:3210-2.
doi:10.1093/bioinformatics/btv351.

27.   Paterson AH. Leafing through the genomes of our major crop plants: strategies
for capturing unique information. Nat Rev Genet. 2006;7 3:174-84.
doi:10.1038/nrg1806.
28.   Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, et al. The B73
maize genome: complexity, diversity, and dynamics. Science. 2009;326
5956:1112-5. doi:10.1126/science.1178534.
29.   Benson G. Tandem repeats finder: a program to analyze DNA sequences.
Nucleic Acids Research. 1999;27 2:573-80.
30.   Tarailo-Graovac M and Chen N. Using RepeatMasker to Identify Repetitive
Elements in Genomic Sequences. Current Protocols in Bioinformatics. 2009;25
1:4.10.1-4..4. doi:doi:10.1002/0471250953.bi0410s25.
31.   Xu Z and Wang H. LTR_FINDER: an efficient tool for the prediction of full-
length LTR retrotransposons. Nucleic Acids Research. 2007;35 suppl_2:W265-
W8. doi:10.1093/nar/gkm286.
32.   Bao W, Kojima KK and Kohany O. Repbase Update, a database of repetitive
elements in eukaryotic genomes. Mobile DNA. 2015;6 1:11.
doi:10.1186/s13100-015-0041-9.
33.   Wu TD and Watanabe CK. GMAP: a genomic mapping and alignment program
for mRNA and EST sequences. Bioinformatics. 2005;21 9:1859-75.
doi:10.1093/bioinformatics/bti310.
34.   Stanke M and Waack S. Gene prediction with a hidden Markov model and a
new intron submodel. Bioinformatics. 2003;19 suppl 2:ii215-ii25.
35.   Initiative IB. Genome sequencing and analysis of the model grass
Brachypodium distachyon. Nature. 2010;463 7282:763.
36.   Du H, Yu Y, Ma Y, Gao Q, Cao Y, Chen Z, et al. Sequencing and de novo assembly
of a near complete indica rice genome. Nature communications. 2017;8:15324.
37.   McCormick RF, Truong SK, Sreedasyam A, Jenkins J, Shu S, Sims D, et al. The
Sorghum bicolor reference genome: improved assembly, gene annotations, a
transcriptome atlas, and signatures of genome organization. The Plant Journal.
2018;93 2:338-54.
38.   Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, et al. Improved maize
reference genome with single-molecule technologies. Nature. 2017;546
7659:524.
39.   Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ. Basic local alignment
search tool. Journal of molecular biology. 1990;215 3:403-10.
doi:10.1016/s0022-2836(05)80360-2.
40.   Birney E, Clamp M and Durbin R. GeneWise and Genomewise. Genome
Research. 2004;14 5:988-95. doi:10.1101/gr.1865504.
41.   Kim D, Langmead B and Salzberg SL. HISAT: a fast spliced aligner with low
memory requirements. Nature Methods. 2015;12:357.
doi:10.1038/nmeth.3317
https://www.nature.com/articles/nmeth.3317#supplementary-information.
42.   Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT and Salzberg SL.
StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. Nature Biotechnology. 2015;33:290. doi:10.1038/nbt.3122
https://www.nature.com/articles/nbt.3122#supplementary-information.

43. Pertea M, Kim D, Pertea GM, Leek JT and Salzberg SL. Transcript-level
expression analysis of RNA-seq experiments with HISAT, StringTie and
Ballgown. Nature Protocols. 2016;11:1650. doi:10.1038/nprot.2016.095
https://www.nature.com/articles/nprot.2016.095#supplementary-information.
44. Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, et al. Automated
eukaryotic gene structure annotation using EVidenceModeler and the
Program to Assemble Spliced Alignments. Genome Biology. 2008;9 1:R7.
doi:10.1186/gb-2008-9-1-r7.
45. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-
length transcriptome assembly from RNA-Seq data without a reference
genome. Nature Biotechnology. 2011;29:644. doi:10.1038/nbt.1883
https://www.nature.com/articles/nbt.1883#supplementary-information.
46. Haas BJ, Delcher AL, Mount SM, Wortman JR, Smith Jr RK, Hannick LI, et al.
Improving the Arabidopsis genome annotation using maximal transcript
alignment assemblies. Nucleic acids research. 2003;31 19:5654-66.
47. Lowe TM and Eddy SR. tRNAscan-SE: a program for improved detection of
transfer RNA genes in genomic sequence. Nucleic acids research. 1997;25
5:955-64.
48. Nawrocki EP, Burge SW, Bateman A, Daub J, Eberhardt RY, Eddy SR, et al. Rfam
12.0: updates to the RNA families database. Nucleic acids research.
2014:gku1063.
49. Skinner ME, Uzilov AV, Stein LD, Mungall CJ and Holmes IH. JBrowse: a next-
generation genome browser. Genome Res. 2009;19 9:1630-8.
doi:10.1101/gr.094607.109.
50. Li C, Lin F, An D, Wang W and Huang R. Genome Sequencing and Assembly by
Long Reads in Plants. Genes. 2017;9 1 doi:10.3390/genes9010006.
488
489
490
491

Tables

493 **Table 1** Comparison of the new genome with previously published assemblies of the Qingke genome.

	This study	**Zeng *et al.*[2]**	**Dai *et al.*[14]**
Sequenced genome size (Gb)	4.00	3.89	3.73
Contig N50 (Kb)	1,563.00	18.07	5.94
Scaffold N50 (Kb)	4,006.00	242.00	171.11
Repeat proportion(%)	87.48	81.39	NA
Annotated protein-coding genes	61,303*	36,151	46,787

494 * Only refer to stringent confident putative protein-coding genes.
495
496
497 **Table 2** Repetitive element annotation statistics.

Type	**Repeat Size**	**% of genome**
Trf	155,813,415	3.89
Repeatmasker	2,935,487,965	73.27
Proteinmask	19,934,506	0.50
De novo	3,359,143,070	83.85
Total	3,504,452,012	**87.48**

499 **Table 3** Protein-coding genes prediction summary.

Method	Software and gene set	Gene number
PacBio	Smrtlink	39,442
	Gmap	14,099
	TransDecoder	13,936
Ab initio	Augustus	128,400
Homology-based	Triticum aestivum A	42,601
	Brachypodium distachyon	33,911
	Hordeum vulgare	46,312
	Oryza sativa	38,426
	Sorghum bicolor	29,909
	Triticum aestivum D	44,897
	Zea mays	29,872
RNA-seq	StringTie	47,490
Integration	EVidenceModeler	129,269
	PASA	129,269
Filter	NR、SwissProt、PFAM	68,018
	After MCScanX moved	65,833
	After TE moved	**61,303**

501 **Table 4** ncRNA genes annotation statistics.

Type		Copy	Average length(bp)	Total length(bp)	% of genome
miRNA		5,530	129.27	714,857	0.0178
tRNA		2,477	74.71	185,064	0.0046
rRNA	**Total**	1,583	151.69	240,131	0.0060
	18S	63	1,076.60	67,826	0.0017
	28S	174	133.20	23,177	0.0006
	5.8S	39	147.33	5,746	0.0001
	5S	1,307	109.70	143,382	0.0036
snRNA	**Total**	1,156	115.14	133,097	0.0033
	CD-box	831	103.44	85,960	0.0021
	HACA-box	132	133.48	17,620	0.0004
	splicing	193	152.94	29,517	0.0007

Figures

Figure 1 Morphological characteristics of the Qingke, Tibetan hulless barley. These pictures show (A),seedling;(B), heading stage; (C), mature stage;(D), filling stage spike and (E), grain of Qingke.

511
512 **Figure 2** Qingke Genome Database(QGD). (A),the home page of QGD;(B), Data download page;(C),gene
513 browse page and (D),sequences BLAST page.