The ortholog of chloroquine resistance transporter (TgCRT) plays a key role in maintaining the integrity of the endolysosomal system in *Toxoplasma gondii* to facilitate host invasion

L. Brock Thornton[1], Paige Teehan[1#a], Katherine Floyd[1¶], Christian Cochrane[1¶], Amy Bergmann[1¶], and Zhicheng Dou[1*]

1. Department of Biological Sciences, Clemson University, Clemson, South Carolina, United States of America

[#a]Current Address: Department of Biochemistry and Molecular Biology, The Pennsylvania State University, Pennsylvania, United States of America

*Corresponding author

E-mail: zdou@clemson.edu (ZD)

[¶]These authors contributed equally to this work.

Short title: Dynamics of the digestive vacuole in *Toxoplasma gondii*

Abstract

Toxoplasma gondii, with the ability to use foodborne, zoonotic, and congenital routes of transmission, is an apicomplexan parasite that can cause severe infectious disease in the immunocompromised human population. The parasites harbor a lysosome-like digestive vacuole, termed Vacuolar Compartment/Plant-Like Vacuole (VAC/PLV), which plays an important role in maintaining the lytic cycle and virulence of *Toxoplasma* parasites. The VAC supplies proteolytic enzymes that are required to maturate the parasite's invasion effectors and digest autophagosomes and endocytosed host proteins. Our previous study identified a *Toxoplasma* ortholog of malarial chloroquine resistance transporter that localized in the VAC. Here, we genetically ablate *TgCRT* and reveal that the TgCRT plays a key role in maintaining the integrity of parasite's endolysosomal system by controlling the morphology of the VAC. The VAC dramatically increases in size by ~15-fold and co-localizes with its adjacent endosome-like compartment when TgCRT is absent. To reduce the aberrant swelling, the parasites decrease the transcript and protein levels of endolysosomal proteases in Δ*crt* parasites. One subtilisin protease residing within the endolysosomal system was significantly reduced at both the transcription and translation stages, which impairs the trimming of micronemal proteins on the surface of the parasites, thereby decreasing parasite invasion. Our study also showed that chemical and genetic inhibition of proteolysis within the VAC reduces its size and partially restores the endolysosomal system, as well as invasion and micronemal protein trimming. In summary, these findings reveal for the first time a native role of this CRT ortholog in controlling the size of the VAC and integrity of the endolysosomal system in *Toxoplasma* parasites to facilitate parasite invasion of host cells. These observations also indicate the function of TgCRT in mediating the transport of small solutes to regulate osmotic pressure within the VAC. Our findings will help expand understanding of CRT orthologs in other apicomplexan parasites.

42 **Author Summary**

43   *Toxoplasma gondii* is an obligate intracellular protozoan parasite that belongs to the phylum

44   Apicomplexa and infects virtually all warm-blooded organisms. Approximately one-third of the human

45   population is infected with *Toxoplasma*. The parasites invade host cells by using processed invasion

46   effectors to disseminate infection. A lysosome-like digestive vacuole is involved in refining these invasion

47   effectors to reach their final forms. One malarial chloroquine resistance transporter-like protein was found

48   to span the membrane of the digestive vacuole in *Toxoplasma*. Although the mutated version of the

49   malarial chloroquine resistance transporter has been shown to confer resistance of chloroquine treatment

50   in malaria parasites, its native role in apicomplexan parasites remains unknown. In this study, we report

51   that this transmembrane protein plays a key role in regulating the integrity of the digestive vacuole and its

52   adjacent organelles in *Toxoplasma* parasites, further affecting proteolytic physiology within these

53   organelles, the secretion of invasion effectors, and parasite invasion and virulence. To relieve osmotic

54   stress caused by swelling within the digestive vacuole, the parasites take a strategy of repression of

55   proteolytic activities within this organelle to decrease the accumulation of small solutes in the lumen. Our

56   findings suggest a native function of this chloroquine resistance transporter ortholog in *Toxoplasma*

57   parasites. Further understanding the molecular mechanism used by *Toxoplasma* to maintain a functional

58   digestive vacuole could shed light on the identification of therapeutic targets for managing toxoplasmosis.

59

Introduction

Toxoplasma gondii, a protozoan pathogen, uses its polypeptide invasion factors to efficiently invade host cells. These proteins are stored in two unique organelles in *Toxoplasma* parasites, termed the microneme and rhoptry. The micronemal proteins need to undergo a series of proteolytic cleavages within the parasite's endosomal system, followed by further intramembrane cleavage and trimming on the parasite's cell membrane before secretion [1,2]. Therefore, proper maturation and secretion of micronemal proteins are crucial for efficient invasion of parasites [3-5].

Micronemal protein maturation and secretion are regulated by several proteases. During intracellular trafficking, the micronemal proteins are first cleaved by the aspartyl protease 3 (TgASP3) in a post-Golgi compartment for removal of their propeptide [5]. A cathepsin L-like protease (TgCPL) was also shown to process some micronemal proteins in the endosome-like compartment (ELC) of the parasites [4]. The mature micronemal proteins pass through the microneme and are subjected to further intramembrane cleavage and trimming on the parasite's surface. A plasma membrane-bound protease, rhomboid 4 (TgROM4), is a major protease required to process some micronemal proteins and release them from the cell surface of the parasites, including micronemal protein 2 (TgMIC2) and apical membrane antigen (TgAMA1) [6-9]. Subsequently, one subtilisin ortholog, TgSUB1, was shown to trim some micronemal proteins into their final forms, such as TgMIC2 and TgMIC2-associated protein (TgM2AP) [3]. Therefore, the precise control of proteolytic activities within the parasite's endosomal system and on the cell membrane is critical for the processing of parasite's invasion effectors and invasion.

Among these proteases, TgCPL and TgSUB1 are both localized in the parasite's endolysosomal system. TgCPL is located in an acidic digestive vacuole, termed the Vacuolar Compartment/Plant-like Vacuole (VAC/PLV, VAC hereafter for convenience) in *Toxoplasma* parasites [4,10]. Our previous studies showed that the genetic ablation of *TgCPL* causes defects in parasite invasion and acute virulence [4,11]. TgCPL becomes activated in the VAC, and a portion of TgCPL is delivered to the juxtaposed ELC for maturation of some micronemal proteins [4]. TgSUB1 is a micronemal protease and contains a GPI anchor to help it stay on the surface of the cell membrane of parasites [3]. TgSUB1 was shown to be activated in a post-ER compartment and enter the parasite's endolysosomal system before arrival to microneme [12].

The deletion of *TgSUB1* led to inefficient trimming of micronemal proteins on the parasite's surface, thereby

leading to invasion and virulence defects [3]. Hence, the maintenance of the integrity of the parasites'

endolysosomal system is critical for regulating the distribution and activity of endolysosomal proteases.

The VAC is an important organelle in the parasite's endolysosomal system. A dysfunctional VAC

results in the reduced invasion, replication, and virulence in acute infection of *Toxoplasma* [4,11]. Moreover,

the parasites are unable to turn over autophagosomes during chronic infection to sustain their survival in

the host's brain [13]. Despite its importance, the VAC has not been well characterized. Only 4 proteins

have been localized in the VAC, including TgCPL, TgCPB (a cathepsin B-like protein), TgCRT (a

Toxoplasma ortholog of chloroquine resistance transporter), and TgVP1 (a pyrophosphatase) [4,14-16].

The VAC forms an intact organelle during the initial infection and fragments throughout subsequent

intracellular replication [4]. It is still unknown how the parasites regulate the morphological changes of their

endolysosomal system. In a previous study, the *TgCRT* was knocked down in type I *Toxoplasma* parasites

using a tetracycline-inducible system [15], and it was observed that the VAC becomes swollen when

TgCRT is absent. Furthermore, the mutant parasites display fitness defects when TgCRT is repressed.

These findings indicate that the TgCRT is involved in controlling the morphology of the VAC.

Interestingly, the swollen phenotype of the TgCRT-deficient VAC mirrors the enlarged digestive

vacuole in chloroquine-resistant *Plasmodium falciparum*. An *in vitro* assay utilizing recombinant

Plasmodium CRT ortholog (PfCRT), reconstituted in proteoliposomes showed that the recombinant PfCRT

transports basic amino acids, synthesized hemoglobin-derived oligopeptides, and small drug molecules,

at the expense of a proton gradient potential [17]. It was also documented that the PfCRT[3D7] derived from

a chloroquine (CQ)-sensitive strain exhibits better transport efficiency than PfCRT[Dd2] and PfCRT[7G8] derived

from the CQ-resistant malaria strains. These findings suggest that the PfCRT could regulate the transport

of small solutes from the digestive vacuole to control its size and morphology. However, the inability to

successfully ablate the *PfCRT* gene in *Plasmodium* limits the understanding of its native function *in vivo*.

Here, we successfully delete the *TgCRT* gene in a Type I *Toxoplasma* parasite strain by double

crossover homologous recombination. The resulting mutant, Δ*crt,* displayed a swollen digestive vacuole,

which overlaps with the neighboring ELC. This aberrant organelle organization led to the reduction of

transcript and protein levels of several proteases residing in the parasite's endolysosomal system, further altering microneme secretion and resulting in defective parasite invasion and acute virulence. Collectively, these findings determine a novel role of TgCRT in regulating the size of the digestive vacuole to maintain the integrity of the parasite's endolysosomal system in *Toxoplasma* parasites. Our results may also lead to a better understanding of the functions of CRT orthologs in other apicomplexan parasites.

Results

1. The *TgCRT* is genetically ablated in *Toxoplasma gondii*.

TgCRT gene was documented to be refractory for genetic disruption and conditionally knocked down by using an anhydrotetracycline-regulated system in a Type I *Toxoplasma* RH strain [15]. The resulting inducible *TgCRT* knockdown parasites showed enlarged VAC and fitness defects when the transcription of *TgCRT* is repressed [15]. However, the incomplete depletion of *TgCRT* limits the further characterization of its role in the parasites. Here, we adopted a genetically tractable RH-derived strain, termed RHΔ*ku80* (hereafter referred to as WT), to produce a complete knockout of *TgCRT*. The RHΔ*ku80* strain represses the non-homologous end-joining DNA repair pathway to facilitate homology dependent DNA recombination [18]. Due to its increased efficiency of DNA homologous recombination, this strain has been widely used as a wild type strain in the *Toxoplasma* research field. We PCR amplified both the 5'- and 3'-untranslated regions (UTRs) (~1.5kb each) of the *TgCRT* gene, and flanked both at the 5'- and 3'-ends of a bleomycin (*BLE*) resistance cassette to assemble a *TgCRT* knockout plasmid. The resulting construct was transfected into WT parasites to replace the entire *TgCRT* gene with a *BLE* resistance cassette by double crossover homologous recombination (**Fig 1A**). To confirm the proper integration, PCR analysis was performed to detect the presence of the *BLE* cassette and the loss of *TgCRT* gene by using the primers shown in the schematic for the generation of the *TgCRT* knockout in Fig 1A. Amplification of ~1.6 kb fragments at both the 5'- and 3'-end integration regions (5'- and 3'-ARMs) were observed in the *TgCRT*-deletion mutant (Δ*crt*), and a ~1.5 kb *TgCRT* coding region was missing in the Δ*crt* strain (**Fig 1B**).

To complement the Δ*crt* parasites, we modified the pTub-TgCRT-mCherry-3xmyc plasmid (a gift from Dr. G. van Dooren, Australian National Laboratory), which over-expresses an mCherry-3xmyc epitope-

tagged *TgCRT* under the *Toxoplasma* tubulin promoter. A 1 kb DNA region containing the cognate promoter of the *TgCRT* gene was PCR-amplified and used to replace the tubulin promoter by molecular cloning to keep the transcription level of the complemented *TgCRT* the same as the endogenous *TgCRT*. The same pair of primers that were used to detect the loss of *TgCRT* in the Δ*crt* strain in Fig 1B were used to confirm the integration of the exogenously introduced *TgCRT* gene. Since the complemented *TgCRT* gene lacks introns, a ~0.2 kb PCR product was observed in the Δ*crtCRT* complementation strain, whereas a ~1.5 kb fragment was yielded in the WT strain (**Fig 1C**). To confirm that the transcript levels of *TgCRT* in the Δ*crtCRT* strain were comparable to the levels in WT parasites, a SYBR® Green-based quantitative PCR (qPCR) was used to quantify messenger RNA of *TgCRT* in WT, Δ*crt*, and Δ*crtCRT* strains. No *TgCRT* transcripts were observed in Δ*crt*, further validating the successful gene disruption in the Δ*crt* mutant. The transcript levels of *TgCRT* were similar between the WT and Δ*crtCRT* strains (**Fig 1D**). Collectively, we successfully ablated the *TgCRT* gene in *Toxoplasma* parasites, and properly restored the expression of the exogenously introduced *TgCRT* in the Δ*crtCRT* complementation strain in order to analyze the role of TgCRT in *Toxoplasma* parasites.

2. The TgCRT-deficient parasites lose the integrity of their endolysosomal system by altering the morphology of the VAC.

After obtaining the Δ*crt* mutant, we observed that purified extracellular Δ*crt* parasites exhibited large concave subcellular structures under differential interference contrast (DIC) microscopy, whereas WT and Δ*crtCRT* strains did not display this striking phenotype (**Fig 1E**). This enlarged subcellular structure was also observed in Δ*crt* parasites during pulse invasion (**Fig 1F**). To identify the swollen structure as the VAC, we stained the WT, Δ*crt*, and Δ*crtCRT* parasites with anti-TgCPL antibodies. TgCPL is a major luminal endoprotease in the VAC of *Toxoplasma* [4,19]. Immunofluorescence microscopy showed that the TgCPL staining nicely co-localized with the concave subcellular structure in the Δ*crt* mutant **(Fig 1F)**. The TgCPL staining in the Δ*crt* mutant was bigger than that in WT and Δ*crtCRT* parasites, indicating that the VAC becomes swollen when TgCRT is absent. We also quantified the sizes of the VACs based on the TgCPL staining as described previously [13,15]. The diameter of the VAC in the Δ*crt* parasites (1.12 ± 0.07 µm, mean ± standard deviation) is approximately 2.6-fold larger than that in WT parasites (0.43 ± 0.03 µm),

while the ∆crtCRT strain had a reduced VAC size (0.46 ± 0.02 µm) compared to that of WT parasites **(Fig**

1F). If we assume the shape of the VAC is a sphere, the VAC in ∆crt parasites is approximately 15-fold

larger in volume than that of WT parasites. In contrast to pulse invasion, the swollen concave structure

was not observed in ∆crt parasites after replication (**Fig 1G**). However, the TgCPL staining showed

differences between WT and ∆crt parasites (**Fig 1G**). As a dynamic organelle, the VAC fragments during

replication and TgCPL exhibits fragmented puncta staining by immunofluorescence microscopy [4].

However, TgCPL did not efficiently fragment and formed a major punctum in replicated ∆crt parasites.

Therefore, the loss of TgCRT altered the morphology and the dynamic property of the VAC in *Toxoplasma*

parasites.

The VAC is a lysosome-like organelle, participating in the parasite's endolysosomal system. It provides

a place for the maturation of TgCPL and delivers activated TgCPL to its adjacent endosome-like

compartment (ELC) to help process certain micronemal proteins required for parasite invasion [4]. It also

serves as a digestive vacuole to digest endocytosed proteins [4,11] [20]. Therefore, the dramatic swelling

of the VAC could affect the integrity of the parasite's endolysosomal system. We stained WT, ∆crt, and

∆crtCRT parasites with antibodies recognizing markers of the VAC (anti-TgCPL) and the ELC (anti-

proTgM2AP or TgVP1) [4]. During pulse invasion, the VAC and ELC displayed distinct subcellular staining

in WT and ∆crtCRT strains, whereas both markers partially co-localized in ∆crt parasites **(Fig 1G).** Similarly,

the major punctum of TgCPL in replicated ∆crt parasites also showed partial co-localization with both

proTgM2AP and TgVP1 (**Fig 1G**). These findings suggest that the TgCRT plays an essential role in

regulating the morphology of the VAC in *T. gondii*, further affecting the integrity of the parasite's

endolysosomal system.

3. TgCRT-null mutant displays reduced invasion and acute virulence.

Toxoplasma parasites utilize their endolysosomal system to perform exocytosis and endocytosis to

release invasion effectors, such as microneme and rhoptry proteins, for their host entry and ingest host

proteins for their intracellular growth, respectively [11,21-23]. We characterized invasion, replication, and

egress steps individually to understand how the arrested co-localization of the VAC and ELC affects the

lytic cycle of *Toxoplasma* parasites. First, we measured the invasion efficiency of parasites at 30 - 120 min

post-infection. At 30 min post-infection, the Δcrt mutant showed ~50% reduction in invasion compared to WT and $\Delta crtCRT$ **(Fig 2A)**. The differences in invasion efficiency between WT and Δcrt were reduced to ~20% at 60 min post-infection, and were not seen at 120 min post-infection **(Fig 2A)**, suggesting that Δcrt parasites have slower kinetics in invasion than the WT strain, but are still able to invade. Second, we conducted immunofluorescence microscopy to quantify parasite replication by counting the numbers of different-sized parasitophorous vacuoles (PVs). The infected cells were stained with DAPI and anti-TgGRA7 antibodies to help recognize the nuclei of individual parasites and the membranes of parasitophorous vacuoles, respectively. The average number of parasites per PV was also calculated in each strain to compare replication rates. There were no significant differences in parasite replication between WT and Δcrt parasites at 28 and 40 hrs post-infection **(Fig 2B)**. We also introduced NanoLuc® luciferase into WT, Δcrt, and $\Delta crtCRT$ parasites, and measured the fold-change of their luciferase activities at 72 hrs post-infection to calculate their growth rates. Similarly, we did not observe growth differences between WT and Δcrt at 24, 48, and 72 hrs post-infection **(Fig 2B)**. Third, the egress efficiency of each strain was determined by a lactate dehydrogenase release-based assay. The parasites were incubated with 0.5 mM Zaprinast for 5 min to induce egress. The egressed parasites disrupted host cell membranes to release lactate dehydrogenase that was subsequently quantified to reflect the number of egressed PVs. We did not observe egress defects in the TgCRT-deficient parasites **(Fig 2C)**. Last, we determined the acute virulence of Δcrt parasites in a murine model. The outbred CD-1 mice were infected with a subcutaneous or intravenous inoculum of 100 WT, Δcrt, or $\Delta crtCRT$ parasites. Thirty percent of mice infected with the Δcrt mutant survived when mice were infected by subcutaneous injection, while the WT and $\Delta crtCRT$ infections led to the mortality of mice in 10-12 days post-infection **(Fig 2D)**. Similarly, the mice receiving WT parasites by intravenous inoculation showed mortality starting at 13 days post-infection and all were dead in 20 days. The Δcrt and $\Delta crtCRT$ parasites caused death in 40% and 80% of infected mice, respectively **(Fig 2D)**. The ability of Δcrt parasites to survive in the host was statistically higher than that of WT parasites, regardless of the mode of inoculation. The statistical analysis showed that the mice infected with WT and Δcrt parasites showed significantly different survival curves. Seroconversion of the surviving mice was confirmed by ELISA. We also challenged the surviving mice with 1000 WT parasites

by subcutaneous injection, and did not observe lethality after 30 days post-challenge. These findings

indicate that the pre-inoculation of Δ*crt* parasites conferred immunological protection to the subsequent

acute toxoplasmosis. Collectively, our findings revealed that *Toxoplasma* parasites require the TgCRT

protein for optimal invasion and acute virulence but not for replication and egress.

4. Microneme secretion is impaired in the Δ*crt* mutant.

During infection, *Toxoplasma* parasites sequentially secrete proteins to facilitate host invasion. The

micronemal proteins are the first secreted among these invasion effectors. Micronemal proteins traffic

through the parasite's endolysosomal system and undergo intracellular maturation, intramembrane

cleavage, and cell surface trimming before reaching their final forms, which facilitate invasion [3-9]. To test

which step(s) was/were affected in the Δ*crt* parasites, we probed cell lysates and excretory secretory

antigen (ESAs) fractions of each strain with anti-TgMIC2, TgM2AP, and TgMIC5 (three representative

micronemal proteins) by immunoblotting to measure their steady abundances and secretion patterns. The

migration patterns of these three micronemal proteins in cell lysates were very similar among the strains.

The abundances of the individual micronemal proteins in cell lysates were normalized against the protein

level of *Toxoplasma* actin protein by densitometry and plotted for quantification. The expression levels of

these micronemal proteins were comparable in the cell lysates of individual strains **(Fig 3A)**. To evaluate

the patterns and abundances of the secreted forms, we probed constitutive and induced ESAs with anti-

TgMIC2, TgM2AP, and TgMIC5 antibodies. The constitutive and induced ESAs were generated by

incubating purified parasites in D3 medium (DMEM medium supplemented with 3% (v/v) cosmic calf serum)

for 30 min at 37ºC or D3 medium supplemented with 1% (v/v) ethanol for 2 min at 37ºC, respectively. In

the ESAs secreted by WT parasites, TgMIC2 exhibited two bands migrating at 100 kDa and 95 kDa, while

TgM2AP showed 4 proteolytically processed polypeptides in addition to the pro- and mature forms.

However, TgMIC2 only existed as a 100 kDa band in Δ*crt* parasites. Furthermore, mature TgM2AP was

not processed in the constitutive ESAs of the Δ*crt* strain and showed significantly reduced processing in

the induced ESAs of Δ*crt* parasites **(Fig 3B)**. The secreted TgMIC5 protein displayed similar migration

patterns among all three strains **(Fig 3B)**. The secretion of these micronemal proteins was also quantified

by normalizing the relative abundances of the proteins against the amount of secreted TgGRA7, a dense

granule protein. The secretion of TgMIC2, TgM2AP, and TgMIC5 were reduced by ~80%, 35%, and 40%,

respectively, in the induced ESAs of Δ*crt* parasites, compared to the WT strain. The differences in the

amount of microneme secretion were less significant in the constitutive ESAs. Only TgMIC2 secretion was

decreased by ~55% in Δ*crt* parasites compared to the WT strain, while TgM2AP and TgMIC5 did not show

differences **(Fig 3B)**. To examine whether the abnormal secretion of micronemal proteins alters their

intracellular trafficking patterns, following pulse invasion and replication, we stained parasites with TgMIC2

and TgM2AP antibodies. Both microneme proteins trafficked to the apical end of the parasites and showed

normal staining patterns (**Fig 3C**).

Before secretion, some membrane-anchored micronemal proteins are released via proteolytic

cleavage by intramembrane rhomboid proteases, such as TgROM4. The deletion of TgROM4 leads to the

retention of some micronemal proteins on the parasite's plasma membrane, such as TgMIC2 and TgAMA1

(*Toxoplasma* apical membrane antigen 1) [6-9]. To test whether the aberrant endolysosomal system alters

the retention of micronemal proteins on the surface of parasites, we stained purified, non-permeabilized,

extracellular parasites with anti-TgMIC2 antibody. Immunofluorescence microscopy did not reveal excess

TgMIC2 on the plasma membrane of Δ*crt* parasites **(Fig S1)**, suggesting that the reduced secretion of

micronemal proteins is not due to inefficient intramembrane cleavage of micronemal proteins on the

parasite's plasma membrane.

The endosome-like compartment is involved not only in the trafficking of micronemal proteins, but also

rhoptry contents [24]. We stained newly invaded and replicated parasites with anti-TgROP7 antibodies to

examine the trafficking of rhoptry proteins and the morphology of the rhoptry. The TgROP7 staining

revealed typical rhoptry patterns located at the apical end of the parasites (**Fig 3D**), excluding the possibility

of aberrant trafficking of rhoptry contents and possible defects in the biogenesis of the rhoptry. Taken

together, our data suggest that the invasion defects in Δ*crt* parasites are caused by the inefficient secretion

and defective trimming of micronemal proteins, while the abnormal morphology of the VAC does not affect

the morphology of the rhoptry and the intracellular maturation, trafficking, and intramembrane cleavage of

micronemal proteins.

5. Transcript and protein levels of TgSUB1 are decreased in the TgCRT-deficient parasites.

The inefficient proteolytic processing of TgMIC2 and TgM2AP in the ESAs led us to investigate the

abundance of *Toxoplasma* subtilisin 1 (TgSUB1) in the Δ*crt* parasites. A previous publication reported that

parasites lacking TgSUB1 showed defective trimming patterns of secreted micronemal proteins, such as

TgMIC2 and TgM2AP [3], which are very similar to the secretion patterns seen in the Δ*crt* mutant. Therefore,

we quantified the abundance of secreted TgSUB1 in both constitutive and induced ESAs by probing them

with anti-PfSUB1 antibody, which was generated against *Plasmodium* SUB1 proteins but cross-reacts with

TgSUB1 [25]. Immunoblotting analysis revealed that there was no detectable TgSUB1 in the ESAs of Δ*crt*

parasites **(Fig 4A).** Non-permeabilized, extracellular parasites were also stained with anti-PfSUB1 to

evaluate the amount of surface-anchored TgSUB1. Similarly, there was no detectable TgSUB1 staining

on the plasma membrane of Δ*crt* parasites (**Fig 4B**). These data suggest that TgSUB1 is not efficiently

delivered to the surface of parasites in the Δ*crt* mutant.

TgSUB1 is a micronemal protein. It passes through the parasite's endolysosomal system and becomes

activated during trafficking [3,12]. The aberrant endolysosomal system in Δ*crt* parasites could potentially

alter the intracellular trafficking of TgSUB1 to prevent its delivery to the   microneme. In addition, the

aberrant endolysosomal system could alter the maturation process of TgSUB1 and reduce its expression.

To test both possibilities, first, following pulse invasion or replication, we stained parasites with anti-PfSUB1

antibodies to examine the intracellular trafficking pattern of TgSUB1. TgMIC5 was also stained as a

reference for typical microneme staining. Interestingly, we observed much less TgSUB1 staining in Δ*crt*

parasites compared to the WT strain (**Fig 4C**). Next, we quantified the steady protein abundance of

TgSUB1 in the cell lysates of parasites by immunoblotting, and found that the total abundance of TgSUB1

was decreased by approximately 90% in Δ*crt* parasites compared to WT parasites **(Fig 4D)**. To further

understand how TgSUB1 expression was suppressed in the Δ*crt* mutant, we performed qPCR to measure

the messenger RNA levels of *TgSUB1* in WT, Δ*crt*, and Δ*crtCRT* parasites. Not only was there a reduction

of protein expression in Δ*crt* parasites, but the transcripts of *TgSUB1* were also reduced ~10-fold in

Toxoplasma parasites lacking *TgCRT* **(Fig 4E)**. Collectively, our findings suggest that the arrested co-

localization of the VAC and ELC dramatically decreases the abundance of TgSUB1 protein, which further

alters the proteolytic processing of micronemal proteins on the surface of parasites during secretion to

ultimately reduce invasion efficiency.

6. The morphological alteration of the VAC led to reduced transcript and protein levels of several

endolysosomal proteases.

The swollen VAC, and its aberrant co-localization with the ELC in Δ*crt* parasites, could result in the

global alteration of gene transcription in order to help the parasites adapt to the sub-optimal

microenvironment within their endolysosomal system. We conducted transcriptome sequencing to detect

global alteration of gene transcription in WT and Δ*crt* parasites. The differential gene expression analysis

identified 102 genes whose transcript levels changed greater than 1.5-fold in the Δ*crt* strain. Forty-six and

fifty-six genes had increased and reduced transcripts, respectively (**Fig 5A and Table S1**).

Four proteases were among the list of genes showing reduced transcripts in the Δ*crt* mutant, including

one putative aminopeptidase N protein (TgAMN, TGGT1_221310), one putative Pro-Xaa serine

carboxypeptidase (TgSCP, TGGT1_254010), aspartyl protease 1 (TgASP1, TGGT1_201840), and an ICE

family protease-like protein (TgICEL, TGGT1_243298). We validated the transcript levels of these

proteases in WT, Δ*crt*, and Δ*crtCRT* strains by quantitative PCR (qPCR), and also quantified the transcript

levels of two known VAC luminal proteases, TgCPL and TgCPB. The qPCR analysis showed that the

transcript levels of TgAMN, TgSCP, TgASP1, and TgCPB were decreased by 50%, 20%, 47%, and 14%,

respectively, in Δ*crt* parasites (**Fig 5B**). The protein levels of TgCPL and TgCPB were quantified and

compared in WT, Δ*crt*, and Δ*crtCRT* parasites by immunoblotting analysis. Although the transcript level of

TgCPL did not show any difference between the WT and Δ*crt* strains, the protein abundance of TgCPL

decreased by ~25% in the Δ*crt* mutant (**Fig 5C**). The expression level of TgCPB protease was reduced

significantly in Δ*crt* parasites. Both the pro- and mature forms of TgCPB in the Δ*crt* mutant showed reduced

abundance compared to WT parasites. The level of mature TgCPB was quantified by densitometry and

was found to decrease by ~60% in the TgCRT-null mutant (**Fig 5C**).

To determine the subcellular locations of these down-regulated putative proteases, we endogenously

tagged TgAMN and TgSCP with 3xHA and 3xmyc epitope tags at their C-termini, respectively, in WT

parasites (**Fig S2**). After drug-selection, we probed the cell lysates of the tagged strains with anti-HA and

anti-myc antibodies, respectively, to confirm their expression. Immunoblotting analysis revealed that the

observed molecular weights of both proteins were similar to the predicted sizes based on their primary

sequences (**Fig 5D**). Next, the tagged strains were stained with antibodies recognizing their respective

epitope tags along with anti-TgCPL and anti-VP1 antibodies to determine their subcellular locations.

Immunofluorescence microscopy revealed the subcellular locations of both TgSCP and TgAMN to be in

the VAC/ELC (**Fig 5D**). The subcellular location of TgASP1 has been determined in a separate study. By

probing WT parasites with polyclonal anti-TgASP1 (recently made in our lab) and anti-TgCPL antibodies,

we identified TgASP1 as a new VAC-localizing protease (data not shown since this work is being prepared

for another publication). Collectively, our data suggest that the swollen VAC in Δ*crt* parasites causes

reduced expression of several endolysosomal proteases at both the transcription and translation levels.

Reduced proteolytic activities within the VAC could decrease the production of degraded protein products,

thereby limiting the swelling of the VAC.

7. Suppression of proteolysis within the swollen VAC in Δ*crt* parasites partially restores the size of

the VAC and the separation of organelles within the parasite's endolysosomal system, thereby

moderately restoring invasion.

Since the parasites lacking TgCRT reduced proteolytic activities within the VAC in order to decrease

the production of small solutes, it is expected that the inhibition of proteolysis in the swollen VAC would

reduce its size. We tested this hypothesis by chemically and genetically suppressing the proteolysis of the

VAC. First, we treated WT, Δ*crt*, and Δ*crtCRT* parasites with 1 μM LHVS, an irreversible inhibitor of the

TgCPL protease [19], or with the vehicle control DMSO. TgCPL is a major endopeptidase and is involved

in the maturation of micronemal proteins and digestion of host proteins [4,11]. Infected host cells were

incubated with LHVS for 48 hrs to allow full inhibition of the TgCPL protease in the parasites. Treated

parasites were liberated from host cells and used to infect new host cells for 30 minutes, followed by TgCPL

staining to quantify the size of the VAC. As expected, the LHVS-treated Δ*crt* parasites displayed smaller

VACs than DMSO-treated Δ*crt* parasites **(Fig 6A)**. The cathepsin B-like protease (TgCPB) is another

known VAC-localizing protease, displaying both endo- and exo-peptidase activities [14,19]. Due to its

carboxypeptidase activity, it is expected that TgCPB contributes to the generation of small solutes to a

greater extent than TgCPL. We genetically deleted *TgCPB* in Δ*crt* parasites using a CRISPR-Cas9 editing

tool to generate a Δ*crt*Δ*cpb* double knockout **(Fig 6B)**. The replacement of *TgCPB* with a pyrimethamine

resistance cassette was confirmed by PCR and immunoblotting **(Fig 6B)**. The resulting Δ*crt*Δ*cpb* mutant

showed a smaller concave subcellular structure compared to the Δ*crt* mutant **(Fig S3)**. The size of the VAC

in WT, Δ*crt*, and Δ*crt*Δ*cpb* was quantified based on the TgCPL staining as described above. The size of

the VACs in Δ*crt*Δ*cpb* parasites (0.85 ± 0.15 μm) were reduced by ~30% compared to that of Δ*crt* parasites

(1.15 ± 0.10 μm) **(Fig 6C)**. The moderate decrease in the size of the VAC in the Δ*crt*Δ*cpb* strain also

reduced the number of parasites showing partial overlapping between the VAC and ELC. Approximately

44% of Δ*crt*Δ*cpb* parasites showed partial co-localization between TgCPL and proTgM2AP staining

compared to 62% in the Δ*crt* strain (**Fig 6D**), but this was still significantly higher than 19% and 25% in WT

and Δ*crt*CRT strains, respectively. Moreover, we detected that TgSUB1 showed comparable expression

in both the WT and Δ*crt*Δ*cpb* strains (**Fig 6E**). Similarly, TgSUB1 was observed in both constitutive and

induced ESA fractions in the Δ*crt*Δ*cpb* parasites (**Fig 6F**). TgM2AP and TgMIC2 were cleaved by TgSUB1

in Δ*crt*Δ*cpb*, and their secretion patterns were similar to those seen in the WT strain (**Fig 6F**). These

restored phenotypes in the Δ*crt*Δ*cpb* mutant improved the invasion efficiency by ~60% compared to the

Δ*crt* strain, although the invasion efficiency was still significantly lower than that of WT parasites **(Fig 6G)**.

This set of data exhibits a closely linked association among the size of the VAC, altered morphology of the

parasite's endolysosomal system, protein abundance of TgSUB1, and parasite invasion. In summary, our

findings strongly suggest a role of TgCRT in the transport of small solutes across the VAC membrane to

regulate the osmotic pressure within this organelle and govern the proper segregation of other organelles

within the endolysosomal system, which facilitates microneme secretion and parasite invasion. It also

indicates that the invasion deficiency exhibited in the Δ*crt* mutant is likely caused by multiple determinants,

since the recovery of TgSUB1 expression and micronemal trimming in Δ*crt*Δ*cpb* did not fully reverse the

invasion defects.

Discussion

As a eukaryotic, single-celled pathogen, *Toxoplasma* parasites utilize their endolysosomal system to

secrete invasion effectors to help disseminate infection. These invasion effectors undergo a series of

proteolytic cleavage and trimming to reach their final forms. Therefore, maintenance of the integrity of the

endolysosomal system is critical for controlling the secretion of invasion effectors in *Toxoplasma*. The

Vacuolar Compartment (VAC) is a lysosome-equivalent structure in the parasites, which harbors an acidic

environment and exhibits proteolytic activity. The deletion of a cathepsin L-like protease (TgCPL), a major

VAC luminal endopeptidase, leads to invasion, replication, and virulence defects [4,11]. The compromised

proteolytic activities within the VAC also result in the inefficient degradation of endocytosed host proteins,

suggesting its involvement in endocytic activities within parasites [11]. Sally *et al.* previously reported that

a *Toxoplasma* ortholog of the chloroquine resistance transporter (TgCRT) resides in the VAC [15]. The

knockdown of TgCRT leads to swelling of the VAC [15]. It is still unknown how the swelling of the VAC

affects the parasite's endolysosomal system. The incomplete depletion of the TgCRT protein in the

knockdown system limits the ability to fully understand its function in the VAC and its role in parasite

infection.

In this study, we genetically ablated *TgCRT* in *Toxoplasma*, and the resulting Δ*crt* had a dramatic

increase in the size of its VAC, which also aberrantly co-localizes with its adjacent endolysosome-like

compartment **(Fig 7)**. A previous study reported that the parasites deliver a minute amount of TgCPL from

the VAC to the ELC which contributes to the maturation of some micronemal proteins [4]. The aberrant co-

localization of the VAC and ELC could potentially bring excess TgCPL into the ELC, which might interfere

with the cleavage of micronemal proteins. Our findings did not reveal abnormal intracellular cleavage or

trafficking of micronemal proteins in Δ*crt* parasites, indicating that the ELC may not be a major site for the

removal of the propeptide of micronemal proteins. Dogga *et al.* recently documented that aspartic acid

protease 3 (TgASP3) localizes in a post-Golgi compartment and serves as a major maturase for invasion

effectors [5], suggesting that the cleavage of micronemal proteins by TgCPL in the ELC plays only a minor

role in their maturation. We also measured the retention and secretion of micronemal proteins on the

parasite's surface and in the medium, respectively, which is executed by TgROM4 and TgSUB1,

respectively. We found that the micronemal proteins were not properly trimmed on the surface of the Δ*crt*

parasites during their secretion. The patterns of the secreted micronemal proteins that were observed in

the Δ*crt* mutant were similar to those seen in Δ*sub1* parasites, which led us to examine the expression of

TgSUB1 in Δ*crt* parasites and ESAs. As expected, the protein level of TgSUB1 was decreased on the

surface of Δ*crt* parasites and in the medium during secretion. Interestingly, the steady state protein

abundance of TgSUB1 was also significantly decreased in the Δ*crt* mutant. Surprisingly, we found that the

reduction of TgSUB1 was due to a decrease in the transcription levels of TgSUB1 in the Δ*crt* strain,

suggesting that the parasites may utilize a feedback mechanism in order to regulate TgSUB1 expression

at the transcriptional stage.

TgSUB1 is a micronemal protein which is anchored on the surface of *Toxoplasma* parasites via a GPI

anchor. It remains unclear how TgSUB1 becomes activated within *Toxoplasma* parasites. Previous pulse-

chase experiments have revealed that TgSUB1 undergoes maturation in a post-ER compartment, and

passes through the endolysosomal system before its arrival at the microneme and subsequent secretion

[12]. The propeptide region of TgSUB1 carries targeting information which helps to guide the protein to the

microneme [26]. The propeptide may also function by binding to active sites of mature TgSUB1 to inhibit

its proteolytic activity during trafficking. Co-localization of the VAC and ELC could bring propeptide-bound

TgSUB1 to a protease-abundant environment, where non-specific digestion of the propeptide could then

lead to increased digestive activities in the VAC and ultimately result in an increase in the osmotic pressure

within the hybrid VAC/ELC organelle **(Fig 7)**. During this scenario, the parasites may utilize a feedback

mechanism to repress the expression of TgSUB1 in order to alleviate the osmotic pressure and avoid

further swelling. Moreover, we also discovered that the Δ*crt* parasites had reduced protein and/or transcript

levels of several endolysosomal proteases, including two known VAC proteases, TgCPL and TgCPB.

Therefore, the parasites down-regulate a number of endolysosomal proteases to suppress proteolytic

activities in the swollen VAC in order to reduce osmotic pressure within the lumen, thereby controlling VAC

size and morphology. Among these proteases, TgSUB1 has been shown to be involved in parasite

invasion and virulence defects but not replication and egress [3]. Additionally, TgCPL plays a role in

parasite invasion by maturating several micronemal proteins [4]. Therefore, the invasion defects exhibited

in the Δ*crt* mutant could be due to a multitude of factors.

The alteration of transcript levels of endolysosomal proteases in Δ*crt* parasites suggests that the

parasites may repress transcription factors or enhance transcription repressors to respond to the increase

of the VAC size. RNA-Seq analysis did not reveal any members of an AP2-family of transcription factors

that had a change in their transcript levels greater than 1.5-fold. Given that the abundance of transcription

factors is generally low in cells, the RNA-seq analysis may not be sensitive enough to detect differences

in the transcriptional changes between the WT and Δ*crt* strains. In mammalian cells, the transcription factor

EB (TFEB) is a master regulator that drives gene expression for autophagy and lysosome biogenesis [27].

The ortholog search against the annotated coding sequences of *Toxoplasma* did not result in the

corresponding counterparts, suggesting that the parasites could adopt an alternative strategy to regulate

lysosomal gene expression. Interestingly, our differential gene expression analysis identified that the

transcript levels of two zinc finger (CCCH) type motif-containing proteins, TGGT1_246200 and

TGGT1_226310, were increased and decreased by 2-fold and 3-fold, respectively, in the Δ*crt* mutant. The

CCCH type zinc finger motif-containing protein is known to regulate the stability of mRNA [28]. For example,

tristetraprolin inhibits the production of tumor necrosis factor-α in macrophages by destabilizing its mRNA

via an interaction with its AU-rich elements at the 3'-untranslated region [29]. Further investigation to

identify transcription factor(s) and regulator(s) that govern the expression of lysosomal genes will help

elucidate how *Toxoplasma* parasites regulate the biogenesis of the VAC.

In this study, we determined that the TgCRT-deficient parasites have reduced expression of several

endolysosomal proteases. We also found that the suppression of proteolytic activities within the swollen

VAC decreases its size. These findings support the idea that CRT functions as an amino acid/peptide

transporter to regulate the osmotic pressure inside the digestive vacuole. When TgCRT is absent on the

membrane of the VAC, decomposed protein products accumulate within the VAC and increase the osmotic

pressure, thereby leading to the swollen phenotype. Chemical inhibition of proteolysis via the small inhibitor

LHVS dramatically reduces the size of the VAC. LHVS principally targets TgCPL in the parasites, but also

binds to TgCPB protease [14]. Additionally, the maturation of TgCPB is dependent upon the presence of

TgCPL [14]. Therefore, the treatment of LHVS blocks both of these VAC proteases. Since the TgCPB

protease exhibits endo- and exo-peptidase activities and the deletion of TgCPB had no effect on the

abundance of TgCPL [14], we genetically ablated *TgCPB* to understand its role in regulating the

morphology of the VAC. The deletion of TgCPB partially restored the size of the VAC, secretion patterns

of micronemal proteins, and invasion defects. These results reveal for the first time that TgCPB plays an

active role in contributing to proteolysis within the VAC in *Toxoplasma* parasites.

Our RNA-Seq analysis also identified several other genes with altered transcription levels, suggesting

that the parasites may utilize other strategies to control the size of the VAC. For example, the transcript

levels of a water channel protein, aquaporin (TGGT1_215450), was reduced in Δ*crt* parasites. A previous

study has reported that this aquaporin protein is localized in the VAC/PLV [30]. Therefore, the Δ*crt*

parasites may express less aquaporin to reduce water intake into the VAC/PLV as a tactic to limit the

swelling of the organelle. We also found that two putative protein phosphatase 2C orthologs

(TGGT1_276920 and TGGT1_201520) became down-regulated in the Δ*crt* mutant. Both of these proteins

carry signal peptides, indicating that they traffic through the parasite's endosomal system. Based on a

sequence-based ortholog search, TGGT1_276920 and TGGT1_201520 are homologous to PTC3 and

PTC1 in *Saccharomyces cerevisiae,* respectively. Interestingly, both PTC1 and PTC3 proteins are involved

in the regulation of osmosensing in yeast. A mitogen-activated protein kinase pathway is activated when

yeast cells experience hyperosmotic conditions. PTC1 and PTC3 were shown to negatively regulate this

pathway [31,32]. Furthermore, PTC1 was found to control the function and morphology of the vacuole in

yeast cells, which further alters its biogenesis [33]. The dramatic change in the volume of the VAC could

induce osmotic stress in the Δ*crt* parasites. The parasites may utilize a similar mechanism to suppress

both of these protein phosphatases, thereby enhancing the activity of a similar signaling pathway to help

them adapt to such stress.

The phenotype of the swollen VAC in the Δ*crt* strain mirrors the enlarged digestive vacuole in

chloroquine (CQ) resistant malaria strains [34]. Thus far, the CRT gene could not be deleted in malaria

parasites. Peptidomics analysis showed that hemoglobin cannot be efficiently degraded within the

digestive vacuole in CQ resistant malaria parasites [35], suggesting that the mutations in malarial CRT

may alter the physiological conditions within the swollen digestive vacuole, thereby also compromising its

proteolytic activities. An *in vitro* assay utilizing recombinant malarial CRT proteins that were reconstituted

in proteoliposomes revealed that PfCRT acts as a proton gradient dependent, poly-specific nutrient

exporter for small solutes, such as amino acids, oligopeptides, glutathione, and small drugs [17]. This study

also demonstrated that the PfCRTs derived from the CQ resistant malaria strains displayed less efficient

transport of small solutes than their counterpart in the CQ sensitive strain [17]. The alteration of proteolytic

activities in the enlarged VAC in the Δ*crt* mutant may reflect a similar scenario that occurred in the digestive

vacuole of the CQ resistant malaria parasites. Given the similarity in the components and functionality of

digestive vacuoles between *Toxoplasma* and *Plasmodium* parasites, the *Toxoplasma TgCRT*-deficient

mutant could be used to study the native role of CRT orthologs in other apicomplexam parasites.

Collectively, our findings revealed that the *Toxoplasma* chloroquine resistance transporter plays an

important role in regulating the size and morphology of the digestive vacuole in the parasites. This

regulation helps to maintain the integrity of the parasite's endolysosomal system, which provides routes

for the trafficking of invasion effectors. The co-localization between the digestive vacuole and endosome-

like compartment also led to a reduction of the transcript and protein levels of several endolysosomal

proteases. We also showed that blocking proteolysis within the swollen digestive vacuole reduced the size

and partially restored the morphology of the organelle. These findings suggest that TgCRT mediates the

transport of small solutes in order to regulate the size and morphology of the digestive vacuole. Our study

provides clear evidence that the integrity of the endolysosomal system in parasites is critical for the

virulence of parasites. Therefore, pharmaceutical modulation of the digestive vacuole could potentially

serve as a novel strategy to manage toxoplasmosis.

Materials and Methods

Ethical statement

This study was performed in compliance with the Public Health Service Policy on Humane Care and Use of Laboratory Animals and Association for the Assessment and Accreditation of Laboratory Animal Care guidelines. The animal protocol was approved by Clemson University's Institutional Animal Care and Use Committee (Animal Welfare Assurance A3737-01, protocol number AUP2016-012). All efforts were made to minimize discomfort. This method of euthanasia is consistent with the recommendations of the Panel on Euthanasia of the American Veterinary Medical Association.

Chemicals and reagents

Morpholine urea-leucyl-homophenyl-vinyl sulfone phenyl (LHVS) was kindly provided by the Bogyo lab at Stanford University. Other chemicals used in this work were analytical grade and were purchased from VWR unless otherwise indicated.

Parasite culture

Toxoplasma gondii parasites were cultured in human foreskin fibroblast (HFF) cells (ATCC, SCRC-1041) in Dulbecco's Modified Eagle Medium (DMEM) media supplemented with 10% cosmic calf serum at 37 °C with 5% CO_2. The parasites were harvested by membrane filtration as described previously [14].

Generation of transgenic parasites

To generate the TgCRT-deficient strain, ~1.5 kilobases (kb) of 5'- and 3'-UTR (untranslated region) of the *TgCRT* gene, respectively, were PCR-amplified and flanked at both ends of the bleomycin resistance cassette (BLE) to assemble a *TgCRT* deletion construct. The resulting plasmids were introduced into WT parasites by electroporation. The transfected parasites were selected with 50 µg/ml bleomycin twice, while in their extracellular stage as described previously [14]. The clones of the TgCRT-deficient parasites were generated by limiting dilution in a 96-well plate. The correct replacement of *TgCRT* with the *BLE* cassette was confirmed by PCR and quantitative PCR.

To complement the TgCRT in Δ*crt* parasites, we modified the plasmid, named pTub-TgCRT-mCherry-3xmyc (a gift from the van Dooren lab), which expresses a C-terminally mCherry-3xmyc epitope-tagged TgCRT under the *Toxoplasma* tubulin promoter. The plasmid was cut with HpaI and MfeI restriction

enzymes to remove the tubulin promoter and a partial sequence of TgCRT. The remaining DNA fragment served as the backbone for the subsequent Gibson assembly to incorporate the PCR amplified ~1 kb region upstream of the *Tgku80* gene, the ~1 kb fragment of the *Tgcrt* 5'-UTR region, and the removed partial *Tgcrt* coding sequence to produce the TgCRT complementation plasmid, pCRT-TgCRT-mCherry-3xmyc. The complemented TgCRT is driven by its cognate promoter to keep its expression level similar to the native TgCRT expression in WT parasites. The 1 kb region located ~6 kb upstream of the *Tgku80* gene was used to facilitate a single integration of the TgCRT complementation plasmid into this specific locus by single crossover homologous recombination. The TgCRT complementation construct was digested with SwaI restriction enzyme, gel-extracted, purified, and transfected into Δ*crt* parasites by the standard electroporation protocol.

To introduce NanoLuc® luciferase (nLuc) into parasites, we PCR-amplified and assembled the *Tgtubulin* promoter, the coding sequence of the nLuc luciferase, and an HXG selection marker into a nLuc expression construct by standard molecular cloning. The resulting plasmids were transfected into WT, Δ*crt*, and Δ*crtCRT*. The transfectants were selected with 25 µg/ml mycophenolic acid and 50 µg/ml xanthine. The stable populations were subjected to limiting dilution to generate individual clones of WT::*nLuc*, Δ*crt*::*nLuc*, and Δ*crtCRT*::*nLuc*. The correct clones were confirmed by their luciferase activities.

To generate the Δ*crt*Δ*cpb* mutant, the *TgCPB* gene was replaced with a pyrimethamine resistance cassette using the CRISPR-Cas9 genome editing system [36,37]. The pyrimethamine resistance cassette was PCR-amplified and flanked by 50 bp regions upstream and downstream of the start and stop codons of the *TgCPB* gene for homologous recombination. A 20 bp region located at the beginning of the coding region of the *TgCPB* gene was used to design guide RNA and replace the guide RNA targeting TgUPRT gene in the plasmid pSAG1-Cas9::UPRTsgRNA using Q5 site-directed mutagenesis kit (NEB). The Cas9-GFP and guide RNA constructs were co-transfected into Δ*crt* parasites with the corresponding repair PCR product. The guide RNA and Cas9 generated a gap within the *TgCPB* gene in the genome to facilitate double crossover homologous recombination. The correct gene replacement was confirmed by PCR.

To endogenously epitope-tag TgAMN, we also used CRISPR-Cas9 editing tools to modify the

corresponding gene in the genome. Guide RNA recognizing the 20 bp region close to the stop codon of

TgAMN was generated using the methods mentioned above.

The 50-bp homologous regions upstream and downstream of the stop codon of the *TgAMN* gene was

cloned at 5'- and 3'-ends of the DNA sequence containing the 3xHA epitope tag and the pyrimethamine

resistance cassette, respectively, by PCR. The plasmid encoding the guide RNA targeting TgAMN and

Cas9-GFP and the PCR product were co-trasfected into WT parasites. The stop codon of TgAMN was

replaced by the 3xHA epitope tag and pyrimethamine resistance cassette. The stable population was

generated after multiple rounds of pyrimethamine selection. The TgAMN-3xHA fusion protein was

confirmed by immunoblotting analysis.

TgSCP was endogenously tagged with a 3xmyc epitope tag via the single crossover recombination.

Approximately a 1 kb upstream region of the stop codon of the *TgSCP* gene was amplified by PCR and

fused in frame with a 3xmyc epitope tag at its 5'-end to assemble a TgSCP-3xmyc tagging construct. A

pyrimethamine resistance cassette was also encoded in the construct for drug selection. The resulting

plasmid was linearized by the restriction endonuclease NarI within the 1 kb region of the *TgSCP* gene prior

to transfection into WT parasites. The 3xmyc epitope tag was incorporated at the end of the *TgSCP* gene.

The correct tagging was confirmed by immunoblotting analysis.

Transfection of *Toxoplasma* parasites

T. gondii parasites were allowed to grow in HFF cells for 48 hrs at 37 °C with 5% CO_2. The freshly

egressed parasites were syringed, filter purified, and harvested in Cytomix buffer (25 mM HEPES, pH 7.6,

120 mM KCl, 10 mM K_2HPO4/ KH_2PO4, 5 mM $MgCl_2$. 0.15 mM $CaCl_2$, and 2 mM EGTA). Parasites were

pelleted at 1,000x *g* for 10 min, washed once in Cytomix buffer, and resuspended in Cytomix buffer at 2.5

x 10^7 parasites per ml. The 400 µL parasite suspension was mixed with 20 µg DNA and 2 mM ATP/5 mM

reduced glutathione for a total volume of 500 µl. The mixture was electroporated at 2 kV and 50 ohm

resistance using the BTX Gemini X2 (Harvard Apparatus). The transfected parasites were inoculated into

a T25 flask pre-seeded with a confluent monolayer of HFF cells. The corresponding drug selection was

applied at 24 hrs post-transfection.

Immunofluorescence assay

To perform immunofluorescence of intracellular parasites, freshly lysed parasites were used to infect confluent HFF cells pre-seeded in an 8-well chamber slide for 1 hr (pulse invasion) or 18-24 hrs (replication). The extracellular parasites were glued onto chamber slides by using 0.1% (w/v) poly-L-lysine before staining. Immunofluorescence staining was performed as described previously [11,14]. Images were viewed and digitally captured using a Leica® CCD camera equipped with a DMi8 inverted epifluorescence microscope and processed with Leica® LAS X software. Adobe Photoshop software was used to assemble the final images.

Excretory secretory antigens (ESAs) preparation

Freshly egressed parasites were syringed, filter purified, and resuspended at 5×10^8 parasites/ml in D3 medium (DMEM medium supplemented with 3% FBS). One hundred microliters of parasite suspension was transferred to a microfuge tube and incubated at 37 °C for 30 min to prepare constitutive ESAs. To make induced ESAs, the parasite resuspension was incubated in D1 medium supplemented with 1% ethanol for 2 min at 37ºC. The ESAs were separated from intact parasites by centrifugation at 1,000 x *g* for 10 min. The ESA fractions were transferred to a new microfuge tube, mixed with SDS-PAGE sample loading buffer, and boiled for 5 min for immunoblotting analysis.

SDS-PAGE electrophoresis and immunoblotting analysis

Parasite lysates and ESA fractions were prepared in 1x SDS-PAGE sample buffer and boiled for 5 min before resolving on SDS-PAGE gels and transferring to PVDF membranes by semi-dry protein transfer methods. Blots were blocked with 5% non-fat milk and incubated with primary antibodies diluted in 1% non-fat milk. Goat anti-mouse or anti-rabbit IgG antibodies conjugated with horseradish peroxidase were used as the secondary antibodies. Immunoblots were developed with SuperSignal™ WestPico chemiluminescent substrate (Thermo). The chemiluminescence signals were captured using the Azure® Imaging System. The densitometry and quantification of the protein bands were quantified by using LI-COR® Image Studio software.

Parasite invasion assay

Red-green invasion assay was used to measure the efficiency of parasite invasion. Freshly purified

parasites were syringed, filter purified, and resuspended at 5×10^7 parasites/ml in invasion medium (DMEM

supplemented with 3% FBS). Two hundred microliters of the parasite resuspension was inoculated into

each well of an 8-well chamber slide pre-seeded with HFF cells, and the parasites were allowed to invade

host cells for 30, 60, and 120 min before fixation with 4% formaldehyde for 20 min. Before membrane

permeabilization, slides were stained with mouse anti-TgSAG1 monoclonal antibody (1:1,000) for 1 hr to

label attached parasites. After treatment with 0.1% Triton X-100 for 10 min, the parasites were stained with

rabbit polyclonal anti-TgMIC5 antibody (1:1,000) for 1 hr to stain both invaded and attached parasites.

Subsequently, slides were stained with goat anti-mouse IgG conjugated with Alexa 594 (red) (Invitrogen,

1:1,000) and goat anti-rabbit IgG conjugated with Alexa 488 (green) (Invitrogen, 1:1,000) along with DAPI

for nucleus staining. After staining, slides were mounted with anti-fade Mowiol solution and observed by

immunofluorescence microscopy. Extracellular parasites only showed red fluorescence, whereas

intracellular parasites exhibited both red and green fluorescence. Six fields of view from individual invasion

experiments were captured by the Leica® DMi8 inverted epifluorescence microscope and enumerated by

ImageJ software. The attachment efficiency of each strain was measured by dividing the total number of

parasites labeled in red by the total number of host nuclei, and normalized against that of WT parasites.

The invasion efficiency of each strain was quantified using the following equation (the sum of parasites

labeled in green − the sum of parasites labeled in red)∕the total number of host nuclei, and normalized

against that of WT parasites.

Immunofluorescence-based replication assay

Freshly egressed parasites were filter-purified and inoculated into individual wells of an 8-well chamber

slide pre-seeded with HFF cells at approximately 1×10^5 cells per well. Non-invaded parasites were

washed off at 4 hrs post-infection. Invaded parasites were allowed to infect host cells for an additional 24

and 36 hrs before fixation. The infected host cells were stained with monoclonal anti-TgGRA7 (1:2,000)

antibody and DAPI to help distinguish individual parasitophorous vacuoles (PVs) and the nuclei of

parasites, respectively. Slides were subjected to standard immunofluorescence microscopy for imaging.

One hundred parasitophorous vacuoles were enumerated for each strain and plotted as the distribution of

different sized PVs. In addition, replication was also expressed as the average number of parasites per

PV.

Luminescence-based growth assay

Parasites expressing NanoLuc luciferase were inoculated into a white 96-well tissue culture plate with

a flat, solid bottom (Greiner Bio-One) pre-seeded with confluent HFF cells at 1.5 x 10^3 cells/well. Each

strain was inoculated into 4 individual wells to monitor the fold-change of luciferase activities over a time

course, which reflects intracellular growth of the parasites. After 4 hrs post-infection, the individual wells

were aspirated to remove non-invaded parasites. The first well was treated with 100 µl of lysis buffer

containing NanoLuc luciferase substrate and incubated for 10 min, and a luminescence reading was taken

by using the BioTek® multimode H1 hybrid plate reader. The rest of the 3 wells were replenished with fresh

D10 medium without phenol red for an additional 24, 48, and 72 hrs. The subsequent luminescence

readings were all determined using the methods mentioned above. The luminescence readings at different

time intervals were normalized against the reading at 4 hrs post-infection to calculate the fold-change of

parasite growth.

Egress assay

Lactate dehydrogenase release assay was used to measure the egress efficiency of parasites. Freshly

lysed parasites were filter-purified and resuspended in D10 medium at 5 x 10^5 parasites/ml. One hundred

microliters of the parasite resuspension was inoculated into each well of a 96-well plate pre-seeded with

HFF cells. The parasites were allowed to replicate for 18-24 hrs, washed, and incubated with 50 µl of

Ringer's buffer (10 mM HEPES, pH 7.2, 3 mM HaH$_2$PO$_4$, 1 mM MgCl$_2$, 2 mM CaCl$_2$, 3 mM KCl, 115 mM

NaCl, 10 mM glucose, and 1% FBS) for 20 min. Subsequently, an equal volume of 1 mM Zaprinast

dissolved in Ringer's buffer was added into the wells and incubated for 5 min at 37ºC and 5% CO$_2$.

Uninfected wells were treated with 50 µl of Ringer's buffer containing 1% Triton X-100 or plain Ringer's

buffer, serving as positive and negative controls, respectively. The released lactate dehydrogenase was

centrifuged at 1,000 *x g* for 5 min twice to pellet insoluble cell debris. Fifty microliters of supernatant was

subjected to the standard lactate dehydrogenase release assay as described previously [38]. The egress

efficiency of each strain was calculated using the following equation, (LDH activity derived from individual

parasites − LDH activity of negative control)∕(LDH activity of positive control − LDH activity of negative

control), and normalized against that of WT parasites.

Size measurement of the VAC

The size of the VAC was quantified based on the staining of TgCPL, a VAC luminal protease. Freshly

purified parasites were inoculated into pre-seeded HFF chamber slides, allowed to invade host cells for 30

min prior to fixation, stained with polyclonal rabbit anti-TgCPL antibody (1:100), and subjected to standard

immunofluorescence microscopy. The distance of the widest diagonal of the TgCPL staining was

determined as the diameter of the VAC and was measured using the Leica® LAS X software.

Measurements of 50 individual parasites were performed for each replicate.

Transcriptome sequencing and quantitative PCR (qPCR) assay

Total RNA was extracted from freshly lysed parasites using the Zymo® Direct-zol^{TM} RNA MiniPrep Plus

kit, and converted to sequencing read libraries by using the TruSeq Stranded mRNA sequencing kit

(Illumina). The prepared libraries were subjected to 2 x 125 bp paired-end Illumina® HiSeq2500 sequencing.

Each sample was sequenced to a depth of at least 20 million reads. Differential expression profiling was

performed by the Clemson University Genomics Computational Lab.

Approximately 500 ng of total RNA was used to measure the steady levels of transcripts for individual

genes by using the Luna® Universal One-Step RT-PCR kit (NEB). The qPCR assay was performed using

the BioRad CFX96 Touch^{TM} Real-Time PCR detection system. The quantification cycle (Cq) values for

individual genes were used for double delta Cq analysis to calculate their relative abundances to that of

WT parasites using the Bio-Rad® CFX Maestro^{TM} software. TgActin was used as the housekeeping gene

for normalization.

Mouse studies

Six- to eight-week-old, outbred CD-1 mice were infected by subcutaneous or intravenous injection with

100 WT or mutant parasites diluted in PBS. The infected mice were monitored for symptoms daily for a

total of 30 days. Mice that appeared moribund were humanely euthanized via CO_2 overdose, in compliance

with IACUC's approved protocol. The seroconversion of the surviving mice was tested by enzyme-linked

immunosorbent assay (ELISA). The surviving mice were allowed to rest for 10 days, prior to subcutaneous

injection with a challenge dose of 1000 WT parasites, and were monitored daily for survival for 30 days.

Cloning and Expression of the recombinant pro-form of TgASP1

A ~1.5-kb DNA fragment encoding the recombinant pro-form of TgASP1 (rproTgASP1) was PCR-

amplified from a *T. gondii* RH cDNA library using Q5® High-Fidelity DNA polymerase (New England Biolabs)

and cloned into a pET22b vector. A 6xHis epitope tag was fused in-frame at the N-terminal end of TgASP1.

The resulting pET22b-6xHis-TgASP1 expression construct was transformed into the *E. coli* protein

expression strain, ER2566 (NEB). Recombinant TgASP1 was induced with 1 mM isopropyl 1-thio-β-D-

galactopyranoside (IPTG) at 37ºC for 4 hrs. The expressed recombinant TgASP1 protein was purified

using a standard denature protein purification procedure of recombinant His-tagged proteins as described

previously [19]. The purified denatured TgASP1 protein was refolded in phosphate buffered saline (PBS)

buffer overnight by dialysis. Mis-aggregated TgASP1 was removed by centrifugation at 20,000 *xg* for 10

min. The soluble fraction was concentrated using an Amicon® Ultra 30-kDa cutoff concentrator (Millipore)

and stored at -20ºC for future use.

Antibody Preparation

710   The purified pro-forms of TgASP1 were shipped to Cocalico Biologicals, Inc. to generate polyclonal

711   antibodies. Two New Zealand white rabbits were initially injected with 200 µg of purified recombinant

712   protein in Freund's complete adjuvant and boosted 3 times at Day 14, 21, and 56 with 50 µg of recombinant

713   protein in Freund's incomplete adjuvant. The specificity of the polyclonal antibodies were assessed by

714   comparing their reactivity in WT and Δ*asp1* parasites by immunoblotting and immunofluorescence assay.

Statistics

716   Statistical analysis was performed using Prism software (GraphPad). The methods used in different

717   assays were indicated in the figure legends.

Data Availability Statement: The raw data of transcriptomic sequencing of WT and Δ*crt* strains have been deposited in the NIH Gene Expression Omnibus (GEO) database. The accession number is GSE116539. https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE116539

Funding: This work was supported by Knights Templar Eye Foundation Pediatric Ophthalmology Career-Starter Research Grant (to Z.D.), a pilot grant of an NIH COBRE grant P20GM109094 (to Z.D.), and the Clemson Startup fund (to Z.D.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Acknowledgments

We thank our colleagues Drs. Michael Blackman, Peter Bradley, Vern Carruthers, Silvia Moreno, and David Sibley for kindly providing key reagents for this study. We also want to thank Drs. Rooksana Noorai and Vijay Shankar at Clemson University Genomics Computational Lab for providing technical assistance and expertise in performing transcriptomic sequencing and differential gene expression analysis and depositing the raw data files of transcriptomic sequencing into the NIH GEO database. We acknowledge Dr. Lesly Temesvari for critically reading this manuscript before submission.

Author Contributions

Conceptualization: Zhicheng Dou.

Formal analysis: L. Brock Thornton, Paige Teehan, Katherine Floyd, Christian Cochrane, Amy Bergmann, Zhicheng Dou.

Funding acquisition: Zhicheng Dou.

Investigation: L. Brock Thornton, Paige Teehan, Katherine Floyd, Christian Cochrane, Amy Bergmann, Zhicheng Dou.

Methodology: L. Brock Thornton, Paige Teehan, Zhicheng Dou.

746   **Project administration:** Zhicheng Dou.

747   **Resources:** Paige Teehan, Amy Bergmann, Zhicheng Dou.

748   **Supervision:** Zhicheng Dou.

749   **Writing – original draft:** Zhicheng Dou.

750   **Writing – review & editing:** L. Brock Thornton, Zhicheng Dou.

Figure legends:

Figure 1. The TgCRT-deficient parasites displayed a swollen digestive vacuole and a disrupted endolysosomal system. A) Schematic illustration of the strategies for the *Tgcrt* deletion and complementation in *Toxoplasma* parasites. The plasmid carrying a bleomycin resistance cassette (*BLE*) flanked by the *TgCRT* targeting sequences was transfected into WT parasites for double crossover replacement of *TgCRT* to produce the Δ*crt* strain. The TgCRT complementation plasmid, containing the coding sequence of TgCRT fused with mCherry and 3xmyc epitope tags at its C-terminus, was introduced into the Δ*crt* strain to produce the Δ*crtCRT* complementation strain. **B)** The primers indicated in panel A were used to verify the correct replacement of *TgCRT* with *BLE* by PCR. **C)** The complemented *TgCRT* gene was transfected into Δ*crt* parasites and was verified by PCR. Since we complemented Δ*crt* with the coding sequence of *TgCRT*, the PCR product was a 0.2 kb fragment in the Δ*crtCRT* strain, whereas the WT strain showed a 1.5 kb product since its *TgCRT* gene contains the introns. **D)** Transcript levels of *TgCRT* in the WT, Δ*crt*, and Δ*crtCRT* strains were evaluated by quantitative PCR. Primers were designed to anneal to the exons of *TgCRT* and are indicated in Panel A. The *Tgactin* gene was included as a control for normalization. The quantification of transcripts was performed in at least three biological replicates and analyzed using unpaired Student's *t*-test. **E)** Extracellular Δ*crt* parasites showed an enlarged concave subcellular structure, indicated by the arrow, under the differential interference contrast (DIC) microscopy. Scale bar = 5 μm. **F)** The swollen subcellular structure indicated by the arrowhead was also observed in Δ*crt* parasites during pulse invasion and co-localized with a major luminal peptidase of the VAC, cathepsin L-like protease (TgCPL). The TgCPL staining was used to assess the morphology of the VAC. The parasites were allowed to invade host cells for 30 min before TgCPL antibody staining. The distance of the widest diagonal of the TgCPL staining was determined to be the diameter of the VAC and was measured using the Leica® LAS X software. The measurements were conducted in three biological replicates. Measurements from 50 individual parasites from one representative assay were shown. The mean VAC size ± SD was calculated for three independent measurements and is listed on the figure. Statistical significance was determined using unpaired Student's *t*-test. Scale bar = 2 μm. **G)** Parasites were co-stained with anti-TgCPL (the marker of the VAC) and anti-proM2AP or anti-TgVP1 (both are markers of

the endosome-like compartment, ELC). During pulse invasion, the VAC and ELC staining were juxtaposed in the WT and Δ*crtCRT* strains, but aberrantly co-localized in the Δ*crt* strain. During replication, the VAC in WT and Δ*crtCRT* parasites became fragmented. However, the abnormal co-localization of the VAC and ELC significantly decreased the extent of VAC fragmentation in the Δ*crt* mutant. A major TgCPL punctum existed in the replicated Δ*crt* parasites, not in WT and Δ*crtCRT* strains, and co-localized with proTgM2AP and TgVP1 (indicated by arrows in the insets). The scale bars in the images of parasites at the stages of pulse invasion and replication are 2 μm and 5 μm, respectively. ****, $p<0.0001$; n.s., not significant.

Figure 2. Parasite invasion and acute virulence were reduced in the Δ*crt* parasites. A) WT, Δ*crt*, and Δ*crtCRT* parasites were allowed to invade host cells for 30, 60, and 120 min prior to fixation and antibody staining. Parasites and host cells within six fields of view were counted for each strain. At least three independent invasion assays were conducted for statistical analysis. The Δ*crt* parasites showed a ~50% reduction in invasion at 30 min post-infection. The extent of the invasion defect in the Δ*crt* mutant was gradually minimized overtime. At 60 min post-infection, there was approximately a 25% reduction in invasion in the Δ*crt* mutant, while there were no significant differences among the three strains at 120 min post-infection. The assay was performed at least in triplicate. Statistical significance was determined using unpaired Student's *t*-test. **B)** We infected confluent HFFs with WT, Δ*crt*, and Δ*crtCRT* parasites for 28 and 40 hrs before fixation and staining. Infected cells were stained with DAPI and anti-TgGRA7 antibodies in order to recognize individual parasites and parasitophorous vacuoles, respectively. One hundred parasitophorous vacuoles (PVs) were enumerated for the number of parasites they contained and the distribution of different sized PVs was plotted. The average number of parasites per PV was also calculated and listed above the plots. In addition, the growth rates of these strains were determined at 24, 48, and 72 hours by using a luminescence-based assay. Parasites were inoculated into a 96-well plate pre-seeded with HFFs prior to lysis and quantification of luminescence activities at pre-determined time intervals. At least three biological replicates were performed for the replication assay. Unpaired Student's *t*-test was performed to calculate the statistical significance of parasite growth between strains. **C)** To measure the egress of the parasites, 5×10^4 parasites were used to infect confluent HFFs in a 96-well plate for 24 hrs.

Replicated parasites were treated with 500 µM Zaprinast for 5 min at 37ºC and 5% CO_2 to induce egress. Disruption of host cell membranes due to parasite egress released lactate dehydrogenase into the medium, which was quantified and plotted. Unpaired Student's *t*-test was used to calculate the statistical significance. There were no differences observed in parasite egress between WT and Δ*crt* parasites. **D)** The acute virulence of TgCRT-deficient parasites was evaluated in a murine model via subcutaneous and intravenous infections. One hundred parasites from each strain were used to infect outbred CD-1 mice (n=10 mice for each strain). The mortality of the mice was monitored for 30 days. Seroconversion of the surviving mice was evaluated by ELISA to confirm successful infection. Additionally, the surviving mice were allowed to rest for 10 days before subsequent challenge with 1,000 WT parasites by subcutaneous inoculation. The Δ*crt* mutant exhibited reduced acute virulence compared to the WT and Δ*crtCRT* strains and conferred immunological protection in the surviving mice. Data were recorded and are presented using the Kaplan-Meier plot. Statistical analysis was performed using the Log-rank (Mantel-Cox) test. For all statistical significance calculations, *, $p<0.05$; **, $p<0.01$; ****, $p<0.0001$; n.s, not significant.

Figure 3. The deletion of *TgCRT* altered microneme secretion, without affecting the microneme steady abundance, intracellular trafficking, and intramembrane cleavage on the parasite surface. A) The steady levels of micronemal proteins were not altered in the Δ*crt* parasites. Freshly lysed parasites were filter-purified, lysed, and subjected to SDS-PAGE electrophoresis and immunoblotting. The blots were probed with anti-TgMIC2, TgM2AP, and TgMIC5 antibodies, along with anti-TgActin as the loading control. Individual micronemal proteins were normalized against the corresponding TgActin to quantify their steady state expression. **B)** Δ*crt* parasites secreted less micronemal proteins than WT and Δ*crtCRT* parasites, and altered the micronemal secretion patterns. Freshly filter-purified parasites were incubated in medium at 37ºC for 30 min to make constitutive ESAs or were treated with 1% (v/v) ethanol in medium to produce induced ESAs. The ESA fractions were separated and probed with anti-TgMIC2, TgM2AP, and TgMIC5 antibodies for quantification of the secreted forms of these micronemal proteins. The ESA fractions were also probed with anti-TgGRA7 antibody, a dense granule protein, as the loading control. Statistical significance was determined using unpaired Student's *t*-test. **C)** Following pulse invasion and replication,

the parasites were stained with anti-TgMIC2 and anti-TgM2AP antibodies to examine their intracellular trafficking. No defects were detected in their intracellular trafficking. **D)** The parasites at the stages of pulse invasion and replication were also stained with anti-TgROP7 to examine the morphology of the rhoptry and intracellular trafficking of TgROP7. Similar morphology and trafficking patterns of the rhoptry were maintained among WT, Δ*crt*, and Δ*crtCRT* strains. The scale bars in the images of parasites during pulse invasion and at extracellular stage are 2 μm, and the scale bar in the images of replicated parasites is 5 μm. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$; n.s., not significant.

Figure 4. The transcript and protein levels of a subtilisin-related protease, TgSUB1, were reduced in Δ*crt* parasites. A) The abundance of secreted TgSUB1 in the constitutive and induced ESAs was measured by immunoblotting. There was no detectable TgSUB1 in the constitutive and induced ESAs secreted by Δ*crt* parasites via immunoblotting. **B)** The abundance of plasma membrane-anchored TgSUB1 was measured by staining non-permeabilized parasites with antibodies recognizing TgSUB1. The Δ*crt* mutant displayed a loss of TgSUB1 on its plasma membrane. **C)** TgSUB1 is a micronemal protein. TgSUB1 and TgMIC5 were both found to localize in the microneme in WT and Δ*crtCRT* parasites. However, TgSUB1 signal was not detected in the Δ*crt* parasites. **D)** The cell lysates of the Δ*crt* parasites showed that the steady level of TgSUB1 was reduced by ~90% in the Δ*crt* strain. **E)** The transcript levels of TgSUB1 were measured by quantitative PCR and were also decreased by approximately 90% in the Δ*crt* mutant. All assays listed in this figure were repeated at least in triplicate. Statistical significance was calculated using unpaired Student's *t*-test: ***, $p<0.001$; ****, $p<0.0001$; n.s., not significant.

Figure 5. The transcript and protein abundances of several VAC-residing proteases were decreased in Δ*crt*. A) RNA-Seq was performed in WT and Δ*crt* parasites. Each sample was sequenced in duplicate for statistical comparison. A volcano plot was used to display a summary of the genes that had altered transcription greater than 1.5-fold with statistical significance less than 0.05 in the Δ*crt* mutant relative to the WT strain. Forty-six and fifty-six genes labeled in blue and red dots became up- and down-regulated in the Δ*crt* mutant, respectively. The blue and red dash lines represented the border line of a 1.5-fold

transcriptional change of genes, and the genes above the black dash line showed *p* values of a statistical significance below 0.05. **B)** qPCR was used to validate 4 down-regulated proteases that were identified by RNA-Seq analysis, along with two known VAC proteases, TgCPL and TgCPB. *TgAMN*, *TgSCP*, *TgASP1*, and *TgCPB* displayed down-regulated transcription in Δ*crt* parasites compared to WT strain. **C)** The steady protein abundances of TgCPL and TgCPB were quantified in the lysates of parasites by immunoblotting. The protein levels of TgCPL and TgCPB in Δ*crt* mutant were reduced by ~25% and 60%, respectively, compared to WT parasites. **D)** TgSCP and TgAMN were endogenously tagged with 3xmyc and 3xHA, respectively, at their C-termini. The expression of the epitope-tagged proteins was confirmed by immunoblotting. The parasites were co-stained with antibodies recognizing their respective epitope tags as well as the VAC and ELC markers. Both TgSCP and TgAMN were localized in the VAC and ELC by immunofluorescence. Statistical significance was performed using unpaired Student's *t*-test. *, *p*<0.05; **, *p*<0.01; n.s., not significant.

Figure 6. Suppression of proteolysis in the VAC reduced VAC size and partially restored integrity of parasite's endolysosomal system, which recovered TgSUB1 expression, micronemal protein trimming in ESAs, and parasite invasion. A) The Δ*crt* parasites were incubated with 1 μM LHVS, an irreversible inhibitor of TgCPL, or with DMSO vehicle control, for 48 hrs, followed by a pulse invasion. Parasites were stained with anti-TgCPL antibodies to determine the size of the VAC. The swollen VAC phenotype was significantly reduced in the LHVS-treated parasites. Scale bar = 2μm. **B)** TgCPB, a VAC-residing endo- and exo-peptidase was genetically deleted in the Δ*crt* strain. A schematic illustration for the creation of the Δ*crt*Δ*cpb* mutant is shown. A PCR product carrying a pyrimethamine resistance cassette (DHFR) flanked by 50 bps of the 5'- and 3'-untranslated regions of *TgCPB* was transfected into WT parasites for double crossover replacement of *TgCPB*. Primers indicated in panel B were used to verify the replacement of *TgCPB* with *DHFR* via PCR and agarose gel electrophoresis. The ablation of *TgCPB* in Δ*crt*Δ*cpb* parasites was also confirmed by immunoblotting. **C)** The sizes of the VAC in WT, Δ*crt*, and Δ*crt*Δ*cpb* parasites were determined by the methods mentioned above. The Δ*crt*Δ*cpb* parasites displayed a partial reduction in the size of the VAC compared to WT parasites. **D)** Following pulse invasion, the

parasites were fixed, stained with anti-TgCPL (VAC marker) and proTgM2AP (ELC marker), and enumerated under immunofluorescence microscope. The parasites showing partial overlapping between the VAC and ELC in WT, Δ*crt*, Δ*crtCRT*, and Δ*crt*Δ*cpb* strains were quantified. At least 100 parasites were quantified for each replicate in a total of three replicates. The Δ*crt*Δ*cpb* parasites showed a significantly lower percentage of parasites having arrested co-localization between the VAC and ELC compared to the Δ*crt* mutant. **E)** The lysates of WT, Δ*crt*, and Δ*crt*Δ*cpb* parasites were probed with antibodies recognizing TgSUB1.The steady expression of TgSUB1 was restored in Δ*crt*Δ*cpb* parasites. **F)** The constitutive and induced ESAs of WT, Δ*crt*, and Δ*crt*Δ*cpb* strains were made and probed with the antibodies indicated in the figure. The secretion and trimming of micronemal proteins were also recovered in the Δ*crt*Δ*cpb* mutant. **G)** The invasion efficiencies of WT, Δ*crt*, and Δ*crt*Δ*cpb* strains were determined using the procedures mentioned above. The Δ*crt*Δ*cpb* parasites showed increased invasion efficiency compared to the Δ*crt* strain, albeit still a lower efficiency than that of WT parasites. Scale bar = 2 μm. Statistical significance was calculated using unpaired Student's *t*-test. *, $p<0.05$; ; **, $p<0.01$; ***, $p<0.001$; n.s., not significant.

Figure 7. A model for the regulation of the endolysosomal system in *Toxoplasma* parasites. *Toxoplasma* parasite contains a separate endosome-like compartment from the digestive vacuole within its endolysosomal system. When the parasite lacks TgCRT, the VAC becomes swollen and cannot separate from its adjacent ELC. This aberrant co-localization leads to reduced transcript and protein abundances of several proteases residing within the endolysosomal system, including TgSUB1. These changes alter the secretion of micronemal proteins, thereby resulting in invasion defects in the TgCRT-null mutant. ELC, endosome-like compartment; ER, endoplasmic reticulum; Go, Golgi apparatus; Mi, microneme; N, nucleus; Rh, rhoptry; TgCRT, *Toxoplasma* chloroquine resistance transporter ortholog; TgSUB1, *Toxoplasma* subtilisin 1; VAC, vacuolar compartment.

Table 1. Strains used in this study.

Name	Genetic background	Comments
WT	RH*Δku80Δhxg*	Requested from the Carruthers Lab, not generated in this study
Δ*crt*	RH*Δku80ΔhxgΔcrt*	*TgCRT* was deleted by double crossover homologous recombination
Δ*crtCRT*	RH*Δku80ΔhxgΔcrt*::*TgCRT-mCherry-3xmyc*	Ectopic expression of a C-termially epitope-tagged TgCRT in Δ*crt* for complementation
Δ*crt*Δ*cpb*	RH*Δku80ΔhxgΔcrtΔcpb*	The entire *TgCPB* gene was ablated by CRISPR-Cas9 based genome editing technique
WT::*nLuc*	RH*Δku80Δhxg*::*nLuc*	Expressed NanoLuc luciferase in WT parasites
Δ*crt*::*nLuc*	RH*Δku80ΔhxgΔcrt*::*nLuc*	Expressed NanoLuc luciferase in Δ*crt* parasites
Δ*crtCRT::nLuc*	RH*Δku80ΔhxgΔcrt*::*TgCRT-mCherry-3xmyc*::*nLuc*	Expressed NanoLuc luciferase in Δ*crtCRT* parasites
TgAMN-3xHA	RH*Δku80Δhxg*TgAMN-3xHA	*TgAMN* gene was endogenously tagged with a 3xHA epitope at its 3'-end by CRISPR-Cas9 based genome editing technique
TgSCP-3xmyc	RH*Δku80Δhxg*TgSCP-3xmyc	*TgSCP* gene was endogenously tagged with a 3xmyc epitope at its 3'-end by single crossover recombination

Supplemental Figure Legends

S1 Figure. No defects in intramembrane proteolytic cleavage of micronemal protein 2 (TgMIC2) were observed in the Δ*crt* mutant. Purified, extracellular parasites that had not been permeabilized were stained with anti-TgMIC2 and anti-TgSAG1 antibodies in order to measure the retention of TgMIC2 on the parasite surface. During secretion, the TgMIC2 protein is cleaved by intramembrane rhomboid proteases, such as TgROM4. The abundance of TgMIC2 on the surface of Δ*crt* parasites was similar to that of the WT and Δ*crtCRT* strains, indicating that there is comparable intramembrane cleavage of TgMIC2 among the parasites with or without TgCRT. Scale bar = 2 μm.

S2 Figure. Schematic of the endogenous epitope-tagging of the *Toxoplasma* putative aminopeptidase N (TgAMN, TGGT1_221310) and the putative Pro-Xaa serine carboxypeptidase (TgSCP, TGGT1_254010). A) The plasmid encoding Cas9 and sgRNA targeting TgAMN were co-transfected into WT parasites with the PCR product carrying a 3xHA epitope tag and a pyrimethamine resistance cassette (DHFR) flanked by 50 bp regions upstream and downstream of the stop codon of TgAMN. The 3xHA tag and the drug resistance cassette were incorporated at the C-terminus of the

Toxoplasma putative aminopeptidase N via double crossover homologous recombination mediated by the CRISPR-Cas9 genome editing tool. **B)** The putative Pro-Xaa serine carboxypeptidase was endogenously tagged with a 3xmyc epitope tag at its C-terminus by single crossover homologous recombination. A 1 kb region upstream of the stop codon of TgSCP was amplified and fused at the 5'-end of the 3xmyc tag to produce the TgSCP-3xmyc tagged plasmid. The 1 kb TgSCP-coding region was cleaved by an endonuclease in the middle prior to transfection to facilitate its integration.

S3 Figure. The subcellular concave structure in the Δ*crt*Δ*cpb* mutant was reduced in size relative to the Δ*crt* strain during the extracellular stage. WT, Δ*crt*, Δ*crtCRT*, and Δ*crt*Δ*cpb* parasites were purified and attached on the surface of the slide for differential interference contrast (DIC) microscopy imaging. The Δ*crt*Δ*cpb* parasites still showed an enlarged concave subcellular structure (indicated by the arrow), however their sizes were significantly smaller than those in the Δ*crt* mutant. Scale bar = 5 μm.

S1 Table. Differential gene expression analysis between WT and *Δcrt* strains. The genes whose fold changes were >1.5 and *p*-values were <0.05 shown in RNA-Seq analysis were listed.

S2 Table. Primers used in this study.

References

1.   Laliberté J, Carruthers VB. Host cell manipulation by the human pathogen *Toxoplasma gondii*. Cell Mol Life Sci. 2008;65: 1900–1915. doi:10.1007/s00018-008-7556-x

2.   Carruthers VB, Tomley FM. Microneme proteins in apicomplexans. Subcell Biochem. 2008;47: 33–45.

3.   Lagal V, Binder EM, Huynh M-H, Kafsack BFC, Harris PK, Diez R, et al. *Toxoplasma gondii* protease TgSUB1 is required for cell surface processing of micronemal adhesive complexes and efficient adhesion of tachyzoites. 2010;12: 1792–1808. doi:10.1111/j.1462-5822.2010.01509.x

4.   Parussini F, Coppens I, Shah PP, Diamond SL, Carruthers VB. Cathepsin L occupies a vacuolar compartment and is a protein maturase within the endo/exocytic system of *Toxoplasma gondii*. Mol Microbiol. 2010;76: 1340–1357. doi:10.1111/j.1365-2958.2010.07181.x

5.   Dogga SK, Mukherjee B, Jacot D, Kockmann T, Molino L, Hammoudi P-M, et al. A druggable secretory protein maturase of Toxoplasma essential for invasion and egress. Elife. 2017;6: 223. doi:10.7554/eLife.27480

6.   Shen B, Buguliskis JS, Lee TD, Sibley LD. Functional analysis of rhomboid proteases during Toxoplasma invasion. MBio. 2014;5: e01795–14. doi:10.1128/mBio.01795-14

7.   Buguliskis JS, Brossier F, Shuman J, Sibley LD. Rhomboid 4 (ROM4) affects the processing of surface adhesins and facilitates host cell invasion by *Toxoplasma gondii.* PLoS Pathog. 2010;6: e1000858. doi:10.1371/journal.ppat.1000858

8.   Santos JM, Ferguson DJP, Blackman MJ, Soldati-Favre D. Intramembrane cleavage of AMA1 triggers *Toxoplasma* to switch from an invasive to a replicative mode. Science. 2011;331: 473–477. doi:10.1126/science.1199284

9.   Parussini F, Tang Q, Moin SM, Mital J, Urban S, Ward GE. Intramembrane proteolysis of Toxoplasma apical membrane antigen 1 facilitates host-cell invasion but is dispensable for replication. Proc Natl Acad Sci USA. National Acad Sciences; 2012;109: 7463–7468. doi:10.1073/pnas.1114661109

10.   Dou Z, Carruthers VB. Cathepsin proteases in *Toxoplasma gondii*. Adv Exp Med Biol. Boston, MA: Springer US; 2011;712: 49–61. doi:10.1007/978-1-4419-8414-2_4

11.   Dou Z, McGovern OL, Di Cristina M, Carruthers VB. *Toxoplasma gondii* ingests and digests host cytosolic proteins. MBio. 2014;5: e01188–14. doi:10.1128/mBio.01188-14

12.   Miller SA, Binder EM, Blackman MJ, Carruthers VB, Kim K. A conserved subtilisin-like protein TgSUB1 in microneme organelles of *Toxoplasma gondii*. J Biol Chem. 2001;276: 45341–45348. doi:10.1074/jbc.M106665200

13.   Di Cristina M, Dou Z, Lunghi M, Kannan G, Huynh M-H, McGovern OL, et al. *Toxoplasma* depends on lysosomal consumption of autophagosomes for persistent infection. Nat Microbiol. 2017;2: 17096. doi:10.1038/nmicrobiol.2017.96

14.   Dou Z, Coppens I, Carruthers VB. Non-canonical maturation of two papain-family proteases in *Toxoplasma gondii*. J Biol Chem. American Society for Biochemistry and Molecular Biology; 2013;288: 3523–3534. doi:10.1074/jbc.M112.443697

15. Warring SD, Dou Z, Carruthers VB, McFadden GI, van Dooren GG. Characterization of the chloroquine resistance transporter homologue in *Toxoplasma gondii*. Euk Cell. 2014;13: 1360–1370. doi:10.1128/EC.00027-14

16. Liu J, Pace D, Dou Z, King TP, Guidot D, Li Z-H, et al. A vacuolar-H^+-pyrophosphatase (TgVP1) is required for microneme secretion, host cell invasion, and extracellular survival of *Toxoplasma gondii*. Mol Microbiol. 2014;93: 698–712. doi:10.1111/mmi.12685

17. Juge N, Moriyama S, Miyaji T, Kawakami M, Iwai H, Fukui T, et al. *Plasmodium falciparum* chloroquine resistance transporter is a H^+-coupled polyspecific nutrient and drug exporter. Proc Natl Acad Sci USA. 2015;112: 3356–3361. doi:10.1073/pnas.1417102112

18. Huynh M-H, Carruthers VB. Tagging of endogenous genes in a *Toxoplasma gondii* strain lacking Ku80. Euk Cell. American Society for Microbiology; 2009;8: 530–539. doi:10.1128/EC.00358-08

19. Larson ET, Parussini F, Huynh M-H, Giebel JD, Kelley AM, Zhang L, et al. *Toxoplasma gondii* cathepsin L is the primary target of the invasion-inhibitory compound morpholinurea-leucyl-homophenyl-vinyl sulfone phenyl. J Biol Chem. 2009;284: 26839–26850. doi:10.1074/jbc.M109.003780

20. McGovern OL, Rivera-Cuevas Y, Kannan G, Narwold AJ, Carruthers VB. Intersection of endocytic and exocytic systems in *Toxoplasma gondii*. Traffic. Wiley/Blackwell (10.1111); 2018;19: 336–353. doi:10.1111/tra.12556

21. Blader IJ, Coleman BI, Chen C-T, Gubbels M-J. Lytic Cycle of Toxoplasma gondii: 15 Years Later. Annu Rev Microbiol. 2015;69: 463–485. doi:10.1146/annurev-micro-091014-104100

22. Dubremetz J-F. Rhoptries are major players in *Toxoplasma gondii* invasion and host cell interaction. Cell Microbiol. Wiley/Blackwell (10.1111); 2007;9: 841–848. doi:10.1111/j.1462-5822.2007.00909.x

23. Hager KM, Carruthers VB. MARveling at parasite invasion. Trends Parasitol. 2008;24: 51–54. doi:10.1016/j.pt.2007.10.008

24. Sloves P-J, Delhaye S, Mouveaux T, Werkmeister E, Slomianny C, Hovasse A, et al. Toxoplasma Sortilin-like Receptor Regulates Protein Transport and Is Essential for Apical Secretory Organelle Biogenesis and Host Infection. Cell Host Microbe. 2012;11: 515–527. doi:10.1016/j.chom.2012.03.006

25. Saouros S, Dou Z, Marchant J, Carruthers VB, Matthews S. Microneme protein 5 regulates the activity of *Toxoplasma* subtilisin 1 by mimicking a subtilisin prodomain. J Biol Chem. American Society for Biochemistry and Molecular Biology; 2012;287: 36029–36040. doi:10.1074/jbc.M112.389825

26. Binder EM, Lagal V, Kim K. The Prodomain of T*oxoplasma gondii* GPI-Anchored Subtilase TgSUB1 Mediates its Targeting to Micronemes. Traffic. 2008;9: 1485–1496. doi:10.1111/j.1600-0854.2008.00774.x

27. Settembre C, Di Malta C, Polito VA, Garcia Arencibia M, Vetrini F, Erdin S, et al. TFEB links autophagy to lysosomal biogenesis. Science. 2011;332: 1429–1433. doi:10.1126/science.1204592

28. Fu M, Blackshear PJ. RNA-binding proteins in immune regulation: a focus on CCCH zinc finger proteins. Nat Rev Immunol. 2017;17: 130–143. doi:10.1038/nri.2016.129

29. Carballo E, Lai WS, Blackshear PJ. Feedback Inhibition of Macrophage Tumor Necrosis Factor-α Production by Tristetraprolin. Science. American Association for the Advancement of Science; 1998;281: 1001–1005. doi:10.1126/science.281.5379.1001

30. Miranda K, Pace DA, Cintron R, Rodrigues JCF, Fang J, Smith A, et al. Characterization of a novel organelle in *Toxoplasma gondii* with similar composition and function to the plant vacuole. Mol Microbiol. 2010;76: 1358–1375. doi:10.1111/j.1365-2958.2010.07165.x

31. Warmka J, Hanneman J, Lee J, Amin D, Ota I. Ptc1, a type 2C Ser/Thr phosphatase, inactivates the HOG pathway by dephosphorylating the mitogen-activated protein kinase Hog1. Mol Cell Biol. 2001;21: 51–60. doi:10.1128/MCB.21.1.51-60.2001

32. Mapes J, Ota IM. Nbp2 targets the Ptc1-type 2C Ser/Thr phosphatase to the HOG MAPK pathway. EMBO J. 2004;23: 302–311. doi:10.1038/sj.emboj.7600036

33. González A, Ruiz A, Serrano R, Ariño J, Casamayor A. Transcriptional profiling of the protein phosphatase 2C family in yeast provides insights into the unique functional roles of Ptc1. J Biol Chem. 2006;281: 35057–35069. doi:10.1074/jbc.M607919200

34. Fidock DA, Nomura T, Talley AK, Cooper RA, Dzekunov SM, Ferdig MT, et al. Mutations in the *P. falciparum* Digestive Vacuole Transmembrane Protein PfCRT and Evidence for Their Role in Chloroquine Resistance. Mol Cell. Cell Press; 2000;6: 861–871. doi:10.1016/S1097-2765(05)00077-8

35. Lewis IA, Wacker M, Olszewski KL, Cobbold SA, Baska KS, Tan A, et al. Metabolic QTL analysis links chloroquine resistance in *Plasmodium falciparum* to impaired hemoglobin catabolism. Neafsey DE, editor. PLoS Genet. 2014;10: e1004085. doi:10.1371/journal.pgen.1004085

36. Shen B, Brown KM, Lee TD, Sibley LD. Efficient Gene Disruption in Diverse Strains of Toxoplasma gondii Using CRISPR/CAS9. MBio. 2014;5: e01114–14–e01114–14. doi:10.1128/mBio.01114-14

37. Sidik SM, Hackett CG, Tran F, Westwood NJ, Lourido S. Efficient Genome Engineering of Toxoplasma gondii Using CRISPR/Cas9. Blader IJ, editor. PLoS ONE. 2014;9: e100450–8. doi:10.1371/journal.pone.0100450

38. Kaja S, Payne AJ, Singh T, Ghuman JK, Sieck EG, Koulen P. An optimized lactate dehydrogenase release assay for screening of drug candidates in neuroscience. Journal of Pharmacological and Toxicological Methods. 2015;73: 1–6. doi:10.1016/j.vascn.2015.02.001

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7