Title: Temperate grass allergy season defined by spatio-temporal shifts in airborne

pollen communities

Authors: Georgina L. Brennan[1*], Caitlin Potter[2], Natasha de Vere[2,3], Gareth W. Griffith[2],

Carsten A. Skjøth[4], Nicholas J. Osborne[5,6], Benedict W. Wheeler[5], Rachel N. McInnes[7],

Yolanda Clewlow[7], Adam Barber[7], Helen M. Hanlon[7], Matthew Hegarty[2], Laura

Jones[1,3], Alexander Kurganskiy[4], Francis M. Rowney [5], Charlotte Armitage[8], Beverley Adams-

Groom[4], Col R. Ford[3], Geoff M. Petch[4], The PollerGEN Consortium, and Simon Creer[1*].

Consortium Angela Elliot[9], Carl A. Frisk[4], Roy Neilson[10], Stephen Potter[11], Abdullah M. Rafiq[1],

David, B. Roy[12], Katherine Selby[13], Natascha Steinberg[9]

Affiliations:

[1]Molecular Ecology and Fisheries Genetics Laboratory, School of Natural Sciences, Bangor

University, Bangor, Gwynedd, Wales, UK

[2]Aberystwyth University, Aberystwyth, Ceredigion, Wales, UK

[3]National Botanic Garden of Wales, Llanarthne, Carmarthenshire, Wales, UK

[4]University of Worcester, Worcester, Worcestershire, UK

[5]University of Exeter, Truro, Cornwall, UK

[6]University of New South Wales, Sydney, New South Wales, Australia

[7]Met Office, Exeter, Devon, UK

[8]The Woodland Trust, Kempton Way, Grantham, Lincolnshire, UK.

[9]College of Life and Environmental Science, University of Exeter, Exeter, UK

[10]Ecological Sciences, The James Hutton Institute, Dundee, Scotland, UK

[11]The David Hide Asthma & Allergy Research Centre, St Mary's Hospital, Newport, Isle of

Wight, UK

[12]Centre for Ecology & Hydrology, Crowmarsh Gifford, Wallingford, Oxfordshire, UK.

[13]Environment Department, University of York, Heslington, York, UK

Corresponding authors: Simon Creer (s.creer@bangor.ac.uk) and Georgina L. Brennan

(g.l.b.doonan@gmail.com)

Abstract

Grass pollen is the world's most harmful outdoor aeroallergen and sensitivity varies between

species. Different species of grass flower at different times, but it is not known how airborne

communities of grass pollen change in time and space. Persistence and high mobility of grass

pollen could result in increasingly diverse seasonal pollen communities. Conversely, if grass

pollen does not persist for an extended time in the air, shifting pollen communities would be

predicted throughout the summer months. Here, using targeted high throughput sequencing,

we tracked the seasonal progression of airborne Poaceae pollen biodiversity across Britain,

throughout the grass allergy season. All grass genera displayed discrete, temporally restricted

peaks of pollen incidence which varied with latitude, revealing that the taxonomic composition

of grass pollen exposure changes substantially across the allergy season. By developing more

refined aeroallergen profiling, we predict that our findings will facilitate the exploration of links

between taxon-specific exposure of harmful grass pollen and disease, with concomitant socio-

economic benefits.

46
47

Introduction

Allergens carried in airborne pollen are associated with both asthma [1] and allergic rhinitis (hay fever), negatively affecting 400 million people worldwide [2]. Pollen from the grass family (Poaceae) constitutes the most significant outdoor aeroallergen [3, 4], and more people are sensitised to grass pollen than to any other pollen type [5]. However, despite the harmful impact of grass pollen on human health, current observational studies and forecasts categorize grass pollen at the family level [Poaceae; 6, 7] due to difficulties in differentiating species and genera of grass pollen based on morphology [8]. Furthermore, we cannot predict seasonal variation in airborne grass pollen from the phenology of local grasses at ground level, since airborne pollen can be highly mobile [9, 10] and often does not directly correlate to local flowering times [9]. Understanding the taxon-specific phenology of airborne pollen would fill a significant knowledge gap in our understanding of allergen triggers, with associated benefits to healthcare providers, pharmaceutical industries and the public.

Many species within the subfamilies Pooideae, Chloridoideae, and Panicoideae release allergenic pollen into the atmosphere [5], including *Phleum* spp. (e.g. Timothy grasses), *Dactylis* spp. (Cocksfoot grasses), *Lolium* spp. (Ryegrasses), *Trisetum* spp. (Oatgrasses), *Festuca* spp. (Fescues), *Poa* spp. (Meadow-grasses and Bluegrasses), and *Anthoxanthum* spp. (Vernal grasses). However, it is unknown whether particular grass species contribute more to the prevalence of allergic symptoms and related diseases than others [11]. Whilst some grasses have been identified as more allergenic than others *in vitro* (triggering higher levels of Immunoglobulin E (IgE) antibody production), there is a high degree of cross-reactivity between grass species [12]. In addition, the allergen profiles and the degree of sensitisation

differ between grass species [12, 13] and the allergenicity of grass pollen varies across

seasons [14]. Family-level estimates of grass pollen concentrations cannot therefore be

considered a reliable proxy for either the concentration of pollen-derived aeroallergens or

pollen-induced public health outcomes.

The identification of biodiversity via the high-throughput analysis of taxonomy marker genes

(popularly termed metabarcoding) provides an emerging solution to semi-quantitatively

identify complex mixtures of airborne pollen grains [15-18]. Previous metabarcoding studies

of airborne pollen have been performed at very limited spatial and temporal scales [e.g. 15,

16]. Recent global DNA barcoding initiatives and co-ordinated regional efforts have now

resulted in near complete genetic databases of national native plants, including grasses in

Great Britain [19].

Here, using two complementary DNA barcode marker genes (*rbcL* and ITS2), we characterise

the spatial and temporal distribution of airborne grass pollen throughout the temperate

summer grass pollen season (May-August) across the latitudinal and longitudinal range of

Great Britain (S1 Fig). We hypothesise that (i) there will be discrete temporal incidences of

pollen from different grasses, linked to Poaceae terrestrial phenology, and (ii) the

composition of grass pollen will be homogenous across the UK due to the potential for long

distance transport of windborne pollen grains.

Results and Discussion

Grass pollen occupied distinct temporal windows across the grass allergy season in 2016

(May to August), thereby supporting our hypothesis (i) that species composition of airborne

grass pollen will change throughout the grass allergy season (Fig 1, Fig 2). Time, measured as

number of days after the first sample was collected, is a good predictor of airborne grass

pollen taxon composition using both markers (Fig 1-2; *ITS2*, $LR_{1, 74}$ = 128.8, *P* = 0.001; *rbcL*,

$LR_{1, 71}$ = 46.71, *P* = 0.001). We found that month (coded as a factor in the models) improves

our ability to predict taxonomic composition across the pollen season (Fig 1-2; *ITS2*, $LR_{1, 70}$ =

319.7, *P* = 0.001; *rbcL*, $LR_{1, 67}$ = 217.25, *P* = 0.001). In addition, community-level ordination

reveals that the community as a whole changed across the allergy season (S2 Fig).

Focusing on the more taxonomically specific ITS2 marker dataset, *Alopecurus* and *Holcus*

typically dominated the early grass pollen season (Fig 1), which coincides with typical peaks in

allergic rhinitis [20], but further research will be required to confirm this association. *Lolium*

featured prominently for the majority of the later grass season. The popularity of *Lolium*

species as forage crop means that many varieties have been bred with the potential to

mature at different times throughout the year [21]. While *Lolium* was the dominant species

in airborne grass pollen from July to the end of the sampling period, the total grass pollen

concentration declined in August, indicating that the absolute number of *Lolium* pollen grains

at this time is low (S3 Fig).

The top five genera contributing to airborne pollen, indicated by the relative abundance of

taxonomy marker genes, were *Alopecurus*, *Festuca*, *Lolium*, *Holcus* and *Poa* (Fig 1; S3 Fig).

Each of these genera are widespread in the UK and have been shown to provoke IgE-

mediated responses in grass-sensitised patients [12], providing candidate species for links

with hay fever and asthma exacerbation. Conversely, less prevalent species in the dataset

could contribute disproportionately to the allergenic load. Species such as *Phleum pratense*

have been identified to be a major source of allergenic pollen [5, 22]. However, we found

that *Phleum* made up a very small proportion of metabarcoding reads (Fig 2), corresponding

with the results of an earlier phenological study [23]. Most genera, such as *Phleum*,

Anthoxanthum and *Dactylis*, show distinct and narrow temporal incidence (Fig 2), and could

allow researchers to identify grass species associated with allergenic windows with greater

accuracy.

Changes in species composition over time were localised. We found that peaks in abundance

of airborne pollen occurred at different times at each location during the summer (Fig 1-2).

For example, the relative abundance of airborne grass pollen from the genus *Poa* peaked in

mid-June in Worcester and Bangor but 6-8 weeks later in Invergowrie (Fig 1), probably due to

latitudinal effects on flowering time [7, 24]. This is supported by a significant interaction

between latitude and time of year for both markers (Fig 1-2; *ITS2*, $LR_{68, 1}$= 46.4, P = 0.001;

rbcL,$LR_{66, 1}$= 59.08, P = 0.001), and between longitude and time of year for the ITS2 dataset

(Fig 1-2; $LR_{67, 1}$= 37.5, P = 0.001). Differences in species composition of airborne grass pollen

between the six sampling sites is supported by a significant effect of latitude (Fig 1-2; *ITS2*,

$LR_{1, 73}$ = 73.2, P = 0.001; *rbcL*, $LR_{1, 70}$ = 26.4, P = 0.025) and longitude (Fig 1-2; *ITS2*, $LR_{1, 69}$ =

36.5, P = 0.005; *rbcL*, $LR_{1, 69}$ = 27.10 P = 0.018). These results do not support our hypothesis

(ii) that the composition of grass pollen will be homogenous across the UK, and instead

suggest taxon-specific effects of regional geography and climate which have been

demonstrated for Poaceae pollen as a whole [7].

Observations of first flowering dates from a citizen science project (UKPN;

www.naturescalendar.org.uk) and metabarcoding data show similar sequences of seasonal

progression (Fig 3). First flowering dates of each genus started almost 3-4 weeks prior to the

observation of peaks of grass pollen in the metabarcoding data (Fig 3). Pollen release

(anthesis) occurs approximately 2-3 weeks after the production of flowering heads (heading)

[25], and this is reflected in the metabarcoding data suggesting that local flowering data are

informative for predicting the composition of airborne pollen. Continuing this study over

multiple years would allow us to track long-term, phenological changes in airborne pollen

communities and improve our ability to forecast the seasonal progression of airborne pollen

[26].

Enabled by contemporary molecular biodiversity analytical approaches and mature, curated

DNA barcoding databases, here we provide a comprehensive taxonomic overview of airborne

grass pollen distribution, throughout an entire allergy season and across large geographic

scales. The grass pollen season is defined by discrete temporal windows of different grass

species, with some grass species displaying geographical variation. Temporal pollen

distributions in metabarcoding data follow observed flowering times. The data provide an

important step towards developing species-level grass pollen forecasting. Additionally, the

research presented here leads the way for future studies facilitating understanding of the

relationships between grass pollen and disease, which have significant global public health

relevance and socioeconomic importance.

Methods

Sampling and Experimental Design

We collected aerial samples from six sites across the UK (S3 Table; S1 Fig) using Burkard

Automatic Multi-Vial Cyclone Samplers (Burkard Manufacturing Co. Ltd. Rickmansworth, UK).

The volumetric aerial sampler uses a turbine to draw in air (16.5 litres/min) and aerial

particles are collected, using mini-cyclone technology into 1.5 ml sterile microcentrifuge

tubes located on a carousel (S5 Fig). Each sampling unit was mounted alongside a seven-day

volumetric trap (Burkard Manufacturing Co. Ltd. Rickmansworth, UK) belonging to the Met

Office UK Pollen Monitoring Network, which provided daily pollen count data. In the seven-

day volumetric trap, aerial particles are collected onto an adhesive coated tape supported on

a clockwork-driven drum. The tape is cut into 24 h sections and pollen are identified and

counted under a microscope [7]. Bangor was the only sampling site which was not part of the

pollen monitoring network, but we deployed the same methodology at the Bangor site.

Sampling began in late May 2016 (S4 Table) and during alternate weeks, aerial samples were

collected for seven days for a total of seven weeks between 25th May and 28th August. Exact

sampling dates varied slightly between sites (S4 Table) and a total of 279 aerial samples were

collected.

DNA Extraction, PCR and Sequencing

From the 279 daily aerial samples, 231 were selected for downstream molecular analysis, as

described below. Within each sampling week, two series of three consecutive days were

pooled. Pooled samples were selected based on grass pollen counts obtained by microscopy.

The final, unselected, day was not used in downstream molecular analysis. In total, seventy-

seven pools of DNA were created. In one instance, three consecutive days of pollen samples

were unavailable (Invergowrie, week 2, pool 2) due to trap errors. For this sample, the next

sampling day was selected for pooling (S4 Table). DNA was extracted from daily samples

using a DNeasy Plant Mini kits (Qiagen, Valencia, CA, USA), with some modifications to the

standard protocol as described by [27]. DNA from daily samples was pooled and eluted into

60 µl of elution buffer at the binding stage of the DNeasy Plant Mini kit.

Illumina MiSeq paired end indexed amplicon libraries were prepared following a two-step

protocol as recommended by the manufacturer [28]. Two marker genes were amplified with

universal primer pairs *rbcL*af and *rbcL*r506 [19, 29], and ITS2 and ITS3 [14] (S6 Table). A 5'

universal tail was added to the forward and reverse primers and a 6N sequence was added

between the forward universal tail and the template-specific primer, which is known to

improve clustering and cluster detection on MiSeq sequencing platforms [30] (Integrated

DNA Technologies, Coralville, USA). Round 1 PCR was carried out in a final volume of 25 µL,

including forward and reverse primers (0.2 µM), 1X Q5 HS High-Fidelity Master Mix (New

England Biolabs) and 1 µL of template DNA. Thermal cycling conditions were an initial

denaturation step at 98 °C for 30s; 35 cycles of 98 °C for 10s, 50 °C for 30s, 72 °C for 30s; and

a final annealing step of 72 °C for 5 minutes. Products from the first PCR were purified using

Agencourt AMPure XP beads (Beckman Coulter) with a 1:0.6 ratio of product to AMPure XP

beads.

The second round PCR added the unique identical i5 and i7 indexes and the P5 and P7

Illumina adaptors, along with universal tails complementary to the universal tails used in

round 1 PCR (S4 Table, S5 Table) (Ultramer, by IDT, Integrated DNA Technologies). Round 2

PCR was carried out in a final volume of 25 µL, including forward and reverse index primers

(0.2 µM), 1X Q5 HS High-Fidelity Master Mix (New England Biolabs) and 5 µL of purified PCR

product. Thermal cycling conditions were: 98 °C for 3 min; 98 °C for 30 s, 55 °C for 30 s, 72 °C

for 30 s (10 cycles); 72 °C for 5 min, 4 °C for 10 min. Both PCRs were run in triplicate. The

same set of unique indices were added to the triplicates which were then pooled following

visual inspection on an agarose gel (1.5%) to ensure that indices were added successfully.

Pooled metabarcoding libraries were cleaned a second time using Agencourt AMPure

magnetic bead purification, run on an agarose gel (1.5%) and quantified using the Qubit high

sensitivity kit (Thermo Fisher Scientific, Massachusetts, USA). Positive and negative controls

were amplified in triplicate with both primer pairs and sequenced alongside airborne plant

community DNA samples using the MiSeq. Sequence data, including metadata, are available

at the Sequence Read Archive (SRA) using the project accession number SUB4136142.

Bioinformatic Analysis

Initial sequence processing was carried out following a modified version of the workflow

described by de Vere *et al.* [27]. Briefly, raw sequences were trimmed using Trimmomatic

v0.33 (*42*) to remove short reads (<200bp), adaptors and low quality regions. Reads were

merged using FLASH v 1.2.11 [27, 31], and merged reads shorter than 450bp were excluded.

Identical reads were merged using fastx-toolkit (v0.0.14), and reads were split into ITS2 and

rbcL based on primer sequences.

To prevent spurious BLAST hits, custom reference databases containing *rbcL* and ITS2

sequences from UK plant species were generated. While all native species of the UK have

234 been DNA barcoded [19], a list of all species found in the UK was generated in order to gain

235 coverage of non-native species. A list of UK plant species was generated by combining lists of

236 native and alien species [32] with a list of cultivated plants obtained from Botanic Gardens

237 Conservation International (BGCI) which represented horticultural species. All available *rbcL*

238 and ITS2 records were downloaded from NCBI Genbank, and sequences belonging to UK

239 species were extracted using the script 'creatingselectedfastadatabase.py', archived on

240 GitHub.

242 Metabarcoding data was searched against the relevant sequence database using blastn [33],

243 via the script 'blast_with_ncbi.py'. The top twenty blast hits were tabulated

244 ('blast_summary.py'), then manually filtered to limit results to species currently present in

245 Great Britain. Reads occurring fewer than four times were excluded from further analysis.

246 All scripts used are archived on GitHub: https://doi.org/10.5281/zenodo.1305767.

248 **Statistical Analysis**

249 To understand how the grass pollen composition changed with space and time, the effect of

250 time (measured as the number of days after the first sampling date), latitude and longitude

251 of sampling location were included in a two-tailed generalized linear model using the

252 'manyglm' function in the package 'mvabund' [34]. The proportion of sequences was set as

253 the response variable; proportion data was used as this has been shown to be an effective

254 way of controlling for differences in read numbers [35]. The effect of time, latitude,

255 longitude, month (coded as a factor), and the interaction between time and latitude were

256 included as explanatory variables in the models. In addition to these explanatory variables,

the interaction between time and longitude was included in a model to analyse the ITS2 data

(S6 Table).

The data best fit a negative binomial distribution, most likely due to the large number of

zeros (zeros indicate that a grass genus is absent from a sampling location), resulting in a

strong mean-variance relationship in the data (S6 Fig). The proportion of sequences was

scaled by 1000 and values were converted to integers so that a generalized linear model with

a negative binomial distribution could be used. Model selection was based by Akaike

Information Criterion (AIC) (S6 Table) and visual inspection of the residuals against predicted

values from the models (S7 Fig).

In order to compare the metabarcoding data with flowering time data, we used phenological

records of first flowering collected in 2016 by citizen scientists from the UK's Nature's

Calendar (www.naturescalendar.org.uk). First flowering time was compared to genus-level

ITS2 metabarcoding data for three species: *Alopecurus pratensis*, *Dactylis glomerata* and

Holcus lanatus. As grass pollen could only be reliably identified to genus level in the

metabarcoding data, the taxa compared may not have been exactly equivalent since both

Alopecurus and *Holcus* contain other widespread species within the UK. However, *Alopecurus*

pratensis and *Holcus lanatus* are the most abundant species within their respective genera.

The comparison was only carried out for ITS2 data because two of the three genera were not

identified by the *rbcL* marker.

NMDS ordination was carried out using package 'VEGAN' in R [36], based on the proportion

of total high-quality reads contributed by each grass genus, using Bray-Curtis dissimilarity (S2

Fig). Ordination is used to reduce multivariate datasets (e.g. abundances of many species)

into fewer variables that reflect overall similarities between samples. A linear model was

carried out using the 'lm' function within the 'stats' package in R, in order to investigate the

relationship between the number of reads obtained for each genus using the rbcL and ITS2

marker.

References

1.   Blackley CH. Experimental Researches on the Causes and Nature of Catarrhus Æstivus
(hay-fever Or Hay-asthma)1873 1873. 202 p.
2.   Marks G, Pearce N, Strachan D, Asher I. Global Burden of Disease Due to Asthma. The
Global Asthma Report 2014. Auckland, New Zealand: The Global Asthma Network; 2014. p.
16-21.
3.   Bauchau V. Prevalence and rate of diagnosis of allergic rhinitis in Europe. Eur Respir J.
2004;24(5):758-64. doi: 10.1183/09031936.04.00013904.
4.   Bousquet PJ, Chinn S, Janson C, Kogevinas M, Burney P, Jarvis D, et al. Geographical
variation in the prevalence of positive skin tests to environmental aeroallergens in the
European Community Respiratory Health Survey I. Allergy. 2007;62(3):301-9. doi:
10.1111/j.1398-9995.2006.01293.x.
5.   García-Mozo H. Poaceae pollen as the leading aeroallergen worldwide: A review.
Allergy. 2017;72(12):1849-58. doi: 10.1111/all.13210.
6.   Emberlin J, Jaeger S, Dominguez-Vilches E, Soldevilla CG, Hodal L, Mandrioli P, et al.
Temporal and geographical variations in grass pollen seasons in areas of western Europe: an
analysis of season dates at sites of the European pollen information system. Aerobiologia.
2000;16(3-4):373-9.
7.   Emberlin J, Jones S, Bailey J, Caulton E, Corden J, Dubbels S, et al. Variation in the start
of the grass pollen season at selected sites in the United Kingdom 1987–1992. Grana.
1994;33(2):94-9.
8.   Mander L, Li M, Mio W, Fowlkes CC, Punyasena SW. Classification of grass pollen
through the quantitative analysis of surface ornamentation and texture. Proc Biol Sci.
2013;280(1770):20131905. doi: 10.1098/rspb.2013.1905.
9.   Estrella N, Menzel A, Krämer U, Behrendt H. Integration of flowering dates in
phenology and pollen counts in aerobiology: analysis of their spatial and temporal coherence
in Germany (1992–1999). Int J Biometeorol. 2006;51(1):49-59. doi: 10.1007/s00484-006-
0038-7.
10.   Skjøth CA, Sommer J, Stach A, Smith M, Brandt J. The long range transport of birch
(Betula) pollen from Poland and Germany causes significant pre-season concentrations in
Denmark. Clinical & Experimental Allergy. 2008;37(8):1204-12.
11.   McInnes RN, Hemming D, Burgess P, Lyndsay D, Osborne NJ, Skjøth CA, et al. Mapping
allergenic pollen vegetation in UK to study environmental exposure and human health. Sci
Total Environ. 2017;599-600:483-99. doi: 10.1016/j.scitotenv.2017.04.136.

12.   van Ree R, van Leeuwen WA, Aalberse RC. How far can we simplify in vitro diagnostics
for grass pollen allergy?: A study with 17 whole pollen extracts and purified natural and
recombinant major allergens. J Allergy Clin Immunol. 1998;102(2):184-90.
13.   Moingeon P, Peltre G, Bergmann KC. Rationale for a five-grass pollen vaccine. Clin Exp
Allergy Rev. 2008;8(1):12-4. doi: 10.1111/j.1472-9733.2008.00124.x.
14.   de Weger LA, Beerthuizen T, Gast-Strookman JM, van der Plas DT, Terreehorst I,
Hiemstra PS, et al. Difference in symptom severity between early and late grass pollen season
in patients with seasonal allergic rhinitis. Clin Transl Allergy. 2011;1(1):18. doi: 10.1186/2045-
7022-1-18.
15.   Kraaijeveld K, de Weger LA, García MV, Buermans H, Frank J, Hiemstra PS, et al.
Efficient and sensitive identification and quantification of airborne pollen using next-
generation DNA sequencing. Mol Ecol Resour. 2014;15(1):8-16. doi: 10.1111/1755-
0998.12288.
16.   Korpelainen H, Pietiläinen M. Biodiversity of pollen in indoor air samples as revealed
by DNA metabarcoding. Nord J Bot. 2017;35(5):602-8. doi: 10.1111/njb.01623.
17.   Deiner K, Bik H, Mächler E, Seymour M, Lacoursière-Roussel A, Altermatt F, et al.
Environmental DNA metabarcoding: Transforming how we survey animal and plant
communities. Molecular Ecology. 2017;26(21):5872-95.
18.   Creer S, Deiner K, Frey S, Porazinksa D, Taberlet P, Kelley Thomas W, et al. The
ecologist's field guide to sequence-based identification of biodiversity. Methods in Ecology
and Evolution. 2016;7(9):1008-18.
19.   de Vere N, Rich TCG, Ford CR, Trinder SA, Long C, Moore CW, et al. DNA barcoding the
native flowering plants and conifers of Wales. PLoS One. 2012;7(6):e37945. doi:
10.1371/journal.pone.0037945.
20.   England PH. GP in-hours consultations bulletin: 25 August 2016 week 33. Birmingham,
UK: PHE Real-time Syndromic Surveillance Team, 2016.
21.   RGCL. Recommended Grass and Clover Lists. In: Society BG, editor.
http://wwwbritishgrasslandcom/system/files/uploads/RGCL%20Handbook%202017%206%20
WEBpdf. Cheshire, UK: British Grassland Society; 2017.
22.   D'Amato G, Vitale C, Sanduzzi A, Molino A, Vatrella A, D'Amato M. Allergenic Pollen
and Pollen Allergy in Europe. Allergy. 2007;62(9):976-90.
23.   Kmenta M, Bastl K, Berger U, Kramer MF, Heath MD, Pätsi S, et al. The grass pollen
season 2015: a proof of concept multi-approach study in three different European cities.
World Allergy Organ J. 2017;10(1):31. doi: 10.1186/s40413-017-0163-2.
24.   Chapman D. Greater phenological sensitivity to temperature on higher Scottish
mountains: new insights from remote sensing. Global Change Biology. 2013;19:3463–71.
25.   Emecz TI. The effect of meteorological conditions on anthesis in agricultural grasses.
Annals of Botany. 1962;26(102):159-72.
26.   Thackeray SJ, Henrys PA, Hemming D, Bell JR, Botham MS, Burthe S, et al.
Phenological sensitivity to climate across taxa and trophic levels. Nature.
2016;535(7611):241-5.
27.   Hawkins J, de Vere N, Griffith A, Ford CR, Allainguillaume J, Hegarty MJ, et al. Using
DNA Metabarcoding to Identify the Floral Composition of Honey: A New Tool for Investigating
Honey Bee Foraging Preferences. PLoS One. 2015;10(8):e0134735. doi:
10.1371/journal.pone.0134735.
28.   Illumina. 16S Metagenomic Library Preparation. San Diego, California, USA: Illumina,
2013.

29.   Kress WJ, Erickson DL. A two-locus global DNA barcode for land plants: the coding
rbcL gene complements the non-coding *trnH-psbA* spacer region. PLoS One. 2007;2(6):e508.
doi: 10.1371/journal.pone.0000508.
30.   Miya M, Sato Y, Fukunaga T, Sado T, Poulsen JY, Sato K, et al. MiFish, a set of universal
PCR primers for metabarcoding environmental DNA from fishes: detection of more than 230
subtropical marine species. Royal Society open science. 2015;2(7):150088.
31.   Magoc T, Salzberg SL. FLASH: fast length adjustment of short reads to improve
genome assemblies. Bioinformatics. 2011;27(21):2957-63. doi:
10.1093/bioinformatics/btr507.
32.   Stace C. New flora of the British Isles. Cambridge, UK: Cambridge University Press;
2010.
33.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC Bioinformatics. 2008;10:421.
34.   Wang YI, Naumann U, Wright ST, Warton DI. mvabund–an R package for model-based
analysis of multivariate abundance data. Methods in Ecology and Evolution. 2012;3(3):471-4.
35.   McMurdie PJ, Holmes S. Waste not, want not: why rarefying microbiome data is
inadmissible. PLoS Computational Biology. 2014;10(4):e1003531.
36.   Dixon P. VEGAN, a package of R functions for community ecology. Journal of
Vegetation Science. 2003;14(6):927-30.
37.   Cope C, Gray A. Grasses of the British Isles. 13 ed. London, UK: Botanical Society of
the British Isles; 2009.

List of Supplementary Materials

391 Supplementary text
392 References
393 S1 – S7 Figs
394 S1 List
395 S1 – S6 Tables

Acknowledgements

We thank John Kenny, Pia Koldkjær, Richard Gregory, and Anita Lucaci of the Liverpool

Centre for Genomic Research for sequencing support. We acknowledge the computational

services & support of the Supercomputing Wales project, which is part-funded by the

European Regional Development Fund (ERDF) via Welsh Government. We thank the Botanic

Gardens Conservation International (BGCI) for access to the list of plant collections in the

National Gardens in the UK and Ireland. We thank the Met Office network for providing

404 additional observational grass pollen count data and Jonathan Winn, UK Met Office for

405 ArcGIS assistance on S1 Fig. We are grateful to the Woodland Trust and Centre for Ecology &

406 Hydrology for supplying the UK Phenology Network data and to the citizen scientists who

407 have contributed to the latter scheme. Final thanks to Wendy Grail and technical support

408 staff at Bangor University.

410 **Author Contributions:** S.C., N.dV., G.W.G., R.N.M., N.J.O., C.A.S., Y.C. and G.L.B. conceived and

411 designed the study; B.A-G., G.L.B., G.P., A.E., R.N., S.P., K.S., and N.S. collected samples and

412 counted pollen; G.L.B. performed laboratory work, supported by S.C.; N.dV., C.R.F., L.J. and

413 S.C. contributed methods; C.P. and G.L.B. analysed the data and G.L.B., C.P. and S.C.

414 produced the first draft of the manuscript. All authors contributed substantially to the final

415 submitted manuscript.

417 **Funding:** This work was supported by the Natural Environment Research Council

418 (https://nerc.ukri.org/), awarded to SC (NE/N003756/1), CS (NE/N002431/1), NO

419 (NE/N002105/1) and NdV and GG (NE/N001710/1). The funders had no role in study design,

420 data collection and analysis, decision to publish, or preparation of the manuscript.

422 **Competing interests:** The authors are not aware of any competing interests.

424 **Data and materials availability:** All sequence data (including metadata) are available at the

425 Sequence Read Archive (SRA) using the project accession number SUB4136142. Archived

426 sequence data was used to generate Fig 1 to 3 (including S2-S4 and S6-S7 Figs). First

427 flowering data used in Fig 3 was obtained from Nature's Calendar, Woodland Trust and is

available upon request. The sequence analysis pipeline is available at

https://github.com/colford/nbgw-plant-illumina-pipeline.

Figure Legends

Fig 1. Abundance of the most common airborne grass pollen taxa throughout the grass allergy

season. The five most abundant grasses (expressed as proportion of total reads), depicted

alongside the total proportion of reads assigned to family Poaceae. Markers used to identify

grass pollen are stated in the top panel label. Due to errors in sampling equipment, only 4

weeks of samples were collected at the York sampling site. Sampling sites are indicated in the

right panel label abbreviated as follows: BNG = Bangor; EXE = Exeter; ING = Invergowrie; IOW

= Isle of Wight; WOR = Worcester; YORK = York. A map of sampling locations can be found in

S1 Fig.

Fig 2. Abundance of airborne grass pollen taxa throughout the grass allergy season.

Abundance of rare grasses (expressed as proportion of total reads). Sampling sites are

indicated in the top panel, followed by the marker used to identify grass pollen. Due to errors

in sampling equipment, only 4 weeks of samples were collected at the York sampling site.

Note that the y axes differ between panels. Refer to Fig 1 for site name abbreviations.

Fig 3. Airborne grass pollen observed 3-4 weeks after first flowering dates. Comparison of

genus incidence in metabarcoding data with records of first flowering dates in 2016 from the

citizen science project Nature's Calendar (www.naturescalendar.org.uk) for (A) *Alopecurus*

pratensis, (B) *Dactylis glomerata* and (C) *Holcus lanatus*. Each grey point represents the

earliest time of flower heading as observed by a participant in the project. Coloured points

represent metabarcoding samples, with the size of the point representing the proportion of

total reads assigned to the relevant genus. Yellow shaded areas represent the expected

flowering period as described in [37], with darker shades showing the 'main' flowering

period.