Genome-wide association studies in Samoans give insight into the genetic

architecture of fasting serum lipid levels

Jenna C. Carlson[1,2]*, Daniel E. Weeks[1,2]¶, Nicola L. Hawley[3], Guangyun Sun[4], Hong

Cheng[4], Take Naseri[5], Muagututi'a Sefuiva Reupena[6]; Ranjan Deka[4]¶, Stephen T.

McGarvey[7,8]¶, Ryan L. Minster[1]¶

[1] Department of Human Genetics, Graduate School of Public Health, University of
Pittsburgh, Pittsburgh, Pennsylvania, United States of America
10
11   [2] Department of Biostatistics, Graduate School of Public Health, University of Pittsburgh,
12   Pittsburgh, Pennsylvania, United States of America
13
14   [3] Department of Epidemiology, School of Public Health, Yale University, New Haven,
15   Connecticut, United States of America
16
17   [4] Department of Environmental Health, College of Medicine, University of Cincinnati,
18   Cincinnati, Ohio, United States of America
19
20   [5] Ministry of Health, Government of Samoa, Apia, Samoa
21
22   [6] Lutia i Puava ae Mapu i Fangalele, Apia, Samoa
23
24   [7] International Health Institute and Department of Epidemiology, School of Public
25   Health, Brown University, Providence, Rhode Island, United States of America
26
27   [8] Department of Anthropology, Brown University, Providence, Rhode Island, United
28   States of America
29

30   * Corresponding Author

31   Email: jnc35@pitt.edu

33   ¶DEW, RD, STM, RLM are Joint Senior Authors

Abstract

The current understanding of the genetic architecture of lipids has largely come from genome-wide association studies. To date, few studies have examined the genetic architecture of lipids in Polynesians, and none have in Samoans, whose unique population history, including many population bottlenecks, may provide insight into the biological foundations of variation in lipid levels. Here we performed a genome-wide association study of four fasting serum lipid levels: total cholesterol (TC), high-density lipoprotein (HDL), low-density lipoprotein (LDL), and triglycerides (TG) in a sample of 2,849 Samoans, with validation genotyping for associations in a replication cohort comprising 1,798 Samoans and American Samoans. We identified multiple genome-wide significant associations ($P < 5 \times 10^{-8}$) previously seen in other populations – *APOA1* with TG, *CETP* with HDL, and *APOE* with TC and LDL – and several suggestive associations ($P < 1 \times 10^{-5}$), including an association of variants downstream of *MGAT1* and *RAB21* with HDL. However, we observed different association signals for variants near *APOE* than what has been previously reported in non-Polynesian populations. The association with several known lipid loci combined with the newly-identified associations with variants near *MGAT1* and *RAB21* suggest that while some of the genetic architecture of lipids is shared between Samoans and other populations, part of the genetic architecture may be Polynesian-specific.

Key words: complex trait genetics; founder population genetics; lipids; genome-wide association study; Polynesia

Introduction

The Samoan Islands, comprising both the U.S. Territory of American Samoa (American Samoan) and the Independent State of Samoa (Samoa), have experienced a rise in prevalence of cardiovascular disease and other non-communicable diseases in the last 30 years partly due to economic modernization, rapid urbanization, and lifestyle changes such as increased caloric intake and sedentary behavior [1–3]. A 2010 population-based survey in Samoa, which gathered the "discovery cohort" studied further here, found that many Samoans are at elevated risk of cardiovascular disease based on known risk factors—increased total cholesterol (TC), low-density lipoprotein cholesterol (LDL), and triglycerides (TG), as well as decreased high-density lipoprotein (HDL) [4–6]—with 47% of the 2,938 adult Samoans examined in the study having elevated TC (≥ 5.2 mmol/L), 88% of men and 91% of women having elevated LDL (> 2.59 mmol/L), and 43% of women having low HDL (< 1.29 mmol/L) [1].

Our current understanding of the genetic component of serum lipid level variation has been largely due to genome-wide association studies (GWAS) [7]. The Global Lipids Genetics Consortium found strong evidence for 157 loci associated with one or more of these traits using a sample of 188,577 individuals of European, East Asian, South Asian, and African ancestry [8]. However, few GWAS of serum lipid levels have been conducted in Pacific Islanders [9, 10] and to our knowledge only one has included a small number of Polynesians [11]. Previous studies have estimated the heritability of serum lipid levels in Samoans, ranging from 16% for HDL to 23% for TG, and have identified genetic susceptibility loci via linkage analysis [12], warranting further study of

80 the genetic architecture of serum lipid levels in Samoans. Samoans are a genetically-

81 isolated founder population, with unique evolutionary history, making them particularly

82 useful in genomic studies [13, 14]. Thus, genomic studies of serum lipid levels could

83 reveal novel lipid-altering loci specific to Pacific Islander populations, as well as highlight

84 susceptibility loci shared with global populations.

86   Here we report the results of a GWAS of fasting TC, LDL, HDL, and TG in up to

87 2,849 individuals from independent Samoa followed by replication in up to 1,798

88 individuals from independent Samoa and American Samoa, as part of ongoing genome-

89 wide association studies of cardiometabolic disease and adiposity-related traits in the

90 Samoan Islands [1]. We identified multiple genome-wide significant associations

91 previously seen in other populations – *APOA1* with TG, *CETP* with HDL, and *APOE*

92 with TC and LDL – and several suggestive associations, including an association

93 between variants downstream of *MGAT1* and *RAB21* with HDL.

Methods

Discovery Cohort and Genotyping

97   The discovery cohort data are available from dbGaP (accession number:

98 phs000914.v1.p1). The discovery cohort of 2,849 individuals is drawn from a

99 population-based sample recruited from Samoa in 2010 (Table 1). The sample

100 selection, data collection methods, and phenotyping, including the laboratory assays for

101 serum lipid and lipoprotein levels, have been previously reported [1, 13]. Briefly, serum

102 lipid levels were derived from fasting whole blood samples collected after a minimum

10-hour overnight fast. Genotyping was performed using Genome-Wide Human SNP

6.0 arrays (Affymetrix). Extensive quality control was conducted on the basis of a

pipeline developed by Laurie et al [15]. Additional details for sample genotyping and

genotype quality control are described in Minster et al [13]. This study was approved by

the institutional review board of Brown University and the Health Research Committee

of the Samoa Ministry of Health. All participants gave informed consent. Imputation was

not performed in this study because prior experience with this population using extant

imputation panels such as the Phase 3 1000 Genomes panel showed that the resulting

imputed genotypes did not correlate well with observed genotypes [13].

Table 1. Demographic, anthropometric, and blood biochemistry statistics of the genotyped discovery and

replication cohorts.

	2010 Discovery Cohort Samoa, n = 2849		2002–2003 Replication Sample Set Samoa, n = 490		1994–1995 Replication Sample Set Samoa, n = 468		2002–2003 Replication Sample Set American Samoa, n = 592		1994–1995 Replication Sample Set American Samoa, n = 248	
	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men
	n = 1703	n = 1146	n = 245	n = 245	n = 243	n = 225	n = 337	n = 255	n = 137	n = 111
age (years)	44.8 (11.10)	45.6 (11.10)	44 (17.00)	40.7 (16.30)	43.3 (8.60)	43.2 (8.90)	43 (16.00)	43.2 (16.60)	43.3 (9.10)	45.5 (10.60)
BMI (kg/m^2)	34.8 (6.70)	31.3 (5.90)	33.2 (7.70)	28.8 (5.40)	31.9 (5.70)	28.8 (5.00)	36.5 (8.40)	33.4 (7.60)	37.4 (6.90)	34.9 (6.00)
total cholesterol (mg/dL)	199.3 (36.10)	200.3 (38.70)	202.3 (35.90)	195.5 (40.60)	199.1 (37.20)	197.4 (35.00)	187.1 (38.60)	189.6 (37.80)	197.8 (31.20)	194.1 (31.00)
HDL (mg/dL)	46.5 (10.80)	43.7 (11.20)	47.1 (10.30)	46.3 (11.20)	43.4 (11.00)	42.5 (11.40)	40.9 (8.80)	40 (8.80)	34.8 (7.60)	32.8 (8.30)
LDL (mg/dL)	130 (32.60)	129.6 (35.30)	130.3 (32.80)	128.6 (37.50)	140.2 (34.80)	133.5 (33.00)	118.5 (33.70)	118.9 (35.10)	136.6 (32.40)	136 (37.50)
trigylcerides (mg/dL)	114.9 (80.50)	139.3 (113.10)	110.9 (58.90)	120.1 (91.30)	85.6 (72.80)	102.8 (97.20)	130.8 (78.10)	200.2 (206.50)	118.9 (63.40)	168.2 (109.00)
fasting glucose (mg/dL)	107.6 (56.20)	107.4 (54.80)	102.2 (46.40)	98.6 (49.80)	90.9 (38.00)	85.3 (18.90)	104.2 (50.30)	108.8 (54.70)	107.3 (49.80)	125.2 (61.70)

Replication Cohort and Genotyping

The replication cohort of 1,798 individuals contains two sample sets recruited

from Samoa and American Samoa (Table 1). Although there is substantial economic

variation across the two polities, with American Samoans generally having a higher

standard of living, the Samoans from both territories form a single socio-cultural unit

with frequent exchange of mates; genetically, they represent a single homogenous

population [3, 16]. The first sample set, referred to as the 1990–95 replication sample

set, contains 716 unrelated individuals derived from a longitudinal study of adiposity and

cardiovascular disease risk factors among adults from American Samoa and Samoa

(Table 1). Detailed descriptions of the sampling and recruitment have been reported

previously [17–19]. Briefly, participants were recruited from 46 villages and worksites in

American Samoa in 1990 and 9 villages in Samoa (Western Samoa, at the time of

recruitment) in 1991 and followed up four years later in 1994 and 1995, respectively. All

participants were, at baseline, free of self-reported history of heart disease,

hypertension, or diabetes. This study was approved by the institutional review board of

the Miriam Hospital, Providence, RI. All participants gave informed consent. The second

sample set, referred to as the 2002–03 replication sample set, contains 1,082

individuals from American Samoa and Samoa and was drawn from an extended family-

based genetic linkage analysis of cardiometabolic traits (Table 1) [12, 20–23]. Probands

and relatives were unselected for obesity or related phenotypes. The recruitment

process and criteria used for inclusion in this study have been described in detail

previously [21, 23]. This study was approved by the institutional review board of Brown

University and research ethics review committees in both Samoa and American Samoa.

All participants gave informed consent. Imputation was not performed in these studies

for the same rationale as the discovery cohort above, but also because genome-wide

marker data was not available for the samples in these studies.

In both replication sample sets, blood samples were collected in the morning

after a minimum of 10 hours fasting, from which serum lipid levels were derived using

assay methods published previously [12, 17]. Genotyping of variants selected for

validation in the replication cohort (described below) was performed using custom-

designed TaqMan OpenArray Real-Time PCR assays (Applied Biosystems). SNPs that

could not be genotyped using OpenArray assays were genotyped individually using

TaqMan SNP Genotyping assays (Applied Biosystems). Eight variants could not be

genotyped due to technical difficulties.

Statistical Analyses

Prior to association analyses, residuals were generated for all four lipid traits.

First, traits were transformed to normality with the Box–Cox power transformation;

secondly, model selection was performed using step-wise linear regression with initial

model covariates previously associated with serum lipid levels: age, age^2, sex, log-

transformed BMI, fasting glucose, smoking status, farming status (as a measure of

physical activity), and interactions between age, age^2, and sex. The final TC model

adjusted for age, age^2, sex, age × sex, and age^2 × sex; the final LDL and TG models

adjusted for age, age^2, sex, and age^2 × sex; the final HDL model adjusted for age and

sex.

Preliminary associations were performed, and variants were selected for

validation without consideration of hypolipidemic medication use, as it was not

measured. However, participants did self-report use of heart disease medication.

Sensitivity analysis revealed that this self-reported use of medication to treat heart

disease was significantly associated with TC and LDL (results not shown); individuals

reporting such medication use (n = 17) were excluded from analyses. The prioritization

of variants for validation genotyping was updated using these analyses, but only after

available resources were fully expended. Unfortunately, not all variants that should have

been prioritized for validation genotyping were successfully genotyped. All results

presented are those of the corrected analyses, removing the individuals with heart

disease medication use.

Additional sensitivity analysis was performed for TG by excluding one outlying

observation (i.e., TG > 4 standard deviations above mean); results did not change

qualitatively, and, since the recorded value was within the range of plausible values for

TG, the individual was retained for presented analyses.

Association between lipid residuals and autosomal genotypes of 659,492 SNPs

with minor allele frequency (MAF) ≥ 0.05 and Hardy-Weinberg Equilibrium (HWE) test P

value ≥ 5×10^{-5} was assessed using linear mixed modelling in GenABEL, including

previously-derived empirical kinship estimates to adjust for subject relatedness [13, 24].

The association between X-chromosome genotypes and the lipid phenotypes were

calculated in GenABEL, without adjustment using the empirical kinship estimates.

Genomic inflation due to population stratification and cryptic relatedness was assessed

by estimating λ_{GC} using the lower 90% of the P value distribution [25]. GWAS P values

in the discovery cohort (P_D) were compared to a threshold for genome-wide significance

of $P_D < 5 \times 10^{-8}$ and a suggestive association threshold of $P_D < 1 \times 10^{-5}$. Statistical

power to detect signals at these thresholds was calculated using the Genetic Power

Calculator [26].

Gene-set enrichment analysis with MAGENTA was also performed to identify any

biological pathways enriched for discovery association signals [27]. Briefly, gene scores

were obtained from the most significant P value among SNPs located within each gene

using the association results from each lipid GWAS. Genes scores were adjusted for

confounding factors including gene size, number of variants, and linkage disequilibrium-

related properties by using step-wise multiple linear regression. The 95th percentile of

all gene scores was used as the enrichment cutoff for each trait [28]. Gene-set

enrichment P values were obtained for highly ranked gene scores. Gene sets were

obtained from Gene Ontology (April 2010), pathway information from the Ingenuity

(June 2008) and KEGG (June 2010), and biological processes and molecular function

from PANTHER (January 2010).

For each of the lipid traits, the INRICH program [29] was used to test for

enrichment of known genes (as constructed from Teslovich et al. [30] and Willer et al

[8]). INRICH tests if more known genes are contained in associated intervals than

expected by chance, using permutation based on 1 million replicates to generate

experiment-wide empirical P values. For each lipid trait, we defined the associated

intervals as 100 kb intervals centered on the most significant SNP within association

peaks with $P_D < 1 \times 10^{-4}$.

We selected 21 regions demonstrating at least suggestive association for

association validation in the replication cohort. An additional 10 regions which should

have selected for validation were not followed-up because their exclusion was based on

preliminary analyses that included 17 participants taking heart disease medication—

participants who were ultimately excluded from these studies. The variant from each

locus with smallest P value across the four lipid scans (defined as 1 Mb windows

surrounding the peak SNP) or a proxy SNP in high linkage disequilibrium with the

lowest-P value SNP was selected as representative of the locus for replication

genotyping.

Statistical association was measured in the 1990–95 and 2002–03 replication

sample sets independently, and results were combined using meta-analysis (see

below). Association analyses for both sample sets were performed using GenABEL [31]

in R [32], using the same regression models as in the discovery cohort but additionally

adjusting for polity (American Samoa or Samoa); the 2002-03 sample set was

additionally adjusted using expected kinship, as derived from familial pedigree

information [33].

Prior to meta-analysis, quality control was performed using EasyQC to check for

strand and allele frequency consistency [34]. *P* value-based meta-analysis using

sample sizes as weights was performed using METAL [35] to generate two *P* values:

one for the meta-analysis of the two replication cohorts (P_R) and one for the replication

cohorts and discovery sample together (P_{DR}). Resulting meta-analysis signals were

evaluated based on genome-wide significance and suggestive thresholds (as described

above) and by the contribution of the replication sample to the signal. Effect directions

for meta-analysis results of peak SNPs were qualitatively compared to those of

previously reported lead SNPs.

For ease of reference, any locus identified here with a corresponding signal

within 1 mega base pairs (Mb) in a prior lipid study is referred to by the previously

prescribed locus name [8, 10]; for loci not previously associated with lipid traits, the

symbol of the gene nearest the peak SNP in the locus or the hyphen-separated symbols

of the nearest two genes is used as the locus label.

Results

The demographic, anthropometric, and biochemical characteristics of the 2,849

participants composing the discovery cohort for this GWAS of serum lipids levels and

the 1,798 participants composing the replication cohort are presented in Table 1. A

detailed description of the discovery cohort and its trends compared to the historical

sample sets making up the replication cohort has been previously reported [1]. Briefly,

the average age was similar for all cohorts; average BMI was higher among women

compared to men and in American Samoa compared to Samoa; average BMI for men

and women in Samoa was higher in more recent studies; average lipid levels are largely

similar across cohorts with minor exceptions.

We assessed 659,492 unique genome-wide markers for association with 4

traits—TC, HDL, LDL and TG—in up to 2,849 Samoans in the discovery cohort.

Relatedness within the discovery cohort was well-controlled using the empirical kinship

coefficients; λ_{GC} ranged between 1.03 and 1.07 for the four lipid traits (Figs S2, S4, S6,

and S8 in S1 Appendix). We observed 38 genome-wide suggestive or significant

associations across 31 loci from the four GWAS (Fig 1; Figs S1, S3, S5, and S7 in S1

Appendix; Tables S1, S7, S10, and S13 in S1 Appendix).

Fig 1. Manhattan plots for GWAS of four lipid traits in the discovery cohort of

2,849 Samoans. The dashed and solid lines denote genome-wide suggestive and

genome-wide significant P value thresholds ($P < 1 \times 10^{-5}$ and $P < 5 \times 10^{-8}$,

respectively). Peaks are labeled with the candidate gene or closest gene in the region if

they have at least suggestive association in the discovery cohort for at least one trait

and demonstrate evidence of replication or have been previously associated.

Genome-wide significant association was observed in the discovery cohort

between all four traits and markers near *APOE*: TC and rs4420638 ($P_D = 2.67 \times 10^{-16}$,

Fig 2E); HDL and rs4420638 ($P_D = 9.07 \times 10^{-9}$, Fig 2F); LDL and rs1160985 ($P_D = 2.61$

$\times 10^{-20}$; Fig 2G); and TG and rs4420638 ($P_D = 7.44 \times 10^{-10}$, Fig 2H). Additionally, HDL

was associated with markers near *CETP* (rs289708, P_D = 1.19 × 10^{-11}), and TG, with

APOA1 (rs6589566, P_D = 3.98 × 10^{-18}, Fig 2C). Suggestive associations were observed

between lipid levels and markers at an additional 28 loci, including the *MGAT1* and

RAB21 loci and HDL (Fig 2A,D), and *APOA1* with TC (rs3741298, P_D = 1.63 × 10^{-7}, Fig

2B). We had 80% power at $α$ = 1 × 10^{-5} and $α$ = 5 × 10^{-8} to detect SNPs that account

for 1.0% and 1.5%, respectively, of the residual variance in a phenotype.

Fig 2. Regional association plots for selected loci. Regional association plots

generated in LocusZoom [36] showing –log10(*P* values) for SNPs in the (A) *MGAT1*

locus and HDL, (B) *APOA1* locus and TC, (C) *APOA1* locus and TG, (D) *RAB21* locus

and HDL, and the *APOE* locus and (E) TC, (F) HDL, (G) LDL, and (H) TG. Points are

color coded within each plot according to pairwise linkage disequilibrium (r^2) with the

labeled SNPs; the saturation of the color of each plotted SNP measures the linkage

disequilibrium (r^2) with the labeled SNP sharing the same color.

Gene-set enrichment analysis with MAGENTA highlighted, at a < 5% false-

discovery rate (FDR), several lipid homeostasis pathways and gene ontologies for HDL

and TG (Tables S8 and S14 in S1 Appendix). Four gene sets were below the FDR for

both HDL and TG: HDL particle remodeling, reverse cholesterol transport, cholesterol

efflux, and phospholipid efflux. An additional 12 gene sets were implicated for HDL and

three gene sets for TG. The HDL particle remodeling and reverse cholesterol transport

gene sets had significant enrichment for TC (Table S3 in S1 Appendix), and a single

gene set was implicated with LDL, the amylase pathway (Table S11 in S1 Appendix). All

four traits had significant enrichment for known TC, HDL, LDL, and TG loci using the

INRICH method (Tables S5, S9, S12, and S15 in S1 Appendix).

Validation of peak SNPs was attempted for 21 loci. At loci with multiple

associated variants, the most significant variant was chosen as representative of the

locus. For some loci, the exclusion of participants using self-reported heart disease

medication resulted in a different peak SNP. Thus, for the *APOE* locus rs1160985 was

genotyped instead of rs4420638 ($P_D = 2.67 \times 10^{-16}$ for TC, $P_D = 9.07 \times 10^{-9}$ for HDL,

and $P_D = 7.44 \times 10^{-10}$ for TG); for the *APOA1* locus rs964184 was genotyped instead of

rs3741298 ($P_D = 1.63 \times 10^{-7}$ for TC) or rs6589566 ($P_D = 3.98 \times 10^{-18}$ for TG); for the

MGAT1 locus rs1038143 was genotyped instead of rs249356 ($P_D = 1.06 \times 10^{-6}$ for

HDL); for the *APOB* locus rs754523 was genotyped instead of rs1469513 ($P_D = 2.71 \times$

10^{-6} for LDL).

We successfully genotyped the peak SNP, or a proxy SNP, in the replication

cohorts for 15 loci. Two loci (*APOA1* with TG, $P_{DR} = 1.81 \times 10^{-29}$; *APOE* with TC, $P_{DR} =$

4.29×10^{-21}, and LDL, $P_{DR} = 1.53 \times 10^{-27}$) demonstrated genome-wide significant

associations in the discovery-replication meta-analysis (Table 2 and Tables S1, S7,

S10, and S13 in S1 Appendix). An additional four associations demonstrated evidence

of replication with consistent directions of effect and suggestive joint P_{DR} values (*GCKR*

with TG, $P_{DR} = 5.62 \times 10^{-8}$; *MGAT1* with HDL, $P_{DR} = 2.91 \times 10^{-7}$; *APOA1* with TC, $P_{DR} =$

1.72×10^{-6}; *RAB21* with HDL, $P_{DR} = 5.92 \times 10^{-7}$). Three associations had suggestive

joint P_{DR} values driven by the discovery associations only (*APOB* with LDL, $P_{DR} = 5.81 \times$

10^{-6}; *LIPC* with HDL, $P_{DR} = 9.15 \times 10^{-7}$; *CDH4* with HDL, $P_{DR} = 8.77 \times 10^{-6}$);

associations at *APOB* and *CDH4* had consistent directions of effect. Among the

remaining loci with at least suggestive association in the discovery sample, but not in

the discovery-replication meta-analysis, consistent effect directions were also seen for

TC and *APOB* and *ZHX2*; LDL and *ALG10* and *CPNE8* (Table 2 and Tables S1, S7,

S10, and S13 in S1 Appendix).

Table 2. Suggestive loci and replication genotyping

Total Cholesterol

Locus	SNP	Chr	BP	EA	OA	Dir	P_D	P_R	P_{DR}	Known gene	Traits	SAM	EAS	SAS	EUR	AMR	AFR
APOB	rs754523	2	21311691	G	A	+++	*6.25E-06*	0.178	1.20E-05	*APOB*	C,H,L,T	0.247	0.265	0.140	0.309	0.306	0.201
PDE4D	rs7711093	5	59593138	G	A	++−	*3.01E-06*	0.747	5.37E-04			0.506	0.643	0.553	0.856	0.775	0.852
LUCAT1	rs10072084	5	90539203	C	T	+··	*9.48E-06*					0.541	0.559	0.422	0.232	0.429	0.822
*FILIP1***	rs2951921	6	76165524	T	C	+··	*9.04E-07*					0.073	0.022	0.063	0.015	0.030	0.293
ZHX2	rs7841763	8	123971081	T	C	+++	*4.82E-06*	0.631	1.03E-04			0.043	0.023	0.146	0.102	0.058	0.169
APOA1	rs964184*	11	116648917	C	G	+++	5.37E-05	0.009	*1.72E-06*	*APOA1*	C,H,L,T	0.560	0.760	0.771	0.838	0.723	0.779
SIRT2	rs10405150	19	39387919	C	T	+··	*6.34E-06*					0.056	0.147	0.144	0.081	0.117	0.774
ZNF283	rs16976816	19	44339377	G	A	+··	*9.78E-06*					0.977	0.970	0.994	0.987	0.976	0.864
APOE	rs1160985*	19	45403412	C	T	+++	**2.13E-13**	**3.36E-09**	**4.29E-21**	*APOE*	C,H,L,T	0.724	0.659	0.590	0.554	0.447	0.378

HDL

Locus	SNP	Chr	BP	EA	OA	Dir	P_D	P_R	P_{DR}	Known gene	Traits	SAM	EAS	SAS	EUR	AMR	AFR
STON1-GTF2A1L	rs6739536	2	48831901	A	G	−··	*1.58E-06*					0.762	0.676	0.837	0.918	0.842	0.609
MGAT1	rs1038143*	5	180213878	T	C	−−−	*3.72E-06*	0.016	*2.91E-07*			0.309	0.110	0.137	0.148	0.081	0.013
AKAP7	rs3777486	6	131584648	G	A	−+−	*3.09E-06*	0.808	4.37E-04			0.976	0.909	0.898	0.858	0.793	0.949
CSMD1	rs1626142	8	4345284	T	C	−··	*7.67E-06*					0.612	0.400	0.435	0.298	0.392	0.654
RAB21	rs328733	12	72197574	T	C	−−−	*2.57E-06*	0.036	*5.92E-07*			0.788	0.626	0.700	0.869	0.738	0.611
ZNF10	rs2292029	12	133734113	A	G	−··	*4.05E-06*					0.179	0.112	0.254	0.245	0.156	0.008
HS6ST3	rs16953620	13	97508453	A	G	−··	*8.48E-06*					0.968	0.950	0.949	0.848	0.911	0.836
LIPC	rs10438284	15	58629424	G	A	−+−	*4.00E-07*	0.133	*9.15E-07*	*LIPC*	C,H,T	0.286	0.326	0.173	0.272	0.199	0.047
CETP	rs289708	16	57038162	T	C	−··	**1.19E-11**			*CETP*	C,H,L,T	0.905	0.815	0.800	0.860	0.823	0.597
LIPG	rs16950739	18	47138509	C	T	−··	*1.07E-07*			*LIPG*	C,H	0.019	0.058	0.190	0.057	0.140	0.004
APOE	rs1160985*	19	45403412	C	T	−+−	0.003	0.342	0.004	*APOE*	C,H,L,T	0.724	0.659	0.590	0.554	0.447	0.378
CDH4	rs817687	20	59753355	A	G	−−−	*2.31E-06*	0.237	*8.77E-06*			0.986	0.924	0.945	0.967	0.855	0.610

LDL

Locus	SNP	Chr	BP	EA	OA	Dir	P_D	P_R	P_{DR}	Known gene	Traits	SAM	EAS	SAS	EUR	AMR	AFR
APOB	rs754523*	2	21311691	G	A	+++	*3.25E-06*	0.158	*5.81E-06*	*APOB*	C,H,L,T	0.247	0.265	0.140	0.309	0.306	0.201
KALRN	rs6789134	3	123942339	G	A	++−	*3.22E-06*	0.925	1.96E-04			0.078	0.161	0.120	0.046	0.118	0.212
ZHX2	rs7841763	8	123971081	T	C	++−	*1.80E-06*	0.557	3.78E-05			0.043	0.023	0.146	0.102	0.058	0.169
SH2D4B	rs10509415	10	82473065	A	C	+··	*7.96E-06*					0.706	0.510	0.735	0.758	0.808	0.708
ALG10	rs3912355	12	34079616	C	T	+++	*2.12E-06*	0.544	3.97E-05			0.855	0.872	0.651	0.605	0.732	0.863
ALG10B	rs10880642	12	38554152	A	G	+··	*5.56E-06*					0.802	0.715	0.555	0.481	0.532	0.299
CPNE8	rs11169807	12	39244161	C	T	+++	*4.77E-06*	0.418	4.13E-05			0.794	0.644	0.536	0.505	0.408	0.898
LINC02408	rs17104016	12	67969929	A	T	+··	*9.29E-06*					0.574	0.734	0.924	0.853	0.906	0.838
LINC00922	rs254371	16	65943650	T	C	+··	*9.04E-06*					0.555	0.698	0.650	0.599	0.693	0.884
ZNF283	rs16976816	19	44339377	G	A	+··	*1.78E-06*					0.977	0.970	0.994	0.987	0.976	0.864
APOE	rs1160985	19	45403412	C	T	+++	**2.61E-20**	**5.07E-09**	**1.53E-27**	*APOE*	C,H,L,T	0.724	0.659	0.590	0.554	0.447	0.378

Triglycerides

Locus	SNP	Chr	BP	EA	OA	Dir	P_D	P_R	P_{DR}	Known gene	Traits	SAM	EAS	SAS	EUR	AMR	AFR
GCKR	rs780094	2	27741237	T	C	+++	9.84E-07	0.01	5.62E-08	GCKR	C,T	0.334	0.476	0.198	0.411	0.360	0.132
CD200	rs2399416	3	112059213	A	G	++−	5.12E-06	0.668	9.29E-04			0.021	0.148	0.140	0.393	0.272	0.101
SPIN1	rs7861888	9	90886340	A	G	+··	4.24E-06					0.706	0.719	0.897	0.921	0.842	0.989
APOA1	rs964184*	11	116648917	C	G	+++	2.37E-17	8.97E-14	1.81E-29	APOA1	C,H,L,T	0.440	0.240	0.229	0.162	0.277	0.221
KIRREL3	rs3018434	11	126805881	G	A	+··	4.16E-06					0.916	0.803	0.920	0.866	0.857	0.974
APOE	rs1160985*	19	45403412	T	C	+−+	0.312	0.836	0.507	APOE	C,H,L,T	0.276	0.341	0.410	0.446	0.553	0.623

EA = effect allele

OA = other allele (reference allele)

Dir = direction of the effect in each of the four samples (+ indicates the effect allele is increasing the trait value on the raw scale)

P_D = discovery cohort GWAS *p*-value

P_R = replication cohort *p*-value

P_{DR} = joint discovery and replication cohorts *p*-value

Known gene = Loci observed in Teslovich et al. 2010 or Willer et al. 2013

Traits = Traits locus associated with in Teslovich et al. 2010 or Willer et al. 2013

C = TC

H = HDL

L = LDL

T = TG

SAM = Samoan effect allele frequency (EAF)

EAS = East Asian EAF

SAS = South Asian EAF

EUR = European EAF

AMR = Admixed American EAF

AFR = African EAF

*The SNP genotyped in the replication population was not the peak SNP at this locus for this trait.

We compared the directions of effect to those previously reported in Willer et al.

[8] and Teslovich et al. [30] for *APOB*, *GCKR*, *APOA1*, *LIPC*, *LIPG*, and *APOE* (i.e.,

genome-wide suggestive loci that have been previously associated with lipid traits). We

observed a consistent direction of effect for the representative SNP for all associations

except for *LIPG* and HDL (Table 2).

The effect allele frequencies in the two samples—discovery and replication—

were largely similar for each of the 15 successfully genotyped SNPs (Tables S1, S7,

S10, and S13 Appendix). However, many loci had markedly different effect allele

frequencies between Samoans and other 1000 Genomes populations (Table 2). For

example, compared to 1000 Genomes populations, there were higher effect allele

frequencies (EAFs) in Samoans for rs964184 near *APOA1* (G allele frequency: 0.440 in

Samoans vs. < 0.277 in 1000 Genomes populations), rs1160985 near *APOE* (C allele

frequency: 0.724 in Samoans vs. < 0.659 in 1000 Genomes populations), and

rs1038143 near *MGAT1* (A allele frequency: 0.309 in Samoans vs. < 0.148 in 1000

Genomes populations).

Discussion

In this study, we examined four measures of fasting lipid levels—TC, HDL, LDL

and TG—for associations with 659,492 SNPs from a genome-wide array in a discovery

cohort of 2,849 Samoans, with follow-up genotyping of significant and suggestive

findings in a replication cohort comprising 1,798 Samoans from Samoa and American

Samoa. Thirty-one loci had at least suggestive evidence of association with one or more

lipid traits in the discovery cohort, of which eight have been reported to be associated

with lipid levels previously: *APOB, GCKR, MGAT1, APOA1*, *LIPC*, *CETP, LIPG,* and

APOE [8, 10, 30] although the direction of effect for the variant near LIPG was in the

opposite direction from previous results. Enrichment analyses highlighted known lipid

metabolism gene sets and previously associated lipid loci.

We observed a difference in the architecture of the statistical association signals

between the four lipid traits and variants near *APOE* (Fig 2E-H). The peak SNP for TC

and LDL was rs1160985, an intronic variant in *TOMM40* upstream of *APOE*; whereas

the peak SNP for HDL and TG was rs4420638, an intergenic variant downstream of

APOC1 and *APOE*. rs1160985 demonstrated evidence of replication for TC and LDL

but not for HDL and TG, consistent with the discovery findings. These markers are in

low linkage disequilibrium with each other (r^2 = 0.093) and may represent distinct

association signals. While this could support a shared genetic architecture for TC and

LDL and for HDL and TG in Polynesians, this study was not positioned to adequately

capture the association signal present at this locus. Future studies with sequencing or

imputation of the 19q13.2 region will be necessary to dissect the genetic architecture of

APOE and lipid levels in Polynesians.

We did not observe suggestive or genome-wide significant association with

several loci which have figured prominently in multiple lipid GWAS (e.g., *LPL*, *LDLR*,

CILP2, FADS1/2/3, ANGPTL3, SORT1, PPP1R3B, MIXIPL, HNF4A, PCSK9, GALNT2,

HMGCR) either because we lacked sufficient power to detect their effects, the effects

are negligible in Samoans, or the allele frequencies of associated variants are different

enough in Samoans to hinder detection. However, it is important to note that this study

was not designed to evaluate the effect of known lipids loci in Samoans, nor were

previously-associated loci examined specifically.

We detected and replicated a suggestive association between HDL and a variant

on 5q35.3 (Fig 2A). While the peak SNP lies within an intron of *BTNL8*, the variant

selected for follow-up genotyping is intergenic, downstream of *MGAT1*. A suggestive

association between variants near *MGAT1* and HDL in a GWAS in the Micronesian

population of Kosrae has been previously reported [10]. Although there is no evidence

of association between *MGAT1* and lipids as reported in prior studies of non-Pacific

Islanders, variation near *MGAT1* has been associated with BMI, serum fatty acid levels

and composition, and glucose response in Europeans [37–39]. The encoded MGAT

enzyme plays a major role in the absorption of dietary fat in the intestine [40]. Due to the

greater frequency of the HDL-associated risk variant observed in Samoans compared to

1000 Genomes populations, it is plausible that variation near *MGAT1* may have a

unique role in or a stronger effect on the lipid metabolism of Pacific Islanders.

We also detected and replicated a novel suggestive association between HDL

and a variant downstream of *RAB21* (Fig 2D). Unlike the variant downstream of

MGAT1, we observed similar allele frequencies between Samoans and 1000 Genomes

populations (i.e., < 20% difference between Samoans and another 1000 Genomes

population) in the HDL-associated variant downstream of *RAB21*. This region, 12q21,

was previously seen in linkage analysis with both univariate and bivariate scans of TC

and LDL [12]. The individuals included in this linkage analysis are also included in our

2002–03 replication sample set, however, they do not appear to be driving the

association signal near *RAB21* (Table S7 in S1 Appendix). Variation near *RAB21* has

been previously associated with obesity [41]. *RAB21* belongs to the family of

monomeric GTPases involved in control of cellular membrane trafficking and is involved

in the targeted trafficking of integrins and the regulation of cell adhesion and migration

[42, 43].

This study is limited in drawing conclusions about the genetic architecture of

lipids in Samoans, as replication genotyping was unavailable for many loci and due to

the lack of genome-wide imputation. Future studies, evaluating the evidence of

association between associations seen here in separate cohorts as well fine-mapping

loci with genotype imputation (given the availability of a relevant reference panel), are

necessary to fully evaluate the genetic architecture of lipids in Samoans.

This is the first GWAS of lipid phenotypes in Samoans, and we observed

association with many known lipid loci, which was further supported by the gene-set

enrichment analysis highlighting lipid metabolism gene sets. However, the difference in

association results near *APOE*, coupled with evidence of Pacific-Islander–specific

associations with *MGAT1* and *RAB21* suggest that some, but not all, of the genetic

architecture of lipids is shared between Samoans and other populations. Given this

evidence of a partially distinct genetic architecture of lipids in Samoans, further

investigation and fine-mapping of lipid loci, especially that across multiple ethnicities, is

warranted.

Acknowledgements

The authors would like to thank the Samoan participants of the study, local

village authorities, and the many Samoan and other field workers over the years. We

acknowledge the Samoan Ministry of Health, the Samoa Bureau of Statistics, and the

American Samoan Department of Health for their support of this research. We give

particular thanks to two research assistants, Melania Selu and Vaimoana Lupematasila,

who contributed to the 2010 recruitment and continue to assist us in our work in Samoa.

References

1. Hawley NL, Minster RL, Weeks DE, Viali S, Reupena MS, Sun G, Cheng H, Deka R, Mcgarvey ST (2014) Prevalence of adiposity and associated cardiometabolic risk factors in the samoan genome-wide association study. Am J Hum Biol 26:491–501

2. Keighley E., McGarvey ST, Quested C, Mccuddin C, Viali S, Maga UA (2007) Nutrition and health in modernizing Samoans: temporal trends and adaptive perspectives. In: Ohtsuka R, Ulijaszek SJ (eds) Heal. Chang. Asia-Pacific Reg. Biocultural Epidemiol. Approaches. Cambridge University Press, Cambridge, NY, pp 147–191

3. McGarvey ST (2001) Cardiovascular disease (CVD) risk factors in Samoa and American Samoa, 1990-95. Pac Health Dialog 8:157–62

4. Kannel WB, Castelli WP, Gordon T, McNamara PM (1971) Serum cholesterol, lipoproteins, and the risk of coronary heart disease. The Framingham study. Ann Intern Med 74:1–12

5. Gordon T, Castelli WP, Hjortland MC, Kannel WB, Dawber TR (1977) High density lipoprotein as a protective factor against coronary heart disease. The Framingham Study. Am J Med 62:707–14

6. Roeters van Lennep J, Westerveld HT, Erkelens DW, van der Wall EE (2002) Risk factors for coronary heart disease: implications of gender. Cardiovasc Res 53:538–549

7. Lange LA, Willer CJ, Rich SS (2015) Recent developments in genome and exome-wide analyses of plasma lipids. Curr Opin Lipidol 26:96–102

8. Willer CJ, Schmidt EM, Sengupta S, et al (2013) Discovery and refinement of loci associated with lipid levels. Nat Genet 45:1274–1283

9. Burkhardt R, Kenny EE, Lowe JK, et al (2008) Common SNPs in HMGCR in micronesians and whites associated with LDL-cholesterol levels affect alternative splicing of exon13. Arterioscler Thromb Vasc Biol 28:2078–2084

10. Lowe JK, Maller JB, Pe'er I, et al (2009) Genome-wide association studies in an isolated founder population from the Pacific Island of Kosrae. PLoS Genet 5:e1000365

11. Dumitrescu L, Carty CL, Taylor K, et al (2011) Genetic determinants of lipid traits in diverse populations from the population architecture using genomics and epidemiology (PAGE) study. PLoS Genet 7:e1002138

12. Åberg K, Dai F, Sun G, et al (2008) A genome-wide linkage scan identifies multiple chromosomal regions influencing serum lipid levels in the population on the Samoan islands. J Lipid Res 49:2169–2178

13. Minster RL, Hawley NL, Su C-T, et al (2016) A thrifty variant in CREBRF strongly influences body mass index in Samoans. Nat Genet 48:1049–1054

14. Kristiansson K, Naukkarinen J, Peltonen L (2008) Isolated populations and complex disease gene identification. Genome Biol 9:109

15. Laurie CC, Doheny KF, Mirel DB, et al (2010) Quality control and quality assurance in genotypic data for genome-wide association studies. Genet Epidemiol 34:591–602

16. Deka R, Mc Garvey ST, Ferrell RE, Kamboh MI, Yu LM, Aston CE, Jin L,
Chakraborty R (1994) Genetic characterization of American and Western
Samoans. Hum Biol 66:805–22
17. McGarvey ST, Levinson PD, Bausser-Man L, Galanis DJ, Hornick CA (1993)
Population change in adult obesity and blood lipids in American Samoa from
1976–1978 to 1990. Am J Hum Biol 5:17–30
18. Chin-Hong P V, McGarvey ST (1996) Lifestyle incongruity and adult blood
pressure in Western Samoa. Psychosom Med 58:131–7
19. Galanis DJ, McGarvey ST, Quested C, Sio B, Afele-Fa'amuli SA (1999) Dietary
intake of modernizing Samoans: implications for risk of cardiovascular disease. J
Am Diet Assoc 99:184–90
20. Åberg K, Dai F, Sun G, et al (2009) Susceptibility Loci for Adiposity Phenotypes
on 8p, 9p, and 16q in American Samoa and Samoa. Obesity 17:518–524
21. Dai F, Keighley ED, Sun G, et al (2007) Genome-wide scan for adiposity-related
phenotypes in adults from American Samoa. Int J Obes 31:1832–1842
22. Åberg K, Dai F, Viali S, Tuitele J, Sun G, Indugula SR, Deka R, Weeks DE,
McGarvey ST (2009) Suggestive linkage detected for blood pressure related traits
on 2q and 22q in the population on the Samoan islands. BMC Med Genet 10:107
23. Dai F, Sun G, Åberg K, et al (2008) A Whole Genome Linkage Scan Identifies
Multiple Chromosomal Regions Influencing Adiposity-Related Traits among
Samoans. Ann Hum Genet 72:780–792
24. Chen W-M, Abecasis GR (2007) Family-Based Association Tests for
Genomewide Association Scans. Am J Hum Genet 81:913–926
25. Devlin B, Roeder K (1999) Genomic control for association studies. Biometrics
55:997–1004
26. Purcell S, Cherny SS, Sham PC (2003) Genetic Power Calculator: design of
linkage and association genetic mapping studies of complex traits. Bioinformatics
19:149–150
27. Carithers LJ, Ardlie K, Barcus M, et al (2015) A Novel Approach to High-Quality
Postmortem Tissue Procurement: The GTEx Project. Biopreserv Biobank 13:311–
319
28. Segrè A V, Groop L, Mootha VK, Daly MJ, Altshuler D (2010) Common Inherited
Variation in Mitochondrial Genes Is Not Enriched for Associations with Type 2
Diabetes or Related Glycemic Traits. PLoS Genet 6:e1001058
29. Lee PH, O'Dushlaine C, Thomas B, Purcell SM (2012) INRICH: interval-based
enrichment analysis for genome-wide association studies. Bioinformatics
28:1797–1799
30. Teslovich TM, Musunuru K, Smith A V., et al (2010) Biological, clinical and
population relevance of 95 loci for blood lipids. Nature 466:707–713
31. Aulchenko YS, Ripke S, Isaacs A, van Duijn CM (2007) GenABEL: an R library for
genome-wide association analysis. Bioinformatics 23:1294–1296
32. R Core Team (2014) R: A language and environment for statistical computing.
33. Therneau T, Atkinson E, Sinnwell J, Schaid D, McDonnell S (2014) kinship2:
Pedigree functions.
34. Winkler TW, Day FR, Croteau-Chonka DC, et al (2014) Quality control and
conduct of genome-wide association meta-analyses. Nat Protoc 9:1192–1212

35. Willer CJ, Li Y, Abecasis GR (2010) METAL: fast and efficient meta-analysis of
genomewide association scans. Bioinformatics 26:2190–2191
36. Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, Boehnke M,
Abecasis GR, Willer CJ (2010) LocusZoom: regional visualization of genome-wide
association scan results. Bioinformatics 26:2336–2337
37. Johansson Å, Marroni F, Hayward C, et al (2010) Linkage and Genome-wide
Association Analysis of Obesity-related Phenotypes: Association of Weight With
the MGAT1 Gene. Obesity 18:803–808
38. Jacobsson JA, Rask-Andersen M, Risérus U, et al (2012) Genetic variants near
the MGAT1 gene are associated with body weight, BMI and fatty acid metabolism
among adults and children. Int J Obes 36:119–129
39. Del-Aguila JL, Beitelshees AL, Cooper-DeHoff RM, Chapman AB, Gums JG,
Bailey K, Gong Y, Turner ST, Johnson JA, Boerwinkle E (2014) Genome-wide
association analyses suggest NELL1 influences adverse metabolic response to
HCTZ in African Americans. Pharmacogenomics J 14:35–40
40. Yen C-LE, Stone SJ, Cases S, Zhou P, Farese R V. (2002) Identification of a
gene encoding MGAT1, a monoacylglycerol acyltransferase. Proc Natl Acad Sci
99:8512–8517
41. Turcot V, Lu Y, Highland HM, et al (2018) Protein-altering variants associated with
body mass index implicate pathways that control energy intake and expenditure in
obesity. Nat Genet 50:26–41
42. Pellinen T, Ivaska J (2006) Integrin traffic. J Cell Sci 119:3723–31
43. Pellinen T, Tuomi S, Arjonen A, et al (2008) Integrin trafficking regulated by
Rab21 is necessary for cytokinesis. Dev Cell 15:371–85
572
573

Supporting Information

S1 Appendix. Supporting Information.

Data Availability Statement

579 The discovery cohort data are available from dbGaP (accession number:

580 phs000914.v1.p1).