The atypical thiol-disulfide exchange protein α-DsbA2 from *Wolbachia pipientis* is a homotrimeric disulfide isomerase

Patricia M. Walden[1], Andrew E. Whitten[1#‡], Premkumar Lakshmanane[1†], Maria Halili[1,3], Begoña Heras[1§], Gordon J. King[1] and Jennifer L. Martin[1,3‡]

[1] Institute for Molecular Bioscience, University of Queensland, Brisbane QLD 4072, Australia

[2] School of Biological Sciences, University of Queensland, Brisbane QLD 4072, Australia

[3] Griffith Institute for Drug Discovery, Griffith University, Nathan, QLD 4111, Australia

Current address:

[#] Australian Nuclear Science and Technology Organisation, Sydney NSW 2232, Australia

[†] Department of Microbiology and Immunology, School of Medicine, University of North Carolina, Chapel Hill, North Carolina 27514, USA

[§] Department of Biochemistry, La Trobe University, Melbourne VIC 3086, Australia

[‡]Correspondence

Professor Jennifer L. Martin, Griffith Institute for Drug Discovery, Griffith University, Nathan, QLD 4111 Australia Email: **jlm@griffith.edu.au**

And Dr Andrew E. Whitten, Australian Nuclear Science and Technology Organisation, Sydney NSW 2232, Australia Email: awh@ansto.gov.au

Abstract

DiSulfide Bond (DSB) oxidative folding enzymes are master regulators of virulence

localized to the periplasm of many Gram-negative bacteria. The archetypal DSB

machinery from *Escherichia coli* K12 has a dithiol oxidizing redox relay pair

(DsbA/B), a disulfide isomerizing redox relay pair (DsbC/D) and specialist reducing

enzymes DsbE and DsbG that also interact with DsbD. By contrast the Gram-negative

bacterium *Wolbachia pipientis* encodes just three DSB enzymes. Two of these α-

DsbA1 and α-DsbB form a redox relay pair analogous to *E. coli* DsbA/B. The third

enzyme α-DsbA2 incorporates a DsbA-like sequence but does not interact with α-

DsbB. In comparison with other DsbA enzymes, α-DsbA2 has ~50 extra N-terminal

residues. The crystal structure of α-DsbA2ΔN, the N-terminally truncated form in

which these residues are removed confirms the DsbA-like nature of this domain.

However, α-DsbA2 does not have DsbA-like activity: it is structurally and

functionally different as a consequence of its N-terminal residues. First, α-DsbA2 is a

powerful disulfide isomerase and a poor dithiol oxidase – *ie* its role is to shuffle rather

than introduce disulfide bonds. Moreover, small-angle X-ray scattering of α-DsbA2

reveals a homotrimeric arrangement. Our results allow us to draw conclusions about

the factors required for functionally equivalent enzymatic activity across structurally

diverse protein architectures.

Introduction

Many secreted and outer membrane proteins of prokaryotes rely on disulfide bonds for stability and function (Feige & Hendershot, 2011). The introduction, isomerization and reduction of protein disulfide bonds in bacteria are controlled by disulfide bond forming (DSB) proteins (Landeta *et al.*, 2018, Inaba, 2009). These DSB proteins are considered master regulators of virulence because they are essential for the folding and activity of diverse virulence factors including bacterial toxins, secretion systems, adhesins, flagella *etc* (Heras *et al.*, 2009).

The classic DSB folding machinery characterised in the model bacterium *Escherichia coli* K-12 comprises two independent periplasmic pathways: the (i) oxidative and (ii) isomerization pathways (Inaba, 2009). In the oxidative pathway, a monomeric thioredoxin (TRX)-fold protein *E. coli* DsbA (EcDsbA) donates its Cys-X-X-Cys active site disulfide bond directly to nascent protein substrates (Zapun *et al.*, 1993, Inaba & Ito, 2002). EcDsbA becomes reduced as a consequence of this reaction and its active site is re-oxidized by a specific interaction with its integral membrane protein partner *E. coli* DsbB (EcDsbB) (Bader *et al.*, 1999).

In the classic isomerization pathway, the *E. coli* homodimeric protein disulfide isomerase DsbC (EcDsbC) reduces and shuffles incorrect disulfide bonds in misfolded proteins to generate correctly folded proteins (Shevchik *et al.*, 1994). Each EcDsbC protomer has a catalytic TRX-fold domain with a characteristic Cys-X-X-Cys active site, and an 87-residue N-terminal region that forms a dimerization domain essential for isomerase activity (McCarthy *et al.*, 2000). EcDsbC forms a redox relay with the integral membrane protein *E. coli* DsbD (EcDsbD) that maintains EcDsbC in

its active reduced state (McCarthy *et al.*, 2000). *E. coli* K12 also encodes two

specialist reducing enzymes EcDsbG and EcDsbE (Depuydt *et al.*, 2009) that interact

with EcDsbD (Missiakas *et al.*, 1995).

Here we focus on one of the DSB proteins encoded by *Wolbachia pipientis w*Mel, a

bacterium from the *Rickettsiaceae* family. The *Rickettsiacea*e family are Gram-

negative bacteria of the α-Proteobacteria class that establish obligate intracellular

infections in arthropods.   *W. pipientis* are widespread, found in ~60% of insect

species, and have an extraordinary impact on host biology. Infection results in

phenotypic alterations such as cytoplasmic incompatibility, feminization or reduction

of lifespan – all of which contribute to the bacterium's own survival (Hilgenboecker

et al., 2008). Several *Wolbachia* strains have been shown to block the transmission of

mosquito-borne viruses and are being trialled as biocontrol agents aimed at

eradicating vector-borne diseases such as dengue, Zika and Chikungunya (reviewed in

(Flores & O'Neill, 2018)).

The *W. pipientis w*Mel strain encodes two DsbA-like proteins, α-DsbA1 and α-

DsbA2, and an integral membrane protein α-DsbB (Walden *et al.*, 2013). Unlike *E.*

coli, this strain does not encode obvious homologues of DsbC or DsbD (though all

other *W. pipientis* strains do encode a DsbD homologue). Of the two encoded DsbAs,

α-DsbA1 has been characterized and shown to form a redox relay with α-DsbB

(Walden *et al.*, 2013) that resembles the redox relay between EcDsbA and EcDsbB.

In contrast, α-DsbA2 – which is highly conserved in *Wolbachia* – does not interact

with α-DsbB and it has a long N-terminal region compared with α-DsbA1 (Walden *et*

al., 2013).

The N-terminal region of EcDsbC forms a dimerization domain essential for disulfide

isomerase activity. Two other DsbA-like proteins with N-terminal extensions are

disulfide isomerases and thought to be dimeric (*Legionella* DsbA2 (Kpadeh *et al.*,

2015), *Caulobacter* ScsC (Cho *et al.*, 2012)). In addition, *P. mirabilis* ScsC is DsbA-

like with an N-terminal extension, and has disulfide isomerase activity. However its

N-terminal residues interact to form a homotrimer (Furlong *et al.*, 2017). In each of

these cases, the N-terminal regions are essential for oligomerisation and protein

disulfide isomerase activity. We hypothesised that the N-terminal region of

Wolbachia α-DsbA2 would also impart disulfide isomerase activity by forming an

oligomerisation domain.

Here we report structural and functional studies of *Wolbachia* α-DsbA2. We used two

constructs: (1) FL α-DsbA2, the full-length mature α-DsbA2 (lacking the signal

peptide that directs the protein to the periplasm) comprising residues 15-252 and (2)

α-DsbA2ΔN a truncated form of α-DsbA2 (lacking both the signal sequence and the

50-residue N-terminal region) comprising residues 64-252. Our results show that FL

α-DsbA2 is a strong protein disulfide isomerase and that removal of the N-terminal

residues eliminates this activity but gives rise to weak dithiol oxidase activity. The

crystal structure of truncated α-DsbA2ΔN reveals a classic monomeric DsbA-like

architecture. However, SAXS models of FL α-DsbA2 are consistent with a trimer

forming through the interaction of the N-terminal residues.

Results

The N-terminus of *Wolbachia* α-DsbA2 is predicted to be helical

The 87-residue N-terminal region of the archetypal disulfide isomerase EcDsbC adopts a β-sheet dimerisation domain with a helix that links to the catalytic domain (McCarthy *et al.*, 2000). The presence of a detectable DsbA-like domain and an extended N-terminal region in the *Wolbachia* α-DsbA2 sequence suggested that – like EcDsbC – the 50 residue N-terminal region might act as a dimerisation domain. However, the predicted secondary structure of the α-DsbA2 N-terminal region has no structural relationship with the equivalent region of EcDsbC (**Figure 1A**).

Recently, three other DsbA-like proteins have been shown by gel filtration to be oligomeric as a consequence of their N-terminal residues, and all three are functional disulfide isomerases. These are *Legionella pneumophila* DsbA2 [LpDsbA2, probably dimeric, with a ~50 residue N-terminal region predicted to be helical (Kpadeh *et al.*, 2015)], *Caulobacter crescentus* ScsC [probably dimeric, ~60 residue N-terminal region predicted to be helical (Cho *et al.*, 2012)] and *Proteus mirabilis* ScsC [confirmed trimeric protein, ~60 residue helical N-terminal region (Furlong *et al.*, 2017)]. An alignment of all five proteins (α-DsbA2, LpDsbA2, CcScsC, PmScsC and EcDsbC) is provided in **Figure 1B**.

α-DsbA2 is redox active and has disulfide isomerase activity as a consequence of its N-terminal residues

We investigated the redox properties of FL α-DsbA2 and truncated α-DsbA2ΔN. The redox potential gives important information about a protein's propensity to acquire

electrons from its substrate, and thereby become reduced. We determined the standard

redox potentials of FL α-DsbA2 and α-DsbA2ΔN relative to the redox potential of

glutathione (-240 mV) (**Figure 2A**). From these data, the K_{eq} for FL α-DsbA2 was

calculated to be $2.16 \pm 0.15 \times 10^{-4}$ M corresponding to a redox potential of -131 mV

at pH 7.0. The calculated K_{eq} for α-DsbA2ΔN was a little more oxidizing, 8.71 ± 0.12

$\times 10^{-5}$ corresponding to a redox potential of -122 mV. By comparison, the redox

potential for *Wolbachia* α-DsbA1 is more reducing, $E^0 = -163$ mV (Kurz *et al.*, 2009).

Thus both the FL and truncated α-DsbA2 proteins have redox potentials that are

similar to monomeric EcDsbA [$E^0 = -122$ mV (Mossner *et al.*, 1998)] and dimeric

EcDsbC [($E^0 = -129$ mV (Zapun *et al.*, 1995)].

DSB enzymes are active to varying degrees in the standard disulfide reductase assay.

We found that half of the insulin in solution was reduced by FL α-DsbA2 (which is in

the reduced form due to DTT in the solution) after ~15 min (**Figure 2B**). By

comparison, EcDsbC reduced half the insulin within ~6 min and the oxidase EcDsbA

(which is in the reduced form due to DTT in the solution), reduced half of the insulin

after ~35 min. The activity of α-DsbA2ΔN in this assay was negligible, and

comparable to the negative control. These data show that FL α-DsbA2 is redox-active

and the N-terminal residues are critical for disulfide reductase activity.

We next assessed the protein disulfide isomerase activity of FL α-DsbA2 and α-

DsbA2ΔN by following the reactivation of scrambled RNase (scRNase) in an *in vitro*

assay. FL α-DsbA2 recovered 73% ± 3% of RNase activity after ~5 hours, compared

to the positive control EcDsbC which recovered 85% ± 5% of RNase activity over the

same time period, though reached that level after 100 min and then plateaued (**Figure**

2C). By comparison, truncated α-DsbA2ΔN recovered just 35% ± 3% of RNase

activity relative to native refolded RNase over that period, similar to that reported for

the oxidase enzyme EcDsbA (Shouldice *et al.*, 2011). Therefore, we conclude that α-

DsbA2 is a protein disulfide isomerase and that its disulfide isomerase activity

requires the presence of the N-terminal residues.

We also investigated whether FL α-DsbA2 or α-DsbA2ΔN demonstrated dithiol

oxidase activity, by measuring their ability to catalyse disulfide bond formation in an

in vitro assay. Specifically, we measured the ability of FL α-DsbA2 or α-DsbA2ΔN to

oxidize the cysteines of a model peptide in the presence of oxidised glutathione

(GSSG). Compared to EcDsbA–mediated peptide oxidation, FL α-DsbA2

demonstrated negligible peptide oxidation (comparable to GSSG control) whereas α-

DsbA2ΔN had weak activity (**Figure 2D**) under the experimental conditions of this

assay.

The crystal structure of α-DsbA2ΔN reveals a canonical DsbA architecture

We were unable to generate crystals of the full-length mature α-DsbA2. However,

crystals of truncated α-DsbA2ΔN did grow and the crystal structure was solved by

molecular replacement to a resolution of 2.25 Å (**Table 1, Figure 3A**).

The final refined structure of α-DsbA2ΔN has four molecules in the asymmetric unit

(chains A, B, C, D) (**Figure 3B**), each of which features the canonical DsbA

architecture of a TRX-fold with an inserted 4-helical domain and a connecting helix.

As in other DsbA-like crystal structures the Cys-Gly-His-Cys active site is located at

the N-terminal end of helix α3 in the TRX domain (present as a mixture of reduced

and oxidized forms, **Figure 3A, dashed line box**).

Using this crystal structure as a probe, the highest-scoring DALI match (1 March

2018) was the trimeric protein PmScsC (5IDR, molecule A) with Z score 25.8,

RMSD 1.5 Å for 178 C_α-atoms, and 24% sequence identity. After PmScsC, the next

highest DALI hit was an uncharacterised monomeric DsbA-like protein from

Silicibacter pomeroyi (from the family *Rhodobacteraceae*) (3GYK molecule A) Z

score 24.8 RMSD 1.5 Å for 169 C_α-atoms and 30% sequence identity. The third

highest hit was the monomeric protein SeScsC (4GXZ, molecule C) Z score 24.1

RMSD 1.6 Å for 166 C_α-atoms, and 25% sequence identity (this was used as the

molecular replacement model to solve the α-DsbA2ΔN crystal structure).

By comparison, superimposition of the α-DsbA2ΔN structure onto the archetypal

disulfide isomerase EcDsbC (1EEJ, molecule A) using DALI gave a Z score of 13.5

and RMSD of 3.3 Å for 147 C_α-atoms (Maiti *et al.*, 2004). These data indicate that

these two disulfide isomerases are structurally very different.

Small angle X-ray scattering shows α-DsbA2 is a homotrimer as a consequence

of its N-terminal region

Although we were unable to crystallise the full-length protein, we were able to obtain

low-resolution structural information using small-angle X-ray scattering (SAXS) data

from both FL α-DsbA2 and α-DsbA2ΔN (**Table 2**). Guinier plots (**Figure 4A, inset**)

reveal a linear trend, consistent with both samples being monodisperse.

Bacterial disulfide isomerases characterised to date are reported to be either dimeric

(EcDsbC, CcScsC, LpDsbA2) or trimeric (PmScsC). We were therefore interested to

determine the molecular mass of the full-length protein and identify whether it is

dimeric or trimeric. Using SAXS data, the molecular mass of truncated α-DsbA2ΔN

estimated to be ~23 kDa from $I(0)$ (Orthaber *et al.*, 2000) and the Porod volume

(Fischer *et al.*, 2010), which is very close to the expected mass for an α-DsbA2ΔN

monomer (21 kDa) and consistent with the crystal structure we report here. However,

the molecular mass of FL α-DsbA2 estimated from $I(0)$ and the Porod volume (75

kDa and 87 kDa respectively) are consistent with the mass of a homotrimer (81 kDa)

rather than a homodimer (54 kDa).

The $p(r)$ for FL α-DsbA2 shows a single peak with a maximum dimension of ~87 Å

(**Figure 4B**) whereas α-DsbA2ΔN demonstrates a single peak with a significantly

smaller maximum dimension of 63 Å. The larger dimension and shifting of the

position of the peak in the $p(r)$ for FL α-DsbA2 are consistent with the formation of a

higher-order oligomer. While these data indicate the FL α-DsbA2 oligomer is

trimeric, this homotrimer differs from that of PmScsC (Furlong *et al.*, 2017), which

has a bimodal pair-distance distribution function (**Figure 4B, dotted line**).

Both dummy atom and rigid body modelling were used to determine low-resolution

solution structures of FL α-DsbA2 and α-DsbA2ΔN. The averaged and filtered

dummy-atom model of α-DsbA2ΔN shows very good agreement with the α-

DsbA2ΔN rigid-body model (composed of the crystal structure plus missing residues

at the N- and C-termini, **Figure 4C**). A comparison of the scattering data with the

rigid-body model scattering profile (**Figure 4A**, red curve and black line) shows good

correspondence, though with a small systematic difference between the two curves

that could be indicative of a low level of sample impurity or possibly a small

difference between the solution and crystal structures.

The averaged and filtered dummy-atom model of FL α-DsbA2 reveals a disk-like

structure with a small protrusion at the centre. Alignment of this model with the FL α-

DsbA2 rigid-body model (**Figure 4D**) shows excellent correspondence; the N-

terminal oligomerisation domain of the rigid-body model coincides with the

protrusion in the dummy-atom model, and the catalytic domains are positioned

around the main disk. The scattering data for FL α-DsbA2 shows excellent

correspondence with the rigid-body model scattering profile (**Figure 4A**, blue curve

and black line).

The two homotrimeric disulfide isomerase enzymes *Wolbachia* FL α-DsbA2 and

PmScsC nevertheless have distinct solution structures. FL α-DsbA2 is disc-like with

the three protomers tightly arranged into a compact shape (**Figure 5A, B**), whereas

PmScsC has a more open arrangement (**Figure 5C, D)** (Furlong *et al.*, 2017).

Discussion

The *w*Mel strain of *W. pipientis* encodes two DsbA-like proteins. One of these, α-DsbA1 has been shown previously to be functionally similar to EcDsbA – it catalyses disulfide formation and forms a redox pair with a membrane protein partner α-DsbB (Walden *et al.*, 2013). Despite sequence similarity, the second protein α-DsbA2, is not DsbA-like. It does not catalyse disulfide formation in the standard assay (this work) and it does not interact with α-DsbB (Walden *et al.*, 2013).

The unusually long N-terminus of *W. pipientis* *w*Mel α-DsbA2 is conserved in DsbA2s encoded in bacterial species from the family *Rickettsiaceae*, class α-Proteobacteria. This includes other members from the same class including *Ehrlichia* and *Anaplasma* that live in hosts such as ticks and cause diseases in animals and humans (Wormser *et al.*, 2006). These organisms all encode α-DsbA1 and α-DsbA2 enzymes similar to those encoded in *Wolbachia*. The high degree of conservation of the N-terminal residues in α-DsbA2 suggests that this region has an important function. Here we showed that these N-terminal residues confer trimerisation and disulfide isomerase properties to *W. pipientis* *w*Mel α-DsbA2.

The enzymatic profile of *W. pipientis* *w*Mel α-DsbA2 overlaps with that of EcDsbC. Indeed, it may replace EcDsbC functionally since *Wolbachia* strains do not encode a DsbC. Most *Wolbachia* strains encode an EcDsbD-like protein that could act as a redox partner for α-DsbA2. Curiously, a DsbD homologue is not present in the genome of the specific *w*Mel *Wolbachia* strain we investigated. We cannot explain why this might be or what other protein might serve as a reducing partner in this organism.

Other organisms encode a disulfide isomerase like that of *Wolbachia* α-DsbA2 rather

than EcDsbC. For example, *Legionella pneumophilia* does not contain a DsbC

homologue (Kpadeh *et al.*, 2013, Kpadeh *et al.*, 2015) though like *Wolbachia* it has

two DsbA-like proteins, one of which, LpDsbA2, has disulfide isomerase activity.

LpDsbA2 has a predicted helical N-terminal extension (**Figure 1B**), and is essential

for assembly of the Type 4b Dot/Icm secretion system (Kpadeh *et al.*, 2013, Kpadeh

et al., 2015).

We have shown that *W. pipientis w*Mel α-DsbA2 is homotrimeric and has disulfide

isomerase activity. Although this is the second example of a homotrimeric TRX-fold

disulfide isomerase – the other being PmScsC (Furlong *et al.*, 2017) - these two

enzymes are quite distinct. First, α-DsbA2 is structurally different from PmScsC in

solution. PmScsC (Furlong *et al.*, 2017) exhibits a bimodal pair-distance distribution

function whereas FL α-DsbA2 has a single peak indicating a more globular and

compact shape. Both dummy atom and rigid body modelling reveal that FL α-DsbA2

is disc-like with no evidence of the flexibility observed for PmScsC (Furlong *et al.*,

2017). Moreover, PmScsC is part of a highly conserved 4-gene *scs* cluster associated

with bacterial copper resistance, including its redox partner PmScsB (Furlong *et al.*,

2018) whereas α-DsbA2 is not part of a gene cluster. Finally, PmScsC is encoded in

organisms that also encode DsbC-like enzymes, whereas α-DsbA2 is not. Presumably,

PmScsC may play a specific role – perhaps it has a specific substrate associated with

copper sensitivity – whereas α-DsbA2 may not.

Although these three structurally characterised disulfide isomerases - EcDsbC,

PmScsC and α-DsbA2 - differ considerably in their structures, we can draw some

broad conclusions about the factors that contribute to their functionally equivalent

enzymatic activity. The present work supports the notion that strong disulfide

isomerase activity requires the presence of at least two catalytic domains in the

enzyme. The way these domains are brought together can vary (dimer/trimer;

sheet/helix) and there is some limited variation in the catalytic active site motif:

CGYC in DsbC and PmScsC; CPYC in CcScsC; CIHC in LpDsbA2; and CGHC in

Wolbachia α-DsbA2. Thus, Gly or Pro predominate in the Cys+1 position, and Tyr or

His predominate in the Cys+2 position. However, this catalytic motif sequence

overlaps with that of monomeric dithiol oxidase DsbAs (eg EcDsbA CPHC). The

second motif that is highly conserved in TRX-like proteins is the *cis*-Pro motif.

Surprisingly, in all five of these disulfide isomerase enzymes the sequence motif is

the same GT*c*P. Monomeric DsbA-like oxidases tend to have more variation, with the

Gly being highly variable and the Thr often replaced with Val. The most telling

sequence feature that discriminates between oxidase and isomerase activity seems to

be the addition of 50 or more residues at the N-terminus of the TRX fold that can

form an oligomerisation domain.

Acknowledgements

Structural data reported in this paper were measured on the MX2 and SAXS/WAXS beamlines at the Australian Synchrotron. We thank the beamline staff for their support. We also thank the University of Queensland Remote Operation Crystallization and X-ray (UQ ROCX) Diffraction Facility for access to crystallization facilities. Additionally, we acknowledge Dr Lachlan Casey for SAXS data measurement, Dr Iñaki Iturbe-Ormaetxe for providing *Wolbachia w*Mel DNA and Dr Mareike Kurz for cloning α-DsbA2 into the original expression vector. This work was supported by an Australian International Postgraduate Research Scholarship to PW, and an ARC Australian Laureate Fellowship (FL0992138) to JLM.

Materials and Methods

Sequence analysis of α-DsbA2

Primary and secondary structure information was obtained from ExPASy (Swiss Institute of Bioinformatics, Switzerland) using the proteomics server ProtParam (Bairoch *et al.*, 2005) and the server PredictProtein (Rost *et al.*, 2004). Sequences of α-DsbA2, EcDsbC, PmScsC, CcScsC and LpDsbA2 were aligned in ClustalOmega (Thompson *et al.*, 1994). Accession codes: EcDsbC P0AEG6; PmScsC B4EV21; CcScsC Q9A747; α-DsbA2 Q73FL6; LpDsbA2 Q5WVK9.

Protein expression and purification

FL α-DsbA2 (locus WD1312; GenBank #AE017196) was amplified from *w*Mel genomic DNA by PCR using primers Forward – 5' TAC TTC CAA TCC AAT GCG

ATG AGC TTG CCG ATA ATA 3' and Reverse – 5' TTA TCC ACT TCC AAT G

CT AGC CTT GCT TGT GAC TTA A 3' which incorporate overhangs for ligation

independent cloning. Truncated α-DsbA2ΔN was amplified using the primers

Forward – 5' TAC TTC CAA TCC AAT GCG GCT CGA GAT AAT GTA ACC 3'

and Reverse – 5' TTA TCC ACT TCC AAT GCT AGC CTT GCT TGT GAC TTA

A 3'. The full-length mature protein without the signal peptide and α-DsbA2ΔN were

cloned into the LIC vector pMCSG7 that incorporates a HIS_6-tag, a linker region

containing 8 amino acids, and a TEV-cleavage site at the N-terminus of the inserted

gene. The construct was transformed into the expression strain BL21(DE3)pLysS

(Life Technologies, USA) to enable over-expression of α-DsbA2 using auto induction

in ZYP-5052 media at 30°C (Studier, 2005). Cells were collected using an Avanti J-

25I centrifuge (Beckman Coulter, Australia) at 12,000 × g at 4°C for 10 min and

frozen at -80°C. α-DsbA2 variants were expressed and purified as described in (Kurz

et al., 2009) with minor variation in the lysis, wash and elution buffer. The lysis

buffer contains 25 mM Tris-HCl pH 7.5 and 150 mM NaCl plus 25 mM imidazole for

the washing buffer, and 250 mM imidazole was included for protein elution.

Protein disulfide reductase assay

The ability of FL α-DsbA2 and α-DsbA2ΔN to catalyse the reduction of insulin in the

presence of DTT was measured *in vitro* (Holmgren, 1979). Insulin comprises two

chains, A and B, which are linked via two disulfide bonds. Upon reduction of the

disulfide bonds by a high reductase-active catalyst, chain B becomes insoluble and

precipitates. FL α-DsbA2, α-DsbA2ΔN, EcDsbC (positive control) or EcDsbA

(negative control) were diluted to a final concentration of 10 μM in buffer containing

100 mM $NaPO_4$ pH 7.0, 1 mM EDTA and 0.33 mM DTT. Insulin (at a final

concentration of 0.13 mM) was added to the cuvette immediately before

measurements were taken, and the extent of insulin reduction was monitored by

measuring the optical density at 650 nm for 50 min. Three experimental replicates

were measured using the same batch of protein and the presented data [mean ±

standard deviation error (SD)] is from three measurements.

Redox potential measurement

2 µM of oxidized FL α-DsbA2 and α-DsbA2ΔN was incubated in fully degassed

buffers containing 100 mM $NaPO_4$ pH 7.0, 1 mM EDTA and 1 mM glutathione

oxidized (GSSG) (Sigma Aldrich, USA) and different concentrations of reduced

glutathione (GSH) (20 µM – 5 mM) for 24 h at RT. After incubation, the reactions

were stopped with 10% trichloroacetic acid (TCA) (Sigma Aldrich, USA) and the

precipitated protein pellets were collected by centrifugation at 16000 x g for 10 min at

4 °C. The pellets were washed with cold acetone and dissolved in buffer containing

50 mM Tris-HCl pH 7.0, 1% SDS and 4 mM 4-acetamide-4'-maleimidylstilbene-2-

2'-disulfonate (AMS) (Molecular Probes, USA). The reduced and oxidized forms

were separated on a 12% SDS Bis-Tris PAGE (Invitrogen, Australia). The fraction of

reduced protein, R, was determined from a scanned image of the stained gel using

ImageJ (Abramoff, 2004). The equilibrium constant K_{eq} was calculated via $R =$

$([GSH]^2/[GSSH])/(K_{eq} + ([GSH]^2/[GSSH]))$, and the redox potential was calculated

via the Nernst equation $E^0 = E^0_{GSH/GSSG} - (RT/nF) \times \ln K_{eq}$, where

$E^0_{GSH/GSSG}$ = -240 mV 240 mV, R = 8.314 J $K^{-1}mol^{-1}$, T = 298 K, n = 2 and F = 9.649

$\times 10^4$ C mol^{-1} (Kurz *et al.*, 2009). Three experimental replicates were measured using

the different batch of protein and data presented are the mean ± SD from duplicate

measurements.

Dithiol oxidase activity assay

Assays were run on a Synergy H1 Multimode plate reader (BioTek, USA) with

excitation at 340nm and emission at 615nm. For time-resolved fluorescence, a 100

μsec delay before reading and 200 msec reading time was employed. The assay was

performed in a white 384-well plate (Perkin Elmer). A 25 μL solution containing 3.2

μM EcDsbA and 2 mM GSSG (positive control), or 3 μM FL α-DsbA2 or α-

DsbA2ΔN and 2 mM GSSG in 50 mM MES, 50 mM NaCl, 2 mM EDTA pH 5.5 was

added to the wells. The assay was initiated by the addition of 25 μL 16 μM peptide (in

50 mM MES, 50 mM NaCl, 2 mM EDTA, pH 5.5) to each well. Measurements were

carried out in triplicate using 3 different protein batches and data shown are the mean

± SD error from these three measurements.

Protein disulfide isomerase assay

The scrambled RNaseA assay was used to detect isomerase activity of FL α-DsbA2,

α-DsbA2ΔN or EcDsbC by monitoring the refolding of scrambled RNaseA (Hillson

et al., 1984). When the four randomly oxidized disulfides are correctly paired, the

RNase is natively folded and active and can convert cyclic cytidine-3,5'-

monophosphate (cCMP) into 3'CMP, which can be monitored colorimetrically. In

these experiments FL α-DsbA2, α-DsbA2ΔN or EcDsbC (10 μM final concentration)

were used in a buffer containing 100 mM sodium phosphate, 1 mM EDTA, pH 7.0,

10 μM dithiothreitol (DTT) and 40 μM of scrambled RNaseA. Scrambled RNaseA

was produced as previously described (Kurz *et al.*, 2009). At various time points, 50

μL of the reaction was mixed with 150 μL of cytidine 3',5'-cyclic monophosphate (3

mM) and hydrolysis activity was monitored using a Biotek H1 plate reader

(Millennium Science, USA) at 296 nm and 298 K. Native RNaseA and scrambled

RNaseA samples without added enzyme served as positive and negative controls,

respectively. Measurements were performed in triplicate using three different batches

of protein and presented data is the mean ± SD from three measurements.

Crystallization and crystal structure determination of α-DsbAΔ65

Crystal screening and optimisation of α-DsbAΔ65 crystals was performed at the UQ

ROCX facility (University of Queensland, Australia). For crystallisation, the hanging

drop vapour diffusion method was used. Crystals of α-DsbAΔ65 were grown by

mixing 1 µL of protein at 20 mg/mL and 1 µL crystallant solution, containing 100

mM Tris pH 7.5, 200 mM NaCl and 20% *w/v* PEG 3350. X-ray data were recorded on

the microcrystallography beamline MX2 at the Australian Synchrotron using the

Blue-Ice software (McPhillips *et al.*, 2002). Reflections were processed in XDS

(Kabsch, 2010), analysed and converted to MTZ in SCALA (Evans, 2006). Phases

were obtained by molecular replacement using remote sequence homologues

identified by Fold and Function Assignment (FFAS) (Jaroszewski *et al.*, 2011). The

top FFAS hit, *Salmonella enterica serovar Typhimurium* ScsC (SeScsC, PDB ID:

4GXZ (Shepherd *et al.*, 2013)) shares 23% sequence identity to α-DsbAΔ65. An

initial search using the complete SeScsC PDB coordinates as a model was

unsuccessful. Instead a trimmed poly-Ser template that retained the side-chains of

residues that were conserved in an alignment of SeScsC with α-DsbA2ΔN was used

to phase the structure using PHASER (McCoy *et al.*, 2007) for α-DsbA2ΔN. The root

mean square deviation between these two structures is 1.8 Å for 124 equivalent C^{α}

atoms. Further refinement was performed using PHENIX (Adams *et al.*, 2010)

(Adams *et al.*, 2011) and COOT. Molecular figures were generated in PyMOL (*The

PyMOL Molecular Graphics System, Version 1.2r3pre, Schrödinger, LLC.*, LLC.

#211). Electrostatic potential figures were generated using APBS (Dolinsky *et al.*,

2004).

RMSD calculations and structural alignments were conducted using PyMOL as well

as DaliLite {Holm, 2008 #39}. The data collection and refinement statistics are given

in (**Table 1**). The refined model of α-DsbA2ΔN has four molecules in the asymmetric

unit with chains A, B, C and D refined with 185, 189, 180 and 180 residues,

respectively. Molecules A and B are both well defined in the electron density, but

poor density regions in molecules C (residues 188-197) and molecule D (residues 92-

98 and 193-197) were not able to improve during refinement with only the Cα atoms

and occupancies set to 0 for any other atoms. The different quality of these molecules

is evident from the average B-factors of A/B (46 and 41 $\mathring{A}^2$ respectively), and

molecules C/D (63 and 70 $\mathring{A}^2$ respectively), (see **Figure 3B**). The catalytic disulfide

bonds in molecules A and molecule B were modeled in a mixed redox state (**Figure**

3), whereas in molecules C/D these are modelled as reduced.

Small angle X-ray scattering of FL α-DsbA2 and α-DsbA2Δ65

SAXS data for α-DsbA2ΔN and FL α-DsbA2 were collected on the SAXS-WAXS

beamline at the Australian Synchrotron (Kirby *et al.*, 2013). Data reduction was

carried out using Scatterbrain software (v 2.71) (Australian Synchrotron), and

corrected for solvent scattering, sample transmission, and detector sensitivity. For α-

DsbA2ΔN, serial dilutions of a ~5 mg/mL stock were loaded into a 96-well plate,

while FL α-DsbA2 was measured using an in-line SEC-SAXS set-up (see **Table 2**).

The estimated molecular mass was calculated using contrast and partial specific

volumes determined from the protein sequences (Whitten *et al.*, 2008). Data

processing and Guinier analysis was performed using Primus (v 3.2) (Konarev *et al.*,

2003). The pair-distance distribution function ($p(r)$) was generated from the

experimental data using *GNOM* (v 4.6) (Svergun, 1992), from which $I(0)$, R_g and

D_{max} were determined. The program *DAMMIN* (v 5.3) (Svergun, 1999) was used to

generate 16 dummy-atom models for each protein, (assuming C_1 symmetry for α-

DsbA2ΔN, and C_3 symmetry for α-DsbA2ΔN), which were averaged using the

program *DAMAVER* (v 2.8.0) (Volkov & Svergun, 2003), and the resolution of the

averaged structures were estimated using SASRES (Tuukkanen *et al.*, 2016). All 16

dummy-atom models were used in the averaging procedure for α-DsbA2ΔN, but only

9 (oblate) of the 16 dummy-atom models were averaged for FL α-DsbA2. Rigid-

body modelling was carried out using the program CORAL (v 1.1) (Petoukhov *et al.*,

2012). For α-DsbA2ΔN, residues D71 – L250 were taken from the crystal structure

and treated as a rigid unit, while 5 additional residues were included at the N- and C-

termini and treated as flexible linkers. For FL α-DsbA2, C_3 symmetry was assumed

and L17 – R27 (a model helical segment), D34 – E57 (a model helical segment) and

A65 – L250 (from the α-DsbA2ΔN crystal structure) were taken as rigid subunits,

while 5 additional residues were added at the N- and C-termini plus the intervening

regions between rigid segments and treated as flexible linkers. As oligomerisation

occurs through the N-terminal region, and the model helices were amphipathic, L17,

I20, W23, I36, L40, I44, F48, V52 and L55 were restrained to be less than 15 Å from

the same residue in a symmetry related chain.

FIGURE LEGENDS

Figure 1. **A) Sequence alignment of the N-terminal residues of α-DsbA2 and**

EcDsbC. Sequences were aligned using Clustal Omega (Sievers *et al.*, 2011).

Secondary structure was determined from the structure of EcDsbC or from predictions

for α-DsbA2 using JPred (Drozdetskiy *et al.*, 2015). α-helices are shown as yellow

cylinders and β-strands as grey boxes. Identical residues are marked with an asterisk.

B) Sequence alignment of α-DsbA2, PmScsC, CcScsC, LpDsbA2 and EcDsbC.

Sequences were aligned using Clustal Omega (Sievers *et al.*, 2011). The sequence

identities for α-DsbA2 are 21%, 19%, 20%, and 21% with PmScsC, CcScsC and

LpDsbA2, and EcDsbC, respectively. The N-terminal residues of each protein

sequence are shown as blue bold letters, α-helices are highlighted in yellow and β-

strands as blue arrows. The catalytic active site and *cis*-Pro loop residues are bold and

underlined. The vertical black arrow indicates the start of the catalytic (TRX) domain

in α-DsbA2.

Figure 2. Redox properties of α-DsbA2. A) Redox potential. The equilibrium

between reduced and oxidized FL α-DsbA2 (triangle) and α-DsbA2ΔN (circle) are

shown. Data are presented as the mean ± SD of two measurements. **B) Protein**

disulfide reductase activity. The insulin reduction assay was performed to measure

the ability of α-DsbA2 variants to reduce insulin. The reduction of insulin catalyzed

by the protein or non-catalyzed (blank) was monitored at 650 nm over 50 min. The

mean (black line) and standard error (light grey lines) of three replicate measurements

are shown. Disulfide oxidase (EcDsbA) and disulfide isomerase (EcDsbC) activities

are shown for comparison. **C) Protein disulfide isomerase activity.** Isomerization of

scRNase A in the presence of EcDsbC (diamond), FL α-DsbA2 (triangle) or α-

DsbA2ΔN (circle), positive control (RNase, inverse triangle) and blank (scRNase

without enzyme, square). Data are presented as the mean ± SD of three replicate

measurements. **D) Dithiol oxidase activity**. Fluorescence curves showing dithiol

oxidation (in the presence of GSSG) of a model peptide by: EcDsbA (positive control,

square), α-DsbA2ΔN (circle), FL α-DsbA2 (triangle), GSSG (without enzyme *ie*

negative control, star), and α-DsbA2 without peptide (Blank, diamond). Data are

presented as the mean ± SD of three replicate measurements.

Figure 3. Structure of α-DsbA2ΔN. A) Ribbon representation of α-DsbA2ΔN chain

B showing the TRX domain in blue and catalytic cysteine sulfur atoms illustrated as

yellow spheres. The helical insertion is highlighted in grey, and the N-terminus of the

truncated protein is shown in light green. **Dashed lined box:** Active site cysteines

Cys107 and Cys110 of chain B were modeled in a mixed redox state (shown for chain

A, with the $2F_o$–F_c electron density map contoured at 1.0 σ). **B)** The four molecules in

the asymmetric unit. The relative crystallographic temperature factors of the refined

coordinates are shown by backbone thickness (low-to-high shown as thin-to-thick)

and colour (low–to–high coloured from blue–to–red).

Figure 4. Small angle X-ray scattering data for α-DsbA2ΔN and FL α-DsbA2.

A) Measured scattering data for α-DsbA2ΔN (red, multiplied by a factor of 10 for

clarity) and FL α-DsbA2 (blue). The scattering profile of rigid-body models are

shown as solid black lines overlaid on the scattering data for α-DsbA2ΔN ($\chi^2 = 3.88$;

CorMap test (Franke *et al.*, 2015), 294 points, C=66, P=0.000) and FL α-DsbA2

($\chi^2 = 1.37$; CorMap test, 320 points, C=13, P=0.037). **Inset:** Guinier plot for α-

DsbA2ΔN (red, $R^2 = 0.999$) and FL α-DsbA2 (blue, $R^2 = 1.000$). **B)** The pair-distance

distribution function, $p(r)$, derived from the scattering data are indicative of a globular

structure with a maximum dimension of ~63 Å for α-DsbA2ΔN (red) and ~87 Å for

FL α-DsbA2 (blue). For reference, the experimental $p(r)$ for the trimeric PmScsC is

also shown (dotted line). **C)** Probable shape of α-DsbA2ΔN (monomeric) obtained

from the filtered average of 16 dummy-atom models (red envelope): $\chi^2 = 1.038 \pm$

0.002; NSD $= 0.446 \pm 0.021$; Resolution $= 17 \pm 2$. **D)** Probable shape of FL α-DsbA2

obtained from the filtered average of 9 dummy-atom models (blue envelope):

$\chi^2 = 1.202 \pm 0.003$; NSD $= 0.602 \pm 0.027$; Resolution $= 30 \pm 2$. Images in **C** and **D**

were generated using PyMol, where the grey shapes represent the total volume

encompassed by the aligned dummy-atom models and the corresponding rigid-body

model is shown aligned to the filtered model (flexible regions are represented by

chains of black spheres).

Figure 5. Comparison of rigid body models of FL α-DsbA2 and PmScsC. Model

of FL α-DsbA2 from **A)** the side and **B)** bottom view. Model of PmScsC (SASDBD

code: SASDB94) from **C)** side and **D)** bottom view. For both FL α-DsbA2 and

PmScsC, the N-terminal trimerisation domain is shown as a green ribbon, the

catalytic domain as a white ribbon, and active site residues as yellow spheres. The

regions treated as flexible linkers in the rigid-body model are not shown, which is the

reason for the apparent gap in the N-terminal trimerisation domain shown in panel A.

Table 1 X-ray data collection and refinement statistics for α-DsbA2ΔN

Data collection	
Wavelength (Å)	0.9537
Resolution range (Å)	19.77 - 2.25
Highest resolution shell (Å)	2.32 – 2.25
Space group	P2$_1$
Unit cell *a, b, c* (Å)	54.4, 104.8, 67.7
α, β, γ (°)	90.0, 96.5, 90.0
No mols in asymm unit	4
Observed reflections	132,099
Unique reflections	35,423 (3209)
R$_{merge}$	0.051 (0.265)
R$_{pim}$	0.038 (0.187)
Completeness (%)	99.04 (97.8)
$<I/\sigma(I)>$	13.6 (3.9)
Redundancy	3.8 (3.6)
Refinement statistics	
R$_{factor}$ (%) (top shell)	22.7 (20.9)
R$_{free}$ (%) (top shell)	27.8 (25.3)
Unique reflections	35,423
No of non-H atoms	6064
Protein	5799
Water	265
B-factors Average	58.0
RMSD from ideal geometry	
Bonds (Å)	0.002
Angles (°)	0.447
Molprobity analysis	
Ramachandran favoured/outliers (%)	97.12/0.0
Clashscore all atoms (percentile)	3.29 (99[th])
Molprobity score (percentile)	1.35 (99[th])

Table 2 SAXS data collection and analysis details

Data Collection Parameters	α-DsbA2ΔN	α-DsbA2
Instrument	SAXS-WAXS (Australian Synchrotron)	
Beam geometry	Point	
Wavelength (Å)	1.033	1.127
Sample to detector distance (m)	1.575	1.575
q-range (Å^{-1})	0.01 – 0.57	0.01-0.48
Exposure time (s)	16 (8×2 s exposures)	40 (8×5 s exposures)
Configuation	Concentration series from 96-well plate	SEC-SAXS Column: WTC-030S5 Flow: 0.5 mL/min Inject:100 μL, 19.8 mg/mL
Protein concentration range (mg/mL)	1.25 – 5.00	0.35 – 2.20
Temperature (K)	283	293
Absolute intensity calibration	Water	
Sample details		
Extinction coefficient (A_{280}, 0.1% *w/v*)	0.494	0.762
Partial specific volume (cm^{3}g^{-1})	0.735	0.735
Contrast, $\Delta\rho$ (10^{10} cm^{-2})	2.88	2.89
Molecular mass [from sequence] (kDa)	21.1	27.1
Protein concentration (mg/mL)*	1.25	0.75
Structural parameters		
$I(0)$ (cm^{-1}) [from Guinier]	0.0215 ± 0.0001	0.04846 ± 0.00003
R_g (Å) [from Guinier]	18.7 ± 0.1	28.0 ± 0.1
$I(0)$ (cm^{-1}) [from $p(r)$]	0.0215 ± 0.0001	0.04845 ± 0.00002
R_g (Å) [from $p(r)$]	18.8 ± 0.1	27.9 ± 0.1
D_{max} (Å)	63 ± 3	87 ± 3
Porod volume (Å^3)	27500 ± 1500	91300 ± 4500
Molecular mass determination		
Molecular mass [from $I(0)$] (kDa)	23 ± 1	87 ± 5
Molecular mass [from Porod] (kDa)	23 ± 1	75 ± 5

*α–DsbA2Δ65: $I(0) = 0.0427 \pm 0.0001$ cm^{-1}, $R_g = 18.9 \pm 0.1$ Å, $M = 23$ kDa (2.5 mg/mL); $I(0) = 0.0890 \pm 0.0002$ cm^{-1}, $R_g = 19.2 \pm 0.1$ Å, $M = 24 \pm 2$ kDa (5.0 mg/mL); There is a small but significant systematic change in R_g and M in the concentration range measured, but based on the trend, 1.25 mg/mL was deemed to be free of concentration dependent effects; FL αDsbA2: At the peak concentration of ~2.2 mg/mL, $R_g = $ ~27.6 Å, which increases and plateaus at $R_g = $ ~28.0 Å at a concentration of between 0.35 – 1.15 mg/mL. This range was deemed to be free of concentration dependent effects and all eight frames collected over these concentrations were combined, where the average concentration was 0.75 mg/mL.

References

References

Abramoff, M. D., Magelhaes, P.J., Ram, S.J. (2004). *Biophotonics International* **11**,
36-42.

Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Davis, I. W., Echols, N.,
Headd, J. J., Hung, L. W., Kapral, G. J., Grosse-Kunstleve, R. W., McCoy, A. J.,
Moriarty, N. W., Oeffner, R., Read, R. J., Richardson, D. C., Richardson, J. S.,
Terwilliger, T. C. & Zwart, P. H. (2010). *Acta Crystallogr D Biol Crystallogr* **66**,
213-221.

Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Echols, N., Headd, J. J.,
Hung, L. W., Jain, S., Kapral, G. J., Grosse Kunstleve, R. W., McCoy, A. J.,
Moriarty, N. W., Oeffner, R. D., Read, R. J., Richardson, D. C., Richardson, J. S.,
Terwilliger, T. C. & Zwart, P. H. (2011). *Methods* **55**, 94-106.

Australian Synchrotron, C., VIC *Scatterbrain—Software for acquiring, processing*
and viewing SAXS/WAXS data at the Australian Synchrotron. Version 2.71.

Bader, M., Muse, W., Ballou, D. P., Gassner, C. & Bardwell, J. C. (1999). *Cell* **98**,
217-227.

Bairoch, A., Apweiler, R., Wu, C. H., Barker, W. C., Boeckmann, B., Ferro, S.,
Gasteiger, E., Huang, H., Lopez, R., Magrane, M., Martin, M. J., Natale, D. A.,
O'Donovan, C., Redaschi, N. & Yeh, L. S. (2005). *Nucleic Acids Res* **33**, D154-159.

Cho, S. H., Parsonage, D., Thurston, C., Dutton, R. J., Poole, L. B., Collet, J. F. &
Beckwith, J. (2012). *MBio* **3**.

Depuydt, M., Leonard, S. E., Vertommen, D., Denoncin, K., Morsomme, P., Wahni,
K., Messens, J., Carroll, K. S. & Collet, J. F. (2009). *Science* **326**, 1109-1111.

Dolinsky, T. J., Nielsen, J. E., McCammon, J. A. & Baker, N. A. (2004). *Nucleic*
Acids Res **32**, W665-667.

Drozdetskiy, A., Cole, C., Procter, J. & Barton, G. J. (2015). *Nucleic Acids Res* **43**,
W389-394.

Evans, P. (2006). *Acta Crystallogr D Biol Crystallogr* **62**, 72-82.

Feige, M. J. & Hendershot, L. M. (2011). *Curr Opin Cell Biol* **23**, 167-175.

Fischer, H., Neto, M. D., Napolitano, H. B., Polikarpov, I. & Craievich, A. F. (2010).
Journal of Applied Crystallography **43**, 101-109.

Flores, H. A. & O'Neill, S. L. (2018). *Nat Rev Microbiol*.

Franke, D., Jeffries, C. M. & Svergun, D. I. (2015). *Nature Methods* **12**, 419.

Furlong, E. J., Choudhury, H. G., Kurth, F., Duff, A. P., Whitten, A. E. & Martin, J.
L. (2018). *J Biol Chem* **293**, 5793-5805.
Furlong, E. J., Lo, A. W., Kurth, F., Premkumar, L., Totsika, M., Achard, M. E. S.,
Halili, M. A., Heras, B., Whitten, A. E., Choudhury, H. G., Schembri, M. A. &
Martin, J. L. (2017). *Nat Commun* **8**, 16065.
Heras, B., Shouldice, S. R., Totsika, M., Scanlon, M. J., Schembri, M. A. & Martin, J.
L. (2009). *Nat Rev Microbiol* **7**, 215-225.
Hilgenboecker, K., Hammerstein, P., Schlattmann, P., Telschow, A. & Werren, J. H.
(2008). *FEMS microbiology letters* **281**, 215-220.
Hillson, D. A., Lambert, N. & Freedman, R. B. (1984). *Methods Enzymol* **107**, 281-
294.
Holmgren, A. (1979). *J Biol Chem* **254**, 9113-9119.
Inaba, K. (2009). *J Biochem* **146**, 591-597.
Inaba, K. & Ito, K. (2002). *EMBO J* **21**, 2646-2654.
Jaroszewski, L., Li, Z., Cai, X. H., Weber, C. & Godzik, A. (2011). *Nucleic Acids Res*
39, W38-44.
Kabsch, W. (2010). *Acta Crystallogr D Biol Crystallogr* **66**, 125-132.
Kirby, N. M., Mudie, S. T., Hawley, A. M., Cookson, D. J., Mertens, H. D. T.,
Cowieson, N. & Samardzic-Boban, V. (2013). *Journal of Applied Crystallography*
46, 1670-1680.
Konarev, P. V., Volkov, V. V., Sokolova, A. V., Koch, M. H. J. & Svergun, D. I.
(2003). *Journal of Applied Crystallography* **36**, 1277-1282.
Kpadeh, Z. Z., Day, S. R., Mills, B. W. & Hoffman, P. S. (2015). *Mol Microbiol* **95**,
1054-1069.
Kpadeh, Z. Z., Jameson-Lee, M., Yeh, A. J., Chertihin, O., Shumilin, I. A., Dey, R.,
Day, S. R. & Hoffman, P. S. (2013). *J Bacteriol* **195**, 1825-1833.
Kurz, M., Iturbe-Ormaetxe, I., Jarrott, R., Shouldice, S. R., Wouters, M. A., Frei, P.,
Glockshuber, R., O'Neill, S. L., Heras, B. & Martin, J. L. (2009). *Antioxidants &*
Redox Signaling **11**, 1485-1500.
Landeta, C., Boyd, D. & Beckwith, J. (2018). *Nat Microbiol* **3**, 270-280.
Maiti, R., Van Domselaar, G. H., Zhang, H. & Wishart, D. S. (2004). *Nucleic Acids*
Res **32**, W590-594.
McCarthy, A. A., Haebel, P. W., Torronen, A., Rybin, V., Baker, E. N. & Metcalf, P.
(2000). *Nature Structural Biology* **7**, 196-199.

McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C. &

Read, R. J. (2007). *J Appl Crystallogr* **40**, 658-674.

McPhillips, T. M., McPhillips, S. E., Chiu, H. J., Cohen, A. E., Deacon, A. M., Ellis,

P. J., Garman, E., Gonzalez, A., Sauter, N. K., Phizackerley, R. P., Soltis, S. M. &

Kuhn, P. (2002). *J Synchrotron Radiat* **9**, 401-406.

Missiakas, D., Schwager, F. & Raina, S. (1995). *EMBO J* **14**, 3415-3424.

Mossner, E., Huber-Wunderlich, M. & Glockshuber, R. (1998). *Protein Sci* **7**, 1233-

1244.

Orthaber, D., Bergmann, A. & Glatter, O. (2000). *Journal of Applied Crystallography*

33, 218-225.

Petoukhov, M. V., Franke, D., Shkumatov, A. V., Tria, G., Kikhney, A. G., Gajda,

M., Gorba, C., Mertens, H. D., Konarev, P. V. & Svergun, D. I. (2012). *J Appl*

Crystallogr **45**, 342-350.

The PyMOL Molecular Graphics System, Version 1.2r3pre, Schrödinger, LLC.

Rost, B., Yachdav, G. & Liu, J. (2004). *Nucleic Acids Res* **32**, W321-326.

Shepherd, M., Heras, B., Achard, M. E., King, G. J., Argente, M. P., Kurth, F.,

Taylor, S. L., Howard, M. J., King, N. P., Schembri, M. A. & McEwan, A. G. (2013).

Antioxid Redox Signal **19**, 1494-1506.

Shevchik, V. E., Condemine, G. & Robert-Baudouy, J. (1994). *EMBO J* **13**, 2007-

2012.

Shouldice, S. R., Heras, B., Walden, P. M., Totsika, M., Schembri, M. A. & Martin, J.

L. (2011). *Antioxid Redox Signal* **14**, 1729-1760.

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R.,

McWilliam, H., Remmert, M., Soding, J., Thompson, J. D. & Higgins, D. G. (2011).

Mol Syst Biol **7**, 539.

Studier, F. W. (2005). *Protein Expression and Purification* **41**, 207-234.

Svergun, D. I. (1992). *Journal of Applied Crystallography* **25**, 495-503.

Svergun, D. I. (1999). *Biophys J* **76**, 2879-2886.

Thompson, J. D., Higgins, D. G. & Gibson, T. J. (1994). *Nucleic Acids Res* **22**, 4673-

4680.

Tuukkanen, A. T., Kleywegt, G. J. & Svergun, D. I. (2016). *IUCrJ* **3**, 440-447.

Volkov, V. V. & Svergun, D. I. (2003). *Journal of Applied Crystallography* **36**, 860-

864.

Walden, P. M., Halili, M. A., Archbold, J. K., Lindahl, F., Fairlie, D. P., Inaba, K. &

Martin, J. L. (2013). *PLoS One* **8**, e81440.

Whitten, A. E., Cai, S. Z. & Trewhella, J. (2008). *Journal of Applied Crystallography*

41, 222-226.

Wormser, G. P., Dattwyler, R. J., Shapiro, E. D., Halperin, J. J., Steere, A. C.,

Klempner, M. S., Krause, P. J., Bakken, J. S., Strle, F., Stanek, G., Bockenstedt, L.,

Fish, D., Dumler, J. S. & Nadelman, R. B. (2006). *Clin Infect Dis* **43**, 1089-1134.

Zapun, A., Bardwell, J. C. & Creighton, T. E. (1993). *Biochemistry* **32**, 5083-5092.

Zapun, A., Missiakas, D., Raina, S. & Creighton, T. E. (1995). *Biochemistry* **34**,

5075-5089.

Figure 1

A

```
α-DsbA2  ---------------------SLPI--INNWLSNRSQML-NDD-----------YIGEKLDNYISRNFDKIVKTLKEESIKSSY----  49

EcDsbC   DDAAIQQTLAKMGIKSSDIQPAPVAGMKTVLTNSGVLYITDDGKHIIQGPMYDVSGTAPVNVTNKMLLKQLNALEKEMIVYKAPQEK  87
                             *:  ::.  *:*  .  :   .**              *  :   *  .   :  *:::*::*  *   .
```

B

```
α-DsbA2  ----------SLPIINN-WLSNRSQMLNDDYIGEKLD----NYISRNFDKIVKTLKEESI  60
PmScsC   --------------------AALNAAQEKEVRALVR----DTLVSNPEILEEAIMALQT  37
CcScsC   -------------------LAGCDQSKPDKAFGEKVR----AYLLEHPEVLMEASQKLQE  40
LpDsbA2  ----------------ADTASASLSDAQKKEIEKVIH----DYLINNPEVLLEASQALQQ  51
EcDsbC   DDAAIQQTLAKMGIKSS-----DIQPAPVAGMKTVLTNSGVLYITDDGKHIIQGPMYDV-  54
                                                  :          :    . :   . . :

α-DsbA2  KSSYAARDNVTKSKISQYKDQIFDLTY-PYSGNENSSVIAVGFLDYSCGHCKAIKNDIKQ  119
PmScsC   K-KADEQQAQFRQALASEHDALYNDAASPRIGAKDAKLVLVSFTDYNCPYCKRFDPLLEK  96
CcScsC   K-QAAQQAVSSQKAIGEYRQAIERDPR-DIVINPAGTITVTEFFDYFCGYCRQATPAVLE  98
LpDsbA2  K-QQQNMQQQAQAAIQENAEQVFQGKL-TTVGNPKGNVTLVEFFDYCIHCKKMASTIEN  109
EcDsbC   --SGTAPVNVTNKMLLKQLNALEKEM--IVYKAPQEKHVITVFTDITCGYCHKLHEQMAD  110
                .      :   : :    : :               .  . ** * * :*:   :.  

α-DsbA2  LIN-DGKIKYIFKDAPILGNASLKAAKSALAVYF-LDKEKYFDFHHAALSHKGEFSDESI  177
PmScsC   ITEQYPDVAVIIKPLPPKGESSAKASQAVLSVWK-EDPKAFLALHQRLMQKKTMLDNASI  155
CcScsC   LVQKNPDIRLVLKDFVIFGNDSEAAARIALGA---KDQGKSLELHKALMA-ENALDARGA  154
LpDsbA2  LVKKDSGLRVIYKFPIFGKTSDLASRVALAA---GMQGKYQAMHNALITIDKRLDEKTV  166
EcDsbC   YNA--LGITVRYLAFPRQGLDSDAEKEM-KAIWCAKDKNKA-------------FDDVMA  154
                :          *  *    .   .                          :.  

α-DsbA2  LDIVKNIGIDEDDFNDSIKDNADKIEQMINNSRLLVRDLGVGGTPFLIIGD-SLFVGATD  236
PmScsC   EDAMKSTNTSKIKL------TDDSLKTLQNNLELSRKLGIQGTPATVIGD-TILPGAVD  207
CcScsC   LRIAERLGIDMDKAKAVG--ESQAITQHLADTDALARALNLSGTPAFIVGD-TLVPG---  208
LpDsbA2  MDAAKSIGLDMQKLKKDM--DSQEVTDILDANRQLAEKLHLMGTPFIIGS-TPDGQYKK  223
EcDsbC   GKSV---------------APASCDVDIADHYALGVQLGVSGTPAVVLSNGTLVPGYQP  198
                                  :      *    * : *  ***  ::.. :

α-DsbA2  LN---VLRKKVDELSH------KQG-     252
PmScsC   YDQLEII---VKEQL----AKVKK--    224
CcScsC   -ADIDALKLAIEQT--R-AARAKAG-    229
LpDsbA2  GSEISFIPGATSEQSLRELIKKAAGN    249
EcDsbC   PKEMKEF---LDEH-----QKMTSGK    216
                :      .:
```

Figure 2

Figure 3

Figure 4

Figure 5

A

72 Å

B

C

78 Å

D