Diversity and composition of methanotroph communities in caves

Kevin D. Webster[a,,b]# Arndt Schimmelmann[c], Agnieszka Drobniak[d], Maria Mastalerz[d], Laura

Rosales Lagarde[e], Penelope J. Boston[f], Jay T. Lennon[g]#

[a]Diné College, School of Science, Technology, Engineering and Math, Tsaile, Arizona, USA

86556

[b]Planetary Science Institute, Tucson, Arizona, USA 85719

[c]Department of Earth and Atmospheric Sciences, Indiana University, Bloomington, Indiana,

USA 47405

[d]Indiana Geological and Water Survey, Indiana University, Bloomington, Indiana, USA 47405

[e]Department of Physical and Life Sciences, Nevada State College, School of Liberal Arts and

Sciences, Henderson, Nevada, USA.

[f]NASA Astrobiology Institute, NASA Ames Research Center, Moffett Field, California, USA.

[g]Department of Biology, Indiana University, Bloomington, Indiana, USA 47405

#Address correspondence to Kevin D. Webster, webster@psi.edu, or Jay T. Lennon,

lennonj@indiana.edu.

ABSTRACT Methane oxidizing microorganisms (methanotrophs) are ubiquitous in the environment and represent a major sink for the greenhouse gas methane (CH_4). Recent studies have demonstrated that methanotrophs are abundant and contribute to CH_4 dynamics in caves. However, very little is known about what controls the distribution and abundance of methanotrophs in subterranean ecosystems. Here, we report a survey of soils collected from > 20 caves in North America to elucidate the factors shaping cave methanotroph communities. Using 16S rRNA sequencing, we recovered methanotrophs from nearly all (98 %) of the samples, including cave sites where CH_4 concentrations were at or below detection limits (≤ 0.3 ppmv). We identified a core methanotroph community among caves that was comprised of high-affinity methanotrophs. Although associated with local-scale mineralogy, methanotroph composition did not systematically vary between the entrances and interior of caves, where CH_4 concentrations varied. We also observed that methanotrophs are able to disperse readily between cave systems showing these organisms have low barriers to dispersal. Last, the relative abundance of methanotrophs was positively correlated with cave-air CH_4 concentrations suggesting that these microorganisms contribute to CH_4 flux in subterranean ecosystems.

IMPORTANCE Recent observations have shown that the atmospheric greenhouse gas methane (CH_4) is consumed by microorganisms (methanotrophs) in caves at rates comparable to CH_4 oxidation in surface soils. Caves are abundant in karst landscapes that comprise 14 % of Earth's land surface area, and therefore may represent a potentially important, but overlooked CH_4 sink. We sampled cave soils to gain a better understand the community composition and structure of cave methanotrophs. Our results show that the members of the USC-γ clade are dominant in cave communities and can easily disperse through the environment, that methanotroph relative

42 abundance was correlated with local scale mineralogy of soils, and that the relative abundance of

43 methanotrophs was positively correlated with CH_4 concentrations in cave air.

45 **KEYWORDS:** Biogeography; Cave; Greenhouse Gas; Karst; Methane; Methanotroph

INTRODUCTION

Methane (CH_4) is as major greenhouse gas in Earth's atmosphere, and an energy source

for humanity (1, 2). Because of this, it is important to identify the pathways that produce and

consume CH_4. Two major processes are responsible for the removal of CH_4 from Earth's

atmosphere, oxidation by atmospheric hydroxyl radicals, and by methane-consuming

microorganisms (methanotrophs) in surface soils and waters. Top-down and bottom up estimates

of yearly atmospheric CH_4 removal indicate hydroxyl radicals account for 90 % of the sink while

methanotrophs consume roughly 5 % (3). Despite the influence that methanotrophs have on

regulating Earth's climate, gaps remain in our understanding of methanotrophs, like how they

respond to changing CH_4 concentrations or how they disperse through the environment (4, 5).

Rocks that host caves are widespread on Earth, and are emerging as an environment that

consumes atmospheric CH_4 (6–11) (Figure 1). Recent studies of cave microbial communities

have revealed the presence of atmospheric-CH_4 consuming methanotrophs in caves (8, 12–15).

The methanotrophs that are thought to be responsible for this are termed high-affinity

methanotrophs, and have been previously observed from upland soils. These groups are referred

to as the Upland Soil Cluster (USC) -α and -γ clades. Intermittent or unsustained atmospheric

CH_4 oxidation has also been observed in methanotrophs among the families of the

Alphaproteobacteria (the *Methylocystaceae* and *Beijernickiaceae*), but it is generally thought that

these organisms contribute little to atmospheric CH_4 consumption (16). Both high-affinity and

low-affinity alphaproteobacterial methanotrophs tend to be the most prominent members of the

methanotrophic community in environments with low CH_4 concentrations and high O_2

concentrations. They are less common in environments with high CH_4 and low O_2

concentrations, like lakes or geological seeps, which are dominated by low-affinity

methanotrophs found among families belonging to the Gammaproteobacteria (Methylococcaceae

and Methylothermaceae) (17, 18).

The factors that control the distribution and abundance of methanotrophs in caves are not

well understood. Previous work has shown that methanotrophs are frequently related to CH_4

concentrations, soil texture, and the abundances of other microorganisms (19), but it is unclear

how these factors, or the local mineralogy influences methanotroph communities in caves. For

example, low-affinity methanotrophs exhibit niche partitioning between environments with high

and low CH_4 concentrations (18, 20). However, it is unknown if methanotrophic communities

exhibit similar changes in community composition along atmospheric to subatmospheric

gradients of CH_4.

In addition, very little is known about the spatial distribution of methanotrophs, even

though such information could shed light on the biogeography of microbes within and among

cave ecosystems. While the community composition of methanotrophs may be influenced by

environmental conditions of caves, their abundances may also be influenced by their capacity to

disperse. Caves are unique and insular habitats that have a patchy distribution in the terrestrial

landscape. However, cells may move through cracks and pores from surface environments or

aeolian transport into nearby cave systems. These potentially stochastic modes of dispersal could

explain be important in structuring cave methanotroph communities and may help explain the

global distribution of methanotrophs. For example, similar methanotrophs have been observed in

both Hawai'i and the Arctic (4) suggesting that the movement of these organisms may be

relatively unhindered. However, biota in caves are typically unique due to their isolation from

other environments (21, 22). Thus, caves provide an ideal environment to test whether

methanotrophs are able to disperse readily across the environment or are limited by their ability

to disperse.

In this study, we sampled 42 cave soils from 21 limestone caves in North America to

examine the factors regulating the community composition of cave methanotrophs (Figure 1).

First, we hypothesized that the methanotroph composition should be similar to what has been

reported for methanotrophic communities in other low CH_4 environments, and that the relative

abundance of methanotrophs should be correlated with cave-air CH_4 concentrations.

Additionally, we hypothesized that, on the local scale, methanotrophs would be affected by the

soil texture and mineral composition of cave soils since these features influence the diffusion of

air and micronutrient abundances. Finally, on the regional scale we hypothesized that high-

affinity methanotrophs should be able to disperse easily between caves.

RESULTS

Methanotroph community structure. Methanotrophs were present in nearly all caves.

Based on 16s rRNA sequencing we identified methanotrophs in 98 % of the cave-soil samples,

including locations where CH_4 concentrations were at or below the analytical detection limit ($\leq$

0.3 ppmv). The maximum relative abundance of methanotrophs in the cave soil samples was

6.25 % and the median relative abundance was 0.88 %. The average site contained 6 OTUs

belonging to methanotrophs. High-affinity methanotrophs comprised a median of 99.0 % of the

total methanotrophic community at locations where methanotrophs were observed. Members of

the USC-γ clade represented 47.7 % of the community on average (median), while the members

of the USC-α clade represented 43.1 % of the community on average (median). Low-affinity

methanotrophs represented 0.09 % of the observed methanotroph community on average

(median). Members of the USC-γ and -α were represented by 22 and nine OTUs, repectively,

whereas low-affinity methanotrophs belonging to the Alphaproteobacteria and

Gammaproteobacteria were represented by 18 and 22 unique OTUs, respectively. Members of

the USC-γ clade were dominant in the sampled caves (Figure 2). The members of

methanotrophic community from Cave 38 (Mexico) were solely high-affinity methanotophs.

Core Methanotroph Community. We defined taxa that were present in at least 60 % of

samples as the core methanotroph community (23, 24). Based on this, the core methanotroph

community was made up of three OTUs. Two of these OTUs were from members of the USC-α

and one was from the USC-γ. These taxa were found together in 79 % of samples, and the most

widely distributed OTU was observed in 93 % of samples. The core methanotroph community

accounted for 97 % of the total methanotroph sequences recovered across all cave samples.

Environmental influences on community structure. Methanotroph composition was

related to the mineralogy of cave soils (Figure 3). The total methanotrophic community was

related to the abundance of clinochlorite (multiple regression, $P = 0.008$), muscovite (multiple

regression , $P = 0.03$), and microcline (multiple regression, $P = 0.007$). High-affinity

methanotrophs showed relationships with the abundance of clinochlorite (multiple regression, P

= 0.03), and microcline (multiple regression, $P = 0.02$), while the low-affinity methanotrophs

were not related to the abundance of any observed mineral ($P > 0.05$). In contrast to

mineralogical associations, the composition of the methanotrophic community was not related to

the distributions of grain size that comprised a soil (regression, $P > 0.05$).

Biogeography. In ecology, the distance decay relationship (DDR) describes how

community similarity changes with increasing geographic distance and environmental similarity

(25, 26). Briefly, as the space between two points increases, both biological communities and the

environment tend to become less similar. The degree to which species can disperse through the

environment can alter how quickly the community assemblage changes across space.

Communities that are hindered by dispersal tend to change more quickly than measures of

environmental similarity, while communities that have have a high capacity to disperse tend to

change as quickly as measures of environmental similarity. Within the methanotroph

assemblages, we found that the composition of high-affinity taxa changed at the same rate as

geographic distance and measures of environmental similarity ($m_{high\text{-}afinnity}$ = -0.09; m_{env} = -0.05;

Difference of slopes test, $P = 0.10$) (Figure 4). Meanwhile, the composition of low-affinity

methanotrophs and the measures of environmental similarity changed little with geographic

distance ($m_{low\text{-}affinity}$ = 0.10 ± 0.06; Linear regression, $P = 0.09$; m_{env} = −0.04 ± 0.05; Linear

regression, $P = 0.43$). At smaller spatial scales, the relative abundance of methanotrophs was not

related to the distance from a cave entrance (High-affinity: Spearman's rank correlations, ρ =

−0.15, S = 4195, $P = 0.45$; Low-affinity: Spearman's rank correlations, $\rho = 0.31$, S = 2488, P =

0.09).

Relationships with cave air CH$_4$ concentrations. The CH$_4$ concentration in cave air from caves in the USA was positively correlated with the relative abundance of methanotrophs (Figure 5, Spearman's test, $\rho = 0.40$, S = 5451, $P < 0.01$). This relationship was largely driven by high-affinity methanotrophs (Spearman's test, $\rho = 0.42$, S = 5778, $P < 0.01$). The relative abundances of low-affinity methanotrophs were not correlated with the CH$_4$ concentration (Low-affinity methanotrophs: Spearman's test, $\rho = 0.03$, S = 9601, $P = 0.86$). The highest relative abundance of methanotrophs at any site occurs in a sample from Cave 38 which has internal sources of CH$_4$ and hydrogen sulfide (H$_2$S) (27), however samples from this cave also contained low relative abundances of methanotrophs.

DISCUSSION

Methanotroph community composition and diversity in caves. Methanotrophs are widespread in cave environments, with high-affinity methanotrophs dominating these ecosystems. The dominance of methanotrophs from the high-affinity clades in the sampled caves is consistent with observations that have been made in other low CH$_4$ environments. For example, high-affinity taxa dominate methanotroph assemblages in forest soils, Arctic soils, and soils associated with basaltic rocks (4, 27, 28). Additionally, the high relative abundance of methanotrophs in the cave soils from this study, and in particular of the USC-γ clade, agrees with observations from other caves where relative abundances of high-affinity methanotrophs can exceed 10 % of the entire bacterial community (8, 12, 14, 15). The relative abundances of methanotrophs observed in our study are higher than the observations of methanotrophs from caves in Vietnam (7) and contrasts with the relatively depauperate assemblage found in a Spanish cave, which reportedly consisted of only three species (29). The highest relative

abundance of methanotrophs at any site occurs in a sample from Cave 38 which has an internal

CH_4 source (30).

Although less abundant, low-affinity methanotrophs from the Alphaproteobacteria and

Gammproteobacteria were also commonly observed in our cave survey. Low-affinity

alphaproteobacterial methanotrophs are typically more abundant in aerated soils where they have

access to oxygen, whereas low-affinity gammaproteobacterial methanotrophs are more often

found in environments with high CH_4 concentrations and low oxygen availability (17, 18). Cave

soils are typically nutrient poor (31, 32), implying that anoxic conditions created by the decay of

organic matter are not likely to be widespread. Thus, the presence of low-affinity

gammaproteobacterial methanotrophs in cave soils may be limited to local pockets of anoxia

which have been suggested based on observations of limited methanogenesis in cave systems

(10, 11).

We identified a core group of three OTUs that were present in the studied caves. This

core community was derived entirely of high-affinity methanotrophs, suggesting that these taxa

are well attuned to subterranean environments that they are able to disperse between cave

ecosystems. This interpretation is consistent with previous observations suggesting that high-

affinity methanotrophs have high-dispersal capabilities. For example, methanotrophs from a

newly formed soil in Hawai'i showed close taxonomic affiliations with Arctic methanotrophs

(4). Given that microbes are known to travel intercontinental distances on windblown dust (33),

it is possible that surface dwelling high-affinity methanotrophs (27, 34) may be carried into

caves on windblown dust or other passive vectors.

It is important to note that our survey presents a fairly conservative estimate for diversity

and abundance of methanotrophs in cave ecosystems. In the bacteria, methanotrophy is carried

out through the presence of the enzyme methane monooxygenase. Methanotrophs are viewed as

specialists with regard to their metabolism (35), and, as a result, once the enzymatic machinery

has been established in a lineage, it is likely to be conserved since there are barriers to evolving

back into generalist species (36). Recent findings suggest that some high-affinity methanotrophs

may be able to use other trace gases for carbon and energy, however (37). Often, diversity

inventories of methanotrophs focus on sequences of the alpha subunit of the particulate methane

monooxygenase enzyme (34). 16S rRNA analysis of methanotrophs may therefore miss some

individuals in a community. However, in some studies, it has been shown that the two

approaches yield similar estimates of diversity (38).

Microenvironmental influences on methanotroph communities. Our results indicate

that soil properties can play an important role in structuring methanotroph communities. The

relationship between soil mineralogy and methanotrophs is most likely related to the abundance

of trace metals. For example, copper is the trace metal responsible for the oxidation of CH_4 in

particulate methane monooxygenase enzyme. The methanotroph community also showed

relationships with the abundance of clinochlorite, muscovite, and microcline. These minerals are

associated with felsic metamorphic and igneous rocks, and metals like copper are associated with

these kinds of rocks as well (39).

In contrast, the grain size of the cave soils did not influence the abundance or

composition of methanotrophs, even though these edaphic characteristics strongly affect the

physical and chemical properties that influence metabolism. Previous research observed that

methanotrophs tend to be more abundant in silt and clay fractions of soils because these fractions

made up the bulk of the soils themselves (40, 41). Despite some previous observations showing

relationships between methanotrophs and soil characteristics, other studies have also shown

absences of relationships between methanotroph community composition and soil characteristics

(42). The importance of soil grain size in structuring methanotroph communities remains an open

question.

Biogeography of methanotroph communities. We observed regional changes in

methanotrophic communities in caves. The community composition of high-affinity

methanotrophs changed at the same rate as measures of environmental similarity compared to

changes in the distance between the samples. This suggests that process like mass effects—

which are dominant when organisms are able to actively disperse through their environment—

are operating on this community (43). This observation aligns with past work indicating that high

affinity methanotrophs are able to disperse through the environment (4), and the observation that

three OTU accounted for 97 % of the total methanotrophic community. Dispersal through

surface soils (27, 34) to caves in drip waters, and transport into caves on windblown dust (30)

represent two possible ways for methanotrophs to disperse into cave systems.

The community composition of low-affinity methanotrophs did not change appreciably

with geographic distance nor did measures of environmental similarity for these samples. This

suggests that neutral processes are operating on these communities (43) and that random birth-

death processes, or stochastic dispersal event are important in determining the distribution and

abundance of methanotroph popoulations. These processes are thought to operate more strongly

on communities that are not strongly influenced by the environment and are the strongest in

small populations (44). This suggests that these communities may be being passively dispersed

into the caves.

Biogeographic patterns at the local scale were also relatively weak. The composition and

relative abundance of methanotrophic communities at the entrance versus the interior of caves

were statistically indistinguishable. This suggests that environmental conditions at the entrance

and interior of the cave are similar in ways the select for similar methanotroph communities.

Alternatively, similarity of methanotrophs at the relatively small spatial scale may reflect

repeated dispersal events in and out of the cave owing to air exchange that accompanies

temperature fluctuations over daily and seasonal periods. Cave size and shape may also

contribute to the exchange of microbial propagules between above ground and subterranean

habitats.

Methanotroph relative abundance and cave air CH_4 concentrations. The relative abundance

of methanotrophs was positively correlated with CH_4 concentrations in cave air. This pattern was

driven by the high-affinity methanotrophs. High-affinity methanotrophs are thought to derive a

significant portion of their energy from CH_4 at atmospheric concentrations and less energy is

available for these organisms at locations in caves with sub-atmospheric CH_4 concentrations.

Some of the variability observed in the relationship between high-affinity methanotrophs and

cave-air CH_4 concentrations may be explained by daily and seasonal variation in cave air flow

patterns. Many caves are known to exhibit faster air flow when the external atmospheric

temperature is lower than the temperature of cave air. This leads to higher CH_4 concentrations in

caves during the winter (10, 45). One result of this is that the measured CH_4 concentration at a

particular point in a cave may or may not be representative of the CH_4 flux at that location, and

the CH_4 flux at a particular site is likely a strong driver of methanotroph relative abundance.

CONCLUSIONS

The structure of cave methanotroph communities appears to be best understood in terms

of how cave methanotrophs access CH_4. Cave methanotrophic communities showed relationships

with cave-air CH_4 concentrations. Additionally, high-affinity methanotrophs were numerically

dominant. This community structure mirrors that of other cave communities, and other

methanotroph communities that appear to consume atmospheric CH_4 (8, 12, 14, 15, 38). High-

affinity methanotrophs also comprised a core methanotrophic community with three members

present together in 79 % of samples. Cave methanotroph communities showed relationships to

minerals in the environment that are known to be related to the abundance of Cu, an essential

trace metal for methane oxidation. These lines of evidence suggest these communities are

actively consuming CH_4 within the cave environment. The cave methanotrophic community

assemblage appears to be actively dispersing across cave environments, suggesting that the

dispersal barrier for high-affinity methanotrophs from cave to cave may be low. Questions

remain about the relative abundance of methanotrophs in cave communities and the CH_4 flux at a

particular site. Further research may be inclined to investigate how USC-γ and USC-α

methanotrophs partition themselves across low CH_4 availability niches.

MATERIALS AND METHODS

Data and software availability. This Targeted Locus Study project has been deposited

at DDBJ/EMBL/GenBank under the accession KCRG00000000. The version described in this

paper is the first version, KCRG00000000.1. All code and data used in this study can be found in

a public GitHub repository (https://github.com/websterkgd/CaveMethanotrophs).

Microbial sampling. We sampled microbial communities from limestone caves along

the western front of the Appalachian fold and thrust belt, in intracratonic settings of the USA,

and in the Sierra Madre of Mexico. We obtained 42 cave soil samples along transects from their

entrances to interiors (Data available at: https://github.com/websterkgd/CaveMethanotrophs).

Cave soil samples were only collected from locations that had accumulations of sediments and

included walls and floors. Samples were collected using a spatula that had been sterilized with 70

volume % ethanol in water from locations that were 0.1 m^2 in area (46). Samples were stored on

ice until they could be transferred to a −80 °C freezer.

Environmental variables. We measured multiple environmental variables to assess

factors that potentially influence the composition of cave methanotrophic communities. We

measured CH_4 concentrations *in-situ* with Fourier Transform Infrared spectroscopy (FTIR)

(Gasmet DX4030 – Milton Keynes, United Kingdom) or we collected discrete air samples

collected in the field for measurement the laboratory with FTIR spectroscopy or gas-

chromatography (Varian – Agilent Technologies, Palo Alto, California). Some of the CH_4

concentrations listed in this study have been previously published (11, 30), and new data were

collected according to methods reported in the same publications. In cases where cave maps

were available, the distance from the nearest entrance to the sampling location was calculated

along the length of the cave passages. We measured cave soil grain-size distributions with a

Malvern Mastersizer 3000 (Malvern Instruments Inc., Westborough Massachusetts). Raw data

from the Mastersizer were converted to % gravel, % sand, % silt, and % clay sized particles by

volume using the GRADISTAT software package (47). Data used for statistical analyses are

available in a public GitHub repository (https://github.com/websterkgd/CaveMethanotrophs).

Latitude and longitude information, however, have been removed from the data to protect the

individual caves. This information can be obtained by contacting the authors directly.

Cave soil mineralogy. We randomly selected 14 samples for mineralogical analysis

using X-ray powder diffraction. Soil sediments were powdered to < 5 μm using a mortar, pestle,

and acetone. Following the methods of Furmann et al. (48), we used Bruker D8 Advance X-ray

diffractometer with a Sol-X solid-state detector and a W X-ray tube operated at 30 mA and 40

kV to identify mineral phases. We placed cave sediment powders in an aluminum large

frontpacked mount. Packed mounts were scanned from 2° to 70° using a count time of 2 s per

0.02° step. Rietveld refinements were used to determine the abundances of the minerals present

in the sample and were quantified with TOPAS software. Proportional abundances of the

minerals from the cave soils are reported in Table 1.

Molecular techniques. We extracted genomic DNA from cave soil samples with a

MoBio PowerSoilTM extraction kit (MoBio, Carlsbad, California USA). About 10 ng of

extracted DNA was used as a template for amplification by polymerase chain reaction (PCR).

The V4 region of the 16S rRNA gene was amplified using 515F and 806R with barcoded primers

designed to work with the Illumina MiSeq platform (49). PCR reactions were as follows: 3 min

denaturing step at 94 °C, followed by 30 cycles of 94 °C for 45 s, 50 °C for 60 s, and 72 °C for

90 s. A final 10-minute extension was carried out at 72 °C. Quality of the PCR products was

checked by gel electrophoresis. Amplified DNA was cleaned using a commercial kit (Beckman

Coulter Agencourt AMPure XP, Indianapolis, Indiana, USA) before quantification using the

QuanIt PicoGreen kit (Invitrogen) and after which aliquots were pooled at a final concentration

of 20 ng per sample.

We sequenced the pooled sample containing PCR products using Illumina MiSeq

technology (Illumina Reagent Kit v2, 500 reaction kit) at the Center for Bioinformatics and

Genomics at Indiana University. Data quality and unique sequences obtained from the PCR

amplifications were analyzed using mothur (50). DNA sequence data were aligned using the

Needleman algorithm and read lengths were limited to 197 base pairs. Sequences matching

chimeras were removed using VSEARCH (51). Sequences that matched chloroplasts, Archaea,

and other non-bacterial sequences were also removed. OTUs were created by binning the data at

97 % sequence similarity using the opticlust algorithm (52). OTUs were identified using the

SILVA reference database (version 132).

Classification of methanotrophs. We identified methanotrophs from the recovered 16S

rRNA sequences using a three step process. First, we compiled a database of organisms known

methanotrophs through literature surveys. This included assembling lists of low-affinity

methanotrophs from the Methylococcaceae, Methylocystaceae, and Beijernickiaceae (34, 53).

We used the genera *Methylacidiphilum*, *Methylacidimicrobium* to identify methanotrophs

belonging to the Verrucomicrobia (34, 54). High-affinity methanotrophs were defined as

organisms belonging to the USC-α and -γ clades, and *Methylocapsa gorgona* MG08 (13, 34,

53). Second, organisms from this database were checked against the identifications assigned to

the OTUs during the initial analysis from the SILVA database (version 132). Third, we compiled

a FASTA file of the 16S rRNA gene of 71 known methanotrophs. 16S rRNA sequences for

USC-α and USC-γ methanotrophs were obtained using published 16S rRNA sequences for USC-

α from (12), for USC-γ from (13), and for *Methylocapsa gorgona* MG08 from (37).. This file

was compared against the FASTA file of all organisms in the dataset using the *usearch_global*

command to identify putative methanotrophs that may have been missed through the SILVA

analysis (usearch v.11.0.667) (55). This list of putative methanotophs was then compared against

the NCBI database. If the sequences did not match an organism capable of the function of

interest at 98 % similarity they were excluded from downstream analysis. The process just

described was intended to prevent the false classification of an organism as a methanotroph.

The compiled methanotroph FASTA file used in our analysis is available in a public GitHub

repository (https://github.com/websterkgd/CaveMethanotrophs).

Statistical analyses. We assessed the relationships between methanotrophs and

environmental conditions through a series of statistical analyses. First, to avoid complication

arising from unavoidable variation in the total number of reads between samples, we calculated

the relative abundance for each taxon within a sample. Next, to quantify among cave variation,

we calculated pairwise dissimilarities using the Bray-Curtis metric. The core methanotroph

community was characterized by including taxa that were found in 60 % or more of our samples.

We used this cutoff because it is in the middle of the range of what has been used in other studies

(23, 24). We evaluated the effect of sediment mineralogy and grain size on the multivariate

composition of methanotroph communities using canonical correspondence analysis (CCA) with

the *vegan* package in R (56). If a sample did not have associated mineralogical data, it was not

included in the CCA analysis.

To gain insight into the biogeography of cave methanotrophs, we examined distance

decay relationships (DDR) while also comparing community composition along gradients from

cave exteriors to cave interiors. We analyzed DDR by plotting Bray-Curtis similarity of

methanotroph composition against environmental and geographic distance calculated as

standardized Euclidean distances. Samples from Cave 38 (Cueva de Villa Luz) were excluded in

DDR analyses since they were roughly 2000 km away from the nearest sample. Slopes for

geographic and environmental DDRs were calculated using ordinary least squares (23). We then

tested for differences in the slopes using the statistical test for a difference of slope between

regression lines (57). We used Spearman's test to examine how methanotrophic communities

changed along gradients from cave exteriors to interiors. Finally, we tested for a relationship

between methanotroph relative abundance and CH_4 concentrations using Spearman's test.

ACKNOWLEDGMENTS

This material is based upon work supported by the U.S. Department of Energy, Office of

Science, Office of Basic Energy Sciences under Award Number DE-SC0006978 to AS.

Additional support was provided by a National Speleological Society research grant awarded to

KDW, the National Geographic Society Expeditions Council (Grant #EC0644-13 to PJB), the

National Science Foundation (1442246 to JTL), the US Army Research Office (W911NF-14-1-

0411 to JTL) and the National Aeronautics and Space Administration (80NSSC20K0618 to

JTL). We thank Brent Lehmkuhl for assistance in the laboratory. Members of the Bloomington

Indiana Grotto facilitated access to caves in Indiana and Kentucky. We thank Rodolfo Gómez

Cruz for his support at Cueva de Villa Luz (Cave 38). ML Larsen, MM Muscarella, and KJ

Locey aided in data analysis and interpretation. KDW thanks A Barberán for discussions on

genetic analysis that resulted in development of the methods used in this paper. KDW thanks the

Planetary Science Institute Mars Laboratory for providing computing resources. The authors

declare no conflicts of interest.

REFERENCES

1. Ciais P, Sabine C, Bala G. 2013. Carbon and other biogeochemical cycles, p. 465–570. *In*
Stocker, TF, Qin, D, Pattner, G-K, Tignor, M, Allen, SK, Boschung, J, Nauels, A, Xia, X,
Bex, V, Midgley, PM (eds.), Climate Change 2013: The Physcial Science Basis.
Contribution of the Intergovernmental Panel on Climate Change. Cambridge University
Press, Cambride, United Kindom and New York, NY, USA.

2. Schimmelmann A, Ensminger SA, Drobniak A, Mastalerz M, Etiope G, Jacobi RD,
Frankenberg C. 2018. Natural geological seepage of hydrocarbon gas in the Appalachian
Basin and Midwest USA in relation to shale tectonic fracturing and past industrial
hydrocarbon production. Sci Total Environ 644:982–993.

3. Kirschke S, Bousquet P, Ciais P, Saunois M, Canadell JG, Dlugokencky EJ, Bergamaschi
P, Bergmann D, Blake DR, Bruhwiler L, Cameron-Smith P, Castaldi S, Chevallier F, Feng
L, Fraser A, Heimann M, Hodson EL, Houweling S, Josse B, Fraser PJ, Krummel PB,
Lamarque J-F, Langenfelds RL, Le Quéré C, Naik V, O'Doherty S, Palmer PI, Pison I,
Plummer D, Poulter B, Prinn RG, Rigby M, Ringeval B, Santini M, Schmidt M, Shindell
DT, Simpson IJ, Spahni R, Steele LP, Strode SA, Sudo K, Szopa S, van der Werf GR,
Voulgarakis A, van Weele M, Weiss RF, Williams JE, Zeng G. 2013. Three decades of
global methane sources and sinks. Nat Geosci 6:813–823.

4. King GM, Nanba K. 2008. Distribution of atmospheric methane oxidation and
methanotrophic communities on hawaiian volcanic deposits and soils. Microbes Environ
23:326–330.

5. Degelmann DM, Borken W, Drake HL, Kolb S. 2010. Different atmospheric methane-
oxidizing communities in european beech and norway spruce soils. Appl Environ
Microbiol 76:3228–3235.

6.   Ford, Derek C; Williams P. 2013. Karst Hydrogeology and Geomorphology. Jon Wiley &
Sons, London, England.

7.   Lennon JT, Nguyễn-Thùy D, Phạm TM, Drobniak A, Tạ PH, Phạm NĐ, Streil T, Webster
KD, Schimmelmann A. 2017. Microbial contributions to subterranean methane sinks.
Geobiology 15:254–258.

8.   Waring CL, Hankin SI, Griffith DWT, Kertesz MA, Kobylski V, Wilson NL, Coleman N
V., Kettlewell G, Zlot R, Bosse M, Bell G. 2017. Seasonal total methane depletion in
limestone caves. Sci Rep 7.

9.   Mattey DP, Fisher R, Atkinson TC, Latin J-P, Durrell R, Ainsworth M, Lowry D,
Fairchild IJ. 2013. Methane in underground air in Gibraltar karst. Earth Planet Sci Lett
374:71–80.

10.   Webster KD, Mirza A, Deli JM, Sauer PE, Schimmelmann A. 2016. Consumption of
atmospheric methane in a limestone cave in Indiana, USA. Chem Geol 443:1–9.

11.   Webster KD, Drobniak A, Etiope G, Mastalerz M, Sauer PE, Schimmelmann A. 2018.
Subterranean karst environments as a global sink for atmospheric methane. Earth Planet
Sci Lett 485:9–18.

12.   Pratscher J, Vollmers J, Wiegand S, Dumont MG, Kaster A-K. 2018. Unravelling the
identity, metabolic potential and global biogeography of the atmospheric methane-
oxidizing upland soil cluster α. Environ Microbiol https://doi.org/10.1111/1462-
2920.14036.

13.   Edwards CR, Onstott TC, Miller JM, Wiggins JB, Wang W, Lee CK, Cary SC, Pointing
SB, Lau MCY. 2017. Draft genome sequence of uncultured upland soil cluster
Gammaproteobacteria gives molecular insights into high-affinity methanotrophy.
Genome Announc 5:e00047-17.

14.   Zhao R, Wang H, Cheng X, Yun Y, Qiu X. 2018. Upland soil cluster γ dominates the
methanotroph communities in the karst Heshang Cave. FEMS Microbiol Ecol 94:1–13.

15.   Cheng X, Liu X, Lu X, Wang H, Su C, Zhao R, Bodelier PLE, Wang W. 2021. USCγ
Dominated Community Composition and Cooccurrence Network of Methanotrophs and
Bacteria in Subterranean Karst. Microbiol Spectr 9:e00820-21.

16.   Knief C, Dunfield PF. 2005. Response and adaptation of different methanotrophic bacteria
to low methane mixing ratios. Environ Microbiol 7:1307–1317.

17.   Bowman JP. 2005. Methylococcales *ord. nov.*, p. 248–270. *In* Brenner, D, Krieg, N,
Staley, J, Garrity, G, Boone, D, De Vos, P, Goodfellow, M, Rainey, F, Schleifer, K-H
(eds.), Bergey's Manual of Systematic Bacteriology. Springer, Boston, MA.

18. Mills CT, Slater GF, Dias RF, Carr SA, Reddy CM, Schmidt R, Mandernack KW. 2013. The relative contribution of methanotrophs to microbial communities and carbon cycling in soil overlying a coal-bed methane seep. FEMS Microbiol Ecol 84:474–494.

19. He R, Wooller MJ, Pohlman JW, Quensen J, Tiedje JM, Leigh MB. 2012. Shifts in identity and activity of methanotrophs in Arctic Lake sediments in response to temperature changes. Appl Environ Microbiol 78:4715–4723.

20. Krause S, Meima-Franke M, Hefting MM, Bodelier PLE. 2013. Spatial patterns of methanotrophic communities along a hydrological gradient in a riparian wetland. FEMS Microbiol Ecol 86:59–70.

21. Gordon MS, Rosen DE. 1962. A cavernicolous form of the poeciliid fish *Poecilia sphenops* from Tabasco, Mexico. Copeia 1962:360–368.

22. Howarth FG. 1993. High-stress subterranean habitats and evolutionary change in cave-inhabing arthropods. Am Nat 142:S65–S77.

23. Björk JR, O'Hara RB, Ribes M, Coma R, Björk JR, Montoya JM. 2017. The dynamic core microbiome: Structure, dynamics and stability. bioRxiv 13.

24. Hernandez-Agreda A, Gates RD, Ainsworth TD. 2017. Defining the Core Microbiome in Corals' Microbial Soup. Trends Microbiol 25:125–140.

25. Nekola JC, White PS. 1999. The distance decay of similarity in biogeography and ecology. J Biogeogr 26:867–878.

26. Locey KJ, Muscarella ME, Larsen ML, Bray SR, Jones SE, Lennon JT. 2020. Dormancy dampens the microbial distance-decay relationship. Philos Trans R Soc B Biol Sci 375.

27. Kolb S. 2009. The quest for atmospheric methane oxidizers in forest soils. Environ Microbiol Rep 1:336–346.

28. Lau MCY, Stackhouse BT, Layton AC, Chauhan A, Vishnivetskaya TA, Chourey K, Ronholm J, Mykytczuk NCS, Bennett PC, Lamarche-Gagnon G, Burton N, Pollard WH, Omelon CR, Medvigy DM, Hettich RL, Pfiffner SM, Whyte LG, Onstott TC. 2015. An active atmospheric methane sink in high Arctic mineral cryosols. ISME J 9:1880–1891.

29. Fernandez-Cortes A, Cuezva S, Alvarez-Gallego M, Garcia-Anton E, Pla C, Benavente D, Jurado V, Saiz-Jimenez C, Sanchez-Moral S. 2015. Subterranean atmospheres may act as daily methane sinks. Nat Commun 6:7003.

30. Webster KD, Rosales Lagarde L, Sauer PE, Schimmelmann A, Lennon JT, Boston PJ. 2017. Isotopic evidence for the migration of thermogenic methane into a sulfidic cave, Cueva de Villa Luz, Tabasco, Mexico. J Cave Karst Stud 79:24–34.

31. Levy DB. 2007. Oxidation-reduction chemistry of Lechuguilla Cave seepage. J Cave
Karst Stud 69:351–358.

32. Laiz L, Groth I, Gonzalez I, Saiz-Jimenez C. 1999. Microbiological study of the dripping
water in Altamira cave (Santillana del Mar, Spain). Jounal Microbiol Methods 36:129–
138.

33. Barberán A, Ladau J, Leff JW, Pollard KS, Menninger HL, Dunn RR, Fierer N. 2015.
Continental-scale distributions of dust-associated bacteria and fungi. Proc Natl Acad Sci
112:5756–5761.

34. Knief C. 2015. Diversity and habitat preferences of cultivated and uncultivated aerobic
methanotrophic bacteria evaluated based on pmoA as molecular marker. Front Microbiol
6.

35. Wood AP, Aurikko JP, Kelly DP. 2004. A challenge for 21st century molecular biology
and biochemistry: What are the causes of obligate autotrophy and methanotrophy? FEMS
Microbiol Rev 28:335–352.

36. Sriswasdi S, Yang C, Iwasaki W. 2017. Generalist species drive microbial dispersion and
evolution. Nat Commun 8.

37. Tveit AT, Hestnes AG, Robinson SL, Schintlmeister A, Dedysh SN, Jehmlich N, von
Bergen M, Herbold C, Wagner M, Richter A, Svenning MM. 2019. Widespread soil
bacterium that oxidizes atmospheric methane. Proc Natl Acad Sci U S A 116:8515–8524.

38. Cai Y, Zhou X, Shi L, Jia Z. 2020. Atmospheric Methane Oxidizers Are Dominated by
Upland Soil Cluster Alpha in 20 Forest Soils of China. Microb Ecol 80:859–871.

39. Müller D, Groves DI. 2019. Tectonic Settings of Potassic Igneous Rocks, p. 31–71. *In*
Mineral Resource Reviews. Springer International Publishing AG.

40. Bender M, Conrad R. 1994. Methane oxidation activity in various soils and freshwater
sediments: Occurrence, characteristics, vertical profiles, and distribution on grain size
fractions. J Geophys Res 99:16531–16540.

41. Jensen S, Øvreås L, Daae FL, Torsvik V. 1998. Diversity in methane enrichments from
agricultural soil revealed by DGGE separation of PCR amplified 16S rDNA fragments.
FEMS Microbiol Ecol 26:17–26.

42. Conrad R. 1995. Soil microbial processes involved in production and consumption of
atmospheric trace gases, p. 207–250. *In* Advances in Microbial Ecology.

43. Tornero I, Boix D, Bagella S, Pinto-Cruz C, Caria MC, Belo A, Lumbreras A, Sala J,
Compte J, Gascón S. 2018. Dispersal mode and spatial extent influence distance-decay
patterns in pond metacommunities. PLoS One 13:e0203119.

44.   Vellend M. 2010. Synthesis in community Ecology. Q Rev Biol 85:183–206.

45.   Gregorič A, Zidanšek A, Vaupotič J. 2011. Dependence of radon levels in Postojna Cave
537   on outside air temperature. Nat Hazards Earth Syst Sci 11:1523–1528.

538   46.   Pašić L, Kovče B, Sket B, Herzog-Velikonja B. 2010. Diversity of microbial communities
539   colonizing the walls of a Karstic cave in Slovenia. FEMS Microbiol Ecol 71:50–60.

540   47.   Blott SJ, Pye K. 2001. Gradistat: A grain size Ddstribution and statistics package for the
541   analysis of unconcolidated sediments. Earth Surf Process Landforms 26:1237–1248.

542   48.   Furmann A, Mastalerz M, Schimmelmann A, Pedersen PK, Bish D. 2014. Relationships
543   between porosity, organic matter, and mineral matter in mature organic-rich marine
544   mudstones of the Belle Fourche and Second White Specks formations in Alberta, Canada.
545   Mar Pet Geol 54:65–81.

546   49.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM,
547   Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012. Ultra-
548   high-throughput microbial community analysis on the Illumina HiSeq and MiSeq
549   platforms. ISME J 6:1621–1624.

550   50.   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,
551   Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ,
552   Weber CF. 2009. Introducing mothur: Open-source, platform-independent, community-
553   supported software for describing and comparing microbial communities. Appl Environ
554   Microbiol 75:7537–7541.

555   51.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: A versatile open
556   source tool for metagenomics. PeerJ 2016:4:e2584.

557   52.   Westcott SL, Schloss PD. 2017. OptiClust, an Improved Method for Assigning Amplicon-
558   Based Sequence Data to Operational Taxonomic Units. mSphere 2.

559   53.   Nazaries L, Murrell JC, Millard P, Baggs L, Singh BK. 2013. Methane, microbes and
560   models: fundamental understanding of the soil methane cycle for future predictions.
561   Environ Microbiol 15:2395–2417.

562   54.   Carere CR, Hards K, Houghton KM, Power JF, McDonald B, Collet C, Gapes DJ,
563   Sparling R, Boyd ES, Cook GM, Greening C, Stott MB. 2017. Mixotrophy drives niche
564   expansion of verrucomicrobial methanotrophs. ISME J 11:2599–2610.

565   55.   Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST.
566   Bioinformatics 26:2460–2461.

56.   Oksanen, J.; Blanchet, F.; Kindt, R.; Legendere, P.; Minchin, P.R.; O'Hara, R.B.O.;
Simpson, G.L; Solymos, P.; Stevens, H.H.; Wagner H. 2018. Vegan: Community ecology
package. R package version v 2.4-6.

57.   Paternoster R, Brame R, Mazerolle P, Piquero A. 1998. Using the correct statistical test
for the equality of regression coefficients. Criminology 36:859–866.

58.   Weary DJ, Doctor DH. 2014. Karst in the United States: a digital map compilation and
database. United States Geol Surv Open-File Rep 1156:1–23.

Figure 1: (A) The distributions of karst landscapes at global scale. Inserts show the occurrence

of karst (B) in the eastern United States (C) and in Mexico. Numbers represent the locations of

sampled caves in this study. Karst land cover data were obtained from (6, 58).

Figure 2: The relative abundance of A) USC-*γ* methanotrophs, B) USC-*α* methanotrophs, and C)

Low-affinity methanotrophs in the sampled caves.

Figure 3: The total methanotrophic community showed relationships with the abundance of

muscovite, clinochlorite, and microcline detected in the samples. The vectors show increasing

abundances of minerals the samples.

Figure 4: Distance decay analyses of the high affinity methanotrophs. The change in

community similarity with geographic distance was statistically indistinguishable from the

change in environmental similarity with distance suggesting the organisms are adept at

dispersing through the environment. Note that the samples from Cave 38 have been removed

from this analysis due to the large geographic separation of these samples from the rest of the

samples.

Figure 5: The relative abundance of members of the methanotrophic community plotted against

the CH$_4$ concentration at each sample location ($\rho = 0.40$, S-statistic: S = 5451, $p < 0.01$).

Table 1: The proportional abundance of minerals from the sampled locations

Sample	Cave	Quartz	Muscovite	Clinochlorite	Albite	Orthoclase	Calcite	Microcline	Dolomite	Anorthite	Gypsum
21-10d	Cave 21	0.72	0.14	0.01	0.06	0.04	0.00	0.00	0.00	0.03	0.00
22-11d	Cave 22	0.65	0.16	0.01	0.07	0.00	0.01	0.11	0.00	0.00	0.00
17-1b	Cave 17	0.48	0.28	0.01	0.00	0.00	0.23	0.00	0.00	0.00	0.00
17-1c	Cave 17	0.89	0.09	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00
27-5b	Cave 27	0.33	0.19	0.01	0.06	0.00	0.07	0.28	0.05	0.00	0.00
27-5d	Cave 27	0.50	0.15	0.01	0.00	0.00	0.16	0.14	0.04	0.00	0.00
30-8d	Cave 30	0.40	0.13	0.00	0.02	0.00	0.19	0.09	0.18	0.00	0.00
31-9b	Cave 31	0.40	0.20	0.01	0.00	0.00	0.05	0.16	0.19	0.00	0.00
31-9c	Cave 31	0.14	0.20	0.00	0.00	0.00	0.06	0.08	0.52	0.00	0.00
35-1	Cave 35	0.70	0.09	0.00	0.08	0.04	0.00	0.08	0.00	0.00	0.00
38-9	Cave 38	0.38	0.29	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.33
10-1	Cave 10	0.76	0.11	0.00	0.06	0.00	0.00	0.06	0.01	0.00	0.00
4-1	Cave 4	0.21	0.24	0.00	0.00	0.00	0.55	0.00	0.00	0.00	0.00
2-3	Cave 2	0.82	0.11	0.00	0.00	0.00	0.00	0.07	0.00	0.00	0.00
2-2	Cave 2	0.75	0.12	0.00	0.04	0.00	0.02	0.07	0.00	0.00	0.00
2-1	Cave 2	0.74	0.10	0.00	0.06	0.00	0.00	0.10	0.00	0.00	0.00