A tale of two species: human and peafowl interactions in human dominated landscape

influence each other's behaviour

Wild life near human habitation

Dhanashree Paranjpe* and Priyanka Dange

Annasaheb Kulkarni Department of Biodiversity, MES Abasaheb Garware College, Karve

Road, Pune-411004.

*Corresponding author

E-mail: dhana4shree@gmail.com

Telephone No : +91-20-41038237

ORCID ID*: https://orcid.org/0000-0003-4787-2052

Priyanka Dange: ORCID ID: http://orcid.org/0000-0002-2799-2156

Abstract:

Wild life is increasingly coming in contact with humans in many parts of the world. Human perception of wild life may be an important factor in population management and conservation practices in urban and semi-urban areas. Human attitude towards bird species may vary from peaceful coexistence to a perception of birds as pests. Based on the data collected from survey interviews, we studied the perceptions of local communities in Rajasthan, India towards Indian Peafowl. Local communities in Rajasthan reported Indian Peafowl as crop pest and this perception varied across seasons. The crop loss incurred due to Indian Peafowl also varied across seasons according to the respondents. Despite reporting peafowls as a crop pest, locals regularly offered grains for them as a traditional practice. Thus, at our study sites locals have mostly positive perception about Indian peafowl around them.

Food provisioning by local human population influenced diet composition and time-budget of Indian Peafowls at food provision and non-provision sites. Sites at which food provisioning was less or absent, peafowl spent more time in walking in search for food and more than half of their diet consisted of natural food. In contrast, the sites at which plenty of grains were available, time spent in walking was significantly less, while time spent in feeding was significantly more; and over 70% of their diet consisted of carbohydrate and protein-rich provisioned grains. Food provisioning changed the benefit: cost ratio (measured as time spent in feeding to time spent in all other behaviours) between provision and non-provision sites. Thus, food provisioning by humans can change feeding ecology of native Indian peafowl populations, while the presence of peafowl in human dominated landscape changes how humans interact with wild life around them.

Introduction:

Wild animals are increasingly finding themselves near human habitation due to

scarcity of natural habitats, high human population density and changing land usage

(Ayyappan et al. 2016; Pathak et al. 2002; Vijayan and Pati, 2001; Meena 2010). As a result,

human-animal interactions and conflicts have risen. The impact of wild animals on local

human population has mostly been studied in the context of human-wildlife conflicts

(Agrawal et al. 2016) while, impact of local human population on animal populations living

close to or within rural, semi-urban, and urban areas has rarely been studied.

There are potentially diverse types of interactions between humans and wildlife. The

impact of human induced ecological stressors on animal populations in human dominated

landscapes is becoming an important area of research (Sih et al. 2011; Agrawal et al. 2016).

Some studies predict large scale extinctions of Earth's species in next 50-100 years due to

human induced rapid environmental change (HIREC) (Tilamn et al. 1994; Stork 2010); while

many other species (e.g. urbanised pests and invasive species) seem to be doing better than

ever in face of the same environmental changes. Wild animals can also influence local human

population socially, economically through crop loss, property damage, life loss, ecosystem

services and aesthetic/ religious value. They can become sources of income through

ecotourism, trade of animal derived products and so on.

Vicinity to human population can change natural behaviour and life history of animals

in multiple ways. Human habitation or agricultural landscapes can serve as safer refuge areas

or even new habitats for wildlife (Horgan et al. 2017). Humans often provide food to wildlife

voluntarily around their homes or in temples as a ritual or in wild life tourism spots/ feeding

stations (Robb et al. 2008; Sengupta et al. 2015). In addition to this, indirect food

provisioning in the form of crops, ornamental plants, livestock or waste food thrown in open

becomes easy and reliable source of food for wild life. Food provided by humans tends to be

calorie rich, easily digestible, available at predictable times and places. It is known that food

provisioning might change feeding habits, diet preference or diet composition of wild

mammals (Robb et al. 2008; Sih et al. 2011; Sengupta et al. 2015; Ayyappan et al. 2016;

Sengupta and Radhakrishna 2017). Reduced predatory pressure and regular availability of

nutrient rich food (in the form of crops) round the year are likely to provide greater resilience

and adoptability of wild animals. It also allows them to live successfully close to agricultural

landscape and human habitation (Gilroy and Sutherland 2007; Sih et al. 2011; Ayyappan et

al. 2016).

There are many avian species like crows, sparrows, cranes, egrets that inhabit human-

dominated landscapes or stay very close to / within human habitation, however, they are not

typically considered as pests. Appreciation for birds and even individual species may vary

from peaceful coexistence (including feeding them) to a perception of birds as pests

(Clergeau et al. 1996). Very little work has been done on how human perception/ attitude

towards avian species can influence the outcome of their interactions with humans. A multi-

dimensional approach to study such interactions is necessary in order to understand the

capacity of local stakeholders to accept wild species (Carpenter et al., 2000) as well as the

potential effect of these interactions on both humans and wildlife who share common spaces.

Indian peafowl (*Pavo cristatus*) is a species that has co-habited human dominated

landscapes for centuries in its native geographical range. This avian species is native to

Indian subcontinent and has been introduced in many parts of the world relatively recently.

Although, it's native habitat is undergrowth in open forest and woodlands near a waterbody,

it is also known to occur near farmlands, villages and increasingly becoming common in

urban and semi-urban areas (Burton and Burton 2002). In some places, such as Morachi

Chincholi in Maharashtra, India, "eco-tourism" based on Indian peafowl is flourishing and it

is an additional or alternative source of income for local community. In contrast to this

"positive" impact, anecdotal reports suggest that the Indian peafowl can cause substantial

crop losses in the areas where their population density is high. There are no known studies

from India that estimate the crop losses due to peafowls. It is not known whether peafowl

really affect crops like other known crop pests (e.g. wild boar, elephants, deer, etc.).

Numerous reports can be found in newspapers in India about deaths of individuals or groups

of peacocks. The deaths may be due to natural reasons such as predation, water scarcity or

may be the result of unintentional poisoning due to insecticides/ pesticides sprayed on crops

that the peafowl later feed on. Sometimes intentional killing of Indian peafowl by certain

tribal communities for their ornamental feathers and meat has also been suspected/ reported.

In contrast to this, in some parts of India (e.g. State of Rajasthan), peafowl are believed to be

sacred and people actively offer grains for them as part of their daily ritual. These varied

interactions between Indian peafowl and local human population make it an interesting study

system to understand consequences of human- wild life interactions. It remains to be seen

how common are the positive perceptions and associated beliefs/ rituals throughout the region

in which the population density of Indian peafowl is higher. Can these perceptions result in

curbing population decline or effective management/ conservation of the species? To address

these questions we studied three aspects of human- peafowl interactions in details. The

objectives of this study were: (i) To understand the perceptions of local community about

Indian peafowl, (ii) To estimate impact of Indian peafowl populations on local agriculture

and (iii) To estimate the impact of food provisioning by local community on peafowl

populations.

Materials and methods:

We used questionnaire based interviews to estimate impact of Indian peafowl

populations on local agriculture and to understand perceptions of local people about Indian

peafowl in four villages located on the periphery of Ranthambhore Tiger Reserve (RTR), in

the state of Rajasthan, India (N = 203). Interviews were conducted by taking family as a unit

respondent. The questionnaire included fixed and open-ended questions. Interviews started

with the interviewer clearly explaining objectives of the study and the interview was

continued only if the respondent(s) gave consent. The questionnaire consisted of 3 parts: (i)

basic information about the respondent such as name, age, address, number of people in

family unit, source(s) of income (agriculture/ others). (ii) Questions related to perceived

impact of peafowls on agriculture (iii) questions related to social/ cultural/ traditional

perceptions about Indian peafowl. Interviews were conducted in three different seasons:

November 2016 (harvest time for kharif crops, winter season, N = 55), March-April 2017

(Harvest time for rabbi crops, beginning of dry summer season, N = 86), July-August 2017

(standing crops, monsoon season, N = 55) to check if responses to some of the questions

related to agriculture differed according to stage of crop/ season.

To estimate the impact of food provisioning by local community on peafowl

populations, we selected three field sites- Morachi Chincholi, Nashik and Rajasthan (Table 1).

Selection of study areas was based on several criteria such as their closeness to human

habitation, accessibility throughout the year, history of peafowl populations in the area,

potential interactions between humans and peafowl (food provisioning, peafowl visiting/

roosting on agricultural land and village homes, sell of peafowl feathers or related "products",

tourism, cultural significance), etc. Areas located outside protected areas and close to human

habitation were given preference. Behaviours of the Indian peafowl were studied throughout

the year using direct field observations during morning (6 am to 10 am) and evening (4 pm to

7 pm) at various field sites. Behaviours were also observed in the late- afternoons if peafowl

were found to be active. The behavioural observations of peafowl were noted while patrolling

the study area using continuous scan and focal animal sampling. Focal animals were chosen

randomly from a group (if the peafowl were seen in a group). Additionally, opportunistic

observations were recorded for individuals sighted during the walk around the study sites.

Behaviours of focal animal were recorded continuously without disturbing or going too close

to focal animal (typically from a distance of 10-100m) until they were out of the sight. During

feeding bouts, location of feeding was recorded. Gender of the focal animal was noted along

with the date, place and the total duration of observations. Season of observation was noted as

Monsoon, pre-monsoon and post-monsoon. Arrival of Monsoon rains marks a remarkable

transition of seasons on Indian subcontinent. Monsoon season is associated with greater

availability of water, food and sowing of Kharif crops. Monsoon season for Morachi Chincholi

and Nashik sites is June to September while for Rajasthan it is July through September as

monsoon rains start in Rajasthan later than the other two sites. Post monsoon season is October

through January, while pre-monsoon season is February through May for Morachi Chincholi

and Nashik, February to June for Rajasthan. Data were also collected about the type of food

the peafowl ate during the observations. Observations were stopped when the focal animal

flew/ walked out of observer's sight.

For calculating time budget of the Indian peafowl populations, the observer(s) documented

behaviours of peafowl using handheld digital audio recorder (SONY ICD-UX560F).

Occasionally, data were recorded as videos using a Panasonic camcorder (Full HD, 29.8mm

wide 20.4 megapixel HC-X920) if a large group of peafowl was found near feeding site.

Data processing

The types of food items peafowls ate were listed using direct observations (Table 2).

Cereals, pulses, vegetables and processed food is available to peafowl only around human

habitation and hence was considered as provisioned food, while all other food items such as

grass, leaves, sprouts, fruits, seeds, insects, worms were considered as natural food items (Table

2). Based on direct observations of peafowl eating various food items, proportion of natural

and provisioned food items in peafowl diet was calculated. The food items were further

categorised as cereals (grains such as sorghum, wheat, bajra, rice, corn, etc.), pulses (split

pigeon peas, black gram lentils, black-eyed pea, Indian brown lentils, green gram, peanuts,

etc.), non-grains (fruits, grass, seeds, leaves, insects, vegetables, etc.) and processed food.

Number of times peafowls were seen eating each food type was counted. Based on this

frequency, the percentage of each food type in their diet was calculated.

The ethogram of the peafowl comprised of following behaviours: feeding,

vigilant/alert, walking, preening, scare, scrape, aggressive, jump, sit, flap, shake, call, display,

turn, stand, perch, scratch, pause, allo-grooming, head shake and fly. For calculating time

budget of the Indian peafowl populations, data about the duration of various activities was

extracted for each recorded individual from audio records. Number of pecks recorded during

the feeding bouts was used for estimating how much food was eaten, while timed audio records

of observations were used to calculate time spent in various behavioural states. Fraction of time

spent in each activity was summed across all individuals in the population. Time budget of the

whole population was calculated as time spent in each activity divided by total time of

observation for the population. Time budget was represented as % of time spent in three major

activities by individuals in the population. Rest of the behaviours (<10% of total time budget)

were pooled together as "other behaviours".

For calculating benefit : cost ratio, time spent in feeding was considered as "benefit"

while time spent in all other activities (except standing inactive) was considered as "cost".

Statistical analysis

For statistical analysis, STATISTICA^{TM} software version 13.2 (Dell Inc. 2016) was used. As the data did not follow normal distribution, non-parametric tests were used for further analysis. Kruskal-Wallis ANOVA was used to see the effect of season (Pre-monsoon, monsoon, post-monsoon), field sites (Morachi Chincholi, Nashik, Rajasthan) and sex (male, female) on time spent in feeding, time spent in vigilance, walking and other behaviours. Mann-Whitney U test was used to estimate the effect of proximity to food provision site on time spent in feeding, vigilance, walking and other behaviours.

Benefit: cost ratio was compared across three field sites (Morachi Chincholi, Nashik, Rajasthan), across seasons (Pre-monsoon, monsoon, post-monsoon) and between food provision and non-provision sites using non-parametric tests.

Results

Perceptions about Indian peafowl and their relation to agriculture

The age of survey respondents varied from 15- 65yrs, all age classes in this range were represented almost equally. Respondents included 90.73% males and 9.27% females. Sources of income for the respondents included agriculture (51.32% of respondents), animal herding (17.88%), labour (farm or other labour 20.2%) and other sources (10. 6%) such as contractor, company job, mechanic, shop owner, artisan, teacher etc. A majority of farmers in the area around RTR are small farmers with 84.11% farmers owning less than 5 acres of land, while rest of the farmers (15.89%) owned 5+ acres to 20+ acres of land. Farmers take two to sixteen types of crops in a year depending on size of farm, availability of labour, water, etc. Most common crops in this region include wheat, bajra, chilly, mustard, sorghum, sesame. All farmers in this region reported crop losses irrespective of time of year during which survey

was conducted. The reasons for crop loss differed according to season during which

interviews were conducted. In November 2016 survey, 100% of respondents reported crop

loss due to crop pests only. During survey conducted in March 2017, 59.82% respondents

reported loss due to crop pests, 31.25% reported loss due to untimely rains, while 8.93% said

there was crop loss due to disease. In the months of July and August, 62.34% respondents

said that the reason for crop loss was crop pests, 36.37% respondents reported crop loss due

to untimely rains, while 1.3% respondents said that the loss was due to crop disease. Major

crop pests reported in the survey are Wild boar (*Sus scrofa*), Nilgai/ Blue bull (*Boselaphus*

tragocamelus), Indian peafowl (*Pavo cristatus*) and Deer. However, number of respondents

reporting particular types of crop pests changed according to season in which survey was

conducted (Fig 1). Although Indian peafowl is the third common crop pest reported, only

2.84% of respondents in November 2016 survey said it was a crop pest compared to 15.16%

respondents in March 2017 and 19.7% respondents in July 2017. On the other hand wild boar

and Nilgai were reported as crop pests throughout the year by 19-28% respondents.

When asked specifically if they think that Indian peafowls affect crops, 87.63-

95.45% responders replied positive during the year around survey. According to respondents,

peafowl eat harvest (41.94- 43.27%), leaves (6.73- 23.66%), insects (16.58- 21.51%), flowers

(0-16.35%), seeds (0-2.15%), saplings (0.52- 8. 65%), and drink water (0-10.36%) when they

visit farms. Only 0.96% respondents said they spread disease, while 0-2.88% respondents

said that the peafowl does no harm to the crops. Between 67.65- 100% respondents reported

that the long train of feathers causes some damage to crops when the males with full grown

train walk through the standing crops. Estimates of total loss due to peafowl varied between

0% to more than 20% according to season in which the survey was conducted (Fig 2).

Chickpea, chilli, wheat and mustard were the major crops that incurred loss due to peafowl.

Loss was also reported occasionally for coriander, sorghum, sesame, tomatoes, green peas

and moong (all reported by 10% or less respondents). Indian peafowl ate crops at post growth

(around harvest) stage according to 60-95% of respondents, next most vulnerable stage was

pre-growth stage (freshly sown, sprouted or sapling stage) according to 2.5-29.27% of

responses. Damage to crops was reported least in the growth stage (0-9.76% respondents, Fig

3). Between 64.65%-100% farmers reported that they changed crop pattern to reduce the loss

due to peafowl. Many farmers chose to take a different variety of crop that is less preferred

by peafowl (e.g. hybrid chilli instead of indigenous variety) or avoided taking certain crops

(e.g. chickpea, ground nut, chilli) or change the place where they take the crop if they own

patches of farm lands in more than one locations within the village. Interestingly, the estimate

of loss (estimated % loss) due to Indian peafowl showed positive correlation with size of

farm (Spearman rank order correlation $r = 0.24$, $p < 0.05$). Farmers with more farm land

estimated more crop losses due to peafowl compared to farmers who owned less land.

Food provisioning for Indian peafowl

In spite of the crop damage the peafowl cause, local people offer grains to peafowl

throughout the year. All households (100% respondents) in the village offer grains to peafowl

except in the month of November slightly less number of respondents (94.87%) reported

giving grains for peafowl. Most common grains offered to peafowl include bajra (35-45%

respondents), sorghum (27.5-41.25% respondents), wheat (10-25.95% respondents), and

lentils (0-7.57% respondents) while some people also offer rice, maize, sesame when

available. People typically offer a handful (100 – 130 g depending on type of grains offered)

to up to 3 Kg grains at a time. In some villages, grains are offered from each household every

day, in some cases they are offered in temple once or twice a week or on special days/

occasions. So year around there are on average ~15 Kg grains available for wild life every

day in and around temple premises in RTR region.

Impact of food provisioning on peafowl populations

Impact of food provisioning was studied in three study areas- Morachi Chincholi,

Nashik and Rajasthan (RTR). Diet composition of the peafowl was different in the selected

study areas. In Morachi Chincholi, peafowl diet consisted of 71% food provisioned by humans

compared to 61% provisioned food in Rajasthan, while just 40% peafowl diet in Nashik

consisted of provisioned food (Fig 4). Peafowl in Nashik ate almost double the amount of

natural (non-grain) food items (60%) compared to peafowl in Morachi Chincholi (29%). Major

portion of food provided to peafowl populations in Morachi Chincholi and Rajasthan consisted

of grains (65% and 58%, respectively). Peafowl in Rajasthan were not seen eating processed

food as opposed to peafowl in Morachi Chincholi and Nashik study area (Fig 5).

There was lot of variation in the total duration of observations for each individual

(Range: 1 min-29 min.), however, total observation time did not differ significantly across field

sites (Median Test, *Chi-Square* = 1.75, *df* = 2, *p* = 0.42) which confirms that equal efforts were

put into observations at the selected field sites. However, frequency of sighting males was

significantly greater compared to frequency of sighting females (Z = -3.06 p = .002). The most

frequent behaviours observed (N= 304) were feeding, vigilance and walking. Less frequent

behaviours (typically shown less than 10% of total observation time) were categorized together

as "other behaviours". The relative proportion of time spent in different behaviours at food

provision sites (where humans put grains for peafowl) and away from provision site (non-

provision sites) is shown in Figure 6. Time spent in vigilance (23.06%) and other behaviours

(9.28%) at provision sites was comparable to time spent in those behaviours at non-

provisioning sites (29.66%, and 12.51%, respectively) (Fig. 6). At the provision sites,

individuals spent significantly less time in walking (22.89%) compared to non-provision sites

(32.35%, Mann- Whitney U test, *p*= 0.003) (Fig. 7). Peafowls spent significantly more time in

feeding (49.30%) at the provision sites as compared to non-provision sites (35.24%, Mann-

Whitney U test, $p = 0.0003$) (Fig 7). These time budget trends were similar across all three study

sites.

Benefit: cost ratio measured in terms of time spent in feeding (benefit) versus other

behaviours (cost) was significantly different between food provision and non-provision sites

(Mann-Whitney U test, $p = 0.00015$). The ratio was similar across field sites (Kruskal-Wallis

ANOVA, $p = 0.22$) and marginally different across pre-monsoon, monsoon and post-monsoon

season (Kruskal-Wallis ANOVA, $p = 0.06$).

Discussion:

Human- wild life interactions have mostly been studied so far in the context of human-

animal conflict which has drastic consequences such as loss of property, crops or even death.

All human- wild life interactions do not necessarily lead to obvious conflicts, nonetheless,

they might impact both sides in significant ways. Peoples' perception about wild life can

potentially influence the long-term outcome of such interactions. Our study investigated in

detail some aspects of non-violent human- wildlife interactions by focusing on Indian

peafowl populations living in human dominated landscape. Our survey covered the

perceptions about Indian Peafowl in a wide range of age groups (15-65 yrs) in a geographic

area with high population density of peafowls. Most of the surveys were conducted in remote

rural areas of Rajasthan by volunteers going door-to-door for collecting data. Survey was

conducted considering family as a unit respondent. In most cases, head of the family along

with one or two other members (male/ female) present in the household answered the

questions. Therefore, number of female respondents in our survey remained below 10%.

However, looking at the responses, the unbalanced gender ratio of respondents may not have

much bearing on the results of the survey.

Most of the respondents in the survey were small farmers with less than five acres of land.

The source(s) of income, in most cases (89.4% respondents) were related to agriculture

(crops + farm animals + labour).

Perceptions about Indian peafowl and their relation to agriculture

We did not want to bias the respondents by directly asking them if peafowl damaged

their crops. Hence, we asked questions step-by-step about types of crops and typical reasons

for crop loss in RTR area. Losses due to erratic rain patterns (deficient rainfall, wet drought,

untimely rains) is the main reason for crop losses for Indian agriculture which depends

largely on Monsoon rains from Indian ocean. Surprisingly, damage due to crop pests emerged

as the top most reason across all seasons for loss to farmers in RTR area, followed by loss

due to untimely rains, crop disease, or any other reasons. The area in which we conducted

surveys includes buffer zone and villages just outside the Ranthambhore National park. As a

result, wild herbivores from the protected area frequently enter the villages, farmlands and

surrounding areas. This can create serious problems for the farmers whose livelihood depends

on agricultural produce. Similar issues of crop raiding by wild herbivores have been reported

all over the world (Naughton-Treves 1997; Treves et al. 2006; Graham et al. 2010; Mathur et

al. 2015; Perera and Vandercone 2017) including India (Chauhan and Singh, 1990; Chauhan

et al. 2009; Ramkumar et al. 2014; Meena 2017). Wild boar (*Sus scrofa*), Nilgai/ blue bull

(*Boselaphus tragocamelus*), Indian peafowl (*Pavo cristatus*) and deer were reported as the

major crop pests in RTR area. While wild boar and blue bull are some of the commonly

known crop raiders in India, ours is first systematic survey in which Indian peafowl has been

reported as a crop pest in India. There are sporadic newspaper reports and anecdotal

information from different parts of India about Indian peafowl damaging crops, however,

there has been no systematic survey or documentation of extent and type of crop damage

done by Indian peafowl as we have done in this survey. Recently, Indian peafowl was

reported as a crop pest by farmers in Sri Lanka (Horgan et al. 2017; Perera and Vandercone

2017). One study reported that Maavee paddies and surrounding habitats offer refuges for

wildlife including Indian peafowl; and Indian peafowl damaged rice at the seedling and grain

filling stage irrespective of the farming method used (Horgan et al. 2017).

A majority of respondents (87.63-95.45%) in our survey reported that peafowl do

affect crops in an adverse way. So Indian peafowl is not only visiting their farms seeking

refuge or using farms as new feeding habitat, in most cases they do have perceivable negative

effects on crops. Although they were not reported to spread disease (only 0.96% respondents

think they spread crop disease), their long train was found to cause some damage as they

walked through the standing crop. They also eat various parts and stages of the crop.

Estimates of total crop loss due to Indian peafowl ranged from negligible (0%) to over 20%.

Typically, figures between 5-20% losses were mentioned by respondents for various crops. It

is important to note that these are "perceived" estimates of loss given by farmers and the

actual figures need to be verified through more detailed 'on field' data collection. Other wild

herbivores such as elephants can cause up to 35% crop damage (Ramakumar et al. 2014),

while wild boars can destroy 5-36% of crops (Chauhan et al. 2009). Blue bull (*Boselaphus*

tragocamelus) seems to have the highest impact with 58% losses (rarely less than 10%) in

Harayana, India (Chauhan and Singh 1990). The losses due to blue bull can be as high as

70% in high density areas (Meena 2017). Comparatively, 5-20% loss due to Indian peafowl

seems less, nevertheless, it may still be considerable amount of loss to small farmers in RTR

area.

Chickpea, chilli, wheat, mustard (17-50% respondents for each crop) are the crops

preferred by Indian peafowl compared to tomatoes, sesame, coriander, sorghum, green peas,

moong (less than 10% respondents).

Peoples' perception about Indian peafowl being a crop pest changed according to

season in which they were interviewed. While wild boar or blue bull were consistently

reported as crop pests throughout the year, only 2.84% respondents mentioned Indian

peafowl as crop pest in November and as many as 19.7% respondents called it a crop pest in

July 2017. Actually, in July and August various crops are in growth stage. Our survey also

indicates that crops around harvest stage (not growth stage) are most likely to be eaten by

Indian peafowl followed by pre-growth stage (freshly sown, sprouted or sapling stage). Least

damage is incurred due to peafowl visiting the growing crops (between saplings to seed set)

(Fig 3).

The crop damage at different stages of growth may lead to different amount of loss.

Some of the damaged parts may regenerate but there is nevertheless a substantial amount of

loss in terms of total yield (Bayani et al. 2016). In contrast, for some crops such as chickpea,

herbivory in first 20 days after sowing may actually lead to greater branching resulting in

more number of seeds. However, herbivory beyond a threshold in pre-flowering stage is

known to decrease crop yield even in chickpea by as much as 67% (Bayani et al. 2016).

Many farmers have adopted different strategies to reduce the crop loss due to peafowl.

The strategies include guarding the farms during most vulnerable stages, changing crop

variety or place. The effectiveness of these strategies needs to be checked thoroughly. None-

the-less the strategies to reduce crop damage/ loss themselves cost the farmers in terms of

time, resources, money invested not to mention the inconvenience and loss incurred due to

choosing less than optimal choice of crop or farming place.

Overall, the effects of crop damage due to Indian peafowl may vary with type of crop

taken, season, etc. There is no denying that Indian peafowl seems to be a crop pest at least in

some parts of India. Surprisingly, most respondents in our survey did not have negative views

or resentment about Indian peafowl damaging/ eating their crops. Many of them, in fact,

mentioned that even peafowl have "right-to-live" and farmers do not consider it as "loss" if

peafowl come and eat in their farm. This is in sharp contrast to farmers' perceptions when other

crop pests such as wild boar, elephants or blue bulls visit their crops. Most of the households

in RTR area in fact pro-actively offer grains to peafowl throughout the year (94.87-100%

respondents). Typically bajra, sorghum, wheat or lentils are offered by each household in a

designated place around temple premises in RTR region. The quantity or frequency with which

each household offers grains may vary from place to place but on average about 15 Kg of grains

are offered daily at these designated places in RTR region as part of temple ritual/ traditional

belief. As a result, such places have become a reliable source of food for the wildlife including

Indian peafowl. Thus, people's positive/ neutral perception about Indian peafowl, traditional

beliefs and ritual practices such as active food provisioning at temples may be important for

maintaining high population numbers of peafowl in RTR area. One can conclude that human-

wildlife interactions and people's perception/ attitude towards wild life can impact not only

food provisioning but also wild life management (Jacobs et al. 2014a, b), conservation efforts

(Jacob and Harms 2014) and ultimately human–wildlife relationships (Jacobs et al. 2012).

Effects of food provisioning on peafowl populations

All around the world humans have provided many novel and reliable food resources to

herbivores in the form of crops and ornamental plants. Some species exploit these resources

quite well and can become economically important crop pests (Sih et al. 2011). Generalist

herbivores (those who have broad range of plant species they like to feed on) are more likely

to exploit the novel food resources (crops, ornamental plants) present in the human dominated

landscapes. Peafowls generally have a broad range of diet options including plant parts (fresh

sprouts, leaves, flowers, fruits, seeds) as well as insects, worms, sometimes small reptiles

(Sathyakumar and Kaul 2007). In spite of having most of these diet options available in our

study areas, peafowls were found eating food provided by humans (cereals, pulses, processed

food). According to our estimate, ~30-35 Kg food grains are offered to peafowl per day in the

village and surrounding areas of Morachi Chincholi, while ~15Kg grains are offered per day

around homes and temple premises in the villages in Rajasthan included in this study. In both

these study areas, grains are offered at designated places everyday throughout the year. Thus,

Indian peafowl population in Morachi Chincholi has access to diet rich in carbohydrates

(cereals) and proteins (pulses) throughout the year (Table 2). As much as 71% of their diet

consisted of food provisioned indirectly in the form of crops or directly as grains offered in the

village surroundings (Fig. 4). Villages in Rajasthan offer relatively less variety of cereals and

very few pulses, yet the diet of peafowl in Rajasthan has up to 61% of food provisioned by

humans versus 39% natural food. Peafowl population in Nashik, on the other hand has access

to less reliable and less varied food offered by humans (~ 3-5 Kg per day) (Table 2), as a result

only about 40% of their diet consists of provisioned food. It is argued that crop varieties are

rich in nutritional quality and poorer in secondary metabolites compared to their wild

counterparts (Rode et al., 2006). If this is true, it could explain why peafowl are choosing

harvested/ unharvested crops and ready to eat grains/ food, wherever they are available, over

their "natural food". The quantity, variety and nutritional quality of food available to peafowl

plus the reliability of such food resources might be crucial factor in making their food choices

and overall diet composition (Fig. 5).

Food provisioning not only influenced their diet composition, but also the time budget.

At food provision sites, the peafowls could spend up to 49.30% of their time just eating, which

is significantly higher than the time spent in feeding at other sites (Fig. 6). They could get high

quality food in one place without spending much time in walking (22.89%). In contrast, the

peafowl had to spend more time in walking (32.35%) in search of food if they were feeding at

non-provision sites, while they could spend only 35.24% of their time in feeding (Fig. 6).

Considering that most behaviours in the peafowl behavioural repertoire except feeding and

standing inactive consume energy, the benefit: cost ratio in terms of time/ energy spent in

various behaviours is higher at food provision sites compared to non-provision sites. This trend

is similar across the field sites, and marginally different according to the season. Peafowl spend

more time in reproduction related behaviours (display, vocal communication, mating, egg

laying, parental care etc.) during monsoon season, hence the benefit: cost ratio is slightly lower

compared to pre- and post-monsoon season as expected.

Effects of food provisioning on time budgets, diet and activity seen in Indian peafowl

populations are very similar to effects seen in primate populations (Kamal et al. 1997; Berman

and Li 2002; Ram et al. 2003; Hadi et al. 2007; McKinney 2010; El-Alami et al. 2012).

Interestingly, these effects extended to the ranging patterns, foraging success, and consequently

fitness of primates (Oro et al. 2013; Becker and Hall 2014). Food provisioned primate troops

also had higher growth rates and population densities (Altmann and Muruthi 1988; Jaman and

Huffman 2013). Comparable studies on birds, however, are rare especially in Indian

subcontinent. Thus, our study for the first time has documented effects of food provisioning on

feeding ecology of big bird such as Indian peafowl. The method(s) used in this study for

quantifying the changes in diet and activity budget may be generally applicable to understand

the changes in feeding ecology of other large birds found in human dominated landscapes.

Supplementary food available at bird feeders (typically in western countries) is known

to have far reaching consequences on avian ecology in terms of increasing survival during

overwintering, enhanced breeding success, changing sex-ratios of offspring in smaller avian

species, range expansion of species (Robb et al. 2008 and references therein). Although, most

of these consequences are positive, some negative consequences of food provisioning are also

possible in avian as well as primate systems. If animals start depending on food provided by

humans it may lead to behavioural changes as illustrated in Finland where some great-tit

populations are so dependent on supplementary food during winter that they can no longer be

sustained by natural food resources alone (Orell 1989). In Morachi Chincholi, where large

amounts of grains are provisioned throughout the year, somewhat parallel scenario is seen

during summer months. In the months of April, May and part of June, there are high

temperatures (up to 42-45^{0}C), water is scarce and if there are no crops during summer months

in a particular year, the peafowl concentrate around remaining waterbodies (village wells,

ponds) and wherever humans provide them grains and water (personal observations). Although

it is not surprising that peafowl incorporated nutrient rich resources in their diet wherever such

option was available, it is interesting that these resources are being actively provided to them

by humans in at least some places. It may, thus, be important to factor in human attitude towards

this species in population management or future conservation efforts especially considering

how frequently they are found in human dominated landscape.

The dynamic interplay of experiential, ecological, socioeconomic and cultural factors

can influence perception and local human behaviour towards wild-species with significant

implications for the management of wild species and the ecosystems they inhabit (D'Lima et

al., 2013). Studies such as ours may help in designing effective conservation strategies for a

magnificent bird that intrigued even Charles Darwin.

Acknowledgements

Thanks to Maharashtra Education Society's Abasaheb Garware College, Pune for hosting the

Ramalingaswami Re-entry Fellowship to DAP and providing the infrastructure and

administrative support. DAP and PD are thankful to Late Mr. Biswaroop Raha for suggesting

field sites in Nashik, Dr. Dharmendra Khandal and volunteers of Tiger Watch who extended

support for field work in Rajasthan. Authors also appreciate the support and hospitality of

Morachi Chincholi residents during field work. We are thankful to Rupesh Gawade, Pranav

Mhaisalkar and Ankita Divekar for helping with data collection on field.

Funding- This study was funded through Ramalingaswami Re-entry Fellowship to DAP by

Dept. of Biotechnology, Govt. of India.

Declarations

Authors contributions- DAP conceived the study, designed methodology, collected data,

analysed the data and prepared the manuscript. PD collected data, helped with analysis and

manuscript preparation.

Conflict of interest- Authors declare no conflict of interest.

Permissions- No live animals/ samples were handled during this study. The study was

conducted outside protected areas. Therefore, the work did not require special permissions.

References:

Agrawal, P.K., Vergese, A., Radhakrishna, S., & Subaharan K, Human-wildlife conflict. Curr

Sci, 2016, 111, 1581-1582.

Altmann, J., & Muruthi, P., Differences in daily life between semi-provisioned and wild-

feeding baboons. Am J Primatol, 1988, 15, 213–221.

Ayyappan, S., Agrawal, P.K., Rao, V.V., & Tripathi, R.S., Human-Wildlife 'conflict to

coexistence' in Agro-pastoral landscapes. In Human Animal Conflict in Agro-Pastoral

Context: Issues & Policies, (ed. Agarwal, P.K., Vergese, A., Radhakrishna, S. & Subaharan,

K.) ICAR, New Delhi, 2016, Pp. 1-9.

Bayani, A., Tiwade, D., Dongre, A., Phatak, R, & Watve, M., Assessment of Crop Damage

by Protected Wild Mammalian Herbivores on the Western Boundary of Tadoba-Andhari

Tiger Reserve (TATR), Central India. PLOS One, 2016, 11, 1-18.

Becker, D.J., & Hall, R.J., Too much of a good thing: Resource provisioning alters infectious

disease dynamics in wildlife. Biol Lett, 2014, 10, 1-5.

Berman, C.M., & Li, J.H., Impact of translocation, provisioning and range restriction on a

group of *Macaca thibetana*. Intl J Primatol, 2002, 23, 283–397.

Burton, M., & Burton, R., International Wildlife Encyclopedia. Marshall Cavendish, New

York, 2002.

Chauhan, N.P.S., & Singh, R., Crop Damage by over abundant populations of Nilgai and

Blackbuck in Haryana (India) and its management, Proceedings of the Fourteenth Vertebrate

Pest Conference, University of Nebraska-Lincoln, 1990.

Carpenter, L.H., Decker, D.J., & Lipscomb, J.F., Stakeholder acceptance capacity in wildlife

management. Human Dimensions of Wildlife, 2000, 5, 5–19.

Chauhan, N.P.S., Barwal, K.S., & Kumar, D., Human-wild pig conflict in selected states in

India and mitigation strategies. Acta Silvatica et Lignaria Hungarica, 2009, 5, 189-197.

Clergeau, P., Mennechez, G., Sauvage, A., & Lemoine, A., Human perception and

appreciation of birds: A motivation for wildlife conservation in urban environments of

France, In Avian Ecology and Conservation in an Urbanizing World, (eds. Marzluff, J.M.,

Bowman, R., & Donnelly, R.), Springer, Boston, Massachusetts, 2001, pp. 77-96.

D'Lima, C., Marsh, H., Hamann, M., Sinha, A., & Arthur R., Positive interactions between

Irrawaddy dolphins and artisanal fishers in the Chilika Lagoon of Eastern India are driven by

ecology, socioeconomics, and culture. J Human Environ, 2013, 43, 614-624.

El-Alami, A., Lavieren, E.V., Rachida, A., & Chait, A., Differences in activity budgets and

diet between semiprovisioned and wild-feeding groups of the Endangered Barbary macaque

(*Macaca sylvanus*) in the Central High Atlas Mountains, Morocco. Am J Primatol, 2012, 74,

210–216.

Graham, D.M., Notter, B., Adams, W.M., & Lee, P.C., & Ochieng, T.N., Patterns of crop-

raiding by elephants, *Loxodonta africana,* in Laikipia,Kenya, and the management of human–

elephant conflict. Systematics & Biodiversity, 2010, 8: 435–445.

Gilroy, J.J., and Sutherland W. J., Beyond ecological traps: perceptual errors and undervalued

resources, Trends Ecol Evol, 2007, 22, 351–356.

Hadi, I., Suryobroto, B., & Perwitasari-Farajallah, D.Y.A.H., Food preference of semi-

provisioned macaques based on feeding duration and foraging party size, HAYATI J Biosci,

2007, 14, 13 -17.

Horgan, F.G., Kudavidanage, E.P., Weragodaarachchi, A., & Ramp, D., Traditional 'maavee'

rice production in Sri Lanka: environmental, economic and social pressures revealed through

stakeholder interviews, Pawees, 2017.

Jacobs, M.H., Vaske, J.J., & Roemer J. M., Toward a mental systems approach to human

relationships with wildlife: the role of emotional dispositions, Human Dimensions of

Wildlife, 2012, 17, 4-15.

Jacobs, M.H. and Harms, M., Influence of interpretation on conservation intentions of whale

tourists, Tourism Management, 2014, 42, 123-131.

Jacobs, M.H., Vaske, J.J., & Sijtsma, M.T.J., Predictive potential of wildlife value

orientations for acceptability of management interventions, J Nature Conservation, 2014a, 22,

377-383.

Jacobs, M.H., Vaske, J.J., Dubois, S., & Fehres, P., More than fear: role of emotions in

acceptability of lethal control of wolves, Euro J Wildlife Res, 2014b, 60, 589-598.

Jaman, M.F., & Huffman, M.A., The effect of urban and rural habitats and resource type on

activity budgets of commensal rhesus macaques *Macaca mulatta* in Bangladesh, Primates,

2013, 54, 49–59.

Kamal, K.B., Boug, A., and Brain, P.F., Effects of food provisioning on the behaviour of

commensal Hamadryas baboons, *Papio hamadryas*, at Al Hada mountain in Western Saudi

Arabia, 1997, Zool Middle East, 14, 11–22.

Mathur, V.B., Kaushik, M., Bist, S.S., Mungi, N.A., & Qureshi, Q., Management of human–

wildlife interaction and invasive alien species in India, Report no. TR 2015/004. Wildlife

Institute of India, Dehradun, 2015, pp. 1–235.

McKinney, T., The effects of provisioning and crop-raiding on the diet and foraging activities

of human-commensal white-faced capuchins (*Cebus capucinus*), Am J Primatol, 2010, 71: 1–

10.

Meena, V., 'Site'ing the right reasons: critical evaluation of conservation planning for the

Asiatic lion. Euro J Wildlife Res, 2010, 56, 209–213.

Meena, M., Agriculture crop damage by antelope (*Boselaphus tragocamelus*) and

management strategies: Challenges in India, Intl J Zool Studies, 2017, 2, 157-160.

Naughton-Treves, L., 1997, Farming the forest edge: Vulnerable places and people around

Kibale National Park, Uganda, Geographical Rev, 87, 27-46.

Orell, M., Population fluctuations and survival of Great Tits (*Parus major)* dependent on

food supplied by man in winter. IBIS, 1989, 131, 112-127.

Oro, D., Genovart, M., Tavecchia, G., Fowler, M.S., & Martınez-Abrain, A., Ecological and

evolutionary implications of food subsidies from humans, Ecol Lett, 2013, 16, 1501–1514.

Pathak, B.J., Pati, B.P., Kumar, R., Kumar, A., Raval, P.P., Patel, V.S., & Rana, V.J.,

Biodiversity Conservation Plan for Gir. Supplementary Management Plan, 2002, pp. 407.

Perera, M., Vandercone, R., Patterns and perceptions of wildlife crop raiding in and around

the Kaludiyapokuna Forest Reserve in the dry zone of Sri Lanka, Proceedings of the Third

Undergraduate Research Symposium on Zoology & Environmental Management, 2017.

Ram, S., Venkatachalam, S., Sinha, A., Changing social strategies of wild female Bonnet

Macaques during natural foraging and on provisioning, Curr Sci India, 2003, 84, 780–790.

Ramkumar, K., Balasundaram, R., & Sarvanamuthu, R., Crop Damage by Asian Elephants

Elephas maximus and Effectiveness of Mitigating Measures in Coimbatore Forest Division,

South India, Intl Res J Biol Sci, 2014, 3, 1-11.

Robb, G. N., McDonald, R. A., Chamberlain, D. E., & Bearhop, S., Food for thought:

supplementary feeding as a driver of ecological change in avian populations, Front Ecol

Environ, 2008, 6, 476–484.

Rode, K.D., Chiyo, P.I, Chapman, C.A., McDowells, L.R., Nutritional ecology of elephants

in Kibale National Park, Uganda, and its relationship with crop-raiding behaviour, J Tropical

Ecol, 2006, 22, 441–449.

Sathyakumar, S. and Kaul, R., Pheasants, In: ENVIS Bulletin: Wildlife and Protected

Areas, *Galliformes of India*. (ed. Sathyakumar S., Sivakumar K.), Wildlife Institute of

India, Dehradun, India, 2007, pp. 33-52.

Sengupta, A., McConkey, K.R., & Radhakrishna, S., Primates, Provisioning and Plants:

Impacts of Human Cultural Behaviours on Primate Ecological Functions, PLOS ONE, 2015,

10(11), e0140961.

Sengupta, A., & Radhakrishna, S., The Hand That Feeds the Monkey: Mutual Influence of

Humans and Rhesus Macaques (*Macaca mulatta*) in the Context of Provisioning, Intl J

Primatol, 2018, 39, 817-830.

Sih, A., Ferrari, M.C.O., & Harris, D.J., Evolution and behavioural responses to human-

induced rapid environmental change, Evolutionary Applications, 2011, 4, 367-387.

Stork, N., Re-assessing current extinction rates. Biodiversity Conservation, 2010, 19, 357–

371.

Tilamn, D., May, R.M., Lehman, C.L., & Nowak, M.A., Habitat destruction and extinction

debt, Letters to Nature, 1994, 371, 65-66.

Treves, A., Wallace, R.B., Naughton-Treves, L., & Morales, A., Co-Managing Human–

Wildlife Conflicts: A Review, Human Dimensions of Wildlife, 2006, 11, 383–396.

Vijayan, S., and Pati, B.P., Impact of changing cropping pattern on man-animal conflicts

around Gir P.A. with specific reference to Talala Taluka, Indian Forester, 2001, 127, 1121-

1133.

Figure Captions:

Fig1 Percentage of respondents reporting crop pests during the months of November, July-

August and March in Rajasthan

Fig2 Percentage of respondents reporting crop loss due to Peafowl during the months July-

August, March and November in Rajasthan. Bars indicate percentage of crop loss reported by

respondents in various months of the year.

Fig3 Percentage of Respondents reporting stage of crop affected due to Peafowl during the

months July- August, March and November in Rajasthan

Fig4 Proportion of natural versus provisioned food items in the Indian Peafowl diet across

study areas.

Fig5 Diet composition of the Indian Peafowl populations across selected field areas

Fig6 Time budget of Indian Peafowl populations at food provision and non-provision sites.

Fig7 Comparison of time spent in pecking and walking at food provision and non-provision

sites. Error bars indicate 95% confidence intervals.

Figure 1

640   **Figure 2**

643   **Figure3**

647 **Figure 4**

Figure 5

Figure 6

Figure 7

696 **Table 1:** Brief description of study areas

Sr.No.	Name of the study area	GPS co-ordinates	Brief site description
1	Morachi Chincholi	18° 48' 26.9" N, 74° 09' 29.3" E, elevation 641m	Village, agricultural lands, agro-eco-tourism resorts (artificial habitat), dry deciduous hillock
2	Nashik a. Ganagpur Nursery, Gangapur dam	20° 02.109'N, 73° 40.970'E, elevation 611m	Plantation, agricultural fields, big water reservoir
	b. MERI campus, Nashik City	20° 01'11.0"N, 73° 48'04.0"E, elevation 404m	Plantation
3	Sawai Madhopur, Rajasthan		
	a. Kutthalpura	26° 03' 58.5" N, 76° 26' 13.4" E, elevation 261m	Dry deciduous forest, agricultural farms. Tourism for tigers.
	b. Bangadda-Kalan	26° 3'57.78"N, 76°42'35.76"E	Village, agricultural fields, No tourism
	c. Sanwata-Kalakhora	26° 6'41.46"N, 76°38'33.81"E elevation 267m	Remote village with agricultural fields, dry deciduous forest. No tourism
	d. Govindpura	26° 04.826" N, 76° 48.097" E, elevation 227m	Village, agricultural fields. No tourism

Table 2: Diet composition of Indian Peafowl *Pavo cristatus* at three field sites based on direct field observations.

Diet Item	Food Category	Natural/ provisioned food	Morachi Chincholi	Nashik	Rajasthan
Sorghum (*Sorghum spp*)	Cereals	P	1	0	0
Wheat (*Triticum spp.*)	Cereals	P	1	1	1
Bajra (*Sorghum spp*)	Cereals	P	1	0	1
Rice (*Oryza sativum*)	Cereals	P	1	1	1
Corn (*Zea mays*)	Cereals	P	1	0	0
Unknown grains	Cereals	P	1	0	0
Vilayati Chinch (*Pithecellobium dulce*)	Non-grain	N	0	1	0
Methi Grass	Non-grain	N	1	0	0
Insects	Non-grain	N	1	1	1
Leaves/ grass	Non-grain	N	1	1	1
Seeds	Non-grain	N	0	1	0
Chillies (*Capsicum annum)*	Non-grain	P	1	0	1
Vegetables	Non-grain	P	1	0	0
Unknown (non-grain)	Non-grain	N	1	1	1
Processed food	Processed food	P	1	1	0
Indian Brown lentils	Pulses	P	1	0	1

Green gram (*Vigna radiata*)	Pulses	P	1	0	1
Split pigeon peas (*Cajanus cajan)*	Pulses	P	0	0	1
Black gram lentils (*Vigna mungo)*	Pulses	P	1	0	0
Moth beans (*Vigna aconitifolia)*	Pulses	P	1	0	0
Black eyed peas (*Vigna unguiculata)*	Pulses	P	1	0	0
Chickpeas (*Cicer arietum)*	Pulses	P	1	0	0
Peanuts	Pulses	P	1	0	0