A tale of two species: human and peafowl interactions in human dominated landscape

influence each other's behaviour

Short Title: Wild life near human habitation

Dhanashree Paranjpe* and Priyanka Dange

Annasaheb Kulkarni Department of Biodiversity, MES Abasaheb Garware College, Karve

Road, Pune-411004.

*Corresponding author

E-mail: dhana4shree@gmail.com

Telephone No: +91-20-41038237

ORCID ID*: https://orcid.org/0000-0003-4787-2052

Priyanka Dange: ORCID ID: http://orcid.org/0000-0002-2799-2156

Abstract:

Wild life is increasingly coming in contact with humans in many parts of the world. Human perception of wild life may be an important factor in population management and conservation practices in urban and semi-urban areas. Human attitude towards bird species may vary from non-violent coexistence to a perception of birds as pests. Based on the data collected from survey interviews, we studied the perceptions of local communities in Rajasthan, India towards Indian Peafowl. Local communities in Rajasthan reported Indian Peafowl as crop pest and this perception varied across seasons. The crop loss incurred due to Indian Peafowl also varied across seasons according to the respondents. Despite reporting peafowls as a crop pest, locals regularly offered grains for them as a traditional practice. Thus, at our study sites locals have mostly positive perception about Indian peafowl around them.

Food provisioning by local human population influenced diet composition and time-budget of Indian Peafowls at food provision and non-provision sites. Sites at which food provisioning was less or absent, peafowl spent more time in walking in search of food and more than half of their diet consisted of natural food. In contrast, the sites at which plenty of grains were available, time spent in walking was significantly less, while time spent in feeding was significantly more; and over 70% of their diet consisted of carbohydrate and protein-rich provisioned grains. Food provisioning changed the benefit: cost ratio (measured as time spent in feeding to time spent in all other behaviours) between provision and non-provision sites. Thus, food provisioning by humans can change feeding ecology of native Indian peafowl populations, while the presence of peafowl in human dominated landscape changes how humans interact with wild life around them.

Introduction:

Wild animals are increasingly finding themselves near human habitation due to

scarcity of natural habitats, high human population density and changing land usage

(Ayyappan et al. 2016; Pathak et al. 2002; Vijayan and Pati, 2001; Meena 2010). As a result,

human-animal interactions and conflicts have risen. The impact of wild animals on local

human population has mostly been studied in the context of human-wildlife conflicts

(Agrawal et al. 2016) while, impact of local human population on animal populations living

close to or within rural, semi-urban, and urban areas has rarely been studied.

There are potentially diverse types of interactions between humans and wildlife. The

impact of human induced ecological stressors on animal populations in human dominated

landscapes is becoming an important area of research (Sih et al. 2011; Agrawal et al. 2016).

Wild animals can influence local human population socially, economically through crop loss,

property damage, life loss, ecosystem services and aesthetic/ religious value. They can

become sources of income through ecotourism, trade of animal derived products and so on.

Vicinity to human population can change natural behaviour and life history of animals

in multiple ways. Human habitation or agricultural landscapes can serve as safer refuge areas

or even new habitats for wildlife (Horgan et al. 2017). Humans often provide food to wildlife

voluntarily around their homes, in religious places as a ritual or in wild life tourism spots/

feeding stations (Robb et al. 2008; Sengupta et al. 2015). In addition to this, indirect food

provisioning in the form of crops, ornamental plants, livestock or waste food thrown in open

becomes easy and reliable source of food for wild life. Food provided by humans tends to be

calorie rich, easily digestible, available at predictable times and places. It is known that food

provisioning might change feeding habits, diet preference or diet composition of wild

mammals (Robb et al. 2008; Sih et al. 2011; Sengupta et al. 2015; Ayyappan et al. 2016;

Sengupta and Radhakrishna 2018). Reduced predatory pressure and regular availability of

nutrient rich food (in the form of crops) round the year are likely to provide greater resilience

of wild animals. It also allows them to live successfully close to agricultural landscape and

human habitation (Gilroy and Sutherland 2007; Sih et al. 2011; Ayyappan et al. 2016).

There are many avian species like crows, sparrows, pigeons, cranes, egrets that

inhabit human- dominated landscapes or stay very close to / within human habitation.

Appreciation for birds and even individual species may vary from nonviolent coexistence

(including feeding them) to a perception of birds as pests (Clergeau et al. 2001). Very little

work has been done on how human perception/ attitude towards avian species can influence

the outcome of their interactions with humans. A multi-dimensional approach to study such

interactions is necessary in order to understand the capacity of local stakeholders to accept

wild species (Carpenter et al., 2000) as well as the potential effect of these interactions on

both humans and wildlife who share common spaces.

Indian peafowl (*Pavo cristatus*) is a species that has co-habited human dominated

landscapes for centuries in its native geographical range. This avian species is native to

Indian subcontinent and has been introduced in many parts of the world relatively recently.

Although, it's native habitat is undergrowth in open forest and woodlands near a waterbody,

it is also known to occur near farmlands, villages and increasingly becoming common in

urban and semi-urban areas (Burton and Burton 2002). In some places, such as Morachi

Chincholi in Maharashtra, India, "eco-tourism" based on Indian peafowl is flourishing and it

is an additional or alternative source of income for local community (personal observations).

In contrast to this "positive" impact, anecdotal reports suggest (Ganesan, R. 2016) that the

Indian peafowl can cause substantial crop losses in the areas where their population density is

high. There are no known studies from India that estimate the crop losses due to peafowls. It

is not known whether peafowl really affect crops like other known crop pests (e.g. wild boar,

elephants, deer, etc.). Numerous reports can be found in newspapers in India about deaths of

individuals or groups of peacocks. The deaths may be due to natural reasons such as

predation, water scarcity or may be the result of unintentional poisoning due to insecticides/

pesticides sprayed on crops that the peafowl later feed on. Sometimes intentional killing of

Indian peafowl by certain tribal communities for their ornamental feathers and meat has also

been suspected/ reported. In contrast to this, in some parts of India (e.g. State of Rajasthan),

peafowl are believed to be sacred and people actively offer grains for them as part of their

daily ritual. These varied interactions between Indian peafowl and local human population

make it an interesting study system to understand consequences of human- wild life

interactions. It remains to be seen how common are the positive perceptions and associated

beliefs/ rituals throughout the region in which the population density of Indian peafowl is

higher. Can these perceptions result in curbing population decline or effective management/

conservation of the species? To address these questions we studied three aspects of human-

peafowl interactions in details. The objectives of this study were: (i) To understand the

perceptions of local community about Indian peafowl, (ii) To estimate impact of Indian

peafowl populations on local agriculture and (iii) To estimate the impact of food provisioning

by local community on peafowl populations.

Materials and methods:

Study area

We selected three field sites- Morachi Chincholi, Nashik and Rajasthan (Table 1).

Selection of study areas was based on several criteria such as their closeness to human

habitation, accessibility throughout the year, history of peafowl populations in the area,

potential interactions between humans and peafowl (food provisioning, peafowl visiting/

roosting on agricultural land and village homes, sell of peafowl feathers or related

"products", tourism, cultural significance), etc. Areas located outside protected areas and

close to human habitation were given preference.

We used questionnaire based interviews (N = 196) to estimate impact of Indian

peafowl populations on local agriculture and to understand perceptions of local people about

Indian peafowl in four villages located on the periphery of Ranthambhore Tiger Reserve

(RTR), in the state of Rajasthan, India. Before actual survey, a pilot survey was conducted to

standardize the protocol and contents of the questionnaire. Based on the results of the pilot

survey some questions were modified or added to reduce ambiguity in questions and get more

accurate data. Interviews were conducted by taking family as a unit respondent. The

questionnaire included fixed and open-ended questions (See supplementary material for

complete questionnaire). Interviews started with the interviewer clearly explaining objectives

of the study and the interview was continued only if the respondent(s) gave consent. The

questionnaire consisted of 3 parts: (i) basic information about the respondent such as name,

age, address, number of people in family unit, source(s) of income (agriculture/ others). (ii)

Questions related to perceived impact of peafowls on agriculture (iii) questions related to

social/ cultural/ traditional perceptions about Indian peafowl. Interviews were conducted in

three different seasons: November 2016 (harvest time for kharif crops, winter season, N =

55), March-April 2017 (Harvest time for rabbi crops, beginning of dry summer season, N =

86), July-August 2017 (standing crops, monsoon season, N = 55) to check if responses to

some of the questions related to agriculture differed according to stage of crop/ season. For

each survey equal number of respondents were sampled randomly from the local village

populations.

To estimate the impact of food provisioning by local community on peafowl

populations, behaviours of the Indian peafowl were studied at Morachi Chincholi, Nashik and

Rajasthan throughout the year using direct field observations. Observations were recorded

during morning (6 am to 10 am) and evening (4 pm to 7 pm). Behaviours were also observed

in the late- afternoons if peafowl were found to be active. The behavioural observations of

peafowl were noted while patrolling the study area using continuous scan and focal animal

sampling. Focal animals were chosen randomly from a group (if the peafowl were seen in a

group). Additionally, opportunistic observations were recorded for individuals sighted during

the walk around the study sites. Behaviours of focal animal were recorded continuously without

disturbing or going too close to focal animal (typically from a distance of 10-100m) until they

were out of the sight. During feeding bouts, location of feeding was recorded and was later

classified as provisioned and non-provisioned sites. Gender of the focal animal was noted along

with the date, place and the total duration of observations. Data were also collected about the

type of food the peafowl ate during the observations using binoculars to get clear view of the

type of food they were eating. Observations were stopped when the focal animal flew/ walked

out of observer's sight.

For calculating time budget of the Indian peafowl populations, the observer(s)

documented start/ end/ transition of behaviours of focal animal as a running commentary

using handheld digital audio recorder (SONY ICD-UX560F). Occasionally, data were

recorded as videos using a Panasonic camcorder (Full HD, 29.8mm wide 20.4 megapixel HC-

X920) if a large group of peafowl was found near feeding site.

Data processing

The types of food items peafowls ate were listed using direct observations (Table 2). Cereals, pulses, vegetables and processed food is available to peafowl only around human habitation and hence was considered as provisioned food, while all other food items such as grass, leaves, sprouts, fruits, seeds, insects, worms were considered as natural food items (Table 2). Based on direct observations of peafowl eating various food items, proportion of times it was seen eating natural and provisioned food items in peafowl diet was calculated. If it was not clear what the peafowl were eating, such observations were not considered for analysis of diet composition. The food items were further categorised as cereals (grains such as sorghum, wheat, bajra, rice, corn, etc.), pulses (split pigeon peas, black gram lentils, black-eyed pea, Indian brown lentils, green gram, peanuts, etc.), non-grains (fruits, grass, seeds, leaves, insects, vegetables, etc.) and processed food. Number of times peafowls were seen eating each food type was counted. Based on this frequency, the percentage of each food type in their diet was calculated.

The ethogram of the peafowl comprised of following behaviours: alert, walk, peck, preen, jump, run, scratch, calls, shake, scrape, search, turn, flap, fly, stand, display related, climb, headshake, perch, chase away, sit, pause, poop, drink, aggressive and shiver. (Fig.1). Relative frequency of each behaviour recorded for 293 individuals was calculated as (# of times a behaviour is shown/total no. of observations)*100 to identify the major and minor behaviours of Indian Peafowl from the ethogram. Peck, Alert and Walk were the major behaviours while the minor (<10% frequency) were pooled as other behaviours. For calculating time budget of the Indian peafowl populations, data about time of beginning of a behaviour and end of behaviour/ transition to other behavioural state was noted down from the audio records. This was used to calculate the time spent in various activities/ behaviours for each recorded individual. Fraction of time spent in each activity was summed across all individuals in the

population. Time budget of the whole population was calculated as time spent in each activity

divided by total time of observation for the population.

For calculating benefit : cost ratio, time spent in feeding was considered as "benefit"

while time spent in all other activities (except standing inactive) was considered as "cost".

Statistical analysis

For statistical analysis, STATISTICA^{TM} software version 13.2 (Dell Inc. 2016) was

used. As the data did not follow normal distribution, non-parametric tests were used for further

analysis. Kruskal-Wallis ANOVA was used to see the effect of field sites (Morachi Chincholi,

Nashik, Rajasthan) on time spent in feeding, time spent in vigilance, walking and other

behaviours. Mann-Whitney U test was used to estimate the effect of proximity to food

provision site on time spent in feeding, vigilance, walking and other behaviours.

Benefit: cost ratio was compared across three field sites (Morachi Chincholi, Nashik,

Rajasthan) and between food provision and non-provision sites using non-parametric tests such

as Kruskal- Wallis ANOVA.

Results

Perceptions about Indian peafowl and their relation to agriculture

The age of survey respondents varied from 15- 65yrs, all age classes in this range were

represented almost equally. Respondents included 90.73% males and 9.27% females. Sources

of income for the respondents included agriculture (51.32% of respondents), animal herding

(17.88%), labour (farm or other labour 20.2%) and other sources (10. 6%) such as contractor,

company job, mechanic, shop owner, artisan, teacher etc. A majority of farmers in the area

around RTR are small farmers with 84.11% farmers owning less than 5 acres of land, while

rest of the farmers (15.89%) owned 5+ acres to 20+ acres of land. Farmers take two to sixteen

types of crops in a year depending on size of farm, availability of labour, water, etc. Most

common crops in this region include wheat, bajra, chilly, mustard, sorghum, sesame. All

farmers in this region reported crop losses irrespective of time of year during which survey

was conducted. The reasons for crop loss differed according to season during which

interviews were conducted. Table 3 gives details of season-wise reasons of crop loss. Major

crop pests reported in the survey are Wild boar (*Sus scrofa*), Nilgai/ Blue bull (*Boselaphus*

tragocamelus), Indian peafowl (*Pavo cristatus*) and Deer. However, number of respondents

reporting particular types of crop pests changed according to season in which survey was

conducted (Fig 2). Although Indian peafowl is the third common crop pest reported, only 3 %

of respondents in November 2016 survey said it was a crop pest compared to 15 %

respondents in March 2017 and 20% respondents in July 2017. On the other hand wild boar

and blue-bull were reported as crop pests throughout the year by 19-28% respondents. The

range reflects variation in data collected across different seasons in the study area.

According to the respondents, peafowl used the agricultural land for feeding and as a

source of water (Table 4). Some of the respondents (0.33%) also mentioned that they spread

diseases. Between 67.65- 100% respondents reported that the long train of feathers causes

some damage to crops when the males with full grown train walk through the standing crops.

Estimates of total loss due to peafowl varied between 0% to more than 20% according to

season in which the survey was conducted (Fig 3). Chickpea, chilli, wheat and mustard were

the major crops that incurred loss due to peafowl. Loss was also reported occasionally for

coriander, sorghum, sesame, tomatoes, green peas and moong (all reported by 10% or less

respondents). Indian peafowl ate crops at post growth (around harvest) stage according to 60-

95% of respondents, next most vulnerable stage was pre-growth stage (freshly sown, sprouted

or sapling stage) according to 3-29 % of responses. Damage to crops was reported least in the

growth stage (8 to 10 % respondents). Very few respondents reported damage to crops at all

stages (2 %) and no stages (2 %). (Fig 4). Between 64.65%-100% farmers reported that they

changed crop pattern to reduce the loss due to peafowl. Many farmers chose to take a

different variety of crop that is less preferred by peafowl (e.g. hybrid chilli instead of

indigenous variety) or avoided taking certain crops (e.g. chickpea, ground nut, chilli) or

change the place where they take the crop if they own patches of farm lands in more than one

locations within the village. Interestingly, the estimate of loss (estimated % loss) due to

Indian peafowl showed positive correlation with size of farm (Spearman rank order

correlation $r = 0.24$, $p < 0.05$). Farmers with more farm land estimated more crop losses due

to peafowl compared to farmers who owned less land.

Food provisioning for Indian peafowl

In spite of the crop damage the peafowl cause, local people offer grains to peafowl

throughout the year. All households (100% respondents) in the village offer grains to peafowl

except in the month of November slightly less number of respondents (94.87%) reported

giving grains for peafowl. Most common grains offered to peafowl include bajra (35-45%

respondents), sorghum (27.5-41.25% respondents), wheat (10-25.95% respondents), and

lentils (0-7.57% respondents) while some people also offer rice, maize, sesame when

available. People typically offer a handful (100 – 130 g depending on type of grains offered)

to up to 3 Kg grains at a time. In some villages, grains are offered from each household every

day, in some cases they are offered in temple once or twice a week or on special days/

occasions. So year around there are on average ~15 Kg grains available for wild life every

day in and around temple premises in RTR region.

Impact of food provisioning on peafowl populations

Impact of food provisioning was studied in three study areas- Morachi Chincholi,

Nashik and Rajasthan (RTR). Diet composition of the peafowl was different in the selected

study areas. In Morachi Chincholi, peafowl diet consisted of 71% food provisioned by humans

compared to 61% provisioned food in Rajasthan, while just 40% peafowl diet in Nashik

consisted of provisioned food (Fig 5). Peafowl in Nashik ate almost double the amount of

natural (non-grain) food items (60%) compared to peafowl in Morachi Chincholi (29%). Major

portion of food provided to peafowl populations in Morachi Chincholi and Rajasthan consisted

of grains (66% and 58%, respectively). Peafowl in Rajasthan were not seen eating processed

food as opposed to peafowl in Morachi Chincholi and Nashik study area (Table 5).

There was lot of variation in the total duration of observations for each individual

(Range: 1 min-29 min.), however, total observation time did not differ significantly across field

sites (Median Test, *Chi-Square* $= 1.75$, *df* $= 2$, *p* $= 0.42$) which confirms that equal efforts were

put into observations at the selected field sites. The relative proportion of time spent in different

behaviours at food provision sites (where humans put grains for peafowl) and away from

provision site (non-provision sites) is shown in Figure 6. Time spent in vigilance (21.88%) and

other behaviours (8.98%) at provision sites was comparable to time spent in those behaviours

at non-provisioning sites (30.43 %, and 14.26 %, respectively) (Fig. 6). At the provision sites

(N=181), individuals spent significantly less time in walking (18.41%) compared to non-

provision sites (N=99, 22.99%, Mann- Whitney U test, *p*$= 0.003$) (Fig. 6 and Fig. 7). Peafowls

spent significantly more time in feeding (50.73%) at the provision sites (N=183) as compared

to non-provision sites (N=92, 32.32%, Mann-Whitney U test, *p*$= 0.0003$) (Fig. 6 and fig. 7).

Time spent in other behaviours was significantly more at Nashik (N=59) as compared to

Morachi Chincholi (N=139) and Rajasthan (N=88), respectively (Kruskal-Wallis ANOVA,

Multiple comparison of means, p$= 0.03$).

Benefit: cost ratio measured in terms of time spent in feeding (benefit) versus other

behaviours (cost) was significantly different between food provision and non-provision sites

(Mann-Whitney U test, $p = 0.00015$). The ratio was similar across field sites (Kruskal-Wallis

ANOVA, $p = 0.22$).

Discussion:

Human- wild life interactions have mostly been studied so far in the context of human-

animal conflict which has drastic consequences such as loss of property, crops or even death.

All human- wild life interactions do not necessarily lead to obvious conflicts, nonetheless,

they might impact both sides in significant ways. Peoples' perception about wild life can

potentially influence the long-term outcome of such interactions. Our study investigated in

detail some aspects of non-violent human- wildlife interactions by focusing on Indian

peafowl populations living in human dominated landscape. Our survey covered the

perceptions about Indian Peafowl in a wide range of age groups (15-65 yrs) in a geographic

area with high population density of peafowls. Most of the surveys were conducted in remote

rural areas of Rajasthan by volunteers going door-to-door for collecting data. Survey was

conducted considering family as a unit respondent. In most cases, head of the family along

with one or two other members (male/ female) present in the household answered the

questions. Number of female respondents in our survey remained below 10%. So the results

may be biased for male gender. Most of the respondents in the survey were small farmers

with less than five acres of land. The source(s) of income, in most cases (89.4% respondents)

were related to agriculture (crops + farm animals + labour).

Perceptions about Indian peafowl and their relation to agriculture

We did not want to bias the respondents by directly asking them if peafowl damaged

their crops. Hence, we asked questions step-by-step about types of crops and typical reasons

for crop loss in RTR area. Losses due to erratic rain patterns (deficient rainfall, wet drought,

untimely rains) is the main reason for crop losses for Indian agriculture which depends

largely on Monsoon rains from Indian ocean. Surprisingly, damage due to crop pests emerged

as the top most reason across all seasons for loss to farmers in RTR area, followed by loss

due to untimely rains, crop disease, or any other reasons (Table 3). The area in which we

conducted surveys includes buffer zone and villages just outside the Ranthambhore National

park. As a result, wild herbivores from the protected area frequently enter the villages,

farmlands and surrounding areas. This can create serious problems for the farmers whose

livelihood depends on agricultural produce. Similar issues of crop raiding by wild herbivores

have been reported all over the world (Naughton-Treves 1997; Treves et al. 2006; Graham et

al. 2010; Mathur et al. 2015; Perera and Vandercone 2017) including India (Chauhan and

Singh, 1990; Chauhan et al. 2009; Ramkumar et al. 2014; Meena 2017). Wild boar (*Sus*

scrofa), Nilgai/ blue bull (*Boselaphus tragocamelus*), Indian peafowl (*Pavo cristatus*) and

deer were reported as the major crop pests in RTR area. While wild boar and blue bull are

some of the commonly known crop raiders in India, ours is first systematic survey in which

Indian peafowl has been reported as a crop pest in India. There are sporadic newspaper

reports and anecdotal information from different parts of India about Indian peafowl

damaging crops (Ganesan, R. 2016), however, there has been no systematic survey or

documentation of extent and type of crop damage done by Indian peafowl as we have done in

this survey. Recently, Indian peafowl was reported as a crop pest by farmers in Sri Lanka

(Horgan et al. 2017; Perera and Vandercone 2017). One study reported that Maavee paddies

and surrounding habitats offer refuges for wildlife including Indian peafowl; and Indian

peafowl damaged rice at the seedling and grain filling stage irrespective of the farming

method used (Horgan et al. 2017).

Peafowl used agricultural land as a source of food, water (Table 4), display sites and

resting sites (personal observations). A majority of respondents (87.63-95.45%) in our

survey reported that peafowl do affect crops in an adverse way. So Indian peafowl is not only

visiting their farms seeking refuge or using farms as new feeding habitat, in most cases they

do have perceivable negative effects on crops. Although they were not reported to spread

disease (only 0.33% respondents think they spread crop disease), their long train was found to

cause some damage as they walked through the standing crop. They also eat various parts and

stages of the crop (Fig. 4). Estimates of total crop loss due to Indian peafowl ranged from

negligible (0%) to over 20%. Typically, figures between 5-20% losses were mentioned by

respondents for various crops. It is important to note that these are "perceived" estimates of

loss given by farmers and the actual figures need to be verified through more detailed 'on

field' data collection. Other wild herbivores such as elephants can cause up to 35% crop

damage (Ramakumar et al. 2014), while wild boars can destroy 5-36% of crops (Chauhan et

al. 2009). Blue bull (*Boselaphus tragocamelus*) seems to have the highest impact with 58%

losses (rarely less than 10%) in Harayana, India (Chauhan and Singh 1990). The losses due to

blue bull can be as high as 70% in high density areas (Meena 2017). Comparatively, 5-20%

loss due to Indian peafowl seems less, nevertheless, it may still be considerable amount of

loss to small farmers in RTR area. Chickpea, chilli, wheat, mustard (17-50% respondents for

each crop) are the crops preferred by Indian peafowl compared to tomatoes, sesame,

coriander, sorghum, green peas, moong (less than 10% respondents).

Peoples' perception about Indian peafowl being a crop pest changed according to

season in which they were interviewed. While wild boar or blue bull were consistently

reported as crop pests throughout the year, only 3 % respondents mentioned Indian peafowl

as crop pest in November, 15% in March and as many as 20 % respondents called it a crop

pest in July 2017. Our survey also indicates that crops around harvest stage (post-growth

stage) are most likely to be eaten by Indian peafowl followed by pre-growth stage (freshly

sown, sprouted or sapling stage) and growth stage (between saplings to leaves set). Least

damage is incurred due to peafowl visiting the growing crops (Fig 4).

The crop damage at different stages of growth may lead to different amount of loss.

Some of the damaged parts may regenerate but there is nevertheless a substantial amount of

loss in terms of total yield (Bayani et al. 2016). In contrast, for some crops such as chickpea,

herbivory in first 20 days after sowing may actually lead to greater branching resulting in

more number of seeds. However, herbivory beyond a threshold in pre-flowering stage is

known to decrease crop yield even in chickpea by as much as 67% (Bayani et al. 2016).

Many farmers have adopted different strategies to reduce the crop loss due to peafowl.

The strategies include guarding the farms during most vulnerable stages, changing crop

variety or place. The effectiveness of these strategies needs to be checked thoroughly. None-

the-less the strategies to reduce crop damage/ loss themselves cost the farmers in terms of

time, resources, money invested not to mention the inconvenience and loss incurred due to

choosing less than optimal choice of crop or farming place.

Overall, the effects of crop damage due to Indian peafowl may vary with type of crop

taken, season, etc. There is no denying that Indian peafowl seems to be a crop pest at least in

some parts of India. Surprisingly, most respondents in our survey did not have negative views

or resentment about Indian peafowl damaging/ eating their crops. Many of them, in fact,

mentioned that even peafowl have "right-to-live" and farmers do not consider it as "loss" if

peafowl come and eat in their farm. It is possible that the 'perceived beauty' of peafowl along

with their association with popular deities in India (Nair, 1974) and relatively less amounts of

crop damage by this species may play a role in biasing the perception of people towards this

species as compared to the other pests. This seemingly positive perception is in sharp contrast

to farmers' perceptions when other crop pests such as wild boar, elephants or blue bulls visit

their crops. Most of the households in RTR area in fact pro-actively offer grains to peafowl

throughout the year (94.87-100% respondents). Typically bajra, sorghum, wheat or lentils are

offered by each household in a designated place around temple premises in RTR region. The

quantity or frequency with which each household offers grains may vary from place to place

but on average about 15 Kg of grains are offered daily at these designated places in RTR region

as part of temple ritual/ traditional belief. As a result, such places have become a reliable source

of food for the wildlife including Indian peafowl. Thus, people's positive/ neutral perception

about Indian peafowl, traditional beliefs and ritual practices such as active food provisioning

at temples may be important for maintaining high population numbers of peafowl in RTR area.

One can conclude that human- wildlife interactions and people's perception/ attitude towards

wild life can impact not only food provisioning but also wild life management (Jacobs et al.

2014a, b), conservation efforts (Jacob and Harms 2014) and ultimately human–wildlife

relationships (Jacobs et al. 2012).

Effects of food provisioning on peafowl populations

All around the world humans have provided many novel and reliable food resources to

herbivores in the form of crops and ornamental plants. Some species exploit these resources

quite well and can become economically important crop pests (Sih et al. 2011). Generalist

herbivores (those who have broad range of plant species they like to feed on) are more likely

to exploit the novel food resources (crops, ornamental plants) present in the human dominated

landscapes. Peafowls generally have a broad range of diet options including plant parts (fresh

sprouts, leaves, flowers, fruits, seeds) as well as insects, worms, sometimes small reptiles

(Sathyakumar and Kaul 2007). In spite of having most of these diet options available in our

study areas, peafowls were found eating food provided by humans (cereals, pulses, processed

food). According to our estimate, ~30-35 Kg food grains are offered to peafowl per day in the

village and surrounding areas of Morachi Chincholi, while ~15Kg grains are offered per day

around homes and temple premises in the villages in Rajasthan included in this study. In both

these study areas, grains are offered at designated places everyday throughout the year. Thus,

Indian peafowl population in Morachi Chincholi has access to diet rich in carbohydrates

(cereals) and proteins (pulses) throughout the year (Table 2). As much as 71% of their diet

consisted of food provisioned indirectly in the form of crops or directly as grains offered in the

village surroundings (Fig. 5). Villages in Rajasthan offer relatively less variety of cereals and

very few pulses, yet the diet of peafowl in Rajasthan has up to 61% of food provisioned by

humans versus 39% natural food. Peafowl population in Nashik, on the other hand has access

to less reliable and less varied food offered by humans (~ 3-5 Kg per day), as a result only

about 40% of their diet consists of provisioned food. It is argued that crop varieties are rich in

nutritional quality and poorer in secondary metabolites compared to their wild counterparts

(Rode et al., 2006). If this is true, it could explain why peafowl are choosing harvested/

unharvested crops and ready to eat grains/ food, wherever they are available, over their "natural

food". The quantity, variety and nutritional quality of food available to peafowl plus the

reliability of such food resources might be crucial factor in making their food choices and

overall diet composition (Table 5).

Food provisioning not only influenced their diet composition, but also the time budget.

At food provision sites, the peafowls could spend up to 50.73% of their time just eating, which

is significantly higher than the time spent in feeding at non-provision sites (Fig. 6 and fig. 7).

They could get high quality food in one place without spending much time in walking. In

contrast, the peafowl had to spend more time in walking in search of food if they were feeding

at non-provision sites, while they could spend only 32.32% of their time in feeding (Fig. 6).

Effect of less amount of food provisioning at Nashik as compared to Morachi Chincholi and

Rajasthan can be seen in the higher time spent by Peafowls in other behaviours. Our results

suggest that the benefit: cost ratio in terms of time spent in behaviours can be influenced due

to food provisioning. We could not calculate benefits and costs in terms of energy budget as it

was difficult to estimate energy gain from non-provisioned food items. It will be interesting to

explore other cost benefits of food provisioning in terms of body condition/ health of Indian

peafowl near human habitation.

Effects of food provisioning on time budgets, diet and activity seen in Indian peafowl

populations are very similar to effects seen in primate populations (Kamal et al. 1997; Berman

and Li 2002; Ram et al. 2003; Hadi et al. 2007; McKinney 2010; El-Alami et al. 2012).

Interestingly, these effects extended to the ranging patterns, foraging success, and consequently

fitness of primates (Oro et al. 2013; Becker and Hall 2014). Food provisioned primate troops

also had higher growth rates and population densities (Altmann and Muruthi 1988; Jaman and

Huffman 2013). Comparable studies on birds, however, are rare especially in Indian

subcontinent. Thus, our study for the first time has documented effects of food provisioning on

feeding ecology of big bird such as Indian peafowl.

Supplementary food available at bird feeders (typically in western countries) is known

to have far reaching consequences on avian ecology in terms of increasing survival during

overwintering, enhanced breeding success, changing sex-ratios of offspring in smaller avian

species, range expansion of species (Robb et al. 2008 and references therein). Although, most

of these consequences are positive, some negative consequences of food provisioning are also

possible in avian as well as primate systems. If animals start depending on food provided by

humans it may lead to behavioural changes as illustrated in Finland where some great-tit

populations are so dependent on supplementary food during winter that they can no longer be

sustained by natural food resources alone (Orell 1989). In Morachi Chincholi, where large

amounts of grains are provisioned throughout the year, somewhat parallel scenario is seen

during summer months. In the months of April, May and part of June, there are high

temperatures (up to 42-45^{0}C), water is scarce and if there are no crops during summer months

in a particular year, the peafowl concentrate around remaining waterbodies (village wells,

ponds) and wherever humans provide them grains and water (personal observations). Although

it is not surprising that peafowl incorporated nutrient rich resources in their diet wherever such

option was available, it is interesting that these resources are being actively provided to them

by humans in at least some places. It may, thus, be important to factor in human attitude towards

this species in population management or future conservation efforts especially considering

how frequently they are found in human dominated landscape.

The dynamic interplay of experiential, ecological, socioeconomic and cultural factors

can influence perception and local human behaviour towards wild-species with significant

implications for the management of wild species and the ecosystems they inhabit (D'Lima et

al., 2013). Studies such as ours may help in designing effective conservation strategies for a

magnificent bird that intrigued even Charles Darwin.

Acknowledgements

Thanks to Maharashtra Education Society's Abasaheb Garware College, Pune for hosting the

Ramalingaswami Re-entry Fellowship to DAP and providing the infrastructure and

administrative support. DAP and PD are thankful to Late Mr. Biswaroop Raha for suggesting

field sites in Nashik, Dr. Dharmendra Khandal and volunteers of Tiger Watch who extended

support for field work in Rajasthan. Authors also appreciate the support and hospitality of

Morachi Chincholi residents during field work. We are thankful to Rupesh Gawade, Pranav

Mhaisalkar, Ankita Divekar, Akash Dubey and Vedanti Mahimkar for helping with data

collection on field.

Funding- This study was funded through Ramalingaswami Re-entry Fellowship to DAP by

Dept. of Biotechnology, Govt. of India.

Declarations

Authors contributions- DAP conceived the study, designed methodology, collected data,

analysed the data and prepared the manuscript. PD collected data, helped with analysis and

manuscript preparation.

Conflict of interest- Authors declare no conflict of interest.

Permissions- No live animals/ samples were handled during this study. The study was

conducted outside protected areas. Therefore, the work did not require special permissions.

References:

Agrawal, P.K., Vergese, A., Radhakrishna, S., & Subaharan K, Human-wildlife conflict. Curr

Sci, 2016, 111, 1581-1582.

Altmann, J., & Muruthi, P., Differences in daily life between semi-provisioned and wild-

feeding baboons. Am J Primatol, 1988, 15, 213–221.

Ayyappan, S., Agrawal, P.K., Rao, V.V., & Tripathi, R.S., Human-Wildlife 'conflict to

coexistence' in Agro-pastoral landscapes. In Human Animal Conflict in Agro-Pastoral

Context: Issues & Policies, (ed. Agarwal, P.K., Vergese, A., Radhakrishna, S. & Subaharan,

K.) ICAR, New Delhi, 2016, Pp. 1-9.

Bayani, A., Tiwade, D., Dongre, A., Phatak, R, & Watve, M., Assessment of Crop Damage

by Protected Wild Mammalian Herbivores on the Western Boundary of Tadoba-Andhari

Tiger Reserve (TATR), Central India. PLOS One, 2016, 11, 1-18.

Becker, D.J., & Hall, R.J., Too much of a good thing: Resource provisioning alters infectious

disease dynamics in wildlife. Biol Lett, 2014, 10, 1-5.

Berman, C.M., & Li, J.H., Impact of translocation, provisioning and range restriction on a

group of *Macaca thibetana*. Intl J Primatol, 2002, 23, 283–397.

Burton, M., & Burton, R., International Wildlife Encyclopedia. Marshall Cavendish, New

York, 2002.

Chauhan, N.P.S., & Singh, R., Crop Damage by over abundant populations of Nilgai and

Blackbuck in Haryana (India) and its management, Proceedings of the Fourteenth Vertebrate

Pest Conference, University of Nebraska-Lincoln, 1990.

Carpenter, L.H., Decker, D.J., & Lipscomb, J.F., Stakeholder acceptance capacity in wildlife

management. Human Dimensions of Wildlife, 2000, 5, 5–19.

Chauhan, N.P.S., Barwal, K.S., & Kumar, D., Human-wild pig conflict in selected states in

India and mitigation strategies. Acta Silvatica et Lignaria Hungarica, 2009, 5, 189-197.

Clergeau, P., Mennechez, G., Sauvage, A., & Lemoine, A., Human perception and

appreciation of birds: A motivation for wildlife conservation in urban environments of

France, In Avian Ecology and Conservation in an Urbanizing World, (eds. Marzluff, J.M.,

Bowman, R., & Donnelly, R.), Springer, Boston, Massachusetts, 2001, pp. 77-96.

D'Lima, C., Marsh, H., Hamann, M., Sinha, A., & Arthur R., Positive interactions between

Irrawaddy dolphins and artisanal fishers in the Chilika Lagoon of Eastern India are driven by

ecology, socioeconomics, and culture. J Human Environ, 2013, 43, 614-624.

El-Alami, A., Lavieren, E.V., Rachida, A., & Chait, A., Differences in activity budgets and

diet between semiprovisioned and wild-feeding groups of the Endangered Barbary macaque

(*Macaca sylvanus*) in the Central High Atlas Mountains, Morocco. Am J Primatol, 2012, 74,

210–216.

Graham, D.M., Notter, B., Adams, W.M., & Lee, P.C., & Ochieng, T.N., Patterns of crop-

raiding by elephants, *Loxodonta africana,* in Laikipia,Kenya, and the management of human–

elephant conflict. Systematics & Biodiversity, 2010, 8: 435–445.

Gilroy, J.J., and Sutherland W. J., Beyond ecological traps: perceptual errors and undervalued

resources, Trends Ecol Evol, 2007, 22, 351–356.

Hadi, I., Suryobroto, B., & Perwitasari-Farajallah, D.Y.A.H., Food preference of semi-provisioned macaques based on feeding duration and foraging party size, HAYATI J Biosci, 2007, 14, 13 -17.

Horgan, F.G., Kudavidanage, E.P., Weragodaarachchi, A., & Ramp, D., Traditional 'maavee' rice production in Sri Lanka: environmental, economic and social pressures revealed through stakeholder interviews, Pawees, 2017.

Jacobs, M.H., Vaske, J.J., & Roemer J. M., Toward a mental systems approach to human relationships with wildlife: the role of emotional dispositions, Human Dimensions of Wildlife, 2012, 17, 4-15.

Jacobs, M.H. and Harms, M., Influence of interpretation on conservation intentions of whale tourists, Tourism Management, 2014, 42, 123-131.

Jacobs, M.H., Vaske, J.J., & Sijtsma, M.T.J., Predictive potential of wildlife value orientations for acceptability of management interventions, J Nature Conservation, 2014a, 22, 377-383.

Jacobs, M.H., Vaske, J.J., Dubois, S., & Fehres, P., More than fear: role of emotions in acceptability of lethal control of wolves, Euro J Wildlife Res, 2014b, 60, 589-598.

Jaman, M.F., & Huffman, M.A., The effect of urban and rural habitats and resource type on activity budgets of commensal rhesus macaques *Macaca mulatta* in Bangladesh, Primates, 2013, 54, 49–59.

Kamal, K.B., Boug, A., and Brain, P.F., Effects of food provisioning on the behaviour of commensal Hamadryas baboons, *Papio hamadryas*, at Al Hada mountain in Western Saudi Arabia, 1997, Zool Middle East, 14, 11–22.

Mathur, V.B., Kaushik, M., Bist, S.S., Mungi, N.A., & Qureshi, Q., Management of human–wildlife interaction and invasive alien species in India, Report no. TR 2015/004. Wildlife Institute of India, Dehradun, 2015, pp. 1–235.

McKinney, T., The effects of provisioning and crop-raiding on the diet and foraging activities

of human-commensal white-faced capuchins (*Cebus capucinus*), Am J Primatol, 2010, 71: 1–

10.

Meena, V., 'Site'ing the right reasons: critical evaluation of conservation planning for the

Asiatic lion. Euro J Wildlife Res, 2010, 56, 209–213.

Meena, M., Agriculture crop damage by antelope (*Boselaphus tragocamelus*) and

management strategies: Challenges in India, Intl J Zool Studies, 2017, 2, 157-160.

Naughton-Treves, L., 1997, Farming the forest edge: Vulnerable places and people around

Kibale National Park, Uganda, Geographical Rev, 87, 27-46.

Nair, Thankappan.P., 1974, The Peacock Cult in Asia, Asian Folkfore Studies, 33 , 93-170

Orell, M., Population fluctuations and survival of Great Tits (*Parus major)* dependent on

food supplied by man in winter. IBIS, 1989, 131, 112-127.

Oro, D., Genovart, M., Tavecchia, G., Fowler, M.S., & Martınez-Abrain, A., Ecological and

evolutionary implications of food subsidies from humans, Ecol Lett, 2013, 16, 1501–1514.

Pathak, B.J., Pati, B.P., Kumar, R., Kumar, A., Raval, P.P., Patel, V.S., & Rana, V.J.,

Biodiversity Conservation Plan for Gir. Supplementary Management Plan, 2002, pp. 407.

Perera, M., Vandercone, R., Patterns and perceptions of wildlife crop raiding in and around

the Kaludiyapokuna Forest Reserve in the dry zone of Sri Lanka, Proceedings of the Third

Undergraduate Research Symposium on Zoology & Environmental Management, 2017.

Ganesan, R., National Bird and a Stated Animal- a nuisance in Goa. 2016. Available from:

http://www.downtoearth.org.in/wildlife-and-biodiversity.

Ram, S., Venkatachalam, S., Sinha, A., Changing social strategies of wild female Bonnet

Macaques during natural foraging and on provisioning, Curr Sci India, 2003, 84, 780–790.

Ramkumar, K., Balasundaram, R., & Sarvanamuthu, R., Crop Damage by Asian Elephants

Elephas maximus and Effectiveness of Mitigating Measures in Coimbatore Forest Division,

South India, Intl Res J Biol Sci, 2014, 3, 1-11.

Robb, G. N., McDonald, R. A., Chamberlain, D. E., & Bearhop, S., Food for thought:

supplementary feeding as a driver of ecological change in avian populations, Front Ecol

Environ, 2008, 6, 476–484.

Rode, K.D., Chiyo, P.I, Chapman, C.A., McDowells, L.R., Nutritional ecology of elephants

in Kibale National Park, Uganda, and its relationship with crop-raiding behaviour, J Tropical

Ecol, 2006, 22, 441–449.

Sathyakumar, S. and Kaul, R., Pheasants, In: ENVIS Bulletin: Wildlife and Protected

Areas, *Galliformes of India*. (ed. Sathyakumar S., Sivakumar K.), Wildlife Institute of

India, Dehradun, India, 2007, pp. 33-52.

Sengupta, A., McConkey, K.R., & Radhakrishna, S., Primates, Provisioning and Plants:

Impacts of Human Cultural Behaviours on Primate Ecological Functions, PLOS ONE, 2015,

10(11), e0140961.

Sengupta, A., & Radhakrishna, S., The Hand That Feeds the Monkey: Mutual Influence of

Humans and Rhesus Macaques (*Macaca mulatta*) in the Context of Provisioning, Intl J

Primatol, 2018, 39, 817-830.

Sih, A., Ferrari, M.C.O., & Harris, D.J., Evolution and behavioural responses to human-

induced rapid environmental change, Evolutionary Applications, 2011, 4, 367-387.

Treves, A., Wallace, R.B., Naughton-Treves, L., & Morales, A., Co-Managing Human–

Wildlife Conflicts: A Review, Human Dimensions of Wildlife, 2006, 11, 383–396.

Vijayan, S., and Pati, B.P., Impact of changing cropping pattern on man-animal conflicts

around Gir P.A. with specific reference to Talala Taluka, Indian Forester, 2001, 127, 1121-

1133.

Figure Captions:

Fig1. Ethogram of behaviours of Indian Peafowl across all field sites.

Fig2. Percentage of respondents reporting crop pests during the months of November, July- August and March in Rajasthan

 Fig3. Percentage of respondents reporting crop loss (in %) due to Peafowl during the months July- August, March and November in Rajasthan.

Fig4. Percentage of Respondents reporting stage of crop affected due to Peafowl during the months July- August, March and November in Rajasthan

Fig5. Proportion of natural versus provisioned food items in the Indian Peafowl diet across study areas.

Fig6. Time budget of Indian Peafowl populations at food provision and non-provision sites.

Fig7. Comparison of time spent in pecking and walking at food provision and non-provision sites. Error bars indicate 95% confidence intervals.

Fig.1

Fig. 2

Crop pests in July-August

24 23
11 22
20

⧅Wildboar ⧆Nilgai ▨Peafowl ⬚Deer ▩Others

Crop pests in March

31 26
8 20
15

⧅Wildboar ⧆Nilgai ▨Peafowl ⬚Deer ▩Others

Crop pests in November

21 28
20 28
3

⧅Wildboar ⧆Nilgai ▨Peafowl ⬚Deer ▩Others

Fig. 3

Fig. 4

649   **Fig. 5**

670
671

Fig. 6

Fig. 7

693 **Table 1:** Brief description of study areas

Sr.No.	Name of the study area	GPS co-ordinates	Brief site description
1	Morachi Chincholi	18° 48' 26.9" N, 74° 09' 29.3" E, elevation 641m	Village, agricultural lands, dry deciduous hillock
2	Nashik a. Ganagpur Nursery, Gangapur dam	20° 02.109'N, 73° 40.970'E, elevation 611m	Plantation, agricultural fields, big water reservoir
	b. MERI campus, Nashik City	20° 01'11.0"N, 73° 48'04.0"E, elevation 404m	Plantation
3	Sawai Madhopur, Rajasthan		
	a. Kutthalpura	26° 03' 58.5" N, 76° 26' 13.4" E, elevation 261m	Dry deciduous forest, agricultural farms.
	b. Bangadda-Kalan	26° 3'57.78"N, 76°42'35.76"E	Village, agricultural fields,
	c. Sanwata-Kalakhora	26° 6'41.46"N, 76°38'33.81"E elevation 267m	Remote village with agricultural fields, dry deciduous forest.
	d. Govindpura	26° 04.826" N, 76° 48.097" E, elevation 227m	Village, agricultural fields.

695 **Table 2:** Diet composition (presence/ absence of provisioned and non-provisioned food) of

696 Indian Peafowl *Pavo cristatus* at three field sites based on direct field observations.

Diet Item	Food Category	Natural/ provisioned food	Morachi Chincholi	Nashik	Rajasthan
Sorghum (*Sorghum spp*)	Cereals	P	1	0	0
Wheat (*Triticum spp.*)	Cereals	P	1	1	1
Bajra (*Sorghum spp*)	Cereals	P	1	0	1
Rice (*Oryza sativum*)	Cereals	P	1	1	1
Corn (*Zea mays*)	Cereals	P	1	0	0
Unknown grains	Cereals	P	1	0	0
Vilayati Chinch (*Pithecellobium dulce*)	Non-grain	N	0	1	0
Methi Grass	Non-grain	N	1	0	0
Insects	Non-grain	N	1	1	1
Leaves/ grass	Non-grain	N	1	1	1
Seeds	Non-grain	N	0	1	0
Chillies (*Capsicum annum)*	Non-grain	P	1	0	1
Vegetables	Non-grain	P	1	0	0

Unknown (non-grain)	Non-grain	N	1	1	1
Processed food	Processed food	P	1	1	0
Indian Brown lentils	Pulses	P	1	0	1
Green gram (*Vigna radiata*)	Pulses	P	1	0	1
Split pigeon peas (*Cajanus cajan*)	Pulses	P	0	0	1
Black gram lentils (*Vigna mungo*)	Pulses	P	1	0	0
Moth beans (*Vigna aconitifolia*)	Pulses	P	1	0	0
Black eyed peas (*Vigna unguiculata*)	Pulses	P	1	0	0
Chickpeas (*Cicer arietum*)	Pulses	P	1	0	0
Peanuts	Pulses	P	1	0	0

Table 3: Typical reasons of crop loss in RTR region of Rajasthan

Typical reasons of crop -loss	July-August (%)	March (%)	November (%)
Crop pests	62.34	59.82	100.00
Untimely rains	36.36	31.25	0.00
Disease	1.30	8.93	0.00

Table 4: Agricultural land usage by Indian Peafowl in RTR region of Rajasthan

Peafowl's Activities in farm	% of Respondents
Eat harvest	42.67
Eat leaves	17.00
Eat insects	19.67
Drink water	6.67
Eat flowers	8.00
Eat seeds	0.67
Eat sapplings	3.67
Nothing	1.33
Spread disease	0.33

725 **Table 5: Diet composition (in %) of Indian Peafowl across field sites**

Field Site	Cereals (%) (provisioned)	Pulses (%) (provisioned)	Processed (%) (provisioned)	Non-grain (%) (non-provisioned)
Morachi Chincholi	66	3	1	30
Nashik	38	0	2	60
Rajasthan	58	3	0	39