1

# Genetic characterization of outbred Sprague Dawley rats and utility for genome-wide association studies

Alexander F. Gileta[1,3], Christopher J. Fitzpatrick[2], Apurva S. Chitre[3], Celine L. St. Pierre[1,4], Elizabeth V. Joyce[4], Rachael J. Maguire[4], Africa M. McLeod[4], Natalia M. Gonzales[1], April E. Williams[3], Jonathan D. Morrow[2,5], Terry E. Robinson[6], Shelly B. Flagel[2,5,7], Abraham A. Palmer[3, 8*]

[1] Department of Human Genetics, University of Chicago, Chicago, Illinois, United States of America

[2] Neuroscience Graduate Program, University of Michigan, Ann Arbor, Michigan, United States of America

[3] Department of Psychiatry, University of California, San Diego, La Jolla, California, United States of America

[4] Biological Sciences Collegiate Division, University of Chicago, Chicago, Illinois, United States of America

[5] Department of Psychiatry, University of Michigan, Ann Arbor, Michigan, United States of America

[6] Department of Psychology and Neuroscience Program, University of Michigan, Ann Arbor, Michigan, United States of America

[7] Michigan Neuroscience Institute, University of Michigan, Ann Arbor, Michigan, United States of America

[8] Institute for Genomic Medicine, University of California, San Diego, La Jolla, California, United States of America

* Corresponding author

Email: aapalmer@ucsd.edu

## Abstract

Sprague Dawley (**SD**) rats are among the most widely used outbred laboratory rat populations. Despite this, the genetic characteristics of SD rats have not been clearly described, and SD rats are rarely used for experiments aimed at exploring genotype-phenotype relationships. In order to use SD rats to perform a genome-wide association study (**GWAS**), we collected behavioral data from 4,625 SD rats that were predominantly obtained from two commercial vendors, Charles River Laboratories and Harlan Sprague Dawley Inc. Using double-digest genotyping-by-sequencing (**ddGBS**), we obtained dense, high-quality genotypes at 291,438 SNPs across 4,061 rats. This genetic data allowed us to characterize the variation present in Charles River vs. Harlan SD rats. We found that the two populations are highly diverged ($F_{ST} > 0.4$). Furthermore, even for rats obtained from the same vendor, there was strong population structure across breeding facilities and even between rooms at the same facility. We performed multiple separate GWAS by fitting a linear mixed model that accounted population structure and using meta-analysis to jointly analyze all cohorts. Our study examined Pavlovian conditioned approach (**PavCA**) behavior, which assesses the propensity for rats to attribute incentive salience to reward-associated cues. We identified 46 significant associations for the various metrics used to define PavCA. The surprising degree of population structure among SD rats from different sources has important implications for their use in both genetic and non-genetic studies.

## Author Summary

Outbred Sprague Dawley rats are among the most commonly used rats for neuroscience, physiology and pharmacological research; in the year 2020, 4,188 publications contained the

keyword "Sprague Dawley". Rats identified as "Sprague Dawley" are sold by several

commercial vendors, including Charles River Laboratories and Harlan Sprague Dawley Inc.

(now Envigo). Despite their widespread use, little is known about the genetic diversity of SD.

We genotyped more than 4,000 SD rats, which we used for a genome-wide association study

(**GWAS**) and to characterize genetic differences between SD rats from Charles River

Laboratories and Harlan. Our analysis revealed extensive population structure both between and

within vendors. The GWAS for Pavlovian conditioned approach (**PavCA**) identified a number of

genome-wide significant loci for that complex behavioral trait. Our results demonstrate that,

despite sharing an identical name, SD rats that are obtained from different vendors are very

different. Future studies should carefully define the exact source of SD rats being used and may

exploit their genetic diversity for genetic studies of complex traits.

## Introduction

Rats are among the most commonly used organisms for experimental psychology and

biomedical research. Whereas research using mice makes extensive use of inbred strains,

research conducted in rats typically uses commercially available *outbred* rats. Sprague Dawley

(SD) rats are very commonly used; we identified 4,188 publications in the year 2020 that

contained the key word "Sprague Dawley". Rats identified as "Sprague Dawley" are distributed

by several vendors. Each vendor has multiple breeding locations, and each breeding location has

one or more rooms or "barriers facilities" which may represent population isolates. Prior studies

have identified numerous physiological differences between SD rats obtained from different

vendors [1–5]. Despite these observations, many researchers may assume that SD rats obtained

from different vendors, breeding facilities or barrier facilities are largely interchangeable. There

has been little research into the genetic diversity and population structure that exists among

commercially available outbred rats [6]. Prior rat genetic studies have used $F_2$ and more

complex, multi-parental crosses of inbred strains for quantitative trait loci (QTL) mapping [7–9]

and GWAS [10–13]; however, we are unaware of any such studies that have employed

commercially available outbred rats. We and others have demonstrated the potential benefits and

challenges associated with the use of commercially available outbred mice for GWAS [14–16],

suggesting that similar studies in rats might also be of value.

SD rats originated in 1925 at the Sprague-Dawley Animal Company (Madison, WI),

where they were created from a cross between a hooded male hybrid of unknown origin and an

albino Wistar female [17]. In 1950, Charles River Inc. began to distribute SD rats commercially.

In 1980, Harlan Inc. (now Envigo, Inc.) also began to distribute SD rats after their acquisition of

Sprague-Dawley, Inc. [18]. In 1992, Charles River reestablished a foundation colony of SD rats,

using 100 breeder pairs from various existing colonies [19]. The resulting litters were used to

populate SD colonies globally and have since been bred using a mating system that minimizes

inbreeding. Every 3 years, Charles River replaces 25% of their male breeders in each production

colony with rats from a single foundation colony. Charles River also replaces 5% of their

foundation colony breeding pairs with rats from the production colonies on a yearly basis. These

practices are intended to reduce genetic drift between the production colonies [20]. Harlan also

reported using a rotational breeding system to limit inbreeding [21]; however, more detailed

information was not publicly available. Since Harlan's acquisition by Envigo, the process has

become more transparent [22]. Envigo follows a Poiley rotational breeding scheme [23],

whereby animals are cycled through different sections of the colony with each generation,

reducing genetic drift and inbreeding.

We used SD rats from multiple vendors, breeding locations, and barrier facilities to

elucidate the genetic background of SD and to perform a GWAS of a complex behavior. DNA

samples and phenotypic information were obtained from rats used in multiple studies as part of

an unrelated Program Project grant (P01DA031656) concerned with individual variation in the

propensity to attribute incentive value to food and drug cues [24,25]. All rats were first screened

for Pavlovian conditioned approach behavior (PavCA) [26], which provides an index of the

degree to which a reward cue has been attributed with incentive salience.  Although the genetic

analyses reported here were not part of the original design, our study took advantage of the

opportunities afforded by that large, behavioral study. We extracted genomic DNA from

available tissue samples and then applied double digest genotype-by-sequencing (ddGBS) to

obtain dense genotypes for 4,625 SD rats. We used these genotypes to genetically characterize

different populations of SD. Next, we used those genotype data and behavioral phenotypic data

to perform what we believe is the largest rodent GWAS ever undertaken. Our results provide

insights about the population structure and suitability of SD for GWAS and the genetic

underpinnings of Pavlovian conditioned approach.

**Results**

**Phenotype**

The final quality controlled dataset consisted of 4,061 genotyped male SD rats that were

also phenotyped for Pavlovian conditioned approach; 2,281 from Harlan and 1,780 from Charles

River, from 5 and 4 different breeding locations, respectively. As noted previously [26], we

found that the metrics used to describe performance in PavCA are highly correlated (S1 Fig).

Additionally, several of the base and composite PavCA metrics had tail-heavy distributions due

to biased responding in sign- and goal-tracking from the animals during the testing time periods

(S2 Fig). For this reason, we chose to quantile normalize all measurements prior to mapping. The

PavCA index score [26], which has been used previously to categorize rats into sign-trackers

(**STs**), goal-trackers (**GTs**), and intermediate responders (**IRs**), showed the expected divergence

and stabilization for STs and GTs over the five days of testing (Fig 1A) [26].

125            Charles River and Harlan rats had significantly different average PavCA index score

distributions (Fig 1B; Welch's 2 -sample t-test, p-value $< 2.2 \times 10^{-16}$), with Charles River rats

biased towards goal-tracking and Harlan rats biased towards sign-tracking. We divided the

samples further into the breeding locations that the rats originated from to investigate intra-

vendor behavioral variance (S1 Table). The differences in PavCA index score between breeding

locations within vendor were smaller, but still significant (S3 Fig) [6].

**Fig 1. PavCA index score progression across days and distribution between Charles River**

**and Harlan.**

(A) Samples were classified as ST, GT, or IR based on their average day 4 and 5 PavCA index

score. PavCA index scores for each day of training (1-5) were averaged across all STs, GTs, and

IRs. (B) Density curves of average day 4 and 5 PavCA index score in Harlan (n=2,281) vs.

Charles River (n=1,780) SD rats. (C) Filtered SNP density per 1Mb window in Charles River vs.

Harlan samples for all 20 autosomes.

**Genotyping and genetic characterization of SD rats**

Variant calling was performed on the full sample of genotyped SD rats. Subsequent

quality and minor allele frequency filtering of the variants was initially completed for each

vendor separately, under the notion that the observed differences in PavCA performance between

vendors may reflect underlying genetic variation. We identified more single nucleotide

polymorphisms (SNPs) for the rats from Charles River compared to Harlan (214,309 vs 114,568;

S2 Table). Fig 1C compares the distribution of SNPs across each chromosome for each vendor.

There were some regions where Harlan had few SNPs, but Charles River had many, and other

regions where both Harlan and Charles River had few SNPs. The observed jagged distributions

were expected due to the use of double digest genotyping-by-sequencing (ddGBS) to capture the

genotypes genome-wide (S1 Text).  After combining the two SNP sets, we identified a total of

234,887 unique, high-quality, bi-allelic SNPs. Using 381 samples with two replicates of

sequencing, we were able to evaluate our genotyping accuracy by looking at the discordance

between calls made in the replicates. The discordance rate between replicates was calculated to

be ~1.7%. However, since discordance putatively reflects random errors in either replicate, it is

expected to be approximately twice the true error rate, suggesting that our genotypes were >99%

accurate (0.85% error rate; although we cannot exclude the possibility that certain errors were

consistent across samples).

In addition to the variants described above, which were obtained using ANGSD/Beagle,

we also used STITCH as an alternative genotyping pipeline [27]. This approach identified a

larger total number of variants (S2 Table); however, after pruning SNPs with high linkage

disequilibrium (LD; $r^2 \geq 0.95$), STITCH produced fewer SNPs compared to ANGSD/Beagle.

Preliminary GWAS suggested the two SNP sets produced broadly similar results. Ultimately, we

chose to use the ANGSD/Beagle variants for all subsequent analyses.

To further investigate potential genetic divergence between Charles River and Harlan, we

performed principal component analysis (PCA) on a set of 4,502 LD-pruned ($r^2 < 0.5$) SNPs with

MAF $> 0.05$ across both vendor populations (Fig 2B). The first PC corresponded to vendor

(Charles River or Harlan) and accounted for ~33.7% of the variance. The second PC accounted

for just ~0.9% of the variance and reflected population structure within Harlan SD rats. To

investigate the within-vendor structure further (Fig 2A), we performed PCA on the samples from

each vendor separately using the same set of SNPs. Fig 2C and 2D show evidence of

substructure at both the level of breeding location (i.e. the city) and barrier facility (i.e., the

segregated breeding areas within the building). Interestingly, rats from some barrier facilities

showed greater divergence from barrier facilities within the same breeding location than barriers

at other locations.

**Fig 2. Genetic architecture of SD rats from Charles River vs. Harlan.**

(A) Map of the nine vendor breeding locations and the number of SD rats obtained from each

location. (B) A summary of the genetic data from all 4,061 SD rats based on principal

components 1 and 2 from PCA. Each point represents a sample. The left cluster is composed of

samples from Charles River and the right clusters are composed of samples from Harlan. (C-D)

Repeated PCA analyses on subsets of the samples from Harlan and Charles River, colored by

barrier facility of origin.

Due to the stark clustering of samples within and across certain barriers (Fig 2C & Fig

2D), we opted to divide the sample set up based on PCA clustering for all subsequent analyses,

including the GWAS for PavCA. This heuristic sample division resulted in seven subgroups,

each containing genetically-similar samples: SD rats originating from Harlan 202A/202C/208A,

Harlan 206, Harlan 217, Charles River R09/P03/P07/P10/C71/K92, Charles River R04, Charles

River P09, and Charles River C72. Variant filtering steps were reapplied to each subgroup

individually to reach final sets of SNPs (S4 Table). A more lenient minor allele frequency

(MAF) threshold was applied to permit more variants to pass filtering.

We observed a large difference in the minor allele frequency (MAF) distributions for

subgroups of Charles River vs Harlan origin (Fig 3A), with Charles River rats harboring more

variation and a far greater proportion of SNPs with high MAF (>0.05). This observation could

reflect the fact that Charles River has adhered to their International Genetics Standardization

Protocol for 25+ years, whereas Harlan appears to have focused on maintaining diversity within

breeding colonies and may have allowed for a moderate degree of drift between them. We went

on to examine the levels of linkage disequilibrium (LD) in the subgroups by constructing LD

decay curves (Fig 3B). These curves show the rate at which LD between two genetic loci

dissipates as a function of the distance between the loci. Harlan rats had more extensive LD

compared to Charles River. For contrast, we included the decay curve for Swiss Webster (CFW)

mice, a commercially available outbred mouse population that has been successfully used for

GWAS in the past [14,15].

**Fig 3. Fig 3. SNP MAF distributions and comparison of linkage disequilibrium decay rates.**

(A) Density curves of minor allele frequencies for 214,309 SNPs in Charles River and 114,568

SNPs in Harlan, after removing SNPs with MAF < 0.01. (B) Linkage disequilibrium decay rates

in SD rats from both vendors and outbred Swiss Webster (CFW) mice.

The fixation index ($F_{ST}$) is a statistic widely employed by population geneticists to

measure the level of structure in populations [28]. It is calculated using the variance in allele

frequencies among populations; values closer to 0 indicate genetic homogeneity, and values

closer to 1 indicate genetic differentiation. We calculated the pairwise $F_{ST}$ between all seven

subgroups (Table 1). $F_{ST}$ values between vendors were ~0.423, which is over 2-fold greater than

corresponding estimates for major human lineages [29], whereas the values for different

subgroups within a vendor were substantially lower.

**Table 1. Pairwise $F_{ST}$ statistics for Harlan and Charles River breeding locations.**

| | Charles River | | | | Harlan | | |
|---|---|---|---|---|---|---|---|
| **Subgroups** | R04 | R09 + | P09 | C72 | 202A + | 206 | 217 |
| R04 | 0 | 0.019 | 0.028 | 0.024 | 0.422 | 0.426 | 0.419 |
| R09 + | | 0 | 0.011 | 0.013 | 0.419 | 0.423 | 0.416 |
| P09 | | | 0 | 0.022 | 0.422 | 0.426 | 0.419 |
| C72 | | | | 0 | 0.426 | 0.431 | 0.423 |
| 202A + | | | | | 0 | 0.024 | 0.013 |
| 206 | | | | | | 0 | 0.028 |
| 217 | | | | | | | 0 |

We speculated that some of the rats in our sample might also share close genetic

relationships with one another. We used *plink 1.9* [30,31] to estimate the pairwise proportions of

identity-by-descent (IBD; Panels A and C in S4 Fig), which showed that while most rats were

only distantly related, a subset shared closer familial relationships. We removed several rats that

showed high levels of relatedness with many other samples (presumably due to technical error),

as well as any with unreasonably high levels of IBD (S5 Table; Panels B and D in S4 Fig).

**SNP heritability and genome-wide association analyses**

Although evidence from selective breeding studies has suggested that behavior in the

Pavlovian conditioned approach procedure is heritable [32], we are not aware of any specific

heritability estimates for PavCA. We used GCTA to calculate the proportion of the variance in

the base and composite PavCA metrics that could be explained by the set of SNPs unique to each

subgroup and the within-vendor SNP sets (S6 Table). Due to the low subgroup sample sizes, the

heritability estimates generally only reached significance when considering the vendor as a

whole. The SNP heritability estimates for all PavCA metrics in Harlan ranged from ~4-11%,

whereas they were ~4-21% for Charles River; on average the estimated heritability was about

~1.9-fold greater in Charles River. Importantly, some of the highest heritabilities were for

metrics used to designate sign-trackers vs. goal-trackers, such as the average of day 4 and day 5

response bias, probability difference, and PavCA index score. However, even the heritability

estimates from Charles River were lower than SNP heritability estimates for many other

behavioral traits [12,14,15,33].

Next, we performed GWAS for various metrics derived from Pavlovian conditioned

approach by using GEMMA to fit a linear mixed model that allowed us to account for population

structure with a relatedness matrix. Initially, we planned on performing the GWAS separately for

rats from Charles River vs. Harlan to maximize sample size; however, we observed marked

inflation in our Q-Q plots for a number of PavCA metrics (S5 Fig). This pattern of inflation

could reflect true signal that is amplified by the large LD-blocks characteristic of small breeding

populations -- if a locus shows significant association with a trait, all SNPs co-occurring in the

same LD-block will have inflated test statistics, thereby artificially enriching the Q-Q plot with

mid-range p-values. However, inflation in a Q-Q plot might also reflect a failure to account for

population structure Given the degree of the observed inflation, we suspected that the LMM

might have been insufficient to account for the residual subgroup structure within each vendor.

Therefore, we chose to perform individual GWAS on each subgroup followed by meta-analysis

of the results.

For each of the seven subgroups, all 11 PavCA metrics were examined for all five days of

training. Although PavCA data are often analyzed such that rats are categorized as sign-trackers,

goal-trackers, and intermediate responders, based on their PavCA performance on days 4 and 5

[26], we chose to analyze behavior on all 5 days since those results could yield insights into loci

involved in learning this complex behavior. Significance thresholds were determined using

MultiTrans, a parametric bootstrapping resampling approach [34]. Genome-wide significant

results from the individual subgroup GWAS are available in S1 File, and the Manhattan plots for

each analysis can be viewed in S2 File. The sample sizes for the subgroup analyses were small

and the GWAS yielded a marginal number of genome-wide significant results. In total, we

discovered six independent genome-wide significant hits across five of the seven subgroups,

representing PavCA metrics from days 3, 4, and 5 of training. All six significant hits were unique

to the subgroup they were identified in. This could be due to incomplete power, highly variable

allele frequencies between the groups, or the existence of unique modifier loci. Alternatively,

these hits may be artifactual and occurred by chance due to the number of analyses that were run.

Importantly though, we did not observe any significant inflation in the Q-Q plots for the GWAS

in these 7 subgroups (S3 File). To further assess how well we controlled population structure, we

used LD score regression [35]. We found that the average deviation from the expected intercept

was only 0.027 for the subgroups (S7 Table), indicating that dividing the samples into seven

subgroups was sufficient to control population structure.

To increase our power and improve our ability to identify true signals of association, we

meta-analyzed the results for all seven subgroups together, as well as in smaller sets of

subgroups derived from either Charles River or Harlan. The meta-analyses were run with MR-

MEGA [36], a program originally designed to accomplish trans-ethnic meta-analyses in

populations with potentially heterogeneous allelic effects, which is conceptually similar to this

dataset. Genome-wide significant results from the meta-analyses are presented in S1 File,

Manhattan plots in S4 File, and Q-Q plots in S5 File. There were five genome-wide significant

loci identified in the Harlan-specific meta-analyses, two in the Charles River-specific meta-

analyses, and five in the Harlan plus Charles River meta-analyses. Due to the sparsity of SNP

genotypes assayed by the ddGBS approach, these loci are generally composed of only a handful

of SNPs and their boundaries are not well defined. Notably, of the seven loci identified in the

vendor-specific meta-analyses, three were replicated in the Harlan plus Charles River meta-

analyses. The four loci that did not replicate were composed of SNPs that were unique to either

Charles River or Harlan and therefore could not have been replicated in the Harlan plus Charles

River meta-analyses.

One noteworthy finding was a locus identified in both the Charles River-specific and all-

subgroup meta-analyses. The two significant SNPs are located on chromosome 9 and are

associated with the training day 5 PavCA index score (Fig 4), which measures a rat's overall

propensity of a rat to attribute motivational value to a conditioned stimulus, and the day 5

probability difference, which directly measures the observed skew in responding to a conditioned

stimulus vs. a reward delivery cup (S4 File, page 35). Interestingly, the directions of effect for

the associated alleles were contrasting between the two vendors. In Charles River subgroups, the

associated allele had a negative effect on the trait, whereas in Harlan subgroups, the effect was

positive. This observation emphasizes the necessity of using a meta-analysis approach that

accounts for heterogenous allelic effects, as the association likely would not have replicated in

the broader subgroup meta-analysis with standard meta-analysis approaches. Presumably the

effects are opposite because the relationship between the SNP and the causal allele is reversed in

SD rats from Harlan relative to Charles River.

**Fig 4. Manhattan plots for the GWAS on PavCA Index Score on day 5 in each subgroup**

**and the meta-analysis.**

The titles above each of the eight Manhattan plots indicate the population, SNP count, and

sample size for each depicted GWAS. The y-axis is the $-\log_{10}$ of the p-value from the likelihood-

ratio test performed by GEMMA. The x-axis is the genomic coordinate of the variants within

each chromosome, in ascending order of chromosome.

Although the gene nearest the most associated SNP is frequently not the causal gene; we

examined the nearest gene to the chromosome 9 associated SNPs, which was *Efna5*, located

~500kb downstream. The gene *Efna5* codes for an ephrin-A ligand for EphA receptor tyrosine

kinases and is expressed in the brain. Ephrin-A5 has long been appreciated for its role in

guidance of retinal, cortical, and hippocampal axons during development [37–39]. However, this

gene continues to be expressed in the adult brain and has been shown to contribute to synaptic

plasticity of the mature hippocampus and long-term potentiation [38]. Experiments in mice have

shown that increased activation of EphA5, the receptor for Ephrin-A5, can lead to improved

memory [40] and that knocking out *Efna5* reduces anxious and aggressive behaviors [41,42].

In addition to *Efna5*, another late-stage PavCA training day association was identified on

chromosome 1 and was linked to the average latency to a lever contact (S4 File, page 4). The

associated SNP is located in an intron of *Park2*, a large 1.4Mb gene. This gene is most frequently

thought of as the causal locus for autosomal recessive Parkinson's disease in humans. However,

in rodents, *Park2* knockout lines do not display the same patterns of motor impairment and

neurodegeneration [43]. *Park2* rat knockout lines show reduced expression of dopaminergic

receptors, such as trace-amine receptor TAAR-1 and post-synaptic dopamine D2 receptor, in the

striatum [44]. Dopaminergic signaling is paramount in the establishment of incentive salience in

the Pavlovian conditioned approach paradigm [45,46], making this association an attractive

target for further investigation.

**Discussion**

Numerous behavioral and physiological studies use outbred Sprague Dawley rats

obtained from commercial breeders [47]. The vast majority of these studies are not designed with

genetics in mind. While SD and other outbred rats have been used for genetic selection [48–50],

ours is the first to explore the use of SD for GWAS. We adapted our genotyping methods to SD

and used them to densely genotype more than 4,000 SD rats. These data allowed us to

characterize the genetic background of SD rats and to perform GWAS for Pavlovian conditioned

approach behavior. This represents the largest rodent GWAS ever undertaken, and the first

performed using a commercially available outbred rat population.

We found dramatic genetic differences between SD rats obtained from Harlan versus

Charles River. $F_{ST}$ estimates show that SD rats from Harlan and Charles River are more diverged

than the major human ancestry groups [29] and nearly as diverged as some subspecies of mice

[51]. There was also strong evidence of population structure among the various breeding

locations and barrier facilities for each vendor. We found that SD rats from both Harlan and

Charles River both showed a rapid decay in LD.  However, at the time this study was performed,

SD from Charles River appear to have more polymorphisms and more favorable MAF and LD

profiles, suggesting that they would be preferable for future GWAS.

Although this study is the first to carefully document population structure within SD rats,

it is not the first to highlight phenotypic differences among SD from different vendors. In 1973,

Prejean et al. reported that the incidence of endocrine tumors varied among SD rats from

different vendors [1]. Then Clark et al. (1991) [52] reported differences in noradrenergic neural

projections among SD rats from different vendors. Subsequently, Turnbull & Rivier (1999) [4]

reported vendor-specific differences in the response to inflammatory stimuli. Then Fuller et al.

(2001) [53] reported vendor-specific differences in hypoxic response among SD rats. Even more

recently, there have been additional publications reporting differences for a variety of traits

among SD rats obtained from different vendors [2,5,54,55], and even suggesting that these

phenotypic differences may extend to differences among a single vendor's breeding facility [3].

Our own studies have previously reported both behavioral and genetic differences among SD rats

obtained from different sources [6], observations that are much more comprehensively explored

in the present dataset. Specifically, in addition to wide-spread genetic differences, we have also

shown that the SD rats obtained from Harlan for these studies show a much higher proclivity to

become sign-trackers compared to SD rats from Charles River. However, neither these prior

publications nor the current one can differentiate between two possibilities: that the observed

behavioral differences are the result of the different environment in which these animals are

raised versus the genetic difference that we have clearly demonstrated. This question could be

addressed by future studies in which SD rats are bred in the same facility and the offspring tested

in the same manner.

Using genome-wide genotype data, we have provided the first quantitative estimates of

the SNP heritability of the component measurements of PavCA. The highest heritabilities (16-

20% in Charles River) were seen for measures typically used to assess the propensity to attribute

incentive salience to reward cues, including the averages of the day 4 and day 5 PavCA index

score, response bias, and latency score. Previous work had shown that SD rats selectively bred

for ~15 generations for high or low responses to a novel environment were also highly divergent

for behaviors in PavCA [32]. Those selection studies demonstrate that SD rats have alleles that

influence PavCA. The present results are consistent with this conclusion. We obtained lower

heritability estimates for SD rats from Harlan compared to Charles River, further emphasizing

the genetic differences between SD rats from the two vendors.

As this was the first genome-wide association study to examine PavCA, we had to choose

from among many possible ways of summarizing the phenotype. Ultimately, we decided that the

analyses should be comprehensive. Therefore, we ran GWAS for all possible PavCA component

metrics since it was unclear which had the most predictive power with regard to the overarching

behavioral paradigm. In the individual subgroup GWAS, which had limited power due to sample

size, we identified only six genome-wide significant loci (alpha = 0.05) despite performing 385

total tests (5 training days x 11 PavCA metrics x 7 groups). Additionally, despite high

correlations between many of the metrics, some of the loci we detected were not found for any

other PavCA metric or training day. Furthermore, nearly half of the QTL identified in the

subgroup and meta-analysis GWAS occurred on earlier training days (days 1-3), while the

literature surrounding PavCA focuses on behavior after training (days 4/5). Fortunately, we did

observe replication of several associations in the meta-analyses across multiple PavCA metrics

(S1 File).

Utilizing the cost-efficient ddGBS to genotype SD rats allowed us to gather variant data

on several thousand samples. However, the power of this approach relies on the reasonably large

LD blocks that still exist in most commercially-available outbred rodent populations (Figure 3B).

Due to the sparsity and uneven distribution of genotypes assayed by ddGBS (Figure 1C), it is

difficult to pinpoint the exact source of the signals discovered via GWAS. Although the majority

of identified variants do not have clear functional implications, it is possible that these loci are

tightly linked to variants that influence the expression or function of nearby protein-coding

genes. Among the 46 significant associations from the full sample set meta-analyses (S1 File),

we have highlighted two loci. The first locus was discovered on chromosome 9 and was

associated with the PavCA index score on the final day of testing, when the rats' behaviors have

typically plateaued (Figure 1A). This locus was located ~500kb downstream of *Efna5* which

codes for an ephrin-A ligand for EphA receptor tyrosine kinases and has known ties learning and

memory through its effects on hippocampal development and synaptic plasticity [37–39]. The

second was an intronic variant in *Park2*. In rats, knocking out this gene has been shown to

reduce expression of dopaminergic receptors, notably TAAR-1, a post-synaptic dopamine D2

receptor. Seeing as dopamine is essential for the attribution incentive salience to reward-

associated cues [45,46,56], this result is an enticing one for future follow-up studies.

Because our project used samples from an ongoing, non-genetically focused project, we

inherited a design that used Sprague Dawley from many different sources. The full impact of this

was not clear until we completed genotyping the rats. The population structure that we

encountered was similar to human genetic studies that include individuals from multiple

ancestries, and we sought to address it by analyzing each group separately (with an LMM) and

then combining the results using meta-analysis (e.g. [57]). In humans, different ancestry groups

(e.g. East Asian vs. European) often do not share the same causal variants. Similarly, our power

was likely reduced because causal variants were not shared between the subgroups. Modifiers of

causal variants might also be dissimilar between the various subgroups, further hindering our

meta-analyses. [58]

Regardless of whether differences in SD rats obtained from different vendors are due to

genetic or environmental differences, our results demonstrate the need for much greater care in

the use of SD rats. Even when the primary research question is non-genetic, researchers should

carefully consider whether to use rats from a single vendor and/or breeding facility. Ignoring the

genetic background of test subjects can easily confound experimental results. The exact sources

(vendor, breeding facility, barrier within each breeding facility) should also be carefully

documented and clearly reporting in the results section. Failure to do so can lead to problems

with replication, since observations made in SD rats from one source may not extend to SD rats

from other sources. In addition to non-replication, a more subtle consequence of using SD rats

from multiple vendors or breeding facilities is that spurious correlations can occur. For example,

if SD rats from Charles River were higher for traits A and B, compared to SD rats from Harlan, a

heterogeneous cohort of SD rats from both vendors would show a significant positive correlation

between traits A and B. Such a correlation may not be due to any shared biological mechanisms

but could instead be the result of either genetic population structure or environmental differences

between SD rats from the two vendors. Thus, even studies that are not intended to address

genetics questions could lead to incorrect inferences because of genetic differences among SD

rats obtained from different sources.

Our study is the first to use SD rats for GWAS of a complex trait and is the largest rodent

GWAS ever undertaken. We have exposed extensive population structure among SD rats that has

important implications for genetic and non-genetic uses of SD rats. Based on our data, we

suggest that Charles River SD rats have greater genetic diversity, finer scale LD and more

favorable MAF, and thus are an attractive choice for future GWAS using outbred, commercially

available rats. Although our GWAS was hampered by population structure, our results have

identified several genome-wide significant loci that provide the first insights into the genetic

basis for PavCA, a trait that is relevant to understating motivational processes, especially in the

context of substance abuse.

## Data Availability

Data files that were too large to be included as supplemental content are to be uploaded to a

central online repository through the UCSD Library System: https://doi.org/10.6075/J0XS5V8K.

Data will be available after publication. The DOI will be able to be followed for access to (1) a

VCF file containing all raw SNP calls with dosage data in the original set of 4,625 Sprague

Dawley rats, (2) filtered variants in BIM/BED/FAM format for each of the seven SD subgroups,

and (3) summary files for the GWAS results for all 55 metric x day combinations for all

analyses, individual subgroups and meta-analyses. In addition, FASTQ files from the original

ddGBS and light WGS sequencing runs will be submitted to the NCBI SRA. Please contact the

authors for more information on accessing this sequencing data. These data, along with the

supplemental files and detailed methods included, are sufficient to replicate the analyses

performed in this publication.

## Methods

**Sprague Dawley samples**

Tissue samples from 5,206 male Sprague Dawley rats were obtained, predominantly from

Charles River and Harlan, with a few samples from Taconic. A subset of 4,625 of these rats went

on to be genotyped by ddGBS and/or WGS. After sample filtering, a final set of 4,061 genotyped

SD rats were used for the population genetic and association analyses. S1 Table lists the number

of many samples that came from each vendor, breeding location, and barrier facility. Detailed

information about these 4,061 rats is available in S6 File. Behavioral testing was performed

between February 2012 and August 2015 as part of work for multiple studies [25,59–65,65–71].

All experiments were approved by the University of Michigan IACUC. Housing, feeding,

lighting and other relevant environmental conditions have been previously described. Following

sacrifice at the University of Michigan, tissue samples were shipped to the University of

Chicago; subsequent processing of those samples is described in the following sections.

**Pavlovian conditioned approach**

Pavlovian conditioned approach procedures have been thoroughly described previously

[72,73] as a means to assess the tendency to attribute incentive motivational value or incentive

salience to a cue that has been repeatedly paired with a noncontingent reward. Briefly, rats are

placed into a testing chamber in which an illuminated lever (conditioned stimulus; CS) enters the

chamber and after 8 seconds the lever-CS retracts and a food pellet (unconditioned stimulus; US)

is immediately delivered into an adjacent food cup.  Rats are scored for their three possible

responses to the lever-CS entering the cage: approach and interact with the lever, approach and

interact with the food receptacle (magazine), or make neither approach. Conditioned responses

are captured during the 8-second period during which the lever-CS has entered the chamber, but

before the food reward enters the magazine. The following measures are obtained in automated

fashion: the number of lever contacts as measured by lever depressions, number of magazine

entries as measured by infrared sensor in the food receptacle, and the latency to both during the

8-second lever-CS presentation. The rats are tested in this manner with 25 trials per session, and

one session is conducted per day for 5 consecutive days. For the purposes of this project, the

number of lever contacts and magazine entries are summed across all 25 trials within a given

session, and the latencies are averaged across 25 trials within a session.

Along with response counts and latencies, three additional measurements are recorded: 1)

the proportion of trials in a session during which a rat made a lever contact ("probability" of

lever press), 2) the proportion of trials during which they made a magazine entry ("probability"

of magazine entry), and 3) the number of non-CS (NCS) magazine entries that occurred outside

of the 8 second trials (when the cue was not present during the intertrial interval). We also

calculated composite scores to categorize rats as sign-trackers (ST; defined as rats that

preferentially interacted with the lever-CS), goal-trackers (GT; defined as rats that preferentially

interacted with the food magazine), and intermediate responders (IR; rats that vacillated between

sign- and goal-tracking behavior) [26]. These scores include: response bias ([lever presses –

magazine entries]/[lever presses + magazine entries]), latency score ([average magazine entry

latency – average lever press latency]/8), and probability difference ([lever press probability –

magazine entry probability]). The PavCA index score is the average of the response bias, latency

score, and probability difference. A value of [-1, -0.5] for the PavCA index score indicates a GT,

(-0.5, 0.5) indicates an IR, and [0.5, 1] indicates a ST. We performed a Welch's 2-sample t-test

to show that the PavCA index score distributions differed significantly between Charles River

and Harlan SD rats (t = 20.161, df = 3908.1, p-value $< 2.2 \times 10^{-16}$). In summary, 11 PavCA

metrics were available for analysis, each of which we measured on days 1, 2, 3, 4, and 5 (S8

Table). Phenotypic data is available in S6 File.

**Double digest genotype-by-sequencing (ddGBS)**

To obtain genotypes, we used ddGBS, a genotyping method that reduces the complexity

of the genome by only sequencing regions proximal to restriction enzyme cut sites [6,74]. We

have recently described the technical aspects of this protocol in detail [75]. The ddGBS protocol

used in this paper is a synthesis of the GBS approach described in Graboski et al. [76] and used

more recently by Parker et al. [14] and Gonzales et al. [33], and an analogous approach known as

double digest restriction associated DNA sequencing (ddRADseq) [77]. The full protocol is

available in S1 Text.

DNA was extracted from rat tails using the PureLink® Genomic DNA kit. DNA purity

was assayed using a Nanodrop 8000 (260/280 ≥ 1.8) and DNA integrity by gel electrophoresis

(minimal smearing). Genomic DNA was then digested using two restriction enzymes: *PstI* (6-bp

recognition site) and *NlaIII* (4-bp recognition site). Adapter oligos were annealed to overhangs

left by *Pst1* and *NlaIII*. The *PstI* adapters contained 48 unique 4-8 bp in-line indexes [14,33,76].

A Y-adapter was annealed to the *NlaIII* cut sites, which controlled the direction of the first round

of PCR amplification and thus ensured that the library was primarily composed of fragments

with one of each of the adapters. Post-annealing, sets of 24 individual sample libraries were

quantified and pooled. Pooled libraries were PCR amplified for 12 cycles, size-selected for 300-

450bp using the Pippin Prep, and quality checked by Agilent Bioanalyzer (peak range ~ 300-

500bp and conc. ≥ 20nM). Sequences for the 48 barcoded adapters, Y-adapter, and PCR primers

are provided in S2 Text.

Sequencing of pooled libraries was performed by Beckman Coulter Genomics (now

GENEWIZ). Sequencing was carried out on the Illumina HiSeq 2500 using v4 chemistry and

125-bp single-end reads. Each lane consisted of a pool of 24 samples, resulting in an average of

8.9 million reads per sample. A total of 4,608 unique ddGBS sample libraries were sequenced.

Of these samples, 381 were sequenced twice, resulting in two sets of sequencing data for each

sample from the same library prep that were used for to check genotype concordance (S3 Table).

**Light whole-genome sequencing (WGS)**

To discover new variants and support imputation, we performed low-coverage whole-

genome sequencing of 80 SD rats from this same cohort. The rats were selected to represent

sign-trackers, goal-trackers, and intermediate responders from each of the major barriers within

the 6 major subgroups of Harlan and Charles River. Sample libraries were prepared using the

Illumina TruSeq® PCR-Free Library Prep kit and quality checked using an Agilent Bioanalyzer

and qPCR on an Applied Biosystems StepOne Real-Time PCR System to ensure they met

Illumina quality standards. Sample pooling (10 samples per pool) was performed by Beckman

Coulter Genomics. Each pooled library was sequenced on two lanes of an Illumina 2500 flow

cell with 125-bp single-end reads, resulting in an average of 51.6 million reads per sample per

two lanes. Assuming that the rat genome (rn6) is ~2.87Gb, this provided about 180x coverage of

the rat genome, or about 2.25x coverage per rat.

**ddGBS Sequence data processing**

We have recently described the bioinformatic steps that we use for ddGBS in detail [75].

We follow an analogous approach in this paper, though we deviate at the imputation step due to

our use of SD instead of HS rats. Briefly, raw reads from ddGBS were demultiplexed using

FASTX Barcode Splitter [78], allowing for 1 mismatch. After demultiplexing, barcodes were

trimmed by *cutadapt* [79]. Any reads not matching a sample's barcode within 1-bp were filtered

out. We removed 316 samples for which there were less than 4 million reads, leaving 4,292

samples with ddGBS data. We also used *cutadapt* to trim low-quality base pairs (phred quality

score < 20) at the ends of the reads, and to remove 3' adapter sequences. Reads trimmed to less

than 25-bp were discarded. Next, all reads were aligned to the rat reference genome assembly

(rn6) using *bwa* [80]. All ddGBS reads were realigned at known indel sites by GATK's

IndelRealigner [81]. Because of a lack of SD-specific variant data, we used variant data from 42

whole-genome sequenced rat strains and substrains [82] as the reference set for indels. We then

used GATK to perform base quality score recalibration (BQSR) on the BAM files, using data

(SNP & indel) from the 42 rat genomes as the "known" set of variants. For the ddGBS samples

that were sequenced twice (381 remaining after filtering for read count), we performed all

quality control and variant calling steps in parallel, since our goal was to compare calls made in

these samples as a means of estimating the genotyping error rate.

**Light WGS data processing**

Raw reads from WGS were processed in an analogous manner to the ddGBS data

(detailed above) through the alignment step. After alignment, duplicates reads were removed

using *picard* [83]. Reads were then realigned and underwent BQSR. The final WGS BAM files

from each lane (2 files per sample) were merged. The WGS BAM files for the 63 samples that

had undergone both ddGBS and WGS were then merged.

**Variant discovery and imputation – ANGSD/Beagle**

We found that GATK's HaplotypeCaller tool [81] was ineffectual at making high-

confidence SNP calls in our dataset, likely due to the unusual distribution of reads produced by

ddGBS. Instead, we used the Samtools variant calling model [84] as implemented by ANGSD

[85] to estimate genotype likelihoods from the mapped ddGBS reads. Likelihoods were obtained

in 10Mb chunks of the genome, which were subsequently concatenated. Major and minor alleles

were inferred from the data based on allele frequency estimates made from the genotype

likelihoods. The likelihoods were only estimated at sites where at least 100 samples had reads.

ddGBS data results in low call rates at many loci. However, we retained these loci because we

anticipated they would be useful for imputation. Next, we used the ANGSD genotype likelihoods

to impute missing genotypes (that is for SNPs where only a portion of the rats had genotyping

information) using Beagle [86,87], which produced a VCF file containing hard genotype calls

(0,1, or 2), dosages ([0,2]), and posterior probabilities for each genotype ([0,1]) for 2,274,118

biallelic SNPs in 4,309 rats (ddGBS+WGS). This is the unfiltered set of SNPs and samples we

moved forward with for all subsequent steps. We elected not to pursue variant quality score

recalibration using the GATK VariantRecalibrator algorithm [88], because we did not have the

required "truth" SNP set. Due to the poorly understood population history of SD rats, it was

unclear whether the variation present in the 42 rat genomes would be representative of the

variation present in our sample. Using the 42 genomes as a reference for the VariantRecalibrator

may also have negatively impacted the calling of novel SD variants.

**STITCH (Sequencing To Imputation Through Constructing Haplotypes)**

In addition to variant calling and imputation using ANGSD/Beagle, we also explored the

use of STITCH [27], since it is specifically designed for low-coverage WGS data lacking

haplotype reference panels. However, ddGBS data is higher coverage and sparser than the input

for which STITCH was designed. We used the set of alignment files and known variant sites

from the 42 genomes [82], as described above.  STITCH queries the user for the number of

ancestral haplotypes that exist in the population (K). Due to our lack of knowledge about the

founder population, we ran STITCH on a single chromosome using 5 different values of K: 2, 3,

4, 5, and 6. We found that K values of 3, 4, and 5 worked similarly well and chose K=4 to

maximize the number of SNPs called and minimize error rate as ascertained by comparison of

genotype calls between duplicate samples. STITCH yielded 8,691,886 biallelic SNPs, vastly

more than were called with ANGSD/Beagle. However, after applying filters for dosage $r^2$/INFO

score $\geq 0.9$, MAF $\geq 0.01$, HWE p-value $\leq 1\text{x}10^{-7}$, as well as removing sites in near perfect

pairwise LD $> 0.95$, we found that the genotypes from STITCH contained fewer SNPs compared

to the comparably filtered output from ANGSD/BEAGLE (see S2 Table). For this reason, we did

not use the SNP calls made by STITCH in any of the analyses presented in this paper.

**Genotype concordance check**

Whereas some of our past projects that used GBS were able to determine the accuracy of

GBS genotypes by comparing them to genotypes obtained from SNP microarrays, we did not

have microarray-based genotypes for this cohort. Instead, we relied on the remaining 381

duplicate samples whose genotypes were called in parallel. To estimate genotyping accuracy, we

compared the rate of concordance of hard genotype calls among the duplicate samples. We first

filtered variants by dosage $r^2$ ($DR^2$), a measure of the accuracy of the genotype imputation

performed by Beagle. We tested three different $DR^2$ thresholds ($\geq 0.7$, $\geq 0.8$, $\geq 0.9$). We then

removed variants with MAFs < 1% or that violated Hardy-Weinberg equilibrium at a threshold

of $1 \times 10^{-7}$ in either vendor population. Concordance rates were checked by two methods: 1) by

using hard calls in the RAW format from *plink 1.9* and dividing the number of times a genotype

call matched between duplicate samples by the total number of SNPs and 2) by taking the mean

Pearson correlation of the dosages of the duplicate samples. The results are presented in S3

Table. Similar error estimates were obtained by the hard call and dosage approaches. We chose

to move forward with the $DR^2$ threshold of 0.9 for all subsequent analyses, which yielded an

error rate of ~0.85%.

**Post-genotyping sample filtering**

We removed female rats (n=77) and rats from Taconic Farms (n=4) since they made up a

very small fraction of the total sample. We also removed rats that showed poor clustering in the

PCA analysis, described below. We filtered out individuals with unusually high or low rates of

heterozygosity and high degrees of relatedness as detailed below. Lastly, we excluded a small set

of duplicate samples (n=7) and samples missing phenotype data for mapping (n=10). All filters

and sample numbers are listed in S5 Table. After these steps, 4,061 unique male SD rats from

Charles River (n=1,780) and Harlan (n=2,281) remained.

**Principal component analysis, identity-by-descent, and heterozygosity**

We performed principal component analysis (PCA) on the cohort of 4,228 samples

filtered for low read count, having been received from Taconic, and females. PCA was

performed on genotype calls in R using the *prcomp* function in R [89] on a set of variants pruned

for SNPs with MAF ≤ 0.05 in the combined sample set, SNPs in pairwise LD > 0.5, and SNPs

violating HWE at a p-value $< 1 \times 10^{-7}$ in either Charles River or Harlan. The first PC clearly

separated animals from Harlan and Charles River; however, there was a set of 54 rats that did not

visually cluster as expected at the level of vendor (data not shown). These animals were removed

from all subsequent analyses (we assumed they reflected inaccurate records, sample mix-ups, or

some other technical problems).

With this further reduced set of 4,174 rats, we continued on to assess the genetic

relationships among the rats in our sample. SD rats were ordered in multiple batches over several

years, and we suspected some of these rats would be closely related (siblings, cousins, etc.). We

reapplied the variant filters used for PCA and utilized the *--genome* function in *plink 1.9* on the

resulting SNP set to estimate $\hat{\pi}$ (the proportion of genotypes predicted to be identical by

descent), for all pairs of samples. Panels A and C in S4 Fig show that while most of the animals

were unrelated, there were a significant subset of closely related pairs, as well as some pairs with

exceedingly high IBD1 rates in Harlan.

We used the *plink 1.9* function *--het* to examine possible inflation or deflation of

heterozygosity rates in our samples. Panels A and C in S6 Fig show that a handful of samples in

both populations with uncharacteristically high (> 0.25) or low (< 0.25) rates of heterozygosity.

We filtered out 34 such samples, as we found they drove the majority of the anomalous signal in

our pre-filtering plots in S4 Fig. We also removed 12 samples with more than 30 close relations

(defined as $\hat{\pi} \geq 0.1875$) and 32 samples that had a $\hat{\pi} \geq 0.6$ with another sample (likely sample

contamination/mix-up). After applying these sample filters, we were left with Panels B and D in

S4 Fig and S6 Fig.

After reaching our final set of 4,061 rats, we reapplied the SNP filters on the reduced

sample set, resulting in 4,502 SNPs that were polymorphic in both populations. We ran PCA on

these SNPs as described above. We then repeated PCA on the samples from Charles River and

Harlan separately to examine substructure within each vendor population. Using the results from

the vendor-level PCA analysis, new groupings of barrier facilities were determined based on

genetic similarity. Barrier facilities that clustered most highly in the PCA were grouped together

for subsequent analysis.

**Final variant filtering and minor allele frequency spectrum**

After establishing the final cohort of 4,061 rats, we sought to establish a final set of SNPs

to be used for the association analyses. Starting from the initial full set of 2,274,118 SNPs, we

first removed sites with $DR^2 < 0.9$. Then, based on the PCA results, we separated the samples

into seven subgroups, each containing genetically-similar samples that clustered highly: SD rats

originating from Harlan 202A/202C/208A (Har 202A), Harlan 206, Harlan 217, Charles River

R09/P03/P07/P10/C71/K92 (Charles River R09), Charles River R04, Charles River P09, and

Charles River C72. These seven groupings were used for all subsequent variant filtering

performed in *plink 1.9*, which converts the posterior probabilities we received from Beagle to

hard genotype calls, so long as the probability is ≥ 0.9. In each cluster, we removed SNPs with

MAF < 0.0001 using the *–maf* option in *plink 1.9*. Lastly, we used the *--hardy* function in *plink*

*1.9* [90] to perform tests for SNP Hardy-Weinberg equilibrium and removed all SNPs with an

HWE p-value $< 1\times10^{-7}$ in any of the seven groups. The final counts of samples and SNPs in each

barrier grouping are available in S3 Table. We used the *--freq* option in *plink 1.9* to estimate the

MAFs for these final filtered sets SNPs. The distributions are show in (Fig 3A).

Taking the union of these seven sets of SNPs, there were a total of 291,438 polymorphic

sites passing our filters across all subgroups. Our final set of filtered SNPs contained 75,405

novel SNPs that had not be previously reported by Hermsen et al. [82]. We also compared our

final set of SNPs to the most recent dbSNP release for rats (Build 149, November 7, 2016) and

found that 178,545 of the 291,438 SNPs we discovered were not present in the current dbSNP

build. See the Data Availability section for more information on accessing this data.

**Fixation Index ($F_{ST}$)**

To quantify the population divergence between SD rats from Harlan and Charles River,

we computed $F_{ST}$ for each SNP in the union set using *smartpca* within the *EIGENSOFT* package

[29,91,92]. Due to the substructure within vendor and vendor location that we saw in our PCA

analyses, we chose to estimate the pairwise $F_{ST}$ between all seven sample subgroups defined

above.

**Linkage Disequilibrium**

We plotted the decay of linkage disequilibrium (LD) using the *--$r^2$* utility in *plink 1.9* and

the procedures described in Parker et. al 2016 [14]. Briefly, each population's curve included all

SNPs with MAF > 20% and pairwise LD comparisons were restricted to SNPs with allele

frequencies within 5% of each other. An average $r^2$ estimate was obtained using 10,000

randomly selected SNP pairs from each 100kb interval for the distance between two SNPs,

starting with 0-100kb and end at 9.9-10Mb. The procedure was repeated for cluster grouping.

The CFW mice curve was downloaded from a repository for Parker et al.

(http://datadryad.org/resource/doi:10.5061/dryad.2rs41) [14] and used for comparison as another

commercially available, outbred rodent stock.

**LMM covariates and phenotype data pre-processing**

To select covariates for the GWAS, we performed univariate linear regression for each

potential covariate for each PavCA metric. This was done separately for rats from each

genetically-similar cluster defined previously. Any covariates that accounted for 1% or more of

the variance for at least one PavCA metric were passed into multivariate model selection with

the R package *leaps* [93]. Model selection with *leaps* was performed for all metrics for all days

of testing, as well as the average of days 4 and 5. Out of the 66 models, all covariates that had

surpassed the 1% threshold to reach this step were ultimately selected in at least 40% of the *leaps*

models. The covariates included in the for the downstream GWAS analyses were rat's age at

testing, housing (binary – single or multiple), light cycle (binary – standard or reverse), and a set

of binary 'indicator' variables to model the effects of different experimenters/technicians

responsible for the phenotypic testing (S9 Table). All covariates were included in the LMMs for

association testing by GEMMA, rather than being regressed out prior to GWAS.

Many of the PavCA metrics were exceedingly non-normally distributed. In most cases,

this was expected due to how the behaviors were measured and defined. For example, the rats

only had a window of 8 seconds in each trial during which to contact the lever and/or make a

nose poke into the food magazine. All values for "average latency to lever press" or "average

latency to magazine entry" were therefore necessarily between 0 and 8 seconds. As is typical for

latency traits, many of the values were near 0 or exactly 8. Similarly, the "probability" of lever

press or magazine entry were very skewed towards the limits 0 and 1, especially after

conditioned responding had been established on the later test days of training. Given these

unusual distributions, we chose to quantile normalize all metrics within each subgroup prior to

association testing, accepting a possible loss in power since samples with identical values are

ranked randomly during the quantile normalization procedure.

**SNP-based heritabilities**

Heritabilities were estimated separately for each vendor and subgroup using their sets of filtered SNPs. We used the SNP sets to construct genetic relationship matrices (GRMs) for each cluster using GCTA [94]. We then used the restricted maximum likelihood (REML) approach within GCTA on the GRMs, covariates, and quantile normalized PavCA data to calculate the SNP-based heritabilities for each metric on each testing day.

**GWAS**

We used GEMMA [95], which implements an LMM for GWAS analysis. We included a GRM as a random term to account for relatedness and population structure. Though beneficial for preventing false positive associations, GRMs can also reduce power to detect QTLs in populations with greater levels of LD; this is due to proximal contamination [96,97]. To avoid this reduction in power, we used the leave-one-chromosome-out (LOCO) approach when constructing the GRMs for each cluster of samples [33,98,99]. As described above, we selected covariates that were included as fixed effects in our model (S9 Table). Any samples missing measurements for a given metric were removed from that analysis. For all GWAS, genotypes were represented as dosages (continuous [0,2]) in lieu of 'hard' genotype calls [0, 1, 2] to account for uncertainty in the genotype calls. Reported p-values come from the likelihood-ratio test (LRT) performed by GEMMA. Results were plotted using a custom R script.

**GWAS meta-analysis**

We performed meta-analyses of the GWAS results across all seven subgroups, across only the four Charles River subgroups, and across only the three Harlan subgroups. Due to the significant genetic differentiation between certain subgroups in this meta-analysis, it is possible that tested loci would have differing effect sizes and directions of effect. To account for this and

maximize our power, we utilized the software MR-MEGA, originally designed to perform trans-

ethnic meta-regression in humans [36]. A large number of the SNPs called in each subgroup

were either unique to that singular group or a subset of the groups, making them of low utility in

a meta-analysis of all seven subgroups. Therefore, the seven subgroup meta-analyses were

performed on the intersection SNPs set for all seven clusters, amounting to 64,442 SNPs. We

repeated the meta-analyses for the only the subgroups derived from Charles River (R04, R09,

P09, C72) and only those from Harlan (202A, 206, 217) to allow for more SNPs to pass through

to the meta-analysis. There were a total of 198,817 shared SNPs between the Charles River

subgroups, but only 83,120 shared between the Harlan subgroups. Genome-wide significant

results from these meta-analyses are presented in S1 File. For all significant hits, we also report

the directionality and strength of effect of the SNPs in all the separate subgroups GWAS.

**P-value significance thresholds**

In human GWAS, $5 \times 10^{-8}$ is widely used as a significance threshold [100]. However,

model organisms have vastly different levels of LD, meaning that the effective number of

independent tests differs between studies. Therefore, many prior studies have used permutation

testing [101,102], where phenotype data is shuffled with respect to fixed genotypes. The

computational load of such methods becomes intractable for studies with large sample sizes and

several traits being mapped [96]. Thus, we used the sequence of SLIDE [103,104] and

MultiTrans [34] to obtain significance thresholds. We used separate thresholds for each subgroup

because the number of SNPs, the LD structure, and the marker-based heritabilities were

different, all of which affect the estimated number of independent tests performed as part of the

GWAS. An advantage of this approach is that only one threshold was needed per cluster for all

normalized metrics. We used a sliding window of 1000 SNPs and sampled from the multivariate

normal 10 million times to obtain a 0.05 significance threshold. The subgroup-level $-\log_{10}(p)$

thresholds for Charles River R04, R09, P09, and C72 were determined to be 6.14, 6.13, 6.11, and

6.11, respectively. Similarly, the Harlan 202A, 206, and 217 thresholds were 6.09, 6.00, and

5.89, respectively. Since there is no clear extension of this approach for a meta-analysis with

largely varying sets of SNPs and LD patterns, we ultimately obtained the meta-analysis

thresholds by taking the subgroup from each vendor with the lowest level of LD (Charles River

R04 & Harlan 202A) and filtering their SNP set to contain only the SNPs included in each meta-

analysis. We then reran MultiTrans using these pruned sets of SNPs, providing us with a $-\log_{10}$

threshold of 5.574 for the over-arching meta-analyses, 5.574 for the within-Harlan meta-

analyses, and 5.993 for within-Charles River.

**LD score regression**

We utilized LD score regression to more rigorously test whether after dividing samples

up into seven subgroups, we were still observing inflation of the GWAS test statistics. LDSC is

software designed to determine if observed inflation is due to cryptic relatedness and population

stratification or true polygenic signal spread across regions of high LD [35]. Since there is no

genetic map available for Sprague Dawley rats, we used a map created for heterogeneous stock

(HS) rats [105], a similarly outbred population that shares some ancestry with SD. An LD

window size of 5 centimorgans was used, corresponding to the point at which SD LD decayed

sufficiently in most subgroups for loci to be considered unlinked. The results of running LDSC

for all GWAS in each subgroup are available in S7 Table.

**Acknowledgements**

We would like to thank John Novembre, Mark Abney, and Dan Nicolae at the University

of Chicago for their advice concerning the statistical analyses involved in this publication.

**References**

1. Prejean JD, Peckham JC, Casey AE, Griswold DP, Weisburger EK, Weisburger JH. Spontaneous tumors in Sprague-Dawley rats and Swiss mice. Cancer Res. 1973;33: 2768–2773.

2. Bodnar TS, Hill LA, Taves MD, Yu W, Soma KK, Hammond GL, et al. Colony-Specific Differences in Endocrine and Immune Responses to an Inflammatory Challenge in Female Sprague Dawley Rats. Endocrinology. 2015;156: 4604–4617. doi:10.1210/en.2015-1497

3. Brower M, Grace M, Kotz CM, Koya V. Comparative analysis of growth characteristics of Sprague Dawley rats obtained from different sources. Lab Anim Res. 2015;31: 166. doi:10.5625/lar.2015.31.4.166

4. Turnbull AV, Rivier CL. Sprague-Dawley Rats Obtained from Different Vendors Exhibit Distinct Adrenocorticotropin Responses to Inflammatory Stimuli. Neuroendocrinology. 1999;70: 186–195. doi:10.1159/000054475

5. Weber K. Differences in Types and Incidence of Neoplasms in Wistar Han and Sprague-Dawley Rats. Toxicol Pathol. 2017;45: 64–75. doi:10.1177/0192623316672075

6. Fitzpatrick CJ, Gopalakrishnan S, Cogan ES, Yager LM, Meyer PJ, Lovic V, et al. Variation in the Form of Pavlovian Conditioned Approach Behavior among Outbred Male Sprague-Dawley Rats from Different Vendors and Colonies: Sign-Tracking vs. Goal-Tracking. Campolongo P, editor. PLoS ONE. 2013;8: e75042. doi:10.1371/journal.pone.0075042

7. Dwinell MR, Worthey EA, Shimoyama M, Bakir-Gungor B, DePons J, Laulederkind S, et al. The Rat Genome Database 2009: variation, ontologies and pathways. Nucleic Acids Res. 2009;37: D744-749. doi:10.1093/nar/gkn842

8. Drinkwater NR, Gould MN. The long path from QTL to gene. PLoS Genet. 2012;8: e1002975. doi:10.1371/journal.pgen.1002975

9. Bryant CD, Smith DJ, Kantak KM, Nowak TS, Williams RW, Damaj MI, et al. Facilitating Complex Trait Analysis via Reduced Complexity Crosses. Trends Genet TIG. 2020;36: 549–562. doi:10.1016/j.tig.2020.05.003

10. Woods LCS, Mott R. Heterogeneous Stock Populations for Analysis of Complex Traits. In: Schughart K, Williams RW, editors. Systems Genetics. New York, NY: Springer New York; 2017. pp. 31–44. doi:10.1007/978-1-4939-6427-7_2

11. Parker CC, Chen H, Flagel SB, Geurts AM, Richards JB, Robinson TE, et al. Rats are the smart choice: Rationale for a renewed focus on rats in behavioral genetics. Neuropharmacology. 2014;76: 250–258. doi:10.1016/j.neuropharm.2013.05.047

12. Rat Genome Sequencing and Mapping Consortium, Baud A, Hermsen R, Guryev V, Stridh P, Graham D, et al. Combined sequence-based and genetic mapping analysis of complex traits in outbred rats. Nat Genet. 2013;45: 767–775. doi:10.1038/ng.2644

13. Chitre AS, Polesskaya O, Holl K, Gao J, Cheng R, Bimschleger H, et al. Genome-Wide
Association Study in 3,173 Outbred Rats Identifies Multiple Loci for Body Weight,
Adiposity, and Fasting Glucose. Obes Silver Spring Md. 2020;28: 1964–1973.
doi:10.1002/oby.22927

14. Parker CC, Gopalakrishnan S, Carbonetto P, Gonzales NM, Leung E, Park YJ, et al.
Genome-wide association study of behavioral, physiological and gene expression traits in
outbred CFW mice. Nat Genet. 2016;48: 919–926. doi:10.1038/ng.3609

15. Nicod J, Davies RW, Cai N, Hassett C, Goodstadt L, Cosgrove C, et al. Genome-wide
association of multiple complex traits in outbred mice by ultra-low-coverage sequencing. Nat
Genet. 2016;48: 912–918. doi:10.1038/ng.3595

16. Zou J, Gopalakrishnan S, Parker CC, Nicod J, Mott R, Cai N, et al. Analysis of independent
cohorts of outbred CFW mice reveals novel loci for behavioral and physiological traits and
identifies factors determining reproducibility. Bioinformatics; 2021 Feb.
doi:10.1101/2021.02.05.429998

17. Charles River. CD® (Sprague Dawley) IGS Rat Details. In: CD® (Sprague Dawley) IGS Rat
[Internet]. [cited 2 Jul 2018]. Available: https://www.criver.com/products-services/find-
model/cd-sd-igs-rat?region=3611

18. Envigo, Inc. Sprague Dawley® outbred rat | Envigo. In: Envigo [Internet]. [cited 2 Jul 2018].
Available: https://www.envigo.com/products-services/research-models-
services/models/research-models/rats/outbred/sprague-dawley-outbred-rat/

19. Charles River Laboratories International, Inc. International Genetic Standardization (IGS)
Program. 2016. Available:
https://www.criver.com/sites/default/files/Technical%20Resources/Charles%20River%20Int
ernational%20Genetic%20Standardization%20Program.pdf

20. White WJ, Lee CS. The Development and Maintenance of the Crl:CD(SD) IGS BR Rat
Breeding System. 1998. Available:
https://www.crj.co.jp/cms/cmsrs/pdf/company/rm_rm_a_igs_rat_breeding_system.pdf

21. Weber K, Razinger T, Hardisty JF, Mann P, Martel KC, Frische EA, et al. Differences in Rat
Models Used in Routine Toxicity Studies. Int J Toxicol. 2011;30: 162–173.
doi:10.1177/1091581810391818

22. Envigo, Inc. Genetic integrity assurance program | Envigo. In: Envigo [Internet]. 2 Jul 2018
[cited 2 Jul 2018]. Available: https://www.envigo.com/products-services/research-models-
services/resources/genetic-integrity-assurance-program/

23. Poiley SM. A systematic method of breeder rotation for non-inbred laboratory colonies. Proc
Anim Care Panel. 1960;10: 159–166.

24. Flagel SB, Akil H, Robinson TE. Individual differences in the attribution of incentive
salience to reward-related cues: Implications for addiction. Neuropharmacology. 2009;56:
139–148. doi:10.1016/j.neuropharm.2008.06.027

25. Robinson TE, Yager LM, Cogan ES, Saunders BT. On the motivational properties of reward
cues: Individual differences. Neuropharmacology. 2014;76: 450–459.
doi:10.1016/j.neuropharm.2013.05.040

26. Meyer PJ, Lovic V, Saunders BT, Yager LM, Flagel SB, Morrow JD, et al. Quantifying
Individual Variation in the Propensity to Attribute Incentive Salience to Reward Cues.
Zhuang X, editor. PLoS ONE. 2012;7: e38987. doi:10.1371/journal.pone.0038987

27. Davies RW, Flint J, Myers S, Mott R. Rapid genotype imputation from sequence without
reference panels. Nat Genet. 2016;48: 965–969. doi:10.1038/ng.3594

28. Holsinger KE, Weir BS. Genetics in geographically structured populations: defining,
estimating and interpreting FST. Nat Rev Genet. 2009;10: 639–650. doi:10.1038/nrg2611

29. Bhatia G, Patterson N, Sankararaman S, Price AL. Estimating and interpreting FST: The
impact of rare variants. Genome Res. 2013;23: 1514–1521. doi:10.1101/gr.154831.113

30. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: A
Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. Am J
Hum Genet. 2007;81: 559–575. doi:10.1086/519795

31. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. GigaScience. 2015;4.
doi:10.1186/s13742-015-0047-8

32. Flagel SB, Robinson TE, Clark JJ, Clinton SM, Watson SJ, Seeman P, et al. An Animal
Model of Genetic Vulnerability to Behavioral Disinhibition and Responsiveness to Reward-
Related Cues: Implications for Addiction. Neuropsychopharmacology. 2010;35: 388–400.
doi:10.1038/npp.2009.142

33. Gonzales NM, Seo J, Hernandez-Cordero AI, St. Pierre CL, Gregory JS, Distler MG, et al.
Genome wide association study of behavioral, physiological and gene expression traits in a
multigenerational mouse intercross. 2017 [cited 17 Jul 2018]. doi:10.1101/230920

34. Joo JWJ, Hormozdiari F, Han B, Eskin E. Multiple testing correction in linear mixed models.
Genome Biol. 2016;17. doi:10.1186/s13059-016-0903-6

35. Bulik-Sullivan BK, Loh P-R, Finucane HK, Ripke S, Yang J, Schizophrenia Working Group
of the Psychiatric Genomics Consortium, et al. LD Score regression distinguishes
confounding from polygenicity in genome-wide association studies. Nat Genet. 2015;47:
291–295. doi:10.1038/ng.3211

36. Mägi R, Horikoshi M, Sofer T, Mahajan A, Kitajima H, Franceschini N, et al. Trans-ethnic
meta-regression of genome-wide association studies accounting for ancestry increases power

for discovery and improves fine-mapping resolution. Hum Mol Genet. 2017;26: 3639–3650.
doi:10.1093/hmg/ddx280

37. Drescher U, Kremoser C, Handwerker C, Löschinger J, Noda M, Bonhoeffer F. In vitro
guidance of retinal ganglion cell axons by RAGS, a 25 kDa tectal protein related to ligands
for Eph receptor tyrosine kinases. Cell. 1995;82: 359–370. doi:10.1016/0092-
8674(95)90425-5

38. Gao WQ, Shinsky N, Armanini MP, Moran P, Zheng JL, Mendoza-Ramirez JL, et al.
Regulation of hippocampal synaptic plasticity by the tyrosine kinase receptor, REK7/EphA5,
and its ligand, AL-1/Ephrin-A5. Mol Cell Neurosci. 1998;11: 247–259.
doi:10.1006/mcne.1998.0696

39. Gerlai R, Shinsky N, Shih A, Williams P, Winer J, Armanini M, et al. Regulation of learning
by EphA receptors: a protein targeting study. J Neurosci Off J Soc Neurosci. 1999;19: 9538–
9549.

40. Gerlai R, McNamara A. Anesthesia induced retrograde amnesia is ameliorated by ephrinA5-
IgG in mice: EphA receptor tyrosine kinases are involved in mammalian memory. Behav
Brain Res. 2000;108: 133–143. doi:10.1016/s0166-4328(99)00139-4

41. Sheleg M, Yochum CL, Richardson JR, Wagner GC, Zhou R. Ephrin-A5 regulates inter-
male aggression in mice. Behav Brain Res. 2015;286: 300–307.
doi:10.1016/j.bbr.2015.03.001

42. Sheleg M, Yu Q, Go C, Wagner GC, Kusnecov AW, Zhou R. Decreased maternal behavior
and anxiety in ephrin-A5-/- mice. Genes Brain Behav. 2017;16: 271–284.
doi:10.1111/gbb.12319

43. E.M. S, M. J, J. P. Parkinson's Disease and Parkin: Insights from Park2 Knockout Mice. In:
Dushanova J, editor. Mechanisms in Parkinson's Disease - Models and Treatments. InTech;
2012. doi:10.5772/18148

44. Gemechu JM, Sharma A, Yu D, Xie Y, Merkel OM, Moszczynska A. Characterization of
Dopaminergic System in the Striatum of Young Adult Park2-/- Knockout Rats. Sci Rep.
2018;8: 1517. doi:10.1038/s41598-017-18526-0

45. Flagel SB, Clark JJ, Robinson TE, Mayo L, Czuj A, Willuhn I, et al. A selective role for
dopamine in stimulus-reward learning. Nature. 2011;469: 53–57. doi:10.1038/nature09588

46. Saunders BT, Robinson TE. The role of dopamine in the accumbens core in the expression
of Pavlovian-conditioned responses. Eur J Neurosci. 2012;36: 2521–2532.
doi:10.1111/j.1460-9568.2012.08217.x

47. Foster JR, Frost D. The History of the Rat. Boorman's Pathology of the Rat. Elsevier; 2018.
pp. 7–12. doi:10.1016/B978-0-12-391448-4.00003-4

48. Scott PA, Cierpial MA, Kilts CD, Weiss JM. Susceptibility and resistance of rats to stress-
induced decreases in swim-test activity: a selective breeding study. Brain Res. 1996;725:
217–230.

49. Stead JDH, Clinton S, Neal C, Schneider J, Jama A, Miller S, et al. Selective Breeding for
Divergence in Novelty-seeking Traits: Heritability and Enrichment in Spontaneous Anxiety-
related Behaviors. Behav Genet. 2006;36: 697–712. doi:10.1007/s10519-006-9058-7

50. Bertholomey ML, West CHK, Jensen ML, Li T-K, Stewart RB, Weiss JM, et al. Genetic
propensities to increase ethanol intake in response to stress: studies with selectively bred
swim test susceptible (SUS), alcohol-preferring (P), and non-preferring (NP) lines of rats.
Psychopharmacology (Berl). 2011;218: 157–167. doi:10.1007/s00213-011-2381-6

51. Geraldes A, Basset P, Smith KL, Nachman MW. Higher differentiation among subspecies of
the house mouse (Mus musculus) in genomic regions with low recombination:
RECOMBINATION AND SPECIATION IN MICE. Mol Ecol. 2011;20: 4722–4736.
doi:10.1111/j.1365-294X.2011.05285.x

52. Clark FM, Yeomans DC, Proudfit HK. The noradrenergic innervation of the spinal cord:
differences between two substrains of Sprague-Dawley rats determined using retrograde
tracers combined with immunocytochemistry. Neurosci Lett. 1991;125: 155–158.

53. Fuller DD, Baker TL, Behan M, Mitchell GS. Expression of hypoglossal long-term
facilitation differs between substrains of Sprague-Dawley rat. Physiol Genomics. 2001;4:
175–181. doi:10.1152/physiolgenomics.2001.4.3.175

54. Langer M, Brandt C, Löscher W. Marked strain and substrain differences in induction of
status epilepticus and subsequent development of neurodegeneration, epilepsy, and
behavioral alterations in rats. [corrected]. Epilepsy Res. 2011;96: 207–224.
doi:10.1016/j.eplepsyres.2011.06.005

55. Segerström L, Roman E. Response: Commentary: Supplier-dependent differences in
intermittent voluntary alcohol intake and response to naltrexone in Wistar rats. Front
Neurosci. 2016;10. doi:10.3389/fnins.2016.00442

56. Berridge KC, Robinson TE. Liking, wanting, and the incentive-sensitization theory of
addiction. Am Psychol. 2016;71: 670–679. doi:10.1037/amp0000059

57. Sanchez-Roige S, Palmer AA, Fontanillas P, Elson S, Adams MJ, Howard DM, et al.
Genome-wide association study meta-analysis of the Alcohol Use Disorder Identification
Test (AUDIT) in two population-based cohorts (N=141,958). 2018 [cited 23 Jul 2018].
doi:10.1101/275917

58. Sittig LJ, Carbonetto P, Engel KA, Krauss KS, Barrios-Camacho CM, Palmer AA. Genetic
Background Limits Generalizability of Genotype-Phenotype Relationships. Neuron.
2016;91: 1253–1259. doi:10.1016/j.neuron.2016.08.013

59. Pitchers KK, Flagel SB, O'Donnell EG, Solberg Woods LC, Sarter M, Robinson TE.
Individual variation in the propensity to attribute incentive salience to a food cue: Influence
of sex. Behav Brain Res. 2015;278: 462–469. doi:10.1016/j.bbr.2014.10.036

60. Pitchers KK, Wood TR, Skrzynski CJ, Robinson TE, Sarter M. The ability for cocaine and
cocaine-associated cues to compete for attention. Behav Brain Res. 2017;320: 302–315.
doi:10.1016/j.bbr.2016.11.024

61. Kawa AB, Bentzley BS, Robinson TE. Less is more: prolonged intermittent access cocaine
self-administration produces incentive-sensitization and addiction-like behavior.
Psychopharmacology (Berl). 2016;233: 3587–3602. doi:10.1007/s00213-016-4393-8

62. Ahrens AM, Meyer PJ, Ferguson LM, Robinson TE, Aldridge JW. Neural Activity in the
Ventral Pallidum Encodes Variation in the Incentive Value of a Reward Cue. J Neurosci.
2016;36: 7957–7970. doi:10.1523/JNEUROSCI.0736-16.2016

63. Ahrens AM, Singer BF, Fitzpatrick CJ, Morrow JD, Robinson TE. Rats that sign-track are
resistant to Pavlovian but not instrumental extinction. Behav Brain Res. 2016;296: 418–430.
doi:10.1016/j.bbr.2015.07.055

64. Singer BF, Guptaroy B, Austin CJ, Wohl I, Lovic V, Seiler JL, et al. Individual variation in
incentive salience attribution and accumbens dopamine transporter expression and function.
Dalley J, editor. Eur J Neurosci. 2016;43: 662–670. doi:10.1111/ejn.13134

65. Yager LM, Pitchers KK, Flagel SB, Robinson TE. Individual Variation in the Motivational
and Neurobiological Effects of an Opioid Cue. Neuropsychopharmacology. 2015;40: 1269–
1277. doi:10.1038/npp.2014.314

66. Meyer PJ, Cogan ES, Robinson TE. The Form of a Conditioned Stimulus Can Influence the
Degree to Which It Acquires Incentive Motivational Properties. Le Foll B, editor. PLoS
ONE. 2014;9: e98163. doi:10.1371/journal.pone.0098163

67. Yager LM, Robinson TE. A classically conditioned cocaine cue acquires greater control over
motivated behavior in rats prone to attribute incentive salience to a food cue.
Psychopharmacology (Berl). 2013;226: 217–228. doi:10.1007/s00213-012-2890-y

68. Saunders BT, Yager LM, Robinson TE. Cue-Evoked Cocaine "Craving": Role of Dopamine
in the Accumbens Core. J Neurosci. 2013;33: 13989–14000.
doi:10.1523/JNEUROSCI.0450-13.2013

69. Paolone G, Angelakos CC, Meyer PJ, Robinson TE, Sarter M. Cholinergic Control over
Attention in Rats Prone to Attribute Incentive Salience to Reward Cues. J Neurosci. 2013;33:
8321–8335. doi:10.1523/JNEUROSCI.0709-13.2013

70. Morrow JD, Saunders BT, Maren S, Robinson TE. Sign-tracking to an appetitive cue
predicts incubation of conditioned fear in rats. Behav Brain Res. 2015;276: 59–66.
doi:10.1016/j.bbr.2014.04.002

71. Singer BF, Bryan MA, Popov P, Scarff R, Carter C, Wright E, et al. The sensory features of
a food cue influence its ability to act as an incentive stimulus and evoke dopamine release in
the nucleus accumbens core. Learn Mem. 2016;23: 595–606. doi:10.1101/lm.043026.116

72. Fitzpatrick CJ, Morrow JD. Pavlovian Conditioned Approach Training in Rats. J Vis Exp
JoVE. 2016; e53580. doi:10.3791/53580

73. Robinson TE, Flagel SB. Dissociating the Predictive and Incentive Motivational Properties
of Reward-Related Cues Through the Study of Individual Differences. Biol Psychiatry.
2009;65: 869–873. doi:10.1016/j.biopsych.2008.09.006

74. Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, et al. A Robust,
Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. Orban L,
editor. PLoS ONE. 2011;6: e19379. doi:10.1371/journal.pone.0019379

75. Gileta AF, Gao J, Chitre AS, Bimschleger HV, St Pierre CL, Gopalakrishnan S, et al. G3
Adapting Genotyping-by-Sequencing and Variant Calling for Heterogeneous Stock Rats. G3
Bethesda Md. 2020;10: 2195–2205. doi:10.1534/g3.120.401325

76. Grabowski PP, Morris GP, Casler MD, Borevitz JO. Population genomic variation reveals
roles of history, adaptation and ploidy in switchgrass. Mol Ecol. 2014;23: 4059–4073.
doi:10.1111/mec.12845

77. Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. Double Digest RADseq: An
Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-Model
Species. Orlando L, editor. PLoS ONE. 2012;7: e37135. doi:10.1371/journal.pone.0037135

78. Hannon Lab. FASTX-Toolkit. 2010. Available:
http://hannonlab.cshl.edu/fastx_toolkit/index.html

79. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. 2011;17: 10. doi:10.14806/ej.17.1.200

80. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009;25: 1754–1760. doi:10.1093/bioinformatics/btp324

81. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 2010;20: 1297–1303. doi:10.1101/gr.107524.110

82. Hermsen R, de Ligt J, Spee W, Blokzijl F, Schäfer S, Adami E, et al. Genomic landscape of
rat strain and substrain variation. BMC Genomics. 2015;16. doi:10.1186/s12864-015-1594-1

83. Broad Institute. Picard Tools. 2018. Available: http://broadinstitute.github.io/picard/

84. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. Bioinformatics. 2011;27:
2987–2993. doi:10.1093/bioinformatics/btr509

85. Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation
Sequencing Data. BMC Bioinformatics. 2014;15. doi:10.1186/s12859-014-0356-4

86. Browning BL, Browning SR. A Unified Approach to Genotype Imputation and Haplotype-
Phase Inference for Large Data Sets of Trios and Unrelated Individuals. Am J Hum Genet.
2009;84: 210–223. doi:10.1016/j.ajhg.2009.01.005

87. Browning BL, Browning SR. Genotype Imputation with Millions of Reference Samples. Am
J Hum Genet. 2016;98: 116–126. doi:10.1016/j.ajhg.2015.11.020

88. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework
for variation discovery and genotyping using next-generation DNA sequencing data. Nat
Genet. 2011;43: 491–498. doi:10.1038/ng.806

89. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria:
R Foundation for Statistical Computing; 2017. Available: http://www.R-project.org/

90. Wigginton JE, Cutler DJ, Abecasis GR. A Note on Exact Tests of Hardy-Weinberg
Equilibrium. Am J Hum Genet. 2005;76: 887–893. doi:10.1086/429864

91. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal
components analysis corrects for stratification in genome-wide association studies. Nat
Genet. 2006;38: 904–909. doi:10.1038/ng1847

92. Patterson N, Price AL, Reich D. Population Structure and Eigenanalysis. PLoS Genet.
2006;2: e190. doi:10.1371/journal.pgen.0020190

93. Thomas Lumley based on Fortran code by Alan Miller. leaps: Regression Subset Selection.
R   package version 3.0. 2017. Available: https://CRAN.R-project.org/package=leaps

94. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait
analysis. Am J Hum Genet. 2011;88: 76–82. doi:10.1016/j.ajhg.2010.11.011

95. Zhou X, Stephens M. Genome-wide efficient mixed-model analysis for association studies.
Nat Genet. 2012;44: 821–824. doi:10.1038/ng.2310

96. Cheng R, Palmer AA. A Simulation Study of Permutation, Bootstrap, and Gene Dropping
for Assessing Statistical Significance in the Case of Unequal Relatedness. Genetics.
2013;193: 1015–1018. doi:10.1534/genetics.112.146332

97. Listgarten J, Lippert C, Kadie CM, Davidson RI, Eskin E, Heckerman D. Improved linear
mixed models for genome-wide association studies. Nat Methods. 2012;9: 525–526.
doi:10.1038/nmeth.2037

98. Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. Advantages and pitfalls in the
application of mixed-model association methods. Nat Genet. 2014;46: 100–106.
doi:10.1038/ng.2876

99. Loh P-R, Tucker G, Bulik-Sullivan BK, Vilhjálmsson BJ, Finucane HK, Salem RM, et al.
Efficient Bayesian mixed-model analysis increases association power in large cohorts. Nat
Genet. 2015;47: 284–290. doi:10.1038/ng.3190

100.	Risch N, Merikangas K. The future of genetic studies of complex human diseases.
Science. 1996;273: 1516–1517.

101.	Churchill GA, Doerge RW. Empirical threshold values for quantitative trait mapping.
Genetics. 1994;138: 963–971.

102.	Cheng R, Lim JE, Samocha KE, Sokoloff G, Abney M, Skol AD, et al. Genome-Wide
Association Studies and the Problem of Relatedness Among Advanced Intercross Lines and
Other Highly Recombinant Populations. Genetics. 2010;185: 1033–1044.
doi:10.1534/genetics.110.116863

103.	Han B, Kang HM, Eskin E. Rapid and Accurate Multiple Testing Correction and Power
Estimation for Millions of Correlated Markers. Storey JD, editor. PLoS Genet. 2009;5:
e1000456. doi:10.1371/journal.pgen.1000456

104.	Han B, Eskin E. Interpreting Meta-Analyses of Genome-Wide Association Studies. Kerr
K, editor. PLoS Genet. 2012;8: e1002555. doi:10.1371/journal.pgen.1002555

105.	Littrell J, Tsaih S-W, Baud A, Rastas P, Solberg-Woods L, Flister MJ. A High-
Resolution Genetic Map for the Laboratory Rat. G3amp58 GenesGenomesGenetics. 2018;
g3.200187.2018. doi:10.1534/g3.118.200187

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

# Supporting information

**S1 Fig. Correlation heatmap of PavCA metrics across days 1-5.** The heatmap displays the

absolute value of the Pearson correlation coefficient between each pair of PavCA metrics across

all 5 days of testing.

1100    **S2 Fig. Distributions of the average of day 4 and day 5 measurements for 10 PavCA**

1101    **metrics.** Histograms were constructed using measurements from the combined Harlan and

1102    Charles River sample set. Not shown is the PacCA index score.

1103

1104    **S1 Table. Sample origins for all 4,061 SD rats in final filtered set.**

| | # Samples (Barrier Facility) | # Samples (Breeding Location) |
|---|---|---|
| **Total samples post-filtering** | 4,061 | 4,061 |
| Harlan – Dublin, VA – Barrier 231 | 5 | 5 |
| Harlan – Frederick, MD – Barrier 208A | 465 | 465 |
| Harlan – Haslett, MI – Barrier 206 | 762 | 762 |
| Harlan – Houston, TX – Barrier 211 | 14 | 14 |
| Harlan – Indianapolis, IN – Barrier 202A | 546 | |
| Harlan – Indianapolis, IN – Barrier 202C | 97 | 994 |
| Harlan – Indianapolis, IN – Barrier 217 | 351 | |
| Charles River – Kingston, NY – Barrier K92 | 4 | 4 |
| Charles River – Portage, MI – Barrier P03 | 32 | |
| Charles River – Portage, MI – Barrier P07 | 121 | |
| Charles River – Portage, MI – Barrier P09 | 327 | 558 |
| Charles River – Portage, MI – Barrier P10 | 78 | |
| Charles River – Raleigh, NC – Barrier R04 | 650 | |
| Charles River – Raleigh, NC – Barrier R09 | 144 | 794 |
| Charles River – Saint Constant, CAN – Barrier C71 | 46 | |
| Charles River – Saint Constant, CAN – Barrier C72 | 358 | 404 |
| Unknown Barrier | 61 | 61 |

1105

1106

1107

1108

1109

1110 **S3 Fig. Distributions of the average of day 4 and day 5 PavCA index scores for 6 major**

1111 **breeding locations.** Sample numbers for each breeding location can be found in S1 Table. The

1112 three locations in the left column are from Harlan, and the three locations in the right column

1113 belong to Charles River.

| Breeding Location | Frederick | Haslett | Indianapolis | Portage | Raleigh | St. Constant |
|---|---|---|---|---|---|---|
| Frederick | 1 | 1.955e-10 | 4.645e-06 | <2.2e-16 | <2.2e-16 | <2.2e-16 |
| Haslett | | 1 | 0.0131 | <2.2e-16 | <2.2e-16 | <2.2e-16 |
| Indianapolis | | | 1 | <2.2e-16 | <2.2e-16 | <2.2e-16 |
| Portage | | | | 1 | 0.004155 | 6.657e-06 |
| Raleigh | | | | | 1 | 0.03225 |
| St. Constant | | | | | | 1 |

Welch's 2-sample t-test p-values from pairwaise comparisons of day4 and day 5 average PavCA index score distributions between different breeding locations.

1114

1115 **S2 Table. List of variant filtering steps and the numbers of SNPs remaining after each step**

1116 **for both ANGSD/Beagle and STITCH.** The values highlighted in green are the final SNP totals

1117 used for GWAS in Charles River and Harlan. The values highlighted in orange are the counts we

1118 used as criteria for choosing ANGSD/Beagle over STITCH.

| Sequential Filtering Steps | ANGSD/Beagle | | STITCH | |
|---|---|---|---|---|
| | Number of SNPs Harlan | Number of SNPs Charles River | Number of SNPs Harlan | Number of SNPs Charles River |
| Total called/imputed | 2,274,118 | 2,274,118 | 8,691,886 | 8,691,886 |
| SNPs with dosage $r^2 \geq 0.9$ | 305,779 | 305,779 | 1,992,919 | 1,992,919 |
| SNPs with MAF $\geq 0.01$ | 235,583 | 128,572 | 1,058,812 | 1,843,240 |
| SNPs with HWE p-value $< 1 \times 10^{-7}$ | 214,309 | 114,568 | 930,685 | 1,718,475 |
| SNPs after LD pruning $r^2 < 0.95$ | 44,721 | 93,692 | 25,009 | 75,503 |
| Union set of SNPs | 234,887 | | | |
| Novel SNPs compared to dbSNP (not found in Build 149 - Nov 7, 2016) | 195,299 | | | |
| Novel SNPs compared to 42 genomes ref (not found in Hermsen et al. 2015) | 105,638 | | | |

1119

1120

1121

1122 **S1 Text. ddGBS protocol used to sequence SD rats.**

1123

1124

1125

1126

1127

1128   **S3 Table. Concordance and error rates for ANGSD/Beagle genotypes at different dosage r2**

1129   **thresholds.** Rates of concordance and genotyping error were calculated by comparing genotypes

1130   for 381 duplicate samples called in parallel. Each of the two replicates of the sample was

1131   assumed to contribute half the discordant genotypes. Therefore, the per sample genotyping error

1132   rate was calculated as half of the observed rate of discordance.

| | $DR^2>=0.9$ | $DR^2>=0.8$ | $DR^2>=0.7$ |
|---|---|---|---|
| Total called and imputed by ANGSD/Beagle | 2,274,118 | 2,274,118 | 2,274,118 |
| SNPs passing dosage $r^2$ filter | 305,779 | 427,123 | 541,622 |
| SNPs passing MAF >= 0.01 filter | 244,586 | 326,550 | 393,569 |
| SNPs passing HWE $10^{-7}$ filter | 204,104 | 259,111 | 305,660 |
| Mean Pearson **correlation of dosages** (n=381) | 0.983 | 0.969 | 0.957 |
| **Rate of concordance** of hard calls (n=381) | 0.983 | 0.972 | 0.963 |
| **Rate of discordance** (1 − concordance) | 0.017 | 0.028 | 0.037 |
| **Error rate** (Rate of discordance/2)*100 | 0.85% | 1.4% | 1.85% |

1133

1134

1135

1136

1137

1138

1139

1140

1141 **S4 Table. List of filtered variant counts and sample numbers in subpopulations.**

| Subpopulations | Number of rats | Number of SNPs |
|---|---|---|
| Harlan – All, union SNP set | 2,221 | 223,467 |
| Harlan – 202A/202C/208A | 1,108 | 197,355 |
| Harlan – 206 | 762 | 173,443 |
| Harlan – 217 | 351 | 171,648 |
| Charles River – All, union SNP set | 1,760 | 273,241 |
| Charles River – R09/P03/P07/P10/C71/K92 | 425 | 256,236 |
| Charles River – R04 | 650 | 252,798 |
| Charles River – P09 | 327 | 253,640 |
| Charles River – C72 | 358 | 250,868 |

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

**S4 Fig. Heatmaps of pairwise identity-by-descent pre- and post-filtering.** Panels A and C

show pre-filtering values of P(IBD) = 0 plotted against P(IBD) = 1 for Harlan and Charles River.

Unrelated samples cluster in the lower right corner. Samples along the diagonal have $2^{nd}$ and $3^{rd}$

degree levels of relatedness, while those clustering around (0.5,0.25) are full-siblings. Panels B

and D demonstrate that the sample filtering steps removed several spurious relations from the

sample.

1158

1159    **S5 Table. List of sample filtering criteria and number of samples removed.**

|  | Number of Samples | |
|---|---|---|
| Total sequenced by ddGBS + WGS | 4,625 (17 WGS only) | |
| Low read count in FASTQ ($<$ 4 million reads) **performed prior to variant calling | 316 | |
| Females samples (only females in sample set) | 77 | |
| Purchased from Taconic | 4 | |
| Lack of phenotype information | 10 | |
| Poor clustering in principal component analysis | 54 | |
| Putative sample mix-up | 18 | |
| Duplicate sample | 7 | |
| Inflated/deflated heterozygosity | 34 | |
| $>=$ 30 samples with $\hat{\pi}$ $>=$ 0.1875 in sample | 12 | |
| $\hat{\pi}$ $>$ 0.6 with another sample | 32 | |
| **Final Sample Set (Harlan + Charles River)** | 4,061 | |
|  | CR – 1,780 | Har – 2,281 |

1160

1161

1162

1163

1164    **S6 Table. Heritability estimates in Charles River, Harlan, and their respective subgroups**

1165    **for 55 mapped PavCA metrics.**

1166

1167

1168

1169

1170 **S5 Fig. QQ-plot inflation observed in vendor-level GWAS**

1171

**S1 File. Table of all significant PavCA GWAS hits.** For each significant GWAS association,

the table contains the position, allele, allele frequency, PavCA metric, sample size, direction of

effect, meta-analysis p-value, percent variance explained, effect size, and standard error. In

addition, the values for all the independent GWAS that went into the meta-analyses are provided.

**S2 File. Stacked subgroup Manhattan plots for all analyzed metrics/traits.** Each page

contains vertically stacked Manhattan plots with the GWAS results for all PavCA metrics in all

seven sample subgroups. The plots are in the following order from top to bottom: Harlan

202A/202C/208A (115k SNPs), Harlan 206 (102k SNPs), Harlan 217 (115k SNPs), Charles

River R09/P03/P07/P10 (221k SNPs), Charles River R04 (214k SNPs), Charles River P09 (223k

SNPs), and Charles River C72 (222k SNPs).

**S3 File. Q-Q plots for all analyzed metrics/traits.** Each page contains Q-Q plots for the GWAS

of a given day/metric for the seven subgroups.

**S4 File. Stacked meta-analyses Manhattan plots for all analyzed metrics/traits.** Each page

contains vertically stacked Manhattan plots with the GWAS results for all PavCA metrics in all

three meta-analysis: Charles River plus Harlan (64k), Charles River only (198k SNPs), and

Harlan only (83k SNPs).

**S5 File. Q-Q plots for all metrics/trait meta-analyses.** Each page contains Q-Q plots for the

meta-analyses of the GWAS results for a given day/metric across (1) all seven subgroups, (2)

Charles River subgroups, and (3) Harlan subgroups.

**S2 Text. Sequences for 48 barcoded adapters, Y-adapter, and PCR primers.**

1192

1193 **S8 Table. List of all PavCA metrics collected on SD rats.** There are 11 total metrics across 5

1194 days of training. The first 7 metrics are direct measurements made during the training periods.

1195 The following 3 metrics are calculated from the base measurements. The final metric, PavCA

1196 index score, is a composite score from the previous 3 metrics.

| Day 1 25 8-second trials | Day 2 25 8-second trials | Day 3 25 8-second trials | Day 4 25 8-second trials | Day 5 25 8-second trials | Data Type |
|---|---|---|---|---|---|
| Lever Presses | Lever Presses | Lever Presses | Lever Presses | Lever Presses | Count (summed over trials) |
| Magazine Entries CS | Magazine Entries CS | Magazine Entries CS | Magazine Entries CS | Magazine Entries CS | Count (summed over trials) |
| Magazine Entries NCS | Magazine Entries NCS | Magazine Entries NCS | Magazine Entries NCS | Magazine Entries NCS | Count (summed over trials) |
| Latency to Lever Press | Latency to Lever Press | Latency to Lever Press | Latency to Lever Press | Latency to Lever Press | Continuous [0,8] secs (averaged over trials) |
| Latency to Magazine Entry | Latency to Magazine Entry | Latency to Magazine Entry | Latency to Magazine Entry | Latency to Magazine Entry | Continuous [0,8] secs (averaged over trials) |
| Probability of Lever Press | Probability of Lever Press | Probability of Lever Press | Probability of Lever Press | Probability of Lever Press | Proportion [0-1] (0/25 to 25/25 trials) |
| Probability of Mag Entry | Probability of Mag Entry | Probability of Mag Entry | Probability of Mag Entry | Probability of Mag Entry | Proportion [0-1] (0/25 to 25/25 trials) |
| Response Bias | Response Bias | Response Bias | Response Bias | Response Bias | Continuous [-1,1] |
| Latency Score | Latency Score | Latency Score | Latency Score | Latency Score | Continuous [-1,1] |
| Probability Difference | Probability Difference | Probability Difference | Probability Difference | Probability Difference | Continuous [-1,1] |
| PavCA Index Score | PavCA Index Score | PavCA Index Score | PavCA Index Score | PavCA Index Score | Continuous [-1,1] |

1197

1198 **S9 Table. List of covariates used for the GWAS LMMs for Harlan and Charles River.** The

1199 first three covariates were used in both Charles River and Harlan analyses. The remaining

1200 covariates were unique to each population.

| Har 202A | Har 206 | Har 217 | CR R04 | CR R09 | CR P09 | CR C72 |
|---|---|---|---|---|---|---|
| Age of rat in days at start of training (continuous integer) | | | | | | |
| Housing condition (binary – single or multiple rats per chamber) | | | | | | |
| Light cycle (binary – standard or reverse 12h lighting) | | | | | | |
| AAK | CJF | AK | AAK | CJF | AK | AAK |
| BFS.AA | JC.JDM | BFS | CJF | CJF.JDM | BFS.AA | BFS |
| CJF | JJ | BTS | CJF.JDM | EGO | EGO | BFS.AA |
| CJF.JDM | KP | CJF | JJ | | LMY | CJF.JDM |
| EGO | | CJF.JDM | | | MLR.LMF | |
| KP | | JC.JDM | | | | |
| LMY | | TW.KP | | | | |

1201

1202

1203

1204 **S6 Fig. Pre- and post-filtering distributions of heterozygosity for Harlan and Charles**

1205 **River.** Panels A and C show pre-filtering distributions of heterozygosity in Harlan and Charles

1206 River, as measured by the method-of-moments F coefficient. Panels B and D show the same

1207 distributions post-filtering. A value above 0 indicates a deflation of heterozygosity, whereas a

1208 value below 0 would be an inflation.

1209

1210 **S6 File. Spreadsheet with detailed sample sequencing and phenotype information.**

 **Figure Captions**

 **Fig 1. PavCA index score progression across days and distribution between Charles River**

 **and Harlan.**

(A) Samples were classified as ST, GT, or IR based on their average day 4 and 5 PavCA index

score. PavCA index scores for each day of training (1-5) were averaged across all STs, GTs, and

IRs. (B) Density curves of average day 4 and 5 PavCA index score in Harlan (n=2,281) vs.

Charles River (n=1,780) SD rats. (C) Filtered SNP density per 1Mb window in Charles River vs.

Harlan samples for all 20 autosomes.

**Fig 2. Genetic architecture of SD rats from Charles River vs. Harlan.**

(A) Map of the nine vendor breeding locations and the number of SD rats obtained from each

location. (B) A summary of the genetic data from all 4,061 SD rats based on principal

components 1 and 2 from PCA. Each point represents a sample. The left cluster is composed of

samples from Charles River and the right clusters are composed of samples from Harlan. (C-D)

Repeated PCA analyses on subsets of the samples from Harlan and Charles River, colored by

barrier facility of origin.

**Fig 3. SNP MAF distributions and comparison of linkage disequilibrium decay rates.**

(A) Density curves of minor allele frequencies for 214,309 SNPs in Charles River and 114,568

SNPs in Harlan, after removing SNPs with MAF < 0.01. (B) Linkage disequilibrium decay rates

in SD rats from both vendors and outbred Swiss Webster (CFW) mice.

**Fig 4. Manhattan plots for the GWAS on PavCA Index Score on day 5 in each subgroup**

**and the meta-analysis.**

The titles above each of the eight Manhattan plots indicate the population, SNP count, and

sample size for each depicted GWAS. The y-axis is the $-\log_{10}$ of the p-value from the likelihood-

ratio test performed by GEMMA. The x-axis is the genomic coordinate of the variants within

each chromosome, in ascending order of chromosome.