1 **Nuclear envelope budding is a response to cellular stress**

Dimitra Panagaki[a1] and Jacob T. Croft[a1], Katharina Keuenhof[a], Lisa Larsson-Berglund[a], Stefanie

Andersson[a], Verena Kohler[c], Sabrina Büttner[c], Markus J. Tamás[a], Thomas Nyström[b], Richard

Neutze[a], Johanna L. Höög[a*]

[a]Department of Chemistry and Molecular Biology, University of Gothenburg, 405 30, Gothenburg, Sweden

[b]Department of Microbiology and Immunology, University of Gothenburg, 405 30, Gothenburg, Sweden

[c]Department of Molecular Bioscienses, The Wenner-Gren Institute, Stockholm University, 106 91,

Stockholm, Sweden

* Johanna L. Höög
Email: johanna.hoog@gu.se

13 **Classification**

14 Biological Sciences; Cell Biology

15
16

---

[1] These authors contributed equally to the manuscript.

Keywords

Nuclear transport; budding; vesicles; electron tomography; protein quality control;

Author Contributions

Conceptualization, D.P., J.T.C., S.B. and J.L.H.; Methodology, D.P., J.T.C., L.L.B., K.K. and

J.L.H.; Validation, D.P., J.T.C. and K.K.; Formal Analysis, D.P., J.T.C., R.N. and K.K.;

Investigation, D.P., J.T.C., K.K., L.L.B., V.K. and S.A.; Resources, S.B., M.J.T., T.N., R.N. and

J.L.H.; Writing-Original Draft, D.P., J.T.C. and J.L.H.; Writing-Review & Editing, D.P., J.T.C., K.K.,

L.L.B., S.A., V.K., S.B., M.J.T., T.N., R.N. and J.L.H.; Visualization, D.P., J.T.C. and J.L.H.;

Supervision, L.L.B., R.N., M.J.T., T.N., S.B. and J.L.H.; Funding Acquisition, V.K., S.B., T.N.,

R.N., M.J.T. and J.L.H.

This PDF file includes:

Main Text

Figures 1 to 7

Abstract

Nuclear envelope budding (NEB) is a recently discovered alternative pathway for

nucleocytoplasmic communication distinct from the movement of material through the nuclear

pore complex. Through quantitative electron microscopy and tomography, we demonstrate how

NEB is evolutionarily conserved from early protists to human cells. In the yeast *Saccharomyces*

cerevisiae, NEB events occur with higher frequency during heat shock, upon exposure to

arsenite or hydrogen peroxide, and when the proteasome is inhibited. Yeast cells treated with

azetidine-2-carboxylic acid, a proline analogue that induces protein misfolding, display the most

dramatic increase in NEB, suggesting a causal link to protein quality control. This link was further

supported by both localization of ubiquitin and Hsp104 to protein aggregates and NEB events,

and the evolution of these structures during heat shock. We hypothesize that NEB is part of

normal cellular physiology in a vast range of species and that in *S. cerevisiae* NEB comprises a

stress response aiding the transport of protein aggregates across the nuclear envelope.

Significance Statement

A defining feature of eukaryotes is the nuclear envelope, a double lipid bilayer that serves to

isolate and protect the cells genetic material. Transport of large molecules over this barrier is

believed to occur almost exclusively *via* the nuclear pores. However, herpes virions and mega

ribonucleoproteins (megaRNPs) use an alternative means of transport – *via* nuclear envelope

budding (NEB). Here, we show NEB is a ubiquitous eukaryotic phenomenon and increases when

exposed to various forms of cellular stress. NEB frequency was maximal when the cell was

challenged with a drug that induces protein misfolding, indicating this transport pathway plays a

role in protein quality control. These results imply that NEB is an underappreciated yet

potentiallyfundamental means of nuclear transport.

Main Text

Introduction

The nucleus is the most prominent organelle in eukaryotic cells, enclosing most of the cellular

genetic material within a double lipid bilayer called the nuclear envelope. These two membranes

arose as a critical evolutionary step distinguishing eukaryotes from prokaryotes and restricting

which molecules come into contact with the cellular DNA. As the nuclear envelope is not

permeable to most of the molecules inside the cell, special structures called nuclear pore

complexes (NPCs) exist on its surface that allow highly selective translocation over the nuclear

membrane (1-3). Molecules smaller than 30-40 kDa can passively penetrate the NPC from the

cytoplasm into the nucleus and *vice versa*, whereas bigger molecules require interaction with

nuclear transport receptors and form importin/exportin complexes that are guided through the

pores (4-8).

Since its discovery in 1954, the NPC has generally been accepted as the only means of

communication between the cytoplasm and the nucleoplasm (9). However, herpes simplex virus

replicates in the nucleoplasm and is released into the cytosol via an outwards budding of the

nuclear envelope (10-13). This demonstrates another pathway for nuclear export and there have

been several observations suggesting that nuclear envelope budding (NEB) also occurs in

healthy cells, with different interpretations of this mechanism being suggested (14-27). These

observations were made in a diverse set of organisms but primarily using cells that are under

differentiation (embryonic cells) and various terminologies were given to describe the NEB

process (Supplementary Table 1). The most recent reports were made in healthy developing

Drosophila melanogaster larvae, in the sea urchin gastrula and in the budding yeast

Saccharomyces cerevisiae (28-32). Previous studies have shown that budding events in *D.*

melanogaster contain large ribonucleoprotein granules but whether this function is consistent in

other cell types remains unknown (28, 31).

Evidence that material can be exported from the nucleus through NEB has sparked speculation

that other large cargoes, such as protein aggregates, could be removed from the nucleus by a

similar mechanism (33). Protein misfolding can be highly toxic to the cell and therefore multiple

stress-induced mechanisms have evolved to cope with proteotoxicity (34-36). Cells experiencing

large-scale protein misfolding exhibit DNA mutagenesis, which is one of the first steps towards

carcinogenesis (37), demonstrating the importance of protein quality control in the nucleus (38).

Despite the cell's primary protein quality control factors, such as chaperones and the ubiquitin-

proteasome system, being active in the nucleus, protein aggregates still form under stressful

conditions such as heat shock, because these processes cannot fully cope with the quantity of

misfolded proteins (39). In most occasions, protein aggregates will enter the nucleus for

degradation but cases of nuclear misfolded proteins being transported out of the nucleus to be

degraded in the cytoplasm have also been reported (39). Nuclear protein aggregates greatly

exceed the 39 nm size limit of active transportation through the NPCs (40, 41), implying that an

alternative pathway for their export is required if they are to be transported over the nuclear

envelope.

As well as transporting material from the nucleus to the cytoplasm, NEB can also result in the

transfer of a portion of the inner nuclear membrane (INM) to the outer nuclear membrane (ONM),

which is continuous with the endoplasmic reticulum (ER). In addition to protein quality control

mechanisms in the nucleus and cytoplasm, the endoplasmic reticulum (ER) is equipped with its

own set of degradation pathways, to contend with the large quantity of newly synthesized proteins

entering the ER (42). Protein degradation in the ER is highly dependent on the ubiquitin

proteasome system and facilitates removal of both misfolded soluble and membrane proteins

(43). Similarly, degradation of INM proteins can occur via biochemically similar pathways to ER-

associated degradation of membrane proteins (44). Several branches of INM-associated

degradation (INMAD) exist, each relying on a different E3 ubiquitin ligase to recognize and tag

misfolded substrates through ubiquitination steps(45-47). These reflections pose the question:

does NEB provide an escape valve needed for clearance of aggregated nuclear proteins? To

address this issue, we have examined the NEB pathway of *S. cerevisiae* under five different

stress conditions, all of which caused an increase in NEB events. We also show the presence of

NEB in five different evolutionarily distant organisms (*Homo sapiens, Caenorhabditis elegans,*

Saccharomyces cerevisiae, Schizosaccharomyces pombe, Trypanosoma brucei) under normal

growth conditions, which reveals the evolutionary conservation of the NEB pathway. Our findings

provide evidence that NEB is part of the cell's natural stress response and an important pathway

to consider when studying nuclear transport, especially with regard to the protein quality control

system.

Results

NEB increases in frequency during heat shock in *S. cerevisiae*

S. cerevisiae provides a well-understood model system, perfectly tailored to investigate if NEB is

an active part of the cellular stress response. An easily applied stress impulse is heat-shock and

we therefore subjected *S. cerevisiae* cells to a mild constant heat stress (38°C) for a time course of

up to 90 minutes (Figure 1**a**). Cells were then cryo-immobilized for electron microscopy studies,

which are able to simultaneously visualize both NEB events and protein aggregates, the latter

appearing as electron dense content (EDC) (48) within the nucleus (Figure 1**b**). For each time point

between 60 to 80 electron micrographs were acquired, each from a randomly chosen cell section

that passed through the cell nucleus. NEB events were identified as electron dense material

localized between the two nuclear envelopes, causing it to deform, with a small number of events

showing budding of the nuclear membrane but seeming to lack an electron dense cargo (S1). A

membrane bilayer was often, but not always, detectable around this material. NEB events

commonly had an internal texture resembling the nucleoplasm (Figure 1**b**),but in some rare events

the bud appeared to contain cytoplasmic material including ribosomes (Figure S1). In one case, a

vacuole was seen engulfing a protrusion of the nucleus . This phenomenon is termed piecemeal

microautophagy of the nucleus and has previously been described in nutrient deprived cells (49).

Piecemeal microautophagy has a distinct morphology from the NEB events and were not included

in our quantification.

Morphological variations of NEB (Figure S1) could represent different stages of the same process

or different processes. Images were scored both for the presence of NEB events and EDC in the

nucleus. Both NEB and EDC increased during heat shock, prior to decreasing near the end of the

time course, which potentially reflects a cellular adaptation to the new temperature (Figure 1**a**).

EDC reaches a maximum (83.3% of sections) after 15 minutes and stays relatively stable at this

level until 90 minutes, at which point it is only present in 60.5% of sections (Figure 1**a**). NEB events

were found in 2.4% of cellular sections in undisturbed cultures but this fraction significantly

increased to 10.3% after 30 minutes of heat shock. The effect of a stronger heat shock treatment

(42°C for 30 minutes) was examined to detect a possible correlation of the NEB frequency to the

level of stress. NEB frequency was also here significantly increased in comparison to the control

as 16.6% of the nuclear sections displayed NEB (Figure S2).

To verify that the observed EDC in the nucleus are protein aggregates, immuno-electron

microscopy against the disaggregase Hsp104 (50) and the model misfolding protein temperature

sensitive mutant guk1-7 (51) were performed. In these experiments, the secondary antibody is

coupled to a gold particle for visualization using electron microscopy. Both Hsp104 and guk1-7

localised to the EDC material upon heat shock (Figure 1**c**, **e-f,** n = number of examined structures).

As a control, an anti-Hsp104 antibody was used to label the EDC in heat shocked wild type and

hsp104Δ cells to validate the specificity of the labeling. High labeling density of the EDC could be

detected in the wild type strain, whereas in the *hsp104Δ* strain almost no unspecific gold particles

were found (Figure **d,g**).

The concurrent increase in frequency of NEB and protein aggregation strengthens the hypothesis

that NEB may indeed have an important function during normal cellular stress response. This

apparent correlation between the frequency of NEB with heat shock motivated us to test if this

pathway is solely affected by heat-shock or if cells react in a similar fashion under different stress

conditions.

Hydrogen peroxide and sodium arsenite stressors also increase NEB frequency

Cells may encounter a variety of stress conditions and each of them has a different impact on the

cell (52). To investigate whether additional cellular stresses also correlated with an increased

frequency of NEB events we challenged *S. cerevisiae* cells with multiple stress stimuli. A natural

stimulus is aging which causes a significant induction of stress response pathways in *S. cerevisiae*

(35, 53), and has previously been seen to trigger an increase in "nuclear herniations" (54). Several

factors are responsible for a measurable decline in cellular and physical functions of aged cells,

with one of them being aggregated misfolded proteins (55). We isolated replicatively aged

biotinylated *S. cerevisiae* cells, using Streptavidin magnetic beads, from the young unbound

population (56) and cryo-immobilized the cells. In old yeast cells, the frequency of NEB events was

found to be 9% (Figure 2**a**, 2**d**; n=200 sections) which is significantly higher than an undisturbed *S.*

cerevisiae culture (2.4%; n=337) (Figure 1**a**). However, the remaining younger population also

displayed a significant increase in NEB events to 6.9% (n=204 sections; Figure 2**d**), indicating that

these cells may also have been stressed by the mechanical handling steps associated with isolating

the old cells. Thus, we could not confirm the previous finding of significantly increased NEB events
in old cells in comparison to young cells (54), in this experimental setup.
187
Cells of *S. cerevisiae* were also subjected to 90 minutes 0.6 mM H_2O_2 (hydrogen peroxide,
oxidative stress) and 60 minutes 0.5 mM $NaAsO_2$ (sodium arsenite, heavy metal stress). Treated
cells, as well as a control cultures for each condition, were cryo-immobilized and prepared for
analysis by electron microscopy. Cells that were treated with sodium arsenite had a NEB frequency
of 6% (n=249 sections) and cells treated with hydrogen peroxide had a frequency of 5.9% (n=204
sections) (Figure 2**b-d**). Both control samples had lower levels of NEB events, as found in
undisturbed *S. cerevisiae* (2.4% for the sodium arsenite control for n= 337 and 1.5% for the H_2O_2
control for n= 197, Figure 2d) . From this set of observations, we conclude that NEB is increased in
hydrogen peroxide and sodium arsenite stress as well. .
197

NEB frequency is highest under induction of protein misfolding
All of the stressors examined above have previously been shown to increase rates of aggregation
of misfolded proteins in the cytoplasm and within the nucleus (35, 57-59). To investigate whether
or not the observed increase in NEB events is indeed correlated with proteotoxic stress, we
examined *S. cerevisiae* cells after treatment with azetidine-2-carboxylic acid (AZC). AZC competes
with the amino acid proline during translation and may be mistakenly incorporated into proteins.
Since AZC has one fewer carbon atoms in its ring (4 carbon atoms instead of 5 as in proline), its
incorporation into a polypeptide causes a different conformation of the amino acid backbone and
this results in aggregation of proteins with non-native structures (60, 61). Cells were grown at
normal temperature (30°C) and were treated with AZC, cryo-immobilized after 30 and 90 minutes
and prepared for analysis by electron microscopy. Cellular viability upon treatment with all of the
stressors mentioned above was not compromised, as validated with propidium iodide (PI) staining,
indicative of plasma membrane rupture and thus cell death (Figure S3).
The NEB frequency was similar in untreated cells and cells after 30 min of AZC exposure (2.4% n=
337 sections and 2.6% n= 114 sections respectively, Figure 2**f**). However, after 90 minutes of
exposure to AZC, the number of NEB events dramatically increased in frequency, reaching 22%
(Figure 2**e-f**; n= 100). This time-dependent increase in NEB after AZC exposure demonstrates a
direct link between the NEB pathway and how the cell accommodates proteotoxic stress.
216

NEB events are ubiquitinated
Most proteins that are destined for degradation through the 26S proteasome system must first bind
to a poly-ubiquitin chain (62). Moreover, ubiquitin signaling is known to target cytosolic proteins and
organelles for degradation by autophagy (63, 64). To probe if the NEB cargo is targeted for protein

degradation via ubiquitin signaling, we performed immuno-EM analysis on *S. cerevisiae* cells with

a polyclonal antibody that has a stronger affinity to poly-ubiquitin chains than to monomeric ubiquitin

(Abcam ab19247). In order to stimulate an increase of the NEB events, cells were subjected to 30

minutes of heat shock. Images of 60 randomly chosen cell sections were recorded and the number

of gold particles per area of various cell structures were quantified (Figure 3**a**). Lipid droplets were

used as a negative control (4 gold particles/μm^2) (Figure 3**a-b**) and autophagosomes were chosen

as a positive control (67 gold particles/μm^2). The nucleoplasm (excluding areas containing EDC)

did not differ from the negative control (4 gold/μm^2) whereas EDC (protein aggregates) were

labeled fivefold more frequently (21 gold/μm^2) (Figure 3**a-b**). Similarly, NEB events (17 gold/μm^2)

were labeled fourfold more frequently than lipid droplets (4 gold/μm^2). In order to achieve a higher

n number for the NEB events, cells were actively traced and selected based on the presence of

NEB events. Most of the gold particles identified as associated with NEB events had a preference

of localizing in close proximity to the membrane of the bud, but in other examples it would localize

more centrally. (Figure 3**b, S3**). These results confirm the presence of ubiquitin in both EDC and

the NEB events recorded during heat shock in *S. cerevisiae*.

Proteasome inhibition increases NEB event frequency

If our hypothesis were correct that NEB could function to remove misfolded proteins from the

nucleus that cannot be accommodated by the proteasome, then the inhibition of the proteasome

should alter the NEB event frequency. To investigate this, the proteasome was inhibited using two

different methods. First, we treated *pdr5Δ* cells (65), deficient in a drug efflux pump with the MG132

drug which is known to partially inhibit proteasomal activity (66). This treatment led to an observed

NEB frequency of 7% (n=200), a significant increase compared to the control group (2.5%, n=200)

(Figure 3**c**). In order to achieve a stronger proteasomal inhibition, the Rpn4 transcriptional activator

of genes encoding proteasomal subunits (43, 67) was deleted and the derived strain was prepared

for analysis. Inhibition of the proteasome was confirmed by performing western blots against

ubiquitin, since decreased proteasome activity will result in an increase of ubiquitinated proteins

(Figure 3**d**). Most of the observed NEB events (11 out of 15) in the *rpn4Δ* cells had a distinct

electron dense appearance and no obvious membrane surrounded this electron dense material

(Figure 3**e**). In comparison to wild type cells which displayed a NEB frequency of 2.4% (n=337),

rpn4Δ cells had NEB events in 15% of the sections (n=100) (Figure 3**c**). These results suggest that

NEB pathway is an alternative to the ubiquitin proteasome system.

NEB structures contain Hsp104-GFP

To probe for the presence of aggregated proteins within NEB, we performed immuno-EM in heat

shocked cells (38°C for 30 minutes) with an anti-GFP antibody in cells expressing Hsp104-GFP.

Even though Hsp104 is not an aggregate per se, it is a well-characterized chaperone which

localizes to aggregated proteins in the nucleus and the cytoplasm, where it acts as a disaggregase,

(68-70) and supports the degradation of ubiquitinated membrane proteins in the ER-associated

degradation pathway(71). In comparison to the negative control (lipid droplets, 1.7 gold/ μm^2), NEB

events were highly labeled by gold particles (122 gold/μm^2) comparable to the positive control

(EDC, 144 gold/μm^2) (Figure 3**f-g**). This finding is consistent with either a function of NEB in

transporting misfolded nuclear proteins to the cytoplasm, or INM proteins to the ER for degradation.

NEB structures are not misassembled NPCs

It has been suggested the NEB events are misassembled NPCs (54, 72). When certain key genes

required for NPC assembly are deleted, herniations of the nuclear envelope similar to NEB events

are observed (73, 74). These types of herniations have a distinct morphology with a 'neck

resembling' formation between the vesicle and the INM which was revealed to be a defective NPC

by subtomogram averaging (75). These NPC-exposing herniae are then degraded by autophagy

(75, 76). In addition, NPCs assemble via an inside-out mechanism during which the INM evaginates

slightly, although functional NPC assembly does not resemble NEB (77).

To clarify if our observed NEB could also be misassembled NPCs, we performed immuno-EM using

an antibody that recognizes four NPC components (Nup62, Nup153, Nup214 and Nup358) on both

undisturbed *S. cerevisiae* cultures as well as the aged cells. In the undisturbed culture , the majority

(67%) of NPCs observed were labeled by gold particles (positive control; n = 95) (Figure 4**a**, 4**e**)

and only 16.7% of lipid droplets (negative control; n=82). Only in one case was a NEB event labeled

(16.6%, n=6) (Figure 4**c**) whereas all other NEB events were unlabeled (Figure 4**b-c**, 4**e**). In the

old yeast cells, 81% of the observed NPCs were labeled by gold particles (positive control; n=100)

(Figure 4**d-e**) whereas only 8% of lipid droplets were labeled (negative control; n=100) (Figure 4**e**).

In contrast, only one atypical NEB event was labelled by a gold particle (10%; n= 10) (Figure 4**e**).

To further support this distinction between NEB and herniation, a comparison of the detailed

morphology of NEB events with the previously reported herniations was accomplished through

dual-axis electron tomography of the nuclear envelope in *S. cerevisiae*. This analysis showed that

the selected NEB event was completely enveloped in a lipid bilayer and wedged in between the
two nuclear membranes (Figure 4**f**-**g**). Moreover, the 3D reconstruction revealed that the vesicle
contained between the lipid membranes was complete and was not attached to either nuclear
membrane by a 'neck' with only the outer nuclear membrane extending towards the cytoplasm. As
such this morphology was strikingly different to that previously identified as associated with the
misassembly of NPCs (77). It is also noteworthy that the 3D reconstruction did not show an obvious
connection of the NEB events with the ER, separating these two compartments. A different
morphology of a nuclear envelope-ER connection in comparison with the NEB events can also be
seen in the acquired thin sections of nuclei (Figure S5). As such, both the immuno-EM results and
the 3D morphology of NEB events suggest that the partial or incomplete assembly of NPC and
NEB are distinct structures. However, the possibility that a small fraction of the protrusions
quantified as NEB events in this study may be better assigned NPC related structures remains.
300
The NEB pathway is part of normal cellular function from humans to protists
To gain insights into the evolutionary conservation of NEB we imaged and quantified NEB events
in thin sections of nuclei in human mast cells (HMC-1), *C. elegans* nematodes, two yeast species
S. cerevisiae and *S. pombe,* as well as the parasitic protozoan *T. brucei* using electron microscopy
(Figure 5**a**). Since the HMC-1 nucleus has a diameter of ~8 µm (Figure 5**b**) we chose to examine
sections corresponding to a little more than one nuclear volume (8 µm/70 nm thick sections ≈ 114
serial sections/nucleus). By the same logic, the number of thin sections examined corresponded to
approximately four total nuclear volumes of *T. brucei* and *S. pombe* and five nuclear volumes of *C.*
elegans and *S. cerevisiae* cells. For *C. elegans*, images of nuclei were acquired from intestinal cells
and oocytes.
311
312
In HMC-1 cells, three morphologically different categories of NEBs were observed (Figure 5**a**). The
most commonly observed morphology was outwards protruding buds, occurring in 7.3% of nuclear
sections (n = 9; Figure 5**a**-**b**), which will hereafter be referred to as type 1 NEB. In a few cases of
type 1 NEB events, both the outer and inner membranes expand towards the cytoplasm with
electron dense material partially wrapped by the nuclear envelope bud and the protrusion is open
to the nucleoplasm (Figure 5**b**). In most of the type 1 NEB events however (n= 6), the electron
dense material is clearly located between the two membranes with only the outer membrane
expanding towards the cytoplasm. The second type of NEB events observed had the inner nuclear
envelope expanding towards the nucleoplasm. These were termed type 2 NEB events, and
occurred in 3.3% of sections. Finally, for type 3 NEB events the transported material is situated
between the nuclear membranes with no clear protrusion in either direction. This occurred in 2.5%

of sections. Some thin sections of nuclear envelopes contained more than one bud, revealing that

NEB can occur at multiple sites on the same nucleus (Figure S1).

NEB events are observed to occur in all five organisms examined (Figure 5**a**) but with varying

frequency. The lowest frequency of NEB events was in *S. cerevisiae* (2.5%, n=161 sections),

followed by *S. pombe* (6.9%, n=101 sections), *T. brucei* (8.4%, n=119 sections), *C. elegans* (11.6%,

n=138 sections) and most frequently in HMC-1 cells (13.1%, n=122). Only type 1 NEB events were

identified in all organisms other than HMC-1 (Figure 5**b-e**), except for a single event in *S. pombe*

that we classified as type 3 (Figure 5**f**). The morphology of the NEB events varied between and

within species, with some buds being small and having indistinct electron dense material whereas

other events were large and had a membrane surrounding the material. The average size of the

NEB events did not show a large variation between species except for HMC-1 cells, which yielded

distinctly larger buds (Figure S6).

The morphology of NEB was compared in more detail through electron tomography reconstructions

of NEB events from both *S. pombe* and *T. brucei*. In *S. pombe*, one event of type 1 NEB was

visualized (Figure 6**a-c**) whereas two type 1 NEB events were reconstructed from *T. brucei* (Figure

6**d-h**). The NEB event that appeared to have progressed the furthest had distinct regions of higher

electron density surrounding the base of the bud (Figure 6**g**, black arrow). For both *S. pombe and*

T. brucei the cargoes of the NEB events were found between, and fully separated from, the nuclear

membranes, and were similar to what was observed in *S. cerevisiae* (Figure 4**g**). Moreover, the

NEB event observed for *S. pombe* clearly displayed the presence of a lipid bilayer that completely

enveloped the cargo (Figure 6**b**). However, a clear bilayer surrounding the transported material

could not be ascertained in the *T. brucei* tomographic reconstruction. Since in many cases the

transported material appeared to have a vesicle-like structure, we tested for an involvement of the

ESCRT pathway known to facilitate the formation of cellular vesicles, and already implicated in the

formation of morphologically similar protrusions of the nuclear envelope (78, 79). However, ESCRT

components did not appear to contribute to the formation of NEB events in our experiments (S7).

We conclude that NEB was observed in all species examined in this study. Our findings combined

with previous published observations are summarized into a phylogenetic tree (Figure 7),

illustrating how the NEB pathway is evolutionarily conserved among eukaryotes and is part of
normal cellular function in an evolutionarily diverse sampling of species.

357
358
359

Discussion

Nuclear envelope budding (NEB) has been observed since 1955 as nuclear herniations across a
diverse range of organisms and at different developmental stages. This phenomenon is well
studied with regards to its role in virus infection and the detailed structure of NEB events have
been revealed using cryo-electron tomography of infected cells (11). Yet, NEB remains viewed as
an irregularity rather than a normal route for transport over the nuclear envelope in healthy
mature cells. We present here three different lines of evidence that NEB is a widely conserved,
physiologically normal cellular process that increases in frequency due to cellular stress and is
specifically activated by an increase in protein aggregation. Firstly, we observed that five distinct
stress conditions: heat shock; hydrogen peroxide; arsenite; proteasome inhibition and AZC
treatment, all led to an increase in the frequency of NEB, with the AZC treatment most
prominently activating the NEB pathway among all stressors. Secondly, immuno-EM revealed the
presence of ubiquitin and Hsp104, a protein disaggregase, in the cargo of the buds, which
supports our hypothesis that this pathway is involved in protein degradation. Thirdly, NEB events
were detected in every species examined, from T. brucei to human cell line HMC-1. Collectively,
these results suggest a role of the NEB pathway in protein quality control and highlight its
evolutionary conservation. These observations shed light on the cell's mechanism to cope with
proteotoxic stress and identify a new module of the nuclear quality control system. Moreover,
since the NEB pathway is evolutionarily conserved and is also present in human cells, this work
provides a new perspective for studying cellular stress response related human diseases (80).

Nuclear budding and the protein quality control system
Although chaperones and the ubiquitin-proteasome system are known to clear misfolded proteins
from the nucleus (81, 82), the nuclear protein quality control system is not as well characterized
as the cytosolic and the ER systems (39). Various cellular events and pathways have been
described whereby aggregated proteins and chaperones are imported from the cytosol and into
the nucleus and these pathways are consistent with observations that 80% of the proteasomes at
steady state are located within the nucleus (83, 84). An alternative pathway, however, has also
been reported for which ubiquitinated proteins in mammalian cells and in the nematode *C.*

elegans are transported from the nucleus to the cytosol via a ubiquitin-associated domain-
containing protein (UBIN) (85).

Some proteins that exceed the size limit of the NPCs may nevertheless enter the complex after a
reversible conformational change in their tertiary structure (86). Although large aggregates could
potentially be transported in a similar manner, it stands to reason that under conditions of cellular
stress, the transportation of many aggregates through NEB events would provide a more efficient
and therefore advantageous gateway into or out of the nucleus. In support of the above
supposition, a process called "piecemeal microautophagy of the nucleus" has been described in
S. cerevisiae in which nuclear envelope buds are released directly into the vacuole lumen for
degradation (49).

Our observations that NEB increases during five different cellular stresses (heat shock, hydrogen
peroxide, arsenite, proteasome inhibition and AZC exposure) in *S. cerevisiae* strongly implicate
that NEB is related to the protein quality control system. Aging has been previously shown to
influence the formation of NEB-like events in the perinuclear space (54), with an apparent
increased in replicatively aged cells of about 17% when compared to the remaining young
population (stated as mixed population). Though the isolation method we used in this study
involved a higher number of isolation steps in comparison with the aforementioned literature, we
were unable to reproduce the difference in NEB frequency between the old and young population.
There is a significant increase in both isolated cell groups with a stronger effect on the aged cells.
Possible reasons for the inability to reproduce what has been shown before could be due to the
differences in the isolation process. Conclusions that can be safely extracted are that mechanical
stress is also able to increase NEB frequency and aging most likely creates a further stressful
environment for the cells.
The presence of ubiquitin and Hsp104 in the NEB events lends further support for this
functionality due to the fact that ubiquitin is involved in many protein quality control pathways
including the ubiquitin-proteasome system and macroautophagy, and Hsp104 is known to
mediate disaggregation of misfolded proteins (87-89). The large increase in NEB events due to
the AZC treatment reinforces this putative role of NEB as a cellular stress response. Since AZC
acts upon protein conformational stability directly without influencing other parameters such as
the external temperature or membrane fluidity, these experiments constitute a more targeted
challenge for the protein quality control system. Additionally, in a previous study where
aggregated proteins were identified after cellular stress induced by AZC, arsenite and hydrogen
peroxide treatment, AZC had the largest number of affected proteins among the stressors (58). A

strong effect of AZC on the folding of proteins could explain why NEB is more prominent in the

AZC-induced stress response when compared to the cellular response to other stressors.

Finally, the connection between proteasomal activity and NEB frequency supports the association

with the protein quality control system. Inhibition of proteasomal activity showed a clear increase

in NEB events which could be translated as a cellular attempt to cope with a large number of

misfolded proteins or proteins which are normally degraded by the proteasome. Most of the NEB

events recorded in the *rpn4Δ* strain, had a distinct morphology compared to NEB events triggered

by other stressors. This observation may reflect a difference in the nature and composition of the

transported material, indicating that NEB may well be involved in more than one cellular activity.

The distinction between NEB events and defective NPCs

Deleting key genes required for NPC assembly can induce nuclear envelope "herniations" that

appear somewhat similar in structure to NEB events (73-76, 90, 91). These herniations are

believed to arise from the outer nuclear membrane sealing above the defective NPC, thereby

capturing the transported material between the NPC and the nuclear envelope and creating a

bulge resembling a budding vesicle. If this phenomenon underpinned our observed NEB events,

the intermembrane vesicle should be connected to the INM where the defective NPC is located. A

connection between the INM and intermembrane vesicle resembling a "neck" would also be a

plausible intermediate state in the formation of a vesicle from the INM that encloses nuclear

material. Such neck-like connections between the INM and contents of the bud were previously

observed in strains deficient in TorA, an ER protein with ATPase activity, and various Nups that

lead to defective NPCs (31, 73, 74). Furthermore, the size of the necks seen in NEB events after

hyperactivation of Chm7, a component of the ESCRT-III like complex, and misassembled NPCs

were compared previously and it was concluded that while appearing similar, both features were

morphologically distinct (79). Neck-like or NPC-like structures connecting the intermembrane

vesicles to the INM were not visible in any of our acquired tomograms, suggesting that our

tomograms contain buds in which the intermembrane vesicle or material has fully formed and

these buds are not the result of NPC malformation. In addition to these morphological differences

revealed by 3D tomographic reconstructions, the immuno-EM assay demonstrated that the NEB

events were not associated with NPCs since only two of the 16 observed events in this assay

were labeled for NPC proteins.

The evolutionarily conserved nature of NEB

We observed NEB events within the cells of all species we examined including animals, fungi and

protozoa. There is a thread stretching back through the scientific literature which shows that NEB

events have repeatedly been observed in animals, plants and protists (Table S1). The genetic

diversity of these observations is summarized in Figure 7, which presents a phylogenetic tree

illustrating that NEB events have been observed for 18 different species within the eukaryotic

domain. Therefore, despite NEB not being widely recognized as a means of nuclear transport,

these observations combine to create a compelling argument that NEB is an evolutionarily

conserved phenomenon of eukaryotic cells.

From our sequence of experiments using the model systems *S. cerevisiae* we concluded that

NEB events facilitate the removal of aggregated proteins from the nucleus. This pathway could

potentially also support the co-transportation of other cellular components required for protein

degradation. Since NEB embodies an alternative route for transport over the nuclear envelope

that has been largely overlooked, our observations highlight an opportunity for discovery in

eukaryotic cell biology where a broad range of interconnected questions may emerge. It will be

necessary to determine if the frequency of NEB is also increased by cellular stressors in human

cells, and in particular additional cellular stress associated with aging. One issue is the important

role of protein aggregation within a variety of neurodegenerative diseases such as Alzheimer's

and ALS (80, 92). Should NEB also be an underappreciated mechanism for the clearance of

protein aggregates in human neurons then this potentially has far-reaching neurological

implications.

Materials and Methods

Antibodies and reagents

Please refer to Supplementary tables 3 and 4 for a detailed information on antibodies, strains,

primers and reagents.

Experimental models

HMC-1 cells

Human mast cell line 1 (HMC-1; (93)) cells were cultured in Iscove's modified Dulbecco's medium

(IMDM). The cells were then treated with 10% exosome-depleted fetal bovine serum (FBS), 100

units/ml streptomycin, 100 units/ml penicillin, 2 mM L-glutamine, and 1.2 U/ml alpha-thioglycerol

in incubators kept at 37°C and 5% carbon dioxide (94, 95).

Trypanosoma brucei

Procyclic *T. brucei* strain 427 was cultured in SDM-79 media with 20% fetal bovine serum.

Cultures were prepared and maintained at a concentration of 5×10^5 and 1×10^7 cells per mL

(96-98).

Schizosaccharomyces pombe

Logarithmically growing wild type fission yeast *S. pombe* were grown at 30°C in YE5S medium

(94, 99, 100).

Caenorhabditis elegans

The *C. elegans* wild type reference strain was the Bristol N2 variety. The worms were cultured on

normal growing media plates (NGM plates) and the *E. coli* strain OP50 was used as a food

source. The worms were maintained at the optimal temperature of 25°C (101, 102). Adult worms

were used for electron microscopy.

Saccharomyces cerevisiae

Wild type cells of *S. cerevisiae* (BY4741) were cultured in YPD media at 30°C (103). A strain with

endogenous *HSP104* C-terminally tagged with GFP (104) was used for old cell isolation, heat

shock and sodium arsenite experiments. The deletion mutants *hsp104Δ* and *rpn4Δ* are from the

YKO collection (EUROSCARF, Frankfurt, Germany). Strains used for investigating involvement of

the ESCRT-pathway in NEB and the *pdr5Δ* strain are from this study (see yeast strains list).

Transformations were performed following standard procedures (105), and gene deletions and

endogenous tags were integrated via homologous recombination (106). For the hydrogen

peroxide experiment, cells were grown in synthetic complete media (with yeast nitrogen base,

without amino acids, pH 5.5, complete supplement mixture of amino acids and 2% glucose).

High-pressure freezing for electron microscopy and tomography

All samples used in this study has been prepared using high-pressure freezing followed by freeze

substitution. HMC-1 cells were loaded into membrane carriers and were high-pressure frozen in a

Leica EM PACT1 (Leica microsystems, Wetzlar, Germany). *T. brucei* and *S. pombe* were loaded

into carriers and were high-pressure frozen in a Leica EM PACT2. *C. elegans* and *S. cerevisiae*

samples were loaded into aluminum specimen carriers and were high-pressure frozen in a

Wohlwend Compact 3 (M. Wohlwend GmbH, Sennwald, Switzerland). Yeast paste from wild type

S. cerevisiae cells was used as a cryoprotectant filling the space surrounding the worms.

For a summary of all high-pressure freezing and freeze substitution experiments see table 1. For

all samples (except *S. pombe*) a short freeze substitution protocol was applied, using 2% uranyl

acetate dissolved into acetone (UA; from 20% UA stock in methanol) for one hour (94, 97, 107).

To increase the penetration into intact worms, the UA solution was left on the samples for 14h (as

the temperature was increased to -50°C). The UA incubation was followed by two washes in

100% acetone for one hour each. Before embedding, the temperature was raised from -90°C to -

50°C overnight with a rate of 3°C/h. Samples were then embedded in K4M or HM20 resin in

increasing concentrations of 20%, 40%, 50%, 80% and finally three times in 100% plastic (2

hours per solution). Polymerization of the plastic occurred over 48h using UV light at -50°C

followed by 48h in room temperature. *S. pombe* cells were high-pressure frozen and fixed

through freeze substitution (long protocol) with anhydrous acetone containing 0.25% UA, 0.1%

dehydrated glutaraldehyde and 0.01% osmium tetroxide (OsO_4) (99, 100, 108, 109).

For tomography, serial semi thick sections of about 210-250 nm (*S. pombe*) and 350 nm (*T.*

brucei and *S. cerevisiae*) were cut and samples were poststained with 2% UA in dH_2O followed

by Reyonold's lead citrate. Gold particles (15 nm) from British Bio Cell were applied to both sides

of the grid to be used as fiducial markers. (96, 97, 99).

All other samples were sectioned in 70 nm thin sections and placed on either copper slot or mesh

grids. Sections were stained with 2% UA for 5 minutes and Reynold's lead citrate for 1 minute

(110). Washing steps after each staining were performed in dH_2O.

Immuno-electron microscopy

For the immuno-labeling experiments, the same high-pressure frozen samples embedded in

HM20 resin were used, a benefit of that short FS protocol (107). For a summary of these

experiments, see table 2. Grids with 70 nm thick sections were fixed in 1% paraformaldehyde

(PFA) in PBS for 10 minutes. After three PBS washes of 1 minute each, samples were blocked

with 0.1% fish skin gelatin and 0.8% BSA in PBS for 1 hour. For detection of NPC proteins, grids

were then incubated in a 1:50 dilution of mAb414 (BioLegend, San Diego, USA) for two hours,

followed by a 1:150 dilution of rabbit anti-mouse immunoglobulins (Agilent/Dako, Glostrup,

Denmark) for an hour, and then a 1:70 dilution of 10 nm gold-conjugated protein A (CMC UMC

Utrecht, The Netherlands) for 30 minutes. For labeling of ubiquitin, grids were incubated in a 1:20

dilution of antibody ab19247 (Abcam, Cambridge, UK) for 2 hours. Detection of GFP was

performed by using a 1:5, 1:10, or 1:30 dilution of ab6556 (abcam, Camebridge, UK) and

detection of Hsp104 by using a 1:100 dilution of ab69549 (abcam, Camebridge, UK) incubated

overnight. Goat-anti-Rabbit IgG 10 nm gold (Electron Microscopy Sciences, Hatfield PA, USA)

was then used at a 1:20 dilution for an hour. All incubations were performed at room

temperature, except for the primary antibody which was kept at 4°C. Three washing steps (20 min

in PBS) were carried out after incubations with each antibody. 2.5% glutaraldehyde was applied

to sections for 1 hour followed by three washes (1 min in dH_2O). Sections were then contrast

stained in 2% UA for 5 minutes (wash 3x 2 min in dH_2O) and 1 minute in Reynold's lead citrate

(washed 5x 1 min in dH_2O).

Image acquisition and electron tomography

All thin sections were imaged at 120 kV either on a LEO 912 OMEGA (Zeiss, Krefeldt, Germany)

equipped with a 2k x 2k VELETA Olympus CCD camera or on a Tecnai T12 transmission

electron microscope equipped with a Ceta CMOS 16M camera (FEI Co., Eindhoven, The

Netherlands). Double axis tilt series of serial sections of *T. brucei* samples were acquired every

degree using the serialEM software (111) on a Tecnai TF30 300 kV IVEM microscope (FEI Co.,

The Netherlands) equipped with a Ultrascan 785 4k x 4k camera binned to 2k x 2k (pixel size 1

nm). For *S. pombe*, digital images (Gatan Ultrascan 890 or 895 camera, pixel size 1.5 nm) single

axis tilt series were taken every 1.5° over a ±60°–65° range operating a Tecnai TF20 electron

microscope (FEI Co., Eindhoven, The Netherlands) (99). For *S. cerevisiae*, double axis tilt series

of serial sections were acquired on a Tecnai TF30 300 kV microscope (FEI Co., The Netherlands)

equipped with a Gatan One View camera (pixel size 1.6 nm, increment 1.5° over a ±60° range).

Tomograms were acquired with the use of serialEM (111). The tomographic reconstruction was

performed using of the IMOD software package (112).

Isolation of old yeast cells

Isolation of old yeast cells was performed according to Smeal et al. with some modifications in

order to reduce mechanical stress and improve the cell morphology for electron microscopy (56).

In brief, the cell surface of exponentially growing cells was labeled with biotin by incubating the

cells with 0.5 mg/ml Sulfo-NHS-LC biotin (#21335, Thermo Fisher scientific) for 20 min at room

temperature. Cells were grown in YPD, harvested prior saturation of culture, and washed in PBS

+ 0.5% glucose. Biotinylated cells were labeled with 17.5 ug/ml Streptavidin magnetic beads

(#21344, Thermo Fisher scientific) for 1.5 hours followed by 3 x 15 min magnetic sorting with PBS

+ 0.5% glucose washes in between. Another two rounds of growth, streptavidin-labeling, and

sorting was performed. The old cells and their unbound daughters were recovered in YPD for 4

hours before high-pressure freezing.

Stress treatments of *S. cerevisiae*

Cells were grown at 30°C to mid exponential phase. For mild heat shock, the culture was shifted

to 38°C and samples were collected after 0, 5, 15, 30, 45, and 90 minutes (113). The more

severe heat shock was performed at 42°C for 30 min. For hydrogen peroxide, sodium arsenite,

and AZC treatments the culture was split into two, one kept as unstressed control and one treated

with the stressor. Cells were exposed to 0.6 mM hydrogen peroxide for 90 minutes, 0.5 mM

sodium arsenite for 60 minutes, or 1 mg/mL AZC for 30 and 90 minutes (corresponding control

culture was grown for 60 minutes after mid exponential phase). Cells were harvested by filtration

followed by high-pressure freezing.

Proteasome inhibition

The *rpn4Δ* strain and a wild type control was grown to mid exponential phase and prepared for

high-pressure freezing. The *pdr5Δ* strain was grown to mid exponential phase prior addition of

either 50 μM MG132 (#C2211, Sigma-Aldrich) dissolved in DMSO or an equivalent volume of

DMSO. A 60 min incubation at 30°C followed before cells were prepared for high-pressure

freezing.

Analysis of polyubiquitination

Cells were cultured and treated as described above. Mid-exponentially grown yeast
corresponding to OD_{600} = 20 was harvested, washed once in distilled water and resuspended in 1
ml lysis buffer (100 mM Tris pH 7.5, 100 mM NaCl, 5 mM ethylenediaminetetraacetic acid, 1 mM
dithiothreitol, 1 mM phenylmethylsulfonyl fluoride, 20 mM N-ethylmaleimide). Homogenisation
was conducted via an Avestin Emulsiflex C-15, applying a homogenization pressure of 18 000
psi. Samples were cleared from cell debris and unlysed cells by centrifugation for 5 min at 3500
rcf and 4°C. 100 µl of cleared lysate was mixed with the same volume of 2x Laemmli buffer (100
mM Tris pH 6.8, 4% SDS, 20% glycerol, 0.2% bromophenol blue, 200 mM 2-mercaptoethanol)
and incubated for 15 min at 95°C. The samples were then applied for SDS-PAGE and
immunoblotting following standard protocols. Equal loading was controlled via Ponceau S staining
directly after wet electrotransfer on PVDF membranes and blots were decorated with an anti-
ubiquitin antibody (1:1000, HRP-conjugated, P4D1, sc8017, Santa Cruz). Clarity Western ECL
Substrate (BIO-RAD, 1705060) and a ChemiDoc XRS+ Imaging System (BIO-RAD, 1708265)
were used for detection

Analysis of cell death

Loss of membrane integrity was assessed with propidium iodide (PI) staining as previously

described (114). Briefly, cells were harvested in 96-well plates after the respective stressor/mock

treatment and resuspended in 250 µl of PI solution (100 µg/l PI in phosphate buffered saline PBS;

25 mM potassium phosphate, 0.9% NaCl; adjusted to pH 7.2). After 10 min incubation in the dark,

cells were washed with 250 µl of PBS and analysed via flow cytometry (Guava easyCyte 5HT;

Merck group). 5000 events were recorded per strain and condition using InCyte software (3.1).

Confocal microscopy

For visualization of nuclei, yeast cells were harvested and resuspended in DRAQ5 staining

solution (5 µM DRAQ5 in PBS). After 10 min incubation in the dark, cells were washed once with

PBS and immobilized on agar slides. Specimen were analyzed with a ZEISS LSM700 microscope

using ZEISS ZEN blue software control. Plan-Apochromat 63x/1.40 Oil M27 objective was

employed. Appropriate filter settings were used to visualize GFP and DRAQ5. Micrographs were

analyzed and processed with the open-source software Fiji (115). To reduce image noise,

Gaussian filtering (σ = 1) was applied, followed by background subtraction (rolling ball radius =

100 pixels). Pictures within an experiment were captured and processed in the same way.

Phylogenetic tree

A phylogenetic tree was constructed showing all the organisms where NEB events have been

observed. The tree was generated based on the NCBI taxonomy browser for scientific names and

visualized by the EvolView online software (http://www.evolgenius.info/evolview/).

Statistics and reproducibility

The frequency of NEB events was achieved by counting the number of events present in

approximately 100-200 thin sections of nuclei. The percentage of sections containing events in

each organism was presented as a bar graph. For the NPC immuno-EM assay, different cell

compartments (NPCs, NEBs and lipid droplets) were categorized as labeled or not labeled based

on the presence or absence of gold particles. Similarly, the percentage of labeled compartments

was presented as a bar graph. For the statistical analysis, different statistical tests were

examined but a non-parametric Wilcoxon test was performed as our data represent frequencies

and the numerical values are not following a normal distribution nor are continuous variants

(values were considered as 0 for the absence of events and 1 for the presence of events).. The

test was performed using the MATLAB multi-paradigm programming language. For further details

on the examined statistic tests see Supplementary table 2. . For the ubiquitin and Hsp104-GFP

immuno-EM, the area and number of gold particles of different cell compartments were measured

using IMOD (https://bio3d.colorado.edu/imod/). The amount of gold particles per area was

presented as bar graph. For cell death analysis,

results are displayed either as dot plots (if n ≤ 5) where mean (square), median (centre line) and

s.e.m. are depicted, or as box plots (if n > 5) with mean (square) and median (centre line) as well

as whiskers presenting minima and maxima within 2.2 interquartile range (IQR). Outliers were

defined by the 2.2-fold IQR labelling rule. Normal distribution of data was examined with a

Shapiro-Wilk's test and homogeneity of variances was evaluated with a Levene's test. A student's

t-test was used to compare between two groups and displayed significances are two-sided, and a

one-way ANOVA with a Bonferroni post hoc test was applied to compare between three and

more groups. Significances are indicated with asterisks: *****p*<.0001, ****p*< .001, ***p* < .01, **p* <

.05, n.s. not significant. Further information on sample size can be found in the respective figure

legends. All figures were processed with Origin Pro 2017 or Adobe Illustrator CS6 (Adobe).

Data availability

Links to the online library for the 2D electron microscopy pictures acquired during this study:

http://cellimagelibrary.org/groups/50813

http://cellimagelibrary.org/images/50818

http://cellimagelibrary.org/groups/50832

http://cellimagelibrary.org/groups/50841

http://cellimagelibrary.org/groups/50849

http://cellimagelibrary.org/groups/50873

http://cellimagelibrary.org/images/50882

http://cellimagelibrary.org/images/50888

http://cellimagelibrary.org/images/50900

http://cellimagelibrary.org/groups/50906

http://cellimagelibrary.org/images/50920

http://cellimagelibrary.org/groups/50935

http://cellimagelibrary.org/images/50945

Summarized protocols

Please refer to supplementary table 5 for a summarized information on thin sections, tomograohy

samples and immuno-EM samples preparation.

Acknowledgments

This project was funded by the Swedish Research Council Grant (2015-00560) and the Knut and

Alice Wallenberg Foundation (grant KAW 2012.0284, KAW 2014.0275) to RN, the Swedish

Research Council Grants (621-2014-4597, 2018-03577) to MJT, the Swedish Research Council

Young Investigator Grant (2015-05427), the Austrian Science Fund (J4342-B21) to VK and

Swedish Research Council grant (2019-04004) to JLH and Knut and Alice Wallenberg grant

(KAW 2017.0091) to TN and JLH.

All authors would like to thank Marc Pilon for providing the *C. elegans* worms and for contributing

intellectually to the manuscript writing. Further on, we would like to thank Rebecca Andersson for

her help in conducting the hydrogen peroxide experiment and Per Widlund for reagents. We

would also like to thank Claes Andreasson, Per Widlund and Martin Ott for helpful discussions.

704
705
706
707 **References**
708

1.   M. L. Watson, Pores in the mammalian nuclear membrane. *Biochim Biophys Acta* **15**, 475-
479 (1954).
2.   G. Kabachinski, T. U. Schwartz, The nuclear pore complex – structure and function at a
glance. *Journal of Cell Science* **128**, 423-429 (2015).

3.   M. Beck, E. Hurt, The nuclear pore complex: understanding its function through structural
insight. *Nature Reviews Molecular Cell Biology* **18**, 73-89 (2017).

4.   Y. M. Chook, G. Blobel, Karyopherins and nuclear import. *Curr Opin Struct Biol* **11**, 703-
715 (2001).

5.   A. Hoelz, E. W. Debler, G. Blobel, The Structure of the Nuclear Pore Complex. *Annual*
Review of Biochemistry **80**, 613-643 (2011).

6.   H. B. Schmidt, D. Görlich, Transport Selectivity of Nuclear Pores, Phase Separation, and
Membraneless Organelles. *Trends in Biochemical Sciences* **41**, 46-61 (2016).

7.   J. Garcia-Bustos, J. Heitman, M. N. Hall, Nuclear protein localization. *Biochimica et*
Biophysica Acta (BBA) - Reviews on Biomembranes **1071**, 83-101 (1991).

8.   L. J. Terry, E. B. Shows, S. R. Wente, Crossing the Nuclear Envelope: Hierarchical
Regulation of Nucleocytoplasmic Transport. *Science* **318**, 1412-1416 (2007).

9.   A. Kohler, E. Hurt, Exporting RNA from the nucleus to the cytoplasm. *Nat Rev Mol Cell Biol*
8, 761-773 (2007).

10.   J. Schwartz, B. Roizman, Concerning the egress of herpes simplex virus from infected cells:
electron and light microscope observations. *Virology* **38**, 42-49 (1969).

11.   C. Hagen *et al.*, Structural Basis of Vesicle Formation at the Inner Nuclear Membrane. *Cell*
163, 1692-1701 (2015).

12.   L. G. Fradkin, V. Budnik, This bud's for you: mechanisms of cellular nucleocytoplasmic
trafficking via nuclear envelope budding. *Curr Opin Cell Biol* **41**, 125-131 (2016).

13.   I. H. Wang, C. J. Burckhardt, A. Yakimovich, U. F. Greber, Imaging, Tracking and
Computational Analyses of Virus Entry and Egress with the Cytoskeleton. *Viruses* **10**
(2018).

14.   A. C. Longwell, G. Yerganian, SOME OBSERVATIONS ON NUCLEAR BUDDING AND
NUCLEAR EXTRUSIONS IN A CHINESE HAMSTER CELL CULTURE. *J Natl Cancer Inst* **34**, 53-
69 (1965).

15.   R. N. Elston, B. Stenkvist, Quantitative estimation of nuclear buds and micronuclei in
bovine cells transformed by Rous sarcoma and SV 40 viruses. *Zeitschrift für Zellforschung*
und Mikroskopische Anatomie **68**, 543-549 (1965).

16.   R. Hadek, H. Swift, Nuclear extrusion and intracisternal inclusions in the rabbit blastocyst.
The Journal of cell biology **13**, 445-451 (1962).

17.   D. Szollosi, Extrusion of nucleoli from pronuclei of the rat. *The Journal of cell biology* **25**,
545-562 (1965).

18.   M. S. Szollosi, D. Szollosi, 'Blebbing' of the nuclear envelope of mouse zygotes, early
embryos and hybrid cells. *J Cell Sci* **91 (Pt 2)**, 257-267 (1988).

19.   H. Gay, NUCLEO-CYTOPLASMIC RELATIONS IN SALIVARY-GLAND CELLS OF Drosophila.
Proceedings of the National Academy of Sciences **41**, 370-375 (1955).

20.   R. H. Mepham, G. R. Lane, Observations on the fine structure of developing microspores
ofTradescantia bracteata. *Protoplasma* **70**, 1-20 (1970).

21.   B. M. Gullvåg, Release of Nuclear Material During the Development of Lycopodium
Annotinum L. Spores. *Grana* **10**, 31-34 (1970).

22.   B. A. Afzelius, THE NUCLEUS OF NOCTILUCA SCINTILLANS: Aspects of Nucleocytoplasmic
Exchanges and the Formation of Nuclear Membrane. *The Journal of cell biology* **19**, 229-
238 (1963).

23.   H. G. Dickinson, Nucleo-Cytoplasmic Interaction following Meiosis in the Young
758   Microspores of Lilium Longiflorum; Events at the Nuclear Envelope. *Grana* **11**, 117-127
759   (1971).
760   24.   H. G. Dickinson, P. R. Bell, Nucleocytoplasmic interaction at the nuclear envelope in post
761   meiotic microspores of Pinus banksiana. *J Ultrastruct Res* **33**, 356-359 (1970).
762   25.   H. C. Aldrich, I. K. Vasil, Ultrastructure of the postmeiotic nuclear envelope in microspores
763   of Podocarpus macrophyllus. *J Ultrastruct Res* **32**, 307-315 (1970).
764   26.   M. Hochstrasser, J. W. Sedat, Three-dimensional organization of Drosophila melanogaster
765   interphase nuclei. II. Chromosome spatial organization and gene regulation. *The Journal*
766   *of cell biology* **104**, 1471-1483 (1987).
767   27.   J. M. Verboon *et al.*, Wash and the wash regulatory complex function in nuclear envelope
768   budding. *Journal of Cell Science* 10.1242/jcs.243576, jcs.243576 (2020).
769   28.   S. D. Speese *et al.*, Nuclear envelope budding enables large ribonucleoprotein particle
770   export during synaptic Wnt signaling. *Cell* **149**, 832-846 (2012).
771   29.   N. LaMassa, C. Arenas-Mena, G. R. Phillips, Electron microscopic characterization of
772   nuclear egress in the sea urchin gastrula. *J Morphol* **279**, 609-615 (2018).
773   30.   E. M. Hatch, M. W. Hetzer, RNP export by nuclear envelope budding. *Cell* **149**, 733-735
774   (2012).
775   31.   V. Jokhi *et al.*, Torsin mediates primary envelopment of large ribonucleoprotein granules
776   at the nuclear envelope. *Cell Rep* **3**, 988-995 (2013).
777   32.   B. Ding, A. M. Mirza, J. Ashley, V. Budnik, M. Munson, Nuclear Export Through Nuclear
778   Envelope Remodeling in Saccharomyces cerevisiae. *bioRxiv* 10.1101/224055,
779   224055 (2017).
780   33.   A. Rose, C. Schlieker, Alternative nuclear transport for cellular protein quality control.
781   *Trends in cell biology* **22**, 509-514 (2012).
782   34.   D. Oling, F. Eisele, K. Kvint, T. Nystrom, Opposing roles of Ubp3-dependent
783   deubiquitination regulate replicative life span and heat resistance. *The EMBO journal* **33**,
784   747-761 (2014).
785   35.   N. Erjavec, L. Larsson, J. Grantham, T. Nystrom, Accelerated aging and failure to segregate
786   damaged proteins in Sir2 mutants can be suppressed by overproducing the protein
787   aggregation-remodeling factor Hsp104p. *Genes & development* **21**, 2410-2421 (2007).
788   36.   T. Dubnikov, T. Ben-Gedalya, E. Cohen, Protein Quality Control in Health and Disease. *Cold*
789   *Spring Harbor perspectives in biology* **9** (2017).
790   37.   A. K. Basu, DNA Damage, Mutagenesis and Cancer. *Int J Mol Sci* **19**, 970 (2018).
791   38.   E. Shor, C. A. Fox, J. R. Broach, The yeast environmental stress response regulates
792   mutagenesis induced by proteotoxic stress. *PLoS genetics* **9**, e1003680 (2013).
793   39.   Y. Shibata, R. I. Morimoto, How the nucleus copes with proteotoxic stress. *Current biology*
794   *: CB* **24**, R463-474 (2014).
795   40.   N. Panté, M. Kann, Nuclear Pore Complex Is Able to Transport Macromolecules with
796   Diameters of ∼39 nm. *Molecular biology of the cell* **13**, 425-434 (2002).
797   41.   S. R. Wente, M. P. Rout, The nuclear pore complex and nuclear transport. *Cold Spring*
798   *Harbor perspectives in biology* **2**, a000562 (2010).
799   42.   A. B. Mehrtash, M. Hochstrasser, Ubiquitin-dependent protein degradation at the
800   endoplasmic reticulum and nuclear envelope. *Semin Cell Dev Biol* **93**, 111-124 (2019).

801   43.   G. Mannhaupt, R. Schnall, V. Karpov, I. Vetter, H. Feldmann, Rpn4p acts as a transcription
802   factor by binding to PACE, a nonamer box found upstream of 26S proteasomal and other
803   genes in yeast. *FEBS Letters* **450**, 27-34 (1999).
804   44.   B. A. Koch, H. Yu, Regulation of inner nuclear membrane associated protein degradation.
805   *Nucleus* **10**, 169 - 180 (2019).
806   45.   M. Deng, M. Hochstrasser, Spatially regulated ubiquitin ligation by an ER/nuclear
807   membrane ligase. *Nature* **443**, 827-831 (2006).
808   46.   M. Boban, M. Pantazopoulou, A. Schick, P. O. Ljungdahl, R. Foisner, A nuclear ubiquitin-
809   proteasome pathway targets the inner nuclear membrane protein Asi2 for degradation. *J
810   Cell Sci* **127**, 3603-3613 (2014).
811   47.   O. Foresti, V. Rodriguez-Vaello, C. Funaya, P. Carvalho, Quality control of inner nuclear
812   membrane proteins by the Asi complex. *Science* **346**, 751-755 (2014).
813   48.   J. O'Driscoll, D. Clare, H. Saibil, Prion aggregate structure in yeast cells is determined by
814   the Hsp104-Hsp110 disaggregase machinery. *The Journal of cell biology* **211**, 145-158
815   (2015).
816   49.   P. Roberts *et al.*, Piecemeal microautophagy of nucleus in Saccharomyces cerevisiae.
817   *Molecular biology of the cell* **14**, 129-141 (2003).
818   50.   R. Lum, J. M. Tkach, E. Vierling, J. R. Glover, Evidence for an unfolding/threading
819   mechanism for protein disaggregation by Saccharomyces cerevisiae Hsp104. *J Biol Chem*
820   **279**, 29139-29146 (2004).
821   51.   S. A. Comyn, B. P. Young, C. J. Loewen, T. Mayor, Prefoldin Promotes Proteasomal
822   Degradation of Cytosolic Proteins with Missense Mutations by Maintaining Substrate
823   Solubility. *PLoS Genet* **12**, e1006184-e1006184 (2016).
824   52.   J. L. Folch-Mallol, A. Garay-Arroyo, F. Lledias, A. A. Covarrubias Robles, [The stress
825   response in the yeast Saccharomyces cerevisiae]. *Revista latinoamericana de
826   microbiologia* **46**, 24-46 (2004).
827   53.   M. C. Haigis, B. A. Yankner, The aging stress response. *Molecular cell* **40**, 333-344 (2010).
828   54.   I. L. Rempel *et al.*, Age-dependent deterioration of nuclear pore assembly in mitotic cells
829   decreases transport dynamics. *eLife* **8**, e48186 (2019).
830   55.   R. Josefson, R. Andersson, T. Nystrom, How and why do toxic conformers of aberrant
831   proteins accumulate during ageing? *Essays in biochemistry* **61**, 317-324 (2017).
832   56.   T. Smeal, J. Claus, B. Kennedy, F. Cole, L. Guarente, Loss of transcriptional silencing causes
833   sterility in old mother cells of S. cerevisiae. *Cell* **84**, 633-642 (1996).
834   57.   T. Jacobson *et al.*, Arsenite interferes with protein folding and triggers formation of
835   protein aggregates in yeast. *Journal of cell science* **125** (2012).
836   58.   A. J. Weids, S. Ibstedt, M. J. Tamas, C. M. Grant, Distinct stress conditions result in
837   aggregation of proteins with similar properties. *Scientific reports* **6**, 24554 (2016).
838   59.   X.-C. Zeng *et al.*, Hsp70 dynamics in vivo: effect of heat shock and protein aggregation.
839   *Journal of Cell Science* **117**, 4991-5000 (2004).
840   60.   L. Fowden, M. H. Richmond, Replacement of proline by azetidine-2-carboxylic acid during
841   biosynthesis of protein. *Biochimica et Biophysica Acta* **71**, 459-461 (1963).
842   61.   E. W. Trotter *et al.*, Misfolded proteins are competent to mediate a subset of the
843   responses to heat shock in Saccharomyces cerevisiae. *The Journal of biological chemistry*
844   **277**, 44817-44825 (2002).

62. S. J. S. Berke, H. L. Paulson, Protein aggregation and the ubiquitin proteasome pathway:
gaining the UPPer hand on neurodegeneration. *Current Opinion in Genetics &*
Development **13**, 253-261 (2003).
63. P. K. Kim, D. W. Hailey, R. T. Mullen, J. Lippincott-Schwartz, Ubiquitin signals autophagic
degradation of cytosolic proteins and peroxisomes. *Proceedings of the National Academy*
of Sciences **105**, 20567-20574 (2008).
64. S. Pankiv *et al.*, p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of
ubiquitinated protein aggregates by autophagy. *The Journal of biological chemistry* **282**,
24131-24145 (2007).
65. G. Leppert *et al.*, Cloning by gene amplification of two loci conferring multiple drug
resistance in Saccharomyces. *Genetics* **125**, 13-20 (1990).
66. G. A. Collins, T. A. Gomez, R. J. Deshaies, W. P. Tansey, Combined chemical and genetic
approach to inhibit proteolysis by the proteasome. *Yeast* **27**, 965-974 (2010).
67. Y. Xie, A. Varshavsky, RPN4 is a ligand, substrate, and transcriptional regulator of the 26S
proteasome: A negative feedback circuit. *Proceedings of the National Academy of*
Sciences **98**, 3056-3061 (2001).
68. D. A. Parsell, A. S. Kowal, M. A. Singer, S. Lindquist, Protein disaggregation mediated by
heat-shock protein Hsp104. *Nature* **372**, 475-478 (1994).
69. J. R. Glover, S. Lindquist, Hsp104, Hsp70, and Hsp40: a novel chaperone system that
rescues previously aggregated proteins. *Cell* **94**, 73-82 (1998).
70. J. M. Tkach, J. R. Glover, Nucleocytoplasmic trafficking of the molecular chaperone
Hsp104 in unstressed and heat-shocked cells. *Traffic* **9**, 39-56 (2008).
71. G. M. Preston, C. J. Guerriero, M. B. Metzger, S. Michaelis, J. L. Brodsky, Substrate
Insolubility Dictates Hsp104-Dependent Endoplasmic-Reticulum-Associated Degradation.
Molecular cell **70**, 242-253.e246 (2018).
72. D. J. Thaller, C. Patrick Lusk, Fantastic nuclear envelope herniations and where to find
them. *Biochemical Society Transactions* **46**, 877-889 (2018).
73. S. R. Wente, G. Blobel, A temperature-sensitive NUP116 null mutant forms a nuclear
envelope seal over the yeast nuclear pore complex thereby blocking nucleocytoplasmic
traffic. *The Journal of cell biology* **123**, 275-284 (1993).
74. S. R. Wente, G. Blobel, NUP145 encodes a novel yeast glycine-leucine-phenylalanine-
glycine (GLFG) nucleoporin required for nuclear envelope structure. *The Journal of cell*
biology **125**, 955-969 (1994).
75. M. Allegretti *et al.*, In-cell architecture of the nuclear pore and snapshots of its turnover.
Nature **586**, 796-800 (2020).
76. C.-W. Lee *et al.*, Selective autophagy degrades nuclear pore complexes. *Nature Cell*
Biology **22**, 159-166 (2020).
77. S. Otsuka *et al.*, Nuclear pore assembly proceeds by an inside-out extrusion of the nuclear
envelope. *eLife* **5** (2016).
78. M. Vietri, M. Radulovic, H. Stenmark, The many functions of ESCRTs. *Nature Reviews*
Molecular Cell Biology **21**, 25-42 (2020).
79. D. J. Thaller *et al.*, An ESCRT-LEM protein surveillance system is poised to directly monitor
the nuclear envelope and nuclear transport system. *eLife* **8**, e45284 (2019).
80. I. Liguori *et al.*, Oxidative stress, aging, and diseases. *Clin Interv Aging* **13**, 757-772 (2018).

81. A. von Mikecz, M. Chen, T. Rockel, A. Scharf, The nuclear ubiquitin-proteasome system:
visualization of proteasomes, protein aggregates, and proteolysis in the cell nucleus.
Methods in molecular biology (Clifton, N.J.) **463**, 191-202 (2008).
82. S. V. Nielsen, E. G. Poulsen, C. A. Rebula, R. Hartmann-Petersen, Protein quality control in
the nucleus. *Biomolecules* **4**, 646-661 (2014).
83. R. Prasad, S. Kawaguchi, D. T. W. Ng, A Nucleus-based Quality Control Mechanism for
Cytosolic Proteins. *Molecular biology of the cell* **21**, 2117-2127 (2010).
84. S. H. Park *et al.*, PolyQ proteins interfere with nuclear degradation of cytosolic proteins
by sequestering the Sis1p chaperone. *Cell* **154**, 134-145 (2013).
85. S. Hirayama *et al.*, Nuclear export of ubiquitinated proteins via the UBIN-POST system.
Proceedings of the National Academy of Sciences **115**, E4199-E4208 (2018).
86. Shige H. Yoshimura, M. Kumeta, K. Takeyasu, Structural Mechanism of Nuclear Transport
Mediated by Importin β and Flexible Amphiphilic Proteins. *Structure* **22**, 1699-1710
(2014).
87. S. Shaid, C. H. Brandts, H. Serve, I. Dikic, Ubiquitination and selective autophagy. *Cell
Death Differ* **20**, 21-30 (2013).
88. R. F. Kalejta, T. Shenk, Proteasome-dependent, ubiquitin-independent degradation of the
Rb family of tumor suppressors by the human cytomegalovirus pp71 protein. *Proceedings
of the National Academy of Sciences* **100**, 3263-3268 (2003).
89. A. L. Yokom *et al.*, Spiral architecture of the Hsp104 disaggregase reveals the basis for
polypeptide translocation. *Nature Structural & Molecular Biology* **23**, 830-837 (2016).
90. J. J. Scarcelli, C. A. Hodge, C. N. Cole, The yeast integral membrane protein Apq12
potentially links membrane dynamics to assembly of nuclear pore complexes. *The Journal
of cell biology* **178**, 799-812 (2007).
91. B. M. Webster *et al.*, Chm7 and Heh1 collaborate to link nuclear pore complex quality
control with nuclear envelope sealing. *The EMBO journal* **35**, 2447-2467 (2016).
92. A. Aguzzi, T. O'Connor, Protein aggregation diseases: pathogenicity and therapeutic
perspectives. *Nature Reviews Drug Discovery* **9**, 237-248 (2010).
93. J. H. Butterfield, D. Weiler, G. Dewald, G. J. Gleich, Establishment of an immature mast
cell line from a patient with mast cell leukemia. *Leuk Res* **12**, 345-355 (1988).
94. D. Zabeo *et al.*, Exosomes purified from a single cell type have diverse morphology. *J
Extracell Vesicles* **6**, 1329476 (2017).
95. H. Xiao *et al.*, Mast cell exosomes promote lung adenocarcinoma cell proliferation – role
of KIT-stem cell factor signaling. *Cell Communication and Signaling* **12**, 64 (2014).
96. J. L. Höög, E. Gluenz, S. Vaughan, K. Gull, Ultrastructural investigation methods for
Trypanosoma brucei. *Methods Cell Biol* **96**, 175-196 (2010).
97. J. L. Höög *et al.*, Modes of flagellar assembly in Chlamydomonas reinhardtii and
Trypanosoma brucei. *eLife* **3**, e01479 (2014).
98. J. L. Höög *et al.*, 3D Architecture of the Trypanosoma brucei Flagella Connector, a Mobile
Transmembrane Junction. *PLOS Neglected Tropical Diseases* **10**, e0004312 (2016).
99. J. L. Höög *et al.*, Organization of Interphase Microtubules in Fission Yeast Analyzed by
Electron Tomography. *Developmental Cell* **12**, 349-361 (2007).
100. J. L. Höög, C. Antony, "Whole-Cell Investigation of Microtubule Cytoskeleton Architecture
by Electron Tomography" in Methods in Cell Biology. (Academic Press, 2007), vol. 79, pp.
145-167.

101.   E. Svensk *et al.*, Caenorhabditis elegans PAQR-2 and IGLR-2 Protect against Glucose
935   Toxicity by Modulating Membrane Lipid Composition. *PLoS genetics* **12**, e1005982 (2016).
936   102.   J. Sulston, J. Hodgkin, *Methods*, 1988 (1988).
937   103.   Sandra M. Hill *et al.*, Asymmetric Inheritance of Aggregated Proteins and Age Reset in
938   Yeast Are Regulated by Vac17-Dependent Vacuolar Functions. *Cell Reports* **16**, 826-838
939   (2016).
940   104.   W. K. Huh *et al.*, Global analysis of protein localization in budding yeast. *Nature* **425**, 686-
941   691 (2003).
942   105.   R. D. Gietz, Yeast transformation by the LiAc/SS carrier DNA/PEG method. *Methods in*
943   *molecular biology (Clifton, N.J.)* **1205**, 1-12 (2014).
944   106.   C. Janke *et al.*, A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent
945   proteins, more markers and promoter substitution cassettes. *Yeast* **21**, 947-962 (2004).
946   107.   P. Hawes, C. L. Netherton, M. Mueller, T. Wileman, P. Monaghan, Rapid freeze-
947   substitution preserves membranes in high-pressure frozen tissue culture cells. *J Microsc*
948   **226**, 182-189 (2007).
949   108.   J. L. Höög *et al.*, Electron tomography reveals a flared morphology on growing
950   microtubule ends. *Journal of Cell Science* **124**, 693-698 (2011).
951   109.   J. L. Höög, S. M. Huisman, D. Brunner, C. Antony, Electron Tomography Reveals Novel
952   Microtubule Lattice and Microtubule Organizing Centre Defects in +TIP Mutants. *PLOS*
953   *ONE* **8**, e61698 (2013).
954   110.   E. S. Reynolds, The use of lead citrate at high pH as an electron-opaque stain in electron
955   microscopy. *The Journal of cell biology* **17**, 208-212 (1963).
956   111.   D. N. Mastronarde, Automated electron microscope tomography using robust prediction
957   of specimen movements. *Journal of Structural Biology* **152**, 36-51 (2005).
958   112.   J. R. Kremer, D. N. Mastronarde, J. R. McIntosh, Computer Visualization of Three-
959   Dimensional Image Data Using IMOD. *Journal of Structural Biology* **116**, 71-76 (1996).
960   113.   K. Keuenhof *et al.*, Large Organellar Changes Occur during Mild Heat Shock in Yeast.
961   *bioRxiv* 10.1101/2021.01.25.428102, 2021.2001.2025.428102 (2021).
962   114.   A. Aufschnaiter *et al.*, The Enzymatic Core of the Parkinson's Disease-Associated Protein
963   LRRK2 Impairs Mitochondrial Biogenesis in Aging Yeast. *Frontiers in Molecular*
964   *Neuroscience* **11** (2018).
965   115.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat*
966   *Methods* **9**, 676-682 (2012).

967   **Figures and Tables**
968
969
970
971
Figure 1. NEB increases in frequency during heat shock in *S. cerevisiae*. (**a**) Percentage of

nuclei exhibiting NEB (left axis, black bars) and EDC (right axis, grey bars) as a function of time

after cells have been subjected to heat shock (38 ^{0}C, n=63-337). NEB peaked at a level of 10.3%

after 30 minutes and EDC exhibited a stable plateau of about 81% between 15 and 45 minutes.

(**b**) Thin section containing NEB and EDC in heat shocked *S. cerevisiae* cells. Black arrow

indicates NEB event, where the outer nuclear membrane clearly protrudes outwards and a

particle resides within the perinuclear space. (**c**) Labeling density of gold particles between

untreated and heat shocked cells (38 °C for 30 minutes) using an anti-GFP antibody. Both

Hsp104-GFP and guk1-7-GFP were preferentially localized at the EDC areas compared to the

rest of the nucleoplasm. (**d**) Labeling density of Hsp104 using an anti-Hsp104 antibody in wild

type and *hsp104Δ* cells subjected to heat shock at 38 °C for 45 minutes. No gold particles were

detected in the mutant strain showing the specificity of the antibodies. (**e**-**g**) Representative

images and respective models of the immuno-stained samples quantified in c and d. Hsp104 and

guk1-7 are significantly localized to the EDC area whereas only one gold is detected in the EDC

of the *hsp104Δ* mutant. Scale bars: 300 nm (**b**) and 500 nm (e, f, g). Abbreviations: NEB, nuclear

envelope budding; EDC, electron dense content; N, nucleus; V, vacuole; . *P<.05,

P<.01,*P<.001 vs. zero time point or control group; ns no significant differences between

groups.

Figure 2. Four different cellular stressors increase NEB frequency in *S. cerevisiae* **cells.**

Thin sections containing NEB events in (**a**) old cells (**b**) cells subjected to oxidative stress with

H_2O_2 (**c**) cells exposed to sodium arsenite . (**d**) Percentage of nuclei containing NEB events in

cells in each stress condition compared to control cultures grown under normal conditions.

Young cells were also examined after separation from old cells and show an increase in NEB

compared to the control, but to a lesser extent than old cells. Cells exposed to H_2O_2 were grown

in different media than the other cultures, and therefore were compared to a different control

culture . n is equal to the number of examined sections. (**e**) Thin section containing a NEB event

in cells treated with AZC, a chemical which causes proteotoxic stress. (**f**) Percentage of nuclei

containing NEB events before as well as 30 and 90 minutes after treatment with AZC . Cells

exhibited a higher percentage of NEB (22%) after 90 minutes of exposure to AZC than in any

other condition tested. All scale bars 200 nm. Abbreviations: NEB, nuclear envelope budding;

N, nucleus; black arrows indicate NEB events. *P<.05, **P<.01, ***P<.001 vs. control; ns no

significant differences between groups.

1005

Figure 3. Association of NEB pathway and protein quality control system.. (**a**) Immuno-gold

labeling with an anti-ubiquitin antibody (ab19247) was quantified as the amount of gold particles

per square micrometer for five different cellular structures. Lipid droplets (n=42) served as a

negative and autophagosomes (n=13) as a positive control. EDC (n=192) and NEB events

(n=12) exhibited an increase in labeling compared to lipid droplets and the nucleoplasm (n=42)

which was defined as area within the nucleus not containing EDC. (**b**) Thin sections labeled with

the anti-ubiquitin antibody ab19247. To the bottom of each micrograph a model is drawn to help

distinguish gold particles. Two labeled NEB events are shown in the first two panels and an EDC

containing gold particles is shown in the third panel . (**c-e**) Proteasomal inhibition triggered a

significant increase in NEB events with 73% of them having a distinct electron dense appearance.

(**f-g**) Immuno-gold labeling of an anti-GFP antibody of an Hsp104-GFP strain was quantified as

the amount of gold particles per square micrometer. Lipid droplets (negative control; n=41) and

EDC (positive control; n=50). NEB events (n=10) exhibited an increase labeling density in

comparison to the negative control. All scale bars 200 nm. Abbreviations: NEB, nuclear

envelope budding; N, nucleoplasm; EDC; electron dense content; LD, lipid droplet; APS,

autophagosomes; white arrow indicates NEB.

1022

Figure 4. NEB structures are not misassembled NPCs. Immuno-gold labeling, with the

Mab414 antibody which recognizes four different NPC proteins, was performed on thin sections

of cells grown under normal conditions (**a-c**) and old cells (**d**). To the right of each micrograph, a

model is drawn to help distinguish the locations of gold particles. (**e**) Percentage of labeled NPCs

(positive control), lipid droplets (negative control) and NEB events are plotted. Data for

unstressed cells is shown in black, and for old cells in grey. Lipid droplets and NEB events were

rarely labeled in contrast to the highly labeled NPCs. (**f**) 25 nm thick slice of NEB in a

tomographic reconstruction of *S. cerevisiae*. Both nuclear membranes are clearly visible

surrounding a vesicle within the perinuclear space. In all slices of the tomogram, the membrane

of this vesicle is complete and not continuous with either nuclear membrane. (**g**) 3D model of the

NEB event contained in **f**. Nuclear membranes are presented in orange color and the transporting

material in pink. n is equal to the number of examined structures. Scale bars: 200 nm (**a-d**), 100

nm (**f-g**). Abbreviations: NEB, nuclear envelope budding; N, nucleus; LD, lipid droplet; APS,

autophagosomes; NPC, nuclear pore complex; black arrows indicate NEB events, asterisks

indicate NPCs. *$P<.05$, ****$P<.001$ vs. NPCs (positive control).

1038

Figure 5. The NEB pathway is part of normal cellular function from humans to protists.

Thin sections containing nuclei were examined for the presence of NEB events in five different

organisms in wild type cells grown under normal conditions. (**a**) Percentage of nuclei containing

NEB events. NEB events were either outwards protruding (type 1), inwards protruding (type 2),

or did not protrude more in either direction but clearly contained a particle in the perinuclear

space (type 3). NEB was observed in (**b**) HMC-1 cells (n=122) (**c**) *C. elegans* (n=138) (**d**) *T.*

brucei (n=119) (**e**) *S. cerevisiae* (n=161) and (**f**) *S. pombe* (n=101) by electron microscopy. Scale

bars: 2 µm (**b**), 200 nm (**c, f**), 500 nm (**d**), and 300 nm (**e**). Abbreviations: NEB, nuclear envelope

budding; HMC-1, human mast cell line 1; N, nucleus; black arrows indicate NEB events.

1048

Figure 6. 3D architecture of NEB events of *S. pombe* and *T. brucei* visualized with electron

tomography. (**a**, **d**) 1.5 nm thick slices of *S. pombe* and *T. brucei* tomographical reconstructions

showing NEB events (black boxes). A second NEB event in *T. brucei,* not clearly visible in the

presented tomographic tilt, is located on the left lower part of the nucleus (dashed black box). (**b**,

e) 15 nm thick zoomed-in views of the NEB events in *S. pombe* and *T. brucei*. In *S. pombe* a lipid

bilayer is surrounding the transported material whereas in *T. brucei* the density is surrounded by

the nuclear membranes only. (**g**) Second NEB event of *T. brucei* (acquired at the tomographic tilt

that allowed best visualization of this NEB event). Electron dense particles are observed around

the neck of the bud (black arrow). (**c**, **f**, **h**) Reconstructed 3D models of the NEB events. Nuclear

membranes are presented in orange color and the transporting material in pink. Scale bars: 500

nm (**a**, **d**) and 100 nm (**b-c**, **e-h**). Abbreviations: NEB, nuclear envelope budding; N, nucleus;

1060

Figure 7. Phylogenetic tree of the species where NEB events have previously been

observed. The phylogenetic relationships between the species are presented as a horizontal

cladogram with the root to the left. Species indicated with blue color are from this study.

References presented from 1 to 13 respectively: (17), (18), (14), (16), (15), (29), (19, 26, 28),

(25), (21), (20), (23), (24), (22). All sketches are drawn by the author and are not subject to

copyright protection.

Figure 1

a Effect of heat shock in *S. cerevisiae*

Nuclei with budding events (%) / Nuclei with electron dense content (%)
EDC
NEB
Time (min)
n=337, n=76, n=63, n=72, n=68, n=70, n=81

b EDC, N

c n=
HSP104-GFP: 90, 12, 84, 72
guk1-7-GFP: 117, 26, 147, 144
Labeling density (# gold/µm2)
No HS, 30 min HS
Nucleoplasm
Nuclear EDC

d Labeling density (# gold/µm2)
Nucleoplasm
Nuclear EDC
WT, *hsp104*Δ
Heat shock

e HSP104-GFP: No HS, 30 min HS

f guk1-7-GFP: No HS, 30 min HS

g WT HS, *hsp104*Δ HS

Figure 2

Old *S. cerevisiae* cell (a)

H2O2 stress (b)

Arsenite stress (c)

AZC treated cells (e)

d Effect of cellular stress on NEB frequency

Nuclei with budding events (%)

***

*

*

n=200
n=204
n=249
n=337

Ctrl   As   Young   Old

*

n=204
n=197

Ctrl   H2O2

f Effect of AZC on NEB Frequency

Nuclei with budding events (%)

***

ns

n=337
n=114
n=100

Ctrl   AZC 30 min   AZC 90 min

Figure 3

a Immunogold labeling of Ubiquitin

Labeling density (# gold/µm2)

LD n=42, Aps n=13, Np n=42, EDC n=192, NEB n=12

b N, EDC, N, LD, EDC, N

c Proteasome inhibition

Nuclei with budding events (%)

DMSO n=200, MG132 n=200, WT n=337, rpn4Δ n=100

pdr5Δ

*   ***

d pdr5Δ

DMSO, MG132, WT, rpn4 Δ

anti-Ubiquitin

Ponceau S

e rpn4Δ

N, N

f Immunogold labeling of Hsp104-GFP

Labeling density (# gold/µm2)

LD n=41, EDC n=50, NEB n=10

g N

Figure 4

Unstressed cells (panels a, b, c)

Old cell (panel d)

e Localization of NPC Proteins

Gold labeled (%)

■ Unstressed
☐ Aged

NPCs: n=95 (Unstressed), n=100 (Aged)
LDs: n=82 (Unstressed), n=100 (Aged)
NEBs: n=6 (Unstressed), n=10 (Aged)

***
*

Electron Tomography (panels f, g)

Figure 5

Figure 7

Rattus norvegicus[1]

Mus musculus[2]

Cricetulus griseus[3]

Oryctolagus cuniculus[4]

Bos taurus[5]

Homo sapiens

Strongylocentrotus purpuratus[6]

Drosophila melanogaster[7]

Caenorhabditis elegans

Trypanosoma brucei

Saccharomyces cerevisiae

Schizosaccharomyces pombe

Podocarpus macrophyllus[8]

Spinulum annotinum[9]

Tradescantia bracteata[10]

Lilium longiflorum[11]

Pinus banksiana[12]

Noctiluca scintillans[13]