Nuclear envelope budding is an evolutionary conserved phenomenon

Dimitra Panagaki[1], Richard Neutze[1], Johanna L. Höög[1]

[1]Department of Chemistry and Molecular Biology, University of Gothenburg, Sweden

Keywords: Nuclear transport, electron tomography, budding, vesicles, exosomes

Running title: Electron Microscopy of Nuclear Budding

Corresponding author: johanna.hoog@gu.se

Abstract

Eukaryotic cells are defined by the compartmentalization of the cytoplasm into organelles, the largest of which is the nucleus, which contains the cellular DNA. Transport into and out of the nucleus is highly regulated and is traditionally thought to occur solely through nuclear pores. However, a small number of papers has repeatedly shown vesicular budding from the nuclear envelopes in different organisms. We used electron microscopy to identify such nuclear envelope budding events in a human cell line, *Caenorhabditis elegans* worms, the two yeasts *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* and the parasitic protist *Trypanosoma brucei*. Progressing to electron tomography, the finer details of the 3D architecture of such budding events was revealed. We summarize all the organisms in which this mode of translocation over the nuclear envelope has been observed and conclude that this may be a fundamental, evolutionary conserved mechanism of transport inside eukaryotic cells.

Introduction

The nucleus is the most prominent organelle of the eukaryotes, enclosing most of the cell's genetic material within their double lipid bilayer, the nuclear envelope. These two membranes were the evolutionary step between prokaryotes and eukaryotes that pose limitations for certain molecules to be in contact with the DNA. As the nuclear envelope is not permeable for most of the molecules inside the cells, special structures on their surfaces, called the nuclear pores, exists for highly selective translocation over the membranes (Watson, 1954). Small molecules can passively penetrate the nuclear pores from the cytoplasm to the nucleus and vice versa, while bigger molecules require an energy-dependent transportation mechanism to enter or leave the nucleus. Different types of proteins are required for guiding the larger sized molecules through

the pores, forming specific complexes for importing or exporting their cargoes (importins and

exportins respectively) (Chook et al., 2001; Hoelz et al., 2011).

Since the discovery of the nuclear pores in 1950, this transporting system has been accepted as

the only known way of communication between the cytoplasm and the nucleoplasm (Kohler et

al., 2007). However, in virus-infected cells, an alternative method of nuclear export has been

described. Herpes simplex virus, which are replicated in the nucleoplasm, are released into the

cytosol via outwards budding of the nuclear envelope (Schwartz et al., 1969; Hagen et al., 2015;

Fradkin et al., 2016; Wang et al., 2018). Since 1955, there have been many observations and

reports suggesting that these nuclear budding events probably occur in healthy cells as well and

different interpretations of their mechanism have been discussed (Gay, 1955; Hadek et al., 1962;

Afzelius, 1963; Elston et al., 1965; Longwell et al., 1965; D. Szollosi, 1965; Aldrich et al., 1970;

Dickinson et al., 1970; Gullvåg, 1970; Mepham et al., 1970; Dickinson, 1971; Hochstrasser et

al., 1987; M. S. Szollosi et al., 1988). These observations were made between a range of

organisms using mostly cells that are under differentiation (embryonic cells) and various

terminologies were given to describe the events (Table 1). The most recent reports were made in

healthy developing *Drosophila melanogaster* embryos and in the sea urchin gastrula (Hatch et

al., 2012; Speese et al., 2012; Jokhi et al., 2013; LaMassa et al., 2018).

This poses the question: is nuclear envelope budding an alternative way within eukaryotic cells

for the transport of material through the nuclear envelope specifically during viral infections and

in the development of some organisms, or is it instead a universal mechanism found in most/all

eukaryotes? In this study, we examine nuclear envelopes in a human cell line (*Homo sapiens*),

the worm (*Caenorhabditis elegans*), the budding yeast (*Saccharomyces cerevisiae*), the fission

yeast (*Schizosaccharomyces pombe*) and the protozoan African sleeping sickness parasite

(*Trypanosoma brucei*), to see if there is an evolutionary conserved alternative path of nuclear

import and export, besides the tightly regulated traffic over nuclear pores, in eukaryotes. We use

cells that are matured and grown under normal conditions to investigate if these events are only

present during cell differentiation or are independent of the organism's developmental cell stage.

Materials and Methods

Preparation of cell cultures and animals

Human mast cell line 1 (HMC-1; (Butterfield et al., 1988) cells were cultured in Iscove's

modified Dulbecco's medium (IMDM) and treated as described previously (Zabeo et al., 2017).

Procyclic *T. brucei* strain 427 was cultured in SDM-79 media with 20% fetal bovine serum.

Cultures were prepared and maintained at a concentration of 5×10^5 and 1×10^7 cells per

milliliter (Höög et al., 2010). Logarithmically growing wild type cells of the fission yeast *S.*

pombe were grown at 30^0C as described (Höög et al., 2007; Zabeo et al., 2017). The *C. elegans*

wild type reference strain was the Bristol variety, N2. The worms were cultured on normal

growing media plates (NGM plates) and the *E. coli* strain OP50 was used as a food source. The

worms were maintained at the optimal temperature of 25 ^{0}C (Sulston et al., 1988; Svensk et al.,

2016). Wild type cells of *S. cerevisiae* (S228C background) were cultured and maintained in

YPD media (Hill et al., 2016).

High-pressure freezing of cells and animals for electron microscopy

All samples used in this study has been prepared using high-pressure freezing followed by freeze

substitution. HMC-1 and *S. cerevisiae* cells were loaded into membrane carriers and were high-

pressure frozen in a Leica EM PACT1 (Leica microsystems, Wetzlar, Germany). *T. brucei* and *S.*

pombe were loaded into carriers and were high-pressure frozen in a Leica EM PACT2. *C.*

elegans samples were loaded into aluminium specimen carriers and were high-pressure frozen in

a Wohlwend Compact 3 (M. Wohlwend GmbH, Sennwald, Switzerland). Yeast paste from wild

type *S. cerevisiae* cells was used as a cryoprotectant. A short freeze substitution protocol was

applied, using 2% uranyl acetate (UA) for one hour (Hawes et al., 2007; Zabeo et al., 2017).

Samples were then embedded in K4M or HM20 resin.

All samples were sectioned in 70 nm thin sections and placed on either copper slot grids or mesh

grids. Sections were stained using 2% UA and Reynold's lead citrate (Reynolds, 1963).

Preparation of cells for electron tomography

For tomography, *S. pombe* cells were high-pressure frozen and fixed through freeze substitution

with anhydrous acetone containing 0.25% UA, 0.1% dehydrated glutaraldehyde and 0.01%

osmium tetroxide (OsO_4). Serial semi thick sections of about 210-250 nm (*S. pombe*) and 350

nm (*T. brucei*) thick sections were cut and samples were treated as described before (Höög et al.,

2007; Höög et al., 2014).

Image acquisition and electron tomography

All thin sections were imaged at 120 kV either on a LEO 912 OMEGA (Zeiss, Krefeldt,

Germany) equipped with a 2k x 2k VELETA Olympus CCD camera or on a Tecnai T12

transmission electron microscope equipped with a Ceta CMOS 16M camera (FEI Co.,

Eindhoven, The Netherlands). Tilt series of serial sections of *T. brucei* samples were acquired

using the serialEM software on a Tecnai TF30 300 kV microscope equipped with a Gatan

OneView 4K camera (FEI Co., The Netherlands). For the *S. pombe*, digital images were taken

every 1.5^o over a $\pm60^o$–65^o range operating a Tecnai TF20 electron microscope (FEI Co.,

Eindhoven, The Netherlands). The tomograms were acquired with the use of the serialEM

(Mastronarde, 2005). The tomographic reconstruction was performed with the use of the IMOD

software package (Kremer et al., 1996).

Phylogenetic tree

108   A phylogenetic tree was made showing all the organisms where nuclear budding events have

109   been observed. The tree was generated based on the NCBI taxonomy browser for scientific

110   names and visualized by the EvolView online software.

Results

Electron microscopy pictures revealed nuclear budding events in HMC-1 cells

To examine if nuclear budding occurs in human cells, we imaged 122 thin sections of human

mast cells (HMC-1) containing a nucleus using transmission electron microscopy. The HMC-1

nucleus has a diameter of ~8 μm. Therefore, the 122 thin sections observed here corresponds to

approximately one nuclear volume (8 μm/70 nm thick sections = 114 serial sections equals one

nucleus). Three morphologically different categories of budding events were observed (Figure

1A-C; n=122 sections). The most commonly observed morphology was outwards directed buds

(Figure 1A). In some cases, the electron dense material is clearly between the two membranes

with the outer leaflet expanding towards the cytoplasm. In other occasions, both the outer and

inner membrane leaflets are expanding towards the cytoplasm with material partially wrapped by

the nuclear envelope bud (Figure 1A). The second form of nuclear envelope extensions observed

was material found in between the nuclear membranes with no clear protrusion in either direction

(Figure 1B). Lastly, budding events with a general morphology of the nuclear envelope

expanding towards the nucleoplasm were also observed (Figure 1C). None of these nuclear

budding events are resembling nuclear envelope and ER contact sites (Figure S1). The budding

events were found in 11.5% of HMC-1 cells with some thin sections of nuclear envelopes

containing more than one bud. The frequency of the different bud morphologies is displayed in

Figure 1D. In order to investigate if nuclear budding events are evolutionary conserved inside

animal somatic cells we progressed to examine the nuclear envelope of the nematode *C. elegans*

in a similar manner.

Nuclear budding in the nematode *C. elegans*

Whole *C. elegans* specimens were high pressure frozen and sectioned for electron microscopy.

Images were taken of somatic cells containing a nucleus (n= 138). This number represents

approximately a total nuclear volume of five nuclei. 12.3% of these images contained a nuclear

envelope budding event. The majority of the nuclear budding events recorded in *C. elegans* have

the general morphology of the outwards protruding buds described for the HMC-1 cells (Figure

2A-C). Only one event fell in the third category of the inward nuclear envelope buds as

described before (Figure 2D). In these budding events, the material was visible between the

membranes, with one occasion where the material appeared to be surrounded by a membrane

(Figure 2C). Similar to the HMC-1 cells, some of the nuclear sections contained more than one

budding event. Having established that nuclear envelope budding is present in somatic animal

cells, we progressed to study species from other kingdoms of life.

Nuclear outwards budding events are also present in the fungi *S. cerevisiae*

The fungi *S. cerevisiae* and their nuclei were visualized in a total of 161 thin sections. This

number of sections corresponds to a total nuclear volume of five nuclei. All the five nuclear

budding events recorded in *S. cerevisiae* have the general morphology of the outwards

protruding buds as described for the HMC-1 cells (Figure 3A-C). No other categories were found

(Figure 3D). In these budding events, the electron dense material was not always visible but, in

some occasions, part of the nucleoplasm seemed to be surrounded by the bud (Figure 3C).

Schizosaccharomyces pombe, another well-studied organism from the Fungi kingdom but with

different properties than the *S. cerevisiae*, was further investigated.

Nuclear budding events in *S. pombe*

The fission yeast *S. pombe* was sectioned and 101 images were observed. This number

corresponds to a total nuclear volume of four nuclei. In these images, 7 nuclear budding events

were observed, of which most had the outwards budding morphology (Figures 4A, 4C) with one

of them being positioned in between the membranes (Figure 4B). Two of the events had a

strikingly bigger size compared to the rest and an extra membrane layer surrounded the electron

dense material (Figure 4B). The morphology of the budding events varied a lot, with some buds

having small sizes with not so obvious electron dense material and others being a lot bigger with

an extra membrane surrounding the material.

The nuclear budding events are conserved between mammals and protozoa

To investigate if nuclear budding events also occur in a unicellular organism of the protozoa

sub-kingdom, thin sections of *T. brucei* cells were prepared as described before and 119 nuclear

slices were observed. This number corresponds to a total nuclear volume of approximately four

nuclei. Nuclear budding events were also observed in these samples. All the budding events

found in the *T. brucei* nuclei had the outwards protruding morphology (Figure 5A-C). In the

majority of the visualized events, the electron dense material was easily observed and had a

distinct round morphology (Figure 5C). In order to investigate the 3D structure of these budding

events, electron tomography was performed on *T. brucei* as well as the yeast *S. pombe* samples,

another well-studied representative of the fungi kingdom.

Electron tomography shows the 3D architecture of nuclear budding events in *S. pombe* and

T. brucei

Using electron tomography, a better z-resolution can be achieved and the detailed membrane

morphology of the nuclear budding events could be revealed. Therefore, montaged dual-axis

electron tomograms of large volumes of high-pressure frozen fission yeast and *T. brucei* cells

were acquired, including approximately three and one complete nuclear volumes respectively

(Figure 6 A and D; n=1 *S. pombe* budding event; n=2 *T. brucei* budding events). In *S. pombe*,

one event of nuclear envelope budding was visualized. The membrane-enclosed vesicle had a

nucleoplasm similar texture and was located in between the two nuclear lipid bilayers (Figure

6B). The budding vesicle dimensions were 88 nm length by 149 nm width. The 3D model of this

nuclear envelope budding event shows that only the outer envelope is being extended around the

vesicle.

In electron tomograms of *T. brucei*, two events of nuclear budding were seen (Figure 6E, 6G).

The most progressed budding event had very distinct electron densities at the neck of the nuclear

envelope (Figure 6G, white arrow). The buds had dimensions of 78 nm length by 75 nm width

and 94 nm length by 69 nm width respectively. Similarly, only the outer nuclear envelope was

extended to surround a particle/vesicle surrounded by an electron dense material, the presence of

a membrane surrounding the transported material could not be ascertained (Figure 6 F, 6H).

Thus, we conclude that nuclear envelope budding was found in between 3 and 12% of the thin

sections observed in all species examined, spanning from human cell line to the protist *T. brucei*.

The nuclear envelope buds were similar in all cells but displayed a greater variability and larger

size in HMC-1 cells (Figure 1 and S2).

Discussion

We have shown that nuclear budding events can be seen in cells of all of the examined species,

stretching from animals, over fungi into protozoa (In total 540 thin sections have been observed).

The frequency of these events corresponds to approximately 2.7 times per nucleus. This shows

that the nuclear envelope budding is an evolutionary conserved feature of eukaryotes, even

though the structure of such budding events appears to change between species. For example, in

human cells, we saw three different kinds of nuclear envelope extensions, described as inwards

protruding, outwards protruding and particles in between the double lipid bilayers. The size of

these nuclear envelope buds had only small variations between the species with HMC-1 cells

having relatively bigger sized buds (Figure S2).

Even though not widely discussed as a means of nuclear transport, there is a trail through the

literature of these events occurring in animals, plants and protists. Figure 8 shows a phylogenetic

tree of all the species where nuclear budding events have been observed. Including the species in

our study, these events have now been reported in 18 different species across the eukaryota

domain. This is the first time all these observations from 1955 until now have been gathered and

presented as an evolutionary conserved process. The two fundamental questions are now: what

do they transport and where does it go? In a relatively recent study in *Drosophila melanogaster*

embryos, it has been shown that ribonucleoproteins (RNPs) are being exported through nuclear

envelope budding (Speese et al., 2012). That groundbreaking study strongly supports the

hypothesis that nuclear envelope budding facilitates the transport of genetic material from the

nucleus to the cytoplasm (Gay, 1955; Hadek et al., 1962; Afzelius, 1963; Longwell et al., 1965;

D. Szollosi, 1965; Aldrich et al., 1970; Gullvåg, 1970; Mepham et al., 1970; Hochstrasser et al.,

1987). Alternative theories argue that these budding events may be a membrane storage

mechanism that the cell will use in later stages of its development or that they play a role in the

formation of the endoplasmic reticulum (ER) and secretion granules (Gay, 1955; Dickinson et

al., 1970). An alternative, maybe more far-fetched, interpretation was that these budding vesicles

enclosing genetic material might be incipient mitochondria. That theory was based on the

increased number of mitochondria in the cytoplasm of pollen grains during the mitotic period

that these budding events were observed (Mepham et al., 1970). The exact role of these events is

not yet determined so none of the above theories can be ruled out. However, viruses are known

to high-jack the cells own functions to produce new virions (Davey et al., 2011). The fact that

viruses such as herpes simplex, Rous sarcoma and SV40 use this mechanism to exit the nucleus

(Epstein, 1962; Elston et al., 1965; Darlington et al., 1968; Hagen et al., 2015; Fradkin et al.,

2016; Roller et al., 2017), and its evolutionary conservation, are strong arguments for that this

mechanism of transport over the nuclear envelope has an important function inside eukaryotic

cells.

That leaves us with the question where are these vesicles transported? Some of our images shows

clearly that the vesicle formed by the budding event will have a double bilayer (Figure 2C, 4B,

6B). Yet, we do not commonly observe double membrane vesicles in the cytoplasm of these cells

(Höög et al., 2007), indicating that these vesicles could be either delivered fast to the plasma

membrane, fused with another compartment, demembranated in the cytoplasm, or that we have

failed to note their presence. A number of cryo-electron microscopy studies of exosomes, small

vesicles released from cells into the extracellular space, have shown a small fraction of the

population to have double and even triple bilayers (Poliakov et al., 2009; Yuana et al., 2013;

Höög et al., 2015; Zabeo et al., 2017). In particular, our previous study of HMC-1 cell derived

exosomes showed vesicles with double membrane bilayers 100-200 nm in size, similar to the

size of nuclear budding events measured in most of these cells (Zabeo et al., 2017). This suggests

that vesicles formed by nuclear envelope budding might be secreted. However, all previous

studies of nuclear envelope derived vesicles argue that the enclosed material should be released

into the cytoplasm at some stage but identification of the released material once in the cytoplasm

has not yet been possible. This obviously limits our ability to understand the function of these

budding events. An interesting suggestion on how they are transferred and released into the

cytoplasm is that of the connection with the endoplasmic reticulum (ER) (Hadek et al., 1962;

Gullvåg, 1970). The idea in this case was that once the buds are located between the two nuclear

membranes, they travel through the ER cisternae from where they are later released into the

cytoplasm as ER budding vesicles. As the ER is a continuation of the outer nuclear membrane

and well known for the vesicles it releases (Shibata et al., 2010; Phillips et al., 2015), this might

be a theory worth considering.

Many of the studies of nuclear envelope budding so far has been made on cells that were found

in developing organisms. Based on these studies, it was assumed that these budding events are

important during specific developmental stages of the cells (D. Szollosi, 1965; Dickinson, 1971;

M. S. Szollosi et al., 1988; LaMassa et al., 2018), none of these studies looked for events of

nuclear envelope budding in mature organisms. In our experiments, we examined adult worms

and cell culture cells under normal conditions. Observing frequent budding events under these

conditions implies that these nucleo-cytoplasmic interactions are also important after the

organisms are fully developed and might be part of normal cellular activity. This mode of

transport across the nuclear envelope may have different cargo and function depending on the

developmental stage of the cell. We conclude that this is a fascinating area of conserved cell

biology that has been largely ignored to date and we hope this paper encourage to new studies in

the field.

Acknowledgements

This project was funded by the Swedish Research Council Grand (2015-00560) and the Knut and

Alice Wallenberg Foundation (grand KAW 2012.0284, KAW 2014.0275) to Richard Neutze, the

Swedish Research Council Young Investigator Grant (2015-05427) and Knut and Alice

Wallenberg grant to Prof. Thomas Nyström and Johanna Höög.

All authors would like to thank Prof. Marc Pilon for providing the *C. elegans* worms and for

contributing intellectually to the manuscript writing. We would also like to thank Prof. Thomas

Nyström for providing the *S. cerevisiae* strain.

References

Afzelius, B. A. (1963). THE NUCLEUS OF NOCTILUCA SCINTILLANS: Aspects of Nucleocytoplasmic
Exchanges and the Formation of Nuclear Membrane. *J Cell Biol, 19*(1), 229-238.
Aldrich, H. C., & Vasil, I. K. (1970). Ultrastructure of the postmeiotic nuclear envelope in microspores of
Podocarpus macrophyllus. *J Ultrastruct Res, 32*(3), 307-315.
Butterfield, J. H., Weiler, D., Dewald, G., & Gleich, G. J. (1988). Establishment of an immature mast cell
line from a patient with mast cell leukemia. *Leuk Res, 12*(4), 345-355.
Chook, Y. M., & Blobel, G. (2001). Karyopherins and nuclear import. *Curr Opin Struct Biol, 11*(6), 703-715.
Darlington, R. W., & Moss, L. H., 3rd. (1968). Herpesvirus envelopment. *J Virol, 2*(1), 48-55.
Davey, N. E., Travé, G., & Gibson, T. J. (2011). How viruses hijack cell regulation. *Trends in Biochemical
Sciences, 36*(3), 159-169. doi:10.1016/j.tibs.2010.10.002
Dickinson, H. G. (1971). Nucleo-Cytoplasmic Interaction following Meiosis in the Young Microspores of
Lilium Longiflorum; Events at the Nuclear Envelope. *Grana, 11*(2), 117-127.
doi:10.1080/00173137109429867
Dickinson, H. G., & Bell, P. R. (1970). Nucleocytoplasmic interaction at the nuclear envelope in post
meiotic microspores of Pinus banksiana. *J Ultrastruct Res, 33*(3), 356-359.
Elston, R. N., & Stenkvist, B. (1965). Quantitative estimation of nuclear buds and micronuclei in bovine
cells transformed by Rous sarcoma and SV 40 viruses. *Zeitschrift für Zellforschung und
Mikroskopische Anatomie, 68*(4), 543-549. doi:10.1007/bf00347715

Epstein, M. A. (1962). Observations on the mode of release of herpes virus from infected HeLa cells. *J*
Cell Biol, 12, 589-597.
Fradkin, L. G., & Budnik, V. (2016). This bud's for you: mechanisms of cellular nucleocytoplasmic
trafficking via nuclear envelope budding. *Curr Opin Cell Biol, 41*, 125-131.
doi:10.1016/j.ceb.2016.05.001
Gay, H. (1955). NUCLEO-CYTOPLASMIC RELATIONS IN SALIVARY-GLAND CELLS OF Drosophila.
Proceedings of the National Academy of Sciences, 41(6), 370-375. doi:10.1073/pnas.41.6.370
Gullvåg, B. M. (1970). Release of Nuclear Material During the Development of Lycopodium Annotinum L.
Spores. *Grana, 10*(1), 31-34. doi:10.1080/00173137009429854
Hadek, R., & Swift, H. (1962). Nuclear extrusion and intracisternal inclusions in the rabbit blastocyst. *J*
Cell Biol, 13, 445-451.
Hagen, C., Dent, K. C., Zeev-Ben-Mordehai, T., Grange, M., Bosse, J. B., Whittle, C., Klupp, B. G., Siebert,
C. A., Vasishtan, D., Bauerlein, F. J., et al. (2015). Structural Basis of Vesicle Formation at the
Inner Nuclear Membrane. *Cell, 163*(7), 1692-1701. doi:10.1016/j.cell.2015.11.029
Hatch, E. M., & Hetzer, M. W. (2012). RNP export by nuclear envelope budding. *Cell, 149*(4), 733-735.
doi:10.1016/j.cell.2012.04.018
Hawes, P., Netherton, C. L., Mueller, M., Wileman, T., & Monaghan, P. (2007). Rapid freeze-substitution
preserves membranes in high-pressure frozen tissue culture cells. *J Microsc, 226*(Pt 2), 182-189.
doi:10.1111/j.1365-2818.2007.01767.x
Hill, Sandra M., Hao, X., Grönvall, J., Spikings-Nordby, S., Widlund, Per O., Amen, T., Jörhov, A., Josefson,
R., Kaganovich, D., Liu, B., et al. (2016). Asymmetric Inheritance of Aggregated Proteins and Age
Reset in Yeast Are Regulated by Vac17-Dependent Vacuolar Functions. *Cell Reports, 16*(3), 826-
838. doi:https://doi.org/10.1016/j.celrep.2016.06.016
Hochstrasser, M., & Sedat, J. W. (1987). Three-dimensional organization of Drosophila melanogaster
interphase nuclei. II. Chromosome spatial organization and gene regulation. *J Cell Biol, 104*(6),
1471-1483.
Hoelz, A., Debler, E. W., & Blobel, G. (2011). The Structure of the Nuclear Pore Complex. *Annual Review*
of Biochemistry, 80(1), 613-643. doi:10.1146/annurev-biochem-060109-151030
Höög, J. L., Gluenz, E., Vaughan, S., & Gull, K. (2010). Ultrastructural investigation methods for
Trypanosoma brucei. *Methods Cell Biol, 96*, 175-196. doi:10.1016/s0091-679x(10)96008-1
Höög, J. L., Lacomble, S., O'Toole, E. T., Hoenger, A., McIntosh, J. R., & Gull, K. (2014). Modes of flagellar
assembly in Chlamydomonas reinhardtii and Trypanosoma brucei. *Elife, 3*, e01479.
doi:10.7554/eLife.01479
Höög, J. L., & Lötvall, J. (2015). Diversity of extracellular vesicles in human ejaculates revealed by cryo-
electron microscopy. *J Extracell Vesicles, 4*, 28680. doi:10.3402/jev.v4.28680
Höög, J. L., Schwartz, C., Noon, A. T., O'Toole, E. T., Mastronarde, D. N., McIntosh, J. R., & Antony, C.
(2007). Organization of Interphase Microtubules in Fission Yeast Analyzed by Electron
Tomography. *Developmental Cell, 12*(3), 349-361. doi:10.1016/j.devcel.2007.01.020
Jokhi, V., Ashley, J., Nunnari, J., Noma, A., Ito, N., Wakabayashi-Ito, N., Moore, M. J., & Budnik, V. (2013).
Torsin mediates primary envelopment of large ribonucleoprotein granules at the nuclear
envelope. *Cell Rep, 3*(4), 988-995. doi:10.1016/j.celrep.2013.03.015
Kohler, A., & Hurt, E. (2007). Exporting RNA from the nucleus to the cytoplasm. *Nat Rev Mol Cell Biol,*
8(10), 761-773. doi:10.1038/nrm2255
Kremer, J. R., Mastronarde, D. N., & McIntosh, J. R. (1996). Computer Visualization of Three-Dimensional
Image Data Using IMOD. *Journal of Structural Biology, 116*(1), 71-76.
doi:https://doi.org/10.1006/jsbi.1996.0013
LaMassa, N., Arenas-Mena, C., & Phillips, G. R. (2018). Electron microscopic characterization of nuclear
egress in the sea urchin gastrula. *J Morphol, 279*(5), 609-615. doi:10.1002/jmor.20796

Longwell, A. C., & Yerganian, G. (1965). SOME OBSERVATIONS ON NUCLEAR BUDDING AND NUCLEAR
EXTRUSIONS IN A CHINESE HAMSTER CELL CULTURE. *J Natl Cancer Inst, 34*, 53-69.
Mastronarde, D. N. (2005). Automated electron microscope tomography using robust prediction of
specimen   movements.   *Journal   of   Structural   Biology,   152*(1),   36-51.
doi:https://doi.org/10.1016/j.jsb.2005.07.007
Mepham, R. H., & Lane, G. R. (1970). Observations on the fine structure of developing microspores
ofTradescantia bracteata. *Protoplasma, 70*(1), 1-20. doi:10.1007/bf01276839
Phillips, M. J., & Voeltz, G. K. (2015). Structure and function of ER membrane contact sites with other
organelles. *Nature Reviews Molecular Cell Biology, 17*, 69. doi:10.1038/nrm.2015.8
Poliakov, A., Spilman, M., Dokland, T., Amling, C. L., & Mobley, J. A. (2009). Structural heterogeneity and
protein composition of exosome-like vesicles (prostasomes) in human semen. *Prostate, 69*(2),
159-167. doi:10.1002/pros.20860
Reynolds, E. S. (1963). The use of lead citrate at high pH as an electron-opaque stain in electron
microscopy. *J Cell Biol, 17*, 208-212.
Roller, R. J., & Baines, J. D. (2017). Herpesvirus Nuclear Egress. *Adv Anat Embryol Cell Biol, 223*, 143-169.
doi:10.1007/978-3-319-53168-7_7
Schwartz, J., & Roizman, B. (1969). Concerning the egress of herpes simplex virus from infected cells:
electron and light microscope observations. *Virology, 38*(1), 42-49.
Shibata, Y., Shemesh, T., Prinz, W. A., Palazzo, A. F., Kozlov, M. M., & Rapoport, T. A. (2010). Mechanisms
determining   the   morphology   of   the   peripheral   ER.   *Cell,   143*(5),   774-788.
doi:10.1016/j.cell.2010.11.007
Speese, S. D., Ashley, J., Jokhi, V., Nunnari, J., Barria, R., Li, Y., Ataman, B., Koon, A., Chang, Y. T., Li, Q., et
al. (2012). Nuclear envelope budding enables large ribonucleoprotein particle export during
synaptic Wnt signaling. *Cell, 149*(4), 832-846. doi:10.1016/j.cell.2012.03.032
Sulston, J., & Hodgkin, J. (1988). *Methods*.
Svensk, E., Devkota, R., Stahlman, M., Ranji, P., Rauthan, M., Magnusson, F., Hammarsten, S., Johansson,
M., Boren, J., & Pilon, M. (2016). Caenorhabditis elegans PAQR-2 and IGLR-2 Protect against
Glucose Toxicity by Modulating Membrane Lipid Composition. *PLoS Genet, 12*(4), e1005982.
doi:10.1371/journal.pgen.1005982
Szollosi, D. (1965). Extrusion of nucleoli from pronuclei of the rat. *J Cell Biol, 25*(3), 545-562.
Szollosi, M. S., & Szollosi, D. (1988). 'Blebbing' of the nuclear envelope of mouse zygotes, early embryos
and hybrid cells. *J Cell Sci, 91 (Pt 2)*, 257-267.
Wang, I. H., Burckhardt, C. J., Yakimovich, A., & Greber, U. F. (2018). Imaging, Tracking and
Computational Analyses of Virus Entry and Egress with the Cytoskeleton. *Viruses, 10*(4).
doi:10.3390/v10040166
Watson, M. L. (1954). Pores in the mammalian nuclear membrane. *Biochim Biophys Acta, 15*(4), 475-
479.
Yuana, Y., Koning, R. I., Kuil, M. E., Rensen, P. C., Koster, A. J., Bertina, R. M., & Osanto, S. (2013). Cryo-
electron microscopy of extracellular vesicles in fresh plasma. *J Extracell Vesicles, 2*.
doi:10.3402/jev.v2i0.21494
Zabeo, D., Cvjetkovic, A., Lasser, C., Schorb, M., Lotvall, J., & Hoog, J. L. (2017). Exosomes purified from a
single   cell   type   have   diverse   morphology.   *J   Extracell   Vesicles,   6*(1),   1329476.
doi:10.1080/20013078.2017.1329476

HMC-1

Saccharomyces cerevisiae

Trypanosoma brucei

Schizosaccharomyces pombe

A

Nucleus

500 nm

B

100 nm

C

100 nm

Trypanosoma brucei

D

Nucleus

500 nm

E

100 nm

F

100 nm

G

100 nm

H

100 nm

Phylogenetic tree of species with observed nuclear budding events

Rattus norvegicus[1]

Mus musculus[2]

Cricetulus griseus[3]

Oryctolagus cuniculus[4]

Bos taurus[5]

Homo sapiens

Strongylocentrotus purpuratus[6]

Drosophila melanogaster[7]

Caenorhabditis elegans

Trypanosoma brucei

Saccharomyces cerevisiae

Schizosaccharomyces pombe

Podocarpus macrophyllus[8]

Spinulum annotinum[9]

Tradescantia bracteata[10]

Lilium longiflorum[11]

Pinus banksiana[12]

Noctiluca scintillans[13]