Chronic *Staphylococcus aureus* lung infection correlates with

proteogenomic and metabolic adaptations leading to an increased

intracellular persistence

Xin Tan, Mathieu Coureuil, Elodie Ramond, Daniel Euphrasie, Marion Dupuis, Fabiola Tros,

Julie Meyer, Ivan Nemanzny, Cerina Chhuon, Ida Chiara Guerrera, Agnes Ferroni, Isabelle

Sermet-Gaudelus, Xavier Nassif, Alain Charbit, Anne Jamet

Affiliations:

Xin Tan, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253, Institut

Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Mathieu Coureuil, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253,

Institut Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Elodie Ramond, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253, Institut

Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Daniel Euphrasie, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253,

Institut Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Marion Dupuis, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253, Institut

Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Fabiola Tros, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253, Institut

Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Julie Meyer, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253, Institut

Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Ivan Nemanzny, Plateforme Métabolomique Institut Necker-Enfants Malades, Structure

Fédérative de Recherche SFR Necker, University Paris Descartes, Paris, France

Cerina Chhuon, Plateforme Protéome Institut Necker-Enfants Malades, PPN, Structure

Fédérative de Recherche SFR Necker, University Paris Descartes, Paris, France

Ida Chiara Guerrera, Proteomics platform 3P5-Necker, Université Paris Descartes -

Structure Fédérative de Recherche Necker, INSERM US24/CNRS UMS3633, Paris, France

Agnes Ferroni, Laboratoire de Microbiologie de l'hopital Necker, University Paris

Descartes, Paris, France

Isabelle Sermet-Gaudelus, Université Paris Descartes, INSERM U1151 - CNRS UMR

8253, Institut Necker-Enfants Malades. Team: Canalopathies épithéliales : la mucoviscidose

et autres maladies, Paris, France

Xavier Nassif, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253, Institut

Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Alain Charbit, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253, Institut

Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Anne Jamet, Université Paris Descartes, INSERM U1151 - CNRS UMR 8253, Institut

Necker-Enfants Malades. Team: Pathogenesis of Systemic Infections, Paris, France

Keywords:

Cystic fibrosis; *Staphylococcus aureus*; intracellular persistence; biofilm; proteogenomics

Running title: Persistent infections in CF lungs

*Corresponding author: Anne Jamet

e-mail: anne.jamet@inserm.fr

Alternate corresponding author: Alain Charbit

alain.charbit@inserm.fr

Bâtiment Leriche. 14 Rue Maria Helena Vieira Da Silva CS 61431 - 75993 PARIS –

FRANCE

Tel: 33 1 – 72 60 65 11 — Fax: 33 1 - 72 60 65 13

40-word summary:

S. aureus persists for years in the lungs of patients with cystic fibrosis despite antibiotic

therapies. We demonstrate that *S. aureus* adaptation leads to increased intracellular

persistence suggesting a key role for intracellular niche during *S. aureus* chronic lung

infection.

Abstract (198 words)

Background: Chronic lung infection of cystic fibrosis (CF) patients by *Staphylococcus*

aureus is a well-established epidemiological fact. Indeed, *S. aureus* is the most commonly

identified pathogen in the lungs of CF patients. Strikingly the molecular mechanisms

underlying *S. aureus* persistency are not understood.

Methods: We selected pairs of sequential *S. aureus* isolates from 3 patients with CF and from

one patient with non-CF chronic lung disease. We used a combination of genomic, proteomic

and metabolomic approaches with functional assays for in-depth characterization of *S. aureus*

long-term persistence.

Results: For the first time, we show that late *S. aureus* isolates from CF patients have an

increased ability for intracellular survival in CFBE-F508del cells compared to ancestral early

isolates. Importantly, the increased ability to persist intracellularly was confirmed for *S.*

aureus isolates within the own patient F508del epithelial cells. An increased ability to form

biofilm was also demonstrated.

Furthermore, we identified the underlying genetic modifications inducing altered protein

expression profiles and notable metabolic changes. These modifications affect several

metabolic pathways and virulence regulators that could constitute therapeutic targets.

Conclusions: Our results strongly suggest that the intracellular environment might constitute

an important niche of persistence and relapse necessitating adapted antibiotic treatments.

Introduction

Staphylococcus aureus and *Pseudomonas aeruginosa* are the most common pathogens

infecting the lungs of patients with a chronic lung disease including cystic fibrosis (CF) [1, 2].

Furthermore, *S. aureus* is one of the earliest bacteria detected in infants with CF. However,

very few studies have addressed the adaptations undergone by *S. aureus* in this context [3, 4].

S. aureus has the ability to form biofilm [5-7] and to survive within a wide range of

eukaryotic host cells [8-17]. These abilities are likely to contribute to the persistence of *S.*

aureus in airways of patients with chronic lung diseases despite appropriate antimicrobial

treatments [18, 19]. *S. aureus* persistence is associated with a drastic decrease in metabolism

[20], a decrease in the expression of virulence factors and an increase in the expression of

bacterial adhesins [21]. Such profile is typical of small-colony variants (SCVs) that are

defined by small-sized colonies [15, 22, 23]. Beside SCVs, strains with normal colony

morphology can exhibit similar patterns of "low toxicity" which allow them to persist

intracellularly without being cleared by host cell defense mechanisms [21]. A "low toxicity"

pattern can be achieved either transiently, following changes in the expression of genes

encoding toxins and/or regulators, or permanently, by mutations in global regulators [24-26].

By studying serial isolates, we show that, during long-term lung infection, *S. aureus*

adaptation occurs through genomic modifications that accumulate over time and lead to major

metabolic modifications and protein expression changes. We also reveal that persistence of *S.*

aureus is associated with convergent evolution responsible for an increased ability to form

biofilm as well as to survive within host cells. These observations should be taken into

account in therapeutic decisions aiming at eradicating *S. aureus* chronic infections by

choosing drugs specifically targeting biofilm-embedded and intracellular bacteria.

Methods

Whole genome sequencing was performed on an Illumina MiSeq instrument (2x150

bp) and the sequences were processed using the Nullarbor bioinformatic pipeline software

v1.20 and RAST server. The sequences reported in this paper are available at NCBI's

BioProject   database   under   accession   number   PRJNA446073

(http://www.ncbi.nlm.nih.gov/bioproject/446073).

Quantification of biofilm formation was assessed with crystal violet staining in

polystyrene 96-well plates. Cystic Fibrosis Bronchial Epithelial cell line CFBE41o- and

primary nasal epithelial cell were infected with a multiplicity of infection of 100 using an

inoculum taken from cultures of *S. aureus* grown in Brain Heart Infusion until exponential

growth phase. Infected cells were kept for 6 days in a medium containing 50 μg/mL

gentamycin to kill extracellular bacteria. For proteomics, proteins were digested and analyzed

by liquid chromatography coupled with tandem mass spectrometry (nanoLC-MS/MS). For

metabolomics, metabolite profiling of *S. aureus* isolates was performed by liquid

chromatography–mass spectrometry (LC-MS). The mass spectrometry proteomics data have

been deposited to the ProteomeXchange Consortium via the PRIDE [27] partner repository

with the dataset identifier PXD011281.

A full description of methods is available in Supplementary Methods.

Statistical analysis

Data were analyzed using R or GraphPad Prism softwares. Results are presented either

with one representative experiment for clarity or as means ± standard deviation (SD). The

number of biological and technical replicates is indicated per figure.

For two-sample comparisons statistical significance was measured using unpaired two-tail

Student's t-test or Wilcoxon rank sum test as indicated in the figure legends. For comparisons

between more than two groups, statistical significance was measured using one-way analysis

of variance (ANOVA) with multiple comparisons (Dunnett's correction) performed, with

each value compared to that of the reference strain.

P values of <0.05 were considered to indicate statistical significance.

Ethics statement

All experiments were performed in accordance with the guidelines and regulations

described by the Declaration of Helsinki and the low Huriet-Serusclat on human research

ethics and informed consent was obtained for all participating subjects. Serial isolates of *S.*

aureus were obtained from airway secretions from four patients with chronic lung infection at

the Necker-Enfants Malades University Hospital, Paris, France. Sputum sampling is part of

routine standard care. The research procedure is validated by Ile de France 2 IRB (ID-

RCB/Eudract: 2016 A00309-42).

Results

Selection of *S. aureus* sequential isolates from patients with chronic lung infection.

Three patients with CF (CF1, CF2 and CF3) and, for comparison purpose, one patient with

non-CF chronic lung disease (CLD) were chosen. For each patient we selected one early and

one late isolate separated by 3 to 9 years intervals. Whole-genome sequencing confirmed that

each pair of isolates belonged to four distinct clones (**Figure 1**). Patient diseases and

treatments are detailed in supplementary Methods.

***S. aureus* clinical isolates from CF patients evolved an increased persistence ability**

within CFBE-F508del epithelial cell line.

Numerous studies have shown that *S. aureus* has the ability to survive within human cells [8-

17]. We subsequently aimed at investigating if during the course of within-lung adaptation, *S.*

aureus isolates have evolved a greater ability to persist within epithelial cells. We infected

bronchial CFBE epithelial cell line (F508del +/+ CFTR mutation) with clinical isolates, the

control strain USA300-LAC and a stable SCV mutant altered in the haemin biosynthetic pathway (hereafter named Δ*hem*). As expected, wild-type bacteria were not able to persist whereas the Δ*hem* mutant was able to persist intracellularly during the whole course of the experiment (**Figure S1**) [13]. All early and late clinical isolates were able to persist at least 2.6-fold, and up to 900-fold, more than the USA300-LAC reference strain at day 3 and 6 post-infection (**Figure 2AB**). Furthermore, at day 3 and 6 post-infection, all the late isolates recovered from CF patients exhibited an improved ability to persist intracellularly within CFBE-F508del epithelial cells compared to cognate early isolates (**Figure 2A**). Interestingly, the CLD_late isolate recovered from the non-CF patient did not exhibit an improved ability to persist within CFBE epithelial cells compared to CLD_early isolate (**Figure 2B**). These data suggest that *S. aureus* adaptation within CF-lungs correlates with an improved ability to persist intracellularly in cells with a CFTR dysfunction.

Late *S. aureus* isolate of CF3 patient exhibits an increased persistence within primary F508del epithelial own patient cells.

To confirm the relevance of the results obtained with bronchial CFBE epithelial cell line, we first assessed the persistency of CF3 isolates within primary epithelial cells isolated from the nose of a healthy donor (**Figure 2C**). In addition, we performed an infection assay with the CF3 primary epithelial own patient cells (F508del +/+ CFTR mutation) to verify the specific within patient-adaptation of *S. aureus* recovered from long-term infection (**Figure 2D**). These experiments confirmed that the late isolate persistence ability is improved compared to early isolate at day 3 and 6 within both primary nasal epithelial cells retrieved from a healthy donor and from the CF3 patient.

***S. aureus* clinical isolates from chronically infected patients evolved high biofilm formation ability.**

Assuming that isolates retrieved from chronic infections might have a high biofilm-forming

capacity, we studied the biofilm formation ability of the pairs of isolates. Remarkably, all

isolates displayed a greater capacity to form biofilms compared to that of the weak biofilm-

producer USA300-LAC reference strain (p-value of <0.001, **Figure S2**). Furthermore, for

three chronically infected patients, the late isolates formed more biofilm than the early

isolates, revealing that long-term adaptation within lungs had improved their biofilm

formation capacity (p-value of <0.001, **Figure 3**).

Late *S. aureus* clinical isolates from chronically infected patients acquired auxotrophies.

Compared to that of USA300-LAC, all patients early isolates and the CF1 late isolate display

similar colony morphology on brain heart infusion (BHI) agar plates and a wild-type growth

in a liquid broth mimicking sputum (Cystic Fibrosis Sputum Medium or CFSM) **(Figure S3**

and Figure 4A). In contrast, the late isolate of CF2 and CLD patients displayed a typical

SCV phenotype with very small colonies on BHI agar (**Figure S4**) and CF2_late, CF3_late

and CLD_late isolates exhibited a growth defect in CFSM broth **(Figure 4BCD)**. Thymidine-

dependent SCVs are frequently isolated from patients treated with sulfamethoxazol (SXT)

[28]. Indeed, supplementation with thymidine restored almost wild-type growth for CF2_late

and CLD_late isolates (**Figure 4BD**). We used genomic data to determine the auxotrophy of

CF3_late isolate and identified a frameshift in *panB* gene, which is involved in de novo

biosynthesis of pantothenic acid (**Table 1**). Accordingly, growth of CF3_late isolate in the

presence of pantothenate restored wild-type growth (**Figure 4C**). Thus, isolates from three out

of four patients with *S. aureus* chronic lung infection acquired auxotrophy during the course

of the disease.

Late *S. aureus* clinical isolates from chronically infected patients acquired antibiotic

resistance.

Genome analysis evidenced mutations in *thyA*, *gyrB,* and *rpsJ* genes, which were associated

with SXT, fluoroquinolones and cyclines resistance, respectively (**Table 1**). Thus, isolates

from three out of four patients with *S. aureus* chronic lung infection acquired antibiotic

resistance consistent with the administration of the corresponding drugs during the course of

the disease.

208
Genomic, proteomic and metabolomic modifications associated with *S. aureus*

adaptation during chronic lung infection.

In order to investigate the underlying genomic, proteomic and metabolic modifications

associated with the observed phenotypic changes we compared genomes, proteomes and

metabolomes of late compared to early isolates. The differences in proteomic and metabolic

profiles between early and late isolates of patients are highlighted by heatmaps shown in

Figure S5.

Genomes of all clinical isolates were de novo assembled and coding DNA sequences (CDSs)

were annotated. Most of the SNPs were missense variants occurring in CDSs (**Table 2**).

Nonsynonymous mutations acquired by late isolates were found mainly in genes involved in

metabolic processes (**Figure 5A**) and more specifically in "amino acid transport and

metabolism" and "carbohydrate transport and metabolism" functional categories. In addition,

the largest category of proteins to be differentially expressed for all pairs also comprised

proteins related to metabolism processes (and more specifically to the "amino acid transport

and metabolism" category) (**Figure 5B**). Concordant with genomic and proteomic results, the

category "amino acids" was the most altered metabolites category in the late isolate of all

patients compared to their cognate early isolates (**Figure 5C**).

Many regulatory proteins were differentially expressed. Indeed, proteins of the Agr, Rot, Sae,

Sar or Fur regulatory networks were differently expressed in all late isolates. In CF1_late

isolate, the *agrC* and the *saeR* genes mutations **(Table 1)** had a pleiotropic effect on the

proteome (down-regulation of delta hemolysin and PSMb1 and upregulation of proteins

encoded by *spa, sbi, fnbA, rot* and *coa* genes). In addition, adhesins encoded by *sasG, efb,*

sdrD and *ecb* were up-regulated. In CF2_late isolate, the *agr* regulon is also downregulated

suggesting an evolution toward low virulent and highly adhesive properties. The metabolite

profiling of CF2_late isolate, showed a decrease in ADP, which is well correlated with the

lack of ThiM (hydroxyethylthiazole kinase) expression found in proteomic analysis due to a

frameshift in *thiM* gene **(Table 1)**. In CF3_late isolate, frameshifts in *fakA* and *panB* genes

(Table 1) were associated with a lack of cognate proteins expression in CF3_late isolate. In

addition, adhesins encoded by *sdrD* and *sasF* were upregulated. Interestingly, an over-

production of penicillin-binding protein 2 encoded by *mecA* is correlated with the *saeR*

mutation **(Table 1)** [29]. The metabolite profiling of CF3_late isolate, revealed a drastic

diminution of pantothenate, coenzyme A and dephospho-coenzyme A, which is in line with

the lack of expression of PanB and PanC proteins [30].

Of note, the non-CF control clone displays a different evolution trajectory. Indeed, AgrA and

AgrC were up-regulated in CLD_late isolate, suggesting that it has retained virulent

properties.

Altogether, the proteogenomic data suggest that all the late isolates recovered from CF

patients (but not CLD patient) have evolved toward highly adhesive and low virulent

properties. Besides, metabolic profiling suggests that all late isolates have evolved a reduced

citric acid cycle activity compared to cognate early isolates.

Discussion

Our study showed that during chronic lung infection, *S. aureus* adapts through the

acquisition of common adaptive traits including antibiotic resistances, auxotrophies, reduced

citric acid cycle activity, increased biofilm and intracellular persistence abilities that occurred

irrespective of the clone type.

Of particular interest, we report mutations in two master regulatory systems, Agr and Sae,

likely to impact multiple proteins expression and metabolites amounts.

agr-defective mutants, such as CF1 late isolate, have been shown to arise during chronic

infections and are better adapted to persistence within the infected host [25, 31, 32].

Genetic alterations directly or indirectly targeting SaeR regulon were identified in the 3 CF

patients. Since SaeR is involved in the regulation of over 20 virulence factor genes [33] and

SaeRS-deficient bacteria are less infective in animal models [34], it is likely that SaeRS is a

key factor in long-term colonization. In the three CF late isolates, we observed an increased in

the expression of the SdrD adhesin belonging to the SaeR regulon and involved in adhesion to

human nasal epithelial cells and to human keratinocytes [35][36]. Our results suggest that

SdrD is also important for long-term lung colonization.

In patients with chronic lung infections, SCVs detection is most often the consequence of a

long-term SXT treatment [37]. Mutations in the *thyA* gene, as found in CF2 and CLD late

isolates, lead to stable clinical SCVs that are no longer susceptible to SXT and are thymidine-

auxotrophic (TA-SCV) [28, 37]. Since thymidine is assumed to be abundant during lung

inflammation, TA-SCVs can still grow in this environment.

In CF3 late isolate, we observed a pantothenate auxotrophy, which has been previously

associated with persistency in *Mycobacterium tuberculosis* [38]. The acquisition of

pantothenate auxotrophy suggests that pantothenate could also be present in CF lungs. Thus,

our data confirm that metabolic specialization is a common phenomenon among long-term

colonizers [39].

Other striking traits of phenotypic convergent evolution of *S. aureus* identified in this work

were the increased ability to form biofilm and to persist in the intracellular niche. For CF2

and CF3 patients, the increased biofilm ability of late isolates could be linked to a mutation in

the *fakA* gene, encoding fatty acid kinase A (FakA). Indeed, several studies showed that

FakA-null strains were proficient in biofilm formation [6] and deficient in the expression of

virulence factors controlled by the SaeRS system [40]. Overexpression of adhesins detected in

proteomic analysis could also ultimately lead to increase biofilm formation in clinical isolates.

Numerous studies have demonstrated *S. aureus* ability to persist within host cells [8-17].

Strikingly, for the three CF patients, the *S. aureus* late isolates showed a greater ability to

persist within CFBE-F508del epithelial cells compared to the early ones at day 3 and 6 post-

infection. Of note, the late isolate of CLD patient did not present an improved ability to

persist intracellularly within CFBE-F508del epithelial cells possibly due the fact that it has

adapted to a non-CF patient.

Our multi-omics approach allowed both confirmation of previously known mechanisms

and identification of novel candidate genes and pathways involved in the persistence ability of

clinical isolates. We now provide evidence that the *saeR/fakA* regulon and the pantothenate

pathway could also be promising therapeutic targets to fight persistent *S. aureus* infections.

Our study suggests that the use of antibiotic with a good intracellular penetration should be

the best therapeutic option in order to eradicate *S. aureus* from chronically infected lungs.

Funding:

This work was supported by Institut national de la Santé et de la Recherche Médicale; Centre

National de la Recherche Scientifique; and Université Paris Descartes Paris Cité Sorbonne. A

scholarship from the China Scholarship Council [n° CSC NO. 201508500097] has been

provided to XT.

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Acknowledgments:

We thank Aurélie Hatton, Charlotte Roy and Zhicheng Zhou for their help with primary nasal

epithelial cells culture.

Author contributions:

XT, MC, XN, AC and AJ conceived and designed the study. XT, ER, MD, DE, FT, JM and AJ

made the experiments and analysis. IN, CC, ICG, AF and ISG contributed with data and

analysis. AJ, AC and XT wrote the manuscript, with contributions and comments from all

authors.

No conflicts exist for the authors

     Tables

318   **Table 1. Mutations linked to phenotypic changes in clinical isolates**

Isolate	Gene	Product	Mutation	Associated phenotype
CF1_late	*agrC*	Accessory gene regulator protein C	Premature stop codon (L193X)	Down-regulation of toxins and up-regulation of adhesins
CF1_late	*saeR*	Staphylococcus exoprotein expression protein R	Missense G179L	Up-regulation of adhesins
CF2_late	*thyA*	Thymidylate synthase	Premature stop codon (W88X)	SXT resistance and thymidine auxotrophy
CF2_late	*rpsJ*	30S ribosomal protein S10	Missense K57M	Cyclines resistance
CF2_late	*thiM*	Hydroxyethylthiazole kinase	Frameshift	Decrease in ADP
CF2_late	*fakA*	Fatty acid kinase A	Missense G187D	Increased biofilm formation and intracellular persistency
CF3_late	*panB*	3-methyl-2-oxobutanoate hydroxymethyltransferase	Frameshift	Pantothenate auxotrophy
CF3_late	*fakA*	Fatty acid kinase A	Frameshift	Increased biofilm formation and intracellular persistency
CF3_late	*gyrB*	DNA topoisomerase subunit B	Missense F226S	Fluoroquinolones resistance
CF3_late	*saeR*	Staphylococcus exoprotein expression protein R	Missense A190T	Over-production of penicillin-binding protein 2
CLD_late	*thyA*	Thymidylate synthase	Missense P48R	SXT resistance and thymidine auxotrophy

323 **Table 2. General features of detected mutations**

	CF1_late	CF2_late	CF3_late	CLD_late
Time since early isolate	2.8 years	6.7 years	9 years	4.4 years
Total polymorphisms	21	34	79	30
SNP[a]	19	25	73	26
INDEL[b]	2	9	6	4
CDS[c]	18	21	62	23
NON-SYN[d] (%)	14 (77.8)	16 (47.1)	51 (64.6)	17 (56.7)
FR[e]	2	4	4	2
MS[f]	10	11	47	14
STOP gained[g]	1	1	0	1
Other	1	0	0	0
SYN[h]	4	5	11	6
IG[i]	3	13	17	7

324 [a]SNP, single nucleotide polymorphism; [b]INDEL, insertion-deletion; [c]CDS, coding sequence;

325 [d]NON-SYN, nonsynonymous mutation; [e]FR, frameshift variant; [f]MS, missense variant;

326 [g]STOP gained, premature stop codon; [h]SYN, synonymous mutation; [i]IG, intergenic

328
329

Figures

Tree scale: 1000 ⊢——⊣

Fig 1. **Selection of four pairs of *S. aureus* isolates belonging to four different STs in four**

patients.

Dendrogram generated by wgsa.net from the genomes of the eight clinical isolates retrieved

from the respiratory samples of three patients with CF, one patient with non-CF chronic lung

infection and four reference genomes from public databases. Branch length is proportional to

the number of variant nucleotide sites within the core genes. For each patient, the isolate

taken first is called "early" while the isolate taken later is named "late". The dates of sampling

and the sequence type (ST) of the isolates are indicated. "TA" and "PA" mean that the isolate

is auxotrophic for thymidine or pantothenate respectively. The name of the isolate is indicated

in red when it is resistant to methicillin (MRSA).

Reference strains included are PFESA1902 (ERR554197), st1307 (ERR158691), PFESA1195

(ERR554722).

Fig 2. Intracellular persistence of *S. aureus* clinical isolates in CFBE-F508del epithelial cell line and within primary patient cells. A and B) Bronchial CFBE epithelial cell line (CFBE-F508del homozygous for the F508del-CFTR mutation) was infected with the control strain USA300-LAC and clinical isolates from CF patients (A) or CLD patient (B). **C and D)** Primary nasal epithelial cells retrieved from a healthy donor ("Primary WT") (C) and from the CF3 patient ("Primary CF3" with F508del +/+ CFTR mutation) (D) were infected with the control strain USA300-LAC and CF3 isolates.

For all experiments, gentamicin was present throughout the experiment to prevent

extracellular bacterial growth and new infection. Bacterial loads inside cells were evaluated

by CFU enumeration at 3 and 6 days after infection. Results are normalized with USA300-

LAC strain as a reference and expressed as a fold change of CFUs.

Results have been obtained from two independent experiments performed in triplicate for

ABC and one experiment for D. Statistical analysis was performed by Wilcoxon rank sum test

*P < 0.05; **P<0.01; ns P>0.05.

Fig 3. Quantification of biofilm formation of *S. aureus* clinical isolates. Biofilm formation

quantification was performed using the crystal violet microtiter assay in BHI medium with

1% glucose. Results shown are the mean ±SD for three independent experiments performed in

triplicate. Statistical significance was measured using a two-tail Student's t-test when biofilm

production of a late isolate was compared with biofilm production of cognate early isolate

from the same patient. ** indicates p-value of <0.001 whereas ns indicates p-values>0.05.

Fig 4. Growth of late *S. aureus* clinical isolates in CFSM. Growth curves were carried out in medium mimicking the respiratory fluid of cystic fibrosis patients (Cystic Fibrosis Sputum Medium or CFSM), with or without the addition of thymidine or pantothenate. The results shown correspond to a representative experiment. The orange and green curves correspond to bacterial growth in media supplemented with either thymidine or pantothenate; the red and blue curves, to bacterial growth in medium without thymidine or pantothenate. WT, USA300-LAC.

Fig 5. Proteogenomic and metabolomic analysis of the four pairs of *S. aureus* isolates. A)
Vertical histograms show the functional classification of proteins encoded by genes with
nonsynonymous mutations in the genomes of late isolates of *S. aureus* compared to early
isolates. B) Horizontal histograms show the functional classification of differentially
expressed annotated proteins in late compared to early isolates of each patient. For each
category, histograms represent the number of down- and up-regulated proteins from
proteomic analysis using the threshold of <2 and >2, respectively. Only genes and proteins
with functional annotation available are included. The "cellular processes and signaling"
category encompasses regulatory proteins and proteins involved in cell wall and capsule
synthesis. The "information storage and processing" category encompasses proteins involved

in replication, translation and repair processes. The "metabolism" category encompasses

proteins involved in metabolism and transport. The "virulence" category encompasses

exotoxins, proteins involved in adhesion, biofilm formation and immunomodulation. **C)**

Categorization in 8 categories of altered amount of metabolites in late compared to early

isolates. Metabolites were detected by carrying out 2 independent experiments performed in

triplicate. TCA, Tricarboxylic acid cycle; PPP, Pentose phosphate pathway; Nucl,

Nucleotides; CoA, Coenzyme A; AA, Amino acids.

References

1.   Wijers CD, Chmiel JF, Gaston BM. Bacterial infections in patients with primary ciliary
dyskinesia: Comparison with cystic fibrosis. Chronic respiratory disease **2017**; 14(4): 392-406.

2.   Jesenak M, Banovcin P, Jesenakova B, Babusikova E. Pulmonary manifestations of
primary immunodeficiency disorders in children. Frontiers in pediatrics **2014**; 2: 77.

3.   McAdam PR, Holmes A, Templeton KE, Fitzgerald JR. Adaptive evolution of
Staphylococcus aureus during chronic endobronchial infection of a cystic fibrosis patient. PloS one
2011; 6(9): e24301.

4.   Lopez-Collazo E, Jurado T, de Dios Caballero J, et al. In vivo attenuation and genetic
evolution of a ST247-SCCmecI MRSA clone after 13 years of pathogenic bronchopulmonary
colonization in a patient with cystic fibrosis: implications of the innate immune response. Mucosal
immunology **2015**; 8(2): 362-71.

5.   Zapotoczna M, O'Neill E, O'Gara JP. Untangling the Diverse and Redundant
Mechanisms of Staphylococcus aureus Biofilm Formation. PLoS pathogens **2016**; 12(7): e1005671.

6.   Sabirova JS, Hernalsteens JP, De Backer S, et al. Fatty acid kinase A is an important
determinant of biofilm formation in Staphylococcus aureus USA300. BMC genomics **2015**; 16: 861.

7.   Schwartbeck B, Birtel J, Treffon J, et al. Dynamic in vivo mutations within the ica
operon during persistence of Staphylococcus aureus in the airways of cystic fibrosis patients. PLoS
pathogens **2016**; 12(11): e1006024.

8.   Garzoni C, Francois P, Huyghe A, et al. A global view of Staphylococcus aureus
whole genome expression upon internalization in human epithelial cells. BMC genomics **2007**; 8: 171.

9.   Garzoni C, Kelley WL. Staphylococcus aureus: new evidence for intracellular
persistence. Trends in microbiology **2009**; 17(2): 59-65.

10.   Kalinka J, Hachmeister M, Geraci J, et al. Staphylococcus aureus isolates from
chronic osteomyelitis are characterized by high host cell invasion and intracellular adaptation, but still
induce inflammation. International journal of medical microbiology : IJMM **2014**; 304(8): 1038-49.

11.   Mitchell G, Grondin G, Bilodeau G, Cantin AM, Malouin F. Infection of polarized airway
epithelial cells by normal and small-colony variant strains of Staphylococcus aureus is increased in
cells with abnormal cystic fibrosis transmembrane conductance regulator function and is influenced by
NF-kappaB. Infection and immunity **2011**; 79(9): 3541-51.

12.   Proctor RA, von Eiff C, Kahl BC, et al. Small colony variants: a pathogenic form of
bacteria that facilitates persistent and recurrent infections. Nature reviews Microbiology **2006**; 4(4):
295-305.

13.   Rollin G, Tan X, Tros F, et al. Intracellular Survival of Staphylococcus aureus in
Endothelial Cells: A Matter of Growth or Persistence. Frontiers in microbiology **2017**; 8: 1354.

14.   Sendi P, Proctor RA. Staphylococcus aureus as an intracellular pathogen: the role of
small colony variants. Trends in microbiology **2009**; 17(2): 54-8.

15.   Tuchscherr L, Heitmann V, Hussain M, et al. Staphylococcus aureus small-colony
variants are adapted phenotypes for intracellular persistence. The Journal of infectious diseases **2010**;
202(7): 1031-40.

16.   Tuchscherr L, Medina E, Hussain M, et al. Staphylococcus aureus phenotype
switching: an effective bacterial strategy to escape host immune response and establish a chronic
infection. EMBO molecular medicine **2011**; 3(3): 129-41.

17.   von Eiff C, Becker K, Metze D, et al. Intracellular persistence of Staphylococcus
aureus small-colony variants within keratinocytes: a cause for antibiotic treatment failure in a patient
with darier's disease. Clinical infectious diseases : an official publication of the Infectious Diseases
Society of America **2001**; 32(11): 1643-7.

18.   Branger C, Gardye C, Lambert-Zechovsky N. Persistence of Staphylococcus aureus
strains among cystic fibrosis patients over extended periods of time. Journal of medical microbiology
1996; 45(4): 294-301.

19.   Kahl BC, Duebbers A, Lubritz G, et al. Population dynamics of persistent
Staphylococcus aureus isolated from the airways of cystic fibrosis patients during a 6-year prospective
study. Journal of clinical microbiology **2003**; 41(9): 4424-7.

20.   Kriegeskorte A, Grubmuller S, Huber C, et al. Staphylococcus aureus small colony
variants show common metabolic features in central metabolism irrespective of the underlying
auxotrophism. Frontiers in cellular and infection microbiology **2014**; 4: 141.

21.   Garcia-Betancur JC, Goni-Moreno A, Horger T, et al. Cell differentiation defines acute
and chronic infection cell types in Staphylococcus aureus. eLife **2017**; 6.

22.   Tuchscherr L, Kreis CA, Hoerr V, et al. Staphylococcus aureus develops increased
resistance to antibiotics by forming dynamic small colony variants during chronic osteomyelitis. The
Journal of antimicrobial chemotherapy **2016**; 71(2): 438-48.

23.   Sadowska B, Bonar A, von Eiff C, et al. Characteristics of Staphylococcus aureus,
isolated from airways of cystic fibrosis patients, and their small colony variants. FEMS immunology
and medical microbiology **2002**; 32(3): 191-7.

24.   Das S, Lindemann C, Young BC, et al. Natural mutations in a Staphylococcus aureus
virulence regulator attenuate cytotoxicity but permit bacteremia and abscess formation. Proceedings
of the National Academy of Sciences of the United States of America **2016**; 113(22): E3101-10.

25.   Suligoy CM, Lattar SM, Noto Llana M, et al. Mutation of Agr Is Associated with the
Adaptation of Staphylococcus aureus to the Host during Chronic Osteomyelitis. Frontiers in cellular
and infection microbiology **2018**; 8: 18.

26.   Tuchscherr L, Loffler B. Staphylococcus aureus dynamically adapts global regulators
and virulence factor expression in the course from acute to chronic infection. Current genetics **2016**;
62(1): 15-7.

27.   Deutsch EW, Csordas A, Sun Z, et al. The ProteomeXchange consortium in 2017:
supporting the cultural change in proteomics public data deposition. Nucleic acids research **2017**;
45(D1): D1100-D6.

28.   Chatterjee I, Kriegeskorte A, Fischer A, et al. In vivo mutations of thymidylate
synthase (encoded by thyA) are responsible for thymidine dependency in clinical small-colony variants
of Staphylococcus aureus. Journal of bacteriology **2008**; 190(3): 834-42.

29.   Nygaard TK, Pallister KB, Ruzevich P, Griffith S, Vuong C, Voyich JM. SaeR binds a
consensus sequence within virulence gene promoters to advance USA300 pathogenesis. The Journal
of infectious diseases **2010**; 201(2): 241-54.

30.   Spry C, Kirk K, Saliba KJ. Coenzyme A biosynthesis: an antimicrobial drug target.
FEMS microbiology reviews **2008**; 32(1): 56-106.

31.   Painter KL, Krishna A, Wigneshweraraj S, Edwards AM. What role does the quorum-
sensing accessory gene regulator system play during Staphylococcus aureus bacteremia? Trends in
microbiology **2014**; 22(12): 676-85.

32.   Shopsin B, Eaton C, Wasserman GA, et al. Mutations in agr do not persist in natural
populations of methicillin-resistant Staphylococcus aureus. The Journal of infectious diseases **2010**;
202(10): 1593-9.

33.   Liu Q, Yeo WS, Bae T. The SaeRS Two-Component System of Staphylococcus
aureus. Genes **2016**; 7(10).

34.   Montgomery CP, Boyle-Vavra S, Daum RS. Importance of the global regulators Agr
and SaeRS in the pathogenesis of CA-MRSA USA300 infection. PloS one **2010**; 5(12): e15177.

35.   Askarian F, Ajayi C, Hanssen AM, et al. The interaction between Staphylococcus
aureus SdrD and desmoglein 1 is important for adhesion to host cells. Scientific reports **2016**; 6:
22134.

36.   Cassat JE, Hammer ND, Campbell JP, et al. A secreted bacterial protease tailors the
Staphylococcus aureus virulence repertoire to modulate bone remodeling during osteomyelitis. Cell
host & microbe **2013**; 13(6): 759-72.

37.   Kriegeskorte A, Lore NI, Bragonzi A, et al. Thymidine-Dependent Staphylococcus
aureus Small-Colony Variants Are Induced by Trimethoprim-Sulfamethoxazole (SXT) and Have
Increased Fitness during SXT Challenge. Antimicrobial agents and chemotherapy **2015**; 59(12): 7265-
72.

38.   Sambandamurthy VK, Wang X, Chen B, et al. A pantothenate auxotroph of
Mycobacterium tuberculosis is highly attenuated and protects mice against tuberculosis. Nature
medicine **2002**; 8(10): 1171-4.

39.   La Rosa R, Johansen HK, Molin S. Convergent Metabolic Specialization through
Distinct Evolutionary Paths in Pseudomonas aeruginosa. mBio **2018**; 9(2).

40.   Ericson ME, Subramanian C, Frank MW, Rock CO. Role of Fatty Acid Kinase in
Cellular Lipid Homeostasis and SaeRS-Dependent Virulence Factor Expression in Staphylococcus
aureus. mBio **2017**; 8(4).

Supplementary figure

Figure S1. Intracellular growth curves of reference *S. aureus* isolates in CFBE

epithelial cell line. Bronchial CFBE epithelial cell line (CFBE-F508del homozygous

for the F508del-CFTR mutation) was infected with the control strain USA300-LAC

(WT) and a stable SCV mutant altered in the haemin pathway (Δ*hem*). Gentamicin

was present throughout the experiment to prevent extracellular bacterial growth and

new infection. Bacterial load inside cells were evaluated by CFU enumeration at 3

and 6 days after infection. Results shown are the mean ±SD for four experiments

performed in triplicate.

Figure S2. Quantification of biofilm formation of *S. aureus* clinical isolates.
Biofilm formation was monitored using the crystal violet microtiter assay in BHI
medium with 1% glucose. USA300-LAC strain (WT) was used as a reference for
weak biofilm production whereas its *agrC* derivative obtained from the Nebraska
Transposon Mutant Library was used as a reference for strong biofilm production.
Results shown are the mean ±SD for three independent experiments performed in
triplicate. Statistical significance was measured using one-way ANOVA with multiple
comparisons (Dunnett's correction) performed on the dataset as a whole, with each
value compared to the WT. ** indicates p-value of <0.001.

Figure S3. Growth curves of early *S. aureus* clinical isolates in CFSM. Growth

curves were carried out in CFSM. The results shown correspond to a representative

experiment. WT, USA300-LAC.

543 **Figure S4. Colony morphology of USA300-LAC (WT), CF2_late and CLD_late**

544 **isolates.** Bacteria were grown on BHI agar incubated 12h at 37°C.

Figure S5. Heatmaps showing comparison of protein and metabolite profiles of early/late *S. aureus* isolates. The isolates were cultured to the stationary phase in medium mimicking the respiratory fluid of cystic fibrosis patients (CFSM) with the addition of thymidine. **A)** Heatmap visualization and hierarchical clustering analysis of the proteomic profiling in the late isolate compared to the early isolate of each patient. One experiment with three biological replicates was performed for each isolate. Rows: proteins; columns: samples; color key indicates protein relative concentration value (yellow: lowest; blue: highest). **B)** Heatmap visualization and hierarchical clustering analysis of the metabolite profiling in the late isolate compared to the early isolate of each patient. The top 50 most changing compounds are presented. Two independent experiments with three biological replicates were performed for each isolate. Rows: metabolites; columns: samples; color key indicates metabolite relative concentration value (blue: lowest; red: highest).