ATM is activated by ATP depletion and modulates mitochondrial function through NRF1

(ATM regulates ATP levels through NRF1)

Hei-Man CHOW[1,2]

Aifang CHENG[1]

Xuan SONG[1]

Mavis R. SWERDEL[3]

Ronald P. HART[3]

Karl HERRUP[1,4]

[1] Division of Life Science, Hong Kong University of Science and Technology
[2] Institute for Advanced Study, Hong Kong University of Science and Technology
[3] Department of Cell Biology and Neuroscience, Rutgers University
[4] Lead contact, to who correspondence should be addressed

Summary

ATM deficiency depletes ATP levels in neurons with high energy demands. Oxidative stress resulting from neuronal activity normally activates ATM, which triggers NRF1 nuclear translocation and the upregulation of mitochondrial gene expression, restoring the neuron's energy supply

Abstract

A newly uncovered role for ATM protein deepens our understanding of the neurological symptoms of Ataxia-telangiectasia (A-T). With high energy demands, neurons of the brain, particularly Purkinje cells, require dynamic maintenance of ATP levels. ATM is the gene mutated in A-T and its deficiency compromises the activity of both the TCA cycle and the electron transport chain (ETC). We show that ATP insufficiency activates ATM via reactive oxygen species. While resting ATP levels are normal, $Atm^{-/-}$ cells have a reduced capacity to respond to surges in energy demands. This blunted response results from impaired ATM-mediated activation of nuclear respiratory factor-1 (NRF1). ATM phosphorylation of NRF1 at T259 dimerizes, translocates to the nucleus and drives mitochondrial gene expression, allowing increased mitochondrial protein synthesis. Our findings thus identify ATM not only as a guardian of genomic integrity bu also of mitochondrial output, suggesting that alternate fuel sources may ameliorate or delay ataxic symptoms in A-T.

Introduction

Mitochondrial diseases commonly involve neurological symptoms, and ataxia resulting from cerebellar atrophy and Purkinje cell loss is the most frequent of these (Bargiela et al., 2015). In one cohort study, 225 out of 345 patients (65%) afflicted with a range of different mitochondrial diseases showed symptoms consistent with ataxia (Bargiela et al., 2015; Lax et al., 2012). The reverse relationship is also found (Bargiela et al., 2015); one-fifth of persons showing symptoms of definitive ataxia also present with features of mitochondrial dysfunction. Thus, ataxia is linked to mitochondrial defects and vice versa (Fang et al., 2014; Scheibye-Knudsen et al., 2013). This bi-directional correlation led us to consider the protein involved in the inherited ataxia known as ataxia-telangiectasia (A-T), a debilitating autosomal recessive multi-system disease caused by a mutation of the *ATM* gene (Watters, 2003). The protein product of the *ATM* gene was originally identified as a large PI3K-kinase family member that functions as a DNA damage response protein. Yet while the prevalence of many A-T symptoms such as cancer or immunodeficiency can vary, pronounced ataxia, associated with a profound loss of cerebellar Purkinje cells, afflicts all patients with A-T. This is intriguing since, while various mechanisms have been proposed to explain the cerebellar focus of A-T neuropathology, the links between the loss of ATM function and the selective susceptibility of cerebellar neurons to neurodegeneration remain largely unknown.

ATP regulation is critical for a nerve cell. A typical resting neuron contains a billion ATP molecules, yet the firing of only a single action potential is estimated to require the hydrolysis of 10-100 million ATPs to fully restore the resting membrane potential (Howarth et al., 2012; Howarth et al., 2010). This type of estimate underscores the dynamic nature of ATP supply in neurons and raises questions as to how the levels of such a critical molecule are regulated. Computational modeling predicts that each cerebellar Purkinje cell, with its large cellular size and numerous synaptic connections (Carter and Bean, 2009; Herzig and Shaw, 2018; Sugimori and Llinas, 1990; Welsh et al., 2002), consumes 62 times more energy than a nearby cerebellar granule cell. The majority of this energy is used for the production of action potentials (52%) and the regulation of postsynaptic receptors (38%) with the remainder going towards

maintaining membrane potential at rest and other minor tasks (Howarth et al., 2012; Howarth et al., 2010). Thus neuronal health and survival is heavily dependent on the constant availability of adequate supplies of ATP. Most ATP is derived from glucose through glycolysis (Bratic and Trifunovic, 2010) although the catabolism of other carbon sources may contribute to the process. The predominant site of ATP production is the mitochondrion, where the tricarboxylic acid (TCA) cycle and oxidative phosphorylation (OXPHOS) reactions of the electron transport chain (ETC) provide the pathways for its genesis (Hall et al., 2012). The TCA cycle in the mitochondrial matrix produces reduced nicotinamide adenine dinucleotide (NADH) and flavin adenine dinucleotide (FADH$_2$). These in turn are used as electron donors by the ETC located in the inner mitochondrial membrane (Owen et al., 2002). The five complexes of the ETC are built from the protein products of literally hundreds of genes, most of which are encoded by the nuclear genome (Dimauro and Rustin, 2009). The highly deleterious effects of mutations in these genes demonstrate that even minor structural changes in ETC proteins disrupt electron transport, ATP production, and can thus cause a range of conditions recognized as mitochondrial diseases that usually have profound impacts on brain functioning.

We report here that a previously unrecognized relationship exists between ATM and the regulation of ATP production in the neuronal mitochondrion. ATM-deficiency results in compromised activities of the TCA cycle and electron transport chain, leading to a reduced capacity to respond to increases in ATP demand. This newly discovered activity of ATM is mediated through nuclear respiratory factor-1 (NRF1). We propose that in the absence of ATM, neurons, in particular mature cerebellar Purkinje cells, cannot respond adequately to the increased in energy demands of neuronal activity. The resulting ATP deficit leads to their degeneration and the observed ataxia and other neurological deficits of A-T.

Results

ATM-related deficits in the respiratory chain and TCA cycle

As predicted from the observed correlation between mitochondrial disease and cerebellar ataxia (Bargiela et al., 2015; Lax et al., 2012), the symptoms of A-T cluster with those typically found in diseases involving the mitochondrion (Fang et al., 2014; Scheibye-Knudsen et al., 2013). To confirm this in an unbiased manner, we used the MITODB web application to screen all reported A-T clinical symptoms—neurological as well as peripheral— for association with mitochondrial function. We found that peripheral symptoms, such as immune deficiency, radiosensitivity, and telangiectasia, failed to show any meaningful mitochondrial association, but CNS phenotypes such as cerebellar atrophy and ataxia showed a strong overlap (Red bars, Fig.1A and Table S1A). This suggests a connection between ATM and mitochondrial function that is most prominent in the nervous system. This is reminiscent of findings from several laboratories, identifying links between mitochondria and A-T (Fang et al., 2016; Patel et al., 2011; Valentin-Vega et al., 2012) and suggested to us that A-T neurodegeneration might be related more specifically to defects in OXPHOS and electron transport chain function. A final hint as to the focus of this connection comes from the observation that of 8 of the 9 features of mitochondrial syndromes that are shared by A-T are associated with defects in the ETC (Fig.1B).

With this in mind, we re-analyzed earlier microarray results (Li et al., 2013) from human A-T and control cerebellar cortex focusing on genes associated with OXPHOS and ETC functions. Of the more than 31,000 transcripts analyzed, 23% showed significant changes in A-T (Fig.2A, Table S1B). Three-quarters were down-regulated; the rest were up-regulated (Fig.2A, Table S1C). The altered transcripts fell most prominently into 30 Gene Ontology (GO) groups (p<0.0001) as identified by the GATHER platform (Fig.2B, Table S1D) (Chang and Nevins, 2006). Enrichment clustering based on the total number of genes associated with a GO term (Fig.2C) identified several primary metabolic pathways. Using the Gene Set Enrichment Analysis (GSEA), we further identified 73 KEGG pathways with nominal p-value<0.01 (Table S1E). Pathways associated with OXPHOS (Fig.2D) and the TCA cycle (Fig.2E) were significantly depleted in A-T

(adjusted $p<0.05$). Together, these results suggest that mitochondrial physiology in the cerebellum is significantly altered in the absence of ATM.

Mitochondria are not autonomous entities; they rely heavily on >1,000 imported nuclear-encoded gene products (Lopez et al., 2000). We therefore use the above dataset to perform sub-analyses on 1,158 genes encoding mitochondria-localized proteins as indicated in the MitoCarta 2.0 inventory (Calvo et al., 2016). We observed a distinct pattern of altered expression in these mitochondrial genes in the A-T cerebellar samples (Fig.2F, Table S1F). Expression of 458 (41%) nuclear-encoded, mitochondrial genes was significantly altered in A-T. Among these, 83% had reduced expression ($p<0.05$). The processes that would be affected by the reduced gene expression included virtually every aspect of mitochondrial function – OXPHOS as well as mitochondrial ribosomes, nucleotide levels, mitochondrial DNA repair and replication, and mitochondrial-specific transcription and translation (Fig.2G, Table S1G). This gene regulatory pattern predicts a generalized loss of nuclear-encoded mitochondrial proteins in A-T and suggests that the normal function of ATM is crucial to sustain their presence.

Reduced expression and function of metabolic enzymes is associated with lack of ATM

Given the conserved nature of most mitochondrial proteins, we sought to validate the findings in $Atm^{-/-}$ mouse. Biogenesis of a functional OXPHOS system requires not only the core structural proteins of Complex I-V, it also requires an array of OXPHOS assembly factors, as well as proteins needed for coenzyme Q10, cytochrome-c and Fe-S-complex biogenesis (Koopman et al., 2013) (Fig.3A, Table S1H). In the $Atm^{-/-}$ mouse cerebellum, as in the A-T human samples (Fig. S1A), expression of genes encoding proteins constituting the OXPHOS core subunits (16/21, $p<0.05$–Fig.3B, S1B), OXPHOS assembly factors (17/21, $p<0.05$–Fig.3C, S1C), OXPHOS coenzymes and cofactor assembly factors (9/11, $p<0.05$–Fig.3D, S1D), mitochondrial ribosomal proteins (8/8, $p<0.05$–Fig.3E, S1E); and factors regulating mitochondrial DNA repair, replication, transcription and translation (5/7, $p<0.05$–Fig.3F, S1F) were significantly reduced. A notable exception to this downward trend was the increased expression of genes for proteins involved in mitochondrial DNA repair (3/3, $p<0.05$) and maintenance of the mitochondrial dNTP pool

(4/5, p<0.05) (Fig.3F, S1F; Table S1H). The latter two groups suggest the cells are mounting a sustained response to enhanced mitochondrial DNA damage. To determine how loss of ATM affected the function of each individual complex, we compared the activities of Complexes I–IV in wild type and $Atm^{-/-}$ cultured cortical neurons. All four showed significantly reduced activity (Fig.3G-J). The F_1F_0-ATP synthase (Complex V) trended lower, but the difference was not significant (Fig.3K). This indicates that in the absence of ATM, the functions of the four key ETC complexes are reduced. Importantly, total mtDNA content was not changed in $Atm^{-/-}$ neurons (Fig.S1G), implying that the effects seen in the mitochondria are secondary to a defect in nuclear gene expression. Together, the clinical symptoms, the patterns of gene expression and the reduced functions of the ETC complexes all point to the conclusion that the A-T neurological phenotype has deep roots in energy metabolism.

Altered metabolic dependence of ATM-deficient cells

Previous studies indicated that TCA cycle metabolites are altered in the cerebellum of $Atm^{-/-}$ mice (Fang et al., 2016). We therefore queried our previously published human cerebellar microarray data (Fig.4A) and found that the expression of hexokinase-1/2 (*HK1/2*), the first rate-limiting enzyme of glycolysis, was significantly reduced in A-T. This suggested that glycolysis might be diminished, and indeed the levels of pyruvate, the end product of glycolysis, were significantly reduced in cultured A-T human fibroblasts (Fig.4B, vehicle condition). We next treated wild type human fibroblasts with 2-deoxyglucose (2-DG), a hexokinase inhibitor (Chen and Gueron, 1992). This caused pyruvate levels in unaffected cells to drop to those seen in A-T fibroblasts (Fig.4B). Pyruvate can either enter mitochondria and be converted to acetyl-CoA by pyruvate dehydrogenase (*PDHA1*) or remain in the cytoplasm where it is converted to lactate by lactate dehydrogenase (*LDH*) (McCommis and Finck, 2015). Since the gene expression array data showed that *LDH* was increased and *PDHA1* was dramatically reduced in $Atm^{-/-}$ cells (Fig.4A), it appears that pyruvate is both reduced and steered toward the production of lactate in the cytoplasm rather than acetyl CoA in the mitochondrion. Further supporting this hypothesis, UK5099, an inhibitor of the mitochondrial pyruvate carrier [MPC (Hildyard et al., 2005)], resulted in a roughly 50% elevation of pyruvate levels in wild type cells, but no

significant change in A-T cells (Fig.4B). This suggests that in ATM-deficient cells pyruvate is converted to lactate in the cytoplasm and thus is unavailable inside mitochondria. Consistent with this idea, treatment with the LDH inhibitor oxamate (Novoa et al., 1959) elevated pyruvate levels in A-T cells while having no significant effect on wild type cells (Fig.4B). Since lactate levels were upregulated by about 3-fold in A-T cells (Fig.4C, vehicle), pCMBS treatment that inhibits lactate efflux increased lactate inside A-T cells, but had no effect on wild type cells. By contrast, blocking lactate production with oxamate returned lactate in A-T cells to normal levels with no difference observed in wild type cells. Mouse primary neurons isolated from $Atm^{-/-}$ knockout mice (Fig.S2A) and wild type neurons silenced by Atm-specific siRNA (Fig.S2B) both gave similar observations. We next followed the dynamic changes of intracellular pyruvate and lactate in Atm knockout neurons over a 24-hour period using FRET nanosensors—Pyronic (PYRuvate Optical Nano Indicator from CECs; San Martin et al., 2014) (Fig.S2C, S2E) and Laconic (LACtate Optical Nano Indicator from CECs; San Martin et al., 2013) (Fig.S2D, S2F). Both sensors were consistent with the above results. Thus, in A-T cells mitochondrial pyruvate is lost while cytosolic lactate is increased.

Once pyruvate enters the mitochondrion it is converted to acetyl-CoA by pyruvate dehydrogenase (PDHA1) before entering the TCA cycle. This conversion would appear to be blocked in A-T cells as there is a significant reduction in PDHA1 (Fig.4A). After this, genes for subsequent the first steps of the TCA cycle prior to the formation of succinate remained unchanged, but for the two TCA cycle steps downstream from succinate – succinate dehydrogenase (SDH) and fumarase (FH) – were significantly reduced (Fig.4A). Consistent with these array results, colorimetric assays revealed that the levels of mitochondrial isocitrate were reduced by more than two-thirds (Fig.4D) while a near doubling of mitochondrial succinate levels was found in A-T cells (Fig.4E). This situation in A-T cells would seem to be energetically precarious. We therefore considered alternative carbon sources like fatty acid and ketone body catabolism. This can generate acyl-CoA, which enters the mitochondria where it is converted to acetyl-CoA by carnitine palmitoyltransferase (CPT). Unfortunately, the mitochondrial isoform of CPT $(CPT2)$ is reduced by two-thirds in A-T cells (Fig.4A). Earlier findings from our lab showed

that dietary supplements of the amino acid glutamine to $Atm^{-/-}$ mice alleviated their symptoms and extended their lifespan (Chen et al., 2016a). As the mitochondrial *GLS* isoform, *GLS2*, was increased 5-fold in A-T cells (Fig.4A) suggesting that A-T cells use glutamine as their source of carbon for the TCA cycle. We confirmed this suggestion using the Mito Fuel Flex test. While glucose was the major fuel for wild type cells A-T cells showed a major dependence on glutamine as an alternative fuel (Fig.4E).

ATM-deficient cells have an impaired response to enhanced energy demand

We next investigated how the observed abnormalities affected overall mitochondrial function in ATM-deficient cells. We first used tetramethylrhodamine methylester (TMRM) fluorescence to assess mitochondrial membrane potential ($\Delta\psi$m). In $Atm^{-/-}$ neurons, steady state TMRM fluorescence was 1.4-fold higher than in wild type controls (Fig.5A). In examining the cells response to stress, however, an opposite relationship was observed. Oligomycin, an inhibitor of the F_1F_0-ATP synthase, caused a rapid reduction in $\Delta\psi$m in $Atm^{-/-}$ cells but a slow sustained increase in wild type, presumably because in oligomycin protons were no longer used to power the ATP-synthase (Fig.5B). The results were similar in human A-T fibroblasts (Fig.S3A-B). The literature suggests that this response reflects a significant problem facing the $Atm^{-/-}$ cells. Because of the importance of the proton gradient, in cells with impaired OXPHOS Complex V runs in reverse – cleaving ATP to act as a proton pump and sustain $\Delta\psi$m (Abramov et al., 2010; McKenzie et al., 2007). The above data suggest that the impaired mitochondrial status ATM-deficient cells makes them less flexible in adapting to changes in energy demand. To validate these interpretations, we measured the dynamic changes in oxygen consumption rate (OCR) – an indicator of mitochondrial respiration. We found that the basal OCR in mouse $Atm^{-/-}$ neurons (Fig.5D) and human A-T fibroblasts (Fig.S3D) were not significantly different from controls. Treatment with oligomycin which determines oxygen consumed to drive ATP synthesis also indicated that ATP-linked OCR was nearly identical in A-T and control cells (Fig.5D, Fig.S3D). By convention, these data indicate that under resting conditions, respiration activity of both wild type and ATM-deficient cell are similar. However, FCCP stimulus, which drives maximal OCR, revealed that ATM-deficient cells had a reduced response compared to wild type. This implies

that the reserve respiratory capacity (RRC) in ATM deficient cells was significantly smaller than in controls (Fig.5D, S3D; dashed lines). This reduced RRC, together with our data which showed that ATM deficient cells had to hydrolyze ATP to maintain their $\Delta\psi m$, potentially mean less ATP is available in response to surges in energy demand. Under equilibrium conditions, we found that, compared to controls, the levels of ATP were reduced by one quarter in mouse $Atm^{-/-}$ cortical neurons (Fig.5E) and by nearly one half in A-T human fibroblasts (Fig.S3E). We synaptically activated the cultured neurons for 30 seconds with glutamate to produce large fluxes of sodium and potassium and thus stimulate the hydrolysis of ATP by the Na^+/K^+-pump. We then monitored the ATP:ADP ratio in real time using the PercevalHR biosensor (Tantama et al., 2013). In wild type cultures, this treatment resulted in a rapid 50% reduction in the ATP:ADP ratio but over the next 30 minutes the ratio returned to pre-stimulus levels (Fig.5F-G). In $Atm^{-/-}$ cultures, while the initial reduction in the ATP:ADP ratio was nearly the same as in controls, the rate of recovery was significantly slower (Fig.5F-G). Indeed, while wild type cells could recover fully after glutamate, the treatment killed nearly a quarter of the $Atm^{-/-}$ cells (Fig.5H). These data show that cells lacking ATM are unable to respond to increased energy demands and are thus more vulnerable during periods of high energy demand.

Nuclear respiratory factor-1 links ATM activity and mitochondrial function

To determine how ATM deficiency led to the observed defects in ATP generation, we returned to the global gene expression data from human cerebellum (Fig.2A). We analyzed all possible transcript isoforms of the genes that were differentially expressed in A-T and control samples (12,931 transcripts from 7,222 genes). ATM itself is a serine/threonine kinase with no known DNA binding capacity; if it were to modulate gene expression it would most likely work through modulating activities of transcription factors. We therefore analyzed the promoter regions (-450 to +50 bp) of all differentially-regulated transcripts for transcription factor binding sequences using the PSCAN tool (Zambelli et al., 2009). This analysis identified 71 candidate transcription factors that were significantly associated with the differentially expressed genes (p<0.05, Z-test; Table S1J). Of these 71, binding motifs for NRF1 ranked highest (p=8.24 x 10^{-114}; Fig.6A); of the 12,931 transcripts, binding sequences with >85% similarity to the NRF1

consensus motif were found in about one-third of them (4,624 promoters belonging to 3,360 genes, Fig.6B-C). Thus, NRF1 emerged as a strong candidate to explain the ATM-mediated changes in ATP. To pursue this prediction, we analyzed the expression patterns of the 3,360 genes identified as potential regulated by NRF1- (Table S1K) and found that over 80% of them were significantly reduced in A-T cerebellar samples (Fig.6D, Table S1L). Further, of 2,470 NRF1 targets identified in both the ENCODE database and a previous NRF1 ChIP-Seq study (Satoh et al., 2013) we found that 440 appeared in our list of genes with altered expression in A-T (Fig.6E). NRF1 is a basic leucine zipper (bZIP) transcription factor (Fig.6B) that functions as a trans-activator of nuclear-encoded respiratory genes (Evans and Scarpulla, 1990; Scarpulla, 2008) as well as performing extra-mitochondrial regulatory roles (Satoh et al., 2013). KEGG pathway analysis of the 440 common NRF1 targets from the A-T regulated genes identified cell cycle, Wnt signaling, oxidative phosphorylation and circadian rhythm pathways as significantly altered (Fig.6F). We next used immunohistochemistry to determine the anatomic location of the NRF1 protein in normal and ATM-deficient brain. In wild-type mouse and control human samples, we found that the NRF1 immunoreactivity was substantially enriched in Purkinje cells compared to other cerebellar cell types. In the absence of ATM, NRF1 immunostaining in Purkinje cells was reduced and this reduction was most dramatic in the cell nucleus (Fig.6G). The implication of these findings is that ATM regulates NRF1 nuclear-cytoplasmic shuttling.

ATP depletion activates ATM with downstream effects on NRF1

Several studies have shown that ATM can function as a redox sensor because it can be activated by oxidative stress independently of its response to DNA damage (Ashley et al., 2012; Guo et al., 2010). We wondered, therefore, whether transient reductions in ATP might lead to ETC imbalances and that the resulting increase in reactive oxygen species (ROS) might activate ATM. To test this, we induced a modest depletion of ATP by treating cells with atractyloside (an inhibitor of the ADP/ATP carrier) or a more dramatic ATP depletion with oligomycin. Oligomycin resulted in significant cell death due to the severe ATP depletion (Fig.S4A-B); yet neither atractyloside nor oligomycin induced a significant DNA damage response (Fig.7A). Confident that our results would not be confounded by an independent response of ATM to DNA damage,

we used the less toxic atractyloside to test primary cultures of mouse neurons. We observed a good correlation between the atractyloside-induced oxidative stress and its progressive depletion of ATP, consistent with our hypothesis that cellular ATP depletion induces oxidative stress (Fig.S4C). To determine whether this ATP depletion-induced oxidative stress could activate ATM and impact NRF1, we tracked the subcellular location of NRF1 and activated (phosphorylated) ATM. Oxidative damage induced either directly with hydrogen peroxide (H_2O_2) or indirectly with atractyloside resulted in the nuclear localization of both phospho-ATM and NRF1 (Fig.7B-C). Significantly, the actions of both drugs were blocked by the anti-oxidant, N-acetyl cysteine (NAC) (Fig.7B-C). We determined that ATM was required for this response as NRF1 nuclear localization in response to either H_2O_2 or atractyloside was lost in $Atm^{-/-}$ neurons (Fig.7B-C). Equally important, induction of DNA damage alone was insufficient to trigger the NRF1 response. Etoposide, a topoisomerase II inhibitor that leads to DNA double strand breaks (Fig.7A), failed to reduce ATP levels (Fig.S4A) or stimulate the translocation of NRF1 to the nucleus (Fig.7B-C).

NRF1 helps to coordinate the expression of nuclear and mitochondrial genomes. We therefore suspected that nuclear localization of NRF1 was predictive of its transcriptional activity. We tested this using the live-cell reporter, NRF1mitoGFP, which contains an NRF1 binding sequence driving GFP protein expression fused with a mitochondrion localization signal; this allows us to simultaneously monitor NRF1-mediated transcriptional activity as well as mitochondrial morphology (Hodneland Nilsson et al., 2015). As predicted from the NRF1 localization, we observed that chronic moderate depletion of ATP induced NRF1-mediated transcription (Fig.7D and S4D). In contrast, when ATM activity was lost, or with NAC co-treatment, induction of NRF1 activity was abolished (Fig.7D and S4D). Significantly, in the presence of ATM, the mitochondria appeared to maintain their normal filamentous morphologies while in ATM-deficient situations, the mitochondria appeared more fragmented (Fig.S4E). Thus, in conditions when cellular ATP is chronically in demand, ATM is required to stimulate NRF1 activity and upregulate mitochondrial capacities. Taken together, our results suggest that the oxidative activation of ATM induced by suboptimal cellular ATP levels results in nuclear localization of NRF1 with important

consequences for nuclear encoded mitochondrial gene expression. The strong implication of the data is that ATM is a transducer of deficits in the energy status of a cell.

Next we asked how the kinase activity of ATM was able to alter the transcriptional activity of NRF1. NRF1 binds to its DNA recognition site as a homodimer whose formation is triggered by phosphorylation of the monomer (Gugneja and Scarpulla, 1997). While NRF1 gene transcription level were not significantly different (Fig.8A), Western blots revealed that in ATM-deficient cells the levels of NRF1 protein was reduced and the fraction of total NRF1 that was present as dimer dropped significantly (Fig.8B). Scrutiny of the NRF1 protein sequences of four mammalian species identified a total of eight well-conserved (S/T)Q sequences – canonical ATM target sites (Fig.8C). To determine whether these sites were ATM-targeted we co-expressed FLAG-tagged wild type (WT) ATM and GFP-tagged NRF1 in $Atm^{-/-}$. Subsequent immunoprecipitation revealed a robust physical interaction between the two (Fig.7D). Further, Western blot of the immunoprecipitates probed with a phospho(S/T)Q antibody revealed a strong band that was missing if a kinase-dead (KD) FLAG-ATM construct was used instead of a wild type one (Fig.7E). (S/T)Q phosphorylation of NRF1 was also lost in cultures treated with an ATM inhibitor (1 μM KU60019) but not with inhibitors of ATR (0.5 μM VE821) or DNA-PKcs (20 nM NU7441) (Fig.7F). These data suggest that specific ATM phosphorylation facilitates the homodimerization of NRF1. Although there are a total of eight potential (S/T)Q phosphorylation sites on NRF1 (Fig.7C), only the T259 residue located within the DNA-binding and dimerization (DIM) domain was previously identified by mass spectroscopy (Hornbeck et al., 2015). In our search for the relevant ATM target we therefore focused on T259 and created mutant clones of the NRF1 protein (Fig.8G). In the presence of wild type ATM, co-expression of the wild type NRF1 protein or the NLS-deletion mutant (NLSdel) resulted in robust phosphorylation of the (S/T)Q site (Fig.7G). However, when the deletion was extended into the DNA-binding and dimerization domain (DIM) (DIMdel: T87-E283), the (S/T)Q phosphorylation signal was lost (Fig.7G). Refining the ATM target residue still further, expression clones that carried threonine-to-alanine triple (T201A, T259A and T287A), double (T201A and T259A) mutations, or the single (T259A)

mutation resulted (Fig.7G) in similar loss of (S/T)Q phosphorylation. This confirms that the DIM domain as the most likely site of ATM-mediated NRF1 phosphorylation.

Although T259 is the putative ATM target, we noticed that ATM-mediated NRF1 dimerization was lost when either the NLS domain was deleted or the T259 residue was mutated. On the basis of these findings we speculated that both the nuclear localization of the protein and T259 phosphorylation are crucial for dimerization. We explored this prediction further with *in silico* analysis. While the structural features of the entire NRF1 protein have not yet been identified, superimposing the structure of the homodimerization regions of the NRF1 protein predicted by *Phyre*[2] on the bromodomain-containing protein-2/4 (BRD2/4 – Garcia-Gutierrez et al., 2012) with *ICM-Pro Browser* suggested that the DIM domain closely resembles a bromodomain (Fig.8H-I). *In silico* phosphorylation of T259 followed by *ZDOCK* docking and contact area analyses with *ICM-Pro Browser* suggested a more favorable interaction between monomers containing phosphorylated T259 than between non-phosphorylated monomers (Fig.7J, Table S1M, N). This predicts that T259 phosphorylation is a post-translational modification that triggers homodimerization. We validated this prediction experimentally by ectopically expressing the phosphomimetic T259E or non-phosphorylatable T259A NRF1 mutants *in vitro.* Western blot analysis showed that NRF1-T259E was able to dimerize efficiently, but T259A could not (Fig.7K). We separately verified the impact of NRF1 phosphorylation on nuclear-cytoplasmic shuttling (Fig.7L-M). Expression of NRF1 (T259E) promoted its nuclear localization while expression of NRF1 (T259A) led to a more prominent cytoplasmic localization in primary neurons. As expected, nuclear localization of wild type NRF1 was lost if cells were pre-treated with ATM inhibitor (Fig.7L-M). These results support a model in which ATM-dependent phosphorylation of NRF1 on T259 results in both NRF1 dimerization and nuclear localization.

Discussion

Our data present an expanded picture of the relationship between the ATM kinase and mitochondrial function and offer a new perspective on the reasons for the particular set of symptoms that are found in persons with A-T. ATM functions as a sensor and regulator of a

broad cellular response that integrates energy demand and mitochondrial biosynthesis. This newly identified ATM function is particularly important in large, energy intensive cells such as Purkinje cells. The key to the new pathway is the ATM-mediated phosphorylation of NRF1, a transcription factor responsible for the regulation of genes involved in mitochondrial structure and the electron transport chain. A linkage between ATM and mitochondria was first proposed by the Kastan lab (Valentin-Vega et al., 2012) who reported that ATM-deficient fibroblasts had defective mitophagy (the clearance of mitochondria by autophagy), and, as a result, accumulated mitochondria with aberrant morphologies. They identified the Beclin-1 protein, an autophagy inhibitor, as the mediator of the influence of ATM (Valentin-Vega et al., 2012). Fang et al. (2016) extended this line of investigation, showing that defective mitophagy was also found in xeroderma pigmentosa (XPA) cells. They suggested that the problem in XPA lay in PARP1 hyperactivation, which depleted the cellular levels of NAD^+, and thus reduced the activity of the NAD^+-responsive histone deacetylase, SIRT1. They proposed that the link between impaired mitophagy and ATM deficiency followed a similar $PARP1/NAD^+/SIRT1$ pathway. Other researchers have also reported on the importance of ATM in the regulation of mitochondrial function (Aird et al., 2015; Fang et al., 2016; Morita et al., 2014; Zakikhani et al., 2012). None of these earlier studies, however, focused on the more fundamental linkage between ATM and cellular energy levels.

Our results indicate that a key linkage between ATM and mitochondrial function is through ATP itself, specifically the oxidative stress generated during ATM insufficiency. If TCA cycle components (e.g. acetyl-CoA) decline, mitochondrial $NADH_2$ levels drop, the ETC becomes inefficient and the mitochondria cannot sustain an adequate proton gradient across their inner membrane. To protect the proton gradient, which is critical for cellular survival, the ATP synthase transiently runs in reverse and uses ATP (from stores or generated directed from cytosolic metabolism) to pump protons back across the inner mitochondrial membrane, thus reestablishing a healthy Δψm. If this situation persists, non-mitochondrial sources of ATP are insufficient, and a cell responds by increasing its mitochondrial content. This requires enhanced transcription from dozens of nuclear-encoded genes whose protein products collaborate to

form the functional electron transport chain. This response is critical to cell viability and thus is regulated by several intersecting pathways. ETC inefficiency also leads to the enhanced generation of reactive oxygen speciies (ROS). For example, chronic insufficiency of ATP also results in intracellular calcium increase (Le Masson et al., 2014), which activates multiple extramitochondrial ROS-generating enzymes (Yan et al., 2006). In a similar vein, cytosolic glutathione, a cellular ROS scavenger (Hargreaves et al., 2005), is synthesized *de novo* by two sequential enzymatic ATP-dependent reactions (Chen et al., 2005). In this and possibly other cases, reduced ATP levels lead to increased ROS. It is these intersecting ROS generating pathways that engage ATM in the regulation of cellular metabolism.

We propose that ATM, in its role as a redox sensor is activated by the increased ROS levels caused by the insufficiency of ATP. Activated in this way, ATM phosphorylates the transcription factor NRF1, a master transcriptional regulator of nuclear genes encoding proteins for the mitochondrial respiratory chain (Kelly and Scarpulla, 2004; Scarpulla, 2002). Activated ATM phosphorylates NRF1 at T259 to facilitate its formation of a transcriptionally active homodimer. It is noteworthy that this pathway of ATM activation is unrelated to its function in the DNA damage response pathway. The downstream targets of the two pathways are different as are the cellular responses. Whether subcellular localization or altered substrate specificity explains these differences remains to be determined. This newly recognized ATM response system is only revealed when the cell is under conditions of stress. While the activities of Complex I-IV are reduced in in the absence of ATM, the affected cells are still able to maintain a nearly normal basal oxygen consumption rate; the steady state leak of protons in $Atm^{-/-}$ neurons is also no different from the levels found wild type cells. It is only when the $\Delta\psi m$ is completely depolarized with FCCP or when the cells are challenged with a massive energy demand, such as that following a prolonged depolarization, that the homeostatic failure of $Atm^{-/-}$ cells with regard to their ATP levels becomes apparent.

This scenario of sustained ATP deficiency causing ATM-dependent activation of NRF1-related gene transcription offers fresh insight into why cerebellar ataxia, with its regional and cell type

specific pathology, manifests as the most prominent neurological phenotype of human A-T. Here the explanation seems to lie in both the physiology of the Purkinje cell and the expression pattern of NRF1. Cerebellar Purkinje cells, with their large cellular size and vast number of synaptic inputs (Carter and Bean, 2009; Sugimori and Llinas, 1990; Welsh et al., 2002), are the only efferent neurons of the cerebellar cortex and are often lost in diseases where ataxia is involved. Computational modeling predicts that each Purkinje cell consumes 62 times more energy than each cerebellar granule cell and 3.9 times more than a cortical pyramidal neuron (Fig.8N, Table S1O). Most of this energy (4.26×10^9 ATP/neuron/s) is used to run the Na^+/K^+-pumps that restore the neuronal membrane potential after an action potential (Fig.8N, Table S1O) (Howarth et al., 2012; Howarth et al., 2010). Furthermore, a Purkinje cell typically spikes at high frequency. During a 'burst', almost twice as much sodium enters during each spike as would be predicted from the kinetics of a single, isolated action potential (Carter and Bean, 2009). Because of this, a slower-spiking cortical pyramidal neuron uses only 25% more sodium than predicted – closer to the theoretical minimum. With their extraordinarily high metabolic/ATP demands, it follows that Purkinje cells are exquisitely vulnerable to oxidative stress and metabolic insufficiency (Sugimori and Llinas, 1990).

This explanation for the unique impact of ATM deficiency on the Purkinje cell is further supported by regional variations in the NRF1 protein itself. Immunocytochemistry has shown that the cerebellum in general and Purkinje cells in particular, have high levels of NRF1 protein. This presumably prepares the Purkinje cell to respond when ATP levels drop, but in the absence of ATM this cellular preparedness cannot be used to increase mitochondrial gene transcription. In keeping with this prediction a CNS-targeted NRF1 knockout produces Purkinje cell loss, motor ataxia and postnatal lethality (Kobayashi et al., 2011). It is also noteworthy that early onset cerebellar ataxia involving extensive Purkinje cell neurodegeneration is a frequent symptom of patients with mitochondrial DNA defects (Scaglia et al., 2005) or others with mutations of nuclear genes encoding proteins crucial for mitochondrial function (Ashley et al., 2012; Chen et al., 2007). Consistent with our model, these mutations all result in ETC defects (Ashley et al., 2012; Chen et al., 2007; Scaglia et al., 2005). Our findings can also be seen in the

context of the more traditional pathway for the regulation of ATP levels by the AMPK kinase. When AMP levels rise (or ATP levels fall) this heterotrimeric kinase is activated and alters the behavior of a number of downstream targets that enhance catabolic and inhibit anabolic functions, thus allowing the cell to restore ATP levels to normal (recently reviewed in Herzig and Shaw, 2018). Comparing the AMPK response to that of ATM suggests that the former is more likely to respond to moment-to-moment fluctuations in ATP levels, while the latter, with its requirement for significant oxidative damage, ensures a structural and more sustained response to large, chronic changes. This suggests that ATM functions as a failsafe mechanism that acts as a backup for the AMPK response. When ATP levels suffer a sustained drop to the point where the ETC is compromised and ROS are increased, a cell would be well served by being able to enhance its overall mitochondrial capacity in addition to adjusting the activity of Complex I-V.

Considered together with recent findings on the role of ATM in the modulation of synaptic vesicle trafficking (Cheng et al., 2018), our findings emphasize the importance of the localization, and independent functions, of ATM in cytoplasm and nucleus. Although it has been suggested that the neurodegenerative phenotype of A-T is mainly attributable to inefficient DSB repair, it has always been challenging to explain the full range of neurological and behavioral symptoms by invoking only this pathway. A-T cells show normal levels of DNA repair (Kastan and Lim, 2000) although the process is slower. Further, since DNA damage is mostly a stochastic event that occurs throughout the genome of all cells, it is challenging to explain how inefficient repair of genome-wide DNA damage explains the selective loss of Purkinje cells in all A-T patients. Combining the predicted energy demands unique to Purkinje cells with a defective capacity to respond to the drop in ATP levels brought on by high levels of neuronal activity, makes the neurological symptoms of A-T seem much less puzzling.

Materials and methods

Mitochondrial database. Associations between various symptoms of ataxia-telangiectasia were generated using the online database http://www.mitodb.com (Scheibye-Knudsen et al., 2013). On the home page of the database, Ataxia-telangiectasia was chosen as the disease to be tested. Once test results were generated, the number of mitochondrial versus non-mitochondrial diseases (including those in the Unknown category) associated with different symptoms of Ataxia-telangiectasia were tabulated on the subpage named "Barcode". Details of the data can be found in Table S1A.

Multiple protein sequence alignment. Protein sequences of NRF1 in *Homo sapiens* (human) [NP_001035199.1], *Pan troglodytes* (chimpanzee) [XP_001155812], *Mus musculus* (mouse) [NP_035068.3] and *Rattus norvegicus* (rat) [NP_001094178.1] were obtained from the NCBI-Gene database https://www.ncbi.nlm.nih.gov/gene . Multiple protein sequence alignment was performed on the Clustal Omega platform https://www.ebi.ac.uk/Tools/msa/clustalo/. Conserved S/TQ sites among all the species analyzed were manually identified for subsequent analysis.

Human microarray data mining. A human expression array dataset (GEO Accession GSE61019), comparing samples of A-T and control cerebella (Jiang et al., 2015), was analyzed with GEO2R. Raw data from each of the samples were extracted. *Gene Ontology (GO) analysis was performed*. All transcripts that reflected significant changes (Adjusted p-value ≤ 0.05) by the GEO2R analyzer were extracted (Table S1B, yellow highlights) and analyzed on the GATHER platform http://changlab.uth.tmc.edu/gather/gather.py (Chang and Nevins, 2006). Among all 30 GO terms that reflected the greatest possible significance (P=0.0001), we performed enrichment analysis (listed in descending order) based on the corresponding Bayes factors and the number of genes indicated in each GO term. We performed a second analysis using the *Gene Set Enrichment analysis (GSEA)* using the KEGG pathway groupings of genes http://software.broadinstitute.org/gsea/index.jsp (Subramanian et al., 2005). The full list of phenotypes enriched in the normal can be found in Table S1E. Pathways with normalized p-

value (NOM p-val) < 0.01 are highlighted in yellow. We performed a third analysis using *Human MitoCarta2.0 analysis*—a collection of 1,158 nuclear and mitochondrial genes encoding proteins with strong evidence for mitochondrial localization https://www.broadinstitute.org/files/shared/metabolism/mitocarta/human.mitocarta2.0.html (Calvo et al., 2016). The matching genes were extracted from the raw dataset of this human array for expression analysis. The full list of extracted data from each individual sample can be found in Table S1F. Table S1G contains the comparison data between the control versus A-T groups. Specific genome localization of each gene was also determined. Note that a list of 42 genes from the MitoCarta2.0 list failed to match probes on the human expression array. To identify potential transcription factors that result in global changes of gene expression, we performed *PSCAN gene promoter analysis*. The PSCAN platform http://159.149.160.88/pscan/ (Zambelli et al., 2009) was used to analyze the genes identified in the human expression array data that showed significant differences of expression in A-T and control cells. The list of gene IDs was first converted to the corresponding RefSeq mRNA IDs using the DAVID platform https://david.ncifcrf.gov (Huang da et al., 2009), and subsequently input into PSCAN using the "human and mouse" species option; a default promoter setting of "-450 to +50" was used to cover the regions located immediately adjacent to the transcription start site. The JASPAR 2014 matrices were also selected for analysis. The results are compiled in Supplementary Table S2. Detailed matrix information, sample mean score and statistics are listed for each site of interest. NRF1 was the gene that ranked highest in this analysis and was thus subjected to further scrutiny. The list of NRF1 binding sites in each input promoter, with corresponding gene name, oligo score, position with respect to transcription start site of the gene, oligo sequence and strand were extracted for further analysis.

In silico protein structure modeling, docking, and contact area analysis. With the protein sequence of the human NRF1 protein [NP_001035199.1] obtained from the NCBI-Gene database, the DNA binding and dimerization domain (aa 116-283) was used to predict the corresponding 3-D protein structure on the Phyre[2] engine http://www.sbg.bio.ic.ac.uk/~phyre2/html/page.cgi?id=index using the intensive modelling

mode (Kelley et al., 2015). The highest-ranking model (with score > 85), was used to overlap the excised bromodomain structures of BRD2 (PDB: 5UEW) and BRD4 (PDB: 4O76) on Molsoft ICM-Pro Browser, or directly adopted for subsequent *in silico* docking analysis for the homodimerized NRF1 structure on the ZDOCK server http://zdock.umassmed.edu (Pierce et al., 2014). The best-ranked model (with the highest ZDOCK score) from each analysis was then subjected to contact area analysis for visualizing interacting amino acid residues with the Molsoft ICM-Pro Browser. To evaluate the monomer structures, T259 was mutated to T259(PO$_4^-$) using Pymol mutagenesis tool prior to the same docking procedures and contact area analysis.

Animal maintenance, brain tissue isolation, primary neuronal and mouse embryonic fibroblast (MEF) culture. C57BL/6J and C57BL/6J and 129S6/SvEvTac-Atmtm1bal/J (*Atm$^{+/-}$*) (heterozygous for an engineered ATM kinase domain mutation Xu et al., 1996) were obtained from The Jackson Laboratory. Colonies were maintained in the Animal and Plant Care Facility of the Hong Kong University of Science and Technology (HKUST). Homozygous mutants (*Atm$^{-/-}$*) were generated by mating heterozygous carriers (*Atm$^{+/-}$*), as per The Jackson Laboratory protocol. All animal experimental protocols were approved by the Animal Ethics Committee at HKUST and their care was in accord with the institutional and Hong Kong guidelines

Brain tissue was isolated by anesthetizing adult mice with intraperitoneal administration of 1.25% (vol/vol) Avertin at a dosage of 30 mL/kg body weight. The heart of each mouse was then surgically exposed, the left chamber was catheterized, and the right atrium was opened. Chilled physiological saline was perfused transcardially for 3 minutes to remove blood from the body. After perfusion, the cranial bones were opened; cortex and cerebellum were harvested, snap-frozen in liquid nitrogen, and stored at -80 ^{o}C until use.

Embryonic cortical neurons were isolated by standard procedures (Chow et al., 2014). Embryos were collected in ice-cold PBS-glucose, and the cortical lobes were excised. Meninges were removed and the cortices were placed in 1X trypsin solution for 10 minutes, with manual

shaking at 5 min. After digestion, an equal volume of DMEM with 10% (vol/vol) FBS was added to inactivate the trypsin. Samples were then centrifuged at 2,500 rpm for 5 min. Supernatant was removed, followed by transferring the pellet to fresh Neurobasal medium supplemented with B-27, penicillin/streptomycin (1X) and L-glutamine (2mM; GlutaMAX, Invitrogen) prior to gentle re-suspension. Tissue was triturated 10 times through a 5-mL pipette and allowed to settle to the bottom of a 15-mL conical tube. Cells in solution above the pellet were removed. Surviving cells were identified by trypan blue exclusion and counted before plating on poly-L-lysine-coated (0.05 mg/mL) glass coverslips. Unless otherwise specified, cells were plated in 24-well plates at 50,000 cells per well and allowed to mature for over 7-10 days in vitro (DIV) before transfection or lentiviral transduction. For other experiments, cells were grown for a minimum of 14 days in vitro (DIV14) before drug-treatment experiments. Every 3 days half of the culture medium was replaced with an equal volume of fresh medium to sustain the culture.

MEF cell cultures were established from the same E16 embryos used for isolation and culture of primary cortical neurons. After the heads (for primary neuronal culture), tails, limbs, and internal organs were removed, the remaining tissue was minced and trypsinized for 20 min, then seeded into P100 culture dishes in 12 mL of the complete MEF medium [Iscove's modified Dulbecco medium (IMDM) containing 10% iron-supplied calf serum (Hyclone), with additional non-essential amino acids (NEAA), 2 mM GlutaMAX^{TM} and 1% Penicillin-Streptomycin.]. When cultures became confluent, the cells were split 1:2 or 1:3 then passaged two or three times to obtain a morphologically homogenous culture. At this point cells were harvested, and separated into aliquots that were either frozen or expanded for further studies.

Cell culture and transfection. Lines of primary human skin fibroblasts from A-T patients and their respective controls were obtained from the NIA Aging Cell Repository of the Coriell Institute for Medical Research and maintained in DMEM/10% FBS. Primary neurons were isolated from E16 embryos of C57BL/6J or $Atm^{+/-}$ pregnant mice and cultured as described above. DNA constructs were transfected with Lipofectamine LTX in the presence of Plus Reagent into both types of primary culture following the manufacturer's protocol. At 5 hours

after transfection, cells were refreshed with new culture medium (or conditioned medium for primary neurons) and further incubated for 48-72 hours to allow recovery and ectopic expression of transfected constructs.

Lentivirus production and transduction. Lentivirus stocks were produced as previously described with slight modifications (Cribbs et al., 2013). Human embryonic kidney 293FT cells (Invitrogen) were transfected using Lipofectamine 2000 (Invitrogen) with the expression of two helper plasmids: psPAX2 and pMD2.G. Ten micrograms of the transfer vector, 5 μg pMD2.G and 5 μg psPAX2 of DNA were used per 10-cm plate. Forty-eight hours after transfection, the supernatants of four plates were pooled, centrifuged at 780 x g for 5 minutes, filtered through a 0.45-μm pore size filter, and further centrifuged at 24,000 rpm for 2 hours. The resulting pellet was re-suspended in 100 μL of PBS. Lentivirus titration was performed with a p24 ELISA (Clontech).

Co-immunoprecipitation, SDS-PAGE-Western Blotting and Native PAGE-Western Blotting. Isolated brain tissues or cell pellets were homogenized in RIPA buffer (Millipore) with 1X complete protease inhibitor mixture (Roche) and 1X PhosSTOP phosphatase inhibitor mixture (Roche) on ice then centrifuged for 10 minutes at 18,400 x g to remove large debris. The protein concentration of the supernatant was determined by Bradford Assay (Bio-Rad). For co-immunoprecipitation, 1 mg of the total cell lysates was first incubated with control IgG (Santa Cruz Biotechnology) for 30 minutes, pre-cleared with 50 μL Dynabeads Protein G (Invitrogen), and then incubated with various antibodies overnight at 4^{o}C, using the suggested dilutions from the product datasheets. Beads bound with immune complexes were collected by DynaMag-2 (Life Technologies) and washed three times before elution in 90 μL of buffer containing 0.2M Glycine-HCl, pH 2.5, which was neutralized with 10 μL of neutralization buffer [1M Tris-HCl (pH 9.0)]. The eluates were subjected to 9-15% SDS-PAGE and Western Blotting analysis.

For SDS-PAGE (polyacrylamide gel electrophoresis), 100 μg of proteins derived from cell or tissue lysates were prepared in 5X sample buffer [10%w/v SDS; 10mM beta-mercaptoethanol;

20%v/v glycerol; 0.2M Tris-HCl, pH6.8; 0.05% Bromophenolblue]. With a Bio-Rad system, separating gels of different acrylamide percentages (6%-15%) were prepared with the following components in double distilled water: acrylamide/bis-acrylamide (30%/0.8% w/v); 1.5M Tris (pH=8.8); 10% (w/v) SDS; 10% (w/v) ammonium persulfate and TEMED) and a 5% stacking gel (5 ml prep: 2.975ml Water; 1.25 ml 0.5M Tris-HCl, pH6.8; 0.05 ml 10% (w/v) SDS; 0.67ml acrylamide/bis-acrylamide (30%/0.8% w/v); 0.05ml 10%(w/v) ammonium persulfate and 0.005 ml TEMED) were prepared. Samples were run in SDS-containing running buffer [25mM Tris-HCl; 200mM Glycine and 0.1% (w/v) SDS] until the dye front and the protein marker reached the foot of the glass plate. Standard immune-blotting procedures were used, which include protein transfer to polyvinylidene difluoride (PVDF) membranes, non-fat milk blocking, primary and secondary antibody incubation, followed by visualization with the SuperSignal West Dura/Femto Chemiluminescent Substrate (ThermoFisher Scientific).

For native-PAGE, the same discontinuous chloride and glycine ion fronts as SDS-PAGE were used to form moving boundaries that stack and then separate polypeptides by their charge to mass ratio. Proteins (100 µg) were prepared in a non-reducing non-denaturing sample buffer (2X: 62.5 mM Tris-HCl, pH 6.8; 25% glycerol; 1% Bromophenol Blue) without heating or boiling to maintain secondary structure and native charge density. With a Bio-Rad system, 10 ml/gel 8% native PAGE separating gel [2.6ml acrylamide/bis-acrylamide (30%/0.8% w/v); 7.29ml 1.5M Tris-HCl (pH8.8); 100 µl 10%(w/v) ammonium persulfate and 10 µl TEMED) and 5 ml/gel native PAGE stacking gel [4.275 ml 0.5M Tris-HCl. pH=6.8; 0.67ml Acrylamide/Bis-acrylamide (30%/0.8% w/v); 0.05ml 10%(w/v) ammonium persulfate; 5µl TEMED] were prepared. Samples were run in a non-denaturing running buffer (25mM Tris, 192 mM glycine, pH 8.3) with the entire system run on ice. After electrophoresis, standard immune-blotting procedures were followed as above.

Quantitative RT-PCR (qPCR) analysis. Total cellular RNA was purified from brain cortex tissues or cultured cells using the RNeasy mini kit (Qiagen) following the manufacturer's protocol. For quantitative real-time PCR (qPCR), RNA was reverse-transcribed using the High-Capacity cDNA

Reverse Transcription Kit (Applied Biosystems) according to the manufacturer's instructions. The resulting cDNA was analyzed by qRT-PCR using SYBR Green PCR Master Mix (Applied Biosystems). All reactions were performed in a Roche LightCycler (LC) 480 instrument using the following protocol: pre-incubation at 95 $^{\circ}$C for 15 min (1 cycle); denaturation at 94 $^{\circ}$C for 15 seconds, annealing and extension at 55 $^{\circ}$C for 30 seconds (40 cycles), melting at 95 $^{\circ}$C for 5 seconds, 65 $^{\circ}$C for 60 seconds and 95 $^{\circ}$C continues (1 cycle) followed by cooling at 40 $^{\circ}$C for 30 seconds. The specificity of the primers was confirmed by observing a single melting point peak. qPCR efficiency was calculated from the slope was between 95 and 105% with co-efficiency of reaction R^2=0.98-0.99. A total of 7-9 biological replicates x 4 technical replicates were performed for each treatment group. Data was analyzed using the comparative Ct method (ΔΔCt method).

Mitochondrial complex activity assays. Activities of Complexes I to IV were measured using its purchased from BioVision (K968-100, K660-100, K520-100 and K287-100) and that of Complex V was measured using a kit from Invitrogen (KHM2051). Measurements were conducted by strictly adhering to the manufacturers' instructions.

Imaging intracellular pyruvate level in single cells with the FRET sensor, Pyronic. Pyruvate sensor, Pyronic (Pyruvate Optical Nano-Indicator from CECs), was obtained from Addgene (Plasmid #51308). The exposure of the sensor to pyruvate causes an increase in mTFP (donor) fluorescence intensity and a decrease in Venus (mCFP, receiver) fluorescence intensity, thus resulting in a decrease in FRET efficiency. As reported previously (San Martin et al., 2014), the Pyronic biosensor was transfected into cortical neurons on DIV9. Twenty-four hours after transfection, cells were exposed to one of the following drug conditions: i) vehicle (saline); ii) 30 mM 2-deoxyglucose (2-DG) (Saraiva et al., 2010); iii) 1 μM UK5099 (Jourdain et al., 2016) or iv) 2.5mM oxamate (Choi et al., 2012)] in KRH-bicarbonate buffer (112mM NaCl, 1.25mM $CaCl_2$, 1.25mM $MgSO_4$, 1-2 mM glucose, 10mM HEPES, 24mM $NaHCO_3$ (pH7.4) and 3mM KCl). Cultures were then live-imaged for another 24 hours with imaging at 6-hour intervals in a 95% air/5% CO2-gassed incubator using a Leica SP8 confocal microscope equipped with a 63X oil-

immersion objective. Image acquisition was controlled by the Leica Application Suite X Microscope Software (LAS X). As nanosensor ratio measurements were performed, cells were excited at 430 nm for 0.2-0.8 seconds. Emission was divided with a Acousto-Optical Beam Splitter (Leica Microsystems Inc.), and recorded at 480±20 (mTFP) and 535±15nm (Venus). The ratio between mTFP and Venus along time was used to estimate pyruvate concentrations inside the cell.

Pyruvate colorimetric assay. Cellular contents of pyruvate were measured using kits purchased from BioVision (K609-100) following the manufacturers' instructions.

Imaging intracellular lactate level with in single cells with the FRET sensor, Laconic. Lactate sensor Laconic (LACtate Optical Nano Indicator from CECs) was obtained from Addgene (Plasmid #44238). Exposure of the sensor to lactate causes an increase in mTFP (donor) fluorescence intensity and a decrease in Venus (mCFP, receiver) fluorescence intensity, resulting in a reduction in FRET intensity. As described previously (San Martin et al., 2013), the Laconic biosensor was transfected into cortical neurons on DIV9. Twenty-four hours after transfection, cells were exposed to one of the following drug conditions: i) vehicle (saline); ii) 500 µM pCMBS (San Martin et al., 2014); or iii) 2.5 mM Oxamate (Choi et al., 2012)] in imaging buffer (112mM NaCl, 1.25mM CaCl2, 1.25mM MgSO4, 1–2mM glucose, 10mM HEPES, 24mM NaHCO3, pH 7.4, with 3 mM KCl). Transfected cells were then live imaged for another 24 hours images captured at 6-hour intervals in a 95% air/5% CO_2-gassed incubator using a Leica SP8 confocal microscope equipped with a 63X oil-immersion objective, controlled by the Leica Application Suite X Microscope Software (LAS X). As nanosensor ratio measurements were performed, cells were excited at 430 nm for 0.2-0.8 seconds. Emission was divided with a Acousto-Optical Beam Splitter (Leica Microsystems Inc.), and recorded at 480±20 (mTFP) and 535±15nm (Venus). The ratio between mTFP and Venus along time was used to estimate dynamics of pyruvate concentration inside the cell.

Lactate colorimetric assay. Cellular contents of lactate were measured using specific kits purchased from BioVision (K607-100), following the manufacturer's instructions.

Cytosol and mitochondria separation followed by isocitrate and succinate colorimetric assays. pMX-3XFLAG-EGFP-OMP25 plasmid was obtained from Addgene (Plasmid #83354). Human primary fibroblasts were first transfected with the plasmid as described above. In cases where the transfection efficiencies were 80% or greater, after 72 hours mitochondria were harvested by immunopurification (Chen et al., 2016b). 3X-FLAG-EGFP-OMP25-expressing cells were gently scraped into 1 ml KPBS (136 mM KCl, 10 mM KH_2PO_4, pH 7.25). The cell suspension was centrifuged at 1000 x g for 2 minutes, supernatant was discarded, and the cell pellet gently re-suspended in 1 ml of the kit-supplied assay buffer. Cells were homogenized with 25 strokes in a 2 ml Dounce homogenizer, taking care not to introduce air bubbles into the solution. The homogenate was then centrifuged and the mitochondria isolated by anti-FLAG IP (the entire procedure was performed on ice in a 4°C cold room with pre-chilled buffers and equipment). To increase the speed of isolation only one sample for one metabolite analysis was processed at a time. Anti-FLAG magnetic beads (200 µl) were pre-washed three times with KPBS by gentle pipetting using a wide-bore pipette tip. Beads were collected using a DynaMag (Thermo Fisher Scientific). Sample was then added to the beads and the mixture incubated overnight. The following day antibody-bounded mitochondria were separated from the cytosol using the DynaMag; the cytosol fraction was saved for metabolite analysis. The isolated beads were quickly washed twice with 1 ml KPBS, re-suspended in 1 ml assay buffer, and transferred to a new tube. An aliquot of the mitochondrial suspension (250 µl) was taken for detergent lysis and protein quantification; the remainder was used for metabolite extraction analyses.

Cytosolic and mitochondrial contents of isocitrate and succinate were measured using kits purchased from BioVision (K656-100 and K649-100 respectively).

Live imaging of cytoplasmic redox status. Cyto-roGFP obtained from Addgene (Plasmid #49435) senses redox changes in a cell (Waypa et al., 2010). The Cyto-roGFP biosensor was

transfected into cortical neurons on DIV9, and 24 hours after transfection, neurons were exposed to one of the following conditions: i) vehicle (saline); ii) 10 nM H_2O_2 (Wang et al., 2007); or iii) 1 µM atractyloside (Schutt et al., 2012)] in Neurobasal culture medium. Cultures were then imaged for 24 hours at hourly intervals in a 95% air/5% CO2-gassed incubator using a Leica SP8 confocal microscope equipped with a 63X oil-immersion objective, controlled by the Leica Application Suite X Microscope Software (LAS X). The redox-sensitive protein reporter has excitation maxima at 400±15nm and 484±15nm, and an emission maximum at 525±15nm. The relative amplitudes of these peaks depends on the state of oxidation. With increased oxidation the 400±15nm excitation peak increases while the 484±15nm peak decreases (Hanson et al., 2004), Importantly, the ratiometric characteristics of the analysis renders the results independent of the expression levels of the plasmid in any one cell. Data were collected with the Leica Application Suite X Microscope Software. The fluorescence excitation ratios were obtained by dividing integrated intensities obtained from manually selected portions of the imaged regions of intact whole cells collected using 400±15nm and 480±15nm excitation filters after appropriate background correction. Background correction was performed by subtracting the intensity of a nearby cell-free region from the signal of the imaged cell.

Imaging intracellular ATP-to-ADP ratio. Perceval High Range (PercevalHR) obtained from Addgene (Plasmid #49082) senses changes in the ATP:ADP ratios in mammalian cells (Tantama et al., 2013). We transfected PercevalHR into cortical neurons on DIV9. Twenty-four hours after transfection, neurons were imaged on a Carl Zeiss AxioOberver Z1 Inverted Microscope with a piezo Z stage glass slide insert. The sensor has two distinct peaks of excitation, ATP binding increases the fluorescence at 500 nm, while ADP increases fluorescence at 420 nm excitation. PercevalHR was excited sequentially using 482±15nm and 445±15nm band pass filters; emission was measured 529±15nm band pass filter. The excitation and emission signals were optically separated with a 490 nm short pass dichroic mirror. Images were taken over a period of 72 minutes (capture time, 2 min). The ratio of the fluorescence signal at the two different excitation wavelengths (F_{high}/F_{low}), reports the ATP occupancy of the reporter, and is independent of the amount of sensor protein in the cell.

During the imaging process, neurons were placed in a custom chamber which was continuously perfused with a bathing solution (2 ml/min; 32-34^{o}C) containing 120mM NaCl, 3mM KCl, 2mM CaCl$_2$, 1mM MgCl$_2$, 3mM NaHCO$_3$, 1.25mM NaH2PO4, 15mM HEPES, 5mM glucose, 0.2mM pyruvate, and 0.5mM GlutaMax adjusted to pH 7.4. We determined a baseline reporter signal by imaging a resting cell for at least 25 minutes. After ensuring a stable signal, 50 µM glutamate was applied for 30 seconds to strongly excite the neurons. Following the depolarization, the cell would need considerable amounts of ATP to drive the Na$^+$/K$^+$-ATPase (sodium pump) to be able to restore the resting ion gradients. We followed the drop in ATP and its subsequent restoration by the mitochondria for 30-45 minutes after the glutamate stimulus.

ATP colorimetric assay. The cellular contents of ATP were measured using kits purchased from BioVision (K354-100).

Determination of mitochondrial membrane potential. Tetramethylrhodamine, methyl ester (TMRM, Thermofisher) is a cell-permeant, cationic, red-orange fluorescent dye that is sensitive to the mitochondrial membrane potential; Mitotracker green (Thermofisher) – a mitochondrial dye that is insensitive to Δψm – was used as a counterstain. Stock solutions of the dyes (10mM) were prepared in anhydrous dimethylsulfoxide (DMSO). Cultured cortical neurons or human fibroblasts were first washed thrice with Tyrode's buffer (145 mM NaCl, 5 mM KCl, 10 mM glucose, 1.5 mM CaCl$_2$, 1 mM MgCl$_2$, and 10 mM HEPES; adjusted to pH 7.4 with NaOH). Working concentrations (20 nM) of TMRM and Mitotracker green were prepared in Tyrode's buffer and applied to the cells. After incubation in dark at room temperature for 45 minutes, imaging was performed on a confocal microscope (Leica, Sp8) using the time-series and sequential scanning programs. TMRM was excited at 514±15nm and and the emmision at 570±15nm was measured. Mitotracker green was excited at 480±15nm and detected at 520±15nm. To test changes in mitochondrial membrane potential, 1 µM carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP) or 2 µg/ml of oligomycin, were applied, which

depolarized or hyperpolarized the mitochondrial membrane potential, respectively (Joshi and Bakowska, 2011).

Assessment of mitochondrial function. The mitochondrial oxygen consumption rate (OCR) in both primary neurons and human fibroblasts were assessed using a Seahorse Bioscience XF^e24 analyzer (Agilent, Massachusetts, USA) in combination with the Seahorse Bioscience XF Cell Mito Stress Test assay kit. In this assay, sequential additions of the ATP synthase inhibitor, oligomycin, the mitochondrial uncoupler, carbonyl cyanide 4-(trifluoromethoxy)-phenylhydrazone (FCCP) and the complex I + II inhibitors rotenone + antimycin A provide insight into different aspects of mitochondrial function as described in the text. Experiments were performed on intact adherent cells.

Coating and cell plating. As noted that the surface area of each well of the 24-well Seahorse XF24 microplate is identical to that of standard 96-well microplates (0.32 cm^2). Poly-L-lysine (0.1 mg/ml in sterile borate buffer [pH8.4], 500 μl/well) was incubated overnight at 37^oC, 5% CO_2, and 95% humidity. The substrate solution was then removed, and the microplates were dried without further rinsing under a sterile laminar flow hood. Dissociated primary cortical neurons were isolated as described above. 100,000 neurons/well or 80,000 primary fibroblasts/well were seeded in a volume of 150 μl. Immediate after seeding, plates were incubated for 1 hour at 37 °C prior bringing the total volume of each well to 500 μl with corresponding culturing medium (Neurobasal medium for neurons; DMEM with 10% FBS for human fibroblast). Fibroblast cultures were maintained for 2 days prior to analysis. For neurons, after 1 night of settling, cultures were treated with 20 μM FDU in Neurobasal medium for 24 hours with medium changed every other day; this method eliminates ~99% of non-neuronal cells. Neurons were maintained until DIV7 prior subsequent analysis.

Cell mito stress test assay. Growth medium was removed, leaving 50 μl in each well after which cells were rinsed twice with 400 μl of pre-warmed assay medium consisting of XF base medium supplemented with 25 mM glucose, 2 mM glutamine and 1 mM sodium pyruvate buffrered to

pH7.4. Following the rinses, 475 μl assay medium and 50 μl conditioned medium (525 μl final) was added to each well. Cells were then incubated in a 37 $^{\circ}$C incubator without CO_2 for 1 hour after which pre-warmed oligomycin, FCCP, rotenone and antimycin A solution were loaded into injector ports A, B and C of the sensor cartridge to achieve final concentrations of 0.5 μM for oligomycin, 1 μM for FCCP and 1 μM for rotenone and antimycin A. The cartridge was calibrated with the XF24 analyzer, and the assay performed as described (Nicholls et al., 2010).

Oxygen consumption rate (OCR) was measured and changes from baseline rates were tracked following the sequential addition of oligomycin, FCCP and the rotenone plus antimycin A mixture. This allowed for calculation of the basal respiration rate, proton leakage, maximal respiration, spare respiratory capacity, non-mitochondrial respiration and ATP production (Nicholls et al., 2010) by Seahorse XF24 software version 1.8. At the end of each assay, cells were washed once with an excess of room temperature PBS, lysed with ice-cold RIPA buffer (0.15 M NaCl, 1 mM EDTA, 1 mM EGTA, 0.5% sodium deoxycholate, 0.1% SDS, 1% Triton X-100, 50 mM Tris-HCl [pH 8] with added protease and phosphatase inhibitor cocktails). Protein content was estimated by the Bio-Rad Dc protein assay (Bio-Rad, Hercules, CA, USA) using a Molecular Devices Softmax M3 microplate reader (Sunnyvale, CA, USA). Data from each well were normalized for total protein content. Normalized Seahorse XFe24 measurements were used to calculate a mean (± SEM) for each density group and each plate using Seahorse XF24 software version 1.8 (Seahorse Bioscience, Billerica, MA, USA).

Metabolic Fuel Flux Assays. The Mito Fuel Flex Tests were performed on Seahorse XFe24 Bio-analyzer (Agilent). All assays were performed following manufacture's protocols. Briefly, the test inhibits import of three major metabolic substrates (pyruvate, fatty acids, and/or glutamine) with mitochondrial pyruvate carrier inhibitor UK5099 (2 μM), carnitine palmitoyltransferase 1A inhibitor etomoxir (4 μM), or glutaminase inhibitor BPTES (3 μM). This test determines cellular dependence on each of the metabolite to fuel mitochondrial metabolism by inhibiting the individual substrate import. Baseline OCR was monitored for 18 minutes followed by sequential

inhibitor injections (i.e. Treatment 1 or Treatment 2) with OCR reading for 1 hour following each treatment. The inhibitor treatments and calculations are shown below:

Dependency (%)=[(Baseline OCR-Target inhibitor OCR)/(Baseline OCR-All inhibitors OCR)]x100%

Metabolic tests and associated inhibitors		
Metabolite test	Treatment 1	Treatment 2
Glucose/Pyruvate dependence	UK5099	Etomoxir + BPTES
Glutamine dependence	BPTES	Etomoxir + UK5099
Fatty acid dependence	Etomoxir	BPTES + UK5099

Immunocytochemistry. For immunostaining, primary neuronal cultures were grown on 13-mm coverslips (Marienfeld) in 24-well plates. At the appropriate times cultures were fixed with fresh 4% (wt/vol) paraformaldehyde (Sigma-Aldrich) for 10 min, and the cells permeabilized by treatment with 0.3% Triton X-100 in PBS for 10 min. After blocking with 5% (wt/vol) BSA in PBS for 1 hour, primary antibodies were added and incubated overnight at 4 °C. The following day, coverslips were washed three times (10 min each) with PBS. After rinsing, secondary antibodies applied for 1 h at room temperature followed by three additional washes with PBS. The coverslips were then inverted and mounted on glass slides with ProLong Gold Antifade Reagent (Life Technologies). Immunofluorescence was analyzed and Z-stacked maximum projected images were photographed using a TCS Sp8 confocal microscope (Leica Microsystems Inc.).

For immunostaining of brain sections, we performed antigen retrieval by treating the mounted cryostat sections with citrate buffer at 95 °C for 15 min followed by cooling to room temperature for 1 hour. Sections were rinsed in PBS, blocked with 5% (vol/vol) donkey serum/PBS for 1 hour before immunostaing as above.

LIVE/DEAD cell viability assays. The LIVE/DEAD viability/cytotoxicity kit for mammalian cells was purchased from Thermofisher. The method relies on two probes — calcein AM and ethidium homodimer-1 (EthD-1). Cultured cortical neurons or human fibroblasts were washed

three times with 500 volumes of Dulbecco's phosphate-buffered saline (D-PBS) to remove or dilute serum esterase activity generally present in serum-supplemented growth media. Two hundred microliters of the calcein/EthD working solution (1 µM and 2 µM respectively in D-PBS) were applied to the surface of a coverslip or 35-mm disposable petri-dish and incubated for 45 minutes at room temperature in the dark. Imaging was performed on a confocal laser microscope (Leica, Sp8), with the sequential scanning program. Calcein was excited at 485±15nm and emission was measured at 530±15 nm; EthD-1 was excited at 530±15nm with emission measured at 645±15nm. Quantification of live (green) and dead (red) cells was performed randomly on 10 fields from each of 3-4 blinded samples. The percentage of dead cells was calculated as (number of dead cells/total number of live and dead cells) x 100%.

Monitoring NRF1 nuclear transcription activity and changes in mitochondrial morphology. The NRF1mitoGFP reporter was a kind gift from Prof. Johan Tronstad (Hodneland Nilsson et al., 2015). Cortical neurons were transfected with reporter-expressing plasmids on DIV9. Twenty-four hours after transfection, cells were exposed to one of the following drug conditions i) vehicle (saline); ii) 1 µM atractyloside alone or iii) atractyloside plus 5 mM N-acetylcysteine (NAC) (Chao et al., 2016) followed by live-imaging for another 24 hours in a 95% air/5% CO_2-gassed chamber mounted on a Leica SP8 confocal microscope equipped with a 63X oil-immersion objective, controlled by the Leica Application Suite X Microscope Software (LAS X). GFP was excited at 480±15nm and emission detected at 520±15nm. To anlayze mitochondrial morphology the GFP signals of each cell (in a different set of cultures) were imaged at 24 hours after drug treatment and later the z-stack images were used to construct the 3D model by the Leica Application Suite X Microscope Software (LAS X) for morphology analysis.

Nrf1 expression plasmid and mutated Nrf1. A mouse cDNA ORF clone encoding NM_001164226 was purchased from OriGene (catalog number MR226726). EGFP coding sequence was fused to the C-terminus by Gibson assembly to make pCMV6-Nrf1-EGFP. Site-directed mutagenesis used the NEBaseChanger kit (NEB), with oligonucleotide primers designed using the manufacturer's website. All mutations and deletions were confirmed by sequencing.

Quantification of the relative abundance of mouse mitochondrial DNA to nuclear DNA. The quantification protocol for mouse mitochondrial DNA without co-amplification of nuclear mitochondrial insertion sequences was adopted (Malik et al., 2016). To prepare genomic DNA, mouse tissue samples were homogenized using TissueLyser (Qiagen) to eliminate cross-contamination. Total genomic DNA was extracted using the DNA easy Blood & Tissue kit according to the manufacturer's instruction (Qiagen). Before proceeding to qPCR, the DNA template was subjected to pre-treatment shearing (Bath sonicator Kerry, Pulsatron 55 – 38kHz for 10 minutes) as described previously in order to avoid dilution bias (Malik et al., 2011). The template concentration was determined using NanoDrop and adjusted to 10 ng/µl. To avoid errors arising from repeated freeze thaw cycles DNA samples were kept at $4^{o}C$ for the duration of study.

The mitochondrial DNA content was assessed by absolute quantification using real time PCR. Primers of mouse mitochondrial DNA (mMito) and mouse single copy nuclear β-2 microglobulin (B2M) were used to amplify the respective products. PCR products were purified and used to prepare dilution standards for both amplicons using range of dilutions from 10^{2}-10^{8} copies per 2 µl for quantification. Mitochondrial copy number per cell was determined by carrying out qPCR in a total volume of 10 µl, containing 5 µl of Quantifast SYBR Master Mix (Qiagen), 0.5 µl of forward and reverse primer (400 nM final concentration each), 2 µl template DNA and 2 µl DNase free water. The reactions were performed in a Roche LightCycler (LC) 480 instrument using the following protocol: pre-incubation at 95 ^{o}C for 5 min (1 cycle); denaturation at 95 ^{o}C for 10 seconds, annealing and extension at 60 ^{o}C for 30 seconds (40 cycles), melting at 95 ^{o}C for 5 seconds, 65 ^{o}C for 60 seconds and 95 ^{o}C continues (1 cycle), cooling at 40 ^{o}C for 30 seconds. The specificity of the primers was confirmed by the presence of a single melt peak. qPCR efficiency calculated from the slope between 95 and 105% with co-efficient of reaction, R^{2}=0.98-0.99. A total of 7-9 biological replicates x 4 technical replicates were performed for each treatment group. Data was analyzed using the comparative Ct method (ΔΔCt method).

Reagents and resources

Antibodies

Anti-Actin (Cell Signaling Technology, #4970)

Anti-ATM (phospho S1981) (Abcam, ab81292)

Anti-Gamma-H2AX (phospho S139) (Abcam, ab2893)

Anti-GFP (Santa Cruz Biotechnology, sc-8334)

Anti- MAP2 (Abcam, ab5392)

Anti-NRF1 (ThermoFisher Scientific, PA5-40912)

Anti-SDHA (Cell Signaling Technology, #5839)

Anti- 53BP1 (Cell Signaling Technology, #4937)

Anti-FLAG M2 (Sigma, F3165)

Chemicals and assay kits

Agilent Seahorse XF Cell Mito Stress Test Kit (103015-100)

Alpha-Ketoglurate Colorimetric/Fluorometric Assay Kit (BioVision, K677-100)

Atractyloside (potassium salt) (Cayman, #14804)

ATP Colorimetric/Fluorometric Assay Kit (BioVision, K354-100)

B27 Supplement (50X), serum free (ThermoFisher Scientific, 17504044)

Carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP) (Sigma, C2920-50MG)

Cytochrome Oxidase Activity Colorimetric Assay Kit (BioVision, K287-100)

Etoposide (Sigma, E1383-25MG)

GlutaMAX supplement (ThermoFisher Scientific, 35050061)

Gibson Assembly Master Mix (NEB, E2611)

High capacity cDNA Reverse Transcription kit (Applied Biosystems)

HyClone^{TM} Calf Serum (FisherScientific, SH3007203)

Hydrogen peroxide solution 30% (w/w) in water (Sigma, H1009-100ML)

Iscove's modified Dulbecco medium (IMDM) (ThermoFisher Scientific, 12440053)

Isocitrate Colorimetric Assay Kit (Biovision, K656-100)

KU60019 (Selleckchem, S1570)

Lactate Colorimetric/Fluorometric Assay Kit (BioVision, K607-100)

L-Glutamic acid (Sigma, G1251-100G)

LIVE/DEAD^{TM} Viability/ Cytotoxicity Kit, for mammalian cells (L3224)

Lipofectamine 2000 (ThermoFisher Scientific, 11668019)

Lipofectamine LTX with Plus reagent (ThermoFisher Scientific, 15338100)

MEM Non-essential amino acid solution (100X) (ThermoFisher Scientific, 11140050)

Mitochondrial Complex I Activity Colorimetric Assay Kit (Biovision, K968-100)

Mitochondrial Complex III Activity Assay Kit (BioVision, K520-100)

MitoProfile® Human Complex V Activity (Invitrogen, KHM2051)

Mitotracker Green FM (ThermoFisher Scientific, M7514)

Neurobasal medium (ThermoFisher Scientific, 21103049)

N-Acetyl-L-cysteine (Sigma, A7250-25G)

NU7441 (Tocris Bioscience, #3712)

Oligomycin (Sigma, 75351-5MG)

Opti-MEM^{TM} reduced serum mdium, GlutaMAX Supplement (ThermoFisher, 51985034)

Penicillin-Streptomycin (10,000U/mL) (ThermoFisher Scientific, 15140122)

pCMBS (Clearsynth, CS-T-50790)

Pyruvate Colorimetric/Fluorometric Assay Kit (BioVision, K609-100)

Sodium Oxamate (Sigma, O2751-5G)

Succinate Dehydrogenase Activity Colorimetric Assay Kit (BioVision, K660-100)

Succinate (Succinic Acid) Colorimetric Assay Kit (BioVision, K649-100)

SYBR Green PCR Master Mix (Applied Biosystems)

Tetramethylrhodamine, Methyl Ester, Perchlorate (TMRM) (ThermoFisher Scientific, T668)

UK5099 (Merck-Calbiochem, CAS56396-35-1)

VE821 (Selleckchem, S8007)

2-Deoxy-D-glucose (Sigma, D8375-1G)

Recombinant DNA

Laconic (Addgene, #44238)

pMX-3XFLAG-EGFP-OMP25 (Addgene, #83354)

pcDNA3.1(+)FLAG-HIS-ATM-wt (FLAG-ATM-WT) (Addgene, #31985)

pcDNA3.1(+)FLAG-HIS-ATM-kd (FLAG-ATM-KD) (Addgene, #31986)

pEGFP-N1 (EGFP-vector) (Clontech)

EGFP-Nrf1-WT (in-house)

EGFP-NRF1-NLSdel (in-house)

EGFP-NRF1-DIMdel (in house)

EGFP-NRF1-3XA (in house)

EGFP-NRF1-2XA (in house)

EGFP-NRF1-T259A (in house)

EGFP-NRF1-T259E (in house)

NRF1mitoGFP (A gift from Prof. Johan Tronstad)

Perceval High Range (Addgene, #49082)

Pyronic (Addgene, #51308)

Cyto-roGFP (Addgene, #49435)

Sequence-Based Reagents

Primers for real time PCR

GENE SYMBOLS	NCBI Accession number (mouse)	Forward Primer (5' to 3')	Reverse Primer (5' to 3')
Ndufv1	NM_133666.3	AAGCCATCGCTCGTCTCATT	TCTCCCTTCACAAATCGGGC
Ndufv2	NM_028388.3	ATGGTGCTGGAGGAGCCTTA	CGGAGATAGGTAGCCATCCA
Ndufs1	NM_145518.2	GTTTTGAGGCACCGCTGTTT	ACCTCGCAGAATGAATGCCT
Ndufs2	NM_153064.5	AAAGCGGTGACCAACATGAC	GAAGGGCCTGCAGATAGGTC
Ndufs3	NM_026688.2	CTGACTTGACGGCAGTGGAT	CATACCAATTGGCCGCGATG
Ndufs4	NM_010887.2	CAGACAACCAGACTCGGGAC	TGCATGTTATTGCGAGCAGG
Ndufs6	NM_010888.2	GCATACCGGCCAGGTGTATG	AGGCTATGATGCGGTGCTC
Ndufs7	NM_029272.3	AGCTCTCTCATCTTCCCCGA	CCGAAGGTCATAGGCCACAG
Ndufs8	NM_001106322.2	TGACAGTGCAGCTCGGATTC	AAGGGGTAGTTGATGGTGGC
Ndufa1	NM_019443.2	CTCGCCATTATGGGGGTGTG	ACTGGTACTGAACACGAGCA
Ndufa2	NM_010885.5	AACGGTACGTGGAGCTGAAG	GCTTTGCCGCTTAGCACATT

GENE SYMBOLS	NCBI Accession number (mouse)	Forward Primer (5' to 3')	Reverse Primer (5' to 3')
Ndufa9	NM_025358.3	GTGTTCTGTGGCTCATCCCA	TGTGACCCCATTCGTCCAAG
Ndufa10	NM_024197.1	TCCAGGACATCGAGAATGCG	TCCAGTCGTCCTGTTTCAGC
Ndufa11	NM_027244.1	AGCGATGTTTGGCCTTACCA	AGTGCCCATGTAGACACAGC
Ndufa12	NM_025551.3	CTCCGACGACTAATCCACCG	GTGTCGAAGGTGGAACCCAT
Ndufb3	NM_025597.3	AGGGACGCCATTAGAAACGG	CCCAAAGCTACCACAAACGC
Sdha	NM_023281.1	AGTGCGGGTCGATGAGTATG	GGATAGCAGGAGGTACGGTG
Ttc19	NM_028360.2	GCTCTCGCCTGGTTCTCG	CACTCTCATACGCAAGGCGA
Uqcrq	NM_025352.3	ACGGCACGTGATCTCCTACA	AGACCACTACAAACGGCGG
Cox6b1	NM_025628.2	GCTGTGAGAAGGCAATGACG	TTCAGCTATGCGGTCATCCC
Atp5e	NM_025983.3	GGACTCAGCTACATCCGGTTT	TGCTGCCCGAAGTCTTCTC
Adck3	NM_023341.3	GACTTCGGCGCAACTAGAGA	TGGCCTTGACCTCATAACCG
Coq2	NM_027978.2	GTCCTTCGTCTTCCTTGGGG	CCAACTGAGGCCAAAACGTG
Coq9	NM_026452.2	CCAGTCCTTGGGTCTCTCCA	ACCTGGTTGAGGCGAGCATT
Hccs	NM_008222.5	TCGTTGGTCCGATTTGGAGG	GATCCAGTCATGGCGGTCAA
Pdss1	NM_019501.3	GCGCGCAGATACCCTATTTT	CAACTGCACATTGCTGGTGT
Pdss2	NM_027772.2	CGCCACTGGTGGTACTTCAG	CCTGCATGGCGATATTGCTG
Abcb7	NM_009592.1	TGCTCGCGATACATTCCTGG	TTCTGTAGTCTGGGACAGCC
Bola3	NM_175277.4	CCTCGAGCCACAGCTATTCA	CGTGCATGCCCTTGATCTCT
Fxn	NM_008044.2	GGGGACATTGGACAACCCAA	CAGCTTAATGGTGAGCACGC
Nubpl	NM_029760.2	TGTGGGCGCCAGTTATTAGG	AAGGTTCACTGCGGTGGTAG
Nfu1	NM_020045.3	ATGTAGCGACTCCACACACC	GGTCCTCGTCTCAAGAACCG
Ndufaf1	NM_027175.4	AACCGAAGTGCCCTGCTTTA	TCAAAAGCACCCCGTGGAAT
Ndufaf2	NM_001127346.1	ACTACGTCGCCGAGTACAAG	CTCGGTTGGGATGTCTCCTG
Ndufaf3	NM_023247.1	GGATTTCGCTGCCTTTTCCG	CTTTGCAGCAGTGAGATGCG
Ndufaf4	NM_026742.4	TTCCAAAGATCCGGTGCCTG	TTGCCTTTGGGAATGTCCGT
C8orf38	NM_001085493.2	TCATGCTGCAAGCCACATTG	TGCTGCACACAGACATCCAT
C20orf7	NM_027093.4	AAAGGTCGTTCTTTTGCGCC	AAATTTCATGGGGTCGGGCT
Acad9	NM_172678.4	TCGGCGGAATCACTAATGGG	ATCGCCCACTGTTCAGGATG
Foxred1	NM_172291.2	CTTTGCTCCAGGGTCTTCGT	GTTCCCCAGTTGGGGTCTTC
Sdhaf1	NM_001033140.3	ACGTGCTGCGTATCGAGTAT	TTCTTTGGGGCACCACCATC
Bcs1l	NM_025784.5	CCTTCTTTATGGGCCCCCTG	ATGGTTGAGCCGGTCATCTG
Cox10	NM_178379.3	TCAGTAATCAGTGGGTCACGC	GAAAGACGGCAGCTCATCCT

GENE SYMBOLS	NCBI Accession number (mouse)	Forward Primer (5' to 3')	Reverse Primer (5' to 3')
Cox14	NM_183256.3	TACAAGACCTTCTCTGCCTCG	GACTCCTGAGGTCTTCTGCTCT
Cox15	NM_144874.4	ATCCCAGAGGACCTCCTGAC	GTGACAGACGTGACTCCCAG
Fastkd2	NM_172422.3	TCCTAGAAGCTCTGGCGAGT	TCACCCGAAGTCAGCAGTTC
Lrpprc	NM_028233.2	AGAAGACACTGCGTTCCAGG	AGCTTCTCAGGGGGCAAATC
Sco1	NM_001040026.1	GGGGGCCATTTTCCCTTACA	AGGGCAATGAGTGAAGCCAA
Sco2	NM_001111288.1	TGGACCACTCCATTGCCATC	AGGGCAAGACACTGTGGAAG
Surf1	NM_013677.2	ATGGTGGATCCTGTCCGAGA	GGTGACTCCCAAGTCAGAGC
Taco1	NM_027346.1	AGAGTTGTGTCGCAGCAAGA	ATGTCCAGATGGCACTTGGG
Atpaf2	NM_145427.2	TGATTGACTTGCGCCTGACT	GACAAAAAGGGTGCCAGCAG
Tmem70	NM_026392.1	TTGGGAACAATGTGTCCGCT	GGCAAGAATGCAAGGCTGAC
Mrpl3	NM_053159.3	TGACCCCGGCTTAGAGAGAA	GTAAAGGCCACGGCTCATCT
Mrps16	NM_025440.2	GTGGCTGCTCACAACAAGTG	CGGATACGGTCCAGGTTGAG
Mrps22	NM_025485.3	AGCTTCATGGGAAGAACGGG	GCTCAAACTGGGCAACACAG
Aars2	NM_198608.2	ACGCCTTCCTGAGCTTCTTC	GGAAGCCTGCCATCTCACTT
Dars2	NM_172644.4	ATCCAGACCCATCGGGAAGA	ACGAACGCAACTCTCCACAT
Gars	NM_180678.3	TCTGGAAGCAAAGGAGCTGG	CTACCGGCCCGAAGTCATAC
Kars	NM_053092.3	GCTGAGGAGGAGACTCTGGA	GGGTGACATCAGTCAGGTGG
Rars2	NM_181406.4	CTGGTTTCCAGCTGTTTGGC	TCGGTGGAAGAACTCGTGTG
Ercc6	NM_001081221.1	GTCCACAGCCCAAGCAAAAG	GTGCACGCCCACTGATTTTT
Ercc8	NM_028042	GCTTTTTGTCCGCTCGACAG	CAAGGTCATAGAGTACAACCACG
Polg	NM_017462	CGGGAGCTGTTCGTCAAAG	CAGGTAGGACACACCCATCT
Gfm1	NM_138591	AGCAGGCTGATTGGAAGACC	CCCGTTCTGTTAGCGTCGT
Mpv17	NM_008622	TGGCGAGCATACCAGAGAG	GGTCTAAAACTTTGTACCAGCCT
Mtfmt	NM_027134	GAAAAATCTGCCCGAAAGTCTGA	ATGGCAAGGTGAAGTCTGAGT
Mtpap	NM_026157	GTGGGGTTATTGACCCGTTTG	GTTCTTCGCTCCCGATGCTC
Polg	NM_017462	GAGCCTGCCTTACTTGGAGG	GGCTGCACCAGGAATACCA
Tsfm	NM_025537	ATCGGCTCTCCACGTACAGTA	CTCTCCCAGTTTACCAATCGC
Tufm	NM_172745	CTCGGGGCATCACCATCAAT	CCACCACCAGGATACAGCC
Dguok	NM_001162521	CGTCTCCGAGCGTCTTTCC	CACAGCGATGTTGCCTTCA
Tk2	NM_021028	AGCAGTGGTTTGTATTGAGGG	ACATGAGGCTCAGAGGGTTATG
Rrm2b	NM_199476	GAGCCACTCCTAAGAAAGAGTTC	GAGGGAGGTCCTTTGACAAGT
Sucla2	NM_011506	ACCCTTTCGCTGCATGAATAC	CCTGTGCCTTTATCACAACATCC

GENE SYMBOLS	NCBI Accession number (mouse)	Forward Primer (5' to 3')	Reverse Primer (5' to 3')
Suclg1	NM_019879	GGATACGACACGGGTCTTACA	GTGGTTCCTCCCACGAGTTTG
Dbt	NM_010022	AGACTGACCTGTGTTCGCTAT	GAGTGACGTGGCTGACTGTA
Hadha	NM_178878	TGCATTTGCCGCAGCTTTAC	GTTGGCCCAGATTTCGTTCA
Opa1	NM_001199177	AACAGCATTTCGAGCAACAGA	CTTCCGCAGCTCTTTGTTCTC
mMito DNA	NC_005089.1	CTAGAAACCCCGAAACCAAA	CCAGCTATCACCAAGCTCGT
mB2M DNA	NC_000068.7	ATGGGAAGCCGAACATACTG	CAGTCTCAGTGGGGGTGAAT

NRF1 mutant clones: Mutagenesis primers

EGFP-tagged expression clone	Forward Primer (5' to 3')	Reverse Primer (5' to 3')
NRF1-DIMdel	CAGGCTGTGGAGGTGGTG	ATTAGCATCTTGGACTCCAGTAAG
NRF1-NLSdel	GCCACGTTGGATGAGTACAC	TGTTGCCACAGCAGCAGC
NRF1-T201A	GGACAAAATGGCTCAGGCTCAGC	ACAGAGACTGGAATCCCATC
NRF1-T259A	GGTTTCATGGGCTCAGGCATTAC	CTTTGTTTTTGCTCTTCTGTG
NRF1-T259E	GGTTTCATGGGAGCAGGCATTACG	CTTTGTTTTTGCTCTTCTG
NRF1-T287A	AGATCAGCAAGCTCAAACTCAGG	TCAAAAGCATATAAAAGATCC

Software and Algorithms

GraphPad Prism 7

Molsoft ICM-Pro Browser

Pymol

Phyre2 http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index

GEO2R https://www.ncbi.nlm.nih.gov/geo/geo2r/

GSEA http://software.broadinstitute.org/gsea/index.jsp

GATHER http://changlab.uth.tmc.edu/gather/gather.py

MitoDB http://www.mitodb.com

MitoCarta2.0

https://www.broadinstitute.org/files/shared/metabolism/mitocarta/human.mitocarta2.0.html

PSCAN http://159.149.160.88/pscan/

ZDOCK http://zdock.umassmed.edu

Supplemental material

Fig. S1 shows impaired expression of OXPHOS genes in human A-T cerebellum. Fig. S2 shows loss of ATM impairs mitochondrial usage of pyruvate but enhanced lactate production in mouse neurons. Fig. S3 shows loss of ATM impairs ATP insufficiency response of human fibroblasts. Fig. S4 shows chronic insufficient ATP induces oxidative stress, activation of NRF1 and mitochondrial biogenesis. Table S1 shows raw data of multiple bioinformatics analyses.

Acknowledgments

This work was financially supported by Research Grants Council, HKSAR (HKUST12/CRF/13G, GRF660813, GRF16101315, GRF16103317 and AoE/M-05/12), the Offices of Provost, VPRG and Dean of Science, HKUST (VPRGO12SC02), and an RGC/HKUST initiation Grant (IGN16SC02), grant C6030-14E, and HKUST Institute for Advanced Study Junior Fellowship and Alzheimer's Association Research Fellowship (AARF-17-531566) to HMC.

Author Contributions

HMC designed the experiments, performed the work, analyzed the data and wrote the manuscript. AC, XS and MS performed essential work, helped in the analysis of the data and helped in the writing of the experiments. RH and KH designed experiments, analyzed data and wrote the manuscript.

Declaration of Interests

All authors declare no conflict of interests relate to the description and publication of the work described.

References

Abramov, A.Y., T.K. Smulders-Srinivasan, D.M. Kirby, R. Acin-Perez, J.A. Enriquez, R.N. Lightowlers, M.R. Duchen, and D.M. Turnbull. 2010. Mechanism of neurodegeneration of neurons with mitochondrial DNA mutations. *Brain*. 133:797-807.

Aird, K.M., A.J. Worth, N.W. Snyder, J.V. Lee, S. Sivanand, Q. Liu, I.A. Blair, K.E. Wellen, and R. Zhang. 2015. ATM couples replication stress and metabolic reprogramming during cellular senescence. *Cell Rep*. 11:893-901.

Ashley, C.N., K.D. Hoang, D.R. Lynch, S.L. Perlman, and B.L. Maria. 2012. Childhood ataxia: clinical features, pathogenesis, key unanswered questions, and future directions. *J Child Neurol*. 27:1095-1120.

Bargiela, D., P. Shanmugarajah, C. Lo, E.L. Blakely, R.W. Taylor, R. Horvath, S. Wharton, P.F. Chinnery, and M. Hadjivassiliou. 2015. Mitochondrial pathology in progressive cerebellar ataxia. *Cerebellum Ataxias*. 2:16.

Bratic, I., and A. Trifunovic. 2010. Mitochondrial energy metabolism and ageing. *Biochim Biophys Acta*. 1797:961-967.

Calvo, S.E., K.R. Clauser, and V.K. Mootha. 2016. MitoCarta2.0: an updated inventory of mammalian mitochondrial proteins. *Nucleic Acids Res*. 44:D1251-1257.

Carter, B.C., and B.P. Bean. 2009. Sodium entry during action potentials of mammalian neurons: incomplete inactivation and reduced metabolic efficiency in fast-spiking neurons. *Neuron*. 64:898-909.

Chang, J.T., and J.R. Nevins. 2006. GATHER: a systems approach to interpreting genomic signatures. *Bioinformatics*. 22:2926-2933.

Chao, M.W., C.P. Chen, Y.H. Yang, Y.C. Chuang, T.Y. Chu, and C.Y. Tseng. 2016. N-acetylcysteine attenuates lipopolysaccharide-induced impairment in lamination of Ctip2-and Tbr1-expressing cortical neurons in the developing rat fetal brain. *Scientific reports*. 6:32373.

Chen, H., J.M. McCaffery, and D.C. Chan. 2007. Mitochondrial fusion protects against neurodegeneration in the cerebellum. *Cell*. 130:548-562.

Chen, J., Y. Chen, G. Vail, H. Chow, Y. Zhang, L. Louie, J. Li, R.P. Hart, M.R. Plummer, and K. Herrup. 2016a. The impact of glutamine supplementation on the symptoms of ataxia-telangiectasia: a preclinical assessment. *Mol Neurodegener*. 11:60.

Chen, W., and M. Gueron. 1992. The inhibition of bovine heart hexokinase by 2-deoxy-D-glucose-6-phosphate: characterization by 31P NMR and metabolic implications. *Biochimie*. 74:867-873.

Chen, W.W., E. Freinkman, T. Wang, K. Birsoy, and D.M. Sabatini. 2016b. Absolute Quantification of Matrix Metabolites Reveals the Dynamics of Mitochondrial Metabolism. *Cell*. 166:1324-1337 e1311.

Chen, Y., H.G. Shertzer, S.N. Schneider, D.W. Nebert, and T.P. Dalton. 2005. Glutamate cysteine ligase catalysis: dependence on ATP and modifier subunit for regulation of tissue glutathione levels. *J Biol Chem*. 280:33766-33774.

Cheng, A., T. Zhao, K.H. Tse, H.M. Chow, Y. Cui, L. Jiang, S. Du, M.M.T. Loy, and K. Herrup. 2018. ATM and ATR play complementary roles in the behavior of excitatory and inhibitory vesicle populations. *Proc Natl Acad Sci U S A*. 115:E292-E301.

Choi, H.B., G.R. Gordon, N. Zhou, C. Tai, R.L. Rungta, J. Martinez, T.A. Milner, J.K. Ryu, J.G. McLarnon, M. Tresguerres, L.R. Levin, J. Buck, and B.A. MacVicar. 2012. Metabolic communication between astrocytes and neurons via bicarbonate-responsive soluble adenylyl cyclase. *Neuron*. 75:1094-1104.

Chow, H.M., D. Guo, J.C. Zhou, G.Y. Zhang, H.F. Li, K. Herrup, and J. Zhang. 2014. CDK5 activator protein p25 preferentially binds and activates GSK3beta. *Proc Natl Acad Sci U S A*. 111:E4887-4895.

Cribbs, A.P., A. Kennedy, B. Gregory, and F.M. Brennan. 2013. Simplified production and concentration of lentiviral vectors to achieve high transduction in primary human T cells. *BMC Biotechnol*. 13:98.

Dimauro, S., and P. Rustin. 2009. A critical approach to the therapy of mitochondrial respiratory chain and oxidative phosphorylation diseases. *Biochim Biophys Acta*. 1792:1159-1167.

Evans, M.J., and R.C. Scarpulla. 1990. NRF-1: a trans-activator of nuclear-encoded respiratory genes in animal cells. *Genes Dev*. 4:1023-1034.

Fang, E.F., H. Kassahun, D.L. Croteau, M. Scheibye-Knudsen, K. Marosi, H. Lu, R.A. Shamanna, S. Kalyanasundaram, R.C. Bollineni, M.A. Wilson, W.B. Iser, B.N. Wollman, M. Morevati, J. Li, J.S. Kerr, Q. Lu, T.B. Waltz, J. Tian, D.A. Sinclair, M.P. Mattson, H. Nilsen, and V.A. Bohr. 2016. NAD+ Replenishment Improves Lifespan and Healthspan in Ataxia Telangiectasia Models via Mitophagy and DNA Repair. *Cell Metab*. 24:566-581.

Fang, E.F., M. Scheibye-Knudsen, L.E. Brace, H. Kassahun, T. SenGupta, H. Nilsen, J.R. Mitchell, D.L. Croteau, and V.A. Bohr. 2014. Defective mitophagy in XPA via PARP-1 hyperactivation and NAD(+)/SIRT1 reduction. *Cell*. 157:882-896.

Garcia-Gutierrez, P., M. Mundi, and M. Garcia-Dominguez. 2012. Association of bromodomain BET proteins with chromatin requires dimerization through the conserved motif B. *J Cell Sci*. 125:3671-3680.

Gugneja, S., and R.C. Scarpulla. 1997. Serine phosphorylation within a concise amino-terminal domain in nuclear respiratory factor 1 enhances DNA binding. *J Biol Chem*. 272:18732-18739.

Guo, Z., S. Kozlov, M.F. Lavin, M.D. Person, and T.T. Paull. 2010. ATM activation by oxidative stress. *Science*. 330:517-521.

Hall, C.N., M.C. Klein-Flugge, C. Howarth, and D. Attwell. 2012. Oxidative phosphorylation, not glycolysis, powers presynaptic and postsynaptic mechanisms underlying brain information processing. *J Neurosci*. 32:8940-8951.

Hanson, G.T., R. Aggeler, D. Oglesbee, M. Cannon, R.A. Capaldi, R.Y. Tsien, and S.J. Remington. 2004. Investigating mitochondrial redox potential with redox-sensitive green fluorescent protein indicators. *J Biol Chem*. 279:13044-13053.

Hargreaves, I.P., Y. Sheena, J.M. Land, and S.J. Heales. 2005. Glutathione deficiency in patients with mitochondrial disease: implications for pathogenesis and treatment. *J Inherit Metab Dis*. 28:81-88.

Herzig, S., and R.J. Shaw. 2018. AMPK: guardian of metabolism and mitochondrial homeostasis. *Nat Rev Mol Cell Biol*. 19:121-135.

Hildyard, J.C., C. Ammala, I.D. Dukes, S.A. Thomson, and A.P. Halestrap. 2005. Identification and characterisation of a new class of highly specific and potent inhibitors of the mitochondrial pyruvate carrier. *Biochim Biophys Acta*. 1707:221-230.

Hodneland Nilsson, L.I., I.K. Nitschke Pettersen, J. Nikolaisen, D. Micklem, H. Avsnes Dale, G. Vatne Rosland, J. Lorens, and K.J. Tronstad. 2015. A new live-cell reporter strategy to simultaneously monitor mitochondrial biogenesis and morphology. *Scientific reports*. 5:17217.

Hornbeck, P.V., B. Zhang, B. Murray, J.M. Kornhauser, V. Latham, and E. Skrzypek. 2015. PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic Acids Res*. 43:D512-520.

Howarth, C., P. Gleeson, and D. Attwell. 2012. Updated energy budgets for neural computation in the neocortex and cerebellum. *J Cereb Blood Flow Metab*. 32:1222-1232.

Howarth, C., C.M. Peppiatt-Wildman, and D. Attwell. 2010. The energy use associated with neural computation in the cerebellum. *J Cereb Blood Flow Metab*. 30:403-414.

Huang da, W., B.T. Sherman, and R.A. Lempicki. 2009. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*. 4:44-57.

Jiang, D., Y. Zhang, R.P. Hart, J. Chen, K. Herrup, and J. Li. 2015. Alteration in 5-hydroxymethylcytosine-mediated epigenetic regulation leads to Purkinje cell vulnerability in ATM deficiency. *Brain*. 138:3520-3536.

Joshi, D.C., and J.C. Bakowska. 2011. Determination of mitochondrial membrane potential and reactive oxygen species in live rat cortical neurons. *J Vis Exp*.

Jourdain, P., I. Allaman, K. Rothenfusser, H. Fiumelli, P. Marquet, and P.J. Magistretti. 2016. L-Lactate protects neurons against excitotoxicity: implication of an ATP-mediated signaling cascade. *Scientific reports*. 6:21250.

Kastan, M.B., and D.S. Lim. 2000. The many substrates and functions of ATM. *Nat Rev Mol Cell Biol*. 1:179-186.

Kelley, L.A., S. Mezulis, C.M. Yates, M.N. Wass, and M.J. Sternberg. 2015. The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc*. 10:845-858.

Kelly, D.P., and R.C. Scarpulla. 2004. Transcriptional regulatory circuits controlling mitochondrial biogenesis and function. *Gene Dev*. 18:357-368.

Kobayashi, A., T. Tsukide, T. Miyasaka, T. Morita, T. Mizoroki, Y. Saito, Y. Ihara, A. Takashima, N. Noguchi, A. Fukamizu, Y. Hirotsu, M. Ohtsuji, F. Katsuoka, and M. Yamamoto. 2011. Central nervous system-specific deletion of transcription factor Nrf1 causes progressive motor neuronal dysfunction. *Genes Cells*. 16:692-703.

Koopman, W.J., F. Distelmaier, J.A. Smeitink, and P.H. Willems. 2013. OXPHOS mutations and neurodegeneration. *EMBO J*. 32:9-29.

Lax, N.Z., P.D. Hepplewhite, A.K. Reeve, V. Nesbitt, R. McFarland, E. Jaros, R.W. Taylor, and D.M. Turnbull. 2012. Cerebellar ataxia in patients with mitochondrial DNA disease: a molecular clinicopathological study. *J Neuropathol Exp Neurol*. 71:148-161.

Le Masson, G., S. Przedborski, and L.F. Abbott. 2014. A computational model of motor neuron degeneration. *Neuron*. 83:975-988.

Li, J., R.P. Hart, E.M. Mallimo, M.R. Swerdel, A.W. Kusnecov, and K. Herrup. 2013. EZH2-mediated H3K27 trimethylation mediates neurodegeneration in ataxia-telangiectasia. *Nature neuroscience*. 16:1745-1753.

Lopez, M.F., B.S. Kristal, E. Chernokalskaya, A. Lazarev, A.I. Shestopalov, A. Bogdanova, and M. Robinson. 2000. High-throughput profiling of the mitochondrial proteome using affinity fractionation and automation. *Electrophoresis*. 21:3427-3440.

Malik, A.N., A. Czajka, and P. Cunningham. 2016. Accurate quantification of mouse mitochondrial DNA without co-amplification of nuclear mitochondrial insertion sequences. *Mitochondrion*. 29:59-64.

Malik, A.N., R. Shahni, A. Rodriguez-de-Ledesma, A. Laftah, and P. Cunningham. 2011. Mitochondrial DNA as a non-invasive biomarker: accurate quantification using real time quantitative PCR without co-amplification of pseudogenes and dilution bias. *Biochem Biophys Res Commun*. 412:1-7.

McCommis, K.S., and B.N. Finck. 2015. Mitochondrial pyruvate transport: a historical perspective and future research directions. *Biochem J*. 466:443-454.

McKenzie, M., D. Liolitsa, N. Akinshina, M. Campanella, S. Sisodiya, I. Hargreaves, N. Nirmalananthan, M.G. Sweeney, P.M. Abou-Sleiman, N.W. Wood, M.G. Hanna, and M.R.

Duchen. 2007. Mitochondrial ND5 gene variation associated with encephalomyopathy and mitochondrial ATP consumption. *J Biol Chem*. 282:36845-36852.

Morita, A., K. Tanimoto, T. Murakami, T. Morinaga, and Y. Hosoi. 2014. Mitochondria are required for ATM activation by extranuclear oxidative stress in cultured human hepatoblastoma cell line Hep G2 cells. *Biochem Biophys Res Commun*. 443:1286-1290.

Nicholls, D.G., V.M. Darley-Usmar, M. Wu, P.B. Jensen, G.W. Rogers, and D.A. Ferrick. 2010. Bioenergetic profile experiment using C2C12 myoblast cells. *J Vis Exp*.

Novoa, W.B., A.D. Winer, A.J. Glaid, and G.W. Schwert. 1959. Lactic dehydrogenase. V. Inhibition by oxamate and by oxalate. *J Biol Chem*. 234:1143-1148.

Owen, O.E., S.C. Kalhan, and R.W. Hanson. 2002. The key role of anaplerosis and cataplerosis for citric acid cycle function. *J Biol Chem*. 277:30409-30412.

Patel, A.Y., T.M. McDonald, L.D. Spears, J.K. Ching, and J.S. Fisher. 2011. Ataxia telangiectasia mutated influences cytochrome c oxidase activity. *Biochem Biophys Res Commun*. 405:599-603.

Pierce, B.G., K. Wiehe, H. Hwang, B.H. Kim, T. Vreven, and Z. Weng. 2014. ZDOCK server: interactive docking prediction of protein-protein complexes and symmetric multimers. *Bioinformatics*. 30:1771-1773.

San Martin, A., S. Ceballo, F. Baeza-Lehnert, R. Lerchundi, R. Valdebenito, Y. Contreras-Baeza, K. Alegria, and L.F. Barros. 2014. Imaging mitochondrial flux in single cells with a FRET sensor for pyruvate. *PLoS One*. 9:e85780.

San Martin, A., S. Ceballo, I. Ruminot, R. Lerchundi, W.B. Frommer, and L.F. Barros. 2013. A genetically encoded FRET lactate sensor and its use to detect the Warburg effect in single cancer cells. *PLoS One*. 8:e57712.

Saraiva, L.M., G.S. Seixas da Silva, A. Galina, W.S. da-Silva, W.L. Klein, S.T. Ferreira, and F.G. De Felice. 2010. Amyloid-beta triggers the release of neuronal hexokinase 1 from mitochondria. *PLoS One*. 5:e15230.

Satoh, J., N. Kawana, and Y. Yamamoto. 2013. Pathway Analysis of ChIP-Seq-Based NRF1 Target Genes Suggests a Logical Hypothesis of their Involvement in the Pathogenesis of Neurodegenerative Diseases. *Gene Regul Syst Bio*. 7:139-152.

Scaglia, F., L.J. Wong, G.D. Vladutiu, and J.V. Hunter. 2005. Predominant cerebellar volume loss as a neuroradiologic feature of pediatric respiratory chain defects. *AJNR Am J Neuroradiol*. 26:1675-1680.

Scarpulla, R.C. 2002. Nuclear activators and coactivators in mammalian mitochondrial biogenesis. *Biochim Biophys Acta*. 1576:1-14.

Scarpulla, R.C. 2008. Transcriptional paradigms in mammalian mitochondrial biogenesis and function. *Physiol Rev*. 88:611-638.

Scheibye-Knudsen, M., K. Scheibye-Alsing, C. Canugovi, D.L. Croteau, and V.A. Bohr. 2013. A novel diagnostic tool reveals mitochondrial pathology in human diseases and aging. *Aging (Albany NY)*. 5:192-208.

Schutt, F., S. Aretz, G.U. Auffarth, and J. Kopitz. 2012. Moderately reduced ATP levels promote oxidative stress and debilitate autophagic and phagocytic capacities in human RPE cells. *Invest Ophthalmol Vis Sci*. 53:5354-5361.

Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, A. Paulovich, S.L. Pomeroy, T.R. Golub, E.S. Lander, and J.P. Mesirov. 2005. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 102:15545-15550.

Sugimori, M., and R.R. Llinas. 1990. Real-time imaging of calcium influx in mammalian cerebellar Purkinje cells in vitro. *Proc Natl Acad Sci U S A*. 87:5084-5088.

Tantama, M., J.R. Martinez-Francois, R. Mongeon, and G. Yellen. 2013. Imaging energy status in live cells with a fluorescent biosensor of the intracellular ATP-to-ADP ratio. *Nature communications*. 4:2550.

Valentin-Vega, Y.A., K.H. Maclean, J. Tait-Mulder, S. Milasta, M. Steeves, F.C. Dorsey, J.L. Cleveland, D.R. Green, and M.B. Kastan. 2012. Mitochondrial dysfunction in ataxia-telangiectasia. *Blood*. 119:1490-1500.

Wang, Z.J., C.L. Liang, G.M. Li, C.Y. Yu, and M. Yin. 2007. Stearic acid protects primary cultured cortical neurons against oxidative stress. *Acta Pharmacol Sin*. 28:315-326.

Watters, D.J. 2003. Oxidative stress in ataxia telangiectasia. *Redox Rep*. 8:23-29.

Waypa, G.B., J.D. Marks, R. Guzy, P.T. Mungai, J. Schriewer, D. Dokic, and P.T. Schumacker. 2010. Hypoxia triggers subcellular compartmental redox signaling in vascular smooth muscle cells. *Circ Res*. 106:526-535.

Welsh, J.P., G. Yuen, D.G. Placantonakis, T.Q. Vu, F. Haiss, E. O'Hearn, M.E. Molliver, and S.A. Aicher. 2002. Why do Purkinje cells die so easily after global brain ischemia? Aldolase C, EAAT4, and the cerebellar contribution to posthypoxic myoclonus. *Adv Neurol*. 89:331-359.

Xu, Y., T. Ashley, E.E. Brainerd, R.T. Bronson, M.S. Meyn, and D. Baltimore. 1996. Targeted disruption of ATM leads to growth retardation, chromosomal fragmentation during meiosis, immune defects, and thymic lymphoma. *Genes Dev*. 10:2411-2422.

Yan, Y., C.L. Wei, W.R. Zhang, H.P. Cheng, and J. Liu. 2006. Cross-talk between calcium and reactive oxygen species signaling. *Acta Pharmacol Sin*. 27:821-826.

Zakikhani, M., M. Bazile, S. Hashemi, S. Javeshghani, D. Avizonis, J. St Pierre, and M.N. Pollak. 2012. Alterations in cellular energy metabolism associated with the antiproliferative effects of the ATM inhibitor KU-55933 and with metformin. *PLoS One*. 7:e49513.

Zambelli, F., G. Pesole, and G. Pavesi. 2009. Pscan: finding over-represented transcription factor binding site motifs in sequences from co-regulated or co-expressed genes. *Nucleic Acids Res.* 37:W247-252.

Figure Legends

Figure 1*.* **The neurological symptoms from A-T associated with mitochondrial diseases**

(A) A summary of mitochondrial and non-mitochondrial disease counts by classes of A-T clinical symptoms using MitoDB. Histograms indicate the number of mitochondrial or non-mitochondrial diseases associated with each A-T symptom and the percentages of A-T patients associated with each symptom are listed at bottom. (B) A list of nine common mitochondrial diseases associated with symptoms of ataxia and cerebellar atrophy. Eight out of nine are mitochondrial diseases with functional defects in the OXPHOS system.

Figure 2. **Loss of metabolic function in mitochondria of A-T patients**

(A) Volcano plot of differentially expressed genes in cerebellar samples of A-T patients (n=3) versus age-matched controls (n=4). The $-\log_{10}$[adjusted p-value (p-value)] was plotted against $\log_2$[Fold Change (FC)]. Genes with significant differences in expression (adjusted p-value<0.05) are above the dotted line. (B and C) Gene Ontology (GO) analysis of differentially expressed genes (Table S1D). (B) Thirty GO terms with p-value <0.0001 were plotted against the corresponding Bayes factor (BF). Enrichment ranking of pathways with significance metrics above the recommended threshold [P<0.0001 and BF>6 (as suggested by the authors)] was performed (C). Pathways significantly enriched, determined by the relative number of genes associated within a GO category. (D and E) Gene Set Enrichment Analysis (GSEA) of significantly different genes, among significant differences were pathways of (D) oxidative phosphorylation and the (E) TCA cycle. The leading edge (most significant genes), shown as vertical bars, accumulated either left and below the peak of the enrichment score plot (green) or right of the valley, indicating the up- or down-regulated genes, respectively. Genes with the highest enrichment score is shown as a gene expression-based heatmap and indicates relatively reduced gene expression (blue) in A-T samples. (F) Heatmap summarizing 1,116 nuclear genes encoding proteins with a high likelihood of mitochondrial localization segregated by function. (G) Significantly changed genes were mostly diminished in A-T (blue gene symbols). Gene symbols marked in red are relatively up-regulated. Note: 42 genes from the original MitoCarta2.0 list failed to match probes from the human expression array.

***Figure 3*. Loss of *Atm* impairs the function of OXPHOS complexes in mouse neurons**

(A) Diagram indicating the assembly of the functional OXPHOS system. (B-F) Heatmaps indicating the general expression trends of 79 nuclear-encoded genes whose loss of function results in neurodegeneration. Each group represents the mean±SEM (see also Fig.S1B-1F; *P<0.05, **P<0.01, ***P<0.001* and *ns*=non-significant, unpaired t-test). (G-K) The activity of each mitochondrial complexes in *Atm*$^{-/-}$ cultured mouse cortical neurons. Each group represents the mean±SEM (Complexes I, III, IV and V: N=30; Complex II: N=12) (****P<0.001* and *ns*=non-significant, unpaired t-test).

***Figure 4*. Loss of *Atm* impairs the TCA cycle and favors the conversion of pyruvate to lactate over acetyl-CoA**

(A) Diagram of the TCA cycle. Enzymes up-regulated in A-T are shown in blue (adj. p>0.05); down-regulated enzymes are shown in red. Those without significant changes are shown in black. (B) Inhibition of mitochondrial usage of pyruvate by 1μM UK5099 or of its cytosolic conversion to lactate by 2.5mM oxamate revealed that the latter reaction is favored in A-T fibroblasts. In addition, impaired glycolysis was also observed in A-T patient fibroblast upon treatment with 30mM 2-DG. Affected pathways are diagramed at left (n=18, ****P<0.001*, One-way ANOVA).(C) Inhibition of lactate production by 2.5mM oxamate or of lactate release from the cytosol to the extracellular space by 500 μM pCMBS confirmed that pyruvate conversion to lactate is enhanced in A-T cells. Pathways involved are diagrammed at left (n=18, ****P<0.001*, One-way ANOVA). (D) Changes in levels of isocitrate and succinate in mitochondria in normal or A-T human fibroblast (n=18, ****P<0.001, ns=non-significant*, unpaired t-test). (E) Seahorse Mito Fuel Flex test to evaluate cellular respiration of normal and A-T human fibroblast in conditions when specific metabolic pathway of glucose, glutamine or fatty acids was inhibited. The relative fuel dependence was calculated. Comparison was made to each fuel source between the normal and patient groups (n=3, ***P<0.001, **P<0.01, ns=non-significant, unpaired t-test).

***Figure 5*. Loss of *Atm* impairs cellular response towards ATP insufficiency in neurons**

(A) Static and (B) dynamic changes in mitochondrial membrane potential in primary cortical neurons were evaluated with TMRM, and counterstained with Mitotracker Green (n=15). (C) The relative changes in TMRM signals, expressed as the ratio of post-oligomycin (85 sec) to pre-oligomycin (35 sec) OCR values (n=15, ***P<0.001, unpaired t-test). (D) Seahorse Mito Stress assay of cellular respiration of neurons isolated from $Atm^{+/+}$ and $Atm^{-/-}$ mice (n=16, ***P<0.001, two-way ANOVA). (E-F) Static and dynamic changes in ATP levels in $Atm^{+/+}$ and $Atm^{-/-}$ primary cortical neurons measured by the Perceval HR reporter. (E) Equilibrium ATP levels at t=75 min (n=20, ***P<0.001, unpaired t-test). (F) Live-imaging of cultured neuron responses to synaptic activity-dependent energy consumption. After establishing a steady baseline, 50 µM glutamate was added for 30 seconds to excite the neurons and cause ATP to be consumed; glutamate was then washed away and intracellular ATP levels allowed to recover (n=30, ***P<0.001, two-way ANOVA). (G) The relative abundance of ATP immediately following the removal of glutamate (Signal ratio: Post-$E_{2135sec}$/Pre-$E_{1993sec}$, left panel) and after recovery (Signal ratio: Endpoint$_{4270sec}$/Post-$E_{2135sec}$, right panel) (n=30, ***P<0.05, ns=non-significant, unpaired t-test). (H) Representative images of the live/dead status of neurons four hours after Perceval HR assay. (Live cell: Green; Dead cell: red). Quantification shown in right panel (n=25, ***P<0.001, unpaired t-test).

Figure 6. Nuclear respiratory factor-1 (NRF1) links ATM activity and mitochondrial function
(A) The top 10 over-represented transcription factors predicted by the PSCAN platform to bind promoters of nuclear-encoded mitochondrial genes, ranked according to their z-test P-value. (B) The consensus NRF1 binding site for these transcription factors. (C) Top panel shows the positions of the best occurrences—location of the best predicted NRF1 binding motifs within the -450 to +50 bp region on the tested promoters with corresponding matching scores (the value representing the likelihood for NRF1 to bind to the promoter) higher than the average genome-wide matching score 0.843. Bottom panel shows the distribution of occurrence scores for tested promoters, with an average matching score values of all predictions equaling 0.862, higher than the average genome-wide matching score 0.843. These, together with the z-test P-value shown in Fig.6A (NRF1 = 8.242 x 10^{-114}) imply that the NRF1 binding motif is non-randomly

enriched in the tested set of promoters, in which the gene expression level was significantly altered in A-T cerebella samples. Predictions on each promoter were colored as straight lines according to their corresponding matching score (red=high; yellow=medium; green=low). (D) Volcano plot showing the relative expression levels of genes predicted to be regulated by NRF1. (E) Venn diagram illustrating the overlap between the set of genes computationally identified as NRF1 targets and the combined sets of observed NRF1 targets found in the ENCODE database and a published NRF1-Chip sequencing study. (F) A table showing the four KEGG pathways identified using genes whose promoters scored as one of the 400 top hits for likelihood as an NRF1 promoter. Genes associated in the pathways as well as the corresponding p-values are shown. (G) Representative images of NRF1 immunostaining in mouse (upper panels) and human (lower panels) cerebellar tissue samples from wild type (left) and ATM-deficient (right) subjects. Quantification of Purkinje cells with cytoplasmic and nuclear NRF1 in cells are shown on the right (n=15, ***$P<0.001$, one-way ANOVA).

Figure 7. ATP depletion activates ATM with downstream effects on NRF1

(A) DIV14 Primary cortical neurons isolated from wild type mice were treated with vehicle, 1 μM etoposide, 0.5 μM oligomycin or 1 μM atractyloside for 24 hours. The number of γH2AX and 53BP1 foci in MAP2$^+$ neurons were then quantified (n=20, ***$P<0.001$, one-way ANOVA). (B) Immunocytochemistry for activated ATM [pATM(S1981)], NRF1 and MAP2. $Atm^{+/+}$ and $Atm^{-/-}$ DIV14 primary cortical neurons treated with either vehicle, 1 μM etoposide, 10 nM $H_2O_2,$ or 1 μM atractyloside in the presence or absence of 5 mM N-acetylcysteine (NAC) for 24 hours. (C) Quantification is shown (n=9, ***$P<0.001$, ns=non-significant, one-way ANOVA). (D) NRF1-mediated transcription activities of wild type DIV14 primary cortical neurons treated with either vehicle or 1μM atractyloside in the presence or absence of 5mM NAC for 24 hours using the NRF1mitoGFP reporter. Intracellular GFP signals resulting from NRF1 transcription activity were monitored for 24 hours after initiation of drug treatments. The ratios of cellular GFP intensities at 24-hour over 0-hour time points are shown. Representative images of the reporter signals at 0- and 24-hour time points can be found in Fig.S4D.

Figure 8. ATM-mediated phosphorylation of the NRF1 T259 residue facilitates homodimerization and nuclear localization

(A) *NRF1* gene expression in mouse cerebellum (n=28-30, n=ns, unpaired t-test). (B) Representative Western blots (left panel) and quantification (right panel) of monomeric and dimeric NRF1 and its target, succinate dehydrogenase in mouse cerebellum (n=4, ****P<0.001*, unpaired t-test). (C) NRF1 protein sequence alignment, performed on the Clustal Omega platform. (D-F) Representative Western blots of immunoprecipitates (IPs) showing interactions between NRF1 and ATM. (D) Reciprocal IPs of EGFP-NRF1 and FLAG-ATM-WT; (E) S/TQ phosphorylation (top) and dimerization of EGFP-NRF1 mediated by wild type (WT) but not kinase dead (KD) ATM (F) Inhibition of ATM, but not ATR or DNA-PKcs, blocks S/TQ phosphorylation of EGFP-NRF1 in mouse embryonic fibroblasts. (G) Western blot images showing that both the nuclear localization of NRF1 and its phosphorylation on T259 are required for NRF1 homodimerization (n=4). (H) NRF1 (aa198-271) structure is predicted to adopt a bromodomain configuration in the region of its DNA-binding and dimerization domain and (I) superimpose of BRD2/4 bromodomains. (J) *in silico* site-directed mutagenesis was performed to obtain the T259(PO4⁻) structure followed by homodimer docking analyses using the ZDOCK platform. Predicted homodimerized best-ranking NRF1 bromodomain structures were obtained and contact area analysis was performed using the ICM-browser pro. Amino acid residues that showed significant interactions between the monomers were ranked according to the percentage of contacting area over total exposed area. (K) Western blots of tagged-NRF1 co-immunoprecipitations.. Quantification of the percentage of dimerized NRF1 is shown (n=6, ****P<0.001*, one-way ANOVA). (L) Nuclear versus cytoplasmic localization of NRF1 mutants in DIV12 primary cortical neurons as shown in a heatmap. (M) Distribution of EGFP-NRF1-WT proteins in neurons treated with KU60019 (n=10, ****P<0.001*, one-way *ANOVA*). (D) Histograms showing ATM requirements of individual Purkinje (red), cerebellar granule (blue) or cortical pyramidal (black) neurons. Individual panels show energy consumption by specific subcellular processes as marked. Parameters and calculations as per Howarth et al (2012), details can be found in Table S1O.

A

Mitochondrial disease (red) | Non-mitochondrial disease (blue)

Lymphatic

	Immune deficiency	Cancer
Red	0	2
Blue	3	13
% Afflicted	61	7

Nervous system (Motor function)

	Ataxia	Chorea	Athetosis	Areflexia	Systonia
Red	27	4	2	10	7
Blue	19	7	2	7	6
% Afflicted	100	94	81	79	79

Nervous system (Eye movement)

	Dysarthria	Oculomotor apraxia	Nystagmus	Strabismus
Red	8	0	12	4
Blue	12	2	16	5
% Afflicted	100	100	59	25

Nervous system (Degeneration and stunted growth)

	Cerebellar atrophy	Demyelination	Neuropathy	Short statue
Red	11	0	17	13
Blue	16	3	9	28
% Afflicted	100	100	71	65

Integumentary

	Radiosensitivity	Telangiectasia
Red	0	0
Blue	1	5
% Afflicted	100	97

Miscellaneous

	Increased blood AFP	Weight loss
Red	0	4
Blue	1	7
% Afflicted	90	77

B

	Mitochondrial disease	Defective gene(s)	Mitochondrial function defect
1	MERRF	Multiple MT-genes	Multiple defects in respiratory chains enzyme activities
2	NARP	*MTATP6*	H(+)-ATPase
3	Co-Q10 deficiency	*COQ10D1±COQ2*	Compromised complex I-III and complex II-III activities
4	IOSCA	*C10orf2*	Deficiency of complexes I and IV
5	SDHA deficiency	*SDHA*	Deficiency of complex II
6	MTDPS11	*MGME1*	Multiple defects in respiratory chain enzyme activities
7	SCA28	*AFG3L2*	Multiple defects in respiratory chain enzyme activities
8	SPG7	*SPG7*	Deficiency of complex IV
9	LBSL	*DARS2*	N/A

A

	Matrix ID	Matrix name	P-value
1	MA0506.1	NRF1	$8.242e^{-114}$
2	MA0516.1	SP2	$5.693e^{-107}$
3	MA0470.1	E2F4	$3.033e^{-101}$
4	MA0162.2	EGR1	$8.754e^{-93}$
5	MA0024.2	E2F1	$7.861e^{-91}$
6	MA0079.3	SP1	$6.302e^{-90}$
7	MA0079.2	SP1	$5.085e^{-81}$
8	MA0527.1	ZBTB33	$1.208e^{-80}$
9	MA0469.1	E2F3	$9.384e^{-80}$
10	MA0162.1	EGR1	$2.581e^{-77}$

B NRF1 consensus binding motif

C Occurances-Position Distribution (Score≥ 0.843)

Occurances-Score Distribution (Average score = 0.862)

D -log₁₀(p-value)

2778 genes   582 genes

log₂FC

E
- ENCODE and NRF1 CHIP-seq
- PSCAN NRF1 targets (Score ≥ 0.85)
- Common hits

2030   440 ↑47 ↓297   2920

F

Rank	Category	*Genes in the pathway	P-value
1	Cell cycle	*ATM, CCNE1, CDC7, HDAC2, HDAC4, MCM4, MCM6, TFDP1*	0.002
2	Wnt signaling pathway	*APC, FZD4, PPP2CA, PPP2R1A, SENP2*	0.01
3	Oxidative phosphoyrlation	*ATP6V1C1, ATP6V1F, LOC308770, SDHA, UQCRB*	0.02
4	Circadian rhythm	*CLOCK, CRY1*	0.04

All genes listed had reduced expression in A-T

G

Mouse: $Atm^{+/+}$ — 200µm; $Atm^{-/-}$ — 200µm

Human: Normal — 200µm; A-T — 200µm

- $Atm^{+/+}$ - $Atm^{-/-}$
- Normal - A-T