Differential impact of endogenous and exogenous attention on activity in human visual cortex

Laura Dugué[1,2,3,4,5], Elisha P. Merriam[1,2,6], David J. Heeger[1,2] & Marisa Carrasco[1,2]

[1] Department of Psychology, New York University
[2] Center for Neural Science, New York University
[3] CNRS (Integrative Neuroscience and Cognition Center, UMR 8002) Paris, France
[4] Université de Paris, Paris, France
[5] Institut Universitaire de France (IUF), Paris, France
[6] Laboratory of Brain and Cognition, NIMH/NIH, Bethesda, MD

Running title: endogenous and exogenous attention

Corresponding author:

Laura Dugué

Current address: 45 rue des Saints-Pères 75006 Paris, FRANCE

laura.dugue@u-paris.fr

ABSTRACT

How do endogenous (voluntary) and exogenous (involuntary) attention modulate activity in visual cortex? Using ROI-based fMRI analysis, we measured fMRI activity for valid and invalid trials (target at cued/un-cued location, respectively), for pre- or post-cueing in endogenous and exogenous conditions, while observers performed the same task. We found stronger modulation in contralateral than ipsilateral visual regions to the attended hemifield, and higher activity in the valid- than invalid-trials. For endogenous, modulation of stimulus-evoked activity due to a pre-cue increased along the visual hierarchy, but was constant due to a post-cue. In contrast, for exogenous, modulation of stimulus-evoked activity due to a pre-cue was constant along the visual hierarchy, but not modulated due to a post-cue. These findings reveal that endogenous and exogenous attention distinctly modulate activity in visual areas during orienting and reorienting of attention; endogenous facilitates both the encoding and the readout of visual information whereas exogenous only facilitates the encoding of information.

Keywords: exogenous attention, endogenous attention, fMRI activity, striate and extra-striate visual areas, top-down and bottom-up processing, visual perception.

INTRODUCTION

Spatial, covert visual attention is the selective processing of visual information in space, without change in gaze. Attention can be allocated voluntarily –endogenous attention– or involuntarily –exogenous attention. Endogenous and exogenous attention have different temporal dynamics; endogenous takes about 300 ms to be deployed and can be sustained at will whereas exogenous attention takes only about 100 ms to be deployed and it is transient (for review see Carrasco, 2011; Carrasco and Barbot, 2015). These two types of attention often have similar perceptual consequences (reviews by Carrasco, 2011; Carrasco and Barbot, 2015), but notable exceptions indicate that whereas endogenous attention acts in a flexible way, exogenous attention acts in an inflexible or automatic way. For instance: (a) the benefits and costs in perception (discriminability and speed of information accrual) scale with cue validity for endogenous but not for exogenous attention (e.g. Giordano et al., 2009); (b) the effects of covert attention on contrast sensitivity often differ for endogenous and exogenous attention (e.g. Dosher and Lu, 2000; Ling and Carrasco, 2006a; Pestilli et al., 2009; Barbot et al., 2012); and (c) for a texture segmentation task in which heightened spatial resolution improves or impairs performance as a function of target eccentricity, endogenous attention improves performance across eccentricity, whereas exogenous attention improves performance at peripheral locations where resolution is poor, but hampers performance where resolution is already high for the task at hand (e.g. Yeshurun and Carrasco, 1998; Carrasco et al., 2006; Yeshurun et al., 2008; Barbot and Carrasco, 2017; Jigo and Carrasco, 2018).

Basic visual processes, such as contrast sensitivity and spatial resolution, are mediated by activity in early visual cortex, and are altered by covert attention (reviews by Carrasco and Yeshurun, 2009; Carrasco, 2011; Anton-Erxleben and Carrasco, 2013; Carrasco and Barbot, 2015). Specifically, single-unit studies in monkeys has demonstrated effects of endogenous attention in occipital areas (e.g. McAdams and Maunsell, 1999; Reynolds et al., 2000; Martínez-Trujillo and Treue, 2002; Williford and Maunsell, 2006; Mitchell et al., 2009; Reynolds and Heeger, 2009; Ruff and Cohen, 2014; Luo and Maunsell, 2015). Additionally, fMRI studies have shown that endogenous attention causes a baseline shift in early visual areas (e.g. Kastner et

al., 1999; Somers et al., 1999; O'Connor et al., 2002; Buracas and Boynton, 2007; Murray, 2008; Herrmann et al., 2010; Pestilli et al., 2011; review by Beck and Kastner, 2014) and increases the dynamic range of fMRI responses (Li et al., 2008; Lu et al., 2011). Comparatively, little is known about the effect of exogenous attention on visual areas both from single-unit (Busse et al., 2008; Wang et al., 2015) and fMRI (Mayer et al., 2004; Peelen et al., 2004; Liu et al., 2005; Kincade et al., 2005; Natale et al., 2006; Müller and Kleinschmidt, 2007; Esterman et al., 2008; Müller and Ebeling, 2008; Heinen et al., 2011; Mulckhuyse et al., 2011) studies.

Since Corbetta and Schulman's seminal review (2002) on the neural bases of endogenous and exogenous attention, there has been emphasis on characterizing networks of brain regions within the frontal and parietal lobes (for reviews see Chica et al., 2013; Beck and Kastner, 2014). Yet, there remains considerable debate regarding the dissociation between dorsal regions, for endogenous attention, and ventral regions, for exogenous attention (e.g. Hahn et al., 2006; Bowling et al., 2019), including the role of the temporo-parietal junction (TPJ; Doricchi et al., 2010; Geng and Vossel, 2013; Silvetti et al., 2015; Dugué et al., 2017a). Classically, researchers have described endogenous attention as a top-down process, and exogenous attention as a bottom-up process (e.g., Posner et al., 1980; Nakayama and Mackeben, 1989; Corbetta and Shulman, 2002; Hahn et al., 2006; Beck and Kastner, 2009; Carrasco, 2011; Chica et al., 2013; Beck and Kastner, 2014; Bowling et al., 2019). This characterization originated in psychophysics experiments, and was then studied using fMRI, in which the two types of attention have been often investigated separately (for reviews Beck and Kastner, 2009; Carrasco, 2011; Chica et al., 2013; Beck and Kastner, 2014).

Studies of endogenous and exogenous attention focusing on parietal and frontal areas have shown that the two types of attention differentially modulate fronto-parietal connectivity (Buschman and Miller, 2009; Bowling et al., 2019). For example, there are critical differences in the temporal order of neural responses in frontal and parietal cortex in monkeys between these attention conditions, i.e. frontal activity precedes parietal activity during endogenous attentional orienting, whereas parietal activity precedes frontal activity during exogenous orienting. Critically, it is often assumed that the effects of endogenous and exogenous attention are the same in

striate and extra-striate areas (Corbetta and Shulman, 2002; Peelen et al., 2004; Corbetta et al., 2008; Chica et al., 2013; Beck and Kastner, 2014).

A number of important considerations limit the conclusions that may be drawn from the few studies that have directly compared independent effects of visual exogenous and endogenous spatial attention (see **Table 1**):

(1) The effect of attention on behavioral performance was assessed in some studies with a detection task using RT as their only dependent variable (Mayer et al., 2004; Peelen et al., 2004; Natale et al., 2006), in which performance may differ due to speed of processing, discriminability or criterion (Reed, 1973; Wickelgren, 1977; Carrasco and McElree, 2001) and/or motor behavior (Correa et al., 2010; Yashar and Lamy, 2011).

(2) In the studies in which performance was assessed in a discrimination task using RT (Kim et al., 1999; Kincade et al., 2005; Esterman et al., 2008), small RT differences were reported, which could have resulted from speed of processing, discriminability, or criterion factors (Reed, 1973; Wickelgren, 1977; Carrasco and McElree, 2001).

(3) In the studies in which accuracy was not assessed (Rosen et al., 1999; Mayer et al., 2004; Peelen et al., 2004; Natale et al., 2006), it is not possible to know whether task difficulty was the same for both types of attention, and task difficulty can interact with the strength of fMRI activity (Culham et al., 2001; Herath et al., 2001).

(4) For the studies in which eye position was not monitored while observers performed the task in the scanner (Kim et al., 1999; Rosen et al., 1999; Mayer et al., 2004; Peelen et al., 2004; Kincade et al., 2005; Natale et al., 2006; Meyer et al., 2018; Bowling et al., 2019), the results could be due to covert attention, to overt attention or to both (Carrasco et al., 1995; Beauchamp et al., 2001; Carrasco, 2011).

(5) Given that exogenous attention is a fast, transient process (for review see Carrasco, 2011; Carrasco and Barbot, 2015), it was inappropriately manipulated in the studies in which long stimulus onset asynchronies (SOA) were used (Rosen et al., 1999; Peelen et al., 2004; Kincade et al., 2005; Natale et al., 2006; Esterman et al., 2008) making the comparison between the two attention conditions problematic.

(6) Except for two studies, one interested in attentional modulation in the Fusiform Face Area
(FFA; Esterman et al., 2008) and the other in the TPJ (Dugué et al., 2017), statistical
parametric mapping was applied to group averaged data to identify regions of the brain
that were active during task performance. Some found significant activity in the occipital
pole (Mayer et al., 2004; Peelen et al., 2004; Kincade et al., 2005; Natale et al., 2006),
but attentional modulation of fMRI activity was not systematically assessed across differ-
ent visual areas.

Given all these methodological limitations, it is unknown how these two types of attention
affect neural activity in each individual striate and extra-striate visual areas, and how modulation
of activity in visual cortex is linked to changes in perceptual performance. (See Dugué and col-
leagues (Dugué et al., 2017a), who published a table summarizing these and other methodolog-
ical problems for studies regarding covert attention and TPJ activation).

Typically, covert attention is manipulated by presenting a pre-cue, prior to the target –and
the aforementioned studies also did so (see **Table 1**). However, endogenous post-cues, pre-
sented after target offset, can also improve performance by affecting the information readout
(Kinchla et al., 1995; Nobre et al., 2004a; Ruff et al., 2007; Hulme et al., 2009) and modulate
fMRI activity in early visual areas (Nobre et al., 2004a; Hulme et al., 2009; Pestilli et al., 2011;
Sergent et al., 2011). Exogenous post-cues also affect performance in some tasks (Sergent et
al., 2013; Thibault et al., 2016), but not in others (Carrasco and Yeshurun, 1998; Gobell and
Carrasco, 2005; Anton-Erxleben et al., 2007; Fuller et al., 2009), and the only study evaluating
post-cues in exogenous attention showed no such modulation (Liu et al., 2005). Critically, no
single study has compared visual cortex activity with post-cues in endogenous and exogenous
attention.

Study	Cueing manipulation		Task	Dependent Variable	Analysis	Cortical areas	Eye tracking	Correlation MRI-behavior
	Endogenous	Exogenous						
Studies including Occipital areas								
Detection tasks								
Mayer et al., 2004	central arrow 70% valid	luminance change of peripheral place-holder 50% valid	X	RT	Group-averaged	**Occipital:** *Cuneus, MOG, SOG* + Frontal Temporo-Parietal	No	No
	SOA=100 or 800 ms							
Peelen et al., 2004	central arrow 75% valid	brightening of peripheral placeholder 50% valid	square	RT	Group-averaged	**Occipital:** *Cuneus* + Frontal Temporo-Parietal	No	No
	SOA=550 ms							
Natale et al., 2006	single peripheral rectangular frame 100% valid	five peripheral rectangular frames 20% valid	a black and white checkerboard	RT *for 4 out of 8 participants*	Group-averaged	**Occipital:** *Fusiform, TOS, LG, SC, LOG* + Frontal Temporo-Parietal	No	No
	SOA=8, 8.15, 8.3, 8.45 or 8.6 s							
Discrimination tasks								
Kincade et al., 2005	brightening of half of central fixation diamond 75% valid	color singleton in an array of colored squares 50% valid	T vs. L	RT (% correct)	Group-averaged	**Occipital:** *LO, MT, Cuneus, Fusiform, SOG* + Frontal Temporo-Parietal	No	No
	SOA=2160 ms							
Esterman et al., 2008	color change of a peripheral place-holder 75% valid	50% valid	face identity	RT (% correct)	ROI-based (FFA) + Group-averaged	**Occipital:** *FFA* + Frontal Temporo-Parietal	Yes	No
	SOA=300 ms							
Current study	central bar 75% invalid SOA=310 ms	peripheral bar 50% valid SOA=110 ms	orientation	d-prime (RT)	ROI-based	**Occipital:** *V1, V2, V3, V3A, hV4, LO1*	Yes	Yes
Studies NOT including Occipital areas								
Detection tasks								
Rosen et al., 1999	central arrow 80% valid	peripheral dot 50% valid	filling of a square	RT	Group-averaged	Frontal Temporo-Parietal	No	No
	SOA=400, 550 or 700 ms							

Discrimination tasks								
Kim et al., 1999	thickening of part of central fixation diamond 80% valid SOA=200, 400 or 800 ms	luminance change of peripheral place-holder 50% valid SOA= 100, 150 or 200 ms	X vs. +	RT (% correct)	Group-averaged	Frontal Temporo-Parietal	No	No
Meyer et al., 2018	color change of central fixation 83% valid SOA=700–1000 ms	whitening of periph-eral placeholder 50% valid SOA=100–300 ms	central color of a check-erboard	Inverse efficiency = RT/%correct	Group-averaged	Frontal Temporo-Parietal	No	No
Dugué et al., 2017	central bar 75% invalid SOA=310 ms	peripheral bar 50% valid SOA=110 ms	orientation	d-prime (RT)	ROI-based	Temporo-Parietal	Yes	No
Bowling et al., 2019	color change of central fixation 83% valid SOA=700–1000 ms	whitening of periph-eral placeholder 50% valid SOA=100–300 ms	central color of a check-erboard	Inverse efficiency = RT/%correct	Group-averaged	Frontal Temporo-Parietal	No	No

Table 1. fMRI studies comparing endogenous and exogenous attention independently in human participants. _For each study, we report the cueing manip-_

ulation for each attention condition (SOA: Stimulus Onset Asynchrony is the duration of the cue + interval before onset of the target), the task performed by the

participants, the dependent variable reported in the publication, whether the analysis was based on group-averaging or ROI-based on single-participants, the

cortical areas reported in the publications and showing significant activation due to cue and/or target, whether or not eye data were monitor in the scanner or used

for the subsequent fMRI analysis, and whether the fMRI BOLD signal was correlated with behavioral measures. IOG: Inferior Occipital Gyrus; LG: Lingual Gyrus;

LO: Lateral Occipital area; LOG: Lateral Occipital Gyrus; MOG: Middle Occipital Gyrus; MT: Middle Temporal area; SC: Striate Cortex; SOG: Superior Occipital

Gyrus; TOS: Temporal-Occipital Sulcus.

Here, with the same participants, task, stimuli and task difficulty for all attention manipu-

lations, we tested the following four predictions: (1) Pre-cueing should induce an attentional mod-

ulation of fMRI activity, higher in the valid than the invalid condition in which attention needs to

be reoriented to the opposite location to perform the task (e.g., Liu et al., 2005). (2) Both endog-

enous and exogenous pre- and post-cueing effects should be stronger in contralateral than ipsi-

lateral visual regions relative to the attended hemifield (e.g., Liu et al., 2005; Serences and Yantis

2007; Pestilli et al., 2011). (3) Pre-cueing endogenous attention, but not exogenous attention,

should increase activity modulations along the visual hierarchy (e.g., higher in V4 than in V1;

Kastner et al., 1999; Pestilli et al., 2011; for reviews Chica et al., 2013; Beck and Kastner, 2014).

For endogenous attention, a top-down process, modulations from higher-order, fronto-parietal

attentional regions would send feedback information to visual cortex with diminishing effects in

earlier visual areas, given the increased distance from the source. (4) Post-cueing endogenous

(Nobre et al., 2004b; Hulme et al., 2009; Pestilli et al., 2011; Sergent et al., 2011), but not exog-

enous (Liu et al., 2005), attention should induce attentional modulation of fMRI activity in early

visual areas. Voluntary, endogenous attention would facilitate reading out perceptual information

(Dugué et al., 2017a), and modulate its processing (Nobre et al., 2004a; Hulme et al., 2009;

Pestilli et al., 2011; Sergent et al., 2011). The only fMRI study assessing the effects of post-

cueing exogenous attention found no attentional modulation of fMRI activity in early visual areas

(Liu et al., 2005); some behavioral studies report post-cueing effects (Sergent et al., 2013; Thi-

bault et al., 2016)) but others found no such effects (Carrasco and Yeshurun, 1998; Gobell and

Carrasco, 2005).

To test these four predictions, we measured fMRI activity and compared the effects of

endogenous and exogenous attention in early visual areas while the same observers performed

the same task –a 2-AFC orientation discrimination task, contingent upon contrast sensitivity

(Nachmias, 1967; Carrasco et al., 2000; Pestilli et al., 2009). We used a fully-crossed design:

two attention conditions –endogenous or exogenous attentional orienting– and two types of cue-

ing –pre- or post-cue. We evaluated fMRI activity at both the attended and the un-attended loca-

tions, given the ubiquitous performance tradeoffs at attended (benefits) and unattended (costs)

locations compared to a neutral condition (e.g. Luck et al., 1994; Lu and Dosher, 1998; Pestilli

and Carrasco, 2005; Giordano et al., 2009; Montagna et al., 2009; Herrmann et al., 2010; Ye-

shurun and Rashal, 2010), and the importance of evaluating both the orienting and reorienting of

attention (Dugué et al., 2016; 2017a; Senoussi et al., 2019), critical in an ever-changing environ-

ment (Dugué et al., 2016; 2017b).

This is the first study to systematically evaluate and directly compare how pre and post-

orienting, and reorienting, of endogenous and exogenous attention modulate neural activity in

visual cortex to affect behavior. The results indicate that these two types of spatial covert attention

distinctly modulate activity in individual retinotopic visual cortical areas. These differences in ac-

tivity are consonant with differential engagement of top-down and bottom-up processes and their

respective temporal dynamics. These results suggest that endogenous attention facilitates both

the encoding and the readout of visual information whereas exogenous attention only facilitates

the encoding of information.

RESULTS

Endogenous and exogenous attention improve performance

Observers performed a 2-AFC orientation-discrimination task under two attentional con-

ditions (exogenous or endogenous attention), when the cue was presented either before (pre-

cue) or after (post-cue) the grating stimuli (see Methods), and while their brain activity was meas-

ured with fMRI (**Figure 1**). The cue was either valid or invalid (50/50% of the time in the exoge-

nous condition, and 75/25% in the endogenous condition, respectively).

We calculated performance accuracy (d') in each condition as the main dependent varia-

ble, for each observer separately (**Figure 1B**, top row). We conducted a three-way repeated

measures 2x2x2 ANOVA (exogenous/endogenous x valid/invalid x pre/post-cue). There was bet-

ter performance for valid than invalid cues ($F(1,4)=23.6$, $p=0.008$). Exogenous versus endoge-

nous cues were statistically indistinguishable ($F(1,4)<1$), and there was no evidence for a

difference between pre- versus post-cues (F(1,4)<1). None of the two or three-way interactions

were significant.

Figure 1. Experimental protocol. A. *Observers performed a 2-AFC orientation-discrimination task. Pre-cues were presented before the gratings. Post-cues were presented after the gratings. Exogenous cues appeared in the periphery, above one of the two grating locations. Endogenous cues appeared at the center of the screen and pointed toward one of the two grating locations (The display is not at scale for the purpose of visibility). The ISI between the cue and the gratings differed between exogenous (50 ms) and endogenous (250 ms) conditions. A response cue indicated the target location and instructed observers to press one of two keys to indicate whether the target grating was tilted clockwise or counterclockwise of vertical. The fixation-cross turned green for a correct answer, and red for an incorrect answer. **B.** Behavioral performance for each observer (n=5) and averaged across them. (Top) Performance accuracy (d') as a function of cueing condition. (Bottom) Median reaction time. V, valid cue (location of pre-cue/post-cue and response cue matched). IN, invalid cue (location of pre-cue/post-cue and response cue mismatched). Pre, pre-cue presented before the grating stimuli. Post, post-cue presented after gratings. Valid cues led to more accurate and faster responses; there was no evidence for a speed-accuracy trade-off. Error bars, ± 1 SEM across observers.*

Reaction time was also calculated in each condition, for each observer separately (**Figure**

1B, bottom row). The corresponding three-way ANOVA revealed that reaction times were faster

for valid than invalid cues (F(1,4)=62.3, p=0.001). There was no evidence for a difference be-

tween exogenous and endogenous cues (F(1,4)=2.7, p=0.17), either for pre-cues or post-cues

(F(1,4)=1.7, p=0.27). There were two significant interactions, indicating that the differences be-

tween pre-cues and post-cues (F(1,4)=8.1, p=0.047) and between valid cues and invalid cues

(F(1,4)=16.2, p=0.02) were more pronounced for endogenous than for exogenous attention.

These behavioral results, which are consistent with previous findings (Lu and Dosher,

1998; Carrasco et al., 2000; Dosher and Lu, 2000; Ling and Carrasco, 2006b; Giordano et al.,

2009; Pestilli et al., 2009; Herrmann et al., 2010), show that attention improved orientation dis-

crimination (both accuracy and reaction time), with no evidence of a speed-accuracy trade-off,

and similarly for both types of attention and for both pre- and post-cues. Thus, the behavioral

effects confirm the successful manipulation of endogenous and exogenous attention, for both

pre- and post-cues, consistently across individuals.

Attentional modulation of perceptual and post-perceptual information processing in vis-

ual cortex

Visual areas were mapped in each participant following retinotopic mapping procedures

(**Figure 2**), and regions of interest (ROIs) were selected based on previous literature: V1, V2, V3

(for V2 and V3, ventral and dorsal ROIs were averaged), V3A, hV4 and LO1 (e.g., Engel et al.,

1994; Sereno et al., 1995; Engel et al., 1997; Larsson and Heeger, 2006; Wandell et al., 2007).

Activity was higher for contralateral than ipsilateral brain areas. For each ROI, we meas-

ured the fMRI response amplitudes for each type of attention – exogenous and endogenous –

and each cueing condition – pre- and post-cueing, for the contralateral and ipsilateral side to the

cued (attended) location. We analyzed the fMRI responses evoked by each type of attention as

a function of the contralateral and ipsilateral brain activity relative to the cue location (**Supple-**

mental Figure 1). ANOVAs indicated that there was higher contralateral than ipsilateral activity

across brain areas (endogenous: $F(1,4)=59.9$, $p=0.0015$; exogenous: $F(1,4)=218.8$, $p=0.0001$).

For both types of attention, this difference was more pronounced for valid than invalid cues (en-

dogenous: $F(1,4)=8.6$, $p=0.04$; exogenous: $F(1,4)=21.1$, $p=0.01$). In the following analyses, we

then concentrate on fMRI activity in the contralateral ROIs to the (attended) location.

Figure 2. Retinotopic mapping and stimulus localizer of a representative observer. *A flat-tened representation of the right hemisphere occipital pole. Left, map of polar angle. The color of each voxel in the map represents the response phase of the time series in response to a rotating wedge stimulus. Right, stimulus localizer. The black outlines correspond to the retinotopic areas (defined for each observer based on their polar angle maps). The color of each voxel indicated the phase of the response to the grating stimuli presented in the lower left visual field.*

fMRI responses correlated with behavioral variability. We evaluated the degree to which inter-individual variability in behavioral performance co-varied with variability in fMRI responses. We computed the correlation (**Figure 3A**), across observers, ROIs, types of attention and cueing conditions, between the fMRI responses (percent change in image intensity) and behavioral performance accuracy (d'). We found a positive correlation between fMRI activity and d' (Pearson correlation r = 0.23, p = 0.003), which was present for both attention types (endogenous: Pearson correlation r = 0.3, p = 0.003; exogenous: r = 0.2, p = 0.02). We further showed that fMRI activity was overall higher for valid than invalid trials (t(4) = 4.0, p = 0.01). This was the case for each observer (**Figure 3B**). Together, these results are consistent with previous findings showing that fMRI response correlates with behavioral variability, and that attentional orienting increases fMRI signal in early visual cortex (for review Beck and Kastner, 2014).

Figure 3. Inter-individual fMRI response. A. *Positive correlation between fMRI response and behavioral performance (Pearson correlation r = 0.23, p = 0.003). Each dot indicates fMRI response amplitude measured from one ROI and one observer, separately for each attentional condition (pre and post-cue, valid and invalid).* **B.** *fMRI response difference between valid and invalid trials combined across all conditions and ROIs for each observer (colored dots), and across observers (black dot; average significantly positive: t(4) = 4.0, p = 0.01). Error bar, ± 1 SEM across observers.*

We then tested our two novel predictions: (1) attentional modulation of fMRI stimulus-evoked activity increases along the visual hierarchy for endogenous, but is constant for exogenous, pre-cueing; and (2) attentional modulation of fMRI stimulus-evoked activity for endogenous is constant along the visual hierarchy, whereas there is no attentional modulation for exogenous, post-cueing. We measured the differences in activity between valid and invalid trials for each type of attention and for pre- and post-cues, across the hierarchy of visual cortical areas (**Figure 4**). Regression analyses showed a significant increase along the hierarchy of the visual areas of the activity difference evoked by valid and invalid trials for the endogenous pre-cueing condition (F = 25.8, p = 0.007, R^2 = 0.87), but not for the exogenous pre-cueing condition, endogenous post-cueing or exogenous post-cueing conditions (p >.1).

Figure 4. Specificity of pre and post-cueing for endogenous and exogenous attention. The difference in fMRI response between valid and invalid is plotted separately for pre- and post-cue conditions for each ROI. *, Statistically significant difference between valid and invalid, separately for pre and post-cueing (p < 0.05). Error bars on plots are ± 1 SEM.

For both endogenous (p < 0.05 for V3a, V4 and LO1; trend (p = 0.077) for V3; t-tests) and exogenous (p < 0.05 for V3, V3A, hV4 and LO1; trend (p = 0.074) for V2; t-tests) attention pre-cues elicited greater fMRI activity for valid than invalid cues (**Figure 5**, top-left and top-right panels). Furthermore, for endogenous attention (p < 0.05 for V2, V3, V3A, hV4 and LO1; trend (p = 0.079) for V1; t-tests; **Figure 5**, bottom-left panel), but not for exogenous attention (all p > 0.1; t-tests; **Figure 5**, bottom-right panel), post-cues elicited greater fMRI activity for valid than invalid cues in occipital areas. Taken together, these results confirm both of our predictions.

313 ***Figure 5. Single ROI responses for pre and post-cueing and for both endogenous and***
314 ***exogenous attention.*** *The difference in fMRI response between valid and invalid is plotted sep-*
315 *arately for pre and post-cue conditions for each ROI. *, Statistically significant difference between*
316 *valid and invalid, separately for pre and post-cueing (p < 0.05). #, trend (p < 0.1). Error bars on*
317 *plots are ± 1 SEM.*
318

319   These effects were consistently observed across participants (**Figure 6**). fMRI responses

320 were significantly larger for valid than invalid cues, for both endogenous and exogenous cues,

321 and for both pre- and post-cues. ANOVAs conducted separately for each ROI revealed a main

322 effect of validity, i.e. higher activity for valid than invalid cued locations in all visual areas (p<0.05;

323 t-tests), except V1.

Figure 6. Specificity of each visual ROI for endogenous and exogenous attention. *fMRI response amplitude was measured for each attentional condition. V: valid cue condition (target location matches the location indicated by pre-cue/post-cue). IN: invalid cue condition (target location at the opposite location relative to the pre-cue/post-cue). Pre: pre-cue presented before the grating stimuli. Post: post-cue presented after gratings. All four conditions in the contralateral ROI. Dist: all conditions averaged in the ipsilateral ROI. Each colored point corresponds to the data of one observer. The black points represent the average across all 5 observers. *, Statistically significant difference between valid and invalid, separately for pre and post-cueing (p < 0.05). #, trend (p < 0.1). Error bars are plotted on the difference between valid and invalid and represent ± 1 SEM across observers.*

DISCUSSION

This is the first study to compare pre and post-orienting, and reorienting, of endogenous

and exogenous attention in visual cortex, while concurrently assessing visual performance using

well-established psychophysical protocols to manipulate attention. The fact that the same ob-

servers performed the same orientation discrimination task with the same stimuli and task diffi-

culty under different attentional manipulations enabled us to isolate the fMRI activity induced by

each type of attention during orienting and reorienting.

Previous studies comparing endogenous and exogenous attention conditions state vari-

ous attentional effects in early visual areas (see **Table 1**). Some report differential effects be-

tween these attention conditions in the right LO and MT (Kincade et al., 2005), others in the

cuneus (Peelen et al., 2004; Mayer et al., 2004; Kincade et al., 2005), the occipital gyrus (SOG,

MOG, LOG; Mayer et al., 2004; Kincade et al., 2005; Natale et al., 2006), the fusiform area (Kin-

cade et al., 2005; Natale et al., 2006; Esterman et al., 2008), and the TOS, LG and SC (Natale

et al., 2006). This lack of clear picture regarding the differential impact of endogenous and exog-

enous spatial attention is likely due to the methodological differences and limitations we dis-

cussed in the Introduction (see also **Table 1**). In any case, it is often assumed that the effects of

endogenous and exogenous attention are the same in striate and extra-striate areas (Corbetta

and Shulman, 2002; Peelen et al., 2004; Corbetta et al., 2008; Chica et al., 2013; Beck and

Kastner, 2014).

To further our knowledge of the neural correlates of attention, we investigated both atten-

tional orienting (valid cueing) and reorienting (invalid cueing), critical in an ever-changing envi-

ronment (Dugué et al., 2016; 2017b; Senoussi et al., 2019). Furthermore, given ubiquitous per-

formance tradeoffs between attended (benefits) and unattended (costs) locations (e.g. Pestilli

and Carrasco, 2005; Giordano et al., 2009; Montagna et al., 2009; Herrmann et al., 2010), we

assessed activity at both attended (contralateral ROI) and unattended (ipsilateral ROI) locations.

Finally, we investigated how attentional effects varied as a function of pre- and post-cueing, thus

contrasting the neural correlates of perceptual and post-perceptual processing of information.

There was an overall positive correlation between performance in the orientation

discrimination task and the degree of attentional modulation in fMRI activity. This result was ex-

pected, indicating that as discriminability increases so does the attentional modulation in fMRI

activity. But only very few studies on the effect of spatial attention in early visual areas have

reported such a correlation (Ress et al., 2000; Giesbrecht et al., 2006; Liu et al., 2007a). The

behavioral effects obtained in the scanner are consistent with psychophysical studies. The en-

hanced performance brought about by the valid, but uninformative peripheral pre-cue is con-

sistent with an automatic, bottom-up involuntary capture of exogenous attention (e.g. Dosher and

Lu, 2000; Carrasco et al., 2000; Carrasco et al., 2004; Pestilli and Carrasco, 2005; Giordano et

al., 2009; Herrmann et al., 2010). The enhanced performance brought about by the valid, in-

formative central precue is consistent with a top-down, voluntary deployment of endogenous at-

tention (e.g. Dosher and Lu, 2000; Ling and Carrasco, 2006c; Giordano et al., 2009; Liu et al.,

2009; Herrmann et al., 2010).

In the endogenous attention condition, there was an increase in attentional modulation of

stimulus-evoked activity along the hierarchy of visual areas. Such a pattern is consistent with

previous studies suggesting that endogenous attention is a top-down modulation from frontal and

parietal areas feeding back to visual cortex, with diminishing effects in earlier visual areas (Kast-

ner et al., 1999; Maunsell and Cook, 2002; Kastner and Pinsk, 2004; Chica et al., 2013). Incon-

sistent with previous studies (e.g. Boynton et al., 1999; Brefczynski and DeYoe, 1999; Somers

et al., 1999; Herrmann et al., 2010; Pestilli et al., 2011), there was no evidence for attentional

modulation in V1. It might be that attentional modulation of V1 activity is more variable than other

visual cortical areas, making it harder to detect (see also Kastner et al., 1999; Liu et al., 2005).

Methodological differences between this and previous studies may have contributed to weaken-

ing the effect of attention in V1. The accrual time in the current endogenous condition was rela-

tively short (1300 ms in the valid condition and 500 ms in the invalid condition) compared to

previous studies investigating endogenous, voluntary attention, in which the cue and/or stimuli

were presented for a long duration to maximize BOLD measurements (e.g. Boynton et al., 1999;

Brefczynski and DeYoe, 1999; Somers et al., 1999; Pestilli et al., 2011). We chose the minimum

SOA (317 ms) at which performance benefits plateau to have as comparable conditions as

possible to the accrual time with the SOA for exogenous attention (117 ms). This short accrual

time may have limited the effects of attentional feedback to V1.

In the exogenous attention condition, in contrast to the endogenous attention, the atten-

tional modulation was approximately constant across the visual hierarchy. Previous studies have

reported a similar effect (Müller and Kleinschmidt, 2007; Müller and Ebeling, 2008), others a

decrease (Heinen et al., 2011), and yet others an increase (Liu et al., 2005; Mulckhuyse et al.,

2011) across the visual areas. This difference might be explained by different task parameters.

For example, in the Liu et al. (2005) study, observers knew which of the two stimuli was the target

they had to discriminate as soon as the stimuli were displayed; one stimulus was vertical and the

other was tilted to the left or the right. In the present study, both stimuli were independently tilted

and observers did not know which one was the target and which one was the distractor until later

when the response cue appeared.

Unlike in the endogenous pre-cueing condition in which the attention effect increased

along the processing stream, for the endogenous post-cueing effect there was no evidence that

it varied across these visual areas. The constant effect in the post-cue condition could be due to

the contribution of two counteracting factors: (1) the fMRI response evoked by the stimulus in

early visual areas may decrease along the visual hierarchy (Kay et al., 2013b); (2) the top-down

modulations from frontal and parietal areas feedback to visual cortex with diminishing effects in

earlier visual areas (Chica and Lupiáñez, 2009; Kastner et al., 1999; Pestilli et al., 2011).

In the exogenous condition, there was no significant post-cueing effect on early visual

areas. This result is consistent with that of Liu et al. (2005), who while evaluating exogenous

attention effects on occipital cortex included a post-cue condition to rule out sensory contamina-

tion of the cue (i.e. sensory response evoked by the cue itself) contributing to the enhanced

BOLD activity found in their pre-cue condition. In addition to ruling out a possible sensory con-

tamination, the present results show that, in contrast to endogenous attention, exogenous atten-

tion does not aid in the selective readout of information.

The ROI-based analysis that we followed here enabled us to compare contralateral and

ipsilateral modulation of BOLD activity, thus providing additional information regarding the

differences in processing dynamics of both types of attention. We observed a larger difference

between contralateral and ipsilateral areas for the valid than the invalid cueing condition. This

effect could be due to the fact that for the former, observers were attending to the same location

throughout the trial, whereas for the latter, when the response cue did not match the pre-cue,

observers had to switch their spatial attention to the opposite stimulus location, thus activity at

that new location would be accumulated for less time. For instance, for endogenous attention, for

the valid condition observers had been processing the target for almost 500 ms before the re-

sponse cue appeared. When the response cue matched the pre-cue, observers continued pro-

cessing and reading out the signal from that location for up to 800 ms (they were not allowed to

give an answer before the end of the response cue period). But when the response cue did not

match the pre-cue, then observers had to switch after 500 ms to the other location (ipsilateral)

thus accumulating less activity. Similarly, the accumulation time for the invalid cue condition in

exogenous attention was only about 300 ms. This accrual time explanation could also account

for the larger difference between contralateral and ipsilateral for pre-cues than post-cues, i.e.

there is a 300 ms accumulation when the exogenous pre-cue is invalid, while only 100 ms when

the post-cue in invalid. Likewise, the larger modulatory effect for endogenous relative to exoge-

nous attention is consistent with the difference in accrual time.

The results of the present study complement our recent study (Dugué et al., 2017a) in

which we demonstrated that sub-regions of the Temporo-Parietal Junction (TPJ) that respond

specifically to visual stimuli are more active when attention needs to be spatially reoriented (in-

valid cueing) than when attention remains at the cued location (valid cueing), and that partially

overlapping specific visual sub-regions mediate reorienting after orienting of endogenous or ex-

ogenous attention. Together, these two studies provide a comprehensive investigation of endog-

enous and exogenous attention, and pave the way for rigorous psychophysics informed, neu-

roimaging studies of covert, spatial attention. Here, we concentrated the analysis on visual corti-

cal areas in the occipital lobe because the slice prescription covered only a limited portion of the

brain. The present findings further our knowledge of the neurophysiological bases of covert at-

tention and have implications for models of visual attention, which should consider not only the

similarities, but also the differences in the orienting and reorienting of endogenous and exoge-

nous attention in occipital areas reported here.

In conclusion, the present results show some similarities and reveal important differences

in the specific neural correlates of endogenous and exogenous attention on early vision: An in-

creasing modulation of fMRI activity for pre-cueing endogenous attention, but constant modula-

tion for exogenous attention, along the hierarchy of visual occipital areas, as well as a reliable

and constant modulation of fMRI activity for post-cueing endogenous attention in occipital areas

but not for exogenous attention. These results suggest that endogenous attention facilitates both

the encoding and the readout of visual information whereas exogenous attention only facilitates

the encoding of information.

MATERIALS and METHODS

The methods employed in this study as well as their description are identical to those we

reported in a recent study, in which we compared activity in TPJ during orienting and reorienting

of endogenous and exogenous attention (Dugué et al., 2017a). We used optimal spatial and

temporal parameters to maximize the effects of these two types of attention, i.e. the benefits at

the attended location and costs at the unattended location, on an orientation discrimination task.

The same observers performed the same discrimination task under endogenous and exogenous

attention to enable direct comparison between these conditions (reviews by Carrasco, 2011; Car-

rasco and Barbot, 2015).

Observers

Five observers (three female, 24-30 years-old) with normal or corrected-to-normal vision

participated in the study. Observers provided written informed consent. The University Commit-

tee on Activities Involving Human Subjects at New York University approved the experimental

protocol. Each observer participated in 10 scanning sessions: one session to obtain a set of three

high-resolution anatomical volumes, two sessions for retinotopic mapping, three sessions for the

exogenous attention condition and three sessions for the endogenous attention condition (order

counterbalanced between observers). Prior to the first scanning session of each attention condi-

tion, observers performed several practice sessions outside the scanner.

Stimuli

Stimuli were generated using MATLAB (MathWorks) and the MGL toolbox (Gardner et

al., 2018b) on a Macintosh computer. Stimuli were displayed on a flat-panel display (NEC, LC-

XG250 MultiSync LCD 2110; resolution: 1024 x 768 pixels; refresh rate: 60 Hz) housed in a Far-

aday box with an electrically conductive glass front, positioned at the rear of the scanner bore.

Observers viewed the display through an angled mirror attached to the head coil, at a viewing

distance of 172 cm. The display was calibrated and gamma corrected using a linearized lookup

table. A white fixation cross (0.3°) was present at the center of the screen throughout the exper-

iment. The stimuli consisted of two gratings (4 cpd) windowed by raised cosines (3° of diame-

ter; 7% contrast). The stimuli were presented in the bottom quadrants (5° horizontal eccentricity;

–2.65° altitude). Endogenous cues were white rectangles (0.7°), positioned adjacent to the fixa-

tion cross indicating one of the two lower quadrants (0.35° horizontal eccentricity from the edge

of the fixation cross, and 0.35° altitude). Exogenous cues were also white rectangles (0.7°), but

were positioned adjacent to an upcoming grating stimulus, above the horizontal merid-

ian (1° away from the edge of the grating stimulus; and the edge of the cue 4.44° horizontal

eccentricity from the edge of the fixation cross) and vertically aligned with the stimulus.

Behavioral procedure

A single trial lasted 1700 ms for the exogenous attention condition and 1900 ms for the

endogenous attention condition, the only difference being the SOA between the cue and the

display; the timing of all the visual stimuli was the same in both attention conditions (**Figure 1**;

note that for illustration purposes, the display is not at scale). In 40% of the trials (pre-cue condi-

tion), a cue was shown first, followed by the pair of gratings. In 40% of the trials (post-cue condi-

tion), the order of presentation of the cue and the gratings was reversed. In 10% of the trials, the

gratings were not presented ('cue-only' trials). In the remaining 10% of the trials, neither a cue

nor the gratings were presented ('blank' trials). These trials were then included in the GLM

analysis to model the contribution of the visual signal produced by the cue (see MRI procedure).

For both pre-cue and post-cue trials, observers were instructed to report the orientation of a target

grating, i.e., clockwise or counter-clockwise compared to vertical, by pressing one of two keys.

For cue-only and blank trials, observers were asked to press a third key.

Cues (both exogenous and endogenous) were presented for 67 ms, indicating either the

left or the right grating location. The inter-stimulus interval (ISI) between the pre-cues and the

gratings was 50 ms for exogenous cues and 250 ms for endogenous cues, resulting in stimulus-

onset asynchronies (SOA) of 117 ms and 317 ms, optimal delays to manipulate exogenous and

endogenous attention respectively and maximize their behavioral consequences (Nakayama and

Mackeben, 1989; Mackeben and Nakayama, 1993; Liu et al., 2007b; Müller, 2014), while keeping

the trial duration as similar as possible. The behavioral effects of endogenous attention are sus-

tained (e.g. (Ling and Carrasco, 2006c)) and can thus still be present in later brain activity, as

shown in ERP studies (e.g. (Seiss et al., 2009)). Further, the brain responses elicited by exoge-

nous and endogenous cues are different during 300 ms following cue onset. The gratings were

shown for 50 ms. For the post-cue trials we used the same timings but inverted the order of

presentation, so that the cues followed the stimuli (e.g. (Kinchla et al., 1995; Carrasco and Ye-

shurun, 1998; Liu et al., 2005; Pestilli et al., 2011). A response cue, presented for 800 ms, at the

end of the trial indicated which of the two gratings was the target (50% of the trials at each loca-

tion). Note that in all four trial conditions (exogenous/endogenous, pre-/post-cueing), the re-

sponse cue appeared after both the cue and the stimuli had disappeared. The maximum delay

between the stimuli offset and the response cue onset was brief (~400 ms max in the endoge-

nous condition). This time interval is less than typically associated with a demand for working

memory (>600 ms; Phillips, 1974). Visual feedback was provided immediately following each

trial. The fixation cross turned green or red to indicate a correct or incorrect response, respec-

tively. If observers had not pressed any key after 530 ms, the fixation cross did not change color

indicating that they missed the response window.

Exogenous, peripheral cues were not informative regarding the target location or orienta-

tion; the cue and the target location matched in 50% of the trials (valid trials), but not in the other

50% of the trials (invalid). Endogenous, central cues were informative of the target location but

not its orientation; cues pointed towards the target in 75% of trials (valid trials), but not in the

remaining 25% of trials (invalid). Participants were informed of this validity. It is important to notice

that cue validity does not affect cueing effectiveness for exogenous attention, although it does so

do endogenous attention (e.g. Giordano et al., 2009). Finally, using a staircase procedure, per-

formance was equated between the two conditions.

Endogenous and exogenous attention conditions were performed in separate sessions to

ensure optimal manipulation of each attention system. Each of the 3 sessions of the exogenous

condition and 3 sessions of the endogenous consisted of 14 runs of 40 trials each, as well as an

additional run of stimulus localizer (see MRI procedure). Prior to the first session of each atten-

tional scanning condition, observers performed two practice sessions outside the scanner. The

tilt of the grating was adjusted so that each observer would achieve ~80% correct performance

in the valid trials. During the scanning sessions, the tilt was adjusted between runs to maintain

this overall performance level. Eye position was monitored using an infrared video camera sys-

tem (Eyelink 1K, SR Research, Ottawa, Ontario, http://www.sr-research.com/EL__1000.html).

Trials in which the observers blinked or broke fixation (1.5° away from fixation) at any point during

the trial sequence (from fixation onset to response cue offset; 13% ± 4% of the trials on average

across all observers) were identified and removed from the behavioral analysis (see Results),

and regressed separately in the MRI analysis (see below).

MRI Procedure

MRI scanning. Imaging was conducted on a 3T Siemens Allegra head-only scanner (Er-

langen, Germany). Padding was used to minimize observers' head movements. Anatomical im-

ages were acquired using a Siemens NM-011 head coil to transmit and receive. In a single scan-

ning session for each observer, three high-resolution anatomic images were acquired using a

T1-weighted magnetization-prepared rapid gradient echo (MPRAGE) sequence (FOV = 256 x

256 mm; 176 sagittal slices; 1 x 1 x 1 mm voxels). These three images were co-registered and

averaged.   We   then   used   the   public domain   software   FreeSurfer

(http://surfer.nmr.mgh.harvard.edu), to segment the gray matter from these averaged anatomical

volumes. All subsequent analyses were constrained only to voxels that intersected gray matter.

Functional images were acquired with a receive-only 8-channel surface coil array (Nova

Medical, Wilmington, MA). T2*-weighted echo-planar imaging sequence (TR = 1750 ms; TE = 30

ms; flip angle = 90°) measured blood oxygen level-dependent (BOLD) changes in image intensity

(Ogawa et al., 1990). One volume contained 28 slices oriented 45° to the calcarine sulcus and

covered the occipital and posterior parietal lobes (FOV = 192 x 192 mm; resolution = 2 x 2 x

2.5 mm; no gap). In each session, we acquired T1-weighted anatomical images in the same

slices as the functional images (spin echo; TR = 600 ms; TE = 9.1 ms; flip angle = 90°; resolution

= 1.5 × 1.5 × 3 mm). The in-plane images were used to align functional images from different

sessions to the same high-resolution anatomical volume for each participant, using a robust im-

age registration algorithm.

Pre-processing of the MRI data. Imaging data were analyzed using mrTools (Gardner et

al., 2018a) and custom software written in MATLAB. The first eight volumes of each run were

discarded to allow longitudinal magnetization to reach steady state. The measurements of the

B0 static magnetic field performed in each session were used to correct for spatial distortion. Pre-

processing of the functional data included motion correction, linear trend removal, and tem-

poral high-pass filtering (cutoff: 0.01 Hz) to remove low-frequency noise and slow drifts in the

fMRI signal.

Retinotopic mapping. Retinotopic mapping procedures followed well-established methods

using conventional traveling-wave, phase-encoded methods. Phase maps of polar angle were

measured using clockwise and counter-clockwise rotating checkerboard wedges, and eccen-

tricity maps were measured using contracting and expending checkerboard rings (Engel et al.,

1994; Sereno et al., 1995; Engel et al., 1997; Larsson and Heeger, 2006; Wandell et al., 2007).

Figure 2 (left panel) shows the borders of visual areas drawn by hand on flattened surface rep-

resentations of the brain following published conventions (Engel et al., 1997; Larsson and Hee-

ger, 2006; Wandell et al., 2007; Liu et al., 2007a).

Stimulus localizer. The stimuli were the same size, spatial frequency, and location as those in the main experiment, but they were at full contrast, and their orientation and phase were randomly changed every 200 ms to avoid adaptation. A localizer run consisted of a two-condition block alternation protocol: 16 cycles, each cycle was 17.5 s (8.75 s with the stimuli on, 8.75 s stimulus off). Observers completed one localizer run in each scanning session of the main experiment (6 runs overall, 4 min each). Observers were instructed to fixate a cross at the center of the screen throughout each run. Data were averaged across the 6 runs and analyzed using the same methods as for the retinotopic mapping scans, to define the cortical representation of the gratings. Each retinotopic ROI was further restricted to voxels that responded positively during the blocks when the grating stimuli were presented. A sinusoid was fit to the fMRI time series from each voxel. To be conservative, voxels were included in the ROI if the best-fit sinusoid had a phase value between 0 and pi, and if the correlation (technically, coherence) between the best-fit sinusoid and the time series was greater than 0.2 (**Figure 2**, right panel). In addition, we conducted the analysis without restricting the ROI to this coherence level, yielding similar results that supported the same conclusions.

Event-related analysis. fMRI time series in the main experiment were averaged across voxels in each ROI (separately for each hemisphere) and then concatenated across runs. The data were denoised using GLMDenoise (Kay et al., 2013a). fMRI response amplitudes were then computed from the denoised data using linear regression, with twelve regressors: right and left valid pre-cue, right and left invalid pre-cue, right and left valid post-cue, right and left invalid post-cue, right and left cue-only, blank (no cue nor stimulus) and eye-movements (trials in which observers broke fixation or blinked). The resulting fMRI response amplitudes (for correct trials only) were then averaged across observers, separately for each ROI and separately for each hemisphere.

Acknowledgments: This work was supported by: NIH RO1-EY019693 to MC and DJH; NIH RO1-EY016200 to MC; the FYSSEN Foundation and the Philippe Foundation to LD; and the Center for Brain Imaging of New York University. We want to thank members of the Carrasco lab for constructive comments on the manuscript. The authors declare no conflicts of interest.

REFERENCES

Anton-Erxleben K, Carrasco M (2013) Attentional enhancement of spatial resolution: linking behavioural and neurophysiological evidence. Nat Rev Neurosci 14:188–200.

Anton-Erxleben K, Henrich C, Treue S (2007) Attention changes perceived size of moving visual patterns. Journal of Vision 7:1–9.

Barbot A, Carrasco M (2017) Attention Modifies Spatial Resolution According to Task Demands. Psychological Science 28:285–296

Barbot A, Landy MS, Carrasco M (2012) Differential effects of exogenous and endogenous attention on second-order texture contrast sensitivity. Journal of Vision 12:1–15.

Beauchamp MS, Petit L, Ellmore TM, Ingeholm J, Haxby JV (2001) A parametric fMRI study of overt and covert shifts of visuospatial attention. Neuroimage 14:310–321.

Beck DM, Kastner S (2009) Top-down and bottom-up mechanisms in biasing competition in the human brain. Vision Res 49:1154–1165.

Beck DM, Kastner S (2014) Neural systems for spatial attention in the human brain: Evidence from neuroimaging in the framework of biased competition. The Oxford Handbook of Attention:253–288.

Bowling JT, Friston KJ, Hopfinger JB (2019) Top-down versus bottom-up attention differentially modulate frontal–parietal connectivity. Hum Brain Mapp 9:1116.

Boynton GM, Demb JB, Glover GH, Heeger DJ (1999) Neuronal basis of contrast discrimination. Vision Res 39:257–269.

Brefczynski JA, DeYoe EA (1999) A physiological correlate of the "spotlight" of visual attention. Nat Neurosci 2:370–374.

Buracas GT, Boynton GM (2007) The effect of spatial attention on contrast response functions in human visual cortex. Journal of Neuroscience 27:93–97.

Buschman TJ, Miller EK (2009) Serial, Covert Shifts of Attention during Visual Search Are Reflected by the Frontal Eye Fields and Correlated with Population Oscillations. Neuron 63:386–396.

Busse L, Katzner S, Treue S (2008) Temporal dynamics of neuronal modulation during exogenous and endogenous shifts of visual attention in macaque area MT. Proc Natl Acad Sci USA 105:16380–16385.

Carrasco M (2011) Visual attention: the past 25 years. Vision Res 51:1484–1525.

Carrasco M, Barbot A (2015) How attention affects spatial resolution. Cold Spring Harbor Symposia on Quantitative Biology 79:149–160.

Carrasco M, Evert DL, Chang I, Katz SM (1995) The eccentricity effect: target eccentricity affects performance on conjunction searches. Perception & Psychophysics 57:1241–1261.

Carrasco M, Giordano AM, McElree B (2006) Attention speeds processing across eccentricity: Feature and conjunction searches. Vision Res 46:2028–2040.

Carrasco M, Marie Giordano A, McElree B (2004) Temporal performance fields: visual and attentional factors. Vision Res 44:1351–1365.

Carrasco M, McElree B (2001) Covert attention accelerates the rate of visual information pro-
cessing. PNAS 98:5363–5367.

Carrasco M, Penpeci-Talgar C, Eckstein M (2000) Spatial covert attention increases contrast
sensitivity across the CSF: support for signal enhancement. Vision Res 40:1203–1215.

Carrasco M, Yeshurun Y (1998) The contribution of covert attention to the set-size and eccen-
tricity effects in visual search. Journal of Experimental Psychology: Human Perception and
Performance 24:673–692.

Carrasco M, Yeshurun Y (2009) Covert attention effects on spatial resolution. Prog Brain Res
176:65–86.

Chica AB, Bartolomeo P, Lupiáñez J (2013) Two cognitive and neural systems for endogenous
and exogenous spatial attention. Behavioural Brain Research 237:107–123.

Chica AB, Lupiáñez J (2009) Effects of endogenous and exogenous attention on visual pro-
cessing: an Inhibition of Return study. Brain Res 1278:75–85.

Corbetta M, Patel G, Shulman GL (2008) The reorienting system of the human brain: from envi-
ronment to theory of mind. Submitted 58:306–324.

Corbetta M, Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the
brain. Nat Rev Neurosci 3:201–215.

Correa Á, Triviño M, Pérez-Dueñas C, Acosta A, Lupiáñez J (2010) Temporal preparation, re-
sponse inhibition and impulsivity. Brain and Cognition 73:222–228.

Culham JC, Cavanagh P, Kanwisher NG (2001) Attention Response Functions: Characterizing
Brain Areas Using fMRI Activation during Parametric Variations of Attentional Load. Neuron
32:737–745.

Doricchi F, Macci E, Silvetti M, Macaluso E (2010) Neural correlates of the spatial and expectancy
components of endogenous and stimulus-driven orienting of attention in the Posner task.
Cereb Cortex 20:1574–1585.

Dosher BA, Lu ZL (2000) Mechanisms of perceptual attention in precuing of location. Vision Res
40:1269–1292.

Dugué L, Merriam EP, Heeger DJ, Carrasco M (2017a) Specific Visual Subregions of TPJ Medi-
ate Reorienting of Spatial Attention. Cereb Cortex:1–16.

Dugué L, Roberts M, Carrasco M (2016) Attention Reorients Periodically. Current Biology
26:1595–1601.

Dugué L, Xue AM, Carrasco M (2017b) Distinct perceptual rhythms for feature and conjunction
searches. Journal of Vision 17:22–22.

Engel SA, Glover GH, Wandell BA (1997) Retinotopic organization in human visual cortex and
the spatial precision of functional MRI. Cereb Cortex 7:181–192.

Engel SA, Rumelhart DE, Wandell BA, Lee AT, Glover GH, ED C, Shadlen MN (1994) fMRI of
human visual cortex. Nature 369:525.

Esterman M, Prinzmetal W, DeGutis J, Landau AN, Hazeltine E, Verstynen T, Robertson L (2008)
Voluntary and involuntary attention affect face discrimination differently. Neuropsychologia
46:1032–1040.

Fuller S, Park Y, Carrasco M (2009) Cue contrast modulates the effects of exogenous attention
on appearance. Vision Res 49:1825–1837.

Gardner JL, Merriam EP, Movshon JA, Heeger DJ (2008) Maps of Visual Space in Human Oc-
cipital Cortex Are Retinotopic, Not Spatiotopic. J Neurosci 28:3988–3999.

Gardner JL, Merriam EP, Schluppeck D, Besle J, Heeger DJ (2018a) mrTools: Analysis and
visualization package for functional magnetic resonance imaging data. Available at:
http://doi.org/10.5281/zenodo.1299483.

Gardner JL, Merriam EP, Schluppeck D, Larsson J (2018b) MGL: Visual psychophysics stimuli
and experimental design package. Available at: http://doi.org/10.5281/zenodo.1299497.

Geng JJ, Vossel S (2013) Re-evaluating the role of TPJ in attentional control: contextual updat-
ing? Neurosci Biobehav Rev 37:2608–2620.

Giesbrecht B, Weissman DH, Woldorff MG, Mangun GR (2006) Pre-target activity in visual cortex
predicts behavioral performance on spatial and feature attention tasks. Brain Res 1080:63–
72.

Giordano AM, McElree B, Carrasco M (2009) On the automaticity and flexibility of covert atten-
tion: a speed-accuracy trade-off analysis. Journal of Vision 9:30.1–.10.

Gobell J, Carrasco M (2005) Attention alters the appearance of spatial frequency and gap size.
Psychological Science 16:644–651.

Hahn B, Ross TJ, Stein EA (2006) Neuroanatomical dissociation between bottom–up and top–
down processes of visuospatial selective attention. Neuroimage 32:842–853.

Heinen K, Ruff CC, Bjoertomt O, Schenkluhn B, Bestmann S, Blankenburg F, Driver J, Chambers
CD (2011) Concurrent TMS-fMRI reveals dynamic interhemispheric influences of the right
parietal cortex during exogenously cued visuospatial attention. European Journal of Neuro-
science 33:991–1000.

Herath P, Klingberg T, Young J, Amunts K, Roland PE (2001) Neural Correlates of Dual Task
Interference Can be Dissociated from Those of Divided Attention: an fMRI Study | Cerebral
Cortex | Oxford Academic. Cereb Cortex 11:796–805.

Herrmann K, Montaser-Kouhsari L, Carrasco M, Heeger DJ (2010) When size matters: attention
affects performance by contrast or response gain. Nat Neurosci 13:1554–1559.

Hulme OJ, Friston KF, Zeki S (2009) Neural correlates of stimulus reportability. Cognitive Neu-
roscience, Journal of 21:1602–1610.

Jigo M, Carrasco M (2018) Attention alters spatial resolution by modulating second-order pro-
cessing. Journal of Vision 18:2.

Kastner S, Pinsk MA (2004) Visual attention as a multilevel selection process. Cognitive, Affec-
tive, & Behavioral Neuroscience 4:483–500.

Kastner S, Pinsk MA, De Weerd P, Desimone R, Ungerleider LG (1999) Increased Activity in
Human Visual Cortex during Directed Attention in the Absence of Visual Stimulation. Neuron
22:751–761.

Kay K, Rokem A, Winawer J, Dougherty R, Wandell B (2013a) GLMdenoise: a fast, automated
technique for denoising task-based fMRI data. Front Neurosci 7:1–15.

Kay KN, Winawer J, Rokem A, Mezer A, Wandell BA (2013b) A two-stage cascade model of
BOLD responses in human visual cortex. Diedrichsen J, ed. PLoS Comput Biol 9:e1003079.

Kim YH, Gitelman DR, Nobre AC, Parrish TB, LaBar KS, Mesulam MM (1999) The large-scale
neural network for spatial attention displays multifunctional overlap but differential asym-
metry. Neuroimage 9:269–277.

Kincade JM, Abrams RA, Astafiev SV, Shulman GL, Corbetta M (2005) An event-related func-
tional magnetic resonance imaging study of voluntary and stimulus-driven orienting of atten-
tion. Journal of Neuroscience 25:4593–4604.

Kinchla RA, Chen Z, Evert D (1995) Precue effects in visual search: data or resource limited?
Perception & Psychophysics 57:441–450.

Larsson J, Heeger DJ (2006) Two retinotopic visual areas in human lateral occipital cortex. Jour-
nal of Neuroscience 26:13128–13142.

Li X, Lu Z-L, Tjan BS, Dosher BA, Chu W (2008) Blood oxygenation level-dependent contrast
response functions identify mechanisms of covert attention in early visual areas. Proc Natl
Acad Sci USA 105:6202–6207.

Ling S, Carrasco M (2006a) Sustained and transient covert attention enhance the signal via dif-
ferent contrast response functions. Vision Res 46:1210–1220.

Ling S, Carrasco M (2006b) Sustained and transient covert attention enhance the signal via dif-
ferent contrast response functions. Vision Res 46:1210–1220.

Ling S, Carrasco M (2006c) When sustained attention impairs perception. Nat Neurosci 9:1243–
1245.

Liu T, Abrams J, Carrasco M (2009) Voluntary attention enhances contrast appearance. Psycho-
logical Science 20:354–362.

Liu T, Larsson J, Carrasco M (2007a) Feature-based attention modulates orientation-selective
responses in human visual cortex. Neuron 55:313–323.

Liu T, Pestilli F, Carrasco M (2005) Transient attention enhances perceptual performance and
FMRI response in human visual cortex. Neuron 45:469–477.

Liu T, Stevens ST, Carrasco M (2007b) Comparing the time course and efficacy of spatial and
feature-based attention. Vision Res 47:108–113.

Lu Z-L, Li X, Tjan BS, Dosher BA, Chu W (2011) Attention Extracts Signal in External Noise: A
BOLD fMRI Study. J Cogn Neurosci 23:1148–1159.

Lu ZL, Dosher BA (1998) External noise distinguishes attention mechanisms. Vision Res
38:1183–1198.

Luck SJ, Hillyard SA, Mouloua M, Woldorff MG, Clark VP, Hawkins HL (1994) Effects of spatial
cuing on luminance detectability: psychophysical and electrophysiological evidence for early
selection. Journal of Experimental Psychology: Human Perception and Performance 20:887–
904.

Luo TZ, Maunsell JHR (2015) Neuronal Modulations in Visual Cortex Are Associated with Only
One of Multiple Components of Attention. Neuron 86:1182–1188.

Mackeben M, Nakayama K (1993) Express attentional shifts. Vision Res 33:85–90.

Martínez-Trujillo JC, Treue S (2002) Attentional Modulation Strength in Cortical Area MT De-
pends on Stimulus Contrast. Neuron 35:365–370.

Maunsell JHR, Cook EP (2002) The role of attention in visual processing. Philosophical Trans-
actions of the Royal Society of London B: Biological Sciences 357:1063–1072.

Mayer AR, Dorflinger JM, Rao SM, Seidenberg M (2004) Neural networks underlying endoge-
nous and exogenous visual–spatial orienting. Neuroimage 23:534–541.

McAdams CJ, Maunsell JHR (1999) Effects of Attention on Orientation-Tuning Functions of Sin-
gle Neurons in Macaque Cortical Area V4. J Neurosci 19:431–441.

Meyer KN, Du F, Parks E, Hopfinger JB (2018) Exogenous vs. endogenous attention: Shifting
the balance of fronto-parietal activity. Neuropsychologia 111:307–316.

Mitchell JF, Sundberg KA, Reynolds JH (2009) Spatial Attention Decorrelates Intrinsic Activity
Fluctuations in Macaque Area V4. Neuron 63:879–888.

Montagna B, Pestilli F, Carrasco M (2009) Attention trades off spatial acuity. Vision Res 49:735–
745.

Mulckhuyse M, Belopolsky AV, Heslenfeld D, Talsma D, Theeuwes J (2011) Distribution of at-
tention modulates salience signals in early visual cortex. Op de Beeck HP, ed. PLoS ONE
6:e20379.

Müller MM (2014) Neural Mechanisms of Feature-Based Attention. In: Cognitive Electrophysiol-
ogy of Attention, pp 123–135. Elsevier.

Müller NG, Ebeling D (2008) Attention-modulated activity in visual cortex—More than a simple
"spotlight." Neuroimage 40:818–827.

Müller NG, Kleinschmidt A (2007) Temporal Dynamics of the Attentional Spotlight: Neuronal Cor-
relates of Attentional Capture and Inhibition of Return in Early Visual Cortex. The Journal of
Cognitive Neuroscience 19:587–593.

Murray SO (2008) The effects of spatial attention in early human visual cortex are stimulus inde-
pendent. Journal of Vision 8:1–11.

Nachmias J (1967) Effect of Exposure Duration on Visual Contrast Sensitivity with Square-Wave
Gratings. J Opt Soc Am, JOSA 57:421–427.

Natale E, Marzi CA, Girelli M, Pavone EF, Pollmann S (2006) ERP and fMRI correlates of en-
dogenous and exogenous focusing of visual-spatial attention. Eur J Neurosci 23:2511–2521.

Nakayama K, Mackeben M (1989) Sustained and transient components of focal visual attention.
Vision Res 29:1631–1647.

Nobre AC, Coull JT, Maquet P, Frith CD, Vandenberghe R, Mesulam MM (2004a) Orienting at-
tention to locations in perceptual versus mental representations. Cognitive Neuroscience,
Journal of 16:363–373.

Nobre AC, Coull JT, Maquet P, Frith CD, Vandenberghe R, Mesulam MM (2004b) Orienting at-
tention to locations in perceptual versus mental representations. Cognitive Neuroscience,
Journal of 16:363–373.

O'Connor DH, Fukui MM, Pinsk MA (2002) Attention modulates responses in the human lateral
geniculate nucleus. Nature 5:1203–1209.

Ogawa S, Lee TM, Kay AR, Tank DW (1990) Brain magnetic resonance imaging with contrast
dependent on blood oxygenation. Proc Natl Acad Sci USA 87:9868–9872.

Peelen MV, Heslenfeld DJ, Theeuwes J (2004) Endogenous and exogenous attention shifts are
mediated by the same large-scale neural network. Neuroimage 22:822–830.

Pestilli F, Carrasco M (2005) Attention enhances contrast sensitivity at cued and impairs it at
uncued locations. Vision Res 45:1867–1875.

Pestilli F, Carrasco M, Heeger DJ, Gardner JL (2011) Attentional enhancement via selection and
pooling of early sensory responses in human visual cortex. Neuron 72:832–846.

Pestilli F, Ling S, Carrasco M (2009) A population-coding model of attention's influence on con-
trast response: Estimating neural effects from psychophysical data. Vision Res 49:1144–
1153.

Phillips WA (1974) On the distinction between sensory storage and short-term visual memory.
Perception & Psychophysics 16:283–290.

Posner MI, Snyder CR, Davidson BJ (1980) Attention and the detection of signals. J Exp Psychol
109:160–174.

Reed AV (1973) Speed-Accuracy Trade-Off in Recognition Memory. Science 181:574–576.

Ress D, Backus BT, Heeger DJ (2000) Activity in primary visual cortex predicts performance in
a visual detection task. Nat Neurosci 3:940–945.

Reynolds JH, Heeger DJ (2009) The Normalization Model of Attention. Neuron 61:168–185.

Reynolds JH, Pasternak T, Desimone R (2000) Attention increases sensitivity of V4 neurons.
Neuron 26:703–714.

Rosen AC, Rao SM, Caffarra P, Scaglioni A, Bobholz JA, Woodley SJ, Hammeke TA, Cunning-
ham JM, Prieto TE, Binder JR (1999) Neural basis of endogenous and exogenous spatial
orienting. A functional MRI study. Cognitive Neuroscience, Journal of 11:135–152.

Ruff CC, Kristjánsson Á, Driver J (2007) Readout from iconic memory and selective spatial at-
tention involve similar neural processes. Psychological Science 18:901–909.

Ruff DA, Cohen MR (2014) Attention can either increase or decrease spike count correlations in
visual cortex. Nat Neurosci 17:1591–1597.

Seiss E, Driver J, Eimer M (2009) Effects of attentional filtering demands on preparatory ERPs
elicited in a spatial cueing task. Clin Neurophysiol 120:1087–1095.

Senoussi M, Moreland JC, Busch NA, Dugué L (2019) Attention explores space periodically at
the theta frequency. Journal of Vision 19:1–17.

Sereno MI, Dale AM, Reppas JB, Kwong KK, Belliveau JW, Brady TJ, Rosen BR, Tootell R (1995)
Borders of multiple visual areas in humans revealed by functional magnetic resonance im-
aging. Science 268:889–893

Sergent C, Ruff CC, Barbot A, Driver J, Rees G (2011) Top-down modulation of human early
visual cortex after stimulus offset supports successful postcued report. The Journal of Cog-
nitive Neuroscience 23:1921–1934.

Sergent C, Wyart V, Babo-Rebelo M, Cohen L, Naccache L, Tallon-Baudry C (2013) Cueing
Attention after the Stimulus Is Gone Can Retrospectively Trigger Conscious Perception. Cur-
rent Biology 23:150–155.

Silvetti M, Lasaponara S, Lecce F, Dragone A, Macaluso E, Doricchi F (2015) The Response of
the Left Ventral Attentional System to Invalid Targets and its Implication for the Spatial Ne-
glect Syndrome: a Multivariate fMRI Investigation. Cereb Cortex 1–2.

Somers DC, Dale AM, Seiffert AE, Tootell RB (1999) Functional MRI reveals spatially specific
attentional modulation in human primary visual cortex. PNAS 96:1663–1668.

Thibault L, van den Berg R, Cavanagh P, Sergent C (2016) Retrospective Attention Gates Dis-
crete Conscious Access to Past Sensory Stimuli. PLoS ONE 11:e0148504.

Wandell BA, Dumoulin SO, Brewer AA (2007) Visual field maps in human cortex. Neuron 56:366–
383.

Wang F, Chen M, Yan Y, Zhaoping L, Li W (2015) Modulation of Neuronal Responses by Exog-
enous Attention in Macaque Primary Visual Cortex. Journal of Neuroscience 35:13419–
13429.

Wickelgren WA (1977) Speed-accuracy tradeoff and information processing dynamics. Acta Psy-
chologica 41:67–85.

Williford T, Maunsell JHR (2006) Effects of Spatial Attention on Contrast Response Functions in
Macaque Area V4. J Neurophysiol 96:40–54.

Yashar A, Lamy D (2011) Refining the dual-stage account of intertrial feature priming: does motor
response or response feature matter? Atten Percept Psychophys 73:2160–2167.

Yeshurun Y, Carrasco M (1998) Attention improves or impairs visual performance by enhancing
spatial resolution. Nature 396:72–75.

Yeshurun Y, Montagna B, Carrasco M (2008) On the flexibility of sustained attention and its ef-
fects on a texture segmentation task. Vision Research 48:80–95.

Yeshurun Y, Rashal E (2010) Precueing attention to the target location diminishes crowding and
reduces the critical distance. Journal of Vision 10:16–16.