1   # Differential impact of endogenous and exogenous attention

2   # on activity in human visual cortex

4   Laura Dugué[1,2,3,4,5], Elisha P. Merriam[1,2,6], David J. Heeger[1,2] & Marisa Carrasco[1,2]

[1] Department of Psychology, New York University

[2] Center for Neural Science, New York University

[3] CNRS (Integrative Neuroscience and Cognition Center, UMR 8002) Paris, France

[4] Université de Paris, Paris, France

[5] Institut Universitaire de France (IUF), Paris, France

[6] Laboratory of Brain and Cognition, NIMH/NIH, Bethesda, MD

Running title: endogenous and exogenous attention

Corresponding author:

Laura Dugué

Current address: 45 rue des Saints-Pères 75006 Paris, FRANCE

laura.dugue@u-paris.fr

ABSTRACT (148 words)

How do endogenous (voluntary) and exogenous (involuntary) attention modulate activity in visual cortex? Using ROI-based fMRI analysis, we measured fMRI activity for valid and invalid trials (target at cued/un-cued location, respectively), pre- or post-cueing endogenous or exogenous attention, while participants performed the same discrimination task. We found stronger modulation in contralateral than ipsilateral visual regions, and higher activity in valid- than invalid-trials. For endogenous attention, modulation of stimulus-evoked activity due to a pre-cue increased along the visual hierarchy, but was constant due to a post-cue. For exogenous attention, modulation of stimulus-evoked activity due to a pre-cue was constant along the visual hierarchy, but not modulated due to a post-cue. These findings reveal that endogenous and exogenous attention distinctly modulate activity in visuo-occipital areas during orienting and reorienting; endogenous attention facilitates both the encoding and the readout of visual information whereas exogenous attention only facilitates the encoding of information.

Keywords: exogenous attention, endogenous attention, fMRI activity, occipital visual areas, top-down and bottom-up processing, visual perception.

INTRODUCTION

Spatial, covert visual attention is the selective processing of visual information in space, without change in gaze. Attention can be allocated voluntarily –endogenous attention– or involuntarily –exogenous attention. Endogenous and exogenous attention have different temporal dynamics; endogenous takes about 300 ms to be deployed and can be sustained at will whereas exogenous attention takes only about 100 ms to be deployed and it is transient (for review see Carrasco, 2011; Carrasco and Barbot, 2015). These two types of attention often have similar perceptual consequences (reviews by Carrasco, 2011; Carrasco and Barbot, 2015), but notable exceptions indicate that whereas endogenous attention acts in a flexible way, exogenous attention acts in an inflexible or automatic way. For instance: (a) The benefits and costs in perception (discriminability and speed of information accrual) scale with cue validity for endogenous but not for exogenous attention (e.g. Sperling and Melchner, 1978; Kinchla, 1980; Giordano et al., 2009); (b) The effects of covert attention on contrast sensitivity often differ for endogenous and exogenous attention (e.g. Dosher and Lu, 2000; Ling and Carrasco, 2006a; Pestilli et al., 2009; Barbot et al., 2012); and (c) For a texture segmentation task in which heightened spatial resolution improves or impairs performance as a function of target eccentricity, endogenous attention improves performance across eccentricity, whereas exogenous attention improves performance at peripheral locations where resolution is poor, but hampers performance where resolution is already high for the task at hand (e.g. Yeshurun and Carrasco, 1998; Carrasco et al., 2006; Yeshurun et al., 2008; Barbot and Carrasco, 2017; Jigo and Carrasco, 2018).

Basic visual processes, such as contrast sensitivity and spatial resolution, are mediated by activity in early visual cortex, and are altered by covert attention (reviews by Carrasco and Yeshurun, 2009; Carrasco, 2011; Anton-Erxleben and Carrasco, 2013; Carrasco and Barbot, 2015). Specifically, single-unit studies in monkeys have demonstrated effects of endogenous attention in occipital areas (e.g. McAdams and Maunsell, 1999; Reynolds et al., 2000; Martínez-Trujillo and Treue, 2002; Williford and Maunsell, 2006; Mitchell et al., 2009; Reynolds and Heeger, 2009; Ruff and Cohen, 2014; Luo and Maunsell, 2015). Additionally, fMRI studies have shown that endogenous attention causes a baseline shift in early visual areas (e.g. Kastner et

al., 1999; Somers et al., 1999; O'Connor et al., 2002; Buracas and Boynton, 2007; Murray, 2008; Herrmann et al., 2010; Pestilli et al., 2011; review by Beck and Kastner, 2014) and increases the dynamic range of fMRI responses (Li et al., 2008; Lu et al., 2011). Comparatively, little is known about the effect of exogenous attention on visual areas both from single-unit studies (Busse et al., 2008; Wang et al., 2015) and fMRI studies (for reviews see Carrasco, 2011; Anton-Erxleben & Carrasco, 2013).

Since Corbetta and Schulman's seminal review (2002) on the neural bases of endoge­nous and exogenous attention, there has been emphasis on characterizing networks of brain regions within the frontal and parietal lobes (for reviews see Chica et al., 2013; Beck and Kastner, 2014). Yet, there remains considerable debate regarding the dissociation between dorsal re­gions, for endogenous attention, and ventral regions, for exogenous attention (e.g. Hahn et al., 2006; Bowling et al., 2019), including the role of the temporo-parietal junction (TPJ; Doricchi et al., 2010; Geng and Vossel, 2013; Silvetti et al., 2015; Dugué et al., 2017a). Classically, research­ers have described endogenous attention as a top-down process, and exogenous attention as a bottom-up process (e.g., Posner et al., 1980; Nakayama and Mackeben, 1989; Corbetta and Shulman, 2002; Hahn et al., 2006; Beck and Kastner, 2009; Carrasco, 2011; Chica et al., 2013; Beck and Kastner, 2014; Bowling et al., 2019). This characterization originated in psychophysics experiments, and was then studied using fMRI, in which the two types of attention have been often investigated separately (for reviews Beck and Kastner, 2009; Carrasco, 2011; Chica et al., 2013; Beck and Kastner, 2014).

Studies of endogenous and exogenous attention focusing on parietal and frontal areas have shown that the two types of attention differentially modulate fronto-parietal connectivity (Buschman and Miller, 2009; Bowling et al., 2019). For example, there are critical differences in the temporal order of neural responses in frontal and parietal cortex in monkeys between these attention conditions, i.e. frontal activity precedes parietal activity during endogenous attentional orienting, whereas parietal activity precedes frontal activity during exogenous orienting. Critically, it is often assumed that the effects of endogenous and exogenous attention are the same in

striate and extra-striate areas (Corbetta and Shulman, 2002; Peelen et al., 2004; Corbetta et al.,

2008; Chica et al., 2013; Beck and Kastner, 2014).

A number of important considerations limit the conclusions that may be drawn from the

few studies that have directly compared independent effects of visual exogenous and endoge-

nous spatial attention (see **Table 1**):

(1) The effect of attention on behavioral performance was assessed in some studies with a

102   detection task using RT as their only dependent variable (Mayer et al., 2004; Peelen et

103   al., 2004; Natale et al., 2006), in which performance may differ due to speed of pro-

104   cessing, discriminability or criterion (Reed, 1973; Wickelgren, 1977; Carrasco and

105   McElree, 2001) and/or motor behavior (Correa et al., 2010; Yashar and Lamy, 2011).

106   (2) In the studies in which performance was assessed in a discrimination task using RT (Kim

107   et al., 1999; Kincade et al., 2005; Esterman et al., 2008), small RT differences were re-

108   ported, which could have resulted from speed of processing, discriminability, or criterion

109   factors (Reed, 1973; Wickelgren, 1977; Carrasco and McElree, 2001).

110   (3) In the studies in which accuracy was not assessed (Rosen et al., 1999; Mayer et al., 2004;

111   Peelen et al., 2004; Natale et al., 2006), it is not possible to know whether task difficulty

112   was the same for both types of attention, and task difficulty can interact with the strength

113   of fMRI activity (Culham et al., 2001; Herath et al., 2001).

114   (4) For the studies in which eye position was not monitored while participants performed the

115   task in the scanner (Kim et al., 1999; Rosen et al., 1999; Mayer et al., 2004; Peelen et al.,

116   2004; Kincade et al., 2005; Natale et al., 2006; Meyer et al., 2018; Bowling et al., 2019),

117   the results could be due to covert attention, overt attention or both (Carrasco et al., 1995;

118   Beauchamp et al., 2001; Carrasco, 2011).

119   (5) Given that exogenous attention is a fast, transient process (for review see Carrasco, 2011;

120   Carrasco and Barbot, 2015), it was inappropriately manipulated in the studies in which

121   long stimulus onset asynchronies (SOA) were used (Rosen et al., 1999; Peelen et al.,

122   2004; Kincade et al., 2005; Natale et al., 2006; Esterman et al., 2008) making the com-

123   parison between the two attention conditions problematic.

(6) Except for two studies, one interested in attentional modulation in the Fusiform Face Area

(FFA; Esterman et al., 2008) and the other in the TPJ (Dugué et al., 2017), statistical

parametric mapping was applied to group averaged data to identify regions of the brain

that were active during task performance. Some found significant activity in the occipital

pole (Mayer et al., 2004; Peelen et al., 2004; Kincade et al., 2005; Natale et al., 2006),

but attentional modulation of fMRI activity was not systematically assessed across differ-

ent visual areas.

Given all these methodological limitations, it is unknown how these two types of attention

affect neural activity in each individual visuo-occipital areas, and how modulation of activity in

visual cortex is linked to changes in perceptual performance. (See Dugué and colleagues (Dugué

et al., 2017a), who published a table summarizing these and other methodological problems for

studies regarding covert attention and TPJ activation).

Typically, covert attention is manipulated by presenting a pre-cue, prior to the target –and

the aforementioned studies also did so (see **Table 1**). However, endogenous post-cues, pre-

sented after target offset, can also improve performance by affecting the information readout

(Kinchla et al., 1995; Nobre et al., 2004a; Ruff et al., 2007; Hulme et al., 2009) and modulate

fMRI activity in early visual areas (Nobre et al., 2004a; Hulme et al., 2009; Pestilli et al., 2011;

Sergent et al., 2011). Exogenous post-cues also affect performance in some tasks (Sergent et

al., 2013; Thibault et al., 2016), but not in others (Carrasco and Yeshurun, 1998; Gobell and

Carrasco, 2005; Anton-Erxleben et al., 2007; Fuller et al., 2009), and the only study evaluating

post-cues in exogenous attention showed no such modulation (Liu et al., 2005). Critically, no

single study has compared visual cortex activity with post-cues in endogenous and exogenous

attention.

Study	Cueing manipulation		Task	Dependent Variable	Analysis	Cortical areas	Eye tracking	Correlation MRI-behavior
	Endogenous	Exogenous						
Studies including Occipital areas								
Detection tasks								
Mayer et al., 2004	central arrow 70% valid SOA=100 or 800 ms	luminance change of peripheral place-holder 50% valid	X	RT	Group-averaged	**Occipital:** *Cuneus, MOG, SOG* + Frontal Temporo-Parietal	No	No
Peelen et al., 2004	central arrow 75% valid SOA=550 ms	brightening of pe-ripheral placeholder 50% valid	square	RT	Group-averaged	**Occipital:** *Cuneus* + Frontal Temporo-Parietal	No	No
Natale et al., 2006	single peripheral rectangular frame 100% valid	five peripheral rec-tangular frames 20% valid SOA=8, 8.15, 8.3, 8.45 or 8.6 s	a black and white check-erboard	RT *for 4 out of 8 participants*	Group-averaged	**Occipital:** *Fusiform, TOS, LG, SC, LOG* + Frontal Temporo-Parietal	No	No
Discrimination tasks								
Kincade et al., 2005	brightening of half of central fixation diamond 75% valid	color singleton in an array of colored squares 50% valid SOA=2160 ms	T vs. L	RT (% correct)	Group-averaged	**Occipital:** *LO, MT, Cuneus, Fusiform, SOG* + Frontal Temporo-Parietal	No	No
Esterman et al., 2008	color change of a peripheral place-holder 75% valid	50% valid SOA=300 ms	face identity	RT (% correct)	ROI-based (FFA) + Group-averaged	**Occipital:** *FFA* + Frontal Temporo-Parietal	Yes	No
Current study	central bar 75% invalid SOA=310 ms	peripheral bar 50% valid SOA=110 ms	orientation	d-prime (RT)	ROI-based	**Occipital:** *V1, V2, V3, V3A, hV4, LO1*	Yes	Yes
Studies NOT including Occipital areas								
Detection tasks								
Rosen et al., 1999	central arrow 80% valid	peripheral dot 50% valid SOA=400, 550 or 700 ms	filling of a square	RT	Group-averaged	Frontal Temporo-Parietal	No	No

Discrimination tasks								
Kim et al., 1999	thickening of part of central fixation diamond 80% valid SOA=200, 400 or 800 ms	luminance change of peripheral place-holder 50% valid SOA= 100, 150 or 200 ms	X vs. +	RT (% correct)	Group-averaged	Frontal Temporo-Parietal	No	No
Meyer et al., 2018	color change of central fixation 83% valid SOA=700–1000 ms	whitening of periph-eral placeholder 50% valid SOA=100–300 ms	central color of a check-erboard	Inverse efficiency = RT/%correct	Group-averaged	Frontal Temporo-Parietal	No	No
Dugué et al., 2017	central bar 75% invalid SOA=310 ms	peripheral bar 50% valid SOA=110 ms	orientation	d-prime (RT)	ROI-based	Temporo-Parietal	Yes	No
Bowling et al., 2019	color change of central fixation 83% valid SOA=700–1000 ms	whitening of periph-eral placeholder 50% valid SOA=100–300 ms	central color of a check-erboard	Inverse efficiency = RT/%correct	Group-averaged	Frontal Temporo-Parietal	No	No

Table 1. fMRI studies comparing endogenous and exogenous attention independently in human participants. For each study, we report the cueing manipulation for each attention condition (SOA: Stimulus Onset Asynchrony is the duration of the cue + interval before onset of the target), the task performed by the participants, the dependent variable reported in the publication, whether the analysis was based on group-averaging or ROI-based on single-participants, the cortical areas reported in the publications and showing significant activation due to cue and/or target, whether or not eye data were monitor in the scanner or used for the subsequent fMRI analysis, and whether the fMRI BOLD signal was correlated with behavioral measures. IOG: Inferior Occipital Gyrus; LG: Lingual Gyrus; LO: Lateral Occipital area; LOG: Lateral Occipital Gyrus; MOG: Middle Occipital Gyrus; MT: Middle Temporal area; SC: Striate Cortex; SOG: Superior Occipital Gyrus; TOS: Temporal-Occipital Sulcus.

Here, with the same participants, task, stimuli and task difficulty for all attention manipu-

lations, we tested the following four predictions: (1) Pre-cueing should induce an attentional mod-

ulation of fMRI activity, higher in the valid than the invalid condition in which attention needs to

be reoriented to the opposite location to perform the task (e.g., Liu et al., 2005). (2) Both endog-

enous and exogenous pre- and post-cueing effects should be stronger in visual regions contra-

lateral to the attended hemifield (e.g., Liu et al., 2005; Serences and Yantis 2007; Pestilli et al.,

2011). (3) Pre-cueing endogenous attention, but not exogenous attention, should increase activ-

ity modulations along the visual hierarchy (e.g., higher in V4 than in V1; Kastner et al., 1999;

Pestilli et al., 2011; for reviews Chica et al., 2013; Beck and Kastner, 2014). For endogenous

attention, a top-down process, modulations from higher-order, fronto-parietal attentional regions

would send feedback information to visual cortex with diminishing effects in earlier visual areas,

given the increased distance from the source. (4) Post-cueing endogenous (Nobre et al., 2004b;

Hulme et al., 2009; Pestilli et al., 2011; Sergent et al., 2011), but not exogenous (Liu et al., 2005),

attention should induce attentional modulation of fMRI activity in early visual areas. Voluntary,

endogenous attention would facilitate reading out perceptual information (Dugué et al., 2017a),

and modulate its processing (Nobre et al., 2004a; Hulme et al., 2009; Pestilli et al., 2011; Sergent

et al., 2011). The only fMRI study assessing the effects of post-cueing exogenous attention found

no attentional modulation of fMRI activity in early visual areas (Liu et al., 2005); some behavioral

studies report post-cueing effects (Sergent et al., 2013; Thibault et al., 2016) but others found no

such effects (Carrasco and Yeshurun, 1998; Gobell and Carrasco, 2005; Anton-Erxleben et al.,

2007; Fuller et al., 2009).

To test these four predictions, we measured fMRI activity and compared the effects of

endogenous and exogenous attention in early visual areas while the same participants performed

the same task –a 2-AFC orientation discrimination task, contingent upon contrast sensitivity

(Nachmias, 1967; Carrasco et al., 2000; Pestilli et al., 2009). We used a fully-crossed design:

two attention conditions –endogenous or exogenous attentional orienting– and two types of cue-

ing –pre- or post-cue. We evaluated fMRI activity at both the attended and the un-attended loca-

tions, given the ubiquitous performance tradeoffs at attended (benefits) and unattended (costs)

locations compared to a neutral condition (e.g. Luck et al., 1994; Lu and Dosher, 1998; Pestilli

and Carrasco, 2005; Giordano et al., 2009; Montagna et al., 2009; Herrmann et al., 2010; Ye-

shurun and Rashal, 2010), and the importance of evaluating both the orienting and reorienting of

attention (Dugué et al., 2016; 2017a; Senoussi et al., 2019), critical in an ever-changing environ-

ment (Dugué et al., 2016; 2017b).

This is the first study to systematically evaluate and directly compare how pre and post-

orienting, and reorienting, of endogenous and exogenous attention modulate neural activity in

visual cortex to affect behavior. The results indicate that these two types of spatial covert attention

distinctly modulate activity in individual retinotopic visual cortical areas. These differences in ac-

tivity are consonant with differential engagement of top-down and bottom-up processes and their

respective temporal dynamics. These results suggest that endogenous attention facilitates both

the encoding and the readout of visual information whereas exogenous attention only facilitates

the encoding of information.

RESULTS

Endogenous and exogenous attention improve performance

Participants performed a 2-AFC orientation-discrimination task under two attentional con-

ditions (exogenous or endogenous attention), when the cue was presented either before (pre-

cue) or after (post-cue) the grating stimuli (see Methods), and while their brain activity was meas-

ured with fMRI (**Figure 1**). The cue was either valid or invalid (50/50% of the time in the exoge-

nous condition, and 75/25% in the endogenous condition, respectively).

In each condition, we calculated performance accuracy (d'), as the main dependent vari-

able, for each participant separately (**Figure 1B**, top row). A three-way repeated measures 2x2x2

ANOVA (exogenous/endogenous x valid/invalid x pre/post-cue) revealed higher performance for

valid than invalid cues ($F(1,4)=23.6$, $p=0.008$), that exogenous and endogenous cues were sta-

tistically indistinguishable ($F(1,4)<1$), and that there was no significant difference between pre-

and post-cues ($F(1,4)<1$). All the two and three-way interactions were not significant ($F<1$).

Figure 1. Experimental protocol. A. *Participants performed a 2-AFC orientation-discrimination task. (The display is not at scale for visibility purposes). Pre-cues and post-cues were presented before and after the stimuli, respectively. Exogenous cues appeared in the periphery, above one of the two stimulus locations. Endogenous cues appeared at the center of the screen and indicated one of the two stimulus locations The ISI between the cue and the gratings was shorter for the exogenous (50 ms) than the endogenous (250 ms) conditions. A response cue indicated the target location and instructed participants to indicate whether the target grating was tilted clockwise or counterclockwise of vertical by pressing one of two keys. To provide feedback, the fixation-cross turned green or red for a correct or an incorrect answer, respectively.* **B.** *Behavioral performance averaged across participants (black dots) and for each of them (colored dots; n=5) for endogenous (left) and exogenous (right) attention. (Top) Performance accuracy (d'; top) and median reaction time (bottom) as a function of cueing condition. V, valid cue (same location of pre-cue/post-cue as response cue). IN, invalid cue (different location of pre-cue/post-cue than response cue). Pre, pre-cue presented before the stimuli. Post, post-cue presented after the stimuli. Valid cues induced more accurate and faster responses (there was no speed-accuracy trade-off). Error bars, ± 1 SEM across participants.*

In each condition, we also calculated reaction time for each participant separately (**Figure 1B**, bottom row). A three-way repeated measures ANOVA revealed faster reaction times for valid than invalid cues (F(1,4)=62.3, p=0.001). There was no significant difference between exogenous and endogenous cues (F(1,4)=2.7, p=0.17), or between pre- and post-cues (F(1,4)=1.7, p=0.27). Two significant interactions indicated that the differences between valid and invalid cues (F(1,4)=16.2, p=0.02) and between pre- and post-cues (F(1,4)=8.1, p=0.047) were more pronounced for endogenous attention than for exogenous attention.

These behavioral results, which are consistent with previous findings (Lu and Dosher, 1998; Carrasco et al., 2000; Dosher and Lu, 2000; Ling and Carrasco, 2006b; Giordano et al., 2009; Pestilli et al., 2009; Herrmann et al., 2010), show that attention improved orientation

discrimination (d' and reaction time), with no evidence of a speed-accuracy trade-off, and similarly for both types of attention and for both pre- and post-cues. Thus, the behavioral effects confirm the successful manipulation of endogenous and exogenous attention, for both pre- and post-cues, consistently across individuals.

Attentional modulation of perceptual and post-perceptual information processing in visual cortex

Visual areas were mapped in each participant following retinotopic mapping procedures (**Figure 2**, left panel) and a targeted stimulus localizer (**Figure 2**, right panel), and regions of interest (ROIs) were selected based on previous literature: V1, V2, V3 (for V2 and V3, ventral and dorsal ROIs were averaged), V3A, hV4 and LO1 (e.g., Engel et al., 1994; Sereno et al., 1995; Engel et al., 1997; Larsson and Heeger, 2006; Wandell et al., 2007).

Activity was higher for contralateral than ipsilateral brain areas. For each ROI, we measured the fMRI response amplitudes for each type of attention – exogenous and endogenous – and each cueing condition – pre- and post-cueing, for the contralateral and ipsilateral side to the cued (attended) location. We analyzed the fMRI responses evoked by each type of attention in the contralateral and ipsilateral brain regions relative to the cue location (**Supplemental Figure 1**). ANOVAs indicated that there was higher contralateral than ipsilateral activity across brain areas (endogenous: $F(1,4)=59.9$, $p=0.0015$; exogenous: $F(1,4)=218.8$, $p=0.0001$). For both types of attention, this difference was more pronounced for valid than invalid cues (endogenous: $F(1,4)=8.6$, $p=0.04$; exogenous: $F(1,4)=21.1$, $p=0.01$). In the following analyses, we then concentrate on fMRI activity in the contralateral ROIs to the attended location.

261 ***Figure 2. Retinotopic mapping and stimulus localizer of a representative participant.*** *A*
262 *flattened representation of the right hemisphere occipital pole.* ***Left,*** *map of polar angle. The color*
263 *of each voxel in the map represents the response phase of the time series elicited by a rotating*
264 *wedge stimulus.* ***Right,*** *stimulus localizer. The black outlines correspond to the retinotopic areas*
265 *(defined for each participant based on their polar angle maps). The color of each voxel indicates*
266 *the phase of the response to the grating stimuli presented in the lower left visual field.*

267

268   fMRI responses correlated with behavioral variability. We evaluated the degree to which

269   inter-individual variability in behavioral performance co-varied with variability in fMRI responses.

270   We computed the correlation (**Figure 3A**), across participants, ROIs, types of attention and cue-

271   ing conditions, between the fMRI responses (percent change in image intensity) and behavioral

272   performance accuracy (d') . We found a positive correlation between fMRI activity and d' (Pearson

273   correlation r = 0.23, p = 0.003), which was present for both attention types (endogenous: Pearson

274   correlation r = 0.3, p = 0.003; exogenous: r = 0.2, p = 0.02). We further showed that fMRI activity

275   was overall higher for valid than invalid trials (t(4) = 4.0, p = 0.0163, Cohen's d = 1.78). This was

276   the case for each participant (**Figure 3B**). Together, these results are consistent with previous

277   findings showing that fMRI response correlates with behavioral variability (Ress et al., 2000;

278   Giesbrecht et al., 2006; Liu et al., 2007a), and that attentional orienting increases fMRI signal in

279   early visual cortex (for review Beck and Kastner, 2014).

Figure 3. Inter-individual fMRI response. A. *Positive correlation between fMRI response and behavioral performance (Pearson correlation r = 0.23, p = 0.003). Each dot indicates fMRI response amplitude measured from one ROI and one participant, separately for each attentional condition (pre and post-cue, valid and invalid).* **B.** *fMRI response difference between valid and invalid trials combined across all conditions and ROIs for each participant (colored dots), and across participants (black dot; average significantly positive: t(4) = 4.0, p = 0.0163, Cohen's d = 1.78). Error bar, ± 1 SEM across participants.*

We then tested our two novel predictions: (1) attentional modulation of fMRI stimulus-evoked activity increases along the visual hierarchy for endogenous pre-cueing, but is constant for exogenous pre-cueing; and (2) attentional modulation of fMRI stimulus-evoked activity for endogenous post-cueing is constant along the visual hierarchy, whereas there is no attentional modulation for exogenous post-cueing. We measured the differences in activity between valid and invalid trials for each type of attention and for pre- and post-cues, across the hierarchy of visual cortical areas (**Figure 4**). Regression analyses showed a significant increase along the hierarchy of the visual areas of the activity difference evoked by valid and invalid trials for the endogenous pre-cueing condition (F = 25.8, p = 0.007, R^2 = 0.87), but not for the exogenous pre-cueing condition (F = 2.8, p = 0.168, R^2 = 0.41), endogenous post-cueing (F = 3.0, p = 0.161, R^2 = 0.42) or exogenous post-cueing conditions (F = 0.1, p = 0.774, R^2 = 0.02). The differential effects of these regression analyses are consistent with post-hoc t-tests performed for each condition and ROI (**Figure 5**).

Figure 4. Specificity of pre and post-cueing for endogenous and exogenous attention. *The difference in fMRI response between valid and invalid is plotted separately for pre- and post-cue conditions for each ROI. *, Statistically significant regression analysis ($p < 0.05$).*

For both endogenous attention ($p < 0.05$ for V3a, V4 and LO1; trend ($p = 0.077$) for V3; t-tests) and exogenous attention ($p < 0.05$ for V3, V3A, hV4 and LO1; trend ($p = 0.074$) for V2; t-tests) pre-cues elicited greater fMRI activity for valid than invalid cues (**Figure 5**, top-left and top-right panels). Furthermore, for endogenous attention ($p < 0.05$ for V2, V3, V3A, hV4 and LO1; trend ($p = 0.077$) for V1; t-tests; **Figure 5**, bottom-left panel), but not for exogenous attention (all $p > 0.1$; t-tests; **Figure 5**, bottom-right panel), post-cues elicited greater fMRI activity for valid than invalid cues in occipital areas (detailed statistics are presented in Supplementary Table 1). Taken together, these results confirm both of our predictions.

Figure 5. Single ROI responses for pre and post-cueing and for both endogenous and exogenous attention. *The difference in fMRI response between valid and invalid is plotted separately for pre and post-cue conditions for each ROI. *, Statistically significant difference between valid and invalid, separately for pre and post-cueing (p < 0.05). #, trend (p < 0.1). Error bars on plots are ± 1 SEM.*

These effects were consistently observed across participants (**Figure 6**). fMRI responses were significantly larger for valid than invalid cues, for both endogenous and exogenous cues, and for both pre- and post-cues. A three-way repeated measures ANOVA (2 exogenous/endogenous x 2 pre/post-cue x 6 ROIs) of the difference between valid and invalid conditions revealed significant main effects of exogenous/endogenous condition (F = 10.0, p = 0.0341) and of ROI (F = 5.5, p = 0.0024), as well as an interaction between pre/post-cue and ROI (F = 4.1, p = 0.0103).

329
Figure 6. Specificity of each visual ROI for endogenous and exogenous attention. *fMRI*
response amplitude was measured for each attentional condition. V: valid cue condition (target
location matches the location indicated by pre-cue/post-cue). IN: invalid cue condition (target
location at the opposite location relative to the pre-cue/post-cue). Pre: pre-cue presented before
the grating stimuli. Post: post-cue presented after gratings. All four conditions in the contralateral
ROI. Distractor (Dist): all conditions averaged in the ipsilateral ROI. Each colored point corre-
sponds to the data of one participant. The black points represent the average across all 5 partic-
*ipants. *, Statistically significant difference between valid and invalid, separately for pre and post-*
cueing (p < 0.05). #, trend (p < 0.1). Detailed statistics are presented in Supplementary Table 1.
Error bars (vertical lines) are plotted on the difference between valid and invalid and represent ±
1 SEM across participant.

DISCUSSION

This is the first study to compare pre and post-orienting, and reorienting, of endogenous and exogenous attention in visual cortex, while concurrently assessing visual performance using well-established psychophysical protocols to manipulate attention. The fact that the same participants performed the same orientation discrimination task with the same stimuli and task difficulty under different attentional manipulations enabled us to isolate the fMRI activity induced by each type of attention during orienting and reorienting.

Previous studies comparing endogenous and exogenous attention conditions state various attentional effects in early visual areas (see **Table 1**). Some report differential effects between these attention conditions in the right LO and MT (Kincade et al., 2005), others in the cuneus (Peelen et al., 2004; Mayer et al., 2004; Kincade et al., 2005), the occipital gyrus (SOG, MOG, LOG; Mayer et al., 2004; Kincade et al., 2005; Natale et al., 2006), the fusiform area (Kincade et al., 2005; Natale et al., 2006; Esterman et al., 2008), and the TOS, LG and SC (Natale et al., 2006). This lack of clear picture regarding the differential impact of endogenous and exogenous spatial attention is likely due to the methodological differences and limitations we discussed in the Introduction (see also **Table 1**). In any case, it is often assumed that the effects of endogenous and exogenous attention are the same in visual occipital areas (Corbetta and Shulman, 2002; Peelen et al., 2004; Corbetta et al., 2008; Chica et al., 2013; Beck and Kastner, 2014).

To further our knowledge of the neural correlates of attention, we investigated both attentional orienting (valid cueing) and reorienting (invalid cueing), critical in an ever-changing environment (e.g. Dugué et al., 2016; 2017b; Senoussi et al., 2019). Furthermore, given ubiquitous performance tradeoffs between attended (benefits) and unattended (costs) locations (e.g. Pestilli and Carrasco, 2005; Giordano et al., 2009; Montagna et al., 2009; Herrmann et al., 2010; Fernández et al., 2019), we assessed activity at both attended (contralateral ROI) and unattended (ipsilateral ROI) locations. Finally, we investigated how attentional effects varied as a function of pre- and post-cueing, thus contrasting the neural correlates of perceptual and post-perceptual processing of information.

There was an overall positive correlation between performance in the orientation discrim-

ination task and the degree of attentional modulation in fMRI activity. This result was expected,

indicating that as discriminability increases so does the attentional modulation in fMRI activity.

But only very few studies on the effect of spatial attention in early visual areas have reported

such a correlation (Ress et al., 2000; Giesbrecht et al., 2006; Liu et al., 2007a). The behavioral

effects obtained in the scanner are consistent with psychophysical studies. The enhanced per-

formance brought about by the valid, but uninformative peripheral pre-cue is consistent with an

automatic, bottom-up involuntary capture of exogenous attention (e.g. Dosher and Lu, 2000; Car-

rasco et al., 2000; Carrasco et al., 2004; Pestilli and Carrasco, 2005; Giordano et al., 2009; Herr-

mann et al., 2010; Fernández et al., 2019). The enhanced performance brought about by the

valid, informative central precue is consistent with a top-down, voluntary deployment of endoge-

nous attention (e.g. Dosher and Lu, 2000; Ling and Carrasco, 2006c; Giordano et al., 2009; Liu

et al., 2009; Herrmann et al., 2010; Poletti et al., 2017; Jigo and Carrasco, 2018).

In the endogenous attention condition, there was an increase in attentional modulation of

stimulus-evoked activity along the hierarchy of visual areas. There is no consensus regarding the

visual hierarchy beyond area V3 (e.g. Harvey and Dumoulin, 2011; Harvey and Dumoulin, 2016;

Michalareas et al., 2016; Yildirim et al., 2018; Fritsche et al., 2020), especially regarding V3A.

However, most authors agree that hV4 precedes LO1 (Harvey and Dumoulin, 2011; Harvey and

Dumoulin, 2016; Michalareas et al., 2016; Yildirim et al., 2018). In any case, our data are con-

sistent with either a strict hierarchy or with V3A and hV4 being at the same level. Such a pattern

is consistent with previous studies suggesting that endogenous attention is a top-down modula-

tion from frontal and parietal areas feeding back to visual cortex, with diminishing effects in earlier

visual areas (Kastner et al., 1999; Maunsell and Cook, 2002; Kastner and Pinsk, 2004; Chica et

al., 2013). Inconsistent with previous studies (e.g. Boynton et al., 1999; Brefczynski and DeYoe,

1999; Somers et al., 1999; Herrmann et al., 2010; Pestilli et al., 2011), there was no evidence for

attentional modulation in V1. It might be that attentional modulation of V1 activity is more variable

than other visual cortical areas, making it harder to detect (see also Kastner et al., 1999; Liu et

al., 2005). Methodological differences between this and previous studies may have contributed

to weakening the effect of attention in V1. The accrual time in the current endogenous condition

was relatively short (1300 ms in the valid condition and 500 ms in the invalid condition) compared

to previous studies investigating endogenous, voluntary attention, in which the cue and/or stimuli

were presented for a long duration to maximize BOLD measurements (e.g. Boynton et al., 1999;

Brefczynski and DeYoe, 1999; Somers et al., 1999; Pestilli et al., 2011). We chose the minimum

SOA (317 ms) at which performance benefits plateau to have as comparable conditions as pos-

sible to the accrual time with the SOA for exogenous attention (117 ms). This short accrual time

may have limited the effects of attentional feedback to V1.

In the exogenous attention condition, in contrast to the endogenous attention, the atten-

tional modulation was approximately constant across the visual hierarchy. Some previous studies

have reported a similar effect (Müller and Kleinschmidt, 2007; Müller and Ebeling, 2008), others

a decrease (Heinen et al., 2011), and yet others an increase (Liu et al., 2005; Mulckhuyse et al.,

2011) across the visual areas. This difference might be explained by different task parameters.

For example, in the Liu et al. (2005) study, participants knew which of the two stimuli was the

target they had to discriminate as soon as the stimuli were displayed; one stimulus was vertical

and the other was tilted to the left or the right. In the present study, both stimuli were inde-

pendently tilted and participants did not know which one was the target and which one was the

distractor until later when the response cue appeared.

Unlike in the endogenous pre-cueing condition in which the attention effect increased

along the processing stream, for the endogenous post-cueing effect there was no evidence that

it varied across these visual areas. The constant effect in the post-cue condition could be due to

the contribution of two counteracting factors: (1) the fMRI response evoked by the stimulus in

early visual areas may decrease along the visual hierarchy (Kay et al., 2013b); (2) the top-down

modulations from frontal and parietal areas feedback to visual cortex with diminishing effects in

earlier visual areas (Chica and Lupiáñez, 2009; Kastner et al., 1999; Pestilli et al., 2011).

In the exogenous condition, there was no significant post-cueing effect on early visual

areas. This result is consistent with that of Liu et al. (2005), who while evaluating exogenous

attention effects on occipital cortex included a post-cue condition to rule out sensory

contamination of the cue (i.e. sensory response evoked by the cue itself) contributing to the enhanced BOLD activity found in their pre-cue condition. In addition to ruling out a possible sensory contamination, the present results show that, in contrast to endogenous attention, exogenous attention does not aid in the selective readout of information.

The ROI-based analysis that we followed here enabled us to compare contralateral and ipsilateral modulation of BOLD activity, thus providing additional information regarding the differences in processing dynamics of both types of attention. We observed a larger difference between contralateral and ipsilateral areas for the valid than the invalid cueing condition. This effect could be due to the fact that for the former, participants were attending to the same location throughout the trial, whereas for the latter, when the response cue did not match the pre-cue, participants had to switch their spatial attention to the opposite stimulus location, thus activity at that new location would be accumulated for less time. For instance, for endogenous attention, for the valid condition participants had been processing the target for almost 500 ms before the response cue appeared. When the response cue matched the pre-cue, participants continued processing and reading out the signal from that location for up to 800 ms (they were not allowed to give an answer before the end of the response cue period). But when the response cue did not match the pre-cue, then participants had to switch after 500 ms to the other location (ipsilateral) thus accumulating less activity. Similarly, the accumulation time for the invalid cue condition in exogenous attention was only about 300 ms. This accrual time explanation could also account for the larger difference between contralateral and ipsilateral for pre-cues than post-cues, i.e. there is a 300 ms accumulation when the exogenous pre-cue is invalid, while only 100 ms when the post-cue in invalid. Likewise, the larger modulatory effect for endogenous relative to exogenous attention is consistent with the difference in accrual time.

The results of the present study complement our recent study (Dugué et al., 2017a) in which we demonstrated that sub-regions of the Temporo-Parietal Junction (TPJ) that respond specifically to visual stimuli are more active when attention needs to be spatially reoriented (invalid cueing) than when attention remains at the cued location (valid cueing), and that partially overlapping specific visual sub-regions mediate reorienting after orienting of endogenous or

exogenous attention. Together, these two studies provide a comprehensive investigation of en-

dogenous and exogenous attention, and pave the way for rigorous psychophysics informed, neu-

roimaging studies of covert, spatial attention. Here, we concentrated the analysis on visual corti-

cal areas in the occipital lobe because the slice prescription covered only a limited portion of the

brain. The present findings further our knowledge of the neurophysiological bases of covert at-

tention and have implications for models of visual attention, which should consider not only the

similarities, but also the differences in the orienting and reorienting of endogenous and exoge-

nous attention in occipital areas reported here.

In conclusion, the present results show some similarities and reveal important differences

in the specific neural correlates of endogenous and exogenous attention on early vision: An in-

creasing modulation of fMRI activity for pre-cueing endogenous attention, but constant modula-

tion for exogenous attention, along the hierarchy of visual occipital areas, as well as a reliable

and constant modulation of fMRI activity for post-cueing endogenous attention in occipital areas

but not for exogenous attention. These results suggest that endogenous attention facilitates both

the encoding and the readout of visual information whereas exogenous attention only facilitates

the encoding of information.

MATERIALS and METHODS

The behavioral methods employed in this study and the behavioral results are the same

as those we reported in a recent study, in which we compared activity in TPJ during orienting and

reorienting of endogenous and exogenous attention (Dugué et al., 2017a). To maximize the ef-

fects of these two types of attention, i.e. the benefits at the attended location and concurrent

costs at the unattended location, we used optimal spatial and temporal parameters (reviews by

Carrasco, 2011; Carrasco and Barbot, 2015). To enable direct comparison between endogenous

and exogenous attention, the same participants performed the same orientation discrimination

task under both types of attention. The fMRI methods employed in this study are the same as

those used in that study (Dugué et al., 2017a), but here, instead of analyzing TPJ activity, we

analyzed activity in occipital areas.

Participants

Five participants (two male and three female, 24-30 years-old) participated in the study. They all had normal or corrected-to-normal vision. The University Committee on Activities Involving Human Subjects at New York University approved the experimental protocol (IRB # 10-7094), and participants provided written informed consent. Our study used single-participant ROI-based analysis, and thus had a small sample size. The same sample size that has been used in many fMRI studies in our labs (e.g. Dugué et al., 2017a; Herrmann et al., 2010; Liu et al., 2005; Liu et al., 2006; Liu et al., 2007a; Pestilli et al., 2011; Freeman et al., 2013; Wang et al., 2014; Brouwer et al., 2015), as well as in other labs (e.g. McMains and Somers, 2004; Moerel et al., 2016; Mackey et al., 2017; Poltoratski et al., 2017; Bloem and Ling, 2019). Each participant performed nine scanning sessions: one session to obtain a set of three high-resolution anatomical volumes, two sessions for retinotopic mapping, three sessions for the exogenous attention condition and three sessions for the endogenous attention condition (with the order counterbalanced among participants). Participants performed several practice sessions outside the scanner prior to the first scanning session of each attention condition.

Stimuli

Stimuli were generated on a Macintosh computer using the MGL toolbox (Gardner et al., 2018b) in MATLAB (MathWorks). Stimuli were presented on a flat-panel display (NEC, LC-XG250 MultiSync LCD 2110; refresh rate: 60 Hz; resolution: 1024 x 768 pixels) positioned at the rear of the scanner bore and housed in a Faraday box with an electrically conductive glass front. The display, calibrated and gamma corrected using a linearized lookup table, was at a viewing distance of 172 cm from the participant, and visible through an angled mirror attached to the head coil. A central, white fixation cross (0.3°) was presented throughout the experiment. The two stimuli were two 4-cpd gratings windowed by raised cosines (3° of diameter; 7% contrast), one in each bottom quadrant (5° horizontal eccentricity; –2.65° altitude). Both endogenous cues and exogenous cues were white rectangles (0.7°). The endogenous cues appeared adjacent to the fixation cross indicating one of the two lower quadrants (0.35° horizontal eccentricity from the edge of the fixation cross, and 0.35° altitude). The exogenous cues appeared adjacent to an

upcoming grating stimulus, vertically aligned with the stimulus and above the horizontal meridian (1° away from the edge of the grating stimulus; and the edge of the cue 4.44° horizontal eccentricity from the edge of the fixation cross).

Behavioral procedure

An exogenous attention condition trial lasted 1700 ms, whereas an endogenous attention condition trial lasted 1900 ms, the only difference being the stimulus-onset asynchronies (SOA) between the cue and the display; the timing of all the visual stimuli was the same in both attention conditions (**Figure 1**; the display is not at scale for illustration purposes). In the pre-cue condition (40% of the trials), a cue preceded the two gratings. In 40% the post-cue condition, the cue followed the presentation of the gratings. In 'cue-only' trials (10% of the trials), the gratings were not presented. In 'blank' trials (10% of the trials), neither a cue nor the gratings were presented. These trials were then included in the GLM analysis to model the contribution of the visual signal produced by the cue (see MRI procedure). For both pre-cue and post-cue trials, participants were asked to press one of two keys to report the orientation of a target grating, i.e., clockwise or counter-clockwise compared to vertical. Participants pressed a third key in the case of cue-only and blank trials.

In both exogenous and endogenous condition, cues were presented for 67 ms, indicating either the bottom left or right quadrant of the screen. The inter-stimulus interval (ISI) between the cue and the grating stimuli was 50 ms for exogenous and 250 ms for endogenous conditions, resulting in SOA of 117 ms and 317 ms. We used the same timings for pre- and post-cue conditions (e.g. Kinchla et al., 1995; Carrasco and Yeshurun, 1998; Liu et al., 2005; Pestilli et al., 2011). These delays are optimal to manipulate exogenous and endogenous attention, while keeping the trial duration as similar as possible, and have been shown to maximize the behavioral consequences of each attention condition (Nakayama and Mackeben, 1989; Mackeben and Nakayama, 1993; Liu et al., 2007b; Müller, 2014).

The behavioral effects of endogenous attention are sustained (e.g. Ling and Carrasco, 2006c) and thus, as shown in ERP studies (e.g. Seiss et al., 2009), are still present in later brain

activity. Additionally, during 300 ms following cue onset, the brain responses elicited by exoge-

nous and endogenous cues differ (for review Carrasco 2011). The two grating stimuli were then

displayed for 50 ms. For the postcue trials we kept the timings of cue and stimuli constant but

inverted the order of their presentation (e.g. Kinchla et al., 1995; Carrasco and Yeshurun, 1998;

Liu et al., 2005; Pestilli et al., 2011). A response cue, presented for 800 ms at the end of the trial

after both the cue and the stimuli had disappeared, indicated which one of the two gratings was

the target (50% of the trials on the right and the remaining 50% and on the left). The maximum

delay between the offset of the grating stimuli and the onset of the response cue was shorter

(~400 ms max in the endogenous condition) than typically associated with a demand for working

memory (>600 ms; Phillips, 1974). Immediately following each trial, a change of color of the fix-

ation cross provided visual feedback to the participants, i.e. green for correct or red for incorrect

responses. The fixation cross did not change color if participants had missed the response win-

dow, i.e. if they had not pressed any key after 530 ms.

In the exogenous attention condition, a peripheral cue was presented, which was not in-

formative regarding the target location or orientation. When the cue location matched the target

location, it was considered a valid trial (50% of the trials), otherwise it was considered an invalid

trial (the remaining 50% of the trials). In the endogenous attention condition, a central cue pointed

to either the left or right quadrant. The cue was informative of the target location but not its ori-

entation (75% valid trials and 25% invalid trials). Participants were informed of this validity. It is

important to notice that cue validity does not affect cueing effectiveness for exogenous attention,

although it does so do endogenous attention (e.g. Sperling and Melchner, 1978; Kinchla, 1980;

Giordano et al., 2009).

Endogenous and exogenous attention conditions were performed in separate sessions to

ensure optimal manipulation of each attention system. Participants first performed two practice

sessions outside the scanner before the first session of each attentional scanning condition. To

equate task difficulty for both attention conditions, using a staircase procedure, the tilt of the

target grating was adjusted for each participant to achieve ~80% correct performance in the valid

trials in each attention condition. In each of the six experimental scanning sessions (three

sessions of exogenous attention and three of endogenous), participants performed 14 runs of 40

trials each, as well as a run of stimulus localizer (see MRI procedure). The tilt was then adjusted

between runs to maintain overall performance at ~80% correct.

Eye position was monitored during all scanning sessions using an infrared video camera

system (Eyelink 1K, SR Research, Ottawa, Ontario, http://www.sr-research.com/EL_1000.html).

Trials in which the participants blinked or broke fixation (1.5° radius from central fixation) at any

point from fixation onset to response cue offset were identified and regressed separately in the

MRI analysis, and removed from the behavioral analysis (13% ± 4% of the trials on average

across all participants).

MRI Procedure

Scanning. Imaging was conducted on a 3T Siemens Allegra head-only scanner (Erlan-

gen, Germany), using a Siemens NM-011 head coil (to transmit and receive) to acquire anatom-

ical images, a receive-only 8-channel surface coil array (Nova Medical, Wilmington, MA) to ac-

quire functional images. To minimize participants' head movements, padding was used.

For each participant, three high-resolution anatomic images were acquired in one scan-

ning session, using a T1-weighted magnetization-prepared rapid gradient echo (MPRAGE) se-

quence (FOV = 256 x 256 mm; 176 sagittal slices; 1 x 1 x 1 mm voxels), and were co-registered

and averaged. Using FreeSurfer (public domain software; http://surfer.nmr.mgh.harvard.edu),

the gray matter was segmented from these averaged anatomical volumes. All subsequent anal-

yses were constrained only to voxels that intersected gray matter.

T2*-weighted echo-planar imaging sequence (i.e. functional images; TR = 1750 ms; TE

= 30 ms; flip angle = 90°) measured blood oxygen level-dependent (BOLD) changes in image

intensity (Ogawa et al., 1990). In each volume, 28 slices covered the occipital and posterior pa-

rietal lobes and were oriented 45° to the calcarine sulcus (FOV = 192 x 192 mm; resolution = 2

x 2 x 2.5 mm; no gap). To align functional images from different sessions to the same high-reso-

lution anatomical volume for each participant, we acquired an additional T1-weighted anatomical

images in the same slices as the functional images (spin echo; TR = 600 ms; TE = 9.1 ms;

flip angle = 90°; resolution = 1.5 × 1.5 × 3 mm) during each scanning session, and used them in

a robust image registration algorithm.

MRI data pre-processing. Imaging data were analyzed in MATLAB, using mrTools (Gard-

ner et al., 2018a) and custom software. To allow longitudinal magnetization to reach steady state,

the first eight volumes of each run were discarded. Spatial distortion was corrected using the B0

static magnetic field measurements performed in each session. The functional data were then

motion corrected, the linear trend was removed, and a temporal high-pass filter was applied (cut-

off: 0.01 Hz) to remove slow drifts and low-frequency noise in the fMRI signal.

Retinotopic mapping. We followed well-established conventional traveling-wave, phase-

encoded methods. Using clockwise and counter-clockwise rotating checkerboard wedges, we

measured phase maps of polar angle. Using contracting and expending checkerboard rings, we

measured eccentricity maps (Engel et al., 1994; Sereno et al., 1995; Engel et al., 1997; Larsson

and Heeger, 2006; Wandell et al., 2007). **Figure 2** (left panel) shows the visual areas that were

drawn by hand on flattened surface of the brain, following published conventions (Engel et al.,

1997; Larsson and Heeger, 2006; Wandell et al., 2007; Liu et al., 2007a).

Stimulus localizer. In each scanning session of the main experiment, participants com-

pleted one stimulus localizer run (6 runs overall, 4 min each). A run consisted of 16 cycles (17.5 s)

of a block alternation protocol between stimulus on (8.75 s) and stimulus off (8.75 s). Participants

only had to fixate the central cross throughout each run. The stimuli were at the same location,

and of the same size and spatial frequency as those in the main experiment, except at full con-

trast and their phase and orientation changed randomly every 200 ms to avoid adaptation. To

define the cortical representation of the gratings, we then averaged the data across the 6 runs

and followed the same methods as for the retinotopic mapping. Voxels that responded positively

during the blocks when the grating stimuli were presented were used to restrict each retinotopic

ROI. The fMRI time series from each voxel were fit to a sinusoid. To be conservative, only voxels

whose best-fit sinusoid had a phase value between 0 and pi, and a coherence between the best-

fit sinusoid and the time series greater than 0.2 were included in the ROI (**Figure 2**, right panel).

Analysis performed without restricting the ROI to this coherence level yielded similar results.

Event-related analysis. fMRI time series were averaged across voxels in each ROI (sep-

arately for each hemisphere) and then concatenated across runs. The data were denoised using

GLMDenoise (Kay et al., 2013a), and fMRI response amplitudes were computed using linear

regression, with twelve regressors: 8 combinations of right and left valid and invalid pre- and

post-cue, right and left cue-only, blank (no cue nor stimulus) and eye-movements (blink or broken

fixation). For each ROI in each hemisphere, the resulting fMRI response amplitudes (for correct

trials only) were then averaged across participants.

Acknowledgments: This work was supported by: NIH RO1-EY019693 to MC and DJH; NIH
RO1-EY027401 to MC; the FYSSEN Foundation and the Philippe Foundation to LD; and the
Center for Brain Imaging of New York University. We want to thank members of the Carrasco lab
for constructive comments on the manuscript. The authors declare no conflicts of interest.

REFERENCES

Anton-Erxleben K, Carrasco M (2013) Attentional enhancement of spatial resolution: linking behavioural and neurophysiological evidence. Nat Rev Neurosci 14:188–200.

Anton-Erxleben K, Henrich C, Treue S (2007) Attention changes perceived size of moving visual patterns. Journal of Vision 7:1–9.

Barbot A, Carrasco M (2017) Attention Modifies Spatial Resolution According to Task Demands. Psychological Science 28:285–296

Barbot A, Landy MS, Carrasco M (2012) Differential effects of exogenous and endogenous attention on second-order texture contrast sensitivity. Journal of Vision 12:1–15.

Beauchamp MS, Petit L, Ellmore TM, Ingeholm J, Haxby JV (2001) A parametric fMRI study of overt and covert shifts of visuospatial attention. Neuroimage 14:310–321.

Beck DM, Kastner S (2009) Top-down and bottom-up mechanisms in biasing competition in the human brain. Vision Res 49:1154–1165.

Beck DM, Kastner S (2014) Neural systems for spatial attention in the human brain: Evidence from neuroimaging in the framework of biased competition. The Oxford Handbook of Attention:253–288.

Bloem IM, Ling S (2019) Normalization governs attentional modulation within human visual cortex. Nature Communications 10:5660–10.

Bowling JT, Friston KJ, Hopfinger JB (2019) Top-down versus bottom-up attention differentially modulate frontal–parietal connectivity. Hum Brain Mapp 9:1116.

Boynton GM, Demb JB, Glover GH, Heeger DJ (1999) Neuronal basis of contrast discrimination. Vision Res 39:257–269.

Brefczynski JA, DeYoe EA (1999) A physiological correlate of the "spotlight" of visual attention. Nat Neurosci 2:370–374.

Brouwer GJ, Arnedo V, Offen S, Heeger DJ, Grant AC (2015) Normalization in human somatosensory cortex. J Neurophysiol 114:2588–2599.

Buracas GT, Boynton GM (2007) The effect of spatial attention on contrast response functions in human visual cortex. Journal of Neuroscience 27:93–97.

Buschman TJ, Miller EK (2009) Serial, Covert Shifts of Attention during Visual Search Are Reflected by the Frontal Eye Fields and Correlated with Population Oscillations. Neuron 63:386–396.

Busse L, Katzner S, Treue S (2008) Temporal dynamics of neuronal modulation during exogenous and endogenous shifts of visual attention in macaque area MT. Proc Natl Acad Sci USA 105:16380–16385.

Carrasco M (2011) Visual attention: the past 25 years. Vision Res 51:1484–1525.

Carrasco M, Barbot A (2015) How attention affects spatial resolution. Cold Spring Harbor Symposia on Quantitative Biology 79:149–160.

Carrasco M, Evert DL, Chang I, Katz SM (1995) The eccentricity effect: target eccentricity affects performance on conjunction searches. Perception & Psychophysics 57:1241–1261.

Carrasco M, Giordano AM, McElree B (2006) Attention speeds processing across eccentricity:
Feature and conjunction searches. Vision Res 46:2028–2040.

Carrasco M, Marie Giordano A, McElree B (2004) Temporal performance fields: visual and at-
tentional factors. Vision Res 44:1351–1365.

Carrasco M, McElree B (2001) Covert attention accelerates the rate of visual information pro-
cessing. PNAS 98:5363–5367.

Carrasco M, Penpeci-Talgar C, Eckstein M (2000) Spatial covert attention increases contrast
sensitivity across the CSF: support for signal enhancement. Vision Res 40:1203–1215.

Carrasco M, Yeshurun Y (1998) The contribution of covert attention to the set-size and eccen-
tricity effects in visual search. Journal of Experimental Psychology: Human Perception and
Performance 24:673–692.

Carrasco M, Yeshurun Y (2009) Covert attention effects on spatial resolution. Prog Brain Res
176:65–86.

Chica AB, Bartolomeo P, Lupiáñez J (2013) Two cognitive and neural systems for endogenous
and exogenous spatial attention. Behavioural Brain Research 237:107–123.

Chica AB, Lupiáñez J (2009) Effects of endogenous and exogenous attention on visual pro-
cessing: an Inhibition of Return study. Brain Res 1278:75–85.

Corbetta M, Patel G, Shulman GL (2008) The reorienting system of the human brain: from envi-
ronment to theory of mind. Submitted 58:306–324.

Corbetta M, Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the
brain. Nat Rev Neurosci 3:201–215.

Correa Á, Triviño M, Pérez-Dueñas C, Acosta A, Lupiáñez J (2010) Temporal preparation, re-
sponse inhibition and impulsivity. Brain and Cognition 73:222–228.

Culham JC, Cavanagh P, Kanwisher NG (2001) Attention Response Functions: Characterizing
Brain Areas Using fMRI Activation during Parametric Variations of Attentional Load. Neuron
32:737–745.

Doricchi F, Macci E, Silvetti M, Macaluso E (2010) Neural correlates of the spatial and expectancy
components of endogenous and stimulus-driven orienting of attention in the Posner task.
Cereb Cortex 20:1574–1585.

Dosher BA, Lu ZL (2000) Mechanisms of perceptual attention in precuing of location. Vision Res
40:1269–1292.

Dugué L, Merriam EP, Heeger DJ, Carrasco M (2017a) Specific Visual Subregions of TPJ Medi-
ate Reorienting of Spatial Attention. Cereb Cortex:1–16.

Dugué L, Roberts M, Carrasco M (2016) Attention Reorients Periodically. Current Biology
26:1595–1601.

Dugué L, Xue AM, Carrasco M (2017b) Distinct perceptual rhythms for feature and conjunction
searches. Journal of Vision 17:22–22.

Engel SA, Glover GH, Wandell BA (1997) Retinotopic organization in human visual cortex and
the spatial precision of functional MRI. Cereb Cortex 7:181–192.

Engel SA, Rumelhart DE, Wandell BA, Lee AT, Glover GH, ED C, Shadlen MN (1994) fMRI of
human visual cortex. Nature 369:525.

Esterman M, Prinzmetal W, DeGutis J, Landau AN, Hazeltine E, Verstynen T, Robertson L (2008)
Voluntary and involuntary attention affect face discrimination differently. Neuropsychologia
46:1032–1040.

Fernández A, Li H-H, Carrasco M (2019) How exogenous spatial attention affects visual repre-
sentation. Journal of Vision 19:4.

Freeman J, Heeger DJ, Merriam EP (2013) Coarse-scale biases for spirals and orientation in
human visual cortex. Journal of Neuroscience 33:19695–19703.

Fuller S, Park Y, Carrasco M (2009) Cue contrast modulates the effects of exogenous attention
on appearance. Vision Res 49:1825–1837.

Gardner JL, Merriam EP, Movshon JA, Heeger DJ (2008) Maps of Visual Space in Human Oc-
cipital Cortex Are Retinotopic, Not Spatiotopic. J Neurosci 28:3988–3999.

Gardner JL, Merriam EP, Schluppeck D, Besle J, Heeger DJ (2018a) mrTools: Analysis and
visualization package for functional magnetic resonance imaging data. Available at:
http://doi.org/10.5281/zenodo.1299483.

Gardner JL, Merriam EP, Schluppeck D, Larsson J (2018b) MGL: Visual psychophysics stimuli
and experimental design package. Available at: http://doi.org/10.5281/zenodo.1299497.

Geng JJ, Vossel S (2013) Re-evaluating the role of TPJ in attentional control: contextual updat-
ing? Neurosci Biobehav Rev 37:2608–2620.

Giesbrecht B, Weissman DH, Woldorff MG, Mangun GR (2006) Pre-target activity in visual cortex
predicts behavioral performance on spatial and feature attention tasks. Brain Res 1080:63–
72.

Giordano AM, McElree B, Carrasco M (2009) On the automaticity and flexibility of covert atten-
tion: a speed-accuracy trade-off analysis. Journal of Vision 9:30.1–.10.

Gobell J, Carrasco M (2005) Attention alters the appearance of spatial frequency and gap size.
Psychological Science 16:644–651.

Hahn B, Ross TJ, Stein EA (2006) Neuroanatomical dissociation between bottom–up and top–
down processes of visuospatial selective attention. Neuroimage 32:842–853.

Heinen K, Ruff CC, Bjoertomt O, Schenkluhn B, Bestmann S, Blankenburg F, Driver J, Chambers
CD (2011) Concurrent TMS-fMRI reveals dynamic interhemispheric influences of the right
parietal cortex during exogenously cued visuospatial attention. European Journal of Neuro-
science 33:991–1000.

Herath P, Klingberg T, Young J, Amunts K, Roland PE (2001) Neural Correlates of Dual Task
Interference Can be Dissociated from Those of Divided Attention: an fMRI Study | Cerebral
Cortex | Oxford Academic. Cereb Cortex 11:796–805.

Herrmann K, Montaser-Kouhsari L, Carrasco M, Heeger DJ (2010) When size matters: attention
affects performance by contrast or response gain. Nat Neurosci 13:1554–1559.

Hulme OJ, Friston KF, Zeki S (2009) Neural correlates of stimulus reportability. Cognitive Neu-
roscience, Journal of 21:1602–1610.

Jigo M, Carrasco M (2018) Attention alters spatial resolution by modulating second-order pro-
cessing. Journal of Vision 18:2.

Kastner S, Pinsk MA (2004) Visual attention as a multilevel selection process. Cognitive, Affec-
tive, & Behavioral Neuroscience 4:483–500.

Kastner S, Pinsk MA, De Weerd P, Desimone R, Ungerleider LG (1999) Increased Activity in
Human Visual Cortex during Directed Attention in the Absence of Visual Stimulation. Neuron
22:751–761.

Kay K, Rokem A, Winawer J, Dougherty R, Wandell B (2013a) GLMdenoise: a fast, automated
technique for denoising task-based fMRI data. Front Neurosci 7:1–15.

Kay KN, Winawer J, Rokem A, Mezer A, Wandell BA (2013b) A two-stage cascade model of
BOLD responses in human visual cortex. Diedrichsen J, ed. PLoS Comput Biol 9:e1003079.

Kim YH, Gitelman DR, Nobre AC, Parrish TB, LaBar KS, Mesulam MM (1999) The large-scale
neural network for spatial attention displays multifunctional overlap but differential asym-
metry. Neuroimage 9:269–277.

Kincade JM, Abrams RA, Astafiev SV, Shulman GL, Corbetta M (2005) An event-related func-
tional magnetic resonance imaging study of voluntary and stimulus-driven orienting of atten-
tion. Journal of Neuroscience 25:4593–4604.

Kinchla, R. A. (1980). The measurement of attention. In R. S. Nikerson (Ed.), Attention and per-
formance *VIII*, 213-238.

Kinchla RA, Chen Z, Evert D (1995) Precue effects in visual search: data or resource limited?
Perception & Psychophysics 57:441–450.

Larsson J, Heeger DJ (2006) Two retinotopic visual areas in human lateral occipital cortex. Jour-
nal of Neuroscience 26:13128–13142.

Li X, Lu Z-L, Tjan BS, Dosher BA, Chu W (2008) Blood oxygenation level-dependent contrast
response functions identify mechanisms of covert attention in early visual areas. Proc Natl
Acad Sci USA 105:6202–6207.

Ling S, Carrasco M (2006a) Sustained and transient covert attention enhance the signal via dif-
ferent contrast response functions. Vision Res 46:1210–1220.

Ling S, Carrasco M (2006b) Sustained and transient covert attention enhance the signal via dif-
ferent contrast response functions. Vision Res 46:1210–1220.

Ling S, Carrasco M (2006c) When sustained attention impairs perception. Nat Neurosci 9:1243–
1245.

Liu T, Abrams J, Carrasco M (2009) Voluntary attention enhances contrast appearance. Psycho-
logical Science 20:354–362.

Liu T, Heeger DJ, Carrasco M (2006) Neural correlates of visual vertical meridian asymmetry. J
Vis 6:1294–1306.

Liu T, Larsson J, Carrasco M (2007a) Feature-based attention modulates orientation-selective
responses in human visual cortex. Neuron 55:313–323.

Liu T, Pestilli F, Carrasco M (2005) Transient attention enhances perceptual performance and
FMRI response in human visual cortex. Neuron 45:469–477.

Liu T, Stevens ST, Carrasco M (2007b) Comparing the time course and efficacy of spatial and feature-based attention. Vision Res 47:108–113.

Lu Z-L, Li X, Tjan BS, Dosher BA, Chu W (2011) Attention Extracts Signal in External Noise: A BOLD fMRI Study. J Cogn Neurosci 23:1148–1159.

Lu ZL, Dosher BA (1998) External noise distinguishes attention mechanisms. Vision Res 38:1183–1198.

Luck SJ, Hillyard SA, Mouloua M, Woldorff MG, Clark VP, Hawkins HL (1994) Effects of spatial cuing on luminance detectability: psychophysical and electrophysiological evidence for early selection. Journal of Experimental Psychology: Human Perception and Performance 20:887–904.

Luo TZ, Maunsell JHR (2015) Neuronal Modulations in Visual Cortex Are Associated with Only One of Multiple Components of Attention. Neuron 86:1182–1188.

Mackeben M, Nakayama K (1993) Express attentional shifts. Vision Res 33:85–90.

Mackey WE, Winawer J, Curtis CE (2017) Visual field map clusters in human frontoparietal cortex. Elife 6:2704.

Martínez-Trujillo JC, Treue S (2002) Attentional Modulation Strength in Cortical Area MT Depends on Stimulus Contrast. Neuron 35:365–370.

Maunsell JHR, Cook EP (2002) The role of attention in visual processing. Philosophical Transactions of the Royal Society of London B: Biological Sciences 357:1063–1072.

Mayer AR, Dorflinger JM, Rao SM, Seidenberg M (2004) Neural networks underlying endogenous and exogenous visual–spatial orienting. Neuroimage 23:534–541.

McAdams CJ, Maunsell JHR (1999) Effects of Attention on Orientation-Tuning Functions of Single Neurons in Macaque Cortical Area V4. J Neurosci 19:431–441.

McMains SA, Somers DC (2004) Multiple spotlights of attentional selection in human visual cortex. Neuron 42:677–686.

Meyer KN, Du F, Parks E, Hopfinger JB (2018) Exogenous vs. endogenous attention: Shifting the balance of fronto-parietal activity. Neuropsychologia 111:307–316.

Mitchell JF, Sundberg KA, Reynolds JH (2009) Spatial Attention Decorrelates Intrinsic Activity Fluctuations in Macaque Area V4. Neuron 63:879–888.

Moerel D, Ling S, Jehee JFM (2016) Perceptual learning increases orientation sampling efficiency. Journal of Vision 16:36–36.

Montagna B, Pestilli F, Carrasco M (2009) Attention trades off spatial acuity. Vision Res 49:735–745.

Mulckhuyse M, Belopolsky AV, Heslenfeld D, Talsma D, Theeuwes J (2011) Distribution of attention modulates salience signals in early visual cortex. Op de Beeck HP, ed. PLoS ONE 6:e20379.

Müller MM (2014) Neural Mechanisms of Feature-Based Attention. In: Cognitive Electrophysiology of Attention, pp 123–135. Elsevier.

Müller NG, Ebeling D (2008) Attention-modulated activity in visual cortex—More than a simple
"spotlight." Neuroimage 40:818–827.

Müller NG, Kleinschmidt A (2007) Temporal Dynamics of the Attentional Spotlight: Neuronal Cor-
relates of Attentional Capture and Inhibition of Return in Early Visual Cortex. The Journal of
Cognitive Neuroscience 19:587–593.

Murray SO (2008) The effects of spatial attention in early human visual cortex are stimulus inde-
pendent. Journal of Vision 8:1–11.

Nachmias J (1967) Effect of Exposure Duration on Visual Contrast Sensitivity with Square-Wave
Gratings. J Opt Soc Am, JOSA 57:421–427.

Natale E, Marzi CA, Girelli M, Pavone EF, Pollmann S (2006) ERP and fMRI correlates of en-
dogenous and exogenous focusing of visual-spatial attention. Eur J Neurosci 23:2511–2521.

Nakayama K, Mackeben M (1989) Sustained and transient components of focal visual attention.
Vision Res 29:1631–1647.

Nobre AC, Coull JT, Maquet P, Frith CD, Vandenberghe R, Mesulam MM (2004a) Orienting at-
tention to locations in perceptual versus mental representations. Cognitive Neuroscience,
Journal of 16:363–373.

Nobre AC, Coull JT, Maquet P, Frith CD, Vandenberghe R, Mesulam MM (2004b) Orienting at-
tention to locations in perceptual versus mental representations. Cognitive Neuroscience,
Journal of 16:363–373.

O'Connor DH, Fukui MM, Pinsk MA (2002) Attention modulates responses in the human lateral
geniculate nucleus. Nature 5:1203–1209.

Ogawa S, Lee TM, Kay AR, Tank DW (1990) Brain magnetic resonance imaging with contrast
dependent on blood oxygenation. Proc Natl Acad Sci USA 87:9868–9872.

Peelen MV, Heslenfeld DJ, Theeuwes J (2004) Endogenous and exogenous attention shifts are
mediated by the same large-scale neural network. Neuroimage 22:822–830.

Pestilli F, Carrasco M (2005) Attention enhances contrast sensitivity at cued and impairs it at
uncued locations. Vision Res 45:1867–1875.

Pestilli F, Carrasco M, Heeger DJ, Gardner JL (2011) Attentional enhancement via selection and
pooling of early sensory responses in human visual cortex. Neuron 72:832–846.

Pestilli F, Ling S, Carrasco M (2009) A population-coding model of attention's influence on con-
trast response: Estimating neural effects from psychophysical data. Vision Res 49:1144–
1153.

Phillips WA (1974) On the distinction between sensory storage and short-term visual memory.
Perception & Psychophysics 16:283–290.

Poletti M, Rucci M, Carrasco M (2017) Selective attention within the foveola. Nat Neurosci
20:1413–1417.

Poltoratski S, Ling S, McCormack D, Tong F (2017) Characterizing the effects of feature salience
and top-down attention in the early visual system. J Neurophysiol 118:564–573.

Posner MI, Snyder CR, Davidson BJ (1980) Attention and the detection of signals. J Exp Psychol
109:160–174.

Reed AV (1973) Speed-Accuracy Trade-Off in Recognition Memory. Science 181:574–576.

Ress D, Backus BT, Heeger DJ (2000) Activity in primary visual cortex predicts performance in
a visual detection task. Nat Neurosci 3:940–945.

Reynolds JH, Heeger DJ (2009) The Normalization Model of Attention. Neuron 61:168–185.

Reynolds JH, Pasternak T, Desimone R (2000) Attention increases sensitivity of V4 neurons.
Neuron 26:703–714.

Rosen AC, Rao SM, Caffarra P, Scaglioni A, Bobholz JA, Woodley SJ, Hammeke TA, Cunning-
ham JM, Prieto TE, Binder JR (1999) Neural basis of endogenous and exogenous spatial
orienting. A functional MRI study. Cognitive Neuroscience, Journal of 11:135–152.

Ruff CC, Kristjánsson Á, Driver J (2007) Readout from iconic memory and selective spatial at-
tention involve similar neural processes. Psychological Science 18:901–909.

Ruff DA, Cohen MR (2014) Attention can either increase or decrease spike count correlations in
visual cortex. Nat Neurosci 17:1591–1597.

Seiss E, Driver J, Eimer M (2009) Effects of attentional filtering demands on preparatory ERPs
elicited in a spatial cueing task. Clin Neurophysiol 120:1087–1095.

Senoussi M, Moreland JC, Busch NA, Dugué L (2019) Attention explores space periodically at
the theta frequency. Journal of Vision 19:1–17.

Sereno MI, Dale AM, Reppas JB, Kwong KK, Belliveau JW, Brady TJ, Rosen BR, Tootell R (1995)
Borders of multiple visual areas in humans revealed by functional magnetic resonance im-
aging. Science 268:889–893

Sergent C, Ruff CC, Barbot A, Driver J, Rees G (2011) Top-down modulation of human early
visual cortex after stimulus offset supports successful postcued report. The Journal of Cog-
nitive Neuroscience 23:1921–1934.

Sergent C, Wyart V, Babo-Rebelo M, Cohen L, Naccache L, Tallon-Baudry C (2013) Cueing
Attention after the Stimulus Is Gone Can Retrospectively Trigger Conscious Perception. Cur-
rent Biology 23:150–155.

Silvetti M, Lasaponara S, Lecce F, Dragone A, Macaluso E, Doricchi F (2015) The Response of
the Left Ventral Attentional System to Invalid Targets and its Implication for the Spatial Ne-
glect Syndrome: a Multivariate fMRI Investigation. Cereb Cortex 1–2.

Somers DC, Dale AM, Seiffert AE, Tootell RB (1999) Functional MRI reveals spatially specific
attentional modulation in human primary visual cortex. PNAS 96:1663–1668.

Sperling G, Melchner MJ (1978) The attention operating characteristic: examples from visual
search. Science 202:315–318.

Thibault L, van den Berg R, Cavanagh P, Sergent C (2016) Retrospective Attention Gates Dis-
crete Conscious Access to Past Sensory Stimuli. PLoS ONE 11:e0148504.

Wandell BA, Dumoulin SO, Brewer AA (2007) Visual field maps in human cortex. Neuron 56:366–
383.

Wang F, Chen M, Yan Y, Zhaoping L, Li W (2015) Modulation of Neuronal Responses by Exog-
enous Attention in Macaque Primary Visual Cortex. Journal of Neuroscience 35:13419–
13429.

Wang HX, Merriam EP, Freeman J, Heeger DJ (2014) Motion direction biases and decoding in human visual cortex. Journal of Neuroscience 34:12601–12615.

Wickelgren WA (1977) Speed-accuracy tradeoff and information processing dynamics. Acta Psychologica 41:67–85.

Williford T, Maunsell JHR (2006) Effects of Spatial Attention on Contrast Response Functions in Macaque Area V4. J Neurophysiol 96:40–54.

Yashar A, Lamy D (2011) Refining the dual-stage account of intertrial feature priming: does motor response or response feature matter? Atten Percept Psychophys 73:2160–2167.

Yeshurun Y, Carrasco M (1998) Attention improves or impairs visual performance by enhancing spatial resolution. Nature 396:72–75.

Yeshurun Y, Montagna B, Carrasco M (2008) On the flexibility of sustained attention and its effects on a texture segmentation task. Vision Research 48:80–95.

Yeshurun Y, Rashal E (2010) Precueing attention to the target location diminishes crowding and reduces the critical distance. Journal of Vision 10:16–16.