Differential impact of endogenous and exogenous attention on activity in human visual cortex

Laura Dugué[1,2,3,4], Elisha P. Merriam[1,2,5], David J. Heeger[1,2] & Marisa Carrasco[1,2]

[1] Department of Psychology, New York University
[2] Center for Neural Science, New York University
[3] Université de Paris, CNRS, Integrative Neuroscience and Cognition Center, F-75006 Paris, France
[4] Institut Universitaire de France, Paris, France
[5] Laboratory of Brain and Cognition, NIMH/NIH, Bethesda, MD

Running title: endogenous and exogenous attention

Corresponding author:
Laura Dugué
Current address: 45 rue des Saints-Pères 75006 Paris, FRANCE
laura.dugue@u-paris.fr

ABSTRACT (150 words)

How do endogenous (voluntary) and exogenous (involuntary) attention modulate activity in visual cortex? Using ROI-based fMRI analysis, we measured fMRI activity for valid and invalid trials (target at cued/un-cued location, respectively), pre- or post-cueing endogenous or exogenous attention, while participants performed the same orientation discrimination task. We found stronger modulation in contralateral than ipsilateral visual regions, and higher activity in valid- than invalid-trials. For endogenous attention, modulation of stimulus-evoked activity due to a pre-cue increased along the visual hierarchy, but was constant due to a post-cue. For exogenous attention, modulation of stimulus-evoked activity due to a pre-cue was constant along the visual hierarchy, but was not modulated due to a post-cue. These findings reveal that endogenous and exogenous attention distinctly modulate activity in visuo-occipital areas during orienting and reorienting; endogenous attention facilitates both the encoding and the readout of visual information whereas exogenous attention only facilitates the encoding of information.

Keywords: exogenous attention, endogenous attention, fMRI activity, occipital visual areas, top-down processing, bottom-up processing.

INTRODUCTION

Spatial, covert visual attention is the selective processing of visual information in space, without change in gaze. Attention can be allocated voluntarily –endogenous attention– or involuntarily –exogenous attention. Endogenous and exogenous attention have different temporal dynamics; endogenous takes about 300 ms to be deployed and can be sustained at will whereas exogenous attention takes only about 100 ms to be deployed and it is transient (for reviews see [1,2]). These two types of attention often have similar perceptual consequences (reviews by [1,2]), but notable exceptions indicate that whereas endogenous attention acts in a flexible way, exogenous attention acts in an inflexible or automatic way. For instance: (a) The benefits and costs in perception (discriminability and speed of information accrual) scale with cue validity for endogenous but not for exogenous attention (e.g. [3]); (b) The effects of covert attention on contrast sensitivity often differ for endogenous and exogenous attention (e.g. [4-8]); and (c) For a texture segmentation task in which heightened spatial resolution improves or impairs performance as a function of target eccentricity, endogenous attention improves performance across eccentricity, whereas exogenous attention improves performance at peripheral locations where resolution is poor, it hampers performance where resolution is already high for the task at hand (e.g. [9-14]).

Basic visual processes, such as contrast sensitivity and spatial resolution, are mediated by activity in early visual cortex, and are altered by covert attention (reviews by [1,2,15,16]). Specifically, single-unit studies in monkeys have demonstrated effects of endogenous attention in occipital areas (e.g. [17-24]). Additionally, fMRI studies have shown that endogenous attention causes a baseline shift in early visual areas (e.g. [25-31]; review by [32]) and increases the dynamic range of fMRI responses [33,34]. Comparatively, little is known about the effect of exogenous attention on visual areas both from single-unit studies [35,36] and fMRI studies (for reviews see [1,15]).

Since Corbetta and Schulman's seminal review (2002) [37] on the neural bases of endogenous and exogenous attention, there has been emphasis on characterizing networks of brain regions within the frontal and parietal lobes (for reviews see [32,38,39]). Yet, there remains considerable debate regarding the dissociation between dorsal regions, for endogenous attention, and

ventral regions, for exogenous attention (e.g. [39,40]), including the role of the temporo-parietal

junction (TPJ; [41-44]). Classically, researchers have described endogenous attention as a top-

down process, and exogenous attention as a bottom-up process [1,37-39,45-47]. This characteriza-

tion originated in psychophysics experiments, and was then studied using fMRI, in which the

two types of attention have been often investigated separately (for reviews [1,32,38,45]).

Studies of endogenous and exogenous attention focusing on parietal and frontal areas

have shown that the two types of attention differentially modulate fronto-parietal connectivity

[39,48]. For example, there are critical differences in the temporal order of neural responses in

frontal and parietal cortex in monkeys between these attention conditions, i.e. frontal activity

precedes parietal activity during endogenous attentional orienting, whereas parietal activity pre-

cedes frontal activity during exogenous orienting. Critically, it is often assumed that the effects

of endogenous and exogenous attention are the same in striate and extra-striate areas

[32,37,38,49,50].

A number of important considerations limit the conclusions that may be drawn from the

few studies that have directly compared independent effects of visual exogenous and endoge-

nous spatial attention [39,50-52]:

(1) The effect of attention on behavioral performance was assessed in some studies with a

detection task using RT as their only dependent variable [50,52], in which performance may

differ due to speed of processing, discriminability or criterion [53-55] and/or motor behavior

[56,57].

(2) In the studies in which performance was assessed in a discrimination task using RT [51],

small RT differences were reported, which could have resulted from speed of pro-

cessing, discriminability, or criterion factors [53-55].

(3) In the studies in which accuracy was not assessed [39,50,52], it is not possible to know

whether task difficulty was the same for both types of attention, and task difficulty can

interact with the strength of fMRI activity [58,59].

(4) For the studies in which eye position was not monitored while participants performed the

task in the scanner [39,50-52], the results could be due to covert attention, overt attention or

both [1,60,61].

(5) Given that exogenous attention is a fast, transient process (for reviews see [1,2,15,16]), it

was inappropriately manipulated in the studies in which long stimulus onset asynchro-

nies (SOA) were used [50,51,62-64] making the comparison between the two attention condi-

tions problematic.

(6) Except for two studies, one interested in attentional modulation in the Fusiform Face

Area (FFA) [64] and the other in the TPJ [41], statistical parametric mapping was applied to

group averaged data to identify regions of the brain that were active during task perfor-

mance. Some found significant activity in the occipital pole [50-52], but attentional modula-

tion of fMRI activity was not systematically assessed across different visual areas.

Given all these methodological limitations, it is unknown how these two types of atten-

tion affect neural activity in each individual visuo-occipital area, and how modulation of activity

in visual cortex is linked to changes in perceptual performance. (See Dugué and colleagues [41],

who published a table summarizing these and other methodological problems for studies re-

garding covert attention and TPJ activation).

Typically, covert attention is manipulated by presenting a pre-cue, prior to the target –

and the aforementioned studies also did so (see **Table 1**). However, endogenous post-cues,

presented after target offset, can also improve performance by affecting the information readout

[65-68] and modulate fMRI activity in early visual areas [25,67-69]. Exogenous post-cues also affect

performance in some tasks [70,71], but not in others [72-75], and the only fMRI study evaluating post-

cues in exogenous attention in occipital cortex showed no such modulation [76]. Critically, no sin-

gle study has compared visual cortex activity with post-cues in endogenous and exogenous at-

tention.

Study	Cueing manipulation		Task	Dependent Variable	Analysis	Cortical areas	Eye tracking
	Endogenous	Exogenous					
Studies including Occipital areas							
Detection tasks							
Mayer et al. (2004)[52]	central arrow 70% valid	luminance change of peripheral placeholder; 50% valid	X	RT	Group-averaged	**Occipital:** *Cuneus, MOG, SOG* + Frontal Temporo-Parietal	No
	SOA=100 or 800 ms						
Peelen et al. (2004)[50]	central arrow 75% valid	brightening of peripheral place-holder; 50% valid;	square	RT	Group-averaged	**Occipital:** *Cuneus* + Frontal Temporo-Parietal	No
	SOA=550 ms						
Natale et al. (2006)[63]	single peripheral rectangular frame; 100% valid;	five peripheral rectangular frames; 20% valid;	a black and white checkerboard	RT *for 4 out of 8 participants*	Group-averaged	**Occipital:** *Fusiform, TOS, LG, SC, LOG* + Frontal Temporo-Parietal	No
	SOA=8, 8.15, 8.3, 8.45 or 8.6 s						
Discrimination tasks							
Kincade et al. (2005)[51]	brightening of half of central fixation diamond 75% valid	color singleton in an array of colored squares 50% valid	T vs. L	RT (% correct)	Group-averaged	**Occipital:** *LO, MT, Cuneus, Fusiform, SOG* + Frontal Temporo-Parietal	No
	SOA=2160 ms						
Esterman et al. (2008)[64]	color change of a peripheral placeholder 75% valid	50% valid	face identity	RT (% correct)	ROI-based (FFA) + Group-averaged	**Occipital:** *FFA* + Frontal Temporo-Parietal	Yes
	SOA=300 ms						
Current study	central bar; 75% invalid SOA=310 ms	peripheral bar; 50% valid; SOA=110 ms	orientation	d' (RT)	ROI-based	**Occipital:** *V1, V2, V3, V3A, hV4, LO1*	Yes
Studies NOT including Occipital areas							
Detection tasks							
Rosen et al. (1999)[62]	central arrow; 80% valid;	peripheral dot; 50% valid;	filling of a square	RT	Group-averaged	Frontal Temporo-Parietal	No
	SOA=400, 550 or 700 ms						
Discrimination tasks							
Kim et al. (1999)[77]	thickening of part of central fixation diamond; 80% valid; SOA=200, 400 or 800 ms	luminance change of peripheral placeholder; 50% valid; SOA= 100, 150 or 200 ms	X vs. +	RT (% correct)	Group-averaged	Frontal Temporo-Parietal	No
Meyer et al. (2018)[78]	color change of central fixation 83% valid SOA=700–1000 ms	whitening of peripheral place-holder 50% valid SOA=100–300 ms	central color of a checkerboard	Inverse effi-ciency = RT/%correct	Group-averaged	Frontal Temporo-Parietal	No
Dugué et al. (2017)[41]	central bar; 75% invalid; SOA=310 ms	peripheral bar; 50% valid; SOA=110 ms	orientation	d' (RT)	ROI-based	Temporo-Parietal	Yes
Bowling et al. (2019)[39]	color change of central fixation; 83% valid; SOA=700–1000 ms	whitening of peripheral place-holder; 50% valid; SOA=100–300 ms	central color of a checkerboard	Inverse effi-ciency = RT/%correct	Group-averaged	Frontal Temporo-Parietal	No

Table 1. fMRI studies comparing endogenous and exogenous attention independently in

human participants. _For each study, we report the cueing manipulation for each attention con-_

dition (SOA: Stimulus Onset Asynchrony is the duration of the cue + interval before onset of the

target), the task performed by the participants, the dependent variable reported in the publica-

tion, whether the analysis was based on group-averaging or ROI-based on single-participants,

the cortical areas reported in the publications and showing significant activation due to cue

and/or target, and whether or not eye data were monitored in the scanner or used for the sub-

sequent fMRI analysis. IOG: Inferior Occipital Gyrus; LG: Lingual Gyrus; LO: Lateral Occipital

area; LOG: Lateral Occipital Gyrus; MOG: Middle Occipital Gyrus; MT: Middle Temporal area;

SC: Striate Cortex; SOG: Superior Occipital Gyrus; TOS: Temporal-Occipital Sulcus.

Here, with the same participants, task, stimuli and task difficulty for all attention manipu-

lations, we systematically tested the following four predictions: (1) Pre-cueing should induce an

attentional modulation of fMRI activity (for reviews see [1,15,32,38,39]), higher in the valid than the

invalid condition in which attention needs to be reoriented to the opposite location (e.g. [41,49,79,80])

to perform the task. (2) Both endogenous and exogenous pre- and post-cueing effects should

be stronger in visual regions contralateral to the attended hemifield (e.g. [25,76]). (3) Pre-cueing

endogenous attention, but not exogenous attention, should increase activity modulations along

the visual hierarchy (e.g. higher in V4 than in V1; [25,30]; for reviews [32,38]). For endogenous atten-

tion, a top-down process, modulations from higher-order, fronto-parietal attentional regions

would send feedback information to visual cortex with diminishing effects in earlier visual areas

[30,31,38,81-84]. (4) Post-cueing endogenous [25,67-69], but not exogenous [76], attention should induce

attentional modulation of fMRI activity in early visual areas. Voluntary, endogenous attention

would facilitate reading out perceptual information [41], and modulate its processing [25,67-69]. The

only fMRI study assessing the effects of post-cueing exogenous attention in occipital cortex

found no attentional modulation of fMRI activity [76]; some behavioral studies report post-cueing

effects [70,71] but others found no such effects [72-75].

To test these four predictions, we measured fMRI activity and compared the effects of

endogenous and exogenous attention in early (V1, V2, V3) and intermediate (V3A, hV4, LO1)

visual areas while the same participants performed the same task –a two-Alternative Forced

Choice (2-AFC) orientation discrimination task, contingent upon contrast sensitivity [4,5,8,79,85-88].

We used a fully-crossed design: two attention conditions –endogenous or exogenous atten-

tional orienting– and two types of cueing –pre- or post-cue. We evaluated fMRI activity at both

the attended and the un-attended locations, given the ubiquitous performance tradeoffs at at-

tended (benefits) and unattended (costs) locations compared to a neutral condition (e.g. [3,27,87-

92]), and the importance of evaluating both the orienting and reorienting of attention, critical in an

ever-changing environment [41,79,80,93].

This is the first study to systematically evaluate and directly compare how pre and post-

orienting, and reorienting, of endogenous and exogenous attention modulate neural activity in

visual cortex to affect behavior. The results indicate that these two types of spatial covert atten-

tion distinctly modulate activity in individual retinotopic visual cortical areas. These differences

in activity are consonant with differential engagement of top-down and bottom-up processes

and their respective temporal dynamics. These results suggest that endogenous attention facili-

tates both the encoding and the readout of visual information whereas exogenous attention only

facilitates the encoding of information.

RESULTS

Endogenous and exogenous attention improve performance

Participants performed a 2-AFC orientation-discrimination task under two attentional con-

ditions (exogenous or endogenous attention), when the cue was presented either before (pre-

cue) or after (post-cue) the grating stimuli (see Methods), while their brain activity was meas-

ured with fMRI (**Figure 1**).

In each condition, we calculated performance accuracy (d'), as the main dependent varia-

ble, for each participant separately (**Figure 1B**, top row). A three-way repeated measures

2x2x2 ANOVA (exogenous/endogenous x valid/invalid x pre/post-cue) revealed higher perfor-

mance for valid than invalid cues (F(1,4)=23.6, p=0.008), that the effect of exogenous and en-

dogenous cues were statistically indistinguishable (F(1,4)<1), and that there was no significant

difference between pre- and post-cues (F(1,4)<1). None of the two and three-way interactions

were significant (F<1).

In each condition, we also calculated reaction time (RT), as a secondary dependent varia-

ble for each participant separately (**Figure 1B**, bottom row). Although the task is not optimal to

asse RT effects due to the long response cue window during which participants are not allowed

to provide their response, we assessed RT to rule out speed-accuracy tradeoffs. A three-way

repeated measures ANOVA revealed faster reaction times for valid than invalid cues

(F(1,4)=62.3, p=0.001). There was no significant difference between exogenous and endoge-

nous cues (F(1,4)=2.7, p=0.17), or between pre- and post-cues (F(1,4)=1.7, p=0.27). Two sig-

nificant interactions indicated that the differences between valid and invalid cues (F(1,4)=16.2,

p=0.02) and between pre- and post-cues (F(1,4)=8.1, p=0.047) were more pronounced for en-

dogenous attention than for exogenous attention.

These behavioral results, which are consistent with previous findings [3-6,8,27,79,86-88,90,94],

show that attention improved orientation discrimination (d' and reaction time), with no evidence

of a speed-accuracy trade-off, and similarly for both types of attention and for both pre- and

post-cues. Thus, the behavioral effects confirm the successful manipulation of endogenous and

exogenous attention, for both pre- and post-cues, consistently across individuals.

Attentional modulation of perceptual and post-perceptual information processing in vis-

ual cortex

Visual areas were mapped in each participant following retinotopic mapping procedures

(**Figure 2**, left panel) and a targeted stimulus localizer (**Figure 2**, right panel), and regions of

interest (ROIs) were selected based on previous literature: V1, V2, V3 (for V2 and V3, ventral

and dorsal ROIs were averaged), V3A, hV4 and LO1 (e.g. [95-98]).

Activity was higher for contralateral than ipsilateral brain areas. For each ROI, we meas-

ured the fMRI response amplitudes for each type of attention – exogenous and endogenous –

and each cueing condition – pre- and post-cueing, for the contralateral and ipsilateral side to

the cued (attended) location. We analyzed the fMRI responses evoked by each type of atten-

tion in the contralateral and ipsilateral brain regions relative to the cue location (**Supplemental**

Figure 1). ANOVAs indicated that there was higher contralateral than ipsilateral activity across

brain areas (endogenous: $F(1,4)=59.9$, $p=0.0015$; exogenous: $F(1,4)=218.8$, $p=0.0001$). For

both types of attention, this difference was more pronounced for valid than invalid cues (endog-

enous: $F(1,4)=8.6$, $p=0.04$; exogenous: $F(1,4)=21.1$, $p=0.01$). In the following analyses, we then

concentrate on fMRI activity in the contralateral ROIs to the attended location. **Supplemental**

Figure 2 shows mean fMRI responses for a representative participant for each attention condi-

tion for two contralateral ROIs (V2 and LO1) relative to the cue. Response amplitude was ob-

tained from the target areas indicated in **Figure 2B**.

We calculated an attentional modulation index: AMI = Vc / average (Vc + Ic + Vi + Ii),

which takes into account the overall response magnitude of each ROI. The AMI corresponds to

the ratio of the activity in contralateral ROIs for valid trials (Vc) over the average activity of con-

tralateral valid (Vc) and invalid trials (Ic), and distractor stimuli for pre and post-cue trials for ip-

silateral valid (Vi) and invalid trials (Ii). (Because some condition values were negative, we

added a fixed constant (0.2) to all condition values of each participant before computing the

AMI so that all resulting values were positive; see original values in **Figure 5.** Below we show

that the pattern of results is the same for different constant values, indicating that our results do

not depend on the specific constant value we selected). A ratio of 1 would indicate no atten-

tional modulation. The larger the AMI ratio, the higher the modulation in the valid trials. First,

we showed that the AMI was significantly higher than 1 when combined across all conditions

and ROIs ($t(4) = 6.7$, $p = 0.0026$, Cohen's $d = 2.99$), and this was the case for each participant.

These results are consistent with previous findings showing that attentional orienting increases

fMRI signal in early visual cortex (for review [32]).

We then tested our two novel predictions: (1) attentional modulation of fMRI stimulus-

evoked activity increases along the visual hierarchy for endogenous pre-cueing, but is constant

for exogenous pre-cueing; and (2) attentional modulation of fMRI stimulus-evoked activity for

endogenous post-cueing is constant along the visual hierarchy, whereas there is no attentional

modulation for exogenous post-cueing. We first calculated the differences in activity between

valid and invalid trials for each type of attention and for pre- and post-cues, across the hierar-

chy of visual cortical areas. fMRI (AMI) responses were significantly larger for valid than invalid

cues, for both endogenous and exogenous cues, and for both pre- and post-cues. A three-way

repeated measures ANOVA (2 exogenous/endogenous x 2 pre/post-cue x 6 ROIs) of the differ-

ence between valid and invalid conditions revealed significant main effects of exogenous/en-

dogenous condition ($F(1,4) = 33.8$, $p = 0.0043$) and of ROI ($F(5,4) = 3.3$, $p = 0.0258$), as well as

an interaction between pre/post-cue and ROI ($F(5,4) = 3.5$, $p = 0.0192$). Thus, we investigated

modulations of fMRI activity separately for pre and post-cueing attention conditions.

Linear mixed models [99] were then used to test: (1) an increase of fMRI activity along the

visual hierarchy for pre-cueing endogenous attention but not for exogenous attention, and (2)

post-cue attentional modulation of fMRI activity for post-cueing endogenous attention but not for

exogenous attention. We implemented two models, one for the pre-cueing and the other for the

post-cueing condition, to test the link between fMRI activity and the interaction between ROI

and endogenous/exogenous attention conditions. In each model, we entered as fixed effect the

ROI and the endogenous/exogenous conditions into the model, as well as their interaction. As

random effects, we had participant's intercepts and slopes for the effect of ROI and endoge-

nous/exogenous conditions. Visual inspection of residual plots did not reveal any obvious devi-

ations from homoscedasticity or normality. P-values were obtained by likelihood ratio tests. For

the pre-cueing condition, we observed a significant effect of ROI ($t(56)=3.5$, $p=0.0009$, esti-

mate=0.05 ± 0.01, standard error), as well as a significant interaction between ROI and endoge-

nous/exogenous conditions ($t(56)=-2.3$, $p=0.025$, estimate=-0.018 ± 0.008, standard error). For

the post-cueing condition, we observed a significant effect of the fixed effect intercept

($t(56)=23.3$, $p<0.0001$, estimate=11 ± 0.05, standard error). Finally, we performed independent

regression analyses on each pre-/post-cueing and endogenous/exogenous attention conditions (**Figure 3**). We observed a significant increase along the hierarchy of the visual areas of the activity difference evoked by valid and invalid trials for the endogenous pre-cueing condition ($F(4)=16.5$, $p=0.015$, $R^2=0.81$), but not for the exogenous pre-cueing condition ($F(4)=1.8$, $p=0.249$, $R^2=0.31$), endogenous post-cueing ($F(4)=2.2$, $p=0.216$, $R^2=0.35$) or exogenous post-cueing conditions ($F(4)=0.3$, $p=0.61$, $R^2=0.07$). See **Supplemental Figure 3** and **Supplemental Table 1 and 2,** which show the same pattern of results for different constant values, indicating that our results do not depend on the specific value of the constant we selected.

Together, the results of the linear mixed models and the regression analyses (**Figure 3**) showed differential effects between pre-/post-cueing and endogenous/exogenous attention conditions. The AMI increases along the hierarchy of the visual areas in pre-cueing of endogenous attention but not for pre-cueing of exogenous attention. Moreover, the AMI with post-cueing of endogenous attention is constant along the hierarchy (significant intercept), but absent for post-cueing of exogenous attention. These analyses are consistent with post-hoc t-tests performed for each condition and ROI in the AMI modulation (**Figure 4**). See **Supplemental Figure 4,** which shows the same pattern of results for the difference in fMRI activity (in percent signal change) between valid and invalid.

For both endogenous attention ($p < 0.05$ for V2, V3, V3a, V4 and LO1; one-tailed t-tests) and exogenous attention ($p < 0.05$ for V2, V3, V3A, hV4 and LO1; one-tailed t-tests) pre-cues elicited greater fMRI activity for valid than invalid cues (**Figure 4**, top-left and top-right panels). Furthermore, for endogenous attention ($p < 0.05$ for V1, V2, V3, V3A, hV4 and LO1; one-tailed t-tests; **Figure 4**, bottom-left panel), but not for exogenous attention (all $p > 0.05$; one-tailed t-tests; **Figure 4**, bottom-right panel), post-cues elicited greater fMRI activity for valid than invalid cues in occipital areas (detailed statistics are presented in **Supplementary Table 3**). Critically, these effects were consistently observed across participants (**Figure 5**). Taken together, these results confirm both of our predictions.

DISCUSSION

This is the first study to compare pre and post-orienting, and reorienting, of endogenous and exogenous attention in visual cortex, while concurrently assessing visual performance using well-established psychophysical protocols to manipulate attention. The fact that the same participants performed the same orientation discrimination task with the same stimuli and task difficulty under different attentional manipulations enabled us to isolate the fMRI activity induced by each type of attention during orienting and reorienting.

Previous studies comparing endogenous and exogenous attention conditions state various attentional effects in early visual areas (see **Table 1**). Some report differential effects between these attention conditions in the right LO and MT [51], others in the cuneus [50-52], the occipital gyrus (SOG, MOG, LOG; [51,52,63]), the fusiform area [51,63,64], and the TOS, LG and SC [63]. This lack of clear picture regarding the differential impact of endogenous and exogenous spatial attention is likely due to the methodological differences and limitations we discussed in the Introduction (see also **Table 1**). In any case, it is often assumed that the effects of endogenous and exogenous attention are the same in visual occipital areas [32,37,38,49,50].

Other regions in striate and extra-striate visual areas could be implicated in both endogenous and exogenous attention. For instance, area V6 has been reported to be involved in the shift of covert attention, in both monkeys [100,101] and humans [102]. Because this area has a large representation of the periphery [103], we would have needed a wider field of view than the one we had to reliably define it. Similarly, higher-level visual areas are also involved in shifts of covert spatial attention, such as IPS 0 to IPS 5 (e.g. [30,38]). However, given the goal of our study, and thus our slice prescription, we could not identify all IPS regions.

To further our knowledge of the neural correlates of attention, we investigated both attentional orienting (valid cueing) and reorienting (invalid cueing), critical in an ever-changing environment [41,79,80,93]. Furthermore, given ubiquitous performance tradeoffs between attended (benefits) and unattended (costs) locations (e.g. [3,27,87-92]), we assessed activity at both attended

(contralateral ROI) and unattended (ipsilateral ROI) locations. Finally, we investigated how at-

tentional effects varied as a function of pre- and post-cueing, thus contrasting the neural corre-

lates of perceptual and post-perceptual processing of information.

The behavioral effects obtained in the scanner are consistent with psychophysical stud-

ies. The enhanced performance brought about by the valid, but uninformative peripheral pre-

cue is consistent with an automatic, bottom-up involuntary capture of exogenous attention (e.g.

[3,4,7-9,12,14,27,46,47,53,76,87,88,91,107,108]). The enhanced performance brought about by the valid, in-

formative central precue is consistent with a top-down, voluntary deployment of endogenous

attention (e.g. [3-8,11,12,25,27,46,47,53,86,91,109-111]).

In the endogenous attention condition, there was an increase in attentional modulation of

stimulus-evoked activity along the hierarchy of visual areas. There is no consensus regarding

the visual hierarchy beyond area V3 (e.g. [104-108]), especially regarding V3A. However, most au-

thors agree that hV4 precedes LO1 [104-107]. In any case, our data are consistent with either a

strict hierarchy or with V3A and hV4 being at the same level. Such a pattern is consistent with

previous studies suggesting that endogenous attention is a top-down modulation from frontal

and parietal areas feeding back to visual cortex, with diminishing effects in earlier visual areas

[30,38,81,82]. Inconsistent with previous studies (e.g. [25,27,31,109,110]), there was no evidence for atten-

tional modulation in V1. It might be that attentional modulation of V1 activity is more variable

than other visual cortical areas, making it harder to detect (see also [30,76]). Methodological differ-

ences between this and previous studies may have contributed to weakening the effect of at-

tention in V1. The accrual time in the current endogenous condition was relatively short (1300

ms in the valid condition and 500 ms in the invalid condition) compared to previous studies in-

vestigating endogenous, voluntary attention, in which the cue and/or stimuli were presented for

a long duration to maximize BOLD measurements (e.g. [25,27,31,109,110]). We chose the minimum

SOA (317 ms) at which performance benefits plateau to have as comparable conditions as pos-

sible to the accrual time with the SOA for exogenous attention (117 ms). This short accrual time

may have limited the effects of attentional feedback to V1.

We note that the fact that we do not see attentional modulation in V1 with our protocol

does not necessarily imply that this area does not mediate covert attention. For instance, the

inconsistent findings in the EEG literature regarding whether spatial attention can modulate V1

activity during the initial wave of visual processing depends on several factors (e.g., location,

presence of distractors, attentional load [111]). Similarly, some fMRI papers have shown signifi-

cant modulation of spatial attention in V1 (e.g. [25,27,31,109,110]). Furthermore, a recent study [88] has

shown that TMS on the occipital pole (V1/V2) after attentional deployment eliminated the typical

benefits and costs brought about by exogenous attention at the respective attended and unat-

tended locations.

In the exogenous attention condition, in contrast to the endogenous attention, the atten-

tional modulation was approximately constant across the visual hierarchy. Some previous stud-

ies have reported a similar effect [112,113], others a decrease [114], and yet others an increase [76,115]

across the visual areas. This difference might be explained by different task parameters. For

example, in the Liu et al. (2005) study [76], participants knew which of the two stimuli was the tar-

get they had to discriminate while the stimuli were presented, as one stimulus was vertical and

the other was tilted to the left or the right. In the present study, both stimuli were independently

tilted and participants had to wait for the response cue to appear to know which one was the

target and which one was the distractor.

Unlike in the endogenous pre-cueing condition in which the attention effect increased

along the processing stream, for the endogenous post-cueing effect there was no evidence that

the attentional modulation varied across these visual areas. The constant effect in the post-cue

condition could be due to the contribution of two counteracting factors: (1) the fMRI response

evoked by the stimulus in early visual areas may decrease along the visual hierarchy [116]; (2) the

top-down modulations from frontal and parietal areas feedback to visual cortex with diminishing

effects in earlier visual areas [25,30,117], [30,38,81,82].

In the exogenous condition, there was no significant post-cueing effect on early visual ar-

eas. This result is consistent with that of Liu et al. (2005) [76], who while evaluating exogenous

attention effects on occipital cortex included a post-cue condition to rule out sensory contamina-

tion of the cue (i.e. sensory response evoked by the cue itself) contributing to the enhanced

BOLD activity found in their pre-cue condition. In addition to ruling out a possible sensory con-

tamination, the present results show that, in contrast to endogenous attention, exogenous at-

tention does not aid in the selective readout of information. Although there was no reliable mod-

ulation of exogenous post-cue on early and intermediate visual areas, in this study, it had an

effect on behavior. Such effect is sometimes [70,71] but not always [72-75] observed. It is thus possi-

ble that our imaging protocol was not sensitive enough or that this behavioral effect is sup-

ported by higher order regions.

The ROI-based analysis that we followed here enabled us to compare contralateral and

ipsilateral modulation of BOLD activity, thus providing additional information regarding the dif-

ferences in processing dynamics of both types of attention. We observed a larger difference be-

tween contralateral and ipsilateral areas for the valid than the invalid cueing condition. This ef-

fect could be due to the fact that for the former, participants were attending to the same location

throughout the trial, whereas for the latter, when the response cue did not match the pre-cue,

participants had to switch their spatial attention to the opposite stimulus location, thus activity at

that new location would be accumulated for less time. For instance, for endogenous attention,

for the valid condition participants could have been processing the target for almost 500 ms be-

fore the response cue appeared. When the response cue matched the pre-cue, participants

continued processing and reading out the signal from that location for up to 800 ms (they were

not allowed to give an answer before the end of the response cue period). But when the re-

sponse cue did not match the pre-cue, then participants had to switch after 500 ms to the other

location (ipsilateral) thus accumulating less activity. Similarly, the accumulation time for the in-

valid cue condition in exogenous attention was only about 300 ms. This accrual time explana-

tion could also account for the larger difference between contralateral and ipsilateral for pre-

cues than post-cues, i.e. there is a 300 ms accumulation when the exogenous pre-cue is inva-

lid, while only 100 ms when the post-cue is invalid. Likewise, the larger modulatory effect for

endogenous than exogenous attention is consistent with the difference in accrual time.

In a recent study [41] we demonstrated that sub-regions of the Temporo-Parietal Junction

(TPJ) that respond specifically to visual stimuli are more active when attention needs to be spa-

tially reoriented (invalid cueing) than when attention remains at the cued location (valid cueing),

and that partially overlapping specific visual sub-regions mediate such reorienting of endoge-

nous or exogenous attention. Both the present and the TPJ studies provide a comprehensive

investigation of endogenous and exogenous attention, and pave the way for rigorous psycho-

physics informed, neuroimaging studies of covert, spatial attention. Here, we concentrated the

analysis on visual cortical areas in the occipital lobe. The present findings further our

knowledge of the neurophysiological bases of covert attention and have implications for models

of visual attention, which should consider not only the similarities, but also the differences in the

orienting and reorienting of endogenous and exogenous attention in occipital areas reported

here.

CONCLUSION

The present results show some similarities and reveal important differences in the spe-

cific neural correlates of endogenous and exogenous attention on early vision: An increasing

modulation of fMRI activity for pre-cueing endogenous attention, but constant modulation for

exogenous attention, along the hierarchy of visual occipital areas, as well as a reliable and con-

stant modulation of fMRI activity for post-cueing endogenous attention in occipital areas but not

for exogenous attention. These results suggest that endogenous attention facilitates both the

encoding and the readout of visual information whereas exogenous attention only facilitates the

encoding of information.

MATERIALS and METHODS

The behavioral methods employed in this study and the behavioral results are the same as those we reported in a recent study, in which we compared activity in TPJ during orienting and reorienting of endogenous and exogenous attention [41]. To maximize the effects of these two types of attention, i.e. the benefits at the attended location and concurrent costs at the un-attended location, we used optimal spatial and temporal parameters (reviews by [1,2]). To enable direct comparison between endogenous and exogenous attention, the same participants per-formed the same orientation discrimination task under both types of attention. The fMRI meth-ods employed in this study are the same as those used in that study [41], but here, instead of an-alyzing TPJ activity, we analyzed activity in occipital areas.

Participants

Five participants (two male and three female, 24-30 years-old) participated in the study. They all had normal or corrected-to-normal vision. The University Committee on Activities In-volving Human Subjects at New York University approved the experimental protocol (IRB # 10-7094), and participants provided written informed consent. All methods were performed in ac-cordance with US regulations and the Declaration of Helsinki. Our study used single-participant ROI-based analysis, and thus had a small sample size. The same sample size that has been used in many fMRI studies in our labs (e.g. [25,27,41,76,118-122]), as well as in other labs (e.g. [123-127]). Each participant performed nine scanning sessions: one session to obtain a set of three high-resolution anatomical volumes, two sessions for retinotopic mapping, three sessions for the ex-ogenous attention condition and three sessions for the endogenous attention condition (with the order counterbalanced among participants). Participants performed several practice sessions outside the scanner prior to the first scanning session of each attention condition.

Stimuli

Stimuli were generated on a Macintosh computer using the MGL toolbox [128] in MATLAB (MathWorks). Stimuli were presented on a flat-panel display (NEC, LC-XG250 MultiSync LCD 2110; refresh rate: 60 Hz; resolution: 1024 x 768 pixels) positioned at the rear of the scanner

bore and housed in a Faraday box with an electrically conductive glass front. The display, cali-

brated and gamma corrected using a linearized lookup table, was at a viewing distance of 172

cm from the participant, and visible through an angled mirror attached to the head coil. A cen-

tral, white fixation cross (0.3°) was presented throughout the experiment. The two stimuli were

two 4-cpd gratings windowed by raised cosines (3° of diameter; 7% contrast), one in each bot-

tom quadrant (5° horizontal eccentricity; –2.65° altitude; 5.66° of eccentricity from the central

fixation to the stimulus center). Both endogenous cues and exogenous cues were white rectan-

gles (0.7°). The endogenous cues appeared adjacent to the fixation cross indicating one of the

two lower quadrants (0.35° horizontal eccentricity from the edge of the fixation cross, and 0.35°

altitude). The exogenous cues appeared adjacent to an upcoming grating stimulus, vertically

aligned with the stimulus and above the horizontal meridian (1° away from the edge of the grat-

ing stimulus; and the edge of the cue 4.44° horizontal eccentricity from the edge of the fixation

cross).

Behavioral procedure

An exogenous attention condition trial lasted 1700 ms, whereas an endogenous atten-

tion condition trial lasted 1900 ms, the only difference being the stimulus-onset asynchronies

(SOA) between the cue and the display; the timing of all the visual stimuli was the same in both

attention conditions (**Figure 1**; the display is not at scale for illustration purposes). In the pre-

cue condition (40% of the trials), a cue preceded the two gratings. In 40% the post-cue condi-

tion, the cue followed the presentation of the gratings. In 'cue-only' trials (10% of the trials), the

gratings were not presented. In 'blank' trials (10% of the trials), neither a cue nor the gratings

were presented. These trials were then included in the GLM analysis to model the contribution

of the visual signal produced by the cue (see MRI procedure). For both pre-cue and post-cue

trials, participants were asked to press one of two keys to report the orientation of a target grat-

ing, i.e., clockwise or counter-clockwise compared to vertical. Participants pressed a third key in

the case of cue-only and blank trials.

In both exogenous and endogenous condition, cues were presented for 67 ms, indicat-

ing either the bottom left or right quadrant of the screen. The inter-stimulus interval (ISI) be-

tween the cue and the grating stimuli was 50 ms for exogenous and 250 ms for endogenous

conditions, resulting in SOA of 117 ms and 317 ms. We used the same timings for pre- and

post-cue conditions (e.g. [25,65,76,129]). These delays are optimal to manipulate exogenous and en-

dogenous attention, while keeping the trial duration as similar as possible, and have been

shown to maximize the behavioral consequences of each attention condition [46,130-132].

The behavioral effects of endogenous attention are sustained (e.g. [133]) and thus, as

shown in ERP studies (e.g. [134]), are still present in later brain activity. Additionally, during 300

ms following cue onset, the brain responses elicited by exogenous and endogenous cues differ

(for review [1]). The two grating stimuli were then displayed for 50 ms. For the postcue trials we

kept the timings of cue and stimuli constant but inverted the order of their presentation (e.g.

[25,65,76,129]). A response cue, presented for 800 ms at the end of the trial after both the cue and

the stimuli had disappeared, indicated which one of the two gratings was the target (50% of the

trials on the right and the remaining 50% and on the left). The maximum delay between the off-

set of the grating stimuli and the onset of the response cue was shorter (~400 ms max in the

endogenous condition) than typically associated with a demand for working memory (>600 ms;

[135]). Immediately following each trial, a change of color of the fixation cross provided visual

feedback to the participants, i.e. green for correct or red for incorrect responses. The fixation

cross did not change color if participants had missed the response window, i.e. if they had not

pressed any key after 530 ms.

In the exogenous attention condition, a peripheral cue was presented, which was not in-

formative regarding the target location or orientation. When the cue location matched the target

location, it was considered a valid trial (50% of the trials), otherwise it was considered an invalid

trial (the remaining 50% of the trials). In the endogenous attention condition, a central cue

pointed to either the left or right quadrant. The cue was informative of the target location but not

its orientation (75% valid trials and 25% invalid trials). Participants were informed of this validity.

It is important to note that: (1) cue validity does not affect cueing effectiveness for exogenous

attention (e.g. [3,136,137], although it does so do endogenous attention (e.g. [3,136,137]); (2) the short

timing and non-informative cue in the exogenous condition ensured that no voluntary compo-

nent could be involved in the exogenous attentional effect.

Endogenous and exogenous attention conditions were performed in separate sessions to

ensure optimal manipulation of each attention system. Participants performed two practice ses-

sions outside the scanner before the first session of each attentional scanning condition. To

equate task difficulty for both attention conditions, using a staircase procedure, the tilt of the tar-

get grating was adjusted for each participant to achieve ~80% correct performance in the valid

trials in each attention condition. In each of the six experimental scanning sessions (three ses-

sions of exogenous attention and three of endogenous), participants performed 14 runs of 40

trials each, as well as a run of stimulus localizer (see MRI procedure). The tilt was then ad-

justed between runs to maintain overall performance at ~80% correct.

Eye position was monitored during all scanning sessions using an infrared video camera

system (Eyelink 1K, SR Research, Ottawa, Ontario, http://www.sr-research.com/EL_

1000.html). Trials in which the participants blinked or broke fixation (1.5° radius from central fix-

ation) at any point from fixation onset to response cue offset were identified and regressed sep-

arately in the MRI analysis, and removed from the behavioral analysis (13% ± 4% of the trials

on average across all participants).

MRI Procedure

Scanning

Imaging was conducted on a 3T Siemens Allegra head-only scanner (Erlangen, Ger-

many), using a Siemens NM-011 head coil (to transmit and receive) to acquire anatomical im-

ages, a receive-only 8-channel surface coil array (Nova Medical, Wilmington, MA) to acquire

functional images. To minimize participants' head movements, padding was used.

For each participant, three high-resolution anatomic images were acquired in one scan-

ning session, using a T1-weighted magnetization-prepared rapid gradient echo (MPRAGE) se-

quence (FOV = 256 x 256 mm; 176 sagittal slices; 1 x 1 x 1 mm voxels), and were co-regis-

tered and averaged. Using FreeSurfer (public domain software; http://surfer.nmr.mgh.har-

vard.edu), the gray matter was segmented from these averaged anatomical volumes. All subse-

quent analyses were constrained only to voxels that intersected gray matter.

T2*-weighted echo-planar imaging sequence (i.e. functional images; TR = 1750 ms; TE =

30 ms; flip angle = 90°) measured blood oxygen level-dependent (BOLD) changes in image in-

tensity [138]. In each volume, 28 slices covered the occipital and posterior parietal lobes and were

oriented 45° to the calcarine sulcus (FOV = 192 x 192 mm; resolution = 2 x 2 x 2.5 mm; no

gap). To align functional images from different sessions to the same high-resolution anatomical

volume for each participant, we acquired an additional T1-weighted anatomical images in the

same slices as the functional images (spin echo; TR = 600 ms; TE = 9.1 ms; flip angle = 90°;

resolution = 1.5 × 1.5 × 3 mm) during each scanning session, and used them in a robust image

registration algorithm.

MRI data pre-processing

Imaging data were analyzed in MATLAB, using mrTools [139] and custom software. To al-

low longitudinal magnetization to reach steady state, the first eight volumes of each run were

discarded. Spatial distortion was corrected using the B0 static magnetic field measurements

performed in each session. The functional data were then motion corrected, the linear trend

was removed, and a temporal high-pass filter was applied (cutoff: 0.01 Hz) to remove slow

drifts and low-frequency noise in the fMRI signal.

Retinotopic mapping

We followed well-established conventional traveling-wave, phase-encoded methods. Us-

ing clockwise and counter-clockwise rotating checkerboard wedges, we measured phase maps

of polar angle. Using contracting and expending checkerboard rings, we measured eccentricity

maps [95-98]. **Figure 2** (left panel) shows the visual areas of interest that were drawn by hand on

flattened surface of the brain, following published conventions [97,98,119,140].

Stimulus localizer

In each scanning session of the main experiment, participants completed one stimulus localizer run (6 runs overall, 4 min each). A run consisted of 16 cycles (17.5 s) of a block alternation protocol between stimulus on (8.75 s) and stimulus off (8.75 s). Participants only had to fixate the central cross throughout each run. The stimuli were at the same location, and of the same size and spatial frequency as those in the main experiment, except at full contrast and their phase and orientation changed randomly every 200 ms to avoid adaptation. To define the cortical representation of the gratings, we then averaged the data across the 6 runs and followed the same methods as for the retinotopic mapping. Voxels that responded positively during the blocks when the grating stimuli were presented were used to restrict each retinotopic ROI. The fMRI time series from each voxel were fit to a sinusoid. To be conservative, only voxels whose best-fit sinusoid had a phase value between 0 and pi, and a coherence between the best-fit sinusoid and the time series greater than 0.2 were included in the ROI (**Figure 2**, right panel). Analysis performed without restricting the ROI to this coherence level yielded similar results.

Event-related analysis

fMRI time series were averaged across voxels in each ROI (separately for each hemisphere) and then concatenated across runs. The data were denoised using GLMDenoise [141], and fMRI response amplitudes were computed using linear regression, with twelve regressors: 8 combinations of right and left valid and invalid pre- and post-cue, right and left cue-only, blank (no cue nor stimulus) and eye-movements (blink or broken fixation). For each ROI in each hemisphere, the resulting fMRI response amplitudes (for correct trials only) were then averaged across participants.

Acknowledgments: This work was supported by: NIH RO1-EY019693 to MC and DJH; NIH RO1-EY027401 to MC; the FYSSEN Foundation and the Philippe Foundation to LD; and the Center for Brain Imaging of New York University. We want to thank members of the Carrasco lab, especially Rachel Denison and Michael Jigo, as well as Stephanie Badde and Jonathan Winawer for constructive comments on the manuscript.

Competing interests: The authors declare no competing interests.

Author contributions: Laura Dugué: Conceptualization, Data curation, Formal analysis,
Funding acquisition, Investigation, Visualization, Methodology, Writing - original draft, Writing -
review and editing. **Elisha Merriam:** Data curation, Formal analysis, Investigation, Visualiza-
tion, Methodology, Writing - review and editing. **David Heeger:** Resources, Supervision, Fund-
ing acquisition, Visualization, Methodology, Writing - review and editing. **Marisa Carrasco:**
Conceptualization, Resources, Supervision, Funding acquisition, Visualization, Methodology,
Writing - original draft, Project administration, Writing - review and editing.

Data availability: Data relative to the main results of this study (i.e. Figures 3, 4 and 5) are
made available via OSF at [link provided upon acceptance of the manuscript].

REFERENCES

1. Carrasco, M. Visual attention: the past 25 years. *Vision Res.* **51,** 1484–1525 (2011).
2. Carrasco, M. & Barbot, A. How attention affects spatial resolution. *Cold Spring Harbor Symposia on Quantitative Biology* (2015).
3. Giordano, A. M., McElree, B. & Carrasco, M. On the automaticity and flexibility of covert attention: a speed-accuracy trade-off analysis. *Journal of Vision* **9,** 30.1–10 (2009).
4. Pestilli, F., Ling, S. & Carrasco, M. A population-coding model of attention's influence on contrast response: Estimating neural effects from psychophysical data. *Vision Res.* **49,** 1144–1153 (2009).
5. Ling, S. & Carrasco, M. Sustained and transient covert attention enhance the signal via different contrast response functions. *Vision Res.* **46,** 1210–1220 (2006).
6. Dosher, B. A. & Lu, Z. L. Mechanisms of perceptual attention in precuing of location. *Vision Res.* **40,** 1269–1292 (2000).
7. Barbot, A., Landy, M. S. & Carrasco, M. Differential effects of exogenous and endogenous attention on second-order texture contrast sensitivity. *Journal of Vision* **12,** 6–6 (2012).
8. Jigo, M. & Carrasco, M. Differential impact of exogenous and endogenous attention on the contrast sensitivity function across eccentricity. *Journal of Vision* **20,** 11–11 (2020).
9. Yeshurun, Y. & Carrasco, M. Attention improves or impairs visual performance by enhancing spatial resolution. *Nature* **396,** 72–75 (1998).
10. Yeshurun, Y., Montagna, B. & Carrasco, M. On the flexibility of sustained attention and its effects on a texture segmentation task. *Vision Res.* **48,** 80–95 (2008).
11. Barbot, A. & Carrasco, M. Attention Modifies Spatial Resolution According to Task Demands. *Psychological Science* **28,** 285–296 (2017).
12. Jigo, M. & Carrasco, M. Attention alters spatial resolution by modulating second-order processing. *Journal of Vision* **18,** 2 (2018).
13. Yeshurun, Y. & Carrasco, M. The effects of transient attention on spatial resolution and the size of the attentional cue. *Perception & Psychophysics* **70,** 104–113 (2008).
14. Carrasco, M., Loula, F. & Ho, Y.-X. How attention enhances spatial resolution: evidence from selective adaptation to spatial frequency. *Perception & Psychophysics* **68,** 1004–1012 (2006).
15. Anton-Erxleben, K. & Carrasco, M. Attentional enhancement of spatial resolution: linking behavioural and neurophysiological evidence. *Nat. Rev. Neurosci.* **14,** 188–200 (2013).
16. Carrasco, M. & Yeshurun, Y. Covert attention effects on spatial resolution. *Prog. Brain Res.* **176,** 65–86 (2009).
17. Reynolds, J. H. & Heeger, D. J. The Normalization Model of Attention. *Neuron* **61,** 168–185 (2009).
18. Martínez-Trujillo, J. C. & Treue, S. Attentional Modulation Strength in Cortical Area MT Depends on Stimulus Contrast. *Neuron* **35,** 365–370 (2002).
19. McAdams, C. J. & Maunsell, J. H. R. Effects of Attention on Orientation-Tuning Functions of Single Neurons in Macaque Cortical Area V4. *J. Neurosci.* **19,** 431–441 (1999).
20. Mitchell, J. F., Sundberg, K. A. & Reynolds, J. H. Spatial Attention Decorrelates Intrinsic Activity Fluctuations in Macaque Area V4. *Neuron* **63,** 879–888 (2009).
21. Reynolds, J. H., Pasternak, T. & Desimone, R. Attention increases sensitivity of V4 neurons. *Neuron* **26,** 703–714 (2000).
22. Williford, T. & Maunsell, J. H. R. Effects of Spatial Attention on Contrast Response Functions in Macaque Area V4. *J. Neurophysiol.* **96,** 40–54 (2006).
23. Luo, T. Z. & Maunsell, J. H. R. Neuronal Modulations in Visual Cortex Are Associated with Only One of Multiple Components of Attention. *Neuron* **86,** 1182–1188 (2015).
24. Ruff, D. A. & Cohen, M. R. Attention can either increase or decrease spike count correlations in visual cortex. *Nat. Neurosci.* **17,** 1591–1597 (2014).
25. Pestilli, F., Carrasco, M., Heeger, D. J. & Gardner, J. L. Attentional enhancement via selection and pooling of early sensory responses in human visual cortex. *Neuron* **72,** 832–846 (2011).

26. Murray, S. O. The effects of spatial attention in early human visual cortex are stimulus
independent. *Journal of Vision* **8,** 2.1–11 (2008).
27. Herrmann, K., Montaser-Kouhsari, L., Carrasco, M. & Heeger, D. J. When size matters:
attention affects performance by contrast or response gain. *Nat. Neurosci.* **13,** 1554–
1559 (2010).
28. Buracas, G. T. & Boynton, G. M. The effect of spatial attention on contrast response
functions in human visual cortex. *Journal of Neuroscience* **27,** 93–97 (2007).
29. O'Connor, D. H., Fukui, M. M. & Pinsk, M. A. Attention modulates responses in the hu-
man lateral geniculate nucleus - ProQuest. *Nature* **5,** 1203–1209 (2002).
30. Kastner, S., Pinsk, M. A., De Weerd, P., Desimone, R. & Ungerleider, L. G. Increased
Activity in Human Visual Cortex during Directed Attention in the Absence of Visual
Stimulation. *Neuron* **22,** 751–761 (1999).
31. Somers, D. C., Dale, A. M., Seiffert, A. E. & Tootell, R. B. Functional MRI reveals spa-
tially specific attentional modulation in human primary visual cortex. *PNAS* **96,** 1663–
1668 (1999).
32. Beck, D. M. & Kastner, S. Neural systems for spatial **attention** in the human brain: Evi-
dence from neuroimaging in the framework of biased competition. *The Oxford Hand-
book of Attention* 253–288 (2014).
33. Lu, Z.-L., Li, X., Tjan, B. S., Dosher, B. A. & Chu, W. Attention Extracts Signal in Exter-
nal Noise: A BOLD fMRI Study. *J Cogn Neurosci* **23,** 1148–1159 (2011).
34. Li, X., Lu, Z.-L., Tjan, B. S., Dosher, B. A. & Chu, W. Blood oxygenation level-depend-
ent contrast response functions identify mechanisms of covert attention in early visual
areas. *Proc. Natl. Acad. Sci. U.S.A.* **105,** 6202–6207 (2008).
35. Busse, L., Katzner, S. & Treue, S. Temporal dynamics of neuronal modulation during
exogenous and endogenous shifts of visual attention in macaque area MT. *Proc. Natl.
Acad. Sci. U.S.A.* **105,** 16380–16385 (2008).
36. Wang, F., Chen, M., Yan, Y., Zhaoping, L. & Li, W. Modulation of Neuronal Responses
by Exogenous Attention in Macaque Primary Visual Cortex. *Journal of Neuroscience*
35, 13419–13429 (2015).
37. Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in
the brain. *Nat. Rev. Neurosci.* **3,** 201–215 (2002).
38. Chica, A. B., Bartolomeo, P. & Lupiáñez, J. Two cognitive and neural systems for en-
dogenous and exogenous spatial attention. *Behavioural Brain Research* **237,** 107–123
(2013).
39. Bowling, J. T., Friston, K. J. & Hopfinger, J. B. Top-down versus bottom-up attention
differentially modulate frontal–parietal connectivity. *Hum Brain Mapp* **9,** 1116 (2019).
40. Hahn, B., Ross, T. J. & Stein, E. A. Neuroanatomical dissociation between bottom–up
and top–down processes of visuospatial selective attention. *Neuroimage* **32,** 842–853
(2006).
41. Dugué, L., Merriam, E. P., Heeger, D. J. & Carrasco, M. Specific Visual Subregions of
TPJ Mediate Reorienting of Spatial Attention. *Cereb. Cortex* 1–16 (2017).
doi:10.1093/cercor/bhx140
42. Geng, J. J. & Vossel, S. Re-evaluating the role of TPJ in attentional control: contextual
updating? *Neurosci Biobehav Rev* **37,** 2608–2620 (2013).
43. Doricchi, F., Macci, E., Silvetti, M. & Macaluso, E. Neural correlates of the spatial and
expectancy components of endogenous and stimulus-driven orienting of attention in
the Posner task. *Cereb. Cortex* **20,** 1574–1585 (2010).
44. Silvetti, M. *et al.* The Response of the Left Ventral Attentional System to Invalid Targets
and its Implication for the Spatial Neglect Syndrome: a Multivariate fMRI Investigation.
Cereb. Cortex 1–12 (2015). doi:10.1093/cercor/bhv208
45. Beck, D. M. & Kastner, S. Top-down and bottom-up mechanisms in biasing competition
in the human brain. *Vision Res.* **49,** 1154–1165 (2009).
46. Nakayama, K. & Mackeben, M. Sustained and transient components of focal visual at-
tention. *Vision Res.* **29,** 1631–1647 (1989).
47. Posner, M. I., Snyder, C. R. & Davidson, B. J. Attention and the detection of signals. *J
Exp Psychol* **109,** 160–174 (1980).

48.   Buschman, T. J. & Miller, E. K. Serial, Covert Shifts of Attention during Visual Search
696   Are Reflected by the Frontal Eye Fields and Correlated with Population Oscillations.
697   *Neuron* **63,** 386–396 (2009).
698   49.   Corbetta, M., Patel, G. & Shulman, G. L. The reorienting system of the human brain:
699   from environment to theory of mind. *Neuron* **58,** 306–324 (2008).
700   50.   Peelen, M. V., Heslenfeld, D. J. & Theeuwes, J. Endogenous and exogenous attention
701   shifts are mediated by the same large-scale neural network. *Neuroimage* **22,** 822–830
702   (2004).
703   51.   Kincade, J. M., Abrams, R. A., Astafiev, S. V., Shulman, G. L. & Corbetta, M. An event-
704   related functional magnetic resonance imaging study of voluntary and stimulus-driven
705   orienting of attention. *Journal of Neuroscience* **25,** 4593–4604 (2005).
706   52.   Mayer, A. R., Dorflinger, J. M., Rao, S. M. & Seidenberg, M. Neural networks underly-
707   ing endogenous and exogenous visual–spatial orienting. *Neuroimage* **23,** 534–541
708   (2004).
709   53.   Carrasco, M. & McElree, B. Covert attention accelerates the rate of visual information
710   processing. *PNAS* **98,** 5363–5367 (2001).
711   54.   Reed, A. V. Speed-Accuracy Trade-Off in Recognition Memory. *Science* **181,** 574–576
712   (1973).
713   55.   Wickelgren, W. A. Speed-accuracy tradeoff and information processing dynamics. *Acta*
714   *Psychologica* **41,** 67–85 (1977).
715   56.   Correa, Á., Triviño, M., Pérez-Dueñas, C., Acosta, A. & Lupiáñez, J. Temporal prepara-
716   tion, response inhibition and impulsivity. *Brain and Cognition* **73,** 222–228 (2010).
717   57.   Yashar, A. & Lamy, D. Refining the dual-stage account of intertrial feature priming:
718   does motor response or response feature matter? *Atten Percept Psychophys* **73,**
719   2160–2167 (2011).
720   58.   Herath, P., Klingberg, T., Young, J., Amunts, K. & Roland, P. E. Neural Correlates of
721   Dual Task Interference Can be Dissociated from Those of Divided Attention: an fMRI
722   Study | Cerebral Cortex | Oxford Academic. *Cereb. Cortex* **11,** 796–805 (2001).
723   59.   Culham, J. C., Cavanagh, P. & Kanwisher, N. G. Attention Response Functions: Char-
724   acterizing Brain Areas Using fMRI Activation during Parametric Variations of Attentional
725   Load. *Neuron* **32,** 737–745 (2001).
726   60.   Beauchamp, M. S., Petit, L., Ellmore, T. M., Ingeholm, J. & Haxby, J. V. A parametric
727   fMRI study of overt and covert shifts of visuospatial attention. *Neuroimage* **14,** 310–321
728   (2001).
729   61.   Carrasco, M., Evert, D. L., Chang, I. & Katz, S. M. The eccentricity effect: target eccen-
730   tricity affects performance on conjunction searches. *Perception & Psychophysics* **57,**
731   1241–1261 (1995).
732   62.   Rosen, A. C. *et al.* Neural basis of endogenous and exogenous spatial orienting. A
733   functional MRI study. *Journal of Cognitive Neuroscience* **11,** 135–152 (1999).
734   63.   Natale, E., Marzi, C. A., Girelli, M., Pavone, E. F. & Pollmann, S. ERP and fMRI corre-
735   lates of endogenous and exogenous focusing of visual-spatial attention. *Eur. J. Neuro-*
736   *sci.* **23,** 2511–2521 (2006).
737   64.   Esterman, M. *et al.* Voluntary and involuntary attention affect face discrimination differ-
738   ently. *Neuropsychologia* **46,** 1032–1040 (2008).
739   65.   Kinchla, R. A., Chen, Z. & Evert, D. Precue effects in visual search: data or resource
740   limited? *Perception & Psychophysics* **57,** 441–450 (1995).
741   66.   Ruff, C. C., Kristjánsson, Á. & Driver, J. Readout from iconic memory and selective
742   spatial attention involve similar neural processes. *Psychological Science* **18,** 901–909
743   (2007).
744   67.   Hulme, O. J., Friston, K. F. & Zeki, S. Neural correlates of stimulus reportability. *Jour-*
745   *nal of Cognitive Neuroscience* **21,** 1602–1610 (2009).
746   68.   Nobre, A. C. *et al.* Orienting attention to locations in perceptual versus mental repre-
747   sentations. *Journal of Cognitive Neuroscience* **16,** 363–373 (2004).
748   69.   Sergent, C., Ruff, C. C., Barbot, A., Driver, J. & Rees, G. Top-down modulation of hu-
749   man early visual cortex after stimulus offset supports successful postcued report. *Jour-*
750   *nal of Cognitive Neuroscience* **23,** 1921–1934 (2011).

70. Thibault, L., van den Berg, R., Cavanagh, P. & Sergent, C. Retrospective Attention
Gates Discrete Conscious Access to Past Sensory Stimuli. *PLoS ONE* **11,** e0148504
(2016).
71. Sergent, C. *et al.* Cueing Attention after the Stimulus Is Gone Can Retrospectively Trig-
ger Conscious Perception. *Current Biology* **23,** 150–155 (2013).
72. Gobell, J. & Carrasco, M. Attention alters the appearance of spatial frequency and gap
size. *Psychological Science* **16,** 644–651 (2005).
73. Anton-Erxleben, K., Henrich, C. & Treue, S. Attention changes perceived size of mov-
ing visual patterns. *Journal of Vision* **7,** 5.1–9 (2007).
74. Fuller, S., Park, Y. & Carrasco, M. Cue contrast modulates the effects of exogenous
attention on appearance. *Vision Res.* **49,** 1825–1837 (2009).
75. Carrasco, M. & Yeshurun, Y. The contribution of covert attention to the set-size and ec-
centricity effects in visual search. *Journal of Experimental Psychology: Human Percep-
tion and Performance* **24,** 673–692 (1998).
76. Liu, T., Pestilli, F. & Carrasco, M. Transient attention enhances perceptual performance
and FMRI response in human visual cortex. *Neuron* **45,** 469–477 (2005).
77. Kim, Y. H. *et al.* The large-scale neural network for spatial attention displays multifunc-
tional overlap but differential asymmetry. *Neuroimage* **9,** 269–277 (1999).
78. Meyer, K. N., Du, F., Parks, E. & Hopfinger, J. B. Exogenous vs. endogenous attention:
Shifting the balance of fronto-parietal activity. *Neuropsychologia* **111,** 307–316 (2018).
79. Dugué, L., Roberts, M. & Carrasco, M. Attention Reorients Periodically. *Current Biology*
26, 1595–1601 (2016).
80. Senoussi, M., Moreland, J. C., Busch, N. A. & Dugué, L. Attention explores space peri-
odically at the theta frequency. *Journal of Vision* **19,** 22–22 (2019).
81. Maunsell, J. H. R. & Cook, E. P. The role of attention in visual processing. *Philosophi-
cal Transactions of the Royal Society of London B: Biological Sciences* **357,** 1063–
1072 (2002).
82. Kastner, S. & Pinsk, M. A. Visual attention as a multilevel selection process. *Cognitive,
Affective, & Behavioral Neuroscience* **4,** 483–500 (2004).
83. Klein, B. P., Harvey, B. M. & Dumoulin, S. O. Attraction of Position Preference by Spa-
tial Attention throughout Human Visual Cortex. *Neuron* **84,** 227–237 (2014).
84. Kay, K. N., Weiner, K. S. & Grill-Spector, K. Attention Reduces Spatial Uncertainty in
Human Ventral Temporal Cortex. *Current Biology* **25,** 595–600 (2015).
85. Nachmias, J. Effect of Exposure Duration on Visual Contrast Sensitivity with Square-
Wave Gratings. *J. Opt. Soc. Am., JOSA* **57,** 421–427 (1967).
86. Carrasco, M., Penpeci-Talgar, C. & Eckstein, M. Spatial covert attention increases con-
trast sensitivity across the CSF: support for signal enhancement. *Vision Res.* **40,** 1203–
1215 (2000).
87. Pestilli, F. & Carrasco, M. Attention enhances contrast sensitivity at cued and impairs it
at uncued locations. *Vision Res.* **45,** 1867–1875 (2005).
88. Fernández, A. & Carrasco, M. Extinguishing exogenous attention via transcranial mag-
netic stimulation. *Curr. Biol.* **30,** 1–7 (2020).
89. Luck, S. J. *et al.* Effects of spatial cuing on luminance detectability: psychophysical and
electrophysiological evidence for early selection. *Journal of Experimental Psychology:
Human Perception and Performance* **20,** 887–904 (1994).
90. Lu, Z. L. & Dosher, B. A. External noise distinguishes attention mechanisms. *Vision
Res.* **38,** 1183–1198 (1998).
91. Montagna, B., Pestilli, F. & Carrasco, M. Attention trades off spatial acuity. *Vision Res.*
49, 735–745 (2009).
92. Yeshurun, Y. & Rashal, E. Precueing attention to the target location diminishes crowd-
ing and reduces the critical distance. *Journal of Vision* **10,** 16–16 (2010).
93. Dugué, L., Xue, A. M. & Carrasco, M. Distinct perceptual rhythms for feature and con-
junction searches. *Journal of Vision* **17,** 22–22 (2017).
94. Ling, S. & Carrasco, M. Sustained and transient covert attention enhance the signal via
different contrast response functions. *Vision Res.* **46,** 1210–1220 (2006).
95. Engel, S. A. *et al.* fMRI of human visual cortex. *Nature* **369,** 525 (1994).

96.   Sereno, M. I. *et al.* Borders of multiple visual areas in humans revealed by functional
magnetic resonance imaging. *Science* **268,** 889–893 (1995).
97.   Engel, S. A., Glover, G. H. & Wandell, B. A. Retinotopic organization in human visual
cortex and the spatial precision of functional MRI. *Cereb. Cortex* **7,** 181–192 (1997).
98.   Larsson, J. & Heeger, D. J. Two retinotopic visual areas in human lateral occipital cor-
tex. *Journal of Neuroscience* **26,** 13128–13142 (2006).
99.   Pinheiro, J. C. & Bates, D. M. Unconstrained parametrizations for variance-covariance
matrices. *Stat Comput* **6,** 289–296 (1996).
100. Galletti, C. *et al.* Covert Shift of Attention Modulates the Ongoing Neural Activity in a
Reaching Area of the Macaque Dorsomedial Visual Stream. *PLoS ONE* **5,** e15078
(2010).
101. Caspari, N., Janssens, T., Mantini, D., Vandenberghe, R. & Vanduffel, W. Covert shifts
of spatial attention in the macaque monkey. *Journal of Neuroscience* **35,** 7695–7714
(2015).
102. Caspari, N., Arsenault, J. T., Vandenberghe, R. & Vanduffel, W. Functional Similarity of
Medial Superior Parietal Areas for Shift-Selective Attention Signals in Humans and
Monkeys. *Cereb. Cortex* **28,** 2085–2099 (2018).
103. Pitzalis, S. *et al.* Wide-Field Retinotopy Defines Human Cortical Visual Area V6. *J.*
Neurosci. **26,** 7962–7973 (2006).
104. Harvey, B. M. & Dumoulin, S. O. The relationship between cortical magnification factor
and population receptive field size in human visual cortex: constancies in cortical archi-
tecture. *Journal of Neuroscience* **31,** 13604–13612 (2011).
105. Harvey, B. M. & Dumoulin, S. O. Visual motion transforms visual space representations
similarly throughout the human visual hierarchy. *Neuroimage* **127,** 173–185 (2016).
106. Michalareas, G. *et al.* Alpha-Beta and Gamma Rhythms Subserve Feedback and
Feedforward Influences among Human Visual Cortical Areas. *Neuron* **89,** 384–397
(2016).
107. Yildirim, F., Carvalho, J. & Cornelissen, F. W. A second-order orientation-contrast stim-
ulus for population-receptive-field-based retinotopic mapping. *Neuroimage* **164,** 183–
193 (2018).
108. Fritsche, M., Lawrence, S. J. D. & de Lange, F. P. Temporal tuning of repetition sup-
pression across the visual cortex. *J. Neurophysiol.* **123,** 224–233 (2020).
109. Boynton, G. M., Demb, J. B., Glover, G. H. & Heeger, D. J. Neuronal basis of contrast
discrimination. *Vision Res.* **39,** 257–269 (1999).
110. Brefczynski, J. A. & DeYoe, E. A. A physiological correlate of the 'spotlight' of visual at-
tention. *Nat. Neurosci.* **2,** 370–374 (1999).
111. Slotnick, S. D. The experimental parameters that affect attentional modulation of the
ERP C1 component. *Cogn Neurosci* **9,** 53–62 (2017).
112. Müller, N. G. & Kleinschmidt, A. Temporal Dynamics of the Attentional Spotlight: Neu-
ronal Correlates of Attentional Capture and Inhibition of Return in Early Visual Cortex. *J*
Cogn Neurosci **19,** 587–593 (2007).
113. Müller, N. G. & Ebeling, D. Attention-modulated activity in visual cortex—More than a
simple 'spotlight'. *Neuroimage* **40,** 818–827 (2008).
114. Heinen, K. *et al.* Concurrent TMS-fMRI reveals dynamic interhemispheric influences of
the right parietal cortex during exogenously cued visuospatial attention. *European Jour-*
nal of Neuroscience **33,** 991–1000 (2011).
115. Mulckhuyse, M., Belopolsky, A. V., Heslenfeld, D., Talsma, D. & Theeuwes, J. Distribu-
tion of attention modulates salience signals in early visual cortex. *PLoS ONE* **6,** e20379
(2011).
116. Kay, K. N., Winawer, J., Rokem, A., Mezer, A. & Wandell, B. A. A two-stage cascade
model of BOLD responses in human visual cortex. *PLoS Comput. Biol.* **9,** e1003079
(2013).
117. Chica, A. B. & Lupiáñez, J. Effects of endogenous and exogenous attention on visual
processing: an Inhibition of Return study. *Brain Res.* **1278,** 75–85 (2009).
118. Liu, T., Heeger, D. J. & Carrasco, M. Neural correlates of the visual vertical meridian
asymmetry. *Journal of Vision* **6,** 1294–1306 (2006).

119. Liu, T., Larsson, J. & Carrasco, M. Feature-based attention modulates orientation-selective responses in human visual cortex. *Neuron* **55,** 313–323 (2007).

120. Freeman, J., Heeger, D. J. & Merriam, E. P. Coarse-scale biases for spirals and orientation in human visual cortex. *Journal of Neuroscience* **33,** 19695–19703 (2013).

121. Wang, H. X., Merriam, E. P., Freeman, J. & Heeger, D. J. Motion direction biases and decoding in human visual cortex. *Journal of Neuroscience* **34,** 12601–12615 (2014).

122. Brouwer, G. J., Arnedo, V., Offen, S., Heeger, D. J. & Grant, A. C. Normalization in human somatosensory cortex. *J. Neurophysiol.* **114,** 2588–2599 (2015).

123. McMains, S. A. & Somers, D. C. Multiple spotlights of attentional selection in human visual cortex. *Neuron* **42,** 677–686 (2004).

124. Bloem, I. M. & Ling, S. Normalization governs attentional modulation within human visual cortex. *Nature Communications* **10,** 5660–10 (2019).

125. Poltoratski, S., Ling, S., McCormack, D. & Tong, F. Characterizing the effects of feature salience and top-down attention in the early visual system. *J. Neurophysiol.* **118,** 564–573 (2017).

126. Moerel, D., Ling, S. & Jehee, J. F. M. Perceptual learning increases orientation sampling efficiency. *Journal of Vision* **16,** 36–36 (2016).

127. Mackey, W. E., Winawer, J. & Curtis, C. E. Visual field map clusters in human frontoparietal cortex. *Elife* **6,** 2704 (2017).

128. Gardner, J. L., Merriam, E. P., Schluppeck, D. & Larsson, J. MGL: Visual psychophysics stimuli and experimental design package. (2018). doi:10.5281/zenodo.1299497

129. Yaffa Yeshurun, M. C. Attention improves or impairs visual performance by enhancing spatial resolution. *Nature* **396,** 72–75 (1998).

130. Müller, M. M. in *Cognitive Electrophysiology of Attention* 123–135 (Elsevier, 2014). doi:10.1016/B978-0-12-398451-7.00010-5

131. Mackeben, M. & Nakayama, K. Express attentional shifts. *Vision Res.* **33,** 85–90 (1993).

132. Liu, T., Stevens, S. T. & Carrasco, M. Comparing the time course and efficacy of spatial and feature-based attention. *Vision Res.* **47,** 108–113 (2007).

133. Ling, S. & Carrasco, M. When sustained attention impairs perception. *Nat. Neurosci.* **9,** 1243–1245 (2006).

134. Seiss, E., Driver, J. & Eimer, M. Effects of attentional filtering demands on preparatory ERPs elicited in a spatial cueing task. *Clin Neurophysiol* **120,** 1087–1095 (2009).

135. Phillips, W. A. On the distinction between sensory storage and short-term visual memory. *Perception & Psychophysics* **16,** 283–290 (1974).

136. Sperling, G. & Melchner, M. J. The attention operating characteristic: examples from visual search. *Science* **202,** 315–318 (1978).

137. Kinchla, R. A. The measurement of attention. *Attention and performance VIII* 213–238 (1980).

138. Ogawa, S., Lee, T. M., Kay, A. R. & Tank, D. W. Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proc. Natl. Acad. Sci. U.S.A.* **87,** 9868–9872 (1990).

139. Gardner, J. L., Merriam, E. P., Schluppeck, D., Besle, J. & Heeger, D. J. mrTools: Analysis and visualization package for functional magnetic resonance imaging data. (2018). doi:10.5281/zenodo.1299483

140. Wandell, B. A., Dumoulin, S. O. & Brewer, A. A. Visual field maps in human cortex. *Neuron* **56,** 366–383 (2007).

141. Kay, K., Rokem, A., Winawer, J., Dougherty, R. & Wandell, B. GLMdenoise: a fast, automated technique for denoising task-based fMRI data. *Front. Neurosci.* **7,** 1–15 (2013).

Figure

Figure 1. Experimental protocol. A. Participants performed a 2-AFC orientation-discrimination

task. Pre-cues and post-cues were presented before and after the stimuli, respectively. Exoge-

nous cues appeared in the periphery, above one of the two stimulus locations. Endogenous

cues appeared at the center of the screen (the display is not at scale for visibility purposes) and

indicated one of the two stimulus locations The ISI between the cue and the gratings was

shorter for the exogenous (50 ms) than the endogenous (250 ms) conditions. A response cue

indicated the target location and instructed participants to indicate whether the target grating

was tilted clockwise or counterclockwise of vertical by pressing one of two keys. To provide

feedback, the fixation-cross turned green or red for a correct or an incorrect answer, respec-

tively. **B.** Behavioral performance averaged across participants (black dots) and for each of

them (colored dots; n=5) for endogenous (left) and exogenous (right) attention. (Top) Perfor-

mance accuracy (d'; top) and median reaction time (bottom) as a function of cueing condition.

V, valid cue (same location of pre-cue/post-cue as response cue). IN, invalid cue (different loca-

tion of pre-cue/post-cue than response cue). Pre, pre-cue presented before the stimuli. Post,

post-cue presented after the stimuli. Valid cues induced more accurate and faster responses

(there was no speed-accuracy trade-off). Error bars, ± 1 SEM across participants.

Figure 2. Retinotopic mapping and stimulus localizer of a representative participant. A flattened representation of the right hemisphere occipital pole. **Left,** map of polar angle. The color of each voxel in the map represents the response phase of the time series elicited by a rotating wedge stimulus. **Right,** stimulus localizer. The black outlines correspond to the retino-topic areas of interest (defined for each participant based on their polar angle maps). The color of each voxel indicates the phase of the response to the grating stimuli presented in the lower left visual field.

Figure 3. Specificity of pre and post-cueing for endogenous and exogenous attention.

AMI, Attentional Modulation Index separately for pre- and post-cue conditions for each ROI:
AMI = Vc / average (Vc + Ic + Vi + Ii). Vc, valid trials in contralateral ROI. Ic, invalid trials in con-
tralateral ROI. Vi, valid pre- and post-cue trials in ipsilateral ROI. Ii, invalid pre- and post-cue
trials in ipsilateral ROI. The larger this ratio, the higher the modulation in the valid trials. *, Sta-
tistically significant regression analysis (p < 0.05).

Figure 4. Single ROI responses for pre and post-cueing and for both endogenous and exogenous attention. The AMI is plotted separately for pre and post-cue conditions for each ROI (same data as in Figure 3). *, Statistically significant difference between valid and invalid, separately for pre and post-cueing ($p < 0.05$). Error bars on plots are ± 1 SEM.

Figure 5. Specificity of each visual ROI for endogenous and exogenous attention. fMRI

response amplitude was measured for each attentional condition. V: valid cue condition (target

location matches the location indicated by pre-cue/post-cue). IN: invalid cue condition (target

location at the opposite location relative to the pre-cue/post-cue). Pre: pre-cue presented before

the grating stimuli. Post: post-cue presented after gratings. All four conditions in the contrala-

teral ROI. Distractor (Dist): pre-cue and post-cue valid and invalid conditions averaged in the

ipsilateral ROI. Each colored point corresponds to the data of one participant. The black points

represent the average across all 5 participants. *, Statistically significant difference between

valid and invalid, separately for pre and post-cueing ($p < 0.05$). #, trend ($p < 0.1$). Detailed sta-

tistics are presented in **Supplementary Table 1**. Error bars (vertical lines) are plotted on the

difference between valid and invalid and represent ± 1 SEM across participant.