1 # Differential impact of endogenous and exogenous attention

2 ## on activity in human visual cortex

4 Laura Dugué[1,2,3,4], Elisha P. Merriam[1,2,5], David J. Heeger[1,2] & Marisa Carrasco[1,2]

[1] Department of Psychology, New York University
[2] Center for Neural Science, New York University
[3] Université de Paris, CNRS, Integrative Neuroscience and Cognition Center, F-75006 Paris,
France
[4] Institut Universitaire de France, Paris, France
[5] Laboratory of Brain and Cognition, NIMH/NIH, Bethesda, MD
16

18 **Running title:** endogenous and exogenous attention

21 **Corresponding author:**
22 Laura Dugué
23 Current address: 45 rue des Saints-Pères 75006 Paris, FRANCE
24 laura.dugue@u-paris.fr

ABSTRACT (148 words)

How do endogenous (voluntary) and exogenous (involuntary) attention modulate activity in visual cortex? Using ROI-based fMRI analysis, we measured fMRI activity for valid and invalid trials (target at cued/un-cued location, respectively), pre- or post-cueing endogenous or exogenous attention, while participants performed the same discrimination task. We found stronger modulation in contralateral than ipsilateral visual regions, and higher activity in valid- than invalid-trials. For endogenous attention, modulation of stimulus-evoked activity due to a pre-cue increased along the visual hierarchy, but was constant due to a post-cue. For exogenous attention, modulation of stimulus-evoked activity due to a pre-cue was constant along the visual hierarchy, but not modulated due to a post-cue. These findings reveal that endogenous and exogenous attention distinctly modulate activity in visuo-occipital areas during orienting and reorienting; endogenous attention facilitates both the encoding and the readout of visual information whereas exogenous attention only facilitates the encoding of information.

Keywords: exogenous attention, endogenous attention, fMRI activity, occipital visual areas, top-down processing, bottom-up processing.

INTRODUCTION

Spatial, covert visual attention is the selective processing of visual information in space, without change in gaze. Attention can be allocated voluntarily –endogenous attention– or involuntarily –exogenous attention. Endogenous and exogenous attention have different temporal dynamics; endogenous takes about 300 ms to be deployed and can be sustained at will whereas exogenous attention takes only about 100 ms to be deployed and it is transient (for review see [1,2]). These two types of attention often have similar perceptual consequences (reviews by [1,2]), but notable exceptions indicate that whereas endogenous attention acts in a flexible way, exogenous attention acts in an inflexible or automatic way. For instance: (a) The benefits and costs in perception (discriminability and speed of information accrual) scale with cue validity for endogenous but not for exogenous attention (e.g. [3]); (b) The effects of covert attention on contrast sensitivity often differ for endogenous and exogenous attention (e.g. [4-7]); and (c) For a texture segmentation task in which heightened spatial resolution improves or impairs performance as a function of target eccentricity, endogenous attention improves performance across eccentricity, whereas exogenous attention improves performance at peripheral locations where resolution is poor, but hampers performance where resolution is already high for the task at hand (e.g. [8-12]).

Basic visual processes, such as contrast sensitivity and spatial resolution, are mediated by activity in early visual cortex, and are altered by covert attention (reviews by [1,2,13,14]). Specifically, single-unit studies in monkeys have demonstrated effects of endogenous attention in occipital areas (e.g. [15-22]). Additionally, fMRI studies have shown that endogenous attention causes a baseline shift in early visual areas (e.g. [23-29]; review by [30]) and increases the dynamic range of fMRI responses [31,32]. Comparatively, little is known about the effect of exogenous attention on visual areas both from single-unit studies [33,34] and fMRI studies (for reviews see [1,13]).

Since Corbetta and Schulman's seminal review (2002) [35] on the neural bases of endogenous and exogenous attention, there has been emphasis on characterizing networks of brain regions within the frontal and parietal lobes (for reviews see [30,36,37]). Yet, there remains considerable debate regarding the dissociation between dorsal regions, for endogenous attention, and ventral regions, for exogenous attention (e.g. [37,38]), including the role of the temporo-parietal

junction (TPJ; [39-42]). Classically, researchers have described endogenous attention as a top-down

process, and exogenous attention as a bottom-up process [1,35,37,43-45]. This characterization origi-

nated in psychophysics experiments, and was then studied using fMRI, in which the two types of

attention have been often investigated separately (for reviews [1,30,36,43]).

Studies of endogenous and exogenous attention focusing on parietal and frontal areas

have shown that the two types of attention differentially modulate fronto-parietal connectivity [37,46].

For example, there are critical differences in the temporal order of neural responses in frontal

and parietal cortex in monkeys between these attention conditions, i.e. frontal activity precedes

parietal activity during endogenous attentional orienting, whereas parietal activity precedes

frontal activity during exogenous orienting. Critically, it is often assumed that the effects of en-

dogenous and exogenous attention are the same in striate and extra-striate areas [30,35,36,47,48].

A number of important considerations limit the conclusions that may be drawn from the

few studies that have directly compared independent effects of visual exogenous and endoge-

nous spatial attention [37,48-50]:

(1) The effect of attention on behavioral performance was assessed in some studies with a

detection task using RT as their only dependent variable [48,50], in which performance may

differ due to speed of processing, discriminability or criterion [51-53] and/or motor behavior

[54,55].

(2) In the studies in which performance was assessed in a discrimination task using RT [49],

small RT differences were reported, which could have resulted from speed of processing,

discriminability, or criterion factors [51-53].

(3) In the studies in which accuracy was not assessed [37,48,50], it is not possible to know whether

task difficulty was the same for both types of attention, and task difficulty can interact with

the strength of fMRI activity [56,57].

(4) For the studies in which eye position was not monitored while participants performed the

task in the scanner [37,48-50], the results could be due to covert attention, overt attention or

both [1,58,59].

(5) Given that exogenous attention is a fast, transient process (for review see [1,2]), it was

inappropriately manipulated in the studies in which long stimulus onset asynchronies

(SOA) were used [48,49,60,61] [62] making the comparison between the two attention conditions

problematic.

(6) Except for two studies, one interested in attentional modulation in the Fusiform Face Area

(FFA) [61] and the other in the TPJ [39], statistical parametric mapping was applied to group

averaged data to identify regions of the brain that were active during task performance.

Some found significant activity in the occipital pole [48-50], but attentional modulation of fMRI

activity was not systematically assessed across different visual areas.

Given all these methodological limitations, it is unknown how these two types of attention

affect neural activity in each individual visuo-occipital area, and how modulation of activity in

visual cortex is linked to changes in perceptual performance. (See Dugué and colleagues [39], who

published a table summarizing these and other methodological problems for studies regarding

covert attention and TPJ activation).

Typically, covert attention is manipulated by presenting a pre-cue, prior to the target –and

the aforementioned studies also did so (see **Table 1**). However, endogenous post-cues, pre-

sented after target offset, can also improve performance by affecting the information readout [63-

66] and modulate fMRI activity in early visual areas [23,65-67]. Exogenous post-cues also affect per-

formance in some tasks [68,69], but not in others [70-73], and the only fMRI study evaluating post-cues

in exogenous attention showed no such modulation [74]. Critically, no single study has compared

visual cortex activity with post-cues in endogenous and exogenous attention.

Study	Cueing manipulation		Task	Dependent Variable	Analysis	Cortical areas	Eye tracking	Correlation MRI-behavior
	Endogenous	**Exogenous**						
Studies including Occipital areas								
Detection tasks								
50	central arrow 70% valid	luminance change of peripheral placeholder 50% valid	X	RT	Group-averaged	**Occipital:** *Cuneus, MOG, SOG* + Frontal Temporo-Parietal	No	No
	SOA=100 or 800 ms							
48	central arrow 75% valid	brightening of peripheral placeholder 50% valid	square	RT	Group-averaged	**Occipital:** *Cuneus* + Frontal Temporo-Parietal	No	No
	SOA=550 ms							
62	single peripheral rectangular frame 100% valid	five peripheral rectangular frames 20% valid	a black and white checkerboard	RT *for 4 out of 8 participants*	Group-averaged	**Occipital:** *Fusiform, TOS, LG, SC, LOG* + Frontal Temporo-Parietal	No	No
	SOA=8, 8.15, 8.3, 8.45 or 8.6 s							
Discrimination tasks								
49	brightening of half of central fixation diamond 75% valid	color singleton in an array of colored squares 50% valid	T vs. L	RT (% correct)	Group-averaged	**Occipital:** *LO, MT, Cuneus, Fusiform, SOG* + Frontal Temporo-Parietal	No	No
	SOA=2160 ms							
61	color change of a peripheral placeholder 75% valid	50% valid	face identity	RT (% correct)	ROI-based (FFA) + Group-averaged	**Occipital:** *FFA* + Frontal Temporo-Parietal	Yes	No
	SOA=300 ms							
Current study	central bar 75% invalid SOA=310 ms	peripheral bar 50% valid SOA=110 ms	orientation	d-prime (RT)	ROI-based	**Occipital:** *V1, V2, V3, V3A, hV4, LO1*	Yes	Yes
Studies NOT including Occipital areas								
Detection tasks								
60	central arrow 80% valid	peripheral dot 50% valid	filling of a square	RT	Group-averaged	Frontal Temporo-Parietal	No	No
	SOA=400, 550 or 700 ms							

Discrimination tasks								
75	thickening of part of central fixation diamond 80% valid SOA=200, 400 or 800 ms	luminance change of peripheral placeholder 50% valid SOA= 100, 150 or 200 ms	X vs. +	RT (% correct)	Group-averaged	Frontal Temporo-Parietal	No	No
76	color change of central fixation 83% valid SOA=700–1000 ms	whitening of peripheral placeholder 50% valid SOA=100–300 ms	central color of a checkerboard	Inverse efficiency = RT/%correct	Group-averaged	Frontal Temporo-Parietal	No	No
39	central bar 75% invalid SOA=310 ms	peripheral bar 50% valid SOA=110 ms	orientation	d-prime (RT)	ROI-based	Temporo-Parietal	Yes	No
37	color change of central fixation 83% valid SOA=700–1000 ms	whitening of peripheral placeholder 50% valid SOA=100–300 ms	central color of a checkerboard	Inverse efficiency = RT/%correct	Group-averaged	Frontal Temporo-Parietal	No	No

Table 1. fMRI studies comparing endogenous and exogenous attention independently in human participants. *For each study, we report the cueing manipulation for each attention condition (SOA: Stimulus Onset Asynchrony is the duration of the cue + interval before onset of the target), the task performed by the participants, the dependent variable reported in the publication, whether the analysis was based on group-averaging or ROI-based on single-participants, the cortical areas reported in the publications and showing significant activation due to cue and/or target, whether or not eye data were monitor in the scanner or used for the subsequent fMRI analysis, and whether the fMRI BOLD signal was correlated with behavioral measures. IOG: Inferior Occipital Gyrus; LG: Lingual Gyrus; LO: Lateral Occipital area; LOG: Lateral Occipital Gyrus; MOG: Middle Occipital Gyrus; MT: Middle Temporal area; SC: Striate Cortex; SOG: Superior Occipital Gyrus; TOS: Temporal-Occipital Sulcus.*

Here, with the same participants, task, stimuli and task difficulty for all attention manipu-

lations, we tested the following four predictions: (1) Pre-cueing should induce an attentional mod-

ulation of fMRI activity (e.g. [74]), higher in the valid than the invalid condition in which attention

needs to be reoriented to the opposite location (e.g. [39,47,77,78]) to perform the task. (2) Both en-

dogenous and exogenous pre- and post-cueing effects should be stronger in visual regions con-

tralateral to the attended hemifield (e.g. [23,74]). (3) Pre-cueing endogenous attention, but not ex-

ogenous attention, should increase activity modulations along the visual hierarchy (e.g. higher in

V4 than in V1; [23,28]; for reviews [30,36]). For endogenous attention, a top-down process, modulations

from higher-order, fronto-parietal attentional regions would send feedback information to visual

cortex with diminishing effects in earlier visual areas [28,29,36,79-82]. (4) Post-cueing endogenous [23,65-

67], but not exogenous [74], attention should induce attentional modulation of fMRI activity in early

visual areas. Voluntary, endogenous attention would facilitate reading out perceptual information

[39], and modulate its processing [23,65-67]. The only fMRI study assessing the effects of post-cueing

exogenous attention found no attentional modulation of fMRI activity in early visual areas [74]; some

behavioral studies report post-cueing effects [68,69] but others found no such effects [70-73].

To test these four predictions, we measured fMRI activity and compared the effects of

endogenous and exogenous attention in early visual areas while the same participants performed

the same task –a 2-AFC orientation discrimination task, contingent upon contrast sensitivity [4,83,84].

We used a fully-crossed design: two attention conditions –endogenous or exogenous attentional

orienting– and two types of cueing –pre- or post-cue. We evaluated fMRI activity at both the

attended and the un-attended locations, given the ubiquitous performance tradeoffs at attended

(benefits) and unattended (costs) locations compared to a neutral condition (e.g. [3,25,85-89]), and

the importance of evaluating both the orienting and reorienting of attention [39,77,90], critical in an

ever-changing environment [77,91].

This is the first study to systematically evaluate and directly compare how pre and post-

orienting, and reorienting, of endogenous and exogenous attention modulate neural activity in

visual cortex to affect behavior. The results indicate that these two types of spatial covert attention

distinctly modulate activity in individual retinotopic visual cortical areas. These differences in

activity are consonant with differential engagement of top-down and bottom-up processes and

their respective temporal dynamics. These results suggest that endogenous attention facilitates

both the encoding and the readout of visual information whereas exogenous attention only facil-

itates the encoding of information.

RESULTS

Endogenous and exogenous attention improve performance

Participants performed a 2-AFC orientation-discrimination task under two attentional con-

ditions (exogenous or endogenous attention), when the cue was presented either before (pre-

cue) or after (post-cue) the grating stimuli (see Methods), and while their brain activity was meas-

ured with fMRI (**Figure 1**).

In each condition, we calculated performance accuracy (d'), as the main dependent varia-

ble, for each participant separately (**Figure 1B**, top row). A three-way repeated measures 2x2x2

ANOVA (exogenous/endogenous x valid/invalid x pre/post-cue) revealed higher performance for

valid than invalid cues ($F_{(1,4)}=23.6$, $p=0.008$), that exogenous and endogenous cues were sta-

tistically indistinguishable ($F_{(1,4)}<1$), and that there was no significant difference between pre-

and post-cues ($F_{(1,4)}<1$). All the two and three-way interactions were not significant ($F<1$).

Figure 1. Experimental protocol. A. *Participants performed a 2-AFC orientation-discrimination*
task. Pre-cues and post-cues were presented before and after the stimuli, respectively.

 Exogenous cues appeared in the periphery, above one of the two stimulus locations. Endoge-
nous cues appeared at the center of the screen (the display is not at scale for visibility purposes)
and indicated one of the two stimulus locations The ISI between the cue and the gratings was
shorter for the exogenous (50 ms) than the endogenous (250 ms) conditions. A response cue
indicated the target location and instructed participants to indicate whether the target grating was
tilted clockwise or counterclockwise of vertical by pressing one of two keys. To provide feedback,
the fixation-cross turned green or red for a correct or an incorrect answer, respectively. **B.** *Be-*
havioral performance averaged across participants (black dots) and for each of them (colored
dots; n=5) for endogenous (left) and exogenous (right) attention. (Top) Performance accuracy
(d'; top) and median reaction time (bottom) as a function of cueing condition. V, valid cue (same
location of pre-cue/post-cue as response cue). IN, invalid cue (different location of pre-cue/post-
cue than response cue). Pre, pre-cue presented before the stimuli. Post, post-cue presented after
the stimuli. Valid cues induced more accurate and faster responses (there was no speed-accu-
racy trade-off). Error bars, ± 1 SEM across participants.

In each condition, we also calculated reaction time, as a secondary dependent variable (note that the task is not optimize for reaction time's effect due to the long response cue window during which participants are not allowed to provide their response), for each participant separately (**Figure 1B**, bottom row). A three-way repeated measures ANOVA revealed faster reaction times for valid than invalid cues ($F_{(1,4)}=62.3$, $p=0.001$). There was no significant difference between exogenous and endogenous cues ($F_{(1,4)}=2.7$, $p=0.17$), or between pre- and post-cues ($F_{(1,4)}=1.7$, $p=0.27$). Two significant interactions indicated that the differences between valid and invalid cues ($F_{(1,4)}=16.2$, $p=0.02$) and between pre- and post-cues ($F_{(1,4)}=8.1$, $p=0.047$) were more pronounced for endogenous attention than for exogenous attention.

These behavioral results, which are consistent with previous findings [3,4,6,25,84,86,92], show that attention improved orientation discrimination (d' and reaction time), with no evidence of a speed-accuracy trade-off, and similarly for both types of attention and for both pre- and post-cues. Thus, the behavioral effects confirm the successful manipulation of endogenous and exogenous attention, for both pre- and post-cues, consistently across individuals.

Attentional modulation of perceptual and post-perceptual information processing in visual cortex

Visual areas were mapped in each participant following retinotopic mapping procedures (**Figure 2**, left panel) and a targeted stimulus localizer (**Figure 2**, right panel), and regions of interest (ROIs) were selected based on previous literature: V1, V2, V3 (for V2 and V3, ventral and dorsal ROIs were averaged), V3A, hV4 and LO1 (e.g. [93-96]).

Activity was higher for contralateral than ipsilateral brain areas. For each ROI, we meas-

ured the fMRI response amplitudes for each type of attention – exogenous and endogenous –

and each cueing condition – pre- and post-cueing, for the contralateral and ipsilateral side to the

cued (attended) location. We analyzed the fMRI responses evoked by each type of attention in

the contralateral and ipsilateral brain regions relative to the cue location (**Supplemental Figure**

1). ANOVAs indicated that there was higher contralateral than ipsilateral activity across brain

areas (endogenous: $F_{(1,4)}=59.9$, $p=0.0015$; exogenous: $F_{(1,4)}=218.8$, $p=0.0001$). For both types

of attention, this difference was more pronounced for valid than invalid cues (endogenous:

$F_{(1,4)}=8.6$, $p=0.04$; exogenous: $F_{(1,4)}=21.1$, $p=0.01$). In the following analyses, we then concen-

trate on fMRI activity in the contralateral ROIs to the attended location.

Figure 2. Retinotopic mapping and stimulus localizer of a representative participant. A
flattened representation of the right hemisphere occipital pole. ***Left,*** *map of polar angle. The color*
of each voxel in the map represents the response phase of the time series elicited by a rotating
wedge stimulus. ***Right,*** *stimulus localizer. The black outlines correspond to the retinotopic areas*
(defined for each participant based on their polar angle maps). The color of each voxel indicates
the phase of the response to the grating stimuli presented in the lower left visual field.

fMRI responses correlated with behavioral variability. We evaluated the degree to which

inter-individual variability in behavioral performance co-varied with variability in fMRI responses.

We computed the correlation (**Figure 3A**), across participants, ROIs, types of attention and cue-

ing conditions, between the fMRI responses (percent change in image intensity) and behavioral

performance accuracy (d'). We found a positive correlation between fMRI activity and d' (Pearson

correlation r = 0.23, p = 0.003), which was present for both attention types (endogenous: Pearson

correlation r = 0.3, p = 0.003; exogenous: r = 0.2, p = 0.02). We further showed that fMRI activity

was overall higher for valid than invalid trials (t(4) = 4.0, p = 0.0163, Cohen's d = 1.78). This was

the case for each participant (**Figure 3B**). Together, these results are consistent with previous

findings showing that fMRI response correlates with behavioral variability [97-99], and that atten-

tional orienting increases fMRI signal in early visual cortex (for review [30]).

*Figure 3. Inter-individual fMRI response. **A.** Positive correlation between fMRI response and*
behavioral performance (Pearson correlation r = 0.23, p = 0.0003). Each dot indicates fMRI re-
sponse amplitude measured from one ROI and one participant, separately for each attentional
*condition (pre and post-cue, valid and invalid). **B.** fMRI response difference between valid and*
invalid trials combined across all conditions and ROIs for each participant (colored dots), and
across participants (black dot; average significantly positive: t(4) = 4.0, p = 0.0163, Cohen's d =
1.78). Error bar, ± 1 SEM across participants.

We then tested our two novel predictions: (1) attentional modulation of fMRI stimulus-

evoked activity increases along the visual hierarchy for endogenous pre-cueing, but is constant

for exogenous pre-cueing; and (2) attentional modulation of fMRI stimulus-evoked activity for

endogenous post-cueing is constant along the visual hierarchy, whereas there is no attentional

modulation for exogenous post-cueing. We measured the differences in activity between valid

and invalid trials for each type of attention and for pre- and post-cues, across the hierarchy of

visual cortical areas. fMRI responses were significantly larger for valid than invalid cues, for both

endogenous and exogenous cues, and for both pre- and post-cues. A three-way repeated

measures ANOVA (2 exogenous/endogenous x 2 pre/post-cue x 6 ROIs) of the difference be-

tween valid and invalid conditions revealed significant main effects of exogenous/endogenous

condition (F = 10.0, p = 0.0341) and of ROI (F = 5.5, p = 0.0024), as well as an interaction

between pre/post-cue and ROI (F = 4.1, p = 0.0103). Thus, we investigated modulations of fMRI

activity separately for pre and post-cueing attention conditions.

Linear mixed models [100] were then used to test our main predictions; namely (1) increase

of fMRI activity along the visual hierarchy for pre-cueing endogenous attention but not exogenous

attention, and (2) post-cue attentional modulation of fMRI activity for post-cueing endogenous

attention but not exogenous attention. We implemented two models, one for the pre-cueing and

the other for the post-cueing condition, to test the link between fMRI activity and the interaction

between ROI and endogenous/exogenous attention conditions. In each model, we entered as

fixed effect the ROI and the endogenous/exogenous conditions into the model, as well as their

interaction. As random effects, we had participant's intercepts and slopes for the effect of ROI

and endogenous/exogenous conditions. Visual inspection of residual plots did not reveal any

obvious deviations from homoscedasticity or normality. P-values were obtained by likelihood ratio

tests. For the pre-cueing condition, we observed a significant effect of ROI (t(56)=3.8, p=0.0004,

estimate=0.02±0.005, standard error), as well as a significant interaction between ROI and en-

dogenous/exogenous conditions (t(56)=-2.4, p=0.018, estimate=-0.007±0.003, standard error).

For the post-cueing condition, we observed a significant effect of the fixed effect intercept

(t(56)=2.2, p=0.03, estimate=0.04±0.02, standard error). Finally, we performed independent re-

gression analyses on each pre-/post-cueing and endogenous/exogenous attention conditions

(**Figure 4**). We observed a significant increase along the hierarchy of the visual areas of the

activity difference evoked by valid and invalid trials for the endogenous pre-cueing condition (F =

25.8, p = 0.007, R^2 = 0.87), but not for the exogenous pre-cueing condition (F = 2.8, p = 0.168,

$R^2 = 0.41$), endogenous post-cueing (F = 3.0, p = 0.161, $R^2 = 0.42$) or exogenous post-cueing

conditions (F = 0.1, p = 0.774, $R^2 = 0.02$).

Together, the results of the linear mixed models and the regression analyses (**Figure 4**)

showed differential effects between pre-/post-cueing and endogenous/exogenous attention con-

ditions. The difference in fMRI activity between valid and invalid trials increases along the hierar-

chy of the visual areas in pre-cueing of endogenous attention but not of exogenous attention.

Moreover, there is a difference in fMRI activity between valid and invalid trials with post-cueing

of endogenous attention that is constant along the hierarchy (significant intercept), but absent for

post-cueing of exogenous attention. These analyses are consistent with post-hoc t-tests per-

formed for each condition and ROI (**Figure 5**).

Figure 4. Specificity of pre and post-cueing for endogenous and exogenous attention. *The*
difference in fMRI response between valid and invalid is plotted separately for pre- and post-cue
*conditions for each ROI. *, Statistically significant regression analysis (p < 0.05).*

For both endogenous attention (p < 0.05 for V3a, V4 and LO1; trend (p = 0.077) for V3; t-

tests) and exogenous attention (p < 0.05 for V3, V3A, hV4 and LO1; trend (p = 0.074) for V2; t-

tests) pre-cues elicited greater fMRI activity for valid than invalid cues (**Figure 5**, top-left and top-

right panels). Furthermore, for endogenous attention (p < 0.05 for V2, V3, V3A, hV4 and LO1;

trend (p = 0.077) for V1; t-tests; **Figure 5**, bottom-left panel), but not for exogenous attention (all

p > 0.1; t-tests; **Figure 5**, bottom-right panel), post-cues elicited greater fMRI activity for valid

than invalid cues in occipital areas (detailed statistics are presented in Supplementary Table 1).

Critically, these effects were consistently observed across participants (**Figure 6**). Taken to-

gether, these results confirm both of our predictions.

Figure 5. Single ROI responses for pre and post-cueing and for both endogenous and exogenous attention. The difference in fMRI response between valid and invalid is plotted sep- arately for pre and post-cue conditions for each ROI (same data as in Fig 4). *, Statistically sig- nificant difference between valid and invalid, separately for pre and post-cueing (p < 0.05). #, trend (p < 0.1). Error bars on plots are ± 1 SEM.

307
Figure 6. Specificity of each visual ROI for endogenous and exogenous attention. *fMRI*
response amplitude was measured for each attentional condition. V: valid cue condition (target
location matches the location indicated by pre-cue/post-cue). IN: invalid cue condition (target
location at the opposite location relative to the pre-cue/post-cue). Pre: pre-cue presented before
the grating stimuli. Post: post-cue presented after gratings. All four conditions in the contralateral
ROI. Distractor (Dist): all conditions averaged in the ipsilateral ROI. Each colored point
corresponds to the data of one participant. The black points represent the average across all 5
*participants. *, Statistically significant difference between valid and invalid, separately for pre and*
post-cueing (p < 0.05). #, trend (p < 0.1). Detailed statistics are presented in Supplementary
Table 1. Error bars (vertical lines) are plotted on the difference between valid and invalid and
represent ± 1 SEM across participant.

DISCUSSION

This is the first study to compare pre and post-orienting, and reorienting, of endogenous

and exogenous attention in visual cortex, while concurrently assessing visual performance using

well-established psychophysical protocols to manipulate attention. The fact that the same partic-

ipants performed the same orientation discrimination task with the same stimuli and task difficulty

under different attentional manipulations enabled us to isolate the fMRI activity induced by each

type of attention during orienting and reorienting.

Previous studies comparing endogenous and exogenous attention conditions state various

attentional effects in early visual areas (see **Table 1**). Some report differential effects between

these attention conditions in the right LO and MT [49], others in the cuneus [48-50], the occipital gyrus

(SOG, MOG, LOG; [49,50,62]), the fusiform area [49,61,62], and the TOS, LG and SC [62]. This lack of clear

picture regarding the differential impact of endogenous and exogenous spatial attention is likely

due to the methodological differences and limitations we discussed in the Introduction (see also

Table 1). In any case, it is often assumed that the effects of endogenous and exogenous attention

are the same in visual occipital areas [30,35,36,47,48].

To further our knowledge of the neural correlates of attention, we investigated both atten-

tional orienting (valid cueing) and reorienting (invalid cueing), critical in an ever-changing envi-

ronment [77,91]. Furthermore, given ubiquitous performance tradeoffs between attended (benefits)

and unattended (costs) locations (e.g. [3,25,87,88]), we assessed activity at both attended (contrala-

teral ROI) and unattended (ipsilateral ROI) locations. Finally, we investigated how attentional

effects varied as a function of pre- and post-cueing, thus contrasting the neural correlates of

perceptual and post-perceptual processing of information.

There was an overall positive correlation between performance in the orientation discrimi-

nation task and the degree of attentional modulation in fMRI activity. This result was expected,

indicating that as discriminability increases so does the attentional modulation in fMRI activity.

But only very few studies on the effect of spatial attention in early visual areas have reported

such a correlation [97-99]. The behavioral effects obtained in the scanner are consistent with psy-

chophysical studies. The enhanced performance brought about by the valid, but uninformative

peripheral pre-cue is consistent with an automatic, bottom-up involuntary capture of exogenous

attention (e.g. [3,6,25,87,101]). The enhanced performance brought about by the valid, informative cen-

tral precue is consistent with a top-down, voluntary deployment of endogenous attention (e.g.

[3,6,25,102,103]).

In the endogenous attention condition, there was an increase in attentional modulation of

stimulus-evoked activity along the hierarchy of visual areas. There is no consensus regarding the

visual hierarchy beyond area V3 (e.g. [104-108]), especially regarding V3A. However, most authors

agree that hV4 precedes LO1 [104-107]. In any case, our data are consistent with either a strict

hierarchy or with V3A and hV4 being at the same level. Such a pattern is consistent with previous

studies suggesting that endogenous attention is a top-down modulation from frontal and parietal

areas feeding back to visual cortex, with diminishing effects in earlier visual areas [28,36,79,80]. In-

consistent with previous studies (e.g. [23,25,29,109,110]), there was no evidence for attentional modu-

lation in V1. It might be that attentional modulation of V1 activity is more variable than other visual

cortical areas, making it harder to detect (see also [28,74]). Methodological differences between this

and previous studies may have contributed to weakening the effect of attention in V1. The accrual

time in the current endogenous condition was relatively short (1300 ms in the valid condition and

500 ms in the invalid condition) compared to previous studies investigating endogenous, volun-

tary attention, in which the cue and/or stimuli were presented for a long duration to maximize

BOLD measurements (e.g. [23,29,109,110]). We chose the minimum SOA (317 ms) at which perfor-

mance benefits plateau to have as comparable conditions as possible to the accrual time with

the SOA for exogenous attention (117 ms). This short accrual time may have limited the effects

of attentional feedback to V1.

In the exogenous attention condition, in contrast to the endogenous attention, the atten-

tional modulation was approximately constant across the visual hierarchy. Some previous studies

have reported a similar effect [111,112], others a decrease [113], and yet others an increase [74,114] across

the visual areas. This difference might be explained by different task parameters. For example,

in the Liu et al. (2005) study [74], participants knew which of the two stimuli was the target they had

to discriminate while the stimuli were presented, as one stimulus was vertical and the other was

tilted to the left or the right. In the present study, both stimuli were independently tilted and par-

ticipants had to wait for the response cue to appear to know which one was the target and which

one was the distractor.

Unlike in the endogenous pre-cueing condition in which the attention effect increased along

the processing stream, for the endogenous post-cueing effect there was no evidence that it varied

across these visual areas. The constant effect in the post-cue condition could be due to the con-

tribution of two counteracting factors: (1) the fMRI response evoked by the stimulus in early visual

areas may decrease along the visual hierarchy [115]; (2) the top-down modulations from frontal and

parietal areas feedback to visual cortex with diminishing effects in earlier visual areas [23,28,116].

In the exogenous condition, there was no significant post-cueing effect on early visual ar-

eas. This result is consistent with that of Liu et al. (2005) [74], who while evaluating exogenous

attention effects on occipital cortex included a post-cue condition to rule out sensory contamina-

tion of the cue (i.e. sensory response evoked by the cue itself) contributing to the enhanced

BOLD activity found in their pre-cue condition. In addition to ruling out a possible sensory con-

tamination, the present results show that, in contrast to endogenous attention, exogenous atten-

tion does not aid in the selective readout of information.

The ROI-based analysis that we followed here enabled us to compare contralateral and

ipsilateral modulation of BOLD activity, thus providing additional information regarding the differ-

ences in processing dynamics of both types of attention. We observed a larger difference be-

tween contralateral and ipsilateral areas for the valid than the invalid cueing condition. This effect

could be due to the fact that for the former, participants were attending to the same location

throughout the trial, whereas for the latter, when the response cue did not match the pre-cue,

participants had to switch their spatial attention to the opposite stimulus location, thus activity at

that new location would be accumulated for less time. For instance, for endogenous attention, for

the valid condition participants had been processing the target for almost 500 ms before the

response cue appeared. When the response cue matched the pre-cue, participants continued

processing and reading out the signal from that location for up to 800 ms (they were not allowed

to give an answer before the end of the response cue period). But when the response cue did

not match the pre-cue, then participants had to switch after 500 ms to the other location (ipsilat-

eral) thus accumulating less activity. Similarly, the accumulation time for the invalid cue condition

in exogenous attention was only about 300 ms. This accrual time explanation could also account

for the larger difference between contralateral and ipsilateral for pre-cues than post-cues, i.e.

there is a 300 ms accumulation when the exogenous pre-cue is invalid, while only 100 ms when

the post-cue in invalid. Likewise, the larger modulatory effect for endogenous relative to exoge-

nous attention is consistent with the difference in accrual time.

In a recent study [39] we demonstrated that sub-regions of the Temporo-Parietal Junction

(TPJ) that respond specifically to visual stimuli are more active when attention needs to be spa-

tially reoriented (invalid cueing) than when attention remains at the cued location (valid cueing),

and that partially overlapping specific visual sub-regions mediate reorienting after orienting of

endogenous or exogenous attention. Both the present and the TPJ studies provide a compre-

hensive investigation of endogenous and exogenous attention, and pave the way for rigorous

psychophysics informed, neuroimaging studies of covert, spatial attention. Here, we concen-

trated the analysis on visual cortical areas in the occipital lobe because the slice prescription

covered only a limited portion of the brain. The present findings further our knowledge of the

neurophysiological bases of covert attention and have implications for models of visual attention,

which should consider not only the similarities, but also the differences in the orienting and reor-

ienting of endogenous and exogenous attention in occipital areas reported here.

CONCLUSION

The present results show some similarities and reveal important differences in the specific

neural correlates of endogenous and exogenous attention on early vision: An increasing modu-

lation of fMRI activity for pre-cueing endogenous attention, but constant modulation for exoge-

nous attention, along the hierarchy of visual occipital areas, as well as a reliable and constant

modulation of fMRI activity for post-cueing endogenous attention in occipital areas but not for

exogenous attention. These results suggest that endogenous attention facilitates both the en-

coding and the readout of visual information whereas exogenous attention only facilitates the

encoding of information.

MATERIALS and METHODS

The behavioral methods employed in this study and the behavioral results are the same as those we reported in a recent study, in which we compared activity in TPJ during orienting and reorienting of endogenous and exogenous attention [39]. To maximize the effects of these two types of attention, i.e. the benefits at the attended location and concurrent costs at the unattended location, we used optimal spatial and temporal parameters (reviews by [1,2]). To enable direct comparison between endogenous and exogenous attention, the same participants performed the same orientation discrimination task under both types of attention. The fMRI methods employed in this study are the same as those used in that study [39], but here, instead of analyzing TPJ activity, we analyzed activity in occipital areas.

Participants

Five participants (two male and three female, 24-30 years-old) participated in the study. They all had normal or corrected-to-normal vision. The University Committee on Activities Involving Human Subjects at New York University approved the experimental protocol (IRB # 10-7094), and participants provided written informed consent. Our study used single-participant ROI-based analysis, and thus had a small sample size. The same sample size that has been used in many fMRI studies in our labs (e.g. [23,25,39,74,99,117-120]), as well as in other labs (e.g. [121-125]). Each participant performed nine scanning sessions: one session to obtain a set of three high-resolution anatomical volumes, two sessions for retinotopic mapping, three sessions for the exogenous attention condition and three sessions for the endogenous attention condition (with the order counterbalanced among participants). Participants performed several practice sessions outside the scanner prior to the first scanning session of each attention condition.

Stimuli

Stimuli were generated on a Macintosh computer using the MGL toolbox [126] in MATLAB (MathWorks). Stimuli were presented on a flat-panel display (NEC, LC-XG250 MultiSync LCD 2110; refresh rate: 60 Hz; resolution: 1024 x 768 pixels) positioned at the rear of the scanner bore and housed in a Faraday box with an electrically conductive glass front. The display,

calibrated and gamma corrected using a linearized lookup table, was at a viewing distance of

172 cm from the participant, and visible through an angled mirror attached to the head coil. A

central, white fixation cross (0.3°) was presented throughout the experiment. The two stimuli were

two 4-cpd gratings windowed by raised cosines (3° of diameter; 7% contrast), one in each bottom

quadrant (5° horizontal eccentricity; –2.65° altitude). Both endogenous cues and exogenous cues

were white rectangles (0.7°). The endogenous cues appeared adjacent to the fixation cross indi-

cating one of the two lower quadrants (0.35° horizontal eccentricity from the edge of the fixation

cross, and 0.35° altitude). The exogenous cues appeared adjacent to an upcoming grating stim-

ulus, vertically aligned with the stimulus and above the horizontal meridian (1° away from the

edge of the grating stimulus; and the edge of the cue 4.44° horizontal eccentricity from the edge

of the fixation cross).

Behavioral procedure

An exogenous attention condition trial lasted 1700 ms, whereas an endogenous attention

condition trial lasted 1900 ms, the only difference being the stimulus-onset asynchronies (SOA)

between the cue and the display; the timing of all the visual stimuli was the same in both attention

conditions (**Figure 1**; the display is not at scale for illustration purposes). In the pre-cue condition

(40% of the trials), a cue preceded the two gratings. In 40% the post-cue condition, the cue

followed the presentation of the gratings. In 'cue-only' trials (10% of the trials), the gratings were

not presented. In 'blank' trials (10% of the trials), neither a cue nor the gratings were presented.

These trials were then included in the GLM analysis to model the contribution of the visual signal

produced by the cue (see MRI procedure). For both pre-cue and post-cue trials, participants were

asked to press one of two keys to report the orientation of a target grating, i.e., clockwise or

counter-clockwise compared to vertical. Participants pressed a third key in the case of cue-only

and blank trials.

In both exogenous and endogenous condition, cues were presented for 67 ms, indicating

either the bottom left or right quadrant of the screen. The inter-stimulus interval (ISI) between the

cue and the grating stimuli was 50 ms for exogenous and 250 ms for endogenous conditions,

resulting in SOA of 117 ms and 317 ms. We used the same timings for pre- and post-cue

conditions (e.g. [23,63,74,127]). These delays are optimal to manipulate exogenous and endogenous

attention, while keeping the trial duration as similar as possible, and have been shown to max-

imize the behavioral consequences of each attention condition [44,128-130].

The behavioral effects of endogenous attention are sustained (e.g. [102]) and thus, as shown

in ERP studies (e.g. [131]), are still present in later brain activity. Additionally, during 300 ms follow-

ing cue onset, the brain responses elicited by exogenous and endogenous cues differ (for review

[1]). The two grating stimuli were then displayed for 50 ms. For the postcue trials we kept the

timings of cue and stimuli constant but inverted the order of their presentation (e.g. [23,63,74,127]). A

response cue, presented for 800 ms at the end of the trial after both the cue and the stimuli had

disappeared, indicated which one of the two gratings was the target (50% of the trials on the right

and the remaining 50% and on the left). The maximum delay between the offset of the grating

stimuli and the onset of the response cue was shorter (~400 ms max in the endogenous condi-

tion) than typically associated with a demand for working memory (>600 ms; [132]). Immediately

following each trial, a change of color of the fixation cross provided visual feedback to the partic-

ipants, i.e. green for correct or red for incorrect responses. The fixation cross did not change

color if participants had missed the response window, i.e. if they had not pressed any key after

530 ms.

In the exogenous attention condition, a peripheral cue was presented, which was not in-

formative regarding the target location or orientation. When the cue location matched the target

location, it was considered a valid trial (50% of the trials), otherwise it was considered an invalid

trial (the remaining 50% of the trials). In the endogenous attention condition, a central cue pointed

to either the left or right quadrant. The cue was informative of the target location but not its ori-

entation (75% valid trials and 25% invalid trials). Participants were informed of this validity. It is

important to note that: (1) cue validity does not affect cueing effectiveness for exogenous atten-

tion, although it does so do endogenous attention (e.g. [3,133,134]); (2) the short timing and non-

informative cue in the exogenous condition ensured that no voluntary component could be in-

volved in the exogenous attentional effect.

Endogenous and exogenous attention conditions were performed in separate sessions to

ensure optimal manipulation of each attention system. Participants first performed two practice

sessions outside the scanner before the first session of each attentional scanning condition. To

equate task difficulty for both attention conditions, using a staircase procedure, the tilt of the

target grating was adjusted for each participant to achieve ~80% correct performance in the valid

trials in each attention condition. In each of the six experimental scanning sessions (three ses-

sions of exogenous attention and three of endogenous), participants performed 14 runs of 40

trials each, as well as a run of stimulus localizer (see MRI procedure). The tilt was then adjusted

between runs to maintain overall performance at ~80% correct.

Eye position was monitored during all scanning sessions using an infrared video camera

system (Eyelink 1K, SR Research, Ottawa, Ontario, http://www.sr-research.com/EL_1000.html).

Trials in which the participants blinked or broke fixation (1.5° radius from central fixation) at any

point from fixation onset to response cue offset were identified and regressed separately in the

MRI analysis, and removed from the behavioral analysis (13% ± 4% of the trials on average

across all participants).

MRI Procedure

Scanning

Imaging was conducted on a 3T Siemens Allegra head-only scanner (Erlangen, Germany),

using a Siemens NM-011 head coil (to transmit and receive) to acquire anatomical images, a

receive-only 8-channel surface coil array (Nova Medical, Wilmington, MA) to acquire functional

images. To minimize participants' head movements, padding was used.

For each participant, three high-resolution anatomic images were acquired in one scanning

session, using a T1-weighted magnetization-prepared rapid gradient echo (MPRAGE) sequence

(FOV = 256 x 256 mm; 176 sagittal slices; 1 x 1 x 1 mm voxels), and were co-registered and

averaged. Using FreeSurfer (public domain software; http://surfer.nmr.mgh.harvard.edu), the

gray matter was segmented from these averaged anatomical volumes. All subsequent analyses

were constrained only to voxels that intersected gray matter.

T2*-weighted echo-planar imaging sequence (i.e. functional images; TR = 1750 ms; TE =

30 ms; flip angle = 90°) measured blood oxygen level-dependent (BOLD) changes in image in-

tensity [135]. In each volume, 28 slices covered the occipital and posterior parietal lobes and were

oriented 45° to the calcarine sulcus (FOV = 192 x 192 mm; resolution = 2 x 2 x 2.5 mm; no gap).

To align functional images from different sessions to the same high-resolution anatomical volume

for each participant, we acquired an additional T1-weighted anatomical images in the same slices

as the functional images (spin echo; TR = 600 ms; TE = 9.1 ms; flip angle = 90°; resolution = 1.5

× 1.5 × 3 mm) during each scanning session, and used them in a robust image registration algo-

rithm.

MRI data pre-processing

Imaging data were analyzed in MATLAB, using mrTools [136] and custom software. To allow

longitudinal magnetization to reach steady state, the first eight volumes of each run were dis-

carded. Spatial distortion was corrected using the B0 static magnetic field measurements per-

formed in each session. The functional data were then motion corrected, the linear trend was

removed, and a temporal high-pass filter was applied (cutoff: 0.01 Hz) to remove slow drifts and

low-frequency noise in the fMRI signal.

Retinotopic mapping

We followed well-established conventional traveling-wave, phase-encoded methods. Using

clockwise and counter-clockwise rotating checkerboard wedges, we measured phase maps of

polar angle. Using contracting and expending checkerboard rings, we measured eccentricity

maps [93-96]. **Figure 2** (left panel) shows the visual areas that were drawn by hand on flattened

surface of the brain, following published conventions [95,96,99,137].

Stimulus localizer

In each scanning session of the main experiment, participants completed one stimulus lo-

calizer run (6 runs overall, 4 min each). A run consisted of 16 cycles (17.5 s) of a block alternation

protocol between stimulus on (8.75 s) and stimulus off (8.75 s). Participants only had to fixate the

central cross throughout each run. The stimuli were at the same location, and of the same size

and spatial frequency as those in the main experiment, except at full contrast and their phase

and orientation changed randomly every 200 ms to avoid adaptation. To define the cortical rep-

resentation of the gratings, we then averaged the data across the 6 runs and followed the same

methods as for the retinotopic mapping. Voxels that responded positively during the blocks when

the grating stimuli were presented were used to restrict each retinotopic ROI. The fMRI time

series from each voxel were fit to a sinusoid. To be conservative, only voxels whose best-fit

sinusoid had a phase value between 0 and pi, and a coherence between the best-fit sinusoid and

the time series greater than 0.2 were included in the ROI (**Figure 2**, right panel). Analysis per-

formed without restricting the ROI to this coherence level yielded similar results.

Event-related analysis

fMRI time series were averaged across voxels in each ROI (separately for each hemi-

sphere) and then concatenated across runs. The data were denoised using GLMDenoise [138], and

fMRI response amplitudes were computed using linear regression, with twelve regressors: 8

combinations of right and left valid and invalid pre- and post-cue, right and left cue-only, blank

(no cue nor stimulus) and eye-movements (blink or broken fixation). For each ROI in each hem-

isphere, the resulting fMRI response amplitudes (for correct trials only) were then averaged

across participants.

Acknowledgments: This work was supported by: NIH RO1-EY019693 to MC and DJH; NIH
RO1-EY027401 to MC; the FYSSEN Foundation and the Philippe Foundation to LD; and the
Center for Brain Imaging of New York University. We want to thank members of the Carrasco lab
and Jonathan Winawer for constructive comments on the manuscript.

Competing interests: The authors declare no competing interests.

Author contributions: Laura Dugué: Conceptualization, Data curation, Formal analysis,
Funding acquisition, Investigation, Visualization, Methodology, Writing - original draft, Writing -
review and editing. **Elisha Merriam:** Data curation, Formal analysis, Investigation, Visualization,
Methodology, Writing - review and editing. **David Heeger:** Resources, Supervision, Funding
acquisition, Visualization, Methodology, Writing - review and editing. **Marisa Carrasco:**
Conceptualization, Resources, Supervision, Funding acquisition, Visualization, Methodology,
Writing - original draft, Project administration, Writing - review and editing.

Data availability: Data relative to the main results of this study (i.e. Figures 4, 5 and 6) are made
available via OSF at [link provided upon acceptance of the manuscript].

REFERENCES

1. Carrasco, M. Visual attention: the past 25 years. *Vision Res.* **51,** 1484–1525 (2011).
2. Carrasco, M. & Barbot, A. How attention affects spatial resolution. *Cold Spring Harbor Symposia on Quantitative Biology* (2015).
3. Giordano, A. M., McElree, B. & Carrasco, M. On the automaticity and flexibility of covert attention: a speed-accuracy trade-off analysis. *Journal of Vision* **9,** 30.1–10 (2009).
4. Pestilli, F., Ling, S. & Carrasco, M. A population-coding model of attention's influence on contrast response: Estimating neural effects from psychophysical data. *Vision Res.* **49,** 1144–1153 (2009).
5. Ling, S. & Carrasco, M. Sustained and transient covert attention enhance the signal via different contrast response functions. *Vision Res.* **46,** 1210–1220 (2006).
6. Dosher, B. A. & Lu, Z. L. Mechanisms of perceptual attention in precuing of location. *Vision Res.* **40,** 1269–1292 (2000).
7. Barbot, A., Landy, M. S. & Carrasco, M. Differential effects of exogenous and endogenous attention on second-order texture contrast sensitivity. *Journal of Vision* **12,** 6–6 (2012).
8. Yeshurun, Y. & Carrasco, M. Attention improves or impairs visual performance by enhancing spatial resolution. *Nature* **396,** 72–75 (1998).
9. Carrasco, M., Giordano, A. M. & McElree, B. Attention speeds processing across eccentricity: Feature and conjunction searches. *Vision Res.* **46,** 2028–2040 (2006).
10. Yeshurun, Y., Montagna, B. & Carrasco, M. On the flexibility of sustained attention and its effects on a texture segmentation task. *Vision Res.* **48,** 80–95 (2008).
11. Barbot, A. & Carrasco, M. Attention Modifies Spatial Resolution According to Task Demands. *Psychological Science* **28,** 285–296 (2017).
12. Jigo, M. & Carrasco, M. Attention alters spatial resolution by modulating second-order processing. *Journal of Vision* **18,** 2 (2018).
13. Anton-Erxleben, K. & Carrasco, M. Attentional enhancement of spatial resolution: linking behavioural and neurophysiological evidence. *Nat. Rev. Neurosci.* **14,** 188–200 (2013).
14. Carrasco, M. & Yeshurun, Y. Covert attention effects on spatial resolution. *Prog. Brain Res.* **176,** 65–86 (2009).
15. Reynolds, J. H. & Heeger, D. J. The Normalization Model of Attention. *Neuron* **61,** 168–185 (2009).
16. Martínez-Trujillo, J. C. & Treue, S. Attentional Modulation Strength in Cortical Area MT Depends on Stimulus Contrast. *Neuron* **35,** 365–370 (2002).
17. McAdams, C. J. & Maunsell, J. H. R. Effects of Attention on Orientation-Tuning Functions of Single Neurons in Macaque Cortical Area V4. *J. Neurosci.* **19,** 431–441 (1999).
18. Mitchell, J. F., Sundberg, K. A. & Reynolds, J. H. Spatial Attention Decorrelates Intrinsic Activity Fluctuations in Macaque Area V4. *Neuron* **63,** 879–888 (2009).
19. Reynolds, J. H., Pasternak, T. & Desimone, R. Attention increases sensitivity of V4 neurons. *Neuron* **26,** 703–714 (2000).
20. Williford, T. & Maunsell, J. H. R. Effects of Spatial Attention on Contrast Response Functions in Macaque Area V4. *J. Neurophysiol.* **96,** 40–54 (2006).
21. Luo, T. Z. & Maunsell, J. H. R. Neuronal Modulations in Visual Cortex Are Associated with Only One of Multiple Components of Attention. *Neuron* **86,** 1182–1188 (2015).
22. Ruff, D. A. & Cohen, M. R. Attention can either increase or decrease spike count correlations in visual cortex. *Nat. Neurosci.* **17,** 1591–1597 (2014).
23. Pestilli, F., Carrasco, M., Heeger, D. J. & Gardner, J. L. Attentional enhancement via selection and pooling of early sensory responses in human visual cortex. *Neuron* **72,** 832–846 (2011).
24. Murray, S. O. The effects of spatial attention in early human visual cortex are stimulus independent. *Journal of Vision* **8,** 2.1–11 (2008).
25. Herrmann, K., Montaser-Kouhsari, L., Carrasco, M. & Heeger, D. J. When size matters: attention affects performance by contrast or response gain. *Nat. Neurosci.* **13,** 1554–1559 (2010).

26. Buracas, G. T. & Boynton, G. M. The effect of spatial attention on contrast response
functions in human visual cortex. *Journal of Neuroscience* **27,** 93–97 (2007).
27. O'Connor, D. H., Fukui, M. M. & Pinsk, M. A. Attention modulates responses in the hu-
man lateral geniculate nucleus - ProQuest. *Nature* **5,** 1203–1209 (2002).
28. Kastner, S., Pinsk, M. A., De Weerd, P., Desimone, R. & Ungerleider, L. G. Increased
Activity in Human Visual Cortex during Directed Attention in the Absence of Visual Stim-
ulation. *Neuron* **22,** 751–761 (1999).
29. Somers, D. C., Dale, A. M., Seiffert, A. E. & Tootell, R. B. Functional MRI reveals spatially
specific attentional modulation in human primary visual cortex. *PNAS* **96,** 1663–1668
(1999).
30. Beck, D. M. & Kastner, S. Neural systems for spatial **attention** in the human brain: Evi-
dence from neuroimaging in the framework of biased competition. *The Oxford Handbook
of Attention* 253–288 (2014).
31. Lu, Z.-L., Li, X., Tjan, B. S., Dosher, B. A. & Chu, W. Attention Extracts Signal in External
Noise: A BOLD fMRI Study. *J Cogn Neurosci* **23,** 1148–1159 (2011).
32. Li, X., Lu, Z.-L., Tjan, B. S., Dosher, B. A. & Chu, W. Blood oxygenation level-dependent
contrast response functions identify mechanisms of covert attention in early visual areas.
Proc. Natl. Acad. Sci. U.S.A. **105,** 6202–6207 (2008).
33. Busse, L., Katzner, S. & Treue, S. Temporal dynamics of neuronal modulation during
exogenous and endogenous shifts of visual attention in macaque area MT. *Proc. Natl.
Acad. Sci. U.S.A.* **105,** 16380–16385 (2008).
34. Wang, F., Chen, M., Yan, Y., Zhaoping, L. & Li, W. Modulation of Neuronal Responses
by Exogenous Attention in Macaque Primary Visual Cortex. *Journal of Neuroscience* **35,**
13419–13429 (2015).
35. Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in
the brain. *Nat. Rev. Neurosci.* **3,** 201–215 (2002).
36. Chica, A. B., Bartolomeo, P. & Lupiáñez, J. Two cognitive and neural systems for en-
dogenous and exogenous spatial attention. *Behavioural Brain Research* **237,** 107–123
(2013).
37. Bowling, J. T., Friston, K. J. & Hopfinger, J. B. Top-down versus bottom-up attention
differentially modulate frontal–parietal connectivity. *Hum Brain Mapp* **9,** 1116 (2019).
38. Hahn, B., Ross, T. J. & Stein, E. A. Neuroanatomical dissociation between bottom–up
and top–down processes of visuospatial selective attention. *Neuroimage* **32,** 842–853
(2006).
39. Dugué, L., Merriam, E. P., Heeger, D. J. & Carrasco, M. Specific Visual Subregions of
TPJ Mediate Reorienting of Spatial Attention. *Cereb. Cortex* 1–16 (2017).
doi:10.1093/cercor/bhx140
40. Geng, J. J. & Vossel, S. Re-evaluating the role of TPJ in attentional control: contextual
updating? *Neurosci Biobehav Rev* **37,** 2608–2620 (2013).
41. Doricchi, F., Macci, E., Silvetti, M. & Macaluso, E. Neural correlates of the spatial and
expectancy components of endogenous and stimulus-driven orienting of attention in the
Posner task. *Cereb. Cortex* **20,** 1574–1585 (2010).
42. Silvetti, M. *et al.* The Response of the Left Ventral Attentional System to Invalid Targets
and its Implication for the Spatial Neglect Syndrome: a Multivariate fMRI Investigation.
Cereb. Cortex 1–12 (2015). doi:10.1093/cercor/bhv208
43. Beck, D. M. & Kastner, S. Top-down and bottom-up mechanisms in biasing competition
in the human brain. *Vision Res.* **49,** 1154–1165 (2009).
44. Nakayama, K. & Mackeben, M. Sustained and transient components of focal visual at-
tention. *Vision Res.* **29,** 1631–1647 (1989).
45. Posner, M. I., Snyder, C. R. & Davidson, B. J. Attention and the detection of signals. *J
Exp Psychol* **109,** 160–174 (1980).
46. Buschman, T. J. & Miller, E. K. Serial, Covert Shifts of Attention during Visual Search
Are Reflected by the Frontal Eye Fields and Correlated with Population Oscillations.
Neuron **63,** 386–396 (2009).
47. Corbetta, M., Patel, G. & Shulman, G. L. The reorienting system of the human brain: from
environment to theory of mind. *Neuron* **58,** 306–324 (2008).

48. Peelen, M. V., Heslenfeld, D. J. & Theeuwes, J. Endogenous and exogenous attention
shifts are mediated by the same large-scale neural network. *Neuroimage* **22,** 822–830
(2004).
49. Kincade, J. M., Abrams, R. A., Astafiev, S. V., Shulman, G. L. & Corbetta, M. An event-
related functional magnetic resonance imaging study of voluntary and stimulus-driven
orienting of attention. *Journal of Neuroscience* **25,** 4593–4604 (2005).
50. Mayer, A. R., Dorflinger, J. M., Rao, S. M. & Seidenberg, M. Neural networks underlying
endogenous and exogenous visual–spatial orienting. *Neuroimage* **23,** 534–541 (2004).
51. Carrasco, M. & McElree, B. Covert attention accelerates the rate of visual information
processing. *PNAS* **98,** 5363–5367 (2001).
52. Reed, A. V. Speed-Accuracy Trade-Off in Recognition Memory. *Science* **181,** 574–576
(1973).
53. Wickelgren, W. A. Speed-accuracy tradeoff and information processing dynamics. *Acta
Psychologica* **41,** 67–85 (1977).
54. Correa, Á., Triviño, M., Pérez-Dueñas, C., Acosta, A. & Lupiáñez, J. Temporal prepara-
tion, response inhibition and impulsivity. *Brain and Cognition* **73,** 222–228 (2010).
55. Yashar, A. & Lamy, D. Refining the dual-stage account of intertrial feature priming: does
motor response or response feature matter? *Atten Percept Psychophys* **73,** 2160–2167
(2011).
56. Herath, P., Klingberg, T., Young, J., Amunts, K. & Roland, P. E. Neural Correlates of
Dual Task Interference Can be Dissociated from Those of Divided Attention: an fMRI
Study | Cerebral Cortex | Oxford Academic. *Cereb. Cortex* **11,** 796–805 (2001).
57. Culham, J. C., Cavanagh, P. & Kanwisher, N. G. Attention Response Functions: Char-
acterizing Brain Areas Using fMRI Activation during Parametric Variations of Attentional
Load. *Neuron* **32,** 737–745 (2001).
58. Beauchamp, M. S., Petit, L., Ellmore, T. M., Ingeholm, J. & Haxby, J. V. A parametric
fMRI study of overt and covert shifts of visuospatial attention. *Neuroimage* **14,** 310–321
(2001).
59. Carrasco, M., Evert, D. L., Chang, I. & Katz, S. M. The eccentricity effect: target eccen-
tricity affects performance on conjunction searches. *Perception & Psychophysics* **57,**
1241–1261 (1995).
60. Rosen, A. C. *et al.* Neural basis of endogenous and exogenous spatial orienting. A func-
tional MRI study. *Journal of Cognitive Neuroscience* **11,** 135–152 (1999).
61. Esterman, M. *et al.* Voluntary and involuntary attention affect face discrimination differ-
ently. *Neuropsychologia* **46,** 1032–1040 (2008).
62. Natale, E., Marzi, C. A., Girelli, M., Pavone, E. F. & Pollmann, S. ERP and fMRI corre-
lates of endogenous and exogenous focusing of visual-spatial attention. *Eur. J. Neurosci.*
23, 2511–2521 (2006).
63. Kinchla, R. A., Chen, Z. & Evert, D. Precue effects in visual search: data or resource
limited? *Perception & Psychophysics* **57,** 441–450 (1995).
64. Ruff, C. C., Kristjánsson, Á. & Driver, J. Readout from iconic memory and selective spa-
tial attention involve similar neural processes. *Psychological Science* **18,** 901–909
(2007).
65. Hulme, O. J., Friston, K. F. & Zeki, S. Neural correlates of stimulus reportability. *Journal
of Cognitive Neuroscience* **21,** 1602–1610 (2009).
66. Nobre, A. C. *et al.* Orienting attention to locations in perceptual versus mental represen-
tations. *Journal of Cognitive Neuroscience* **16,** 363–373 (2004).
67. Sergent, C., Ruff, C. C., Barbot, A., Driver, J. & Rees, G. Top-down modulation of human
early visual cortex after stimulus offset supports successful postcued report. *Journal of
Cognitive Neuroscience* **23,** 1921–1934 (2011).
68. Thibault, L., van den Berg, R., Cavanagh, P. & Sergent, C. Retrospective Attention Gates
Discrete Conscious Access to Past Sensory Stimuli. *PLoS ONE* **11,** e0148504 (2016).
69. Sergent, C. *et al.* Cueing Attention after the Stimulus Is Gone Can Retrospectively Trig-
ger Conscious Perception. *Current Biology* **23,** 150–155 (2013).
70. Gobell, J. & Carrasco, M. Attention alters the appearance of spatial frequency and gap
size. *Psychological Science* **16,** 644–651 (2005).

71. Anton-Erxleben, K., Henrich, C. & Treue, S. Attention changes perceived size of moving
visual patterns. *Journal of Vision* **7,** 5.1–9 (2007).
72. Fuller, S., Park, Y. & Carrasco, M. Cue contrast modulates the effects of exogenous
attention on appearance. *Vision Res.* **49,** 1825–1837 (2009).
73. Carrasco, M. & Yeshurun, Y. The contribution of covert attention to the set-size and ec-
centricity effects in visual search. *Journal of Experimental Psychology: Human Percep-*
tion and Performance **24,** 673–692 (1998).
74. Liu, T., Pestilli, F. & Carrasco, M. Transient attention enhances perceptual performance
and FMRI response in human visual cortex. *Neuron* **45,** 469–477 (2005).
75. Kim, Y. H. *et al.* The large-scale neural network for spatial attention displays multifunc-
tional overlap but differential asymmetry. *Neuroimage* **9,** 269–277 (1999).
76. Meyer, K. N., Du, F., Parks, E. & Hopfinger, J. B. Exogenous vs. endogenous attention:
Shifting the balance of fronto-parietal activity. *Neuropsychologia* **111,** 307–316 (2018).
77. Dugué, L., Roberts, M. & Carrasco, M. Attention Reorients Periodically. *Current Biology*
26, 1595–1601 (2016).
78. Senoussi, M., Moreland, J. C., Busch, N. A. & Dugué, L. Attention explores space peri-
odically at the theta frequency. *Journal of Vision* **19,** 22–22 (2019).
79. Maunsell, J. H. R. & Cook, E. P. The role of attention in visual processing. *Philosophical*
Transactions of the Royal Society of London B: Biological Sciences **357,** 1063–1072
(2002).
80. Kastner, S. & Pinsk, M. A. Visual attention as a multilevel selection process. *Cognitive,*
Affective, & Behavioral Neuroscience **4,** 483–500 (2004).
81. Klein, B. P., Harvey, B. M. & Dumoulin, S. O. Attraction of Position Preference by Spatial
Attention throughout Human Visual Cortex. *Neuron* **84,** 227–237 (2014).
82. Kay, K. N., Weiner, K. S. & Grill-Spector, K. Attention Reduces Spatial Uncertainty in
Human Ventral Temporal Cortex. *Current Biology* **25,** 595–600 (2015).
83. Nachmias, J. Effect of Exposure Duration on Visual Contrast Sensitivity with Square-
Wave Gratings. *J. Opt. Soc. Am., JOSA* **57,** 421–427 (1967).
84. Carrasco, M., Penpeci-Talgar, C. & Eckstein, M. Spatial covert attention increases con-
trast sensitivity across the CSF: support for signal enhancement. *Vision Res.* **40,** 1203–
1215 (2000).
85. Luck, S. J. *et al.* Effects of spatial cuing on luminance detectability: psychophysical and
electrophysiological evidence for early selection. *Journal of Experimental Psychology:*
Human Perception and Performance **20,** 887–904 (1994).
86. Lu, Z. L. & Dosher, B. A. External noise distinguishes attention mechanisms. *Vision Res.*
38, 1183–1198 (1998).
87. Pestilli, F. & Carrasco, M. Attention enhances contrast sensitivity at cued and impairs it
at uncued locations. *Vision Res.* **45,** 1867–1875 (2005).
88. Montagna, B., Pestilli, F. & Carrasco, M. Attention trades off spatial acuity. *Vision Res.*
49, 735–745 (2009).
89. Yeshurun, Y. & Rashal, E. Precueing attention to the target location diminishes crowding
and reduces the critical distance. *Journal of Vision* **10,** 16–16 (2010).
90. Senoussi, M., Moreland, J. C., Busch, N. A. & Dugué, L. **Attention explores space**
periodically at the theta frequency. *Journal of Vision* **19,** 1–17 (2019).
91. Dugué, L., Xue, A. M. & Carrasco, M. Distinct perceptual rhythms for feature and con-
junction searches. *Journal of Vision* **17,** 22–22 (2017).
92. Ling, S. & Carrasco, M. Sustained and transient covert attention enhance the signal via
different contrast response functions. *Vision Res.* **46,** 1210–1220 (2006).
93. Engel, S. A. *et al.* fMRI of human visual cortex. *Nature* **369,** 525 (1994).
94. Sereno, M. I. *et al.* Borders of multiple visual areas in humans revealed by functional
magnetic resonance imaging. *Science* **268,** 889–893 (1995).
95. Engel, S. A., Glover, G. H. & Wandell, B. A. Retinotopic organization in human visual
cortex and the spatial precision of functional MRI. *Cereb. Cortex* **7,** 181–192 (1997).
96. Larsson, J. & Heeger, D. J. Two retinotopic visual areas in human lateral occipital cortex.
Journal of Neuroscience **26,** 13128–13142 (2006).

97.   Ress, D., Backus, B. T. & Heeger, D. J. Activity in primary visual cortex predicts perfor-
mance in a visual detection task. *Nat. Neurosci.* **3,** 940–945 (2000).
98.   Giesbrecht, B., Weissman, D. H., Woldorff, M. G. & Mangun, G. R. Pre-target activity in
visual cortex predicts behavioral performance on spatial and feature attention tasks.
Brain Res. **1080,** 63–72 (2006).
99.   Liu, T., Larsson, J. & Carrasco, M. Feature-based attention modulates orientation-selec-
tive responses in human visual cortex. *Neuron* **55,** 313–323 (2007).
100. Pinheiro, J. C. & Bates, D. M. Unconstrained parametrizations for variance-covariance
matrices. *Stat Comput* **6,** 289–296 (1996).
101. Carrasco, M., Marie Giordano, A. & McElree, B. Temporal performance fields: visual and
attentional factors. *Vision Res.* **44,** 1351–1365 (2004).
102. Ling, S. & Carrasco, M. When sustained attention impairs perception. *Nat. Neurosci.* **9,**
1243–1245 (2006).
103. Liu, T., Abrams, J. & Carrasco, M. Voluntary attention enhances contrast appearance.
Psychological Science **20,** 354–362 (2009).
104. Harvey, B. M. & Dumoulin, S. O. The relationship between cortical magnification factor
and population receptive field size in human visual cortex: constancies in cortical archi-
tecture. *Journal of Neuroscience* **31,** 13604–13612 (2011).
105. Harvey, B. M. & Dumoulin, S. O. Visual motion transforms visual space representations
similarly throughout the human visual hierarchy. *Neuroimage* **127,** 173–185 (2016).
106. Michalareas, G. *et al.* Alpha-Beta and Gamma Rhythms Subserve Feedback and Feed-
forward Influences among Human Visual Cortical Areas. *Neuron* **89,** 384–397 (2016).
107. Yildirim, F., Carvalho, J. & Cornelissen, F. W. A second-order orientation-contrast stim-
ulus for population-receptive-field-based retinotopic mapping. *Neuroimage* **164,** 183–
193 (2018).
108. Fritsche, M., Lawrence, S. J. D. & de Lange, F. P. Temporal tuning of repetition suppres-
sion across the visual cortex. *J. Neurophysiol.* **123,** 224–233 (2020).
109. Boynton, G. M., Demb, J. B., Glover, G. H. & Heeger, D. J. Neuronal basis of contrast
discrimination. *Vision Res.* **39,** 257–269 (1999).
110. Brefczynski, J. A. & DeYoe, E. A. A physiological correlate of the 'spotlight' of visual
attention. *Nat. Neurosci.* **2,** 370–374 (1999).
111. Müller, N. G. & Kleinschmidt, A. Temporal Dynamics of the Attentional Spotlight: Neu-
ronal Correlates of Attentional Capture and Inhibition of Return in Early Visual Cortex. *J
Cogn Neurosci* **19,** 587–593 (2007).
112. Müller, N. G. & Ebeling, D. Attention-modulated activity in visual cortex—More than a
simple 'spotlight'. *Neuroimage* **40,** 818–827 (2008).
113. Heinen, K. *et al.* Concurrent TMS-fMRI reveals dynamic interhemispheric influences of
the right parietal cortex during exogenously cued visuospatial attention. *European Jour-
nal of Neuroscience* **33,** 991–1000 (2011).
114. Mulckhuyse, M., Belopolsky, A. V., Heslenfeld, D., Talsma, D. & Theeuwes, J. Distribu-
tion of attention modulates salience signals in early visual cortex. *PLoS ONE* **6,** e20379
(2011).
115. Kay, K. N., Winawer, J., Rokem, A., Mezer, A. & Wandell, B. A. A two-stage cascade
model of BOLD responses in human visual cortex. *PLoS Comput. Biol.* **9,** e1003079
(2013).
116. Chica, A. B. & Lupiáñez, J. Effects of endogenous and exogenous attention on visual
processing: an Inhibition of Return study. *Brain Res.* **1278,** 75–85 (2009).
117. Liu, T., Heeger, D. J. & Carrasco, M. Neural correlates of the visual vertical meridian
asymmetry. *Journal of Vision* **6,** 1294–1306 (2006).
118. Freeman, J., Heeger, D. J. & Merriam, E. P. Coarse-scale biases for spirals and orien-
tation in human visual cortex. *Journal of Neuroscience* **33,** 19695–19703 (2013).
119. Wang, H. X., Merriam, E. P., Freeman, J. & Heeger, D. J. Motion direction biases and
decoding in human visual cortex. *Journal of Neuroscience* **34,** 12601–12615 (2014).
120. Brouwer, G. J., Arnedo, V., Offen, S., Heeger, D. J. & Grant, A. C. Normalization in hu-
man somatosensory cortex. *J. Neurophysiol.* **114,** 2588–2599 (2015).

121. McMains, S. A. & Somers, D. C. Multiple spotlights of attentional selection in human
visual cortex. *Neuron* **42,** 677–686 (2004).
122. Bloem, I. M. & Ling, S. Normalization governs attentional modulation within human visual
cortex. *Nature Communications* **10,** 5660–10 (2019).
123. Poltoratski, S., Ling, S., McCormack, D. & Tong, F. Characterizing the effects of feature
salience and top-down attention in the early visual system. *J. Neurophysiol.* **118,** 564–
573 (2017).
124. Moerel, D., Ling, S. & Jehee, J. F. M. Perceptual learning increases orientation sampling
efficiency. *Journal of Vision* **16,** 36–36 (2016).
125. Mackey, W. E., Winawer, J. & Curtis, C. E. Visual field map clusters in human frontopa-
rietal cortex. *Elife* **6,** 2704 (2017).
126. Gardner, J. L., Merriam, E. P., Schluppeck, D. & Larsson, J. MGL: Visual psychophysics
stimuli and experimental design package. (2018). doi:10.5281/zenodo.1299497
127. Yaffa Yeshurun, M. C. Attention improves or impairs visual performance by enhancing
spatial resolution. *Nature* **396,** 72–75 (1998).
128. Müller, M. M. in *Cognitive Electrophysiology of Attention* 123–135 (Elsevier, 2014).
doi:10.1016/B978-0-12-398451-7.00010-5
129. Mackeben, M. & Nakayama, K. Express attentional shifts. *Vision Res.* **33,** 85–90 (1993).
130. Liu, T., Stevens, S. T. & Carrasco, M. Comparing the time course and efficacy of spatial
and feature-based attention. *Vision Res.* **47,** 108–113 (2007).
131. Seiss, E., Driver, J. & Eimer, M. Effects of attentional filtering demands on preparatory
ERPs elicited in a spatial cueing task. *Clin Neurophysiol* **120,** 1087–1095 (2009).
132. Phillips, W. A. On the distinction between sensory storage and short-term visual memory.
Perception & Psychophysics **16,** 283–290 (1974).
133. Sperling, G. & Melchner, M. J. The attention operating characteristic: examples from
visual search. *Science* **202,** 315–318 (1978).
134. Kinchla, R. A. The measurement of attention. *Attention and performance VIII* 213–238
(1980).
135. Ogawa, S., Lee, T. M., Kay, A. R. & Tank, D. W. Brain magnetic resonance imaging with
contrast dependent on blood oxygenation. *Proc. Natl. Acad. Sci. U.S.A.* **87,** 9868–9872
(1990).
136. Gardner, J. L., Merriam, E. P., Schluppeck, D., Besle, J. & Heeger, D. J. mrTools: Anal-
ysis and visualization package for functional magnetic resonance imaging data. (2018).
doi:10.5281/zenodo.1299483
137. Wandell, B. A., Dumoulin, S. O. & Brewer, A. A. Visual field maps in human cortex. *Neu-*
ron **56,** 366–383 (2007).
138. Kay, K., Rokem, A., Winawer, J., Dougherty, R. & Wandell, B. GLMdenoise: a fast, au-
tomated technique for denoising task-based fMRI data. *Front. Neurosci.* **7,** 1–15 (2013).
912