Systematic analysis of the *Myxococcus xanthus* developmental gene regulatory network supports

posttranslational regulation of FruA by C-signaling

Running title: C-signaling regulates FruA posttranslationally

Shreya Saha[1], Pintu Patra[2], Oleg Igoshin[2], Lee Kroos[1*]

[1]Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI

48824, USA.

[2]Department of Bioengineering, Rice University, Houston, TX 77030, USA.

For correspondence. [*]E-mail kroos@msu.edu; Tel. (+1) 517 355 9726; Fax (+1) 517 353 9334.

Keywords: gene regulatory networks, signal transduction, transcription factors, developmental genes,

bacterial spores, *Myxococcus xanthus*

Acknowledgements

We thank Monique Floer for advice about high-throughput qPCR and for use of the LightCycler® 480

System. We thank Montserrat Elias-Arnanz and Penelope Higgs for sharing strains. This work was

supported by the National Science Foundation (award MCB-1411272) and by salary support for L.K.

from Michigan State University AgBioResearch.

Systematic analysis of the *Myxococcus xanthus* developmental gene regulatory network supports

posttranslational regulation of FruA by C-signaling

Summary

Upon starvation *Myxococcus xanthus* undergoes multicellular development. Rod-shaped cells move

into mounds in which some cells differentiate into spores. Cells begin committing to sporulation at 24-

30 h poststarvation, but the mechanisms governing commitment are unknown. FruA and MrpC are

transcription factors that are necessary for commitment. They bind cooperatively to promoter regions

and activate developmental gene transcription, including that of the *dev* operon. Leading up to and

during the commitment period, *dev* mRNA increased in wild type, but not in a mutant defective in C-

signaling, a short-range signaling interaction between cells that is also necessary for commitment. The

C-signaling mutant exhibited ~20-fold less *dev* mRNA than wild type at 30 h poststarvation, despite a

similar level of MrpC and only twofold less FruA. Boosting the FruA level twofold in the C-signaling

mutant had little effect on the *dev* mRNA level, and *dev* mRNA was not less stable in the C-signaling

mutant. Neither did high cooperativity of MrpC and FruA binding upstream of the *dev* promoter

explain the data. Rather, our systematic experimental and computational analyses support a model in

which C-signaling activates FruA at least ninefold posttranslationally in order to commit a cell to spore

formation.

Keywords: gene regulatory networks, signal transduction, transcription factors, developmental genes,

bacterial spores, *Myxococcus xanthus*

Introduction

Differentiated cell types are a hallmark of multicellular organisms. Understanding how pluripotent cells become restricted to particular cell fates is a fascinating question and a fundamental challenge in biology. In general, the answer involves a complex interplay between signals and gene regulation. This is true both during development of multicellular eukaryotes (Davidson & Levine, 2008, Frum & Ralston, 2015, Drapek *et al.*, 2017) and during transitions in microbial communities that lead to different cell types (van Gestel *et al.*, 2015, Norman *et al.*, 2015, Bush *et al.*, 2015, Kroos, 2017). Bacterial cells in microbial communities adopt different fates as gene regulatory networks (GRNs) respond to a variety of signals, including some generated by other cells. Moreover, we now understand that microbial communities or microbiomes profoundly impact eukaryotic organisms, and vice versa (Barratt *et al.*, 2017, Jansson & Hofmockel, 2018). Yet the daunting complexity of microbiomes and multicellular eukaryotes impedes efforts to fully understand their interactions in molecular detail. By studying simpler model systems, paradigms can be discovered that can guide investigations of more complex interactions.

A relatively simple model system is provided by the bacterium *Myxococcus xanthus*, which undergoes starvation-induced multicellular development (Yang & Higgs, 2014). In response to starvation, cells generate intracellular and extracellular signals that regulate gene expression (Bretl & Kirby, 2016, Kroos, 2017). The rod-shaped cells alter their movements so that thousands form a mound. Within a mound, cells differentiate into ovoid spores that resist stress and remain dormant until nutrients reappear. The spore-filled mound is called a fruiting body. Other cells adopt a different fate and remain outside the fruiting body as peripheral rods (O'Connor & Zusman, 1991). A large proportion of the cells lyse during the developmental process (Lee *et al.*, 2012). What determines

whether a given cell in the population forms a spore, remains as a peripheral rod, or undergoes lysis?

M. xanthus provides an attractive model system to discover how signaling between cells affects a GRN

and determines cell fate. Here, we focus on a circuit that regulates commitment to sporulation.

In a recent study, cells committed to spore formation primarily between 24 and 30 h poststarvation

(PS), because addition of nutrients to the starving population prior to 24 h PS blocked subsequent

sporulation, addition at 24 h PS allowed a few spores to form subsequently, and addition at 30 h PS

allowed about tenfold more spores to form (Rajagopalan & Kroos, 2014). At the molecular level,

addition of nutrients before or during the commitment period caused rapid proteolysis of MrpC

(Rajagopalan & Kroos, 2014), a transcription factor required for fruiting body formation (Sun & Shi,

2001b, Sun & Shi, 2001a).

MrpC appears to directly regulate more than one hundred genes involved in development

(Robinson *et al.*, 2014), and one well-characterized MrpC target gene, *fruA* (Ueki & Inouye, 2003),

codes for another transcription factor required for fruiting body formation (Ogawa *et al.*, 1996). FruA

and MrpC bind cooperatively to the promoter regions of many genes, and appear to activate

transcription (Campbell *et al.*, 2015, Lee *et al.*, 2011, Mittal & Kroos, 2009a, Mittal & Kroos, 2009b,

Robinson et al., 2014, Son *et al.*, 2011). In particular, transcription of the *dev* operon appears to be

activated by cooperative binding of the two transcription factors at two sites located upstream of the

promoter (Campbell et al., 2015). Because mutations in three genes of the *dev* operon (*devTRS*)

strongly impair sporulation (Boysen *et al.*, 2002, Thony-Meyer & Kaiser, 1993, Viswanathan *et al.*,

2007a), the feed-forward loop involving MrpC and FruA regulation of the *dev* operon is an attractive

molecular mechanism to control spore formation (Fig. 1). Recent work revealed that products of the

dev operon act as a timer for sporulation (Rajagopalan & Kroos, 2017). DevTRS negatively autoregulate

expression of DevI, which inhibits sporulation if overproduced, and delays sporulation by about 6 h

when produced normally (Rajagopalan & Kroos, 2017, Rajagopalan *et al.*, 2015) (Fig. 1).

Expression of the *dev* operon and many other developmental genes depends on C-signaling (Kroos

& Kaiser, 1987), which has been proposed to activate FruA (Ellehauge *et al.*, 1998) and/or MrpC (Mittal

& Kroos, 2009a) (Fig. 1), although the mechanism of C-signal transduction remains a mystery. Null

mutations in the *csgA* gene block C-signaling and sporulation, but the mutants can be rescued by co-

development with *csgA⁺* cells which supply the C-signal (Shimkets *et al.*, 1983). C-signaling appears to

be a short-range signaling interaction that requires cells to move into alignment (Kim & Kaiser, 1990c,

Kim & Kaiser, 1990b, Kroos *et al.*, 1988), as they do during mound formation (Sager & Kaiser, 1993).

Two theories about the identity of the C-signal have emerged. One theory states that the C-signal is a

17-kDa fragment of CsgA produced by the specific proteolytic activity of PopC at the cell surface (Kim &

Kaiser, 1990a, Lobedanz & Sogaard-Andersen, 2003, Rolbetzki *et al.*, 2008). The other theory is that

diacylglycerols released from the inner membrane by cardiolipin phospholipase activity of intact CsgA

are the C-signal (Boynton & Shimkets, 2015). However, in neither case has the signal receptor been

identified, so our understanding of C-signaling is incomplete. Likewise, how C-signaling impacts

recipient cells is unknown.

One way that C-signaling has been proposed to affect recipient cells is to stimulate

autophosphorylation of a histidine protein kinase, which would then transfer the phosphate to FruA

(Ellehauge et al., 1998). This model was attractive because FruA is similar to response regulators of

two-component signal transduction systems (Ellehauge et al., 1998, Ogawa et al., 1996). Typically, a

response regulator is phosphorylated by a histidine protein kinase in response to a signal, thus

activating the response regulator to perform a function (Stock *et al.*, 2000). The effects of substitutions

at the predicted site of phosphorylation in FruA supported the model that FruA is activated by

phosphorylation on D59 (Ellehauge et al., 1998). However, a histidine protein kinase capable of

phosphorylating FruA has not been identified. Also, several observations suggest that FruA may not be

phosphorylated. Most notably, D59 of FruA is present in an atypical receiver domain that lacks a

conserved metal-binding residue normally required for phosphorylation to occur, and treatment of

FruA with small-molecule phosphodonors did not increase its DNA-binding activity (Mittal & Kroos,

2009a). The receiver domain of FruA was shown to be necessary for cooperative binding with MrpC to

DNA, so it was proposed that C-signaling may affect activity of MrpC and/or FruA (Mittal & Kroos,

2009a) (Fig. 1).

The regulation of MrpC has been reported to be complex, involving autoregulation,

phosphorylation, proteolytic processing, binding to a toxin protein, and stability (Sun & Shi, 2001b,

Nariya & Inouye, 2005, Nariya & Inouye, 2006, Nariya & Inouye, 2008, Schramm *et al.*, 2012,

Rajagopalan & Kroos, 2014, McLaughlin *et al.*, 2018). Also, since MrpC is similar to CRP family

transcription factors that bind cyclic nucleotides (Sun & Shi, 2001b), MrpC activity could be modulated

by nucleotide binding, so there are many ways in which C-signaling could affect MrpC activity (Mittal &

Kroos, 2009a).

Here, using synergistic experimental and computational approaches, we investigate the impact of

C-signaling on a circuit that regulates commitment to sporulation by focusing on the feed-forward loop

involving MrpC and FruA control of *dev* operon transcription (Fig. 1). We describe methods to

systematically and quantitatively study the developmental process. Using these methods we measure

the levels of GRN components in wild type and in mutants (e.g., a *csgA* mutant unable to produce C-

signal) during the period leading up to and including commitment to spore formation. We then

formulate a mathematical model for the steady-state concentration of *dev* mRNA and use the model to

computationally predict the magnitude of potential regulatory effects of C-signaling that would be

required to explain our data. By testing the predictions, some potential regulatory mechanisms are

ruled out and at least ninefold activation of FruA by C-signaling is supported.

Results

M. xanthus *development can be studied systematically*

We first established quantitative assays to analyze cellular and molecular changes during *M. xanthus*

development. To facilitate collection of sufficient cell numbers for counting, as well as for RNA and

protein measurements, development was induced by starvation under submerged culture conditions.

Cells adhere to the bottom of a plastic well or dish, and develop under a layer of buffer. Prior to cell

harvest, photos were taken to document phenotypic differences between strains. As expected, wild-

type strain DK1622 formed mounds by 18 h poststarvation (PS) and the mounds matured into

compact, darkened fruiting bodies at 36 to 48 h PS (Fig. 2). In contrast, *csgA* and *fruA* null mutants

failed to progress beyond forming loose aggregates. A *devI* null mutant was similar to wild type (WT),

whereas a *devS* null mutant formed mounds slowly and they failed to darken. Developing populations

were harvested at the times indicated in Figure 2 to measure cellular and molecular changes.

To quantify changes at the cellular level, we counted the total number of cells (after fixation and

dispersal, so that rod-shaped cells, spores, and cells in transition between the two were counted) and

the number of sonication-resistant spores in the developing populations. We also counted the number

of rod-shaped cells at the time when development was initiated by starvation (T_0). By subtracting the

number of sonication-resistant spores from the total cell number, we determined the number of

sonication-sensitive cells. About 30% of the wild-type cells present at T_0 remained as sonication-

sensitive cells at 18 h PS (Fig. S1A), consistent with the suggestion that the majority of cells lyse early

during development under submerged culture conditions, which was based on the decrease in the

total protein concentration of developing cultures (Rajagopalan & Kroos, 2014). The number of

sonication-sensitive cells continued to decline after 18 h PS, reaching ~4% of the T_0 number by 48 h PS

(Fig. S1A). Spores were first observed at 27 h PS and the number rose to ~1% of the T_0 number by 48 h

PS (Fig. S1B). The *devI* mutant was similar to WT, except spores were first observed 6 h earlier at 21 h

PS, as reported recently (Rajagopalan & Kroos, 2017). The *csgA*, *fruA*, and *devS* mutants failed to make

a detectable number of spores (at a detection limit of 0.04% of the T_0 number) and appeared to be

slightly delayed relative to WT and the *devI* mutant in terms of the declining number of sonication-

sensitive cells (Fig. S1 and Table S1). We conclude that at the cellular level during the time between 18

and 30 h PS (when we measured RNA and protein levels as described below), the developing

populations decline from 36 ± 4% to 16 ± 6% of the initial rod number and only 0.5 ± 0.2% (WT, *devI*) or

<0.04% (*csgA*, *fruA*, *devS*) of the cells form sonication-resistant spores (from which the RNAs and

proteins we measured would not be recovered based on control experiments). We stopped collecting

samples at 30 PS because thereafter the number of sonication-sensitive cells continues to decline and

the spore number continues to rise, making RNA and protein more difficult to recover quantitatively,

yet many cells are committed at 30 h PS to make spores by 36 h PS even if nutrients are added

(Rajagopalan & Kroos, 2014). Hence, we focused on changes at the molecular level between 18 and 30

h PS, the period leading up to and including the time that many cells commit to spore formation. We

acknowledge that the populations also contain cells destined to lyse or become peripheral rods, so the

magnitude of molecular changes required to commit a cell to the spore fate may be greater than

revealed by our population measurements. However, methods of rapidly separating all three cell types

from populations or of analyzing molecular changes in individual cells have not yet been reported.

To measure RNA levels of a large number of samples, we adapted methods described previously

(Rajagopalan & Kroos, 2014) to a higher-throughput robotic platform for RT-qPCR analysis.

Reproducibility of the analysis was tested among biological replicates and two types of technical

replicates as illustrated in Figure S2A, for each RNA to be measured, at 24 h PS, the midpoint of our

focal period. No normalization was done in this experiment. Each transcript number was derived from

a standard curve of genomic DNA subjected to qPCR. For each RNA, we found that the average

transcript number and the standard deviation for three cDNA technical replicates from a single RNA

sample, three RNA technical replicates from a single biological replicate, and three biological

replicates, was not significantly different (single factor ANOVA, α = 0.05) (Fig. S2B-S2E). These results

suggest that biological variation in RNA levels at 24 h PS is comparable to technical variation in

preparing RNA and cDNA. In subsequent experiments, we measured RNA for at least three biological

replicates and we did not perform RNA or cDNA technical replicates. We also note the high abundance

of the *mrpC* transcript (~10%) relative to 16S rRNA, and the lower relative abundance of the *fruA* (~1%)

and *dev* (~0.1%) transcripts.

We have typically used 16S rRNA as an internal standard for RT-qPCR analysis during *M. xanthus*

development (Rajagopalan & Kroos, 2014). The high abundance of *mrpC* transcript relative to 16S

rRNA at 24 h PS (Fig. S2B and S2E) raised the possibility that rRNA decreases relative to total RNA at 18

to 30 h PS. To test this possibility, we measured the 16S rRNA level per μg of total RNA from 18 to 30 h

PS. Figure S3A shows that the level does not change significantly (single factor ANOVA, α = 0.05),

validating 16S rRNA as an internal standard for subsequent experiments. We also found that the total

RNA yield per cell does not change significantly from 18 to 30 h PS (single factor ANOVA, α = 0.05) (Fig.

S3B), consistent with the finding that the 16S rRNA level does not change significantly, since the

majority of total RNA is rRNA.

To measure protein levels, a portion of each well-mixed developing population was immediately

added to sample buffer, boiled, and frozen for subsequent semi-quantitative immunoblot analysis

(Rajagopalan & Kroos, 2017). The rest of the population was used for cell counting and RNA analysis as

described above and in the Experimental Procedures.

Levels of MrpC and FruA fail to account for the low level of dev *mRNA in a* csgA *mutant*

By systematically quantifying protein and mRNA levels during the period leading up to and including

the time that cells commit to spore formation, we investigated whether the GRN shown in Figure 1

could account for observed changes over time in WT and in mutants. In particular, we were interested

in whether changes in the levels of MrpC and/or FruA proteins could account for the observed changes

in the level of *dev* mRNA, since MrpC and FruA bind cooperatively to the *dev* promoter region and

activate transcription (Campbell et al., 2015). In WT, we found that the MrpC level decreased about

1.5-fold on average from 18 to 30 h PS (Fig. 3A) and the FruA level rose about 1.5-fold on average (Fig.

3B), whereas the *dev* mRNA level rose about threefold on average (Fig. 4A). In each case, the fold-

change was small and the variation between biological replicates was large, so the result of a single

factor ANOVA (α = 0.05) for each time course did not support a significant difference. We reasoned

that cooperative binding of MrpC and FruA could easily account for the threefold rise on average in *dev*

mRNA. We also measured the levels of *mrpC* and *fruA* mRNA. The *mrpC* mRNA level changed very

little on average (Fig. 4B), but the *fruA* mRNA level decreased about twofold on average after 18 h PS

(Fig. 4C), in contrast to the 1.5-fold rise on average in the FruA protein level (Fig. 3B), suggesting weak

positive posttranscriptional regulation of the FruA level during the period of commitment to spore

formation.

To investigate how C-signaling affects the GRN shown in Figure 1, we measured protein and mRNA

levels in the *csgA* null mutant. In agreement with earlier studies suggesting that C-signaling activates

FruA (Ellehauge et al., 1998) and/or MrpC (Mittal & Kroos, 2009a), we found very little *dev* mRNA in

the *csgA* mutant (Fig. 4A). Notably, the large decrease in the level of *dev* mRNA in the *csgA* mutant

compared with WT could not be accounted for by a large decrease in the level of MrpC or FruA. The

MrpC level was elevated about 1.5-fold on average in the *csgA* mutant relative to WT at most time

points (Fig. 3A), but the differences were not statistically significant ($p > 0.05$ in Student's two-tailed *t*-

tests comparing mutant to WT at each time point). The FruA level was diminished in the *csgA* mutant

relative to WT, but only about twofold on average (Fig. 3B). The differences in the FruA level were

statistically significant ($p < 0.05$ in Student's two-tailed *t*-tests) at each time point except 21 h PS ($p =$

0.12), but alone the twofold lower levels of FruA in the *csgA* mutant fail to account for the very low

levels of *dev* mRNA.

The *mrpC* and *fruA* mRNA levels were diminished about twofold and 1.5-fold on average,

respectively, in the *csgA* mutant relative to WT (Fig. 4B and 4C), but at nearly all time points the

differences were not statistically significant ($p > 0.05$ in Student's two-tailed *t*-tests, except $p = 0.03$ at

27 h for *mrpC* mRNA). The small differences in the level of *fruA* mRNA in the *csgA* mutant relative to

WT are especially noteworthy, since they imply that C-signaling has little or no effect on MrpC activity.

The results of our *fruA* mRNA measurements agree with published reports using *fruA-lacZ* fusions

(Ellehauge et al., 1998, Srinivasan & Kroos, 2004). Furthermore, we found that *fruA* mRNA stability is

similar in the *csgA* mutant and in WT at 30 h PS (Fig. S4), indicating that the similar steady-state *fruA*

mRNA level we observed (Fig. 4C) reflects a similar rate of synthesis, rather than altered synthesis

compensated by altered stability. We conclude that C-signaling does not affect MrpC activity.

Therefore, the low level of *dev* mRNA in a *csgA* mutant (Fig. 4A) could be due to failure to activate FruA

or to *dev*-specific regulatory mechanisms.

To begin to characterize potential *dev*-specific regulatory mechanisms during the period leading up

to and including commitment to sporulation, we measured protein and mRNA levels in the *devS* and

devI null mutants. The MrpC and FruA levels were similar to WT (Fig. 3). The *dev* mRNA level ranged

from 20-fold higher in the *devS* mutant than in WT at 18 h PS, to 10-fold higher at 30 h PS (Fig. 4A),

consistent with negative autoregulation by DevS (and DevT and DevR) reported previously

(Rajagopalan & Kroos, 2017, Rajagopalan et al., 2015). Unexpectedly, the *dev* mRNA level in the *devI*

mutant was about threefold lower than in WT at 30 h PS (Fig. 4A), suggesting that DevI feeds back

positively on accumulation of *dev* mRNA, although the difference was not quite statistically significant

at the 95% confidence level (p = 0.06 in Student's two-tailed t-test). Other differences were that the

fruA mRNA levels in the *devI* and *devS* mutants were about twofold lower than in WT at 27 and 30 h PS

(Fig. 4C), and these were statistically significant (p < 0.05 in Student's two-tailed t-tests comparing

mutant to WT at each time point). Since the FruA levels in these mutants were similar to those in WT

(Fig. 3B), positive posttranscriptional regulation of FruA appeared to occur in the mutants, as well as in

WT.

We also measured the *dev* mRNA level in a *ladA* null mutant, because LadA was reported to be a

strong positive regulator of *dev* expression based on analysis of transcriptional *lacZ* fusions

(Viswanathan *et al.*, 2007b). Although the *dev* mRNA was on average twofold lower in the *ladA* mutant

than in WT at 18 h PS and the difference was statistically significant ($p < 0.05$ in Student's two-tailed t-

tests), significant differences were not observed at 24 or 30 h (Fig. S5A). These results show that LadA

does not control the *dev* transcript level during the period (24 to 30 h PS) when cells commit to spore

formation under our conditions. Differences in the methods of measuring expression and/or the

developmental conditions may account for the different conclusion reached here about the effect of

LadA on *dev* expression, as compared with the previous study (Viswanathan et al., 2007b). We note

that, as in the previous study, the *ladA* mutant was delayed for mound formation and darkening (Fig.

S6A). Formation of sonication-resistant spores was also delayed (Fig. S6B).

To complete our characterization of the GRN shown in Figure 1, we also measured protein and

mRNA levels in the *fruA* and *mrpC* null mutants. We did not collect samples of the *mrpC* mutant at as

many time points since we expected little or no expression of GRN components. As expected, neither

MrpC nor FruA were detected in the *mrpC* mutant (Fig. S7). In the *fruA* mutant, the MrpC level was

similar to WT and, as expected, FruA was not detected (Fig. 3). Also as expected, in the *fruA* mutant

the *fruA* mRNA was not detected, the *dev* mRNA level was very low, and the *mrpC* mRNA level was

similar to WT (Fig. 4). Since the *mrpC* mutant had an in-frame deletion of codons 74 to 229 (Sun & Shi,

2001b), we were able to design primers for RT-qPCR analysis that should detect the shorter *mrpC*

transcript. Surprisingly, the *mrpC* mutant exhibited an elevated level of *mrpC* transcript compared with

WT at 18 and 24 h PS (Fig. S8A). The result was surprising since expression of an *mrpC-lacZ* fusion had

been reported to be abolished in the *mrpC* mutant, which had led to the conclusion that MrpC

positively autoregulates (Sun & Shi, 2001b). We considered the possibility that the shorter transcript in

the *mrpC* mutant is more stable than the WT transcript, but the transcript half-lives after addition of

rifampicin did not differ significantly (Fig. S9). We conclude that MrpC negatively regulates the *mrpC*

transcript level. While this work was in progress, McLaughlin *et al*. reached the same conclusion

(McLaughlin et al., 2018). In all other respects, the *mrpC* mutant yielded expected results. The *fruA*

and *dev* transcripts were very low (Fig. S8B and S8C), consistent with the expectations that MrpC is

required to activate *fruA* transcription (Ueki & Inouye, 2003) and that MrpC and FruA are required to

activate *dev* transcription (Campbell et al., 2015, Ellehauge et al., 1998, Viswanathan et al., 2007b).

Also, the *mrpC* mutant failed to progress beyond forming loose aggregates (Fig. S10), appeared to be

slightly delayed relative to WT in terms of the declining number of sonication-sensitive cells (Fig. S11A),

and failed to make a detectable number of spores (at a detection limit of 0.04% of the T_0 number) (Fig.

S11B).

Taken together, our systematic, quantitative measurements of components of the GRN shown in

Figure 1 imply that failure to activate FruA and/or *dev*-specific regulatory mechanisms may account for

the low level of *dev* mRNA in a *csgA* mutant. Given the complex feedback architecture of *dev*

regulation (i.e., strong negative feedback by DevTRS and weak positive feedback by DevI at 30 h PS),

delineating the effects of C-signaling on the *dev* transcript level requires a mathematical modeling

approach.

Mathematical modeling suggests several mechanisms that could explain the low level of dev mRNA

in the* csgA *mutant

The observed small differences in the levels of MrpC and FruA in the *csgA* mutant relative to WT do not

account for the very low level of *dev* mRNA in the *csgA* mutant. To evaluate plausible mechanisms that

may explain these experimental findings, we quantitatively analyzed transcriptional regulation of *dev*

by formulating a mathematical model that expresses the *dev* mRNA concentration as a function of the

regulators MrpC, FruA, DevI, and DevS. MrpC and FruA bind cooperatively to the *dev* promoter region

and activate transcription (Campbell et al., 2015). Our results suggest that DevI is a weak positive

regulator and DevS is a strong negative regulator of *dev* transcription by 30 h PS (Fig. 4A).

Incorporating these effects into a transcriptional regulation model, we express the concentration of

dev mRNA as a product of three regulation functions (Π_{FM}, Π_I, Π_S) divided by the transcript

degradation rate δ_{dev} (see Experimental Procedures for detailed explanation):

314
$$[\text{mRNA}_{dev}] = \frac{1}{\delta_{dev}} \left(\alpha_{FM} \underbrace{\frac{\left(\frac{[\text{FruA}][\text{MrpC}]}{K_{FM}}\right)^a}{1+\left(\frac{[\text{FruA}][\text{MrpC}]}{K_{FM}}\right)^a}}_{\Pi_{FM}} \right) \left(\underbrace{1 + \alpha_I \frac{\left(\frac{[\text{DevI}]}{K_I}\right)^b}{1+\left(\frac{[\text{DevI}]}{K_I}\right)^b}}_{\Pi_I} \right) \left(\underbrace{\frac{1}{1+\left(\frac{[\text{DevS}]}{K_S}\right)^c}}_{\Pi_S} \right)$$

Here, we use a quasi-steady state approximation for the mRNA levels by taking advantage of the fact

that mRNA decay (with half-lives typically in minutes) is much faster than our experimental

measurement times (in hours). This allows us to assume a rapid equilibrium between the rate of *dev*

transcription and the decay of its mRNA, which leads to the above equation, in which

α_{FM}, α_I, δ_{dev}, a, b, c, K_{FM}, K_I and K_S are parameters characterizing promoter regulation. We assume

that these biochemical parameters are not a function of the genetic background and, therefore, in the

strains in which *dev* mRNA was measured (e.g., the *csgA* mutant), the concentration of *dev* mRNA is

determined by the concentrations of proteins (indicated by square brackets in the equation), more

specifically the concentrations of their transcriptionally active forms (in case there is a

posttranslational regulation). To estimate how the different regulation parameters (such as

transcription rate, degradation rate, cooperativity constant, etc.) affect the *dev* mRNA level, we first

constrain the model parameters by the experimental result shown in Figure 3B, $[\text{FruA}]_{WT}/$

$[\text{FruA}]_{csgA} \cong 2$, and search for parameters that can result in the observed 22-fold difference

in $[\text{mRNA}_{dev}]$ in WT relative to the *csgA* mutant at 30 h PS (Fig. 4A).

To estimate the contribution of autoregulation by Dev proteins to their own transcription (i.e., the

terms $\Pi_\text{I}, \Pi_\text{S}$) in WT and the *csgA* mutant, we employ the data from the *devI* and *devS* mutants (Fig.

4A). Specifically, we take the ratio of the *dev* mRNA level in WT to that in *devI* and *devS* mutants to

estimate the feedback regulation from DevI and DevS, respectively (see Experimental Procedures for

details). We find the contribution from DevI and DevS feedback regulation in WT to be $\Pi_\text{I,WT} = 2.9$

and $\Pi_\text{S,WT} = 0.091$, respectively. Using these values, we find the contribution from FruA and MrpC

regulation to be $\Pi_\text{FM,WT}/\delta_{dev,\text{WT}} = 11$. In the *csgA* mutant, since the *dev* mRNA level is very low, we

assume the DevI and DevS protein levels to be low. This gives the contribution of different regulation

functions as $\Pi_{\text{I},csgA} \approx 1$, $\Pi_{\text{S},csgA} \approx 1$, and $\Pi_{\text{FM},csgA}/\delta_{dev,csgA} = 0.13$. In summary, this analysis reveals

that the twofold reduction of FruA protein observed in the *csgA* mutant (Fig. 3B) leads to a change

of $(\Pi_\text{FM,WT}/\Pi_{\text{FM},csgA})(\delta_{dev,csgA}/\delta_{dev,\text{WT}}) \approx 84$-fold in the FruA- and MrpC-dependent transcript

regulation term. We reasoned that the observed 22-fold reduction in *dev* transcript in the *csgA* mutant

relative to WT at 30 h PS (Fig. 4A) could result from a reduction in the FruA- and MrpC-dependent

activation rate Π_FM and/or an increase in the transcript degradation rate δ_{dev}. In what follows we use

the mathematical model to predict the magnitude of these effects that would be necessary to explain

the observed 22-fold difference in $[\text{mRNA}_{dev}]$.

• **Hypothesis 1:** Increase in *dev* transcript degradation rate in the *csgA* mutant

First, we estimate the difference in *dev* transcript degradation rate necessary to explain the observed

difference in transcript level between WT and the *csgA* mutant. For this, we make two assumptions.

First, we assume that MrpC and FruA bind to the *dev* promoter region with a Hill cooperativity

coefficient $a = 2$ (i.e., the maximum for a single cooperative binding site). Second, we assume that the

observed twofold difference in FruA protein level results in a twofold difference in transcriptionally

active FruA. Under these assumptions, we vary the remaining unknown parameters to compute the

required fold difference in transcript degradation rate for different values of promoter saturation. Our

results plotted in Figure 5A show that at least a 20-fold difference in transcript degradation rate is

required to explain the transcript data. This experimentally testable prediction will be assessed in a

subsequent section. If the results are inconsistent with this prediction, we must conclude that at least

one of the two assumptions above is invalid, resulting in the following two alternative hypotheses: the

Hill coefficient of MrpC and FruA binding to the *dev* promoter region is much higher than $a = 2$ and/or

the amount of transcriptionally active FruA does not scale with the measured FruA protein level (e.g., if

csgA-dependent C-signaling is also involved in posttranslational activation of FruA).

•   **Hypothesis 2:** High cooperativity of MrpC and FruA binding to the *dev* promoter region

Next, we test if a higher binding cooperativity can explain the difference in *dev* transcript level

between WT and the *csgA* mutant. We compute the required cooperativity coefficient by assuming

the degradation rate does not change between the two strains. Our results plotted in Figure 5B show

that the minimum cooperativity coefficient required to explain the experimental results is six for low

promoter saturation. In biologically realistic conditions, where promoter saturation is higher; the

required cooperativity is even higher. Such a large cooperativity can only be explained if there is more

than one site in the promoter region where MrpC and FruA bind with high cooperativity. We know

that the *dev* promoter region has at least two MrpC and FruA cooperative binding sites; one is proximal

upstream, whereas the other is distal upstream (Campbell et al., 2015). The distal upstream binding

site appeared to boost *dev* promoter activity after 24 h PS, based on β-galactosidase activity from a

lacZ reporter. Hence, in a subsequent section, we study the impact of a distal site deletion on different

transcripts (*mrpC, fruA, dev*) and proteins (MrpC, FruA) to test if presence of the distal site contributes

to higher cooperativity. If the results are not consistent with the model predictions, we must conclude

that the fold difference in active FruA exceeds that observed for the total concentration of each

protein (i.e., *csgA*-dependent C-signaling is involved in posttranslational activation of FruA).

• **Hypothesis 3:** Posttranslational regulation of FruA activity

To assess the difference in active FruA level required to explain the observed difference in *dev*

transcript level, in the absence of other effects, we fix the cooperativity coefficient at $a = 2$ and

assume the transcript degradation rate to be unchanged between WT and the *csgA* mutant. We then

use our model to compute the fold difference in active FruA required to achieve a 22-fold reduction in

dev transcript in the *csgA* mutant relative to WT. Our results plotted in Figure 5C show that at least a

ninefold reduction in active FruA is needed in the *csgA* mutant. The reduction in active FruA in the

csgA mutant would presumably be due to the absence of C-signal-dependent posttranslational

activation of FruA, not due to the twofold lower level of FruA protein we observed in the *csgA* mutant

relative to WT (Fig. 3B). The reduction in active FruA may be considerably greater than ninefold if the

dev promoter region approaches saturation (e.g., 20-fold at 80% saturation in Fig. 5C). Also,

mathematical modeling of our data at each time point from 18 to 30 h PS yields a similar result (Fig.

S12), suggesting that in WT, FruA has already been activated by C-signaling at least ninefold by 18 h PS,

and perhaps as much as 30-fold if the *dev* promoter region approaches saturation (righthand panel in

Fig. S12).

Stability of the* dev *transcript is unchanged in a* csgA *mutant

To measure the *dev* transcript degradation rate in WT and the *csgA* mutant, we compared the *dev*

transcript levels after addition of rifampicin to block transcription at 30 h PS. The average half-life of

the *dev* transcript in three biological replicates was 11 ± 6 min in WT and 7 ± 1 min in the *csgA* mutant

(Fig. 6), which is not a statistically significant difference (p = 0.36 in a Student's two-tailed t-test). We

conclude that elevated turnover does not account for the low level of *dev* transcript in the *csgA*

mutant. These results allow us to rule out Hypothesis 1.

The distal upstream binding site for MrpC and FruA has little impact on the* dev *transcript level

In a previous study, weak cooperative binding of MrpC and FruA to a site located between positions -

254 and -229 upstream of the *dev* promoter appeared to boost β-galactosidase activity from a *lacZ*

transcriptional fusion about twofold between 24 and 30 h PS, but deletion of the distal upstream site

did not impair spore formation (Campbell et al., 2015). These findings suggested that the distal site

has a modest impact on *dev* transcription that is inconsequential for sporulation. However, β-

galactosidase activity from *lacZ* fused to *dev* promoter segments with different amounts of upstream

DNA and integrated ectopically may not accurately reflect the contribution of the distal site to the *dev*

transcript level. Therefore, we measured the *dev* transcript level in a mutant lacking the distal site (i.e.,

DNA between positions -254 and -228 was deleted from the *M. xanthus* chromosome). The level of

dev transcript in the distal site mutant was similar to WT measured in the same experiment, in this

case increasing about twofold from 18 to 30 h PS (Fig. 7). Likewise, the levels of *mrpC* and *fruA*

transcripts (Fig. S8) and the corresponding proteins (Fig. S7) were similar in the distal site mutant as

compared with WT (p > 0.05 in Student's two-tailed t-tests comparing mutant to WT at each time

point, except p = 0.02 for *fruA* transcript levels at 18 h, p = 0.03 for MrpC protein levels at 30 h, and p =

0.03 for FruA protein levels at 18 h). The distal site mutant formed mounds by 18 h PS, which matured

into compact, darkened fruiting bodies at later times, similar to WT (Fig. S10), and the percentages of

sonication-sensitive cells and sonication-resistant spores observed for the distal site mutant were

similar to WT (Fig. S11). We conclude that the distal site has little or no impact on the developmental

process. In particular, the distal site does not contribute to high cooperativity of MrpC and FruA

binding to the *dev* promoter region that could explain the higher level of *dev* transcript in WT than in

the *csgA* mutant. These results allow us to rule out Hypothesis 2.

Boosting the FruA level in the* csgA *mutant has no effect on the* dev *transcript level

Having ruled out the first two hypotheses, our modeling predicts that the only viable option to explain

the effect of the *csgA* null mutation on the *dev* transcript level is Hypothesis 3: at least a ninefold

reduction in active FruA is needed in the *csgA* mutant as compared with WT. Specifically, our model

showed that the low *dev* transcript level in the *csgA* mutant is not due to its twofold lower FruA level

(Fig. 3B), but rather due to a failure to activate FruA in the absence of C-signaling (Fig. 5C and S12). As

a result, the model predicts that in the *csgA* mutant most of the FruA remains inactive. To test this

prediction, we integrated *fruA* transcriptionally fused to a vanillate-inducible promoter ectopically in

the *csgA* mutant. Upon induction the *csgA* P$_{van}$-*fruA* strain accumulated a similar level of FruA as WT

(Fig. 8A), but the *dev* transcript level remained as low as in the *csgA* mutant (Fig. 8B). Hence, boosting

the FruA level in the *csgA* mutant had no effect on the *dev* transcript level, consistent with our

prediction and supporting the hypothesis that C-signaling activates FruA at least ninefold.

Additionally, we tested the previously proposed model that C-signal-dependent phosphorylation of

D59 activates FruA (Ellehauge et al., 1998). We created *fruA D59E*, an allele designed to produce FruA

D59E with a phosphomimetic substitution in its receiver domain. In some cases, such a substitution

causes a response regulator to be constitutively active (Domian *et al.*, 1997, Klose *et al.*, 1993, Lan &

Igo, 1998), including response regulators in the NarL/FixJ subfamily to which FruA belongs (Davlieva *et*

al., 2015, Galperin, 2010, Ellehauge et al., 1998). Therefore, if FruA is normally activated by

phosphorylation of D59 in response to C-signaling, expression of *fruA D59E* may restore the *dev*

transcript level in the *csgA* mutant. We integrated *fruA D59E* transcriptionally fused to a vanillate-

inducible promoter ectopically in the *csgA* mutant. Upon induction the *csgA* P$_{van}$-*fruA D59E* strain

accumulated FruA D59E to a similar level as FruA accumulated by WT (Fig. 8A), but the *dev* transcript

level in the *csgA* P$_{van}$-*fruA D59E* strain remained as low as in the *csgA* mutant (Fig. 8B). These results

suggest that D59 phosphorylation may not be the mechanism by which C-signaling activates FruA.

The *fruA* transcript (Fig. S13A), *mrpC* transcript (Fig. S13B), and MrpC protein (Fig. S14) levels were

similar in the *csgA* P$_{van}$-*fruA* and *csgA* P$_{van}$-*fruA D59E* strains as in WT and the *csgA* mutant. Induction

of the *csgA* P$_{van}$-*fruA* and *csgA* P$_{van}$-*fruA D59E* strains did not rescue development, which failed to

progress beyond forming loose aggregates (Fig. S15), and the strains failed to make a detectable

number of spores by 48 h PS (Table S1). The induced *csgA* P$_{van}$-*fruA* strain appeared to be slightly

delayed relative to WT in terms of the declining number of sonication-sensitive cells, like the *csgA*

mutant, whereas the induced *csgA* P$_{van}$-*fruA D59E* appeared to be defective in attachment, resulting in

recovery of less sonication-sensitive cells at 18 h PS (Fig. S16).

As controls, P$_{van}$-*fruA* and P$_{van}$-*fruA D59E* were integrated ectopically in the *fruA* mutant. Upon

induction the *fruA* P$_{van}$-*fruA* strain formed mounds by 18 h PS and the mounds matured into compact,

darkened fruiting bodies at later times, similar to WT without or with vanillate added (Fig. S17). The

induced *fruA* P$_{van}$-*fruA D59E* strain formed mounds by 24 h PS, which matured into compact, darkened

fruiting bodies, similar to WT. Also, the induced *fruA* P*van*-*fruA* and P*van*-*fruA D59E* strains exhibited a

similar number of sonication-resistant spores as WT at 36 h PS (data not shown). These results show

that induction of the *fruA* P*van*-*fruA* and *fruA* P*van*-*fruA D59E* strains rescued development, presumably

because C-signaling activated the FruA and FruA D59E produced, respectively.

Other FruA-dependent transcripts are present at very low levels in the* csgA *mutant, even when FruA

or FruA D59E is produced

In addition to the *dev* operon, four other genes or operons appear to be activated by cooperative

binding of FruA and MrpC in their promoter region (Lee et al., 2011, Mittal & Kroos, 2009a, Mittal &

Kroos, 2009b, Son et al., 2011). These genes, designated *fmg* for "Fru A- and Mrp-regulated gene", also

depend on C-signaling for expression, based on analysis of *lacZ* fusions (Kroos & Kaiser, 1987). We

measured the levels of *fmgA*, *fmgBC*, *fmgD*, and *fmgE* transcripts in the *csgA* mutant and the induced

csgA P*van*-*fruA* and *csgA* P*van*-*fruA D59E* strains during development. Like the *dev* transcript level (Fig.

8B), the *fmg* transcript levels were very low in the *csgA* mutant and in its derivatives induced to

produce FruA or FruA D59E (Fig. S18). These findings support the hypothesis that C-signaling activates

FruA and that it may do so by a mechanism other than D59 phosphorylation.

To investigate whether LadA regulates *fmg* genes, we measured *fmg* transcript levels in the *ladA*

null mutant. Like the *dev* transcript level (Fig. S5A), the *fmgA* and *fmgD* transcript levels were

significantly lower in the *ladA* mutant than in WT at 18 h PS ($p < 0.05$ in Student's two-tailed *t*-tests),

but not at 24 or 30 h (Fig. S5B and S5D). The *fmgBC* and *fmgE* transcript levels were significantly lower

in the *ladA* mutant at 18 and 24 h, and at all three times, respectively (Fig. S5C and S5E). Hence, LadA

does not control the *fmgA* and *fmgD* transcript levels during the period (24 to 30 h PS) when cells

commit to spore formation under our conditions, but LadA appears to positively regulate the *fmgBC*

transcript level twofold on average at 24 h and the *fmgE* transcript level about fivefold on average at

24 to 30 h.

Discussion

Our systematic, quantitative analysis of a key circuit in the GRN governing *M. xanthus* fruiting body

formation implicates posttranslational regulation of FruA by C-signaling as primarily responsible for *dev*

transcript accumulation during the period leading up to and including commitment to spore formation.

Mathematical modeling of the *dev* transcript level allowed us to predict the magnitude of potential

regulatory mechanisms. Experiments ruled out C-signal-dependent stabilization of *dev* mRNA or highly

cooperative binding of FruA and MrpC to two sites in the *dev* promoter region as the explanation for

the much higher *dev* transcript level in WT than in the *csgA* mutant. Although the FruA level was

twofold lower in the *csgA* mutant than in WT (Fig. 3B and 8A), boosting the FruA or FruA D59E level in

the *csgA* mutant had no effect on the *dev* transcript level (Fig. 8B) and did not rescue development

(Fig. S15). Taken together, our experimental and computational analyses provide evidence that C-

signaling activates FruA at least ninefold posttranslationally during *M. xanthus* development (Fig. 9),

and likely by a mechanism other than phosphorylation of D59. The activation of FruA may be

considerably greater than ninefold if the *dev* promoter region approaches saturation (Fig. 5C and S12).

Since efficient C-signaling requires cells to move into close proximity (Kim & Kaiser, 1990c, Kim &

Kaiser, 1990b, Kroos et al., 1988), we propose that activation of FruA by C-signaling acts as a

checkpoint for mound formation during the developmental process (Fig. 9).

Regulation of FruA by C-signaling

If activation of FruA by C-signaling acts as a checkpoint for mound formation, then active FruA should

be present at 18 h PS since mound formation is well underway (Fig. 2). In agreement, mathematical

modeling of our data using the assumptions of hypothesis 3 at each time point from 18 to 30 h PS

yields a similar result (Fig. S12). This analysis implies that FruA has already been activated by C-

signaling at least ninefold by 18 h PS, if the assumptions of hypothesis 3 apply. The assumption that

the distal site does not contribute to high cooperativity of MrpC and FruA binding to the *dev* promoter

region applies since the *dev* transcript level did not differ significantly in the distal site mutant as

compared with WT at 18 or 24 h PS (Fig. 7). We did not measure *dev* transcript stability at 18 to 27 h

PS, but at 30 h PS there was no significant difference between WT and the *csgA* mutant (Fig. 6).

Therefore, C-signaling may have already activated FruA at least ninefold by 18 h PS, and perhaps as

much as 30-fold if the *dev* promoter region approaches saturation (90% saturation in the righthand

panel of Fig. S12). We note that during the period from 18 to 30 h PS, the *dev* transcript level rises, but

the rise is due to positive autoregulation by DevI (Fig. 4A). Hence, active FruA may not be the limiting

factor for *dev* transcription during this period (i.e., the *dev* promoter region may indeed approach

saturation binding of active FruA and MrpC). The proximal upstream site in the *dev* promoter region,

which is crucial for transcriptional activation, exhibits a higher affinity for cooperative binding of FruA

and MrpC than the distal upstream site (Campbell et al., 2015) or several other sites (Robinson et al.,

2014, Son et al., 2011), perhaps conferring on *dev* transcription a relatively low threshold for active

FruA.

The mechanism of FruA activation by C-signaling is unknown. Since FruA is similar to response

regulators of two-component signal transduction systems, phosphorylation by a histidine protein

kinase was initially proposed to control FruA activity (Ellehauge et al., 1998, Ogawa et al., 1996). While

this potential mechanism of posttranslational control cannot be ruled out, a kinase capable of

phosphorylating FruA has not been identified despite considerable effort. Moreover, the atypical

receiver domain of FruA and the inability of small-molecule phosphodonors to increase its DNA-binding

activity suggest that FruA may not be phosphorylated (Mittal & Kroos, 2009a). In further support of

this possibility, we found that expression of FruA D59E, designed to mimic phosphorylation of the

receiver domain, did not appear to constitutively activate the protein in the absence of C-signaling (Fig.

8 and S15). Neither did the D59E substitution render FruA unable to be activated, since expression of

FruA D59E rescued development of the *fruA* mutant (Fig. S17), consistent a previous study (Ellehauge

et al., 1998). Hence, FruA may be an atypical response regulator that is activated by C-signaling *via* a

mechanism other than phosphorylation.

Several atypical response regulators have been shown to be active without phosphorylation and a

few are regulated by ligand binding (Bourret, 2010, Desai *et al.*, 2016). For example, the atypical

receiver domain of *Streptomyces venezuelae* JadR1 is bound by jadomycin B, causing JadR1 to

dissociate from DNA, and the acylated antibiotic undecylprodigiosin of *Streptomyces coelicolor* may

use a similar mechanism to modulate DNA-binding activity of the atypical response regulator RedZ

(Wang *et al.*, 2009). Conceivably, FruA activity could likewise be regulated by binding of *M. xanthus*

diacylglycerols, which have been implicated in C-signaling (Boynton & Shimkets, 2015). Alternatively,

FruA could be regulated by a posttranslational modification other than phosphorylation or by binding

to another protein (i.e., sequestration).

In addition to regulating FruA activity posttranslationally, C-signaling appears to regulate the FruA

level posttranscriptionally. The FruA level was reproducibly twofold lower in the *csgA* mutant than in

WT (Fig. 3B and 8A), but the *fruA* transcript level was not significantly different (Fig. 4C and S11A).

These results suggest that positive posttranscriptional regulation of the FruA level requires C-signaling.

C-signaling may increase synthesis (i.e., increase *fruA* mRNA accumulation slightly and also increase

translation of *fruA* mRNA) and/or decrease turnover of FruA. We did not investigate this further

because the FruA deficit in the *csgA* mutant could be overcome with P$_{van}$-*fruA* or P$_{van}$-*fruA D59E*, yet

there was very little effect on the *dev* transcript level (Fig. 8). This demonstrates that the activity of

FruA, rather than its level, primarily controls the level of *dev* transcript.

Regulation by Dev proteins

DevI inhibits sporulation if overexpressed, as in the *devS* mutant (Rajagopalan et al., 2015) (Fig. 2 and

S1). Deletion of *devI* or the entire *dev* operon allows spores to begin forming about 6 h earlier than

normal, but does not increase the final number of spores (Rajagopalan & Kroos, 2017) (Fig. S1). The

level of MrpC was about twofold higher on average in the *devI* mutant than in WT at 15 h PS, perhaps

accounting for the observed earlier sporulation, although the difference diminished at 18-24 h PS

(Rajagopalan & Kroos, 2017), as reported here (Fig. 3A). It was concluded that DevI may transiently

and weakly inhibit translation of *mrpC* transcripts during the period leading up to commitment,

delaying sporulation (Rajagopalan & Kroos, 2017). As noted above, DevI positively autoregulates,

causing a small rise in the *dev* transcript level by 30 h PS (Fig. 4A, 7, and 8B). Although the mechanism

of this feedback loop is unknown, one possibility is that DevI inhibits negative autoregulation by

DevTRS (Fig. 9).

In previous studies, mutations in *devT*, *devR*, or *devS* relieved negative autoregulation, resulting in

~10-fold higher *dev* transcript accumulation at 24 h PS (Rajagopalan & Kroos, 2017, Rajagopalan et al.,

2015). In this study, a *devS* mutant likewise accumulated ~10-fold more *dev* transcript than WT at 24-

30 h PS, and the difference was ~20-fold at 18 and 21 h PS (Fig. 4A), suggesting that negative

autoregulation mediated by DevS has a stronger effect leading up to the commitment period than

during commitment. Strong negative autoregulation may promote commitment to sporulation by

lowering the level of DevI, which would raise the MrpC level by relieving inhibition of translation of

mrpC transcripts (Rajagopalan & Kroos, 2017). Our data suggest that negative autoregulation by

DevTRS weakens during the commitment period, perhaps accounting for the observed small rise in the

dev transcript level (Fig. 4A, 7, and 8B). If the elevated *dev* transcript level is accompanied by a small

increase in the level of DevI, then DevI may inhibit translation of *mrpC* transcripts, causing the MrpC

level to decrease slightly by 30 h PS in WT (Fig. 3A). DevI is predicted to be a 40-residue polypeptide

(Rajagopalan et al., 2015) and currently no method has been devised to measure the DevI level. This is

a worthwhile goal of future research, as is understanding how cells overcome DevI-mediated inhibition

of sporulation (depicted in Fig. 9 as inhibition of cellular shape change).

In addition to regulating the timing of commitment to spore formation, Dev proteins appear to play

a role in maturation of spores. Mutations in *dev* genes strongly impair expression of the *exo* operon

(Licking *et al.*, 2000, Rajagopalan & Kroos, 2017), which encodes proteins that help form the

polysaccharide spore coat necessary to maintain cellular shape change and form mature spores (Muller

et al., 2012, Ueki & Inouye, 2005).

The role of MrpC

Our results add to a growing list of observations that indicate MrpC functions differently during *M.*

xanthus development than originally proposed. We found that MrpC negatively autoregulates

accumulation of *mrpC* mRNA about twofold at 18 and 24 h PS (Fig. S8A), and it does so at 18 h PS

without significantly altering transcript stability (Fig. S9). This contradicts an earlier study that

concluded MrpC positively autoregulates, based on finding that expression of an *mrpC-lacZ* fusion was

abolished in an *mrpC* mutant (Sun & Shi, 2001b). Recently, and in agreement with our result, it was

reported that MrpC is a negative autoregulator that competes with MrpB for binding to the *mrpC*

promoter region (McLaughlin et al., 2018). MrpB, likely when phosphorylated, binds to two sites

upstream of the *mrpC* promoter and activates transcription. MrpC binds to multiple sites upstream of

the *mrpC* promoter (Nariya & Inouye, 2006, McLaughlin et al., 2018), including two that overlap the

MrpB binding sites (McLaughlin et al., 2018). Purified MrpC competes with the MrpB DNA-binding

domain for binding to the overlapping sites, supporting a model in which MrpC negatively

autoregulates by directly competing with phosphorylated MrpB for binding to overlapping sites

(McLaughlin et al., 2018) (Fig. 9).

The role of MrpC in cellular lysis during development appears to be less prominent than originally

proposed. MrpC was reported to function as an antitoxin by binding to and inhibiting activity of the

MazF toxin protein, an mRNA interferase shown to be important for developmental programmed cell

death (Nariya & Inouye, 2008). However, the effect of a null mutation in *mazF* on developmental lysis

depends on the presence of a *pilQ1* mutation (Boynton *et al.*, 2013, Lee et al., 2012). In *pilQ*$^+$

backgrounds such as our WT strain DK1622, MazF is dispensable for lysis. Here, we found only a slight

delay of the *mrpC* mutant relative to WT in terms of the declining number of sonication-sensitive cells

at 18-48 h PS (Fig. S11A), comparable to other mutants (*csgA*, *fruA*, *devS*, *csgA* P$_{van}$-*fruA*, *csgA* P$_{van}$-*fruA*

D59E) that were unable to form spores (Fig. S1 and S15; Table S1). Under our conditions, MrpC

appears to play no special role in modulating the cell number during development.

Both the synthesis and the degradation of MrpC are regulated. Synthesis is regulated by

phosphorylated MrpB and MrpC acting positively and negatively, respectively, at the level of

transcription initiation as described above (McLaughlin et al., 2018) (Fig. 9). Degradation is regulated

by the complex Esp signal transduction system (Cho & Zusman, 1999, Higgs *et al.*, 2008, Schramm et

al., 2012), which presumably senses a signal and controls the activity of an unidentified protease

involved in MrpC turnover, thus ensuring that development proceeds at the appropriate pace (Fig. 9).

Interestingly, preliminary results suggest that the Esp system does not govern the proteolysis of MrpC

observed when nutrients are added at 18 h PS (Rajagopalan & Kroos, 2014) (Y. Hoang, R. Rajagopalan,

and L. Kroos; unpublished data). This implies that another system senses nutrients and degrades MrpC

to halt development (Fig. 9).

Combinatorial control by MrpC and FruA

Nutrient-regulated proteolysis of MrpC provides a checkpoint for starvation during the period leading

up to and including commitment to sporulation (Rajagopalan & Kroos, 2014) (Fig. 9). If activation of

FruA by C-signaling acts as a checkpoint for mound formation as we propose (Fig. 9), then

combinatorial control by MrpC and activated FruA could ensure that only starving cells in mounds

express genes that commit them to spore formation.

MrpC and FruA bind cooperatively to the promoter regions of five C-signal-dependent genes or

operons (*dev*, *fmgA*, *fmgBC*, *fmgD*, *fmgE*) (Lee et al., 2011, Mittal & Kroos, 2009a, Mittal & Kroos,

2009b, Son et al., 2011, Campbell et al., 2015). In each case, cooperative binding to a site located just

upstream of the promoter appears to activate transcription. Hence, MrpC and FruA form a type 1

coherent feed-forward loop with AND logic (Mangan & Alon, 2003). This type of loop is abundant in

GRNs and can serve as a sign-sensitive delay element (Mangan & Alon, 2003, Mangan *et al.*, 2003).

The sign sensitivity refers to a difference in the network response to stimuli in the "OFF to ON"

direction versus the "ON to OFF" direction. What this means for the feed-forward loop created by

MrpC, FruA, and their target genes is that target gene expression is delayed as MrpC accumulates,

awaiting FruA activated by C-signaling (i.e., the "OFF to ON" direction) (Fig. 9). As cells move into

mounds and engage in short-range C-signaling, activated FruA would bind cooperatively with MrpC,

stimulating transcription of target genes that eventually commit cells to spore formation (depicted in

Fig. 9 as cellular shape change). However, if nutrients reappear prior to commitment, MrpC is

degraded and transcription of target genes rapidly ceases, halting commitment to sporulation (i.e., the

"ON to OFF" direction). The number of target genes may be large since MrpC binds to the promoter

regions of hundreds of developmental genes based on ChIP-seq analysis, and in 13 of 15 cases

cooperative binding of MrpC and FruA was observed (Robinson et al., 2014).

In addition to the feed-forward loop involving cooperative binding of MrpC and FruA to a site

located just upstream of the promoter, the promoter regions of some genes have more complex

architectures that confer greater dependence on C-signaling for transcriptional activation. For

example, in the *fmgD* promoter region, binding of MrpC to an additional site that overlaps the

promoter and the FruA binding site appears to repress transcription, and it has been proposed that a

high level of active FruA produced by C-signaling is necessary to outcompete MrpC for binding and

result in transcriptional activation (Lee et al., 2011) (Fig. S19A). However, the *fmgD* transcript level in

WT did not increase significantly from 18 to 30 h PS (Fig. S5 and S18), suggesting that active FruA is not

the limiting factor for transcription during this period. In the *fmgE* promoter region, a distal upstream

site with higher affinity for cooperative binding of MrpC and FruA appears to act negatively by

competing for binding with the lower affinity site just upstream of the promoter (Son et al., 2011) (Fig.

S19B). The *fmgE* transcript level in WT did increase significantly from 18 to 30 h (Fig. S5 and S18), but

we cannot conclude that active FruA is the limiting factor since LadA also regulates the *fmgE* transcript

level during this period (Fig. S5). In addition to *fmgD* and *fmgE*, other genes depend more strongly on

C-signaling and are expressed later during development than *dev* (Kroos & Kaiser, 1987). Some of

these genes may require a higher level of active FruA than *dev* and *fmgD* in order to be transcribed. In

contrast to the *dev* promoter region, which may have a relatively low threshold for active FruA and

therefore approach saturation binding of active FruA and MrpC at 18 h PS (Fig. S12), we predict that

the promoter regions of genes essential for commitment to sporulation have more complex

architectures and a higher threshold for active FruA. According to this model, C-signal-dependent

activation of FruA continues after 18 h PS and the rising level of active FruA triggers commitment

beginning at 24 h PS. We speculate that genes governing cellular shape change are under

combinatorial control of MrpC and FruA (Fig. 9), and have a high threshold for active FruA.

Experimental Procedures

Bacterial strains, plasmids and primers

The strains, plasmids, and primers used in this study are listed in Table S2. *Escherichia coli* strain DH5α

was used for cloning. To construct pSS10, primers FruA-F-NdeI-Gibson and FruA-R-EcoRI-Gibson were

used to generate PCR products using chromosomal DNA from *M. xanthus* strain DK1622 as template.

The products were combined with NdeI-EcoRI-digested pMR3691 in a Gibson assembly reaction to

enzymatically join the overlapping DNA fragments (Gibson *et al.*, 2009). The cloned PCR product was

verified by DNA sequencing. To construct pSS9, pSS10 was subjected to site-directed mutagenesis

using the Quikchange kit (Stratagene) and primers D59E F and D59E R, and the DNA sequence of the

fruA D59E gene was verified. *M. xanthus* strains with P*van*-*fruA* and P*van*-*fruA D59E* integrated

ectopically were constructed by electroporation (Kashefi & Hartzell, 1995) of pSS10 and pSS9,

respectively, selection of transformants on CTT agar containing 15 µg/ml of tetracycline (Iniesta *et al.*,

2012), and verification by colony PCR using primers pMR3691 MCS G-F and pMR3691 MCS G-R.

***Growth and development of* M. xanthus**

Strains of *M. xanthus* were grown at 32°C in CTTYE liquid medium (1% Casitone, 0.2% yeast extract, 10

mMTris-HCl [pH 8.0], 1 mM KH_2PO_4-K_2HPO_4, 8 mM $MgSO_4$ [final pH 7.6]) with shaking at 350 rpm. CTT

agar (CTTYE lacking yeast extract and solidified with 1.5% agar) was used for growth on solid medium

and was supplemented with 40 µg/ml of kanamycin sulfate or 15 µg/ml of tetracycline as required.

Fruiting body development under submerged culture conditions was performed using MC7 (10 mM

morpholinepropanesulfonic acid [MOPS; pH 7.0], 1 mM $CaCl_2$) as the starvation buffer as described

previously (Rajagopalan & Kroos, 2014). Briefly, log-phase CTTYE cultures were centrifuged and cells

were resuspended in MC7 at a density of approximately 1,000 Klett units. A 100 µl sample (designated

T_0) was removed, glutaraldehyde (2% final concentration) was added to fix cells, and the sample was

stored at 4°C at least 24 h before total cells were quantified as described below. For each

developmental sample, 1.5 ml of the cell suspension plus 10.5 ml of MC7 was added to an 8.5-cm-

diameter plastic petri plate. Upon incubation at 32°C, cells adhere to the bottom of the plate and

undergo development. At the indicated times developing populations were photographed through a

microscope and collected as described below.

Microscopy

Images of fruiting bodies were obtained using a Leica Wild M8 microscope equipped with an Olympus

E-620 digital camera. In order to quantify cells in samples collected and dispersed as described below,

high resolution images were obtained with an Olympus BX51 microscope using a differential

interference contrast filter and a 40× objective lens, and equipped with an Olympus DP30BW digital

camera.

Sample collection

At the indicated times the submerged culture supernatant was replaced with 5 ml of fresh MC7

starvation buffer with or without inhibitors as required. Developing cells were scraped from the plate

bottom using a sterile cell scraper and the entire contents were collected in a 15-ml centrifuge tube.

Samples were mixed thoroughly by repeatedly (three times total) vortexing for 15 s followed by

pipetting up and down 15 times. For quantification of total cells, 100 μl of the mixture was removed,

glutaraldehyde (2% final concentration) was added to fix cells, and the sample was stored at 4°C for at

least 24 h before counting as described below. For measurement of sonication-resistant spores, 400 μl

of the mixture was removed and stored at -20°C. For immunoblot analysis, 100 μl of the mixture was

added to an equal volume of 2× sample buffer (0.125 M Tris-HCl [pH 6.8], 20% glycerol, 4% sodium

dodecyl sulfate [SDS], 0.2% bromophenol blue, 0.2 M dithiothreitol), boiled for 5 min, and stored at -

20°C. Immediately after collecting the three samples just described, the remaining 4.4 ml of the

developing population was mixed with 0.5 ml of RNase stop solution (5% phenol [pH < 7] in ethanol),

followed by rapid cooling in liquid nitrogen until almost frozen, centrifugation at 8,700 × *g* for 10 min at

4°C, removal of the supernatant, freezing of the cell pellet in liquid nitrogen, and storage at -80°C until

RNA extraction. Control experiments with a sample collected at 30 h PS indicated that the majority of

spores remain intact after boiling in 2× sample buffer or RNA extraction as described below (data not

shown), so the proteins and RNAs analyzed are from developing cells that have not yet formed spores.

Quantification of total cells and sonication-resistant spores

During development a small percentage of the rod-shaped cells transition to ovoid spores that become

sonication-resistant. The number of sonication-resistant spores in developmental samples was

quantified as described previously (Rajagopalan & Kroos, 2014). Briefly, each 400-µl sample collected

as described above was thawed and sonicated using a model 450 sonifier (Branson) at output setting 2

for 10-s intervals three times with cooling on ice in between. A 60 µl sample was removed and ovoid

spores were counted microscopically using 10 µl in a Neubauer counting chamber. A remaining

portion of the sample was used to determine total protein concentration as described below. The total

cell number, including rod-shaped cells, ovoid spores, and cells in transition between the two, was

determined using the glutaraldehyde-fixed samples collected as described above. Each sample was

thawed and mixed by vortexing and pipetting, then 10 or 20 µl was diluted with MC7 to 400 µl,

sonicated once for 10 s, and all cells were counted microscopically. The total cell number minus the

number of sonication-resistant spores was designated the number of sonication-sensitive cells

(consisting primarily of rod-shaped cells) and was expressed as a percentage of the total cell number in

the corresponding T_0 sample (consisting only of rod-shaped cells).

RNA extraction and analysis

RNA was extracted using the hot-phenol method and the RNA was digested with DNase I (Roche) as

described previously (Higgs et al., 2008). One μg of total RNA was subjected to cDNA synthesis using

Superscript III reverse transcriptase (InVitrogen) and random primers (Promega), according to the

instructions provided by the manufacturers. Control reactions were not subjected to cDNA synthesis.

One μl of cDNA at the appropriate dilution (as determined empirically) and 20 pmol of each primer

were subjected to qPCR in a 25 μl reaction using 2× reaction buffer (20 mM Tris-HCl [pH 8.3], 13 mM

$MgCl_2$, 100 mM KCl, 400 μM dNTPs, 4% DMSO, 2× SYBR Green I [Molecular Probes], 0.01% Tween 20,

0.01% NP40, and 0.01 μg/μl of Taq polymerase) as described previously (Bryant *et al.*, 2008). qPCR

was done in quadruplicate for each cDNA using a LightCycler® 480 System (Roche). A standard curve

was generated for each set of qPCRs using *M. xanthus* wild-type strain DK1622 genomic DNA and gene

expression was quantified using the relative standard curve method (user bulletin 2; Applied

Biosystems). 16S rRNA was used as the internal standard for each sample. Relative transcript levels

for mutants are the average of three biological replicates after each replicate was normalized to the

transcript level observed for one replicate of WT at 18 h PS in the same experiment. Transcript levels

for WT at other times PS were likewise normalized to that observed for WT at 18 h PS in the same

experiment. For WT at 18 h PS, the transcript levels of at least three biological replicates from

different experiments were normalized to their average, which was set as 1.

Immunoblot analysis

A semi-quantitative method of immunoblot analysis was devised to measure the relative levels of

MrpC and FruA in many samples collected in different experiments. Equal volumes (10 μl for

measurement of MrpC and 15 μl for measurement of FruA) of samples prepared for immunoblot

analysis as described above were subjected to SDS-PAGE and immunoblotting as described previously

(Rajagopalan & Kroos, 2014, Yoder-Himes & Kroos, 2006). Representative immunoblots for each strain

are shown in Figure S20. On each immunoblot, a sample of the wild-type strain DK1622 at 18 h PS

served as an internal control for normalization of signal intensities across immunoblots. Signals were

detected using a ChemiDoc MP imaging system (Bio-Rad), with exposure times short enough to ensure

signals were not saturated, and signal intensities were quantified using Image Lab 5.1 (Bio-Rad)

software. After normalization to the internal control, each signal intensity was divided by the total

protein concentration of a corresponding sample that had been sonicated for 10-s intervals three times

as described above. After removal of a sample for spore quantification, the remaining portion was

centrifuged at 10,000 × g for 1 min and the total protein concentration of the supernatant was

determined using a Bradford (Bradford, 1976) assay kit (Bio-Rad). The resulting values of normalized

signal intensity/total protein concentration were further normalized to the average value for all

biological replicates of WT at 18 h PS, which was set as 1.

Mathematical modeling

Activation of dev transcription by FruA and MrpC

FruA and MrpC bind cooperatively to the _dev_ promoter region and activate transcription (Campbell et

al., 2015). In agreement, no _dev_ mRNA was detected in either the _fruA_ mutant (Fig. 4A) or the _mrpC_

mutant (Fig. 7).   We represent the activation of _dev_ transcript by FruA and MrpC using a

phenomenological Hill's function,

785
$$\Pi_{FM} = \alpha_{FM} \left[\frac{\left(\frac{[\text{FruA}][\text{MrpC}]}{K_{FM}}\right)^{a}}{1 + \left(\frac{[\text{FruA}][\text{MrpC}]}{K_{FM}}\right)^{a}} \right]$$

where α_{FM} denotes the maximal *dev* transcription rate, K_{FM} is the half-saturation constant, and a

denotes the cooperativity of binding. We model the binding cooperativity using a single Hill exponent

for the product of the FruA and MrpC concentrations since one FruA and one MrpC bind cooperatively

to the proximal site upstream of the *dev* promoter, based on mobility shift assays (Campbell et al., 2015).

Note that this expression will give $\Pi_{FM} = 0$ when $[FruA] = 0$ or $[MrpC] = 0$ (i.e., we have neglected

any basal transcription rate as we did not detect *dev* mRNA in the *fruA* or *mrpC* mutant. The expression

in brackets can be thought as the promoter occupancy probability (P in the equation below), a

dimensional parameter telling what fraction of the promoters will be occupied by the transcription

factors for a given value of K_{FM}.

795
$$P = \frac{\left(\frac{[FruA][MrpC]}{K_{FM}}\right)^a}{1 + \left(\frac{[FruA][MrpC]}{K_{FM}}\right)^a}$$

Note that the sensitivity of this expression to changes in the concentrations of FruA and MrpC are

maximal when $P \sim 0$ and minimal near saturation when $P \sim 1$. In Figure 5 we assess how different

hypotheses about the role of C-signaling in *dev* regulation play out at different levels of K_{FM}. To facilitate

the biological interpretation of the findings, we plot these as a function of *dev* promoter saturation.

Feedback regulation by Dev proteins

The *dev* mRNA level is further regulated by Dev proteins DevI and DevS. Our finding that the *dev*

transcript level is lower in the *devI* mutant than in WT (Fig. 4A) indicates that DevI is a positive regulator

of *dev* mRNA accumulation. In contrast, the *dev* transcript level in the *devS* mutant is significantly higher

than in WT (Fig. 4A), indicating that DevS is a negative regulator of *dev* mRNA accumulation. Since the

exact mechanisms of regulation by DevI and DevS are unclear, we assume for simplicity that these

proteins regulate the *dev* transcript level through independent mechanisms. We model these regulation

functions as follows:

$$810 \qquad \Pi_{\mathrm{I}} = \left(1 + \alpha_{\mathrm{I}} \frac{\left(\frac{[\mathrm{DevI}]}{K_{\mathrm{I}}}\right)^{b}}{1 + \left(\frac{[\mathrm{DevI}]}{K_{\mathrm{I}}}\right)^{b}}\right), \qquad \Pi_{\mathrm{S}} = \left(\frac{1}{1 + \left(\frac{[\mathrm{DevS}]}{K_{\mathrm{S}}}\right)^{c}}\right)$$

Here, α_{I} is a dimensionless parameter characterizing the feedback strength (i.e., the fold-increase in

transcription of the *dev* operon due to DevI), K_{I} is the half-saturation constant, and *b* denotes effective

cooperativity (i.e., Hill coefficient) of DevI binding. Likewise, K_{S} is the half-saturation constant and *c*

denotes the effective cooperativity of DevS binding. Note that these functions are normalized so that

$\Pi_{\mathrm{I}} = 1$ for the *devI* mutant and $\Pi_{\mathrm{S}} = 1$ for the *devS* mutant (i.e., when $[\mathrm{DevI}] = 0$ or $[\mathrm{DevS}] = 0$).

We assume that regulation by the Dev proteins is independent of that by FruA and MrpC, and the

effects will be multiplicative:

$$818 \qquad [\mathrm{mRNA}_{dev}] = \frac{\alpha_{\mathrm{FM}}}{\delta_{dev}} \underbrace{\left(\frac{\left(\frac{[\mathrm{FruA}][\mathrm{MrpC}]}{K_{\mathrm{FM}}}\right)^{a}}{1 + \left(\frac{[\mathrm{FruA}][\mathrm{MrpC}]}{K_{\mathrm{FM}}}\right)^{a}}\right)}_{\Pi_{\mathrm{FM}}} \underbrace{\left(1 + \alpha_{\mathrm{I}} \frac{\left(\frac{[\mathrm{DevI}]}{K_{\mathrm{I}}}\right)^{b}}{1 + \left(\frac{[\mathrm{DevI}]}{K_{\mathrm{I}}}\right)^{b}}\right)}_{\Pi_{\mathrm{I}}} \underbrace{\left(\frac{1}{1 + \left(\frac{[\mathrm{DevS}]}{K_{\mathrm{S}}}\right)^{c}}\right)}_{\Pi_{\mathrm{S}}}$$

where, K_{FM}, K_{I}, and K_{S} are the saturation constants for regulation by [FruA][MrpC], [DevI], and [DevS],

respectively.

Numerical procedure to estimate unknown regulation parameters

To explain the difference in the *dev* mRNA level in the *csgA* mutant as compared with WT, in terms of

perturbation of potential regulatory mechanisms, we use a mathematical approach where we constrain

the FruA ratio ($[\mathrm{FruA}]_{\mathrm{WT}}/[\mathrm{FruA}]_{csgA} \cong 2$) and find the regulation parameters that can result in the

observed 22-fold difference in $[\text{mRNA}_{\text{dev}}]$. Specifically, we use the expression of *dev* transcript ratio

between WT and the *csgA* mutant below:

$$\frac{[\text{mRNA}_{dev}]_{\text{WT}}}{[\text{mRNA}_{dev}]_{csgA}}$$

$$= \frac{\delta_{dev,csgA}}{\delta_{dev,\text{WT}}} \frac{1 + \left(\frac{[\text{FruA}]_{csgA}[\text{MrpC}]_{csgA}}{\text{K}_{\text{FM}}}\right)^a}{1 + \left(\frac{[\text{FruA}]_{\text{WT}}[\text{MrpC}]_{\text{WT}}}{\text{K}_{\text{FM}}}\right)^a} \left(\frac{[\text{FruA}]_{\text{WT}}[\text{MrpC}]_{\text{WT}}}{[\text{FruA}]_{csgA}[\text{MrpC}]_{csgA}}\right)^a \left(\frac{\Pi_{\text{I,WT}}\,\Pi_{\text{S,WT}}}{\Pi_{\text{I},csgA}\Pi_{\text{S},csgA}}\right)$$

$$\frac{[\text{mRNA}_{dev}]_{\text{WT}}}{[\text{mRNA}_{dev}]_{csgA}} = \frac{1}{\delta_R} \left(\frac{R^a + \left(\frac{P_{\text{WT}}}{1 - P_{\text{WT}}}\right)}{1 + \left(\frac{P_{\text{WT}}}{1 - P_{\text{WT}}}\right)}\right) \left(\frac{\Pi_{\text{I,WT}}\,\Pi_{\text{S,WT}}}{\Pi_{\text{I},csgA}\Pi_{\text{S},csgA}}\right)$$

where,

$$R = \frac{[\text{FruA}]_{\text{WT}}}{[\text{FruA}]_{csgA}} \frac{[\text{MrpC}]_{\text{WT}}}{[\text{MrpC}]_{csgA}}, \delta_R = \frac{\delta_{dev,\text{WT}}}{\delta_{dev,csgA}} \text{ and } P_{\text{WT}} = \frac{\left(\frac{[\text{FruA}]_{\text{WT}}[\text{MrpC}]_{\text{WT}}}{\text{K}_{\text{FM}}}\right)^a}{1 + \left(\frac{[\text{FruA}]_{\text{WT}}[\text{MrpC}]_{\text{WT}}}{\text{K}_{\text{FM}}}\right)^a} .$$

First, we estimate the contribution from Dev protein regulation terms $(\Pi_{\text{I}}, \Pi_{\text{S}})$ in determining the *dev*

transcript level in WT and the *csgA* mutant. Since we did not measure the Dev proteins explicitly in our

experiments, we estimate their contribution in regulating *dev* transcription in WT by comparing the

changes in transcript level in their absence (i.e., in the *devI* and *devS* mutants). Based on our transcript

data for WT, and the *devI* and *devS* mutants (Fig. 4A), we have the following relations between the

regulation functions; $[\text{mRNA}_{dev}]_{\text{WT}} = \delta_{dev,\text{WT}}^{-1}\Pi_{\text{FM,WT}}\Pi_{\text{I,WT}}\Pi_{\text{S,WT}} = 2.9$, $\delta_{dev,\text{WT}}^{-1}\Pi_{\text{FM,WT}}\Pi_{\text{S,WT}} =$

1 and $\delta_{dev,\text{WT}}^{-1}\Pi_{\text{FM,WT}}\Pi_{\text{I,WT}} = 32$. Using these relations, we obtain $\Pi_{\text{I,WT}} = 2.9$, $\Pi_{\text{S,WT}} = 0.091$. For

the *csgA* mutant, assuming regulation by Dev proteins is absent due to the low *dev* transcript level, we

have $\Pi_{\text{I},csgA} \approx 1$ and $\Pi_{\text{S},csgA} \approx 1$. With these estimates, the above expression for *dev* transcript ratio

has three unknown parameters $\delta_R, a, P_{\text{WT}}$.

Next, we determine the required fold change in degradation rate δ_R for different promoter

saturation probability P_{WT} values that explains the observed 22-fold difference in *dev* transcript. To

estimate this, we set the cooperativity constant (a) to 2 and take the fold change in FruA from the

experiments, while assuming MrpC is unchanged between WT and the *csgA* mutant. The result is

plotted in Fig. 5A. Then, we determine the required cooperativity a for different P_{WT} values with the

FruA fold change from the experiments and assuming no change in the degradation rate ($\delta_R = 1$). The

result is plotted in Fig. 5B. Finally, we compute the fold change in FruA with $\delta_R = 1$ and $a = 2$ for

different P_{WT} values. The result is shown in Fig. 5C.

RNA stability

At the indicated time the submerged culture supernatant was replaced with fresh MC7 starvation

buffer supplemented with 50 µg/ml of rifampicin to inhibit RNA synthesis. Samples were collected

immediately (designated t_0) and 8 and 16 min later for RNA extraction and analysis as described above,

except for each biological replicate the transcript levels after 8 and 16 min were normalized to the

transcript level at t_0, which was set as 1, and the natural log of the resulting values was plotted versus

minutes after rifampicin treatment and the slope of a linear fit of the data was used to compute the

mRNA half-life.

Induction of P*_{van}*-fruA

To induce expression of *fruA* fused to a vanillate-inducible promoter in *M. xanthus*, the CTTYE growth

medium was supplemented with 0.5 mM vanillate when the culture reached 50 Klett units. Growth

was continued until the culture reached 100 Klett units, then the culture was centrifuged and cells

were resuspended at a density of approximately 1,000 Klett units in MC7 supplemented with 0.5 mM

vanillate, followed by submerged culture development as described previously (Rajagopalan & Kroos,

2014).

Acknowledgements

We thank Monique Floer for advice about high-throughput qPCR and for use of the LightCycler® 480

System. We thank Montserrat Elias-Arnanz and Penelope Higgs for sharing strains. This work was

supported by the National Science Foundation (award MCB-1411272) and by salary support for L.K.

from Michigan State University AgBioResearch.

Author contributions

Conception or design of the study: LK, OI, SS, PP

Acquisition of the data: SS, PP

Analysis or interpretation of the data: SS, PP, LK, OI

Writing of the manuscript: LK, SS, PP, OI

Graphical abstract

Abbreviated summary

Starvation promotes MrpC accumulation, whereas nutrients favor proteolysis. MrpC activates

transcription of *fruA*, but FruA protein appears to be activated by short-range C-signaling in a cycle

leading to mound formation and lysis of some cells. Activated FruA* and MrpC are proposed to

cooperatively stimulate transcription of the *dev* operon and genes that commit starving rod-shaped

cells to form spores, while Dev proteins slow commitment, resulting in a spore-filled fruiting body

surrounded by peripheral rods.

References

Barratt, M. J., C. Lebrilla, H. Y. Shapiro & J. I. Gordon, (2017) The gut microbiota, food science, and human

893   nutrition: a timely marriage. *Cell Host Microbe* **22**: 134-141.

894   Bourret, R. B., (2010) Receiver domain structure and function in response regulator proteins. *Curr. Opin.*

895   *Microbiol.* **13**: 142-149.

896   Boynton, T. O., J. L. McMurry & L. J. Shimkets, (2013) Characterization of *Myxococcus xanthus* MazF and

897   implications for a new point of regulation. *Mol. Microbiol.* **87**: 1267-1276.

898   Boynton, T. O. & L. J. Shimkets, (2015) *Myxococcus* CsgA, *Drosophila* Sniffer and human HSD17B10 are

899   cardiolipin phospholipases. *Genes Dev.* **29**: 1903-1914.

900   Boysen, A., E. Ellehauge, B. Julien & L. Sogaard-Andersen, (2002) The DevT protein stimulates synthesis of FruA,

901   a signal transduction protein required for fruiting body morphogenesis in *Myxococcus xanthus*. *J.*

902   *Bacteriol.* **184**: 1540-1546.

903   Bradford, M., (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein

904   utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**: 248-254.

905   Bretl, D. J. & J. R. Kirby, (2016) Molecular mechanisms of signaling in *Myxococcus xanthus* development. *J. Mol.*

906   *Biol.* **428**: 3805-3830.

907   Bryant, G. O., V. Prabhu, M. Floer, X. Wang, D. Spagna, D. Schreiber & M. Ptashne, (2008) Activator control of

908   nucleosome occupancy in activation and repression of transcription. *PLoS Biol.* **6**: 2928-2939.

909   Bush, M. J., N. Tschowri, S. Schlimpert, K. Flardh & M. J. Buttner, (2015) c-di-GMP signalling and the regulation of

910   developmental transitions in streptomycetes. *Nat. Rev. Microbiol.* **13**: 749-760.

911   Campbell, A., P. Viswanathan, T. Barrett, B. Son, S. Saha & L. Kroos, (2015) Combinatorial regulation of the *dev*

912   operon by MrpC2 and FruA during *Myxococcus xanthus* development. *J. Bacteriol.* **197**: 240-251.

913   Cho, K. & D. R. Zusman, (1999) Sporulation timing in *Myxococcus xanthus* is contolled by the *espAB* locus. *Mol.*

914   *Microbiol.* **34**: 714-725.

Davidson, E. H. & M. S. Levine, (2008) Properties of developmental gene regulatory networks. *Proc. Natl. Acad.*

Sci. USA **105**: 20063-20066.

Davlieva, M., Y. Shi, P. G. Leonard, T. A. Johnson, M. R. Zianni, C. A. Arias, J. E. Ladbury & Y. Shamoo, (2015) A

variable DNA recognition site organization establishes the LiaR-mediated cell envelope stress response

of enterococci to daptomycin. *Nucleic Acids Res.* **43**: 4758-4773.

Desai, S. K., R. S. Winardhi, S. Periasamy, M. M. Dykas, Y. Jie & L. J. Kenney, (2016) The horizontally-acquired

response regulator SsrB drives a *Salmonella* lifestyle switch by relieving biofilm silencing. *eLife* **5**:

e10747.

Domian, I. J., K. C. Quon & L. Shapiro, (1997) Cell type-specific phosphorylation and proteolysis of a

transcriptional regulator controls the G1-to-S transition in a bacterial cell cycle. *Cell* **90**: 415-424.

Drapek, C., E. E. Sparks & P. N. Benfey, (2017) Uncovering gene regulatory networks controlling plant cell

differentiation. *Trends Genet.* **33**: 529-539.

Ellehauge, E., M. Norregaard-Madsen & L. Sogaard-Andersen, (1998) The FruA signal transduction protein

provides a checkpoint for the temporal co-ordination of intercellular signals in *Myxococcus xanthus*

development. *Mol. Microbiol.* **30**: 807-817.

Frum, T. & A. Ralston, (2015) Cell signaling and transcription factors regulating cell fate during formation of the

mouse blastocyst. *Trends Genet.* **31**: 402-410.

Galperin, M. Y., (2010) Diversity of structure and function of response regulator output domains. *Curr. Opin.*

Microbiol. **13**: 150-159.

Gibson, D. G., L. Young, R. Y. Chuang, J. C. Venter, C. A. Hutchison, 3rd & H. O. Smith, (2009) Enzymatic assembly

of DNA molecules up to several hundred kilobases. *Nat. Methods* **6**: 343-345.

Hagen, T. J. & L. J. Shimkets, (1990) Nucleotide sequence and transcriptional products of the *csg* locus of

Myxococcus xanthus. *J. Bacteriol.* **172**: 15-23.

Higgs, P. I., S. Jagadeesan, P. Mann & D. R. Zusman, (2008) EspA, an orphan hybrid histidine protein kinase,

regulates the timing of expression of key developmental proteins of *Myxococcus xanthus*. *J. Bacteriol.*

190: 4416-4426.

Iniesta, A. A., F. Garcia-Heras, J. Abellon-Ruiz, A. Gallego-Garcia & M. Elias-Arnanz, (2012) Two systems for

conditional gene expression in *Myxococcus xanthus* inducible by isopropyl-β-D-thiogalactopyranoside or

vanillate. *J. Bacteriol.* **194**: 5875-5885.

Jansson, J. K. & K. S. Hofmockel, (2018) The soil microbiome-from metagenomics to metaphenomics. *Curr. Opin.*

Microbiol. **43**: 162-168.

Kashefi, K. & P. Hartzell, (1995) Genetic supression and phenotypic masking of a *Myxococcus xanthux frzF*[-]

defect. *Molec. Microbiol.* **15**: 483-494.

Kim, S. K. & D. Kaiser, (1990a) C-factor: a cell-cell signaling protein required for fruiting body morphogenesis of

M. xanthus. *Cell* **61**: 19-26.

Kim, S. K. & D. Kaiser, (1990b) Cell alignment required in differentiation of *Myxococcus xanthus*. *Science* **249**:

926-928.

Kim, S. K. & D. Kaiser, (1990c) Cell motility is required for the transmission of C-factor, an intercellular signal that

coordinates fruiting body morphogenesis of *Myxococcus xanthus*. *Genes Dev.* **4**: 896-905.

Klose, K. E., D. S. Weiss & S. Kustu, (1993) Glutamate at the site of phosphorylation of nitrogen-regulatory

protein NTRC mimics aspartyl-phosphate and activates the protein. *J. Mol. Biol.* **232**: 67-78.

Kroos, L., (2017) Highly signal-responsive gene regulatory network governing *Myxococcus* development. *Trends*

Genet. **33**: 3-15.

Kroos, L., P. Hartzell, K. Stephens & D. Kaiser, (1988) A link between cell movement and gene expression argues

that motility is required for cell-cell signaling during fruiting body development. *Genes Dev.* **2**: 1677-

1685.

Kroos, L. & D. Kaiser, (1987) Expression of many developmentally regulated genes in *Myxococcus* depends on a

sequence of cell interactions. *Genes Dev.* **1**: 840-854.

Lan, C. Y. & M. M. Igo, (1998) Differential expression of the OmpF and OmpC porin proteins in *Escherichia coli* K-

12 depends upon the level of active OmpR. *J. Bacteriol.* **180**: 171-174.

Lee, B., C. Holkenbrink, A. Treuner-Lange & P. I. Higgs, (2012) *Myxococcus xanthus* developmental cell fate

production: heterogeneous accumulation of developmental regulatory proteins and reexamination of

the role of MazF in developmental lysis. *J. Bacteriol.* **194**: 3058-3068.

Lee, J., B. Son, P. Viswanathan, P. Luethy & L. Kroos, (2011) Combinatorial regulation of *fmgD* by MrpC2 and

FruA during *Myxococcus xanthus* development. *J. Bacteriol.* **193**: 1681-1689.

Licking, E., L. Gorski & D. Kaiser, (2000) A common step for changing cell shape in fruiting body and starvation-

independent sporulation of *Myxococcus xanthus*. *J. Bacteriol.* **182**: 3553-3558.

Lobedanz, S. & L. Sogaard-Andersen, (2003) Identification of the C-signal, a contact-dependent morphogen

coordinating multiple developmental responses in *Myxococcus xanthus*. *Genes Dev.* **17**: 2151-2161.

Mangan, S. & U. Alon, (2003) Structure and function of the feed-forward loop network motif. *Proc. Natl. Acad.*

Sci. USA **100**: 11980-11985.

Mangan, S., A. Zaslaver & U. Alon, (2003) The coherent feedforward loop serves as a sign-sensitive delay

element in transcription networks. *J. Mol. Biol.* **334**: 197-204.

McLaughlin, P. T., V. Bhardwaj, B. E. Feeley & P. I. Higgs, (2018) MrpC, a CRP/Fnr homolog, functions as a

negative autoregulator during the *Myxococcus xanthus* multicellular developmental program. *Mol.*

Microbiol. **109**: 245-261.

Mittal, S. & L. Kroos, (2009a) A combination of unusual transcription factors binds cooperatively to control

Myxococcus xanthus developmental gene expression. *Proc. Natl. Acad. Sci. USA* **106**: 1965-1970.

Mittal, S. & L. Kroos, (2009b) Combinatorial regulation by a novel arrangement of FruA and MrpC2 transcription

factors during *Myxococcus xanthus* development. *J. Bacteriol.* **191**: 2753-2763.

Muller, F. D., C. W. Schink, E. Hoiczyk, E. Cserti & P. I. Higgs, (2012) Spore formation in *Myxococcus xanthus* is

tied to cytoskeleton functions and polysaccharide spore coat deposition. *Mol. Microbiol.* **83**: 486-505.

Nariya, H. & M. Inouye, (2008) MazF, an mRNA interferase, mediates programmed cell death during

multicellular *Myxococcus* development. *Cell* **132**: 55-66.

Nariya, H. & S. Inouye, (2005) Identification of a protein Ser/Thr kinase cascade that regulates essential

transcriptional activators in *Myxococcus xanthus* development. *Mol. Microbiol.* **58**: 367-379.

Nariya, H. & S. Inouye, (2006) A protein Ser/Thr kinase cascade negatively regulates the DNA-binding activity of

MrpC, a smaller form of which may be necessary for the *Myxococcus xanthus* development. *Mol.*

Microbiol. **60**: 1205-1217.

Norman, T. M., N. D. Lord, J. Paulsson & R. Losick, (2015) Stochastic switching of cell fate in microbes. *Annu. Rev.*

Microbiol. **69**: 381-403.

O'Connor, K. A. & D. R. Zusman, (1991) Development in *Myxococcus xanthus* involves differentiation into two

cell types, peripheral rods and spores. *J. Bacteriol.* **173**: 3318-3333.

Ogawa, M., S. Fujitani, X. Mao, S. Inouye & T. Komano, (1996) FruA, a putative transcription factor essential for

the development of *Myxococcus xanthus*. *Mol. Microbiol.* **22**: 757-767.

Rajagopalan, R. & L. Kroos, (2014) Nutrient-regulated proteolysis of MrpC halts expression of genes important

for commitment to sporulation during *Myxococcus xanthus* development. *J. Bacteriol.* **196**: 2736-2747.

Rajagopalan, R. & L. Kroos, (2017) The *dev* operon regulates the timing of sporulation during *Myxococcus*

xanthus development. *J. Bacteriol.* **199**: e00788-00716.

Rajagopalan, R., S. Wielgoss, G. Lippert, G. J. Velicer & L. Kroos, (2015) *devI* is an evolutionarily young negative

regulator of *Myxococcus xanthus* development. *J Bacteriol* **197**: 1249-1262.

Robinson, M., B. Son & L. Kroos, (2014) Transcription factor MrpC binds to promoter regions of many

developmentally-regulated genes in *Myxococcus xanthus*. *BMC Genomics* **15**: 1123.

Rolbetzki, A., M. Ammon, V. Jakovljevic, A. Konovalova & L. Sogaard-Andersen, (2008) Regulated secretion of a

protease activates intercellular signaling during fruiting body formation in *M. xanthus*. *Dev. Cell* **15**: 627-

634.

Sager, B. & D. Kaiser, (1993) Two cell-density domains within the *Myxococcus xanthus* fruiting body. *Proc. Natl.*

Acad. Sci. USA **90**: 3690-3694.

Schramm, A., B. Lee & P. I. Higgs, (2012) Intra- and inter-protein phosphorylation between two hybrid histidine

kinases controls *Myxococcus xanthus* developmental progression. *J. Biol. Chem.* **287**: 25060–25072.

Shimkets, L. J., R. E. Gill & D. Kaiser, (1983) Developmental cell interactions in *Myxococcus xanthus* and the *spoC*

locus. *Proc. Natl. Acad. Sci. USA* **80**: 1406-1410.

Son, B., Y. Liu & L. Kroos, (2011) Combinatorial regulation by MrpC2 and FruA involves three sites in the *fmgE*

promoter region during *Myxococcus xanthus* development. *J. Bacteriol.* **193**: 2756-2766.

Srinivasan, D. & L. Kroos, (2004) Mutational analysis of the *fruA* promoter region demonstrates that C-box and 5-

base-pair elements are important for expression of an essential developmental gene of *Myxococcus*

xanthus. *J. Bacteriol.* **186**: 5961-5967.

Stock, A. M., V. L. Robinson & P. N. Goudreau, (2000) Two-component signal transduction. *Annu. Rev. Biochem.*

69: 183-215.

Sun, H. & W. Shi, (2001a) Analyses of *mrp* genes during *Myxococcus xanthus* development. *J. Bacteriol.* **183**:

6733-6739.

Sun, H. & W. Shi, (2001b) Genetic studies of *mrp*, a locus essential for cellular aggregation and sporulation of

Myxococcus xanthus. *J. Bacteriol.* **183**: 4786-4795.

Thony-Meyer, L. & D. Kaiser, (1993) *devRS*, an autoregulated and essential genetic locus for fruiting body

development in *Myxococcus xanthus*. *J. Bacteriol.* **175**: 7450-7462.

Ueki, T. & S. Inouye, (2003) Identification of an activator protein required for the induction of *fruA*, a gene

essential for fruiting body development in *Myxococcus xanthus*. *Proc. Natl. Acad. Sci. USA* **100**: 8782-

8787.

Ueki, T. & S. Inouye, (2005) Identification of a gene involved in polysaccharide export as a transcription target of

FruA, an essential factor for *Myxococcus xanthus* development. *J. Biol. Chem.* **280**: 32279-32284.

van Gestel, J., H. Vlamakis & R. Kolter, (2015) Division of labor in biofilms: the ecology of cell differentiation.

Microbiol. Spectr. **3**: MB-0002-2014.

Viswanathan, P., K. Murphy, B. Julien, A. G. Garza & L. Kroos, (2007a) Regulation of *dev*, an operon that includes

genes essential for *Myxococcus xanthus* development and CRISPR-associated genes and repeats. *J.*

Bacteriol. **189**: 3738-3750.

Viswanathan, P., T. Ueki, S. Inouye & L. Kroos, (2007b) Combinatorial regulation of genes essential for

Myxococcus xanthus development involves a response regulator and a LysR-type regulator. *Proc. Natl.*

Acad. Sci. USA **104**: 7969-7974.

Wang, L., X. Tian, J. Wang, H. Yang, K. Fan, G. Xu, K. Yang & H. Tan, (2009) Autoregulation of antibiotic

biosynthesis by binding of the end product to an atypical response regulator. *Proc Natl Acad Sci U S A*

106: 8617-8622.

Yang , Z. & P. Higgs, (2014) Myxobacteria: genomics, cellular and molecular biology. In. Norfolk, UK: Caister

Academic Press, pp.

Yoder-Himes, D. & L. Kroos, (2006) Regulation of the *Myxococcus xanthus* C-signal-dependent Ω4400 promoter

by the essential developmental protein FruA. *J. Bacteriol.* **188**: 5167-5176.

1050

1051

1052

Figure Legends

Fig. 1. Simplified model of the gene regulatory network governing formation of fruiting bodies.

Starvation increases the level of MrpC early in the process (Sun & Shi, 2001b, Sun & Shi, 2001a, Nariya

& Inouye, 2006). MrpC causes an increase in C-signal (Sun & Shi, 2001a), the product of *csgA* (Hagen &

Shimkets, 1990, Kim & Kaiser, 1990a). MrpC activates transcription of the gene for FruA (Ueki &

Inouye, 2003), and C-signal somehow enhances FruA (Ellehauge et al., 1998) and/or MrpC activity

(Mittal & Kroos, 2009a). MrpC and FruA bind cooperatively to the promoter region of the *dev* operon

and activate transcription (Campbell et al., 2015). The resulting DevTRS proteins negatively

autoregulate (Thony-Meyer & Kaiser, 1993, Viswanathan et al., 2007a, Rajagopalan & Kroos, 2017,

Rajagopalan et al., 2015). DevI delays spore formation within nascent fruiting bodies (Rajagopalan &

Kroos, 2017), but if overproduced, DevI inhibits sporulation (Rajagopalan et al., 2015), which is

promoted by MrpC (Sun & Shi, 2001b) and FruA (Ogawa et al., 1996) activity.

1065

Fig. 2. Development of *M. xanthus* strains. Wild-type DK1622 and its indicated mutant derivatives

were subjected to starvation under submerged culture conditions and images were obtained at the

indicated number of hours poststarvation (PS). DK1622 formed mounds by 18 h PS (an arrow points to

one). The *csgA* and *fruA* mutants failed to form mounds, the *devI* mutant was similar to DK1622, and

the *devS* mutant formed mounds later, by 24 h PS, but the mounds failed to darken at later times. Bar,

100 μm. Similar results were observed in at least three biological replicates.

1072

Fig. 3. Levels of MrpC and FruA during *M. xanthus* development. Wild-type DK1622 and its indicated

mutant derivatives were subjected to starvation under submerged culture conditions and samples

were collected at the indicated number of hours poststarvation (PS) for measurement of MrpC (A) and

FruA (B) by immunoblot. Graphs show the data points and average of at least three biological

replicates, relative to wild-type DK1622 at 18 h PS, and error bars show one standard deviation.

Asterisks indicate a significant difference ($p < 0.05$ in Student's two-tailed t-tests) from wild type at the

corresponding time PS.

1080

Fig. 4. Transcript levels during *M. xanthus* development. Wild-type DK1622 and its indicated mutant

derivatives were subjected to starvation under submerged culture conditions and samples were

collected at the indicated number of hours poststarvation (PS) for measurement of *dev* (A), *mrpC* (B),

and *fruA* (C) transcript levels by RT-qPCR. Graphs show the data points and average of at least three

biological replicates, relative to wild-type DK1622 at 18 h PS, and error bars show one standard

deviation. Asterisks indicate a significant difference ($p < 0.05$ in Student's two-tailed t-tests) from wild

type at the corresponding time PS.

1088

1089   **Fig. 5.** Mathematical modeling of different hypotheses to explain the low *dev* transcript level in a *csgA*

1090   mutant. Plots showing the required fold change in *dev* transcript degradation rate in the *csgA* mutant

1091   in comparison to wild type (A), cooperativity coefficient for MrpC and FruA binding to the *dev*

1092   promoter region (B), and reduction in FruA activity in the *csgA* mutant in comparison to wild type (C),

1093   to explain the experimental data for different values of promoter saturation.

1094

Fig. 6. *dev* transcript stability. Wild-type DK1622 and the *csgA* mutant were subjected to starvation

under submerged culture conditions for 30 h. The overlay was replaced with fresh starvation buffer

containing rifampicin (50 µg/ml) and samples were collected immediately (t_0) and at the times

indicated (t_x) for measurement of the *dev* transcript level by RT-qPCR. Transcript levels at t_x were

normalized to that at t_0 for each of three biological replicates and used to determine the transcript

half-life for each replicate. The average half-life and one standard deviation are reported in the text.

The graph shows the average $\ln(t_x/t_0)$ and one standard deviation for the three biological replicates of

wild type (black dashed line) and the *csgA* mutant (gray solid line).

1103

1104   **Fig. 7.** *dev* transcript levels. Wild-type DK1622 and its indicated mutant derivative were subjected to

1105   starvation under submerged culture conditions and samples were collected at the indicated number of

1106   hours poststarvation (PS) for measurement of *dev* transcript levels by RT-qPCR. Graphs show the data

1107   points and average of three biological replicates, relative to wild-type DK1622 at 18 h PS, and error

1108   bars show one standard deviation.

1109

Fig. 8. FruA protein and *dev* transcript levels. Wild-type DK1622 and its indicated mutant derivatives

were subjected to starvation under submerged culture conditions and samples were collected at the

indicated number of hours poststarvation (PS) for measurement of FruA levels by immunoblot (A) and

dev transcript levels by RT-qPCR (B). Graphs show the data points and average of three biological

replicates, relative to wild-type DK1622 at 18 h PS, and error bars show one standard deviation.

Asterisks indicate a significant difference ($p < 0.05$ in Student's two-tailed *t*-tests) from wild type at the

corresponding time PS.

1117

Fig. 9. Updated model of the gene regulatory network governing formation of fruiting bodies. Relative

to the simplified model shown in Figure 1 (see legend), this model also includes phosphorylated MrpB

(MrpB-P) which appears to activate transcription of *mrpC*, and negative autoregulation by MrpC which

appears to involve competition with MrpB-P for binding to overlapping sites in the *mrpC* promoter

region; proteolysis of MrpC, which is regulated by the Esp signal transduction system that normally

slows the developmental process and is regulated by nutrient addition that can halt development;

posttranslational activation of FruA to FruA* by C-signaling and promotion of mound formation by

FruA*, thus enhancing short-range C-signaling by bringing cells into proximity; the possibility that DevI

inhibits negative autoregulation by DevTRS; and speculation that the feed-forward loop involving MrpC

and FruA* not only controls transcription of the *dev* operon, but that of genes involved in cellular

shape change as well, committing cells to spore formation and resulting in spore-filled fruiting bodies.

This model deletes activation of MrpC by C-signaling, which was included as a possibility in Figure 1,

but was not supported by our data. See the text for details and references.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9