Respiration enhances TDP-43 toxicity, but TDP-43

retains some toxicity in the absence of respiration

By Sei-Kyoung Park, Sangeun Park, and Susan W. Liebman

Department of Pharmacology, University of Nevada, Reno NV, USA

Corresponding author: Susan W. Liebman

1664 N. Virginia St. Mail Stop 318, Reno, 89557

sliebman@unr.edu

Telephone: 775-682-7338

FAX: 775-784-1419

Abstract

The trans-activating response DNA-binding protein 43 (TDP-43) is a transcriptional repressor and splicing factor. TDP-43 is normally mostly in the nucleus, although it shuttles to the cytoplasm. Mutations in TDP-43 are one cause of familial amyotrophic lateral sclerosis (ALS). In neurons of these patients, TDP-43 forms cytoplasmic aggregates. In addition, wild-type TDP-43 is also frequently found in neuronal cytoplasmic aggregates in patients with neurodegenerative diseases not caused by TDP-43 mutations. TDP-43 expressed in yeast causes toxicity and forms cytoplasmic aggregates. This disease model has been validated because genetic modifiers of TDP-43 toxicity in yeast have led to the discovery that their conserved genes in humans are ALS genetic risk factors. While how TDP-43 is associated with toxicity is unknown, several studies find that TDP-43 alters mitochondrial function. We now report that TDP-43 is much more toxic when yeast are respiring than when grown on a carbon source where respiration is inhibited. However, respiration is not the unique target of TDP-43 toxicity because we found that TDP-43 retains some toxicity even in the absence of respiration. We found that H_2O_2 increases the toxicity of TDP-43, suggesting that the reactive oxygen species associated with respiration could likewise enhance the toxicity of TDP-43. In this case, the TDP-43 toxicity targets in the presence or absence of respiration could be identical, with the reactive oxygen species produced by respiration activating TDP-43 to become more toxic or making TDP-43 targets more vulnerable.

Glycolysis   **Respiration**

[*RHO*$^+$] yeast
functional
mitochondria

No TDP-43

↑ TDP-43

↑↑ TDP-43

No TDP-43 + H$_2$O$_2$

↑ TDP-43 + H$_2$O$_2$

[*rho*^{0}] yeast
non-functional
mitochondria

No TDP-43

↑ TDP-43

↑↑ TDP-43

No TDP-43 + H$_2$O$_2$

↑ TDP-43 + H$_2$O$_2$

Communication

The trans-activating response DNA-binding protein 43 (TDP-43) is a nucleic acid binding protein that functions as a transcriptional repressor, splicing factor and in translational regulation. TDP-43 is normally found mostly in the nucleus, although it shuttles between the nucleus and the cytoplasm. Mutations in *TARDBP*, the gene encoding TDP-43, are one cause of familial amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD). In neurons of these patients, TDP-43 is no longer found in the nucleus, but instead forms cytoplasmic aggregates. In addition, wild-type TDP-43 is also frequently found in cytoplasmic aggregates in neurons of patients with other neurodegenerative diseases or with ALS/FTD not caused by TDP-43 mutations[1; 2; 3; 4].

How TDP-43 aggregates are associated with toxicity is the subject of intense research. Evidence links TDP-43 toxicity with both inhibition of the ubiquitin proteasome system[5], and inhibition of lysosome and endosomal activity[6; 7]. Also, several studies find that TDP-43 alters mitochondrial function [8; 9; 10; 11; 12; 13; 14; 15]. Overexpression of TDP-43 in a number of model organisms causes neurodegeneration similar to that seen in ALS patients. In motor neurons of TDP-43 transgenic mice, mitochondria were found in cytoplasmic inclusions or in abnormal juxta-nuclear aggregates and were missing in motor axon termini[9; 10; 11]. In mammalian neuron-like cell culture, TDP-43 localized to mitochondria and caused mitophagy[12]. Likewise, in a mouse model, TDP-43 co-localization with motor neuron mitochondria was enhanced by TDP-43 ALS mutations and this co-localization was associated with inhibited mitochondrial function[13; 14]. Also, overexpressed TDP-43 in human cell culture binds to, and inhibits maturation of, mitochondrial RNA causing a phenotype similar to that seen in cells deficient in mitochondrial RNase P[15]. Finally, mutations in the mitochondrial intermembrane protein, CHCHD10 (yeast homolog MIX17) have recently been associated with sporadic and familial ALS and FDT [16; 17; 18; 19; 20; 21; 22] and functional CHCHD10 appears to help keep TDP-43 in the nucleus away from mitochondria[23].

There is also evidence that TDP-43 expression in yeast causes mitochondrion-dependent apoptosis and that TDP-43 toxicity requires respiratory capacity[8]. TDP-43 expressed in yeast causes toxicity and forms cytoplasmic aggregates[24]. Furthermore, conserved genetic modifiers of this toxicity have been validated in higher organisms and have led to the discovery of human ALS genetic risk factors[25]. This confirms the usefulness of the yeast model.

To examine the effect of respiration on TDP-43 toxicity we used the fact that yeast turns off respiration in dextrose media where it instead uses glycolysis to grow[26]. We found that TDP-43-GFP expressed with a *TET* promoter aggregated in yeast cells whether grown on dextrose, where respiration is inhibited; galactose, where respiration is not inhibited and both fermentation and respiration is present [26]; or glycerol or ethanol where only respiration and not fermentation is present. However, the *TET* controlled TDP-43 was only toxic on galactose, glycerol or ethanol where respiration is present. Also, the previously described TDP-43 aggregation and induction of cell elongation[5] was much more pronounced on media with respiration (galactose and glycerol) vs. media without respiration (dextrose) (Fig. 1ab). Despite this, there was no significant difference in the level of soluble TDP-43 found in boiled lysates of dextrose vs. galactose or glycerol grown cells (Fig. 1c). To test if the difference in sensitivity to TDP-43 was caused by the slower cell division rate in galactose vs. dextrose, we compared the toxicity in cells grown on dextrose at room temperature with cells grown on galactose at 30°C. Although cells grew at approximately the same rate under these two conditions, TDP-43 toxicity was pronounced on galactose but not dextrose plates (Supplementary Fig. 1).

One explanation could be that TDP-43 only inhibits respiration, which only affects growth when cells are respiring. Alternatively, respiration could be required for TDP-43 to be toxic, e.g. by modifying the TDP-43 protein. Indeed, oxidative stress has been shown to increase disulfide cross-linking, acetylation, and aggregation of TDP-43 in mammalian cells[27]. Furthermore, oxidative stress has been proposed to

promote acetylation of TDP-43, causing reduced binding of TDP-43 to RNA and increased levels of cytoplasmic, phosphorylated, aggregated TDP-43[28].

However, we eliminated both of the above hypotheses because we were able to detect TDP-43 toxicity in the absence of respiration on dextrose by expressing TDP-43-YFP with the strong *CUP1* promoter [29; 30]. To do this, we obtained an array of integrants of a plasmid expressing *CUP1*-TDP43-YFP in [*PIN*+] 74D-694. As expected, due to differences in the number of tandem plasmid repeats present [31], different integrants showed different levels of toxicity (high, medium and low). The more toxic integrants clearly showed toxicity on dextrose (Supplementary Fig. 2). Each integrant was cured of [*PIN*+] by growth on guanidine HCl [32]. Then, since yeast lacking functional mitochondria are viable we made all the [*PIN*+] and [*pin*-] [*RHO*+] integrants containing functional mitochondria into [*rho*^{0}] cells lacking functional mitochondrial. We then compared the level of toxicity caused by overexpressed TDP-43 in these isogenic [*RHO*+] and [*rho*^{0}] strains grown on dextrose (dex) where respiration is inhibited. We found that TDP-43 was toxic and retained the same level of expression in both [*RHO*+] and [*rho*^{0}] cells (Fig. 2abc). There was also no obvious difference in the appearance of TDP-43 aggregates (Fig. 2d). As we reported previously[5] TDP-43 is more toxic in the presence of the [*PIN*+] prion. This was clear when cells were grown on dextrose or galactose (Fig. 2a), but the effect of [*PIN*+] on TDP-43 toxicity in cells grown on glycerol was minor and not reproducible (not shown). Since the [*rho*^{0}] cells can never respire and the [*RHO*+] cells do not respire on dextrose, this clearly establishes that there is a TDP-43 toxicity target in yeast distinct from respiration and that respiration is not required for this TDP-43 toxicity.

To test the idea that increased levels of reactive oxygen species associated with respiration could be the cause of the observed increase in TDP-43 toxicity in the presence of respiration, we looked at the effects of H_2O_2, a source of reactive oxygen species, and N-Acetyl L-cysteine (NAC) an antioxidant, on TDP-43 toxicity. Using an integrant with low TDP-43 toxicity, we saw no toxicity of cells grown for 24 h with either 2mM H_2O_2 without TDP-43 expression (Fig. 3a left) or with expression of TDP-43 with 250 μM

CuSO$_4$ in dextrose medium without H$_2$O$_2$ (Fig. 3a right rows marked 0). However, the combination of 24

h of 2mM H$_2$O$_2$ and induction of TDP-43 with 250 µM CuSO$_4$ did cause toxicity (Fig. 3a right row marked

2). This was true in both [*RHO*$^+$] or [*rho*^{0}] cells. We also observed a dramatic increase in the size of TDP-

43 aggregates in cells treated with H$_2$O$_2$ (Fig. 3b), and this was not due to an increase in the cellular level

of TDP-43 (Fig. 3c). Possibly, this was caused by free radical oxygen stress, which has been shown to

cause an increase in protein aggregation[33; 34]. Also, we saw reduced toxicity of cells in the presence of

the antioxidant NAC and this was quite dramatic in the presence of TDP-43 (Fig. 4ab). However, the level

of TDP-43 in cells was dramatically reduced in the presence of NAC (Figs. 3c and 4c). If NAC directly

reduced expression of TDP-43 that would explain the reduced toxicity. Likewise if NAC inhibited

aggregation of TDP-43, the unaggregated TDP-43 would be expected to be rapidly degraded again

explaining reduced toxicity.

The prevalence of TDP-43 aggregation in patients with a variety of neurodegenerative diseases makes it

critical to understand how this is associated with toxicity. Our data shows that toxicity is enhanced in the

presence of respiration, but that TDP-43 remains toxic even in the absence of respiration. This is

consistent with the hypothesis that TDP-43 targets the same cellular components in the presence or

absence of respiration, and that the reactive oxygen species produced by respiration either activates

TDP-43 to become more toxic or makes the same TDP-43 targets more vulnerable.

Figure Legends

Fig. 1. TDP-43 overexpression causes more toxicity and cell elongation in cells grown in media that allows respiration vs. cells grown in the absence of respiration. (a) Yeast cells (L2910 [*pin*⁻] [*RHO*⁺] 74D-694) transformed with the doxycycline-repressible *CEN TET-TDP-43-YFP TRP1* (p2223), or vector control (p1576) plasmid were obtained on plasmid selective dextrose plates with 10 µg/ml of doxycycline (dox) to inhibit expression of TDP-43-YFP. Normalized suspensions of cells were serially diluted in water (1/10) and were spotted on plasmid selective 2% dextrose (dex), 2% galactose (gal), 2% glycerol (gly), and 2% ethanol (EtOH) plates without doxycycline (0) and with 10 µg/ml of doxycycline to respectively express or not express TDP-43-YFP. The plates were photographed after 4 (dextrose) or 6 (galactose, glycerol or ethanol) d of incubation at 30°C. (b) TDP-43 aggregates in galactose, glycerol and dextrose but causes more aggregation and cell elongation in galactose or glycerol vs. dextrose media. Cells from TDP-43-YFP inducing plates (↑TDP-43-YFP) were imaged with a fluorescence microscope at the same exposure and are shown at the same magnification. Left images show bright field with fluorescent, right images show just fluorescence. Control cells with no TDP-43-YFP expression (no TDP-43-YFP) are also shown. The increased aggregation and elongation on galactose and glycerol caused by TDP-43-YFP expression was very obvious. (c) Expression of *TET* controlled TDP-43-YFP is not enhanced on glycerol or galactose medium. The level of TDP-43-YFP was compared in L2910 transformants with doxycycline-repressible -*TDP-43-YFP* (p2223) grown in 2% dextrose (dex) vs. 2% glycerol (gly) or 2% raffinose + 2% galactose (gal) plasmid selective media. Cells grown overnight in medium containing 8 µg/ml of doxycycline were diluted to $OD_{600} = 0.5$ in plasmid selective dex, gly or gal without doxycycline, grown for 24 h, harvested, and lysed for immunoblotting[35]. Normalized cell lysates boiled for 5 mins with 2% SDS in 80 mM DTT sample buffer were resolved on 10% SDS-PAGE followed by immunoblotting with rabbit polyclonal α-TDP-43 antibody (1:3000, Proteintech Group). Also, PGK, yeast 3-Phosphoglycerate Kinase, was detected with anti PGK antibodies (1:10,000, Novex) as an internal control (3c upper). Immunoblot signals for

TDP-43-YFP and PGK were quantified and converted into % ratios of TDP43-YFP to PGK (3c lower). Standard error shown was calculated from 3 independent immunoblots. Gateway destination vector p1576 was made by dropping reading frame B gateway cassette (http://www.lifetechnologies.com) into pCM184[36]. TDP-43-YFP from entry vector pDONR TDP-43-YFP (Addgene plasmid # 27470) was then cloned into p1576 using an LR reaction, creating doxycycline-repressible *CEN TET-TDP-43-YFP TRP1* (p2223).

Fig. 2. TDP-43-YFP is still toxic in non-respiring cells and in cells lacking functional mitochondria. (a) Spot test measure of TDP-43-YFP toxicity. *CUP1-TDP-43-YFP* in p2261 (pAG305 *CUP1::TDP43-YFP*) was integrated at the *LEU2* locus in [*RHO+*] [*PIN+*] and [*RHO+*] [*pin-*] 74D-694 cells. Isogenic [*rho0*] versions of the *CUP1-TDP-43-YFP* integrants were acquired by growing [*RHO+*] integrants on YPD with 0.4 mg/ml ethidium bromide. This method has been shown to cause the loss of all mitochondrial DNA[37]. The lack of functional mitochondria was verified as cells failed to grow on glycerol medium. To assay for TDP-43-YFP toxicity, two independent subclones of [*RHO+*] and [*rho0*] *CUP1-TDP-43-YFP* medium (m) and high (h) level toxicity integrants (Supplementary Fig. 2) (↑TDP-43) and control empty integrants (↑no) grown on dextrose medium without $CuSO_4$ were normalized, serially diluted (1/10) and spotted on dextrose (dex) and galactose (gal) plates including 0, 100, and 250 μM $CuSO_4$. Plates with [*rho0*] cells were allowed to grow longer than plates with [*RHO+*] cells and galactose plates were grown longer than dextrose plates prior to scanning. Reduced growth indicated toxicity. As expected [*rho0*] cells failed to grow on gal. Integrants were constructed by transforming cells with purified linear integrating vector p*CUP1-TDP-43-YFP* (p2261) that had been cut in the *LEU2* gene with BstXI and control integration vector without TDP-43-YFP, pAG305 GAL1-ccdB (Addgene plasmid #14137) that was cut similarly, and selecting for transformants on integrant selective dextrose plates. Integration was verified by observing TDP-43-YFP aggregation in transformants in dextrose medium with 50 μM $CuSO_4$. To generate p2261, the HpaI-PmeI

fragment containing the *TET* promoter from pAG305 *TET-TDP-43-YFP* (p2237) was switched with the

CUP1 promoter from p1988 (*pCUP1-SOD1-GFP*) cut with the same restriction site. Plasmid pAG305 *TET-TDP-43-YFP* (p2237) was made by replacing the *GAL* promoter in pAG305 *GAL-ccdB* (Addgene plasmid # 14137) with PCR amplified HpaI-XbaI *TET* promotor from pCM184. Integrants contain different numbers of tandem plasmids which caused different levels of toxicity. (b) Viability measure of TDP-43-YFP toxicity. Cytotoxicity of the *CUP1* driven TDP-43-YFP in both [*RHO*$^+$] and [*rho^0*] cells was determined by colony forming units[8]. Overnight cultures of a TDP-43-YFP integrant associated with a high level of toxicity (supplementary Fig. 2) or vector controls grown without $CuSO_4$ were inoculated into liquid synthetic dextrose media with 0, 50 or 100 μM $CuSO_4$ to a final OD_{600} of 0.5. Samples were taken thereafter at 8, 24, 48, and 72 h and colony forming units determined on synthetic dextrose plates lacking $CuSO_4$. The number of colonies grown after 3 d of incubation at 30°C were counted and converted into viability (%) calculated as ratio of cells grown in TDP-43-YFP inducing media (with 50 or 100 μM $CuSO_4$) over cells grown in non-inducing media (0 μM $CuSO_4$). Error bars present the standard error calculated from 3 independent subclones of integrants. (c) No significant difference in levels of TDP43-YFP expression in [*RHO*$^+$] vs. [*rho^0*] *CUP1-TDP-43-YFP* integrated strains. Isogenic [*RHO*$^+$] vs. [*rho^0*] high toxicity integrants taken from dextrose plates were inoculated into 50 ml dextrose containing 0, 50, or 100 μM $CuSO_4$ to an OD_{600} of 0.5. Cells were harvested and lysed at 24 h. Normalized cell lysates were boiled, resolved and immunoblotted as shown in Fig. 1c. The intensity of each protein band (upper) was scanned and the ratio between TDP-43-YFP and PGK was calculated (lower). Bars represent standard error calculated from 3 independent immunoblots. (d) TDP-43-YFP aggregates similarly in [*RHO*$^+$] and [*rho^0*] *CUP1-TDP-43-YFP* integrated strains. Samples from 2b were visualized with fluorescent microscope at the same exposure. No obvious visual differences were seen.

Fig. 3. H_2O_2 enhances TDP-43 toxicity and aggregation. (a) H_2O_2 enhances TDP-43 toxicity. [RHO^+] *and*

[rho^0] versions of strain 74D-694 with integrated *CUP1-TDP-43-YFP* (↑TDP-43) associated with a low level

of toxicity (Supplementary Fig. 2), or vector control (No TDP-43), were grown to $OD_{600} = 0.5$ in synthetic

dextrose non-inducing medium and then for an additional 24 h after the simultaneous addition of $CuSO_4$

to 250 μM to induce TDP-43-YFP expression and H_2O_2 to the concentrations listed. Serial dilutions of

cells were then spotted on plates lacking $CuSO_4$ and H_2O_2 and photographed after growth. (b) H_2O_2

enhances TDP-43 aggregation. Cells from the liquid cultures used in (a) were examined under a

fluorescent microscope. Left shows bright field, right is fluorescence plus bright field. Since the intensity

of TDP-43-YFP aggregates was much brighter in H_2O_2 treated cells than in cells without H_2O_2 treatment,

images of H_2O_2 treated cells were taken with less exposure than cells without H_2O_2 treatment to allow

visualization of dots. (c) The level of TDP-43 in cells is unchanged by the addition of H_2O_2, but is

dramatically reduced by the addition of NAC. Freshly grown subclones of medium and high level toxicity

CUP-TDP-43-YFP integrants or control (empty vector) were inoculated at $OD_{600} = 0.5$ into 50 mL plasmid

selective dextrose media with 250 μM $CuSO_4$ and either 2 mM H_2O_2 or 1 mM NAC. Cells were harvested

and lysed after 24 h growth. Normalized cell lysates were boiled, resolved and immunoblotted as shown

in Fig. 1c. The protein level was compared between cells treated or not treated with H_2O_2 or NAC. The

ratio between scanned TDP-43-YFP and PGK signals for control (blue) and H_2O_2 (red) cells is shown. Bars

represent standard error calculated from 3 independent immunoblots.

Fig. 4. Antioxidant reduces TDP-43 level and toxicity. (a) Antioxidant reduces TDP-43 growth inhibition.

Isogenic [RHO^+] and [rho^-] versions of *CUP1-TDP-43-YFP* (TDP-43) integrants with medium (m) and high

(h) levels of toxicity and vector (control) grown on plasmid selective medium were normalized, serially

diluted (1/10) and spotted on plates with the indicated levels of $CuSO_4$ and antioxidant, NAC (N-acetyl-L-

cysteine, Sigma). Plates were photographed after 4 days of incubation at 30°C. (b) NAC reduces lethality

due to TDP-43. An integrant of *CUP1-TDP-43-YFP* (TDP-43) with high toxicity and vector (control) grown on synthetic dextrose medium without $CuSO_4$ were inoculated to OD_{600} = 0.5 in dextrose medium with (+) or without (-) 250 µM $CuSO_4$ and 0 mM or 1 mM NAC. After 24 h growth cells were stained with trypan blue to identify dead cells[5] and counted. (c) NAC reduces TDP-43-YFP level. Cells from (b) were photographed with bright field (left), fluorescence and bright field (middle) and just fluorescence (right). TDP-43-YFP fluorescence was barely observed in NAC treated cells even at the very high exposure shown. Black cells indicate trypan blue stained dead cells.

Supplementary Fig. 1. Slow growth does not enhance TDP-43 toxicity. Three independent transformants of [*PIN*+] 74D-694 (L1749) with either *TET-TDP-43-YFP* (TDP-43) or empty vector (v) were taken from plasmid selective media without TDP-43-YFP expression, normalized to OD_{600} = 3 and serially diluted (1/10) and spotted on dextrose (dex) or galactose (gal) plasmid selective inducing media (lacking doxycycline). Dextrose and galactose plates were respectively incubated at room temperature (RT) or 30°C for the number of days indicated prior to being photographed.

 Supplementary Fig. 2. *CUP1* controlled *TDP-43-YFP* integrants show differences in toxicity. Cells of different [*PIN*+] integrants were normalized and spotted on plasmid selective dextrose plates containing 0, 50, and 250 µM $CuSO_4$ as shown. Integrants were scored as having low, medium and high toxicity.

Acknowledgements

We thank Aaron Gitler, Stanford U. for plasmids and Ruben Dagda, U. of Nevada, Reno, and Martin Duennwald, Western U. Ca, for helpful ideas. This work was supported National Institutes of Health Grant R01GM056350 (SWL).

References

1.   Dormann, D. & Haass, C. (2011). TDP-43 and FUS: a nuclear affair. *Trends Neurosci* **34**, 339-48.

2.   Baloh, R. H. (2011). TDP-43: the relationship between protein aggregation and neurodegeneration in amyotrophic lateral sclerosis and frontotemporal lobar degeneration. *FEBS J* **278**, 3539-49.

3.   McKee, A. C., Gavett, B. E., Stern, R. A., Nowinski, C. J., Cantu, R. C., Kowall, N. W., Perl, D. P., Hedley-Whyte, E. T., Price, B., Sullivan, C., Morin, P., Lee, H. S., Kubilus, C. A., Daneshvar, D. H., Wulff, M. & Budson, A. E. (2010). TDP-43 proteinopathy and motor neuron disease in chronic traumatic encephalopathy. *J Neuropathol Exp Neurol* **69**, 918-29.

4.   Robinson, J. L., Geser, F., Stieber, A., Umoh, M., Kwong, L. K., Van Deerlin, V. M., Lee, V. M. & Trojanowski, J. Q. (2013). TDP-43 skeins show properties of amyloid in a subset of ALS cases. *Acta Neuropathol* **125**, 121-31.

5.   Park, S. K., Hong, J. Y., Arslan, F., Kanneganti, V., Patel, B., Tietsort, A., Tank, E. M. H., Li, X., Barmada, S. J. & Liebman, S. W. (2017). Overexpression of the essential Sis1 chaperone reduces TDP-43 effects on toxicity and proteolysis. *PLoS Genet* **13**, e1006805.

6.   Leibiger, C., Deisel, J., Aufschnaiter, A., Ambros, S., Tereshchenko, M., Verheijen, B. M., Buttner, S. & Braun, R. J. (2018). Endolysosomal pathway activity protects cells from neurotoxic TDP-43. *Microb Cell* **5**, 212-214.

7.   Leibiger, C., Deisel, J., Aufschnaiter, A., Ambros, S., Tereshchenko, M., Verheijen, B. M., Buttner, S. & Braun, R. J. (2018). TDP-43 controls lysosomal pathways thereby determining its own clearance and cytotoxicity. *Hum Mol Genet* **27**, 1593-1607.

8. Braun, R. J., Sommer, C., Carmona-Gutierrez, D., Khoury, C. M., Ring, J., Buttner, S. & Madeo, F. (2011). Neurotoxic 43-kDa TAR DNA-binding protein (TDP-43) triggers mitochondrion-dependent programmed cell death in yeast. *J Biol Chem* **286**, 19958-72.

9. Shan, X., Chiang, P. M., Price, D. L. & Wong, P. C. (2010). Altered distributions of Gemini of coiled bodies and mitochondria in motor neurons of TDP-43 transgenic mice. *Proc Natl Acad Sci U S A* **107**, 16325-30.

10. Magrane, J., Cortez, C., Gan, W. B. & Manfredi, G. (2014). Abnormal mitochondrial transport and morphology are common pathological denominators in SOD1 and TDP43 ALS mouse models. *Hum Mol Genet* **23**, 1413-24.

11. Xu, Y. F., Gendron, T. F., Zhang, Y. J., Lin, W. L., D'Alton, S., Sheng, H., Casey, M. C., Tong, J., Knight, J., Yu, X., Rademakers, R., Boylan, K., Hutton, M., McGowan, E., Dickson, D. W., Lewis, J. & Petrucelli, L. (2010). Wild-type human TDP-43 expression causes TDP-43 phosphorylation, mitochondrial aggregation, motor deficits, and early mortality in transgenic mice. *J Neurosci* **30**, 10851-9.

12. Hong, K., Li, Y., Duan, W., Guo, Y., Jiang, H., Li, W. & Li, C. (2012). Full-length TDP-43 and its C-terminal fragments activate mitophagy in NSC34 cell line. *Neurosci Lett* **530**, 144-9.

13. Wang, W., Li, L., Lin, W. L., Dickson, D. W., Petrucelli, L., Zhang, T. & Wang, X. (2013). The ALS disease-associated mutant TDP-43 impairs mitochondrial dynamics and function in motor neurons. *Hum Mol Genet* **22**, 4706-19.

14. Wang, W., Wang, L., Lu, J., Siedlak, S. L., Fujioka, H., Liang, J., Jiang, S., Ma, X., Jiang, Z., da Rocha, E. L., Sheng, M., Choi, H., Lerou, P. H., Li, H. & Wang, X. (2016). The inhibition of TDP-43 mitochondrial localization blocks its neuronal toxicity. *Nat Med* **22**, 869-78.

15. Izumikawa, K., Nobe, Y., Yoshikawa, H., Ishikawa, H., Miura, Y., Nakayama, H., Nonaka, T., Hasegawa, M., Egawa, N., Inoue, H., Nishikawa, K., Yamano, K., Simpson, R. J., Taoka, M.,

Yamauchi, Y., Isobe, T. & Takahashi, N. (2017). TDP-43 stabilises the processing intermediates of mitochondrial transcripts. *Sci Rep* **7**, 7709.

16. Dols-Icardo, O., Nebot, I., Gorostidi, A., Ortega-Cubero, S., Hernandez, I., Rojas-Garcia, R., Garcia-Redondo, A., Povedano, M., Llado, A., Alvarez, V., Sanchez-Juan, P., Pardo, J., Jerico, I., Vazquez-Costa, J., Sevilla, T., Cardona, F., Indakoechea, B., Moreno, F., Fernandez-Torron, R., Munoz-Llahuna, L., Moreno-Grau, S., Rosende-Roca, M., Vela, A., Munoz-Blanco, J. L., Combarros, O., Coto, E., Alcolea, D., Fortea, J., Lleo, A., Sanchez-Valle, R., Esteban-Perez, J., Ruiz, A., Pastor, P., Lopez De Munain, A., Perez-Tur, J. & Clarimon, J. (2015). Analysis of the CHCHD10 gene in patients with frontotemporal dementia and amyotrophic lateral sclerosis from Spain. *Brain* **138**, e400.

17. Johnson, J. O., Glynn, S. M., Gibbs, J. R., Nalls, M. A., Sabatelli, M., Restagno, G., Drory, V. E., Chio, A., Rogaeva, E. & Traynor, B. J. (2014). Mutations in the CHCHD10 gene are a common cause of familial amyotrophic lateral sclerosis. *Brain* **137**, e311.

18. Muller, K., Andersen, P. M., Hubers, A., Marroquin, N., Volk, A. E., Danzer, K. M., Meitinger, T., Ludolph, A. C., Strom, T. M. & Weishaupt, J. H. (2014). Two novel mutations in conserved codons indicate that CHCHD10 is a gene associated with motor neuron disease. *Brain* **137**, e309.

19. Zhang, M., Xi, Z., Zinman, L., Bruni, A. C., Maletta, R. G., Curcio, S. A., Rainero, I., Rubino, E., Pinessi, L., Nacmias, B., Sorbi, S., Galimberti, D., Lang, A. E., Fox, S., Surace, E. I., Ghani, M., Guo, J., Sato, C., Moreno, D., Liang, Y., Keith, J., Traynor, B. J., St George-Hyslop, P. & Rogaeva, E. (2015). Mutation analysis of CHCHD10 in different neurodegenerative diseases. *Brain* **138**, e380.

20. Chio, A., Mora, G., Sabatelli, M., Caponnetto, C., Traynor, B. J., Johnson, J. O., Nalls, M. A., Calvo, A., Moglia, C., Borghero, G., Monsurro, M. R., La Bella, V., Volanti, P., Simone, I., Salvi, F., Logullo, F. O., Nilo, R., Battistini, S., Mandrioli, J., Tanel, R., Murru, M. R., Mandich, P., Zollino, M., Conforti, F. L., Brunetti, M., Barberis, M., Restagno, G., Penco, S. & Lunetta, C. (2015). CHCH10

mutations in an Italian cohort of familial and sporadic amyotrophic lateral sclerosis patients. *Neurobiol Aging* **36**, 1767 e3-1767 e6.

21.   Ronchi, D., Riboldi, G., Del Bo, R., Ticozzi, N., Scarlato, M., Galimberti, D., Corti, S., Silani, V., Bresolin, N. & Comi, G. P. (2015). CHCHD10 mutations in Italian patients with sporadic amyotrophic lateral sclerosis. *Brain* **138**, e372.

22.   Jiao, B., Xiao, T., Hou, L., Gu, X., Zhou, Y., Zhou, L., Tang, B., Xu, J. & Shen, L. (2016). High prevalence of CHCHD10 mutation in patients with frontotemporal dementia from China. *Brain* **139**, e21.

23.   Woo, J. A., Liu, T., Trotter, C., Fang, C. C., De Narvaez, E., LePochat, P., Maslar, D., Bukhari, A., Zhao, X., Deonarine, A., Westerheide, S. D. & Kang, D. E. (2017). Loss of function CHCHD10 mutations in cytoplasmic TDP-43 accumulation and synaptic integrity. *Nat Commun* **8**, 15558.

24.   Johnson, B. S., McCaffery, J. M., Lindquist, S. & Gitler, A. D. (2008). A yeast TDP-43 proteinopathy model: Exploring the molecular determinants of TDP-43 aggregation and cellular toxicity. *Proc Natl Acad Sci U S A* **105**, 6439-44.

25.   Elden, A. C., Kim, H. J., Hart, M. P., Chen-Plotkin, A. S., Johnson, B. S., Fang, X., Armakola, M., Geser, F., Greene, R., Lu, M. M., Padmanabhan, A., Clay-Falcone, D., McCluskey, L., Elman, L., Juhr, D., Gruber, P. J., Rub, U., Auburger, G., Trojanowski, J. Q., Lee, V. M., Van Deerlin, V. M., Bonini, N. M. & Gitler, A. D. (2010). Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. *Nature* **466**, 1069-75.

26.   De Deken, R. H. (1966). The Crabtree effect: a regulatory system in yeast. *J Gen Microbiol* **44**, 149-56.

27.   Cohen, T. J., Hwang, A. W., Unger, T., Trojanowski, J. Q. & Lee, V. M. (2012). Redox signalling directly regulates TDP-43 via cysteine oxidation and disulphide cross-linking. *EMBO J* **31**, 1241-52.

28. Cohen, T. J., Hwang, A. W., Restrepo, C. R., Yuan, C. X., Trojanowski, J. Q. & Lee, V. M. (2015). An acetylation switch controls TDP-43 function and aggregation propensity. *Nat Commun* **6**, 5845.

29. Ecker, D. J., Khan, M. I., Marsh, J., Butt, T. R. & Crooke, S. T. (1987). Chemical synthesis and expression of a cassette adapted ubiquitin gene. *J Biol Chem* **262**, 3524-7.

30. Thiele, D. J. (1988). ACE1 regulates expression of the Saccharomyces cerevisiae metallothionein gene. *Mol Cell Biol* **8**, 2745-52.

31. Plessis, A. & Dujon, B. (1993). Multiple tandem integrations of transforming DNA sequences in yeast chromosomes suggest a mechanism for integrative transformation by homologous recombination. *Gene* **134**, 41-50.

32. Liebman, S. W., Bagriantsev, S. N. & Derkatch, I. L. (2006). Biochemical and genetic methods for characterization of [PIN+] prions in yeast. *Methods* **39**, 23-34.

33. David, D. C., Ollikainen, N., Trinidad, J. C., Cary, M. P., Burlingame, A. L. & Kenyon, C. (2010). Widespread protein aggregation as an inherent part of aging in C. elegans. *PLoS Biol* **8**, e1000450.

34. Weids, A. J., Ibstedt, S., Tamas, M. J. & Grant, C. M. (2016). Distinct stress conditions result in aggregation of proteins with similar properties. *Sci Rep* **6**, 24554.

35. Bagriantsev, S. N., Gracheva, E. O., Richmond, J. E. & Liebman, S. W. (2008). Variant-specific [PSI+] infection is transmitted by Sup35 polymers within [PSI+] aggregates with heterogeneous protein composition. *Mol Biol Cell* **19**, 2433-43.

36. Gari, E., Piedrafita, L., Aldea, M. & Herrero, E. (1997). A set of vectors with a tetracycline-regulatable promoter system for modulated gene expression in Saccharomyces cerevisiae. *Yeast* **13**, 837-48.

37.   Goldring, E. S., Grossman, L. I., Krupnick, D., Cryer, D. R. & Marmur, J. (1970). The petite

        mutation in yeast. Loss of mitochondrial deoxyribonucleic acid during induction of petites with

        ethidium bromide. *J Mol Biol* **52**, 323-35.

(a)

(b)

(c)

(a)

(b)

(c)

(d)

(a)

	no Cu++			250 μM Cu++		
	0	0.1	0	0.05		0.1 mM NAC

[RHO⁺] — control, TDP-43 (m), TDP-43 (h)

[rho⁰] — control, TDP-43 (h), TDP-43 (m)

(b)

TDP-43-YFP	% dead cells (dead/total cells)					
	−	+	+	−	+	+
	[RHO⁺]			[rho⁰]		
0 mM NAC	9.4 (92/982)	28.1 (248/882)	34.3 (282/822)	9.9 (88/886)	15.6 (102/653)	25.3 (251/992)
1 mM NAC	6.5 (71/1096)	10.5 (94/892)	19.0 (185/971)	7.0 (68/970)	7.5 (67/891)	6.7 (89/921)

(c)

	0 mM NAC	1mM NAC (higher exposure)
[RHO⁺]		
[rho⁰]		

Days of growth

	2		3		4		5	
	RT dex	30°C gal	RT dex	30°C gal	RT dex	30°C gal	RT dex	30°C gal

v

TDP-43

toxic level:

	0	50	250 µMCuSO4
medium			
medium			
medium			
medium			
high			
low			