An interaction between Gβγ and RNA polymerase II regulates transcription in cardiac
fibroblasts

Shahriar M. Khan[†], Ryan D. Martin[†], Sarah Gora, Celia Bouazza, Jace Jones-Tabah, Andy
Zhang, Sarah MacKinnon, Phan Trieu, Paul B.S. Clarke, Jason C. Tanny*, and Terence E.
Hébert*

[1]Department of Pharmacology and Therapeutics, McGill University, Montréal, Québec, H3G
1Y6, Canada

†These authors contributed equally to the study.

*To whom correspondence should be addressed.

Dr. Terence E. Hébert, PhD,

Department of Pharmacology and Therapeutics,

McGill University,

3655 Promenade Sir-William-Osler, Room 1303

Montréal, Québec, H3G 1Y6, Canada

Tel: (514) 398-1398

E-mail: terence.hebert@mcgill.ca

OR

Dr. Jason C. Tanny, PhD,

Department of Pharmacology and Therapeutics,

McGill University,

3655 Promenade Sir-William-Osler, Room 1303

Montréal, Québec, H3G 1Y6, Canada

Tel: (514) 398-3608

E-mail: jason.tanny@mcgill.ca

SUMMARY

Gβγ subunits are involved in many different signalling processes in various compartments of the cell, including the nucleus. To gain insight into the functions of nuclear Gβγ, we investigated the functional role of Gβγ signalling in regulation of GPCR-mediated gene expression in primary rat neonatal cardiac fibroblasts. Following activation of the angiotensin II type I receptor in these cells, Gβγ dimers interact with RNA polymerase II (RNAPII). Our findings suggest that $G\beta_1$ recruitment to RNAPII negatively regulates the fibrotic transcriptional response, which can be overcome by strong fibrotic stimuli. The interaction between Gβγ subunits and RNAPII expands the role for Gβγ signalling in cardiac fibrosis. The Gβγ-RNAPII interaction was regulated by signaling pathways in HEK 293 cells that diverged from those operating in cardiac fibroblasts. Thus, the interaction may be a conserved feature of transcriptional regulation although such regulation may be cell-specific.

INTRODUCTION

In recent years, study of the role of paracrine interactions between cardiomyocytes and cardiac fibroblasts in modulating the response to cardiac damage has expanded dramatically. Cardiac fibroblasts, in particular, respond dynamically following damage to the myocardium which is characterized by differentiation into myofibroblasts, increased proliferation and migration to areas of damage (Travers et al., 2016, Fu et al., 2018, Dobaczewski et al., 2010). This fibrotic response is modulated by the renin-angiotensin system, acting predominantly through the peptide ligand angiotensin II (Ang II) (Murphy et al., 2015, Kawano et al., 2000). Ang II drives changes in fibroblast function both directly and indirectly by increasing expression of other pro-fibrotic growth factors, such as transforming growth factor β1 (TGF-β1) (Campbell and Katwa, 1997). Collectively, these factors regulate alterations in cardiac architecture required for tissue repair by modulating the expression of genes encoding extracellular matrix proteins and proteases (Rosenkranz, 2004, Gao et al., 2009). Ang II also promotes cytokine secretion, thereby triggering autocrine and paracrine signalling to elicit further responses (Cheng et al., 2003, Ahmed et al., 2004). These signalling events create a feedforward loop, amplifying the fibrotic response from the initial area of damage to more distal regions of the heart (Ma et al., 2018). While the process initially aids in wound healing, a prolonged, activated fibrotic response worsens adverse cardiac remodelling and accelerates progression to heart failure (Travers et al., 2016, Weber et al., 2013). Importantly, inhibiting aspects of the fibrotic response reduces adverse cardiac remodelling (Fu et al., 2018, Weber and Diez, 2016). Hence, deciphering how Ang II signalling regulates pro-fibrotic gene expression is an important step towards understanding how these processes might be targeted therapeutically.

Cardiac fibroblasts respond to increased Ang II levels through Ang II type I (AT1R) and

type II (AT2R) G protein-coupled receptors (GPCRs). Of these, the AT1R is responsible for

positively regulating the fibrotic response in cardiac fibroblasts (Travers et al., 2016). The AT1R

couples to multiple heterotrimeric G proteins composed of specific combinations of Gα and Gβγ

subunits (Namkung et al., 2018). G proteins serve as signal transducers to relay extracellular

ligands bound to GPCRs into activation of different intracellular signalling pathways (Khan et

al., 2013). Gβγ subunits, like the more extensively studied Gα subunits, modulate a wide variety

of canonical GPCR effectors at the cellular surface such as adenylyl cyclases, phospholipases

and inwardly rectifying potassium channels (Khan et al., 2013, Dupré DJ, 2009, Smrcka, 2008).

However, compared with Gα-mediated events, Gβγ-mediated signalling is relatively

understudied and is complicated by the existence of 5 Gβ and 12 Gγ subunits which can combine

in multiple ways to form obligate dimers. Gβγ subunits also regulate a variety of non-canonical

effectors in distinct intracellular locations, and a number of studies have described roles for Gβγ

signalling in the nucleus (Khan et al., 2013, Campden et al., 2015a). Nuclear Gβγ subunits

modulate gene expression through interactions with a variety of transcription factors, such as

adipocyte enhancer binding protein 1 (AEBP1), the AP-1 subunit c-Fos, HDAC5 and MEF2A

(Park et al., 1999, Robitaille et al., 2010, Spiegelberg and Hamm, 2005, Bhatnagar et al., 2013).

Furthermore, we have detected $Gβ_1$ occupancy at numerous gene promoters in HEK 293 cells

(Khan et al., 2015). While canonical Gβγ signalling has been implicated in both cardiac fibrosis

and heart failure (Kamal et al., 2017, Travers et al., 2017), how nuclear Gβγ signalling impacts

these events is currently unknown.

Here, we describe a novel interaction between Gβγ subunits and RNA polymerase II

(RNAPII) which regulates the cardiac fibrotic response to Ang II activation of AT1R. We

characterize the GPCR-dependent, signalling pathway-specific regulation of this interaction in

primary neonatal rat cardiac fibroblasts and in HEK 293 cells. To understand the potential role of

individual $G\beta\gamma$ subunits, we knocked down $G\beta_1$ and $G\beta_2$ as exemplars of $G\beta$ subunits highly

expressed in these cells and characterized how nuclear $G\beta_1$, in particular, is a key regulator of

AT1R-driven transcriptional changes.

RESULTS

Gβγ interaction with RNAPII following activation of Gαq-coupled GPCRs

As Gβγ interacts with transcription factors and occupies gene promoter regions, we hypothesized that Gβγ subunits interact with a protein complex ubiquitously involved in transcription, and we initially focused on RNAPII. We assessed the potential Gβγ-RNAPII interaction following endogenous M3-muscarinic acetylcholine receptors (M3-mAChRs) activation with carbachol in HEK 293F cells. An initial co-immunoprecipitation time course experiment revealed a carbachol-induced interaction between endogenous Gβγ subunits (Gβ$_{1-4}$ detected with a pan-Gβ antibody) and Rpb1, the largest subunit of RNAPII, peaking between 45 and 120 mins (**Supplemental Figure 1A, B**). Immunoprecipitation of Rpb1 with two different antibodies also co-immunoprecipitated Gβ$_{1-4}$ in an agonist-dependent manner (**Supplemental Figure 1C**). Further, we observed no basal or carbachol-dependent interaction of Rpb1 with Gαq/11 or ERK1/2 (**Supplemental Figure 1D, E**) suggesting that Gβγ was not in complex with these proteins when it was associated with RNAPII in the nucleus. Under similar conditions, we observed no basal or carbachol-dependent interaction of Gβγ subunits with the A194 subunit of RNA polymerase I (**Supplemental Figure 1F**), suggesting Gβγ is not recruited to all RNA polymerases.

We next assessed the whether the Gβγ-RNAPII interaction also occurred in primary rat neonatal cardiac fibroblasts following treatment with Ang II. A time-course co-immunoprecipitation experiment revealed an agonist induced Gβγ-RNAPII interaction with a major peak interaction observed 75 minutes post stimulation (**Figure 1A, B**). As cardiac fibroblasts express both AT1R and AT2R, we next examined which receptor subtype regulated

the response, by pre-treatment with the AT1R-specific antagonist losartan. Pre-treatment of cells

with losartan prior to Ang II treatment abolished the agonist-induced interaction, but preserved

the basal interaction, suggesting that AT1R, and not AT2R, is primarily responsible for

mediating the interaction (**Figure 1C, D**).

Although several Gβγ isoforms have been detected in the nucleus (Bhatnagar et al., 2013,

Campden et al., 2015b, Zhang et al., 2001), the mechanisms leading to entry of Gβγ into the

nucleus remain unknown. Using subcellular fractionation following M3-mAChR activation in

HEK 293F cells, we observed importin-β dependent translocation of Gβγ into the nucleus **(data**

not shown). In addition, the agonist-dependent interaction of Gβ$_{1-4}$ and RNAPII was blocked by

importazole pre-treatment, suggesting that nuclear import of Gβ$_{1-4}$ is required for the interaction

with RNAPII in these cells (**Supplemental Figure 2A, B**). Next, we determined the effect of

importazole pre-treatment on the Ang II-mediated Gβγ-RNAPII interaction in cardiac

fibroblasts. The Gβγ-RNAPII interaction was also ablated when nuclear import via importin-β

was inhibited, suggesting again that Gβγ subunits must translocate to the nucleus for the

interaction with RNAPII to occur (**Figure 1E, F**).

Signalling pathways regulating Gβγ-RNAPII interaction are cell-specific

We next examined signalling events downstream of receptor activation that could

mediate the interaction between Gβγ subunits and RNAPII. To this end, we pursued a

pharmacological and genetic approach using both cardiac fibroblasts (**Figure 2**) and HEK 293F

cells (**Figure 3**). Our data indicated that the pathways responsible for promoting the Gβγ-

RNAPII interaction are cell type specific. Since AT1R couples to both Gq/11 and Gi/o G

proteins (Sauliere et al., 2012), we used FR900359 to inhibit Gαq/11 (Schrage et al., 2015) and

pertussis toxin (PTX) to inhibit Gαi/o. The agonist-induced response was markedly (~80%)

decreased by the Gαq/11 inhibitor, and also decreased (~30%) by the Gαi/o inhibitor,

demonstrating that AT1R signalling through Gαq is the primary pathway leading to increased

Gβγ-RNAPII interaction (**Figure 2A-B, Supplemental Figure 3A-B**). We next used U71322 to

inhibit the activity of phospholipase Cβ (PLCβ), downstream of both Gq/11 and Gi/o (the latter

via Gβγ signalling). In cardiac fibroblasts, pre-treatment of U71322 blocked the agonist-induced

Gβγ-RNAPII interaction with no effect on the basal interaction, suggesting a pivotal role for

PLCβ (**Figure 2C, Supplemental Figure 3C**). Chelation of Ca^{2+} using BAPTA-AM in cardiac

fibroblasts also abrogated the Ang II-induced Gβγ-RNAPII interaction (**Figure 2D,**

Supplemental Figure 3D), as did treatment with the PKC inhibitor Gö6983 and the CaMKII

inhibitor KN-93 (**Figure 2E, F, Supplemental Figure 3E, F**). Conversely, the MEK1 inhibitor

U0126 led to an increased basal Gβγ-RNAPII interaction but abrogated the Ang II-induced

interaction (**Figure 2G, Supplemental Figure 3G**). Lastly, the calcineurin inhibitor cyclosporin

A lead to an increased basal interaction did not prevent further Ang II-dependent increase in

interaction (**Figure 2H**, **Supplemental Figure 3H**).

Extending these studies to HEK 293F cells, we observed a similar reliance on Gαq

signalling for the agonist-induced Gβγ-RNAPII interaction. The carbachol-induced Gβγ-RNAPII

interaction was prevented by pre-treatment with the Gαq inhibitor FR900359 (**Figure 3A and**

Supplemental Figure 4A) and also by CRISPR/Cas9-mediated knockout of Gαq/11/12/13

(**Figure 3B and Supplemental Figure 4B**). However, except for this common event, the

signalling pathways in cardiac fibroblasts and HEK 293F cells diverged substantially. In HEK

293F cells, U71322 also blocked the carbachol-induced Gβγ-RNAPII interaction but there was a

pronounced increase in the basal interaction (**Figure 3C, Supplemental Figure 4C**). Further

differences were observed following chelation of calcium with BAPTA-AM which increased

basal levels of the Gβγ-RNAPII interaction but did not block further carbachol-induced

stimulation of the interaction (**Figure 3D, Supplemental Figure 4D**), suggesting a modulatory

role for calcium in HEK 293F cells rather than the direct role seen in cardiac fibroblasts. HEK

293F cells employed different regulatory mechanisms involving protein kinases activated

downstream of Gαq/11-coupled GPCRs compared to cardiac fibroblasts. For example, the PKC

inhibitor Gö6983 and the CaMKII inhibitor KN-93 both increased basal levels of interaction but

did not block carbachol-induced interactions between Gβγ and Rpb1 (**Figure 3E, F,**

Supplemental Figure 4E, F). Indeed, inhibition of calcineurin with cyclosporin A blocked the

carbachol-mediated increase in interaction between Gβγ and Rpb1, suggesting a role for this

phosphatase in mediating the interaction in response to M3-mAChR activation (**Figure 3G** and

Supplemental Figure 4G). While the requirement for activation of Gαq is common for the Gβγ-

RNAPII interaction in both cell types, the regulation by downstream signalling pathways

diverges.

Roles of individual Gβ subunits in regulating the angiotensin II-activated fibrotic response in rat

neonatal cardiac fibroblasts

The Gβ family is comprised of five members which, with the exception of $G\beta_5$, exhibit

high levels of sequence and structural similarity (Khan et al., 2013). Despite these similarities,

Gβ isoforms differ considerably with respect to their associated receptors and signalling

pathways (Khan et al., 2015, Yim et al., 2019, Greenwood and Stott, 2019). As our above-

reported characterization used a pan-$G\beta_{1-4}$ antibody, we next sought to examine the specificity of

Gβ isoforms interacting with Rpb1 in cardiac fibroblasts. We initially focused on $G\beta_1$ and $G\beta_2$ as

they exhibit the highest expression in cardiac fibroblasts determined by RNA-seq (Shu et al., 2018) and RT-qPCR (**Supplemental Figure 5A**). Immunoprecipation with a $G\beta_1$ specific antibody revealed an increase in the amount of Rpb1 co-immunoprecipitated in response to Ang II treatment, whereas immunoprecipitation of $G\beta_2$ indicated a basal interaction with Rpb1 that was lost in response to Ang II treatment (**Supplemental Figure 5B**). We also assessed $G\beta$ isoform specificity in HEK 293F cells through heterologous expression of FLAG-tagged versions of each $G\beta$ subunit. In response to M3-mAChR activation, FLAG-$G\beta_1$ was the only isoform that showed an increased interaction with Rpb1 (**Supplemental Figure 5C, D**). Hence, an increased interaction between $G\beta_1$ and Rpb1 was seen in both cell types, suggesting that our earlier observations using the pan-$G\beta_{1-4}$ antibody likely reflected increased interactions with $G\beta_1$.

As we observed isoform-specific roles in RNAPII interactions, we next assessed how knockdown of either $G\beta$ isoform affected the interaction. We first validated knockdown conditions for each $G\beta$ subunit by siRNA at the mRNA and protein levels (**Supplemental Figure 6A, B**). We observed a reduction in the Ang II-induced $G\beta\gamma$-RNAPII interaction upon knockdown of $G\beta_1$, supporting $G\beta_1$ as the isoform involved in the increased interaction with Rpb1. Surprisingly, knockdown of $G\beta_2$ also prevented the Ang II-mediated increase in the $G\beta\gamma$-RNAPII interaction (**Figure 4A, B**). The loss of $G\beta\gamma$-RNAPII interaction after $G\beta_2$ knockdown, despite it not being involved in the Ang II-dependent increase, suggested that AT1R signalling could be altered by loss of $G\beta_2$ subunits.

We thus determined whether specific $G\beta$ isoforms were required to initiate signalling cascades proximal to AT1R activation. Following receptor activation, $G\beta\gamma$ subunits regulate intracellular Ca^{2+} mobilization through activation of $PLC\beta$ (Park et al., 1993). As we have previously demonstrated $G\beta$ isoform specificity for $PLC\beta$ signalling in HEK 293F cells (Khan et

al., 2015), we assessed the relative roles of $G\beta_1$ and $G\beta_2$ in AT1R-dependent Ca^{2+} mobilization.

To assess AT1R-dependent intracellular Ca^{2+} mobilization, we used the cell-permeable Ca^{2+} dye

Fura 2-AM. Following AT1R activation, we observed a rapid increase in intracellular Ca^{2+}

mobilization (**Figure 4A**, black, empty triangles) and the quantified area under the curve (**Figure**

4B, black bar). Knockdown of $G\beta_1$ did not alter Ca^{2+} mobilization following stimulation with

Ang II ($8.1 \pm 7.0\%$ decrease, red bar). However, knockdown of $G\beta_2$ resulted in a significant 31.6

$\pm 9\%$ decrease in Ca^{2+} release (**Figure 4A, B**, green bar), suggesting a role for $G\beta_2$-containing

$G\beta\gamma$ dimers in mediating receptor-proximal signalling downstream of AT1R activation. This

suggests Gβ2 knockdown prevented the Ang II-dependent increase in $G\beta\gamma$-RNAPII interaction

through disruption to AT1R Ca^{2+} signalling, aligning with the observed effect of Ca^{2+} chelation

with BAPTA-AM. These results highlight the complex interplay between cell surface receptors

and multiple $G\beta\gamma$ subunits, in modulating both basal and ligand stimulated RNAPII/$G\beta\gamma$

interactions.

$G\beta\gamma$ interacts with transcribing RNAPII

As we demonstrated that $G\beta\gamma$ is recruited to RNAPII following AT1R activation, which

also activates a transcriptional program in fibroblasts, we assessed the relationship between the

transcriptional response and $G\beta\gamma$ recruitment (Shu et al., 2018, Dang et al., 2015). To assess this

potential relationship, we disrupted the transcription cycle at two different regulatory points

through inhibition of Cdk7, a component of the general transcription factor TFIIH, and Cdk9, the

protein kinase subunit of P-TEFb (Zhou et al., 2012). Following RNAPII recruitment, Cdk7

activity stimulates promoter clearance of RNAPII to begin transcription. Soon after RNAPII

pauses at a promoter-proximal region and requires the activity of Cdk9 in order to be released

into productive elongation (Liu et al., 2015). We assessed involvement of both Cdk7 and Cdk9

on the Ang II-induced Gβγ-RNAPII interaction using the selective inhibitors THZ1 and iCdk9,

respectively (Lu et al., 2015, Kwiatkowski et al., 2014). THZ1 abrogated the Ang II-stimulated

Gβγ-RNAPII interaction (**Figure 5A, Supplemental Figure 7A**) while iCdk9 resulted in a loss

of both the basal and Ang II-stimulated Gβγ-RNAPII interaction (**Figure 5B, Supplemental**

Figure 7B). This suggests that the Gβγ-RNAPII interaction requires the transcriptional response

to Ang II in cardiac fibroblasts. As with cardiac fibroblasts, in HEK 293F cells disruption of the

transcriptional cycle through inhibition of Cdk7 and Cdk9 with DRB also blocked the increased

interaction between RNAPII and Gβγ (**data not shown**), showing that the Gβγ/RNAPII

interaction is dependent on an active transcriptional response in both cell types.

The role of Gβγ subunits in fibrotic gene expression

In order to understand the role of Gβγ in Ang II-regulated gene expression, we examined

changes in the levels of 84 genes involved in the fibrotic response using the Qiagen RT^2

Profiler^{TM} PCR array platform. Gene expression changes were assessed following 75 min or 24 h

Ang II treatment alongside Gβ1 or Gβ2 knockdown. These two time points were selected to

investigate the effect of disrupting the Gβγ-RNAPII interaction or, in the longer term, upstream

signalling, respectively. We assessed gene expression changes across all 67 genes remaining

after excluding genes below our chosen threshold of detection (i.e. Ct > 35). After 75 min of Ang

II treatment, we observed a similar upregulation of fibrotic genes in both control and Gβ$_1$

knockdown conditions (**Figure 5C, Supplemental Table 1**). However, Gβ$_1$ knockdown

increased both basal expression and the total number of genes altered by AT1R stimulation

(**Figure 5C, Supplemental Table 1**). Following 24 h Ang II treatment, this effect became more

pronounced. Gβ$_1$ knockdown led to increases in basal gene expression, expression regulated by

Ang II treatment and the overall number of genes upregulated (**Figure 5D, Supplemental Table**

1). The increased expression following Gβ$_1$ knockdown suggests the Gβγ-RNAPII interaction

negatively modulates the Ang II transcriptional response.

Whereas Gβ1 knockdown altered the transcriptional response to Ang II treatment,

disruption of AT1R signalling by Gβ$_2$ knockdown did not significantly alter basal fibrotic gene

expression or the overall response to 24 h Ang II treatment (**Figure 5D, Supplemental Table 1**).

The lack of effect of Gβ$_2$ knockdown suggests that Gβγ signalling through Ca^{2+} is not required

for AT1R-mediated transcriptional changes. To further address the role of Gβγ signalling, we

utilized the small-molecule pan-Gβγ inhibitor gallein (Lehmann et al., 2008). As with Gβ$_2$

knockdown, pre-treatment with gallein did not significantly alter the transcriptional response

following 24 h Ang II treatment (**Figure 5E**). This suggests that Gβγ-dependent signalling

downstream of the AT1R is not a key driver of transcriptional changes. Instead, Gβγ is required

to modulate processes driven by other signalling pathways and dampen the fibrotic response

until such signals rise above a threshold.

Genome-wide recruitment of Gβ$_1$ and the effect on RNAPII occupancy following Ang II

treatment

To assess the possibility of genome-wide Gβ$_1$ recruitment and changes in RNAPII

occupancy following 75 min Ang II treatment in cardiac fibroblasts, we performed chromatin

immunoprecipitation followed by next generation sequencing (ChIP-seq) for heterologously

expressed FLAG-Gβ$_1$ and endogenous Rpb1. We confirmed that, like endogenous Gβ$_1$, the

interaction of Rbp1 with heterologously expressed FLAG-Gβ$_1$ increased following AT1R

activation (**Supplemental Figure 8A, B**). We focused on genes with RNAPII peaks identified by

the peak calling software macs2 and annotated with HOMER (Heinz et al., 2010, Zhang et al.,

2008). The same Gβ_1 knockdown conditions that increased the number of genes upregulated in

response to Ang II (above) also increased the number of genes occupied by RNAPII following

Ang II treatment (**Figure 6A**). To identify groups of genes with similar FLAG-Gβ_1 and RNAPII

occupancy patterns, we performed K-means clustering with genes that RNAPII peaks were

identified in any treatment condition. Two K-means clusters were identified (98 genes in cluster

1 and 806 in cluster 2) with distinct occupancy patterns (**Figure 6B, C**). In cluster 1, FLAG-Gβ_1

occupancy increased within the gene body in response to Ang II. A similar but weaker tendency

was also observed in cluster 2 (**Figure 6B**). The increased FLAG-Gβ1 occupancy in cluster 1

corresponded to Gβ1-dependent changes to the Ang II-induced RNAPII occupancy alterations.

First, Ang II treatment led to increased RNAPII occupancy throughout the gene body under

siRNA control conditions (**Figure 6C**). In the absence of Ang II, Gβ_1 knockdown increased

RNAPII occupancy near transcription start sites (TSSs) which corresponds with increased gene

expression under these conditions (**Figure 6C**). Lastly, there was greater RNAPII occupancy

when Ang II treatment was combined with Gβ1 knockdown than in the absence of knockdown

(**Figure 6C**). Similar RNAPII occupancy patterns were observed in cluster 2, suggesting that

Gβ1 also plays a regulatory role along these genes and our FLAG-Gβ_1 ChIP-seq was not

sensitive enough to reliably detect Gβ_1. We also assessed the functional pathways enriched in

cluster 1, through gene ontology (GO) term enrichment. The top four significant GO terms

identified (corresponding to cellular processes such as inflammation, fibroblast activation and

apoptosis) indicate that Gβ_1 is recruited to genes involved in processes essential to fibrosis

(**Figure 6D**).

The increased number of genes with RNAPII occupancy in the Ang II and Gβ1

knockdown condition suggested that $G\beta_1$ occupancy impairs RNAPII recruitment. As such, we

would expect cluster 1 genes to be more enriched in genes with RNAPII occupancy under Ang II

and Gβ1 knockdown condition than Ang II and siRNA control conditions. Therefore, we

performed a Fisher's exact test to compare the proportion of cluster 1 genes in these treatment

conditions, which demonstrated a significant (p-value < 0.01) enrichment in the Ang II and Gβ1

knockdown condition gene list compared to Ang II and siRNA control condition. This again

suggests $G\beta_1$ functions to suppress RNAPII transcription following AT1R activation.

In order to assess the relationship between $G\beta_1$ occupancy and transcription, we focused

on genes from our fibrosis qPCR array that were also found in cluster 1. Eight genes from the

fibrosis array were identified in cluster 1, which included five of the seven genes upregulated

after 75 min of Ang II treatment such as thrombospondin 1 (Thbs1) and connective tissue growth

factor (Ctgf) (**Figure 6E, F**). We confirmed the Ang II-dependent increase in $G\beta_1$ occupancy

along Ctgf by ChIP-qPCR (**Supplemental Figure 8C**). We also assessed the effect of $G\beta_1$

knockdown on AT1R-dependent changes of RNAPII occupancy along Ctgf by ChIP-qPCR.

Similar to our ChIP-seq analysis, we observed a greater increase in RNAPII along the gene in

response to Ang II under siRNA GNB1 knockdown compared to siRNA control, where we

observed a slight decrease (**Supplemental Figure 8D**). We also validated the change in

expression of Ctgf by RT-qPCR using primers designed in-house (**Supplemental Table 2**).

Under control conditions Ang II had a minor effect on Ctgf expression, however in the absence

of $G\beta_1$ Ang II treatment resulted in a significant upregulation of Ctgf mRNA. (**Supplemental**

Figure 8E). Taken together, our results demonstrate $G\beta_1$ recruitment negatively regulates

expression of genes involved in the fibrotic response to Ang II by inhibiting early stages of the

RNAPII transcription cycle.

DISCUSSION

The functional specificity of Gβ and Gγ subunits has been mostly investigated in the context of signalling proximal to GPCR activation (i.e., the regulation of effector activity downstream of receptor stimulation) (Khan et al., 2013). In contrast, our findings provide new insights regarding non-canonical roles of specific Gβγ dimers in more distal events in the nucleus, particularly in the regulation of gene expression. Here, we demonstrate for the first time an interaction between Gβγ and RNAPII and investigate the regulatory signalling mechanisms in transformed cell lines (HEK 293 cells) and in primary cells (neonatal rat cardiac fibroblasts). The interaction of Gβγ and RNAPII represents a significant addition to the expanding list of Gβγ interactors, and our findings suggest that regulatory mechanisms impacting the interaction are dependent on cellular context. We also show that Gβγ signalling is a critical regulator of the fibrotic response in cardiac fibroblasts.

Our findings suggest that following acute treatment with Ang II, $G\beta_1$ is transiently recruited to pro-fibrotic genes to negatively regulate RNAPII recruitment, thereby limiting the fibrotic response following transient fluctuations in local Ang II concentrations likely seen *in vivo*. This negative RNAPII regulation may potentially occur through direct interactions with RNAPII, preventing its recruitment or other aspects of initiation, or else via an indirect mechanism in which Gβγ would form part of a larger RNAPII-containing complex altering the local chromatin landscape. We cannot currently distinguish between these two possibilities, given that our co-immunoprecipitation assay was performed using whole-cell lysates. On the other hand, chronic stress or damage to the heart leads to a sustained increase of Ang II concentrations in cardiac tissue (Sun and Weber, 1996, Passier et al., 1996). We propose that such sustained AT1R signalling overcomes the transient $G\beta_1$ "brake" to elicit a robust fibrotic

response. Alternatively, pro-fibrotic factors that are upregulated and secreted following AT1R

activation may elicit autocrine signalling pathways that overcome the $G\beta_1$ transcriptional

repression (Lee et al., 1995, Ma et al., 2018). Our gene expression data at 75 min, and more

especially at 24 h, begins to identify the increased number and greater gene expression in the

absence of the proposed negative regulatory mechanism when $G\beta_1$ is knocked down. Further

analysis of the kinetics of the interaction and how this changes the dynamics of chromatin

occupation or gene expression are required as well. Future experiments assessing nascent RNA

production are required to accurately determine gene expression changes at early time points.

We demonstrated that $G\beta_2$, and not $G\beta_1$, was important for proximal signalling

downstream of AT1R activation similar to the requirement of specific $G\beta$ isoforms for activation

of $PLC\beta$ in HEK 293 cells (Khan et al., 2015). Our data suggest that $G\beta_2$ plays a minimal role in

regulating AT1R-dependent gene expression *per se*. Rather, our findings using the broad-

spectrum $G\beta\gamma$ inhibitor gallein suggest that receptor-proximal $G\beta\gamma$ signalling in general is not

required for the transcriptional response and instead it is dependent on $G\alpha q$ signalling and more

distal $G\beta_1$-dependent events. Knockdown of $G\beta_2$ also compromised Ang II-mediated interactions

between $G\beta\gamma$ and RNAPII even though $G\beta_2$ had a limited role in the fibrotic transcriptional

response. This suggests $G\beta_2$ knockdown does not prevent the response but rather alters the

kinetics of $G\beta\gamma$-RNAPII interactions, which then translates into different fibrotic responses over

time. Further, the roles of specific $G\gamma$ subunits in mediating proximal signal transduction must

also be considered as for other $G\beta\gamma$ effectors (Khan et al., 2015), and should be the subject of

future studies. Taken together, our findings suggest that in fibrosis and potentially in other

diseases, the indiscriminate targeting of $G\beta\gamma$ signalling (e.g. with compounds such as gallein)

will result in outcomes that differ considerably from those obtained by targeting particular Gβγ

combinations (Lin and Smrcka, 2011, Kamal et al., 2011, Smrcka et al., 2008).

Analysis of the signalling networks regulating the Gβγ/RNAPII interaction yielded four

main conclusions: (1) different GPCR signalling systems in distinct cell types lead to different

kinetics of the Gβγ-RNAPII interaction, (2) different signalling pathways downstream of GPCR

activation act to both induce or modulate the interaction, (3) Gαq-coupled GPCRs regulate the

interaction in both cell types examined, and (4) signalling ultimately converged on activation of

transcription. Indeed, our results suggest that the cell context plays a critical role in determining

the mechanism by which the Gβγ-RNAPII interaction is regulated. First, in cardiac fibroblasts,

the Gβγ/RNAPII interaction depended on a Gq-PLCβ-Ca^{2+}-CaMKII/PKC/MEK-dependent

pathway downstream of AT1R activation, whereas calcineurin acted as a basal negative regulator

(summarized in **Supplemental Figure 9**). On the other hand, in HEK 293 cells, we observed that

the interaction was reliant on a Gq-PLCβ-Ca^{2+}-calcineurin pathway downstream of M3-mAChR

activation, whereby PKC and CaMKII both negatively regulate this interaction under basal

conditions (summarized in **Supplemental Figure 9**). The involvement of Ca^{2+}, PKC and

ERK1/2 in the induction of the Gβγ/RNAPII interaction in fibroblasts is supported by previous

reports that demonstrate their involvement in Ang II-induced fibrosis (Chintalgattu and Katwa,

2009, Olson et al., 2008).

The different signalling pathways promoting the Gβγ-RNAPII interaction appear to

converge at the point of Cdk7 and Cdk9 activation. In particular, we found that the Cdk7 and

Cdk9 inhibitors (DRB, THZ1 and iCdk9, respectively) inhibited both carbachol-induced Gβγ-

RNAPII interaction in HEK 293 cells and the analogous Ang II-induced interaction in cardiac

fibroblasts. This suggests the differential regulatory signalling pathways identified are due to cell

type- and receptor-specific activation pathways of both Cdk7 and Cdk9. The recruitment of Gβγ

serves as a common negative regulatory mechanism regardless of the pathway leading to

transcriptional activation. Furthermore, a strong connection has been established between the

control of transcriptional pausing and pathological cardiac remodelling, although primarily in the

cardiomyocyte (Yang et al., 2017, Sayed et al., 2013, Anand et al., 2013, Duan et al., 2017,

Stratton et al., 2016, Sano et al., 2002). Our results indicate that regulation of the early stages of

the RNAPII transcription cycle is also an important checkpoint in the fibrotic response mediated

by cardiac fibroblasts.

Taken together, the Gβγ-RNAPII interaction identifies a new mechanism by which Gβγ

modulates gene expression. Our study highlights the complex interplay of different Gβγ subunit

combinations at the cell surface and in the nucleus initiated upon stimulation of Gαq-coupled

receptors. Since Gβ$_1$γ dimers play an important role in regulating the expression of fibrotic genes

in cardiac fibroblasts, the development of selective Gβ$_1$γ inhibitors hold some promise for

preventing the pathological consequences of myocardial damage.

METHODS

Reagents –The following were all purchased from Sigma-Aldrich: carbachol, angiotensin II,

BAPTA-AM, KN-93, Gö6983, PTX, U0126, calyculin A, cyclosporin A, TRI reagent, isopropyl

thiogalactopyranoside (IPTG), protease inhibitor cocktail, triton X-100, bovine serum albumin,

ethylenediaminetetraacetic acid (EDTA), 70% NP-40 (Tergitol), sodium deoxycholate,

magnesium chloride, lithium chloride, anti-rabbit IgG (whole molecule)-agarose antibody, anti-

mouse IgG (whole molecule)-agarose antibody, goat anti-rabbit IgG (whole molecule)

conjugated to peroxidase secondary antibody, goat anti-mouse IgG (Fab specific) conjugated to

peroxidase secondary antibody, anti-FLAG M2 antibody, and rabbit IgG (St. Louis, MO, USA).

U71322 pan-PKC inhibitor was purchased from Biomol International (Plymouth Meeting, PA,

USA). Lysozyme (from hen egg white) and phenylmethylsulfonyl fluoride (PMSF) were

purchased from Roche Applied Sciences (Laval, QC, Canada). Ethylene glycol bis (2-

aminooethyl ether) N,N,N',N' tetraacetic acid (EGTA) and HEPES were purchased from

BioShop (Burlington, ON, Canada). Sodium chloride, glutathione (reduced form), dithiothreitol

(DTT) and Dynabeads protein G were purchased from Fisher Scientific (Ottawa, ON, Canada).

Dulbecco's modified Eagle's medium (DMEM) (supplemented with 4.5 g/L glucose, L-glutamine

and phenol red), DMEM low glucose (supplemented with 1.0 g/L glucose, L-glutamine and

phenol red), Hank's Balanced salt solution (HBSS), HBSS (with no phenol),

Penicillin/Streptomycin solution, Tris base buffer, ampicillin sodium salt, and fetal bovine serum

were purchased from Wisent (St. Bruno, QC, Canada). Glutathione sepharose 4B GST beads was

purchased from GE Healthcare (Mississauga, ON, Canada). Lipofectamine 2000 and Alexa Fluor

488 goat anti-mouse IgG were purchased from Invitrogen (Burlington, ON, Canada). Enhanced

chemiluminescence (ECL) Plus reagent was purchased from Perkin Elmer (Woodbridge, ON, Canada). Moloney murine leukemia virus reverse transcriptase (MMLV-RT) enzyme and recombinant RNasin® ribonuclease inhibitor were purchased from Promega (Madison, WI, USA). Evagreen 2X qPCR MasterMix was purchased from Applied Biological Materials Inc. (Vancouver, BC, Canada) and iQ SYBR Green Supermix was purchased from Bio-Rad Laboratories (Mississauga, ON, Canada). Anti-Gβ1-4 (T-20) antibody, anti-RNA Polymerase I Rpa194 (N-16) antibody, anti-ERK1/2 antibody, anti-Gαq antibody and anti-Rpb1 (N20) were purchased from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA). Anti-RNA polymerase II clone CTD4H8 (Rpb1) antibody was purchased from EMD Millipore (Temecula, CA, USA). Anti-*Schizosaccharomyces pombe* histone H2B (ab188271) antibody was purchased from Abcam Inc. (Toronto, ON, Canada). Polyclonal anti-Gβ_1 and anti-Gβ_2 were a generous gift of Professor Ron Taussig (UT Southwestern). THZ1 was a gift from Nathanael S. Gray (Harvard University) and iCdk9 was a gift from James Sutton (Novartis). FLAG-Gβ_1, FLAG-Gβ_2, FLAG-Gβ_3, FLAG-Gβ_4 and FLAG-Gβ_5 plasmids were obtained from UMR cDNA Resource (www.cdna.org).

Tissue culture, transfection and treatments – Human embryonic kidney 293 (HEK 293), HEK 293T cells and CRISPR/Cas9 generated ΔGαq/11/12/13 knockout HEK 293 cells (quadKO cells) (Devost et al., 2017), a generous gift from Dr. Asuka Inoue (Tohuku University, Sendai, Japan), were grown at 37°C in 5% CO_2 in DMEM supplemented with 5% (v/v) fetal bovine serum and 1% (v/v) penicillin/streptomycin (P/S). HEK 293 cells were transiently transfected with FLAG-Gβ1-5 using Lipofectamine 2000 as per the manufacturer's recommendations. Primary rat neonatal cardiac fibroblasts were isolated from 1-3 day old Sprague-Dawley rat pups (Charles

River Laboratories, St-Constant, Quebec) as previously described (Calderone et al., 1998). All

procedures using animals were approved by the McGill University Animal Care Committee, in

accordance with Canadian Council on Animal Care Guidelines. Two days after isolation, cells

were detached with trypsin/EDTA and plated at a density of ~8 x 10^3 cells/cm^2 in fibroblast

growth medium for 48h. For siRNA transfection, cardiac fibroblasts were plated at a density of

~20 x 10^3 cells/cm^2 and transfected using Lipofectamine 2000 as per the manufacturer's

instructions. For treatment of HEK 293F cells, HEK 293F quadKO cells or cardiac fibroblasts,

cells were serum-deprived for 6 h with DMEM or overnight (~12 h) with DMEM low glucose

(with no FBS and no P/S) respectively, and subsequently treated with pathway inhibitors, 1 mM

carbachol or 1 μM Ang II for the treatment lengths indicated in the various assays.

RT-qPCR – Reverse transcription of RNA isolated from rat neonatal cardiac fibroblasts was

performed as previously described (Khan et al., 2015). Briefly, cells were lysed in TRI reagent

and RNA was extracted using a protocol adapted from Ambion (Burlington, ON, Canada).

Reverse transcription was performed on 1 μg of total RNA using an MMLV-RT platform

according to the manufacturer's protocol. Subsequent qPCR analysis was performed with

Evagreen Dye qPCR master-mixes using a Corbett Rotorgene 6000 thermocycler or Bio-Rad

1000 Series Thermal Cycling CFX96 Optical Reaction module. mRNA expression data were

normalized to housekeeping transcripts for U6 snRNA. Ct values obtained were analyzed to

calculate fold change over respective control values using the $2^{-\Delta\Delta Ct}$ method. Primer sequences

for all primers used are listed in **Supplemental Table 2**.

Ca^{2+} mobilization – Cardiac fibroblasts were cultured as previously described following

transfection with respective siRNA. Cardiac fibroblasts were washed and media replaced with

HBSS (no phenol) and incubated for 1 h at 37°C and 5% CO_2. Media was replaced with Fura 2-

AM in HBSS and incubated for another 1 h at 37°C and 5% CO_2. Fura 2-AM containing media

was replaced with HBSS and Cardiac fibroblasts incubated for another 30 min at 37°C and 5%

CO_2 prior to recordings. Baseline recordings were obtained every 0.7 s for 10 s followed by

injection of Ang II to a final concentration of 1 μM and recordings obtained every 0.7 s for a

total of 1 min. A control well with no Fura-2 AM was included in order to control for

background fluorescence. Fluorescence intensity was recorded using Bio-Tek Synergy 2 Multi-

Mode Microplate Reader with fluorescence excitation at 340 nm or 360 nm and fluorescence

emission at 516 nm. Data is presented as the ratio of fluorescence emission at 516 nm following

340 nm excitation over 360 nm excitation. The ratio was normalized to the mean baseline ratio

from control cells.

Nuclear isolation – Nuclei from HEK 293 cells and cardiac fibroblasts were isolated as

previously described (Campden et al., 2015b). Briefly, cells seeded in T175 flasks (Corning)

were treated as indicated, washed three times with 1X PBS (137 mM NaCl, 2.7 mM KCl, 10 mM

Na_2HPO_4, 1.8 mM KH_2PO_4), and harvested in 1X PBS by centrifugation. Pelleted cells were

lysed in lysis buffer (320mM sucrose, 10 mM HEPES, 5 mM $MgCl_2$, 1 mM DTT, 1 mM PMSF,

1% Triton X-100), added gently on top of a high-sucrose buffer (1.8 M sucrose, 10 mM HEPES,

5 mM $MgCl_2$, 1 mM DTT, 1 mM PMSF), and centrifuged at 4600 g for 30 min at 4°C,

separating unlysed nuclei from the cytosolic fraction. Pelleted nuclei were then resuspended in

resuspension buffer (320 mM sucrose, 10 mM HEPES, 5 mM MgCl$_2$, 1 mM DTT, 1 mM

PMSF), pelleted at 300 g for 5 min and subsequently lysed in 1X RIPA buffer.

Immunoprecipitation and western blotting – Immunoprecipitation (IP) assays of Gβ and Rpb1

pull downs were performed as previously described, with minor alterations (Robitaille et al.,

2010). Protein extracts from treated HEK 293 cells and cardiac fibroblasts lysed in RIPA (1%

NP-40, 50 mM Tris-HCl ph 7.4, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 0.1% SDS, 0.5%

sodium deoxycholate) were quantified by Bradford assay and 500 μg of protein lysate was

precleared with 15 μl of anti-rabbit IgG-agarose beads. Precleared lysates were then incubated

with 1 μg anti-Gβ$_{1-4}$, 2 μg of anti-Rpb1 or anti-Gβ$_1$ serum or anti-Gβ$_2$ serum overnight at 4°C

with end-over mixing. The next day, 40 μl of washed agarose beads were added to each

lysate/antibody mixture, incubated for 3.5 hours at 4°C with end-over mixing, and then beads

were washed 3X with RIPA. Proteins were eluted off the beads by the addition of 4X Laemmli

buffer followed by denaturation at 65°C for 15 min. Protein immunoprecipitation and co-IP were

then assessed by western blot as previously described (Khan et al., 2015). Resulting western blot

images were quantified using ImageJ 1.48v.

Rat Fibrosis qPCR arrays – Fibrosis qPCR arrays were performed as per the manufacturer's

instructions (Qiagen, Toronto, ON, Canada). Briefly, 0.5 μg of isolated total RNA from siRNA

transfected and vehicle or Ang II treated cardiac fibroblasts was subject to genomic DNA

elimination using mixes supplied with the array kit for 5 mins at 42°C. DNA eliminated RNA

was then subject to reverse transcription reactions using Qiagen RT^2 First Strand Kits with

protocols according to the manufacturer's instructions. Qiagen RT^2 SYBR Green MasterMix was

added to the cDNA and subsequently dispensed in wells of a 96-well plate containing pre-loaded

lyophilized primers provided by the manufacturer. Quantitative PCR reactions were then run on

an Applied Biosystems ViiA 7 thermocycler according to the manufacturers cycle

recommendations. Each sample was run on separate individual 96 well plates and Ct values for

each gene assessed were collected and analyzed; Ct values greater than 35 were eliminated from

the overall analysis. Expression data was normalized to levels of two housekeeping genes

contained on each plate – Ldha1 and Hprt.

AAV Production and transduction of cardiac fibroblasts – FLAG-Gβ_1 and FLAG-Gβ_2 were

PCR amplified from a pcDNA3.1+ plasmid and BamHI and EcoRI restrictions sites added to the

5' and 3' end, respectively. These restrictions sites were used to insert each FLAG-Gβ into the

pAAV-CAG plasmid. Adeno-associated viruses were produced as previously described (Burger

and Nash, 2016). Cells were transduced with AAV1-FLAG-Gβ_1 (MOI of 10^3 or 5×10^4) in

DMEM low glucose for 6h. Additional media was added to obtain a final 7% FBS concentration

and incubated for another 24 h. At this point, the cells were detached with trypsin/EDTA and

plated as described for respective experiments.

ChIP-qPCR – Immunoprecipitation in cardiac fibroblasts was performed as previously

described, with minor modifications (Bolli et al., 2013). Isolated nuclei were sonicated with a

Diagenode BioRuptor^{TM} UCD-200 (18 cycles, 30 s on/off, high power) to shear chromatin.

FLAG-Gβ_1 immunoprecipitation was performed with 10 μg sheared rat chromatin alongside 5

μg of *Schizosaccharomyces pombe* yeast chromatin, obtained as previously described (Mbogning

and Tanny, 2017). Chromatin was immunoprecipitated with an anti-FLAG M2 antibody (2 μg)

or equivalent amount of rabbit IgG alongside an anti-*Schizosaccharomyces pombe* H2B

antibody. RNAPII immunoprecipitation was performed with 20 μg of sheared rat chromatin

alongside 0.2 μg of *Schizosaccharomyces pombe* yeast chromatin. Chromatin was

immunoprecipitated with an anti-Rpb1 (8WG16) antibody. Localization was assessed by qPCR

with primers for specific genomic loci (**Supplemental Table 2**). All qPCR reactions were

performed using a Bio-Rad 1000 Series Thermal Cycling CFX96 Optical Reaction module and

iQ SYBR Green Supermix. Data analysis included subtracting the % Input of IgG control for

each treatment from the respective IP, followed by normalization to the % Input yeast $cdc2^+$ of

each FLAG IP or $act1^+$ for each RNAPII IP to account for differences in IP efficiencies.

ChIP-seq immunoprecipitation and data analysis - Immunoprecipitation in Cardiac

fibroblasts was performed as previously described, with minor modifications (Bolli et al., 2013).

Isolated nuclei were sonicated with a Diagenode BioRuptor™ UCD-200 (18 cycles, 30 s on/off,

high power) to shear chromatin. FLAG-Gβ_1 immunoprecipitation was performed with 40 μg of

sheared rat chromatin alongside 0.4 μg of chromatin from a *S. pombe* strain expressing FLAG-

Bdf2. RNAPII immunoprecipitation was performed using 20 μg sheared rat chromatin alongside

0.2 μg wild-type *S. pombe* chromatin. Chromatin was immunoprecipitated with an anti-FLAG

M2 antibody (2 μg) or anti-Rpb1 (8WG16) antibody (2 μg). Two biological replicates of FLAG-

Gβ_1 immunoprecipitation and three biological replicates of Rpb1 immunoprecipitation were

included. Following immunoprecipitation and DNA cleanup, libraries were prepared with the

NEBNext® Ultra™ II DNA Library Prep kit for Illumina and 50 bp single end reads obtained

with an Illumina HiSeq 4000 at the McGill University and Génome Québec Innovation Centre,

Montréal, Canada.

Reads were trimmed with TrimGalore (0.6.0) (Krueger, Martin, 2011) using the

following settings*: --phred33 --length 36 -q 5 --stringency 1 -e 0.1. A Bowtie2 genome

comprised of the Ensembl rat reference genome (Rattus.norvegicus.Rnor.6.0.94) (Zerbino et al.,

2018) and *S. pombe* reference genome (Schizosaccharomyces_pombe.ASM294v2) was built

with the bowtie2-build function. Processed reads were aligned to the custom combined rat and *S.*

pombe genome with Bowtie2 (v2.3.5), followed by removal of low-quality mapped reads

(MAPQ < 10) and reads mapped to non-standard chromosomes with SAMtools (v1.9) (Li et al.,

2009). Duplicate reads were removed with Picard tools (v2.20.6, Broad Institute). Aligned reads

were separated into individual files for the rat or *S. pombe* genome respectively. For RNAPII

ChIP, peaks were called using macs2 (v2.1.1) with settings --broad and --broad-cutoff 0.1

(Zhang et al., 2008), reads extended by the fragment length determined by

phantompeakqualtools (v1.14) (Landt et al., 2012, Kharchenko et al., 2008), a scaling factor

estimated using the NCIS R package (Liang and Keles, 2012) and potential misassembled

regions of the rat genome blacklisted (Ramdas et al., 2019). RNAPII peaks were annotated with

HOMER (v4.11) (Heinz et al., 2010) and those protein-coding genes with RNAPII peaks in two

of three replicates in any treatment were used for subsequent analysis. BAM files of treatment

replicates were combined, input reads subtracted with the deepTools (Ramirez et al., 2014)

function bamCompare (--scaleFactorsMethod SES) and negative values set to 0. Lastly, values

were converted to counts per million mapped reads with library size adjusted by the total number

of reads aligned to the *S. pombe* genome. K-means clustering for genes with identified RNAPII

peaks and data visualization was performed with the deepTool's computeMatrix and plotProfile

functions. Gene ontology enrichment was performed using the R package topGO (v2.36.0).

Statistical Analysis – Statistical tests were performed using GraphPad Prism 8.0 software. For

quantifications of immunoprecipitation experiments, two-way analysis of variance (ANOVA)

followed by post-hoc Dunnett's test was used on quantifications of western blot bands, with all

multiple comparisons being made to vehicle-vehicle conditions. To analyse Ca^{2+} release

experiments, the dependent measure was the area under the curve (AUC), computed from

release-time data sets. AUC data were subjected to ANOVA and Dunnett's tests, using as a point

of comparison the siRNA control condition. For the FLAG-Gβ and RNAPII interaction in HEK

293F cells, one-sample t-tests were performed with a Bonferroni correction. Summary gene

expression of the fibrosis array qPCR was compared with a two-way ANOVA followed by post-

hoc t-tests with a Bonferroni correction. Individual gene expression from the fibrosis qPCR array

and Ctgf gene expression with in-house primers was assessed with a two-way ANOVA followed

by Bonferroni corrected post-hoc t-tests at individual time points. For validation of $G\beta_1$ and $G\beta_2$

knockdown in cardiac fibroblasts, fold changes over siRNA control were compared to siRNA

control using paired Student's t-tests. For FLAG-$G\beta_1$ ChIP-qPCR, independent paired Student t-

tests with a Bonferroni post-hoc correction were performed. A Fisher's exact test was used to

compare the proportion of cluster 1 genes in the list of genes with RNAPII peaks following Ang

II treatment in control or $G\beta_1$ knockdown conditions and to assess GO Term enrichment in

cluster 1 genes. Alpha was set at $p<0.05$ (2-tailed). All results are expressed as mean ± S.E.M,

and data are represented as pooled experiments whose sample sizes are indicated in figure

legends.

FIGURE LEGENDS

Figure 1. Characterization of Gβγ-RNAPII in rat neonatal cardiac fibroblasts. **(A)** Time

course of the Ang II-stimulated interaction between Gβγ and Rpb1. The ratio of Rpb1 co-

immunoprecipitated with $G\beta_{1-4}$ upon treatment of 1 μM Ang II treatment at the indicated

timepoints in cardiac fibroblasts was assessed. **(B)** Densitometry-based quantification of panel A

was used to determine the ratio of Rpb1 to $G\beta_{1-4}$ immunoprecipitated at each time point. The fold

change over the 0 min time point was then calculated. Data is representative of four independent

experiments. **(C)** Effect of AT1R antagonist losartan pre-treatment on the Ang II-mediated

interaction, demonstrating angiotensin receptor subtype selectivity. **(D)** Densitometry-based

quantification of AT1R antagonist effect on Ang II-induced interaction. The ratio of Rpb1

immunoprecipitated with $G\beta_{1-4}$ was determined for each condition and fold change over

DMSO/DMEM was calculated. Data is representative of three independent experiments. **(E)**

Assessment of the necessity of Gβγ import into the nucleus for interaction to occur upon AT1R

stimulation with Ang II. Cardiac fibroblasts were pretreated for 1 h with importazole prior to

Ang II stimulation. Data are representative of four independent experiments. **(F)** Densitometry-

based quantification of the Ang II induced interaction and the effect of nuclear import inhibition.

The ratio of Rpb1 to $G\beta_{1-4}$ immunoprecipitated was determined and normalized to fold change

over DMSO/DMEM treatment. In all panels, data represents mean ± S.E.M, * indicates p<0.05,

** indicates p<0.01.

Figure 2. Mechanistic analysis of Gβγ interactions with Rpb1 in rat neonatal cardiac

fibroblasts. **(A-H)** Assessment of the effect of inhibition of signalling molecules and effectors

implicated in AT1R signalling on the induction of the Gβγ-RNAPII interaction in cardiac

fibroblasts. Concentrations of inhibitors and lengths of pre-treatment are indicated in each panel.

In all experiments, Ang II treatment was applied at a concentration of 1 μM for 75 min in order

to induce the interaction. Data shown is representative of between 3 and 6 independent co-

immunoprecipitation and western blot experiments. Corresponding quantification analyses of

inhibitor co-IP experiments are depicted in **Supplemental Figure 3**.

Figure 3. Mechanistic analysis of carbachol-induced Gβγ interaction occurs in HEK 293

cells. **(A-G)** HEK 293 cells were starved for 10-12 hours in DMEM without FBS and were then

pre-treated with the indicated inhibitors for the indicated times. Cells were subsequently treated

with 1 mM carbachol for 45 min, and the amount of Rpb1 co-immunoprecipitated with $Gβ_{1-4}$

was assessed by western blot. Data is representative of at least 3 independent experiments. The

associated quantifications of the co-IPs are represented in **Supplemental Figure 4**.

Figure 4. Gβ subunit-specific effects on Ang II signalling and induction of Rpb1

interaction. **(A)** Assessment of the effect of Gβ subunit knockdown by siRNA on the Gβγ-

RNAPII interaction upon AT1R stimulation. Cardiac fibroblasts were transfected with siRNA

control or siRNA to knockdown $Gβ_1$ or $Gβ_2$ and were then serum-deprived overnight before

treatment with Ang II for 75 min. Cells were assessed for Gβγ-RNAPII interaction by co-

immunoprecipitation and western blots. Data represents mean ± S.E.M. of 6-7 independent

experiments. **(B)** Densitometry-based quantification of knockdown experiments in (C) were

normalized as fold change over the respective siRNA-DMEM condition; data represents mean ±

S.E.M. of six independent experiments. **(C)** Traces of calcium release upon AT1R stimulation

with Ang II at the 10 s time point, with or without knockdown of either $Gβ_1$ or $Gβ_2$. Data

represents mean ± S.E.M. of fluorescence ratios of 340/516 emission readings to 360/516

emissions readings normalized to basal ratios of three independent experiments. **(D)** Area under

the curve analysis of the data obtained in panel A. * indicates p<0.05.

Figure 5. Requirement of RNAPII transcription for Gβγ-RNAPII interaction in rat

neonatal cardiac fibroblasts. Effect of Cdk7 inhibition with THZ1 **(A)** or Cdk9 inhibition with

iCdk9 **(B)** on Ang II-induced Gβγ-RNAPII interaction. Length of inhibitor pre-treatment is

indicated in each respective panel, and the extent of Gβγ-RNAPII interaction was assessed by

co-immunoprecipitation coupled to western blot analysis. Data is representative of three

independent experiments. Corresponding quantification analyses of inhibitor co-

immunoprecipitation experiments are depicted in **Supplemental Figure 7**. Cumulative $\log_2$(Fold

Change) of all genes detected by qPCR-based fibrosis array following treatment with 1 μM Ang

II lasting either 75 min **(C)** or 24 h **(D and E)**. Cardiac fibroblasts were transfected with 50 nM

of the indicated siRNA, and were serum-deprived for 12 h before Ang II treatment for the

indicated times. Cardiac fibroblasts were pre-treated for 30 min with 10 μM gallein prior to Ang

II. Ct values were normalized to the housekeeping genes Hprt1 and Ldha and the $\log_2$(fold

change) over control was determined. For each gene, the average $\log_2$(fold change) across three

independent experiments was plotted. *** indicates p<0.001, **** indicates p<0.0001.

Figure 6. **ChIP-seq for FLAG-Gβ1 and Rpb1 following 75 min Ang II treatment in cardiac**

fibroblasts. Cardiac fibroblasts were transduced with AAV1-FLAG-Gβ$_1$ or transfected with the

indicated siRNA followed by Ang II treatment (1 μM for 75 min). **(A)** Comparison of genes with

annotated RNAPII peaks following Ang II treatment and siRNA control or Gβ$_1$. FLAG-Gβ$_1$ **(B)**

or the Rpb1 subunit of RNAPII **(C)** were immunoprecipitated from crosslinked and sonicated

chromatin, followed by DNA purification and next-generation sequencing. Reads were

normalized to an exogenous *S. pombe* chromatin spike-in. Genes with a RNAPII peak annotated

by HOMER in two of the three replicates were used to identify two K-means clusters. **(D)** Top

four significant GO terms enriched in cluster 1. Individual FLAG-Gβ_1 or RNAPII tracks for two

genes from cluster 1, **(E)** Thbs1 or **(F)** Ctgf.

Supplemental Figure 1. Induction of the Gβγ-RNAPII interaction in HEK 293 cells. **(A)**

Time-course analysis of the induction of the Gβγ-RNAPII interaction. The amount of Rpb1 co-

immunoprecipitated with Gβ_{1-4} from HEK 293 cells treated for the indicated times with 1 mM

carbachol was assessed by western blot for each time point. Data is representative of three

independent experiments. **(B)** Quantification of Gβγ-RNAPII time-course co-

immunoprecipitation. Densitometry-based analysis of bands corresponding to Rpb1 at each

timepoint was normalized to the band intensity of the amount of Gβ_{1-4} immunoprecipitated to

yield ratios of Rpb1 pulled down with Gβ_{1-4}. **(C)** Assessing the Gβγ and Rpb1 interaction by

immunoprecipitation of Rpb1 with two different antibodies. Western blots are representative of

at least two independent experiments. Immunoprecipitation experiments demonstrating that

carbachol treatment does not induce interaction of Rpb1 with **(D)** G$\alpha_{q/11}$ nor **(E)** ERK1/2 in HEK

293 cells, and also does not alter the amount of G$\alpha_{q/11}$ or ERK1/2 interacting with Gβγ under

such conditions. **(F)** Assessment of interaction between Gβ_{1-4} and Rpa194, the largest subunit of

RNA polymerase I. Data represents analysis of a time course experiment western blot performed

as in **Supplemental Figure 2A**. Data represents mean ± S.E.M; * indicates $p<0.05$, ** indicates

$p<0.01$.

Supplemental Figure 2. Requirement for Gβγ nuclear transport for RNAPII interaction in HEK 293 cells. **(A**) Representative experiment assessing the requirement of importin-β inhibition on the Gβγ-RNAPII interaction by sub-cellular fractionation and co-immunoprecipitation. **(B)** Densitometry-based quantification of the carbachol induced interaction and the effect of nuclear import inhibition on interaction induction. Data represents mean ± S.E.M. of three independent experiments for black bars, and two independent experiments for white bars (nuclear import inhibition conditions).

Supplemental Figure 3. Quantitative analysis of the effects of inhibition of signalling molecules downstream of AT1R activation. **(A-H)** The relative quantities of Rpb1 co-immunoprecipitated with Gβ$_{1-4}$ under different conditions depicted in **Figure 2** were quantified using ImageJ and normalized to DMSO/DMEM control conditions. Data shown is representative of (A) 3, (B) 6, (C) 3, (D) 4, (E) 5, (F) 5, (G) 4 or (H) 3 independent co-immunoprecipitation and western blot experiments. Data is represented as fold change over respective controls and error bars represent S.E.M. * indicates $p < 0.05$, ** indicates $p < 0.01$.

Supplemental Figure 4. Quantitative analysis of the effects of inhibition of signalling molecules downstream of M3-mAChR activation in HEK 293 cells. **(A-G)** The relative quantities of Rpb1 co-immunoprecipitated with Gβ$_{1-4}$ under different conditions depicted in **Figure 3** were quantified using ImageJ and were normalized to amounts pulled down in DMSO/DMEM control conditions. Data shown is representative of (A) 3, (B) 4, (C) 5, (D) 3, (E) 3, (F) 6 or (G) 4 independent co-immunoprecipitation and western blot experiments. Data is

represented as fold change over respective controls and error bars represent S.E.M. * indicates

p<0.05.

Supplemental Figure 5. Assessment of specific Gβ subunits interacting with RNAPII upon

agonist stimulation in rat cardiac fibroblasts or in HEK 293 cells. **(A)** Transcript levels for

GNB1, GNB2, GNB3, and GNB4 were assessed in cardiac fibroblasts by RT-qPCR. The Ct

values for each gene transcript were normalized to the house keeping U6 snRNA gene transcript

for comparison. Data represents mean ± S.E.M for 3-4 independent experiments. **(B)** Gβ$_1$ and

Gβ$_2$ were immunoprecipitated with isoform specific antibodies from cardiac fibroblasts lysates

treated with 1 μM Ang II for 75 min. The amount of Rpb1 pulled down with either Gβ isoform

was assessed by western blot. **(C)** Assessment of specific FLAG-tagged Gβ isoforms interaction

with Rpb1 under conditions of M3-mAChR stimulation with carbachol in HEK 293 cells. The

amount of Rpb1 interacting with each Gβ isoform was assessed by western blot following FLAG

immunoprecipitation. **(D)** Densitometry-based quantification of the ratio of Rpb1 co-

immunoprecipitated with the indicated FLAG-tagged Gβ subunit. The ratio of Rpb1 to FLAG-

Gβ$_x$ immunoprecipitated was determined and normalized to fold change over DMEM treatment.

Data represents mean ± S.E.M for four independent replicates; * indicates p<0.01

Supplemental Figure 6. Validation of RNAi knockdown of Gβ$_1$ and Gβ$_2$. Validation of Gβ$_1$

and Gβ$_2$ mRNA **(A)** and protein **(B)** knockdown with siRNA in rat neonatal cardiac fibroblasts.

Rat neonatal cardiac fibroblasts were transfected with 50 nM siRNA control, Gβ$_1$ or Gβ$_2$ for 72

hours, serum-deprived for 12 h and RNA or protein collected as described in *Methods*. Data in

(A) represents mean ± S.E.M for four independent experiments; * Ct values were normalized to

the housekeeping U6 snRNA transcript and fold change over siRNA control determined using

the $2^{-\Delta\Delta Ct}$. ** indicates $p<0.001$ and **** indicates $p<0.0001$.

Supplemental Figure 7. Quantitative analysis of the effect of transcriptional regulator

inhibition on the Gβγ-RNAPII interaction in cardiac fibroblasts. **(A-B)** The relative

quantities of Rpb1 co-immunoprecipitated with $G\beta_{1-4}$ under conditions depicted in Figure 5A

(THZ1) and B (iCdk9) were quantified and normalized to DMSO/DMEM control conditions.

Data shown is representative of between three to six independent co-immunoprecipitation and

western blot experiments. Data represents mean ± S.E.M. * indicates $p<0.05$.

Supplemental Figure 8. Validation of heterologously expressed FLAG-tagged Gβ₁ in rat

neonatal cardiac fibroblasts. (A) Assessment of Rpb1 co-immunoprecipitated with $FLAG-G\beta_1$

following 75 min treatment of 1 μM Ang II in rat neonatal cardiac fibroblasts. Cardiac

fibroblasts were transduced with $AAV1-FLAG-G\beta_1$ prior to treatment with 1 μM Ang II. **(B)**

Densitometry-based quantification of the ratio of Rpb1 co-immunoprecipitated with $FLAG-G\beta_1$.

The ratio of Rpb1 to $FLAG-G\beta_1$ was calculated and normalized as fold change over DMEM

condition. Data is represented as mean ± S.E.M for four independent experiments. Assessing

changes in $FLAG-G\beta_1$ **(C)** or **(D)** Rpb1 occupancy along Ctgf following 75 min treatment with 1

μM Ang II. $FLAG-G\beta_1$ or Rpb1 was immunoprecipitated from crosslinked and sonicated

chromatin, DNA purified and quantified by qPCR. Data is represented as mean ± S.E.M for 4-6

independent experiments, * indicates $p<0.05$. **(E)** Validation of Ctgf gene expression with

primers distinct from those used in the Qiagen RT^2 Profiler[TM] PCR array. Data represents mean

± S.E.M for four independent experiments, * indicates $p<0.05$.

Supplemental Figure 9. Schema summarizing signalling events regulating the agonist

induced Gβγ interaction with RNAPII. Signalling cascade downstream of AT1R in cardiac

fibroblasts or M3 muscarinic receptors in HEK 293F cells regulating the interaction. Signalling

pathways were determined by assessing Gβγ-RNAPII interactions by co-immunoprecipitation

and western blot as shown in **Figure 2** and **Supplemental Figure 3** for cardiac fibroblasts and

Figure 3 and **Supplemental Figure 4** for HEK 293F cells.

Tables

Supplemental Table 1. Summary of fibrosis RT-qPCR array results. This table summarizes

gene expression changes measured using the Qiagen RT^2 Profiler^{TM} PCR Array at 75 min and 24

h Ang II stimulation. Genes were considered to have altered expression with fold changes ≥ 1.5

or ≤ 0.5 compared to DMEM/siRNA control conditions at the respective time point. In

parenthesis are the number of genes with a significant ($p<0.05$) change in expression compared

to DMEM/siRNA control at the respective time point. Two-way ANOVA followed by post-hoc

t-test comparisons with Bonferroni correction was performed for each gene individually. Data is

representative of three independent biological replicates.

Time	siRNA	Treatment	Upregulated	Downregulated
75 min	Control	DMEM	0	0
		1 µM Ang II	7(5)	0
	Gnb1	DMEM	4(1)	1
		1 µM Ang II	10(6)	2
24 h	Control	DMEM	0	0
		1 µM Ang II	37(7)	0
	Gnb1	DMEM	26(2)	1
		1 µM Ang II	53(13)	0
	Gnb2	DMEM	7	1
		1 µM Ang II	44(11)	0

858 **Supplemental Table 2. List of primers used to assess gene expression by RT-qPCR and**

859 **ChIP-qPCR in cardiac fibroblasts**. Forward and reverse primers were used at a concentration

860 of 300 nM for each qPCR reaction. Primer sequences were designed using NCBI's Primer-

861 BLAST tool and validated by analysis of standard curve qPCR assays performed in-house.

Target	Forward (5' -> 3')	Reverse (5' -> 3')
U6 snRNA	TGGAACGATACAGAGAAGATTAG	GAATTTGCGTGTCATCCTTG
$G\beta_1$	CTCATGACCTACTCCCATGA	TCAGCTTTGAGTGCATCC
$G\beta_2$	CAGCTACACCACTAACAAGG	CTCTCGGGTCTTGAGACTAT
$G\beta_3$	CTCCTTAGGGTCAGTCTTCTAT	AAAGGCACACTCCCATAATC
$G\beta_4$	GGTGGTCAAAGAAACAATCAAG	GTCTGTCGGGATAGGGATAA
Ctgf	TGCATCCTCCTACCGCGTCC	GAGGCTGATGGGACCTGCGA
Ctgf TSS	CAGACCCACTCCAGCTCCGA	GTGGCTCCTGGGGTTGTCCA
Ctgf Exon	TCAAGCTGCCCGGGAAATGC	GCGGTCCTTGGGCTCATCAC
Ctgf 3' End	AATGGCTTGCTCAGGGTAACTGG	AACTGCCTCCCAAACCAGTCATAG
$cdc2^+$	ATCATTCTCGCATCTCTATTA	ATTCTCCATTGCAAACCACTA
$act1^+$	GGTTGCTCAATGTTATCCGTTTC	TGATAAAGCCACACACAGCGTTA

Acknowledgements

This work was supported by a grant from the Heart and Stroke Foundation of Canada (G-

15-0008938) to T.E.H and J.C.T, a grant from Canadian Institute of Health Science (CIHR)

(MOP 130362) to J.C.T. and a grant from CIHR (PJT-159687) to T.E.H. R.M. and S.M.K were

supported by studentships from the McGill-CIHR Drug Development Training Program and the

McGill Faculty of Medicine. We thank Dr. Ron Taussig (UT Southwestern), Dr. Nathanael S.

Gray, and Novartis for providing materials instrumental to this study. Lastly, we thank all the

members of the Hébert and Tanny labs for discussion and critical reading of the manuscript.

Author Contributions

S.M.K, R.D.M, J.C.T, and T.E.H designed the experiments and wrote the manuscript. S.M.K,

R.D.M, S.G, C.B, J.J.T, A.Z, S.M, and P.T conducted experiments. S.M.K, R.DM and C.B

analyzed experiments. R.D.M, J.J.T, J.C.T and T.E.H edited the manuscript.

Declaration of Interests

None

References

AHMED, M. S., ØIE, E., VINGE, L. E., YNDESTAD, A., ØYSTEIN ANDERSEN, G., ANDERSSON, Y., ATTRAMADAL, T. & ATTRAMADAL, H. 2004. Connective tissue growth factor—a novel mediator of angiotensin II-stimulated cardiac fibroblast activation in heart failure in rats. *Journal of Molecular and Cellular Cardiology,* 36**,** 393-404.

ANAND, P., BROWN, J. D., LIN, C. Y., QI, J., ZHANG, R., ARTERO, P. C., ALAITI, M. A., BULLARD, J., ALAZEM, K., MARGULIES, K. B., CAPPOLA, T. P., LEMIEUX, M., PLUTZKY, J., BRADNER, J. E. & HALDAR, S. M. 2013. BET bromodomains mediate transcriptional pause release in heart failure. *Cell,* 154**,** 569-82.

BHATNAGAR, A., UNAL, H., JAGANNATHAN, R., KAVETI, S., DUAN, Z. H., YONG, S., VASANJI, A., KINTER, M., DESNOYER, R. & KARNIK, S. S. 2013. Interaction of G-protein βγ complex with chromatin modulates GPCR-dependent gene regulation. *PLoS One,* 8**,** e52689.

BOLLI, P., VARDABASSO, C., BERNSTEIN, E. & CHAUDHRY, H. W. 2013. Chromatin immunoprecipitation of adult murine cardiomyocytes. *Curr Protoc Cell Biol,* Chapter 17**,** Unit17 14.

BURGER, C. & NASH, K. R. 2016. Small-Scale Recombinant Adeno-Associated Virus Purification. *Methods Mol Biol,* 1382**,** 95-106.

CALDERONE, A., THAIK, C. M., TAKAHASHI, N., CHANG, D. L. & COLUCCI, W. S. 1998. Nitric oxide, atrial natriuretic peptide, and cyclic GMP inhibit the growth-promoting effects of norepinephrine in cardiac myocytes and fibroblasts. *J Clin Invest,* 101**,** 812-8.

CAMPBELL, S. E. & KATWA, L. C. 1997. Angiotensin II stimulated expression of transforming growth factor-β1 in cardiac fibroblasts and myofibroblasts. *J Mol Cell Cardiol,* 29**,** 1947-58.

CAMPDEN, R., AUDET, N. & HÉBERT, T. E. 2015a. Nuclear G Protein Signaling: New Tricks for Old Dogs. *Journal of Cardiovascular Pharmacology,* 65**,** 110-122.

CAMPDEN, R., PÉTRIN, D., ROBITAILLE, M., AUDET, N., GORA, S., ANGERS, S. & HÉBERT, T. 2015b. Tandem Affinity Purification to Identify Cytosolic and Nuclear Gβγ-Interacting Proteins. *In:* ALLEN, B. G. & HÉBERT, T. E. (eds.) *Nuclear G-Protein Coupled Receptors.* Springer New York.

CHENG, T.-H., CHENG, P.-Y., SHIH, N.-L., CHEN, I.-B., WANG, D. L. & CHEN, J.-J. 2003. Involvement of reactive oxygen species in angiotensin II-induced endothelin-1 gene expression in rat cardiac fibroblasts. *Journal of the American College of Cardiology,* 42**,** 1845-1854.

CHINTALGATTU, V. & KATWA, L. C. 2009. Role of protein kinase C-δ in angiotensin II induced cardiac fibrosis. *Biochemical and Biophysical Research Communications,* 386**,** 612-616.

DANG, M. Q., ZHAO, X. C., LAI, S., WANG, X., WANG, L., ZHANG, Y. L., LIU, Y., YU, X. H., LIU, Y., LI, H. H. & XIA, Y. L. 2015. Gene expression profile in the early stage of angiotensin II-induced cardiac remodeling: a time series microarray study in a mouse model. *Cell Physiol Biochem,* 35**,** 467-76.

DEVOST, D., SLENO, R., PETRIN, D., ZHANG, A., SHINJO, Y., OKDE, R., AOKI, J., INOUE, A. & HÉBERT, T. E. 2017. Conformational Profiling of the AT1 Angiotensin II Receptor Reflects Biased Agonism, G Protein Coupling, and Cellular Context. *J Biol Chem,* 292**,** 5443-5456.

DOBACZEWSKI, M., BUJAK, M., LI, N., GONZALEZ-QUESADA, C., MENDOZA, L. H., WANG, X. F. &
FRANGOGIANNIS, N. G. 2010. Smad3 signaling critically regulates fibroblast phenotype
and function in healing myocardial infarction. *Circ Res,* 107**,** 418-28.
DUAN, Q., MCMAHON, S., ANAND, P., SHAH, H., THOMAS, S., SALUNGA, H. T., HUANG, Y.,
ZHANG, R., SAHADEVAN, A., LEMIEUX, M. E., BROWN, J. D., SRIVASTAVA, D., BRADNER,
J. E., MCKINSEY, T. A. & HALDAR, S. M. 2017. BET bromodomain inhibition suppresses
innate inflammatory and profibrotic transcriptional networks in heart failure. *Sci Transl*
Med, 9.
DUPRÉ DJ, R. M., REBOIS RV, HÉBERT TE 2009. The role of Gβγ subunits in the organization,
assembly and function of GPCR signaling complexes. *Annu. Rev. Pharmacol. Toxicol.,* 49**,**
31-56.
FU, X., KHALIL, H., KANISICAK, O., BOYER, J. G., VAGNOZZI, R. J., MALIKEN, B. D., SARGENT, M.
A., PRASAD, V., VALIENTE-ALANDI, I., BLAXALL, B. C. & MOLKENTIN, J. D. 2018.
Specialized fibroblast differentiated states underlie scar formation in the infarcted
mouse heart. *J Clin Invest,* 128**,** 2127-2143.
GAO, X., HE, X., LUO, B., PENG, L., LIN, J. & ZUO, Z. 2009. Angiotensin II increases collagen I
expression via transforming growth factor-β1 and extracellular signal-regulated kinase in
cardiac fibroblasts. *European Journal of Pharmacology,* 606**,** 115-120.
GREENWOOD, I. A. & STOTT, J. B. 2019. The Gβ1 and Gβ3 Subunits Differentially Regulate Rat
Vascular Kv7 Channels. *Front Physiol,* 10**,** 1573.
HEINZ, S., BENNER, C., SPANN, N., BERTOLINO, E., LIN, Y. C., LASLO, P., CHENG, J. X., MURRE, C.,
SINGH, H. & GLASS, C. K. 2010. Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. *Mol Cell,* 38**,** 576-89.
KAMAL, F. A., SMRCKA, A. V. & BLAXALL, B. C. 2011. Taking the heart failure battle inside the
cell: small molecule targeting of Gβγ subunits. *J Mol Cell Cardiol,* 51**,** 462-7.
KAMAL, F. A., TRAVERS, J. G., SCHAFER, A. E., MA, Q., DEVARAJAN, P. & BLAXALL, B. C. 2017. G
Protein-Coupled Receptor-G-Protein βγ-Subunit Signaling Mediates Renal Dysfunction
and Fibrosis in Heart Failure. *J Am Soc Nephrol,* 28**,** 197-208.
KAWANO, H., DO, Y. S., KAWANO, Y., STARNES, V., BARR, M., LAW, R. E. & HSUEH, W. A. 2000.
Angiotensin II has multiple profibrotic effects in human cardiac fibroblasts. *Circulation,*
101**,** 1130-7.
KHAN, S. M., MIN, A., GORA, S., HOURANIEH, G. M., CAMPDEN, R., ROBITAILLE, M., TRIEU, P.,
PÉTRIN, D., JACOBI, A. M., BEHLKE, M. A., ANGERS, S. & HÉBERT, T. E. 2015. Gβ4γ1 as a
modulator of M3 muscarinic receptor signalling and novel roles of Gβ1 subunits in the
modulation of cellular signalling. *Cellular Signalling,* 27**,** 1597-1608.
KHAN, S. M., SLENO, R., GORA, S., ZYLBERGOLD, P., LAVERDURE, J.-P., LABBÉ, J.-C., MILLER, G. J.
& HÉBERT, T. E. 2013. The Expanding Roles of Gβγ Subunits in G Protein–Coupled
Receptor Signaling and Drug Action. *Pharmacological Reviews,* 65**,** 545-577.
KHARCHENKO, P. V., TOLSTORUKOV, M. Y. & PARK, P. J. 2008. Design and analysis of ChIP-seq
experiments for DNA-binding proteins. *Nat Biotechnol,* 26**,** 1351-9.
KRUEGER, F. *Trim Galore!* [Online]. Available:
http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/ [Accessed].

KWIATKOWSKI, N., ZHANG, T., RAHL, P. B., ABRAHAM, B. J., REDDY, J., FICARRO, S. B., DASTUR, A., AMZALLAG, A., RAMASWAMY, S., TESAR, B., JENKINS, C. E., HANNETT, N. M., MCMILLIN, D., SANDA, T., SIM, T., KIM, N. D., LOOK, T., MITSIADES, C. S., WENG, A. P., BROWN, J. R., BENES, C. H., MARTO, J. A., YOUNG, R. A. & GRAY, N. S. 2014. Targeting transcription regulation in cancer with a covalent CDK7 inhibitor. *Nature,* 511**,** 616-20.

LANDT, S. G., MARINOV, G. K., KUNDAJE, A., KHERADPOUR, P., PAULI, F., BATZOGLOU, S., BERNSTEIN, B. E., BICKEL, P., BROWN, J. B., CAYTING, P., CHEN, Y., DESALVO, G., EPSTEIN, C., FISHER-AYLOR, K. I., EUSKIRCHEN, G., GERSTEIN, M., GERTZ, J., HARTEMINK, A. J., HOFFMAN, M. M., IYER, V. R., JUNG, Y. L., KARMAKAR, S., KELLIS, M., KHARCHENKO, P. V., LI, Q., LIU, T., LIU, X. S., MA, L., MILOSAVLJEVIC, A., MYERS, R. M., PARK, P. J., PAZIN, M. J., PERRY, M. D., RAHA, D., REDDY, T. E., ROZOWSKY, J., SHORESH, N., SIDOW, A., SLATTERY, M., STAMATOYANNOPOULOS, J. A., TOLSTORUKOV, M. Y., WHITE, K. P., XI, S., FARNHAM, P. J., LIEB, J. D., WOLD, B. J. & SNYDER, M. 2012. ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia. *Genome Res,* 22**,** 1813-31.

LEE, A. A., DILLMANN, W. H., MCCULLOCH, A. D. & VILLARREAL, F. J. 1995. Angiotensin II stimulates the autocrine production of transforming growth factor-β 1 in adult rat cardiac fibroblasts. *J Mol Cell Cardiol,* 27**,** 2347-57.

LEHMANN, D. M., SENEVIRATNE, A. M. & SMRCKA, A. V. 2008. Small molecule disruption of G protein βγ subunit signaling inhibits neutrophil chemotaxis and inflammation. *Mol Pharmacol,* 73**,** 410-8.

LI, H., HANDSAKER, B., WYSOKER, A., FENNELL, T., RUAN, J., HOMER, N., MARTH, G., ABECASIS, G., DURBIN, R. & GENOME PROJECT DATA PROCESSING, S. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics,* 25**,** 2078-9.

LIANG, K. & KELES, S. 2012. Normalization of ChIP-seq data with control. *BMC Bioinformatics,* 13**,** 199.

LIN, Y. & SMRCKA, A. V. 2011. Understanding molecular recognition by G protein βγ subunits on the path to pharmacological targeting. *Mol Pharmacol,* 80**,** 551-7.

LIU, X., KRAUS, W. L. & BAI, X. 2015. Ready, pause, go: regulation of RNA polymerase II pausing and release by cellular signaling pathways. *Trends Biochem Sci,* 40**,** 516-25.

LU, H., XUE, Y., YU, G. K., ARIAS, C., LIN, J., FONG, S., FAURE, M., WEISBURD, B., JI, X., MERCIER, A., SUTTON, J., LUO, K., GAO, Z. & ZHOU, Q. 2015. Compensatory induction of MYC expression by sustained CDK9 inhibition via a BRD4-dependent mechanism. *Elife,* 4**,** e06535.

MA, Z. G., YUAN, Y. P., WU, H. M., ZHANG, X. & TANG, Q. Z. 2018. Cardiac fibrosis: new insights into the pathogenesis. *Int J Biol Sci,* 14**,** 1645-1657.

MARTIN, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *2011,* 17**,** 3.

MBOGNING, J. & TANNY, J. C. 2017. Chromatin Immunoprecipitation of Histone Modifications in Fission Yeast. *Methods Mol Biol,* 1528**,** 199-210.

MURPHY, A. M., WONG, A. L. & BEZUHLY, M. 2015. Modulation of angiotensin II signaling in the prevention of fibrosis. *Fibrogenesis Tissue Repair,* 8**,** 7.

NAMKUNG, Y., LEGOUILL, C., KUMAR, S., CAO, Y., TEIXEIRA, L. B., LUKASHEVA, V., GIUBILARO, J., SIMOES, S. C., LONGPRE, J. M., DEVOST, D., HÉBERT, T. E., PINEYRO, G., LEDUC, R.,

COSTA-NETO, C. M., BOUVIER, M. & LAPORTE, S. A. 2018. Functional selectivity profiling
of the angiotensin II type 1 receptor using pathway-wide BRET signaling sensors. *Sci*
Signal, 11.
OLSON, E. R., SHAMHART, P. E., NAUGLE, J. E. & MESZAROS, J. G. 2008. Angiotensin II-Induced
Extracellular Signal-Regulated Kinase 1/2 Activation Is Mediated by Protein Kinase Cδ
and Intracellular Calcium in Adult Rat Cardiac Fibroblasts. *Hypertension,* 51**,** 704-711.
PARK, D., JHON, D. Y., LEE, C. W., LEE, K. H. & RHEE, S. G. 1993. Activation of phospholipase C
isozymes by G protein βγ subunits. *J Biol Chem,* 268**,** 4573-6.
PARK, J. G., MUISE, A., HE, G. P., KIM, S. W. & RO, H. S. 1999. Transcriptional regulation by the
γ5 subunit of a heterotrimeric G protein during adipogenesis. *Embo J,* 18**,** 4004-12.
PASSIER, R. C., SMITS, J. F., VERLUYTEN, M. J. & DAEMEN, M. J. 1996. Expression and
localization of renin and angiotensinogen in rat heart after myocardial infarction. *Am J*
Physiol, 271**,** H1040-8.
RAMDAS, S., OZEL, A. B., TREUTELAAR, M. K., HOLL, K., MANDEL, M., WOODS, L. C. S. & LI, J. Z.
2019. Extended regions of suspected mis-assembly in the rat reference genome. *Sci*
Data, 6**,** 39.
RAMIREZ, F., DUNDAR, F., DIEHL, S., GRUNING, B. A. & MANKE, T. 2014. deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic Acids Res,* 42**,** W187-91.
ROBITAILLE, M., GORA, S., WANG, Y., GOUPIL, E., PETRIN, D., DEL DUCA, D., VILLENEUVE, L. R.,
ALLEN, B. G., LAPORTE, S. A., BERNARD, D. J. & HÉBERT, T. E. 2010. Gβγ is a negative
regulator of AP-1 mediated transcription. *Cell Signal,* 22**,** 1254-66.
ROSENKRANZ, S. 2004. TGF-β1 and angiotensin networking in cardiac remodeling. *Cardiovasc*
Res, 63**,** 423-32.
SANO, M., ABDELLATIF, M., OH, H., XIE, M., BAGELLA, L., GIORDANO, A., MICHAEL, L. H.,
DEMAYO, F. J. & SCHNEIDER, M. D. 2002. Activation and function of cyclin T-Cdk9
(positive transcription elongation factor-b) in cardiac muscle-cell hypertrophy. *Nat Med,*
8**,** 1310-7.
SAULIERE, A., BELLOT, M., PARIS, H., DENIS, C., FINANA, F., HANSEN, J. T., ALTIE, M. F.,
SEGUELAS, M. H., PATHAK, A., HANSEN, J. L., SENARD, J. M. & GALES, C. 2012.
Deciphering biased-agonism complexity reveals a new active AT1 receptor entity. *Nat*
Chem Biol, 8**,** 622-30.
SAYED, D., HE, M., YANG, Z., LIN, L. & ABDELLATIF, M. 2013. Transcriptional regulation patterns
revealed by high resolution chromatin immunoprecipitation during cardiac hypertrophy.
J Biol Chem, 288**,** 2546-58.
SCHRAGE, R., SCHMITZ, A. L., GAFFAL, E., ANNALA, S., KEHRAUS, S., WENZEL, D., BULLESBACH,
K. M., BALD, T., INOUE, A., SHINJO, Y., GALANDRIN, S., SHRIDHAR, N., HESSE, M.,
GRUNDMANN, M., MERTEN, N., CHARPENTIER, T. H., MARTZ, M., BUTCHER, A. J.,
SLODCZYK, T., ARMANDO, S., EFFERN, M., NAMKUNG, Y., JENKINS, L., HORN, V.,
STOSSEL, A., DARGATZ, H., TIETZE, D., IMHOF, D., GALES, C., DREWKE, C., MULLER, C. E.,
HOLZEL, M., MILLIGAN, G., TOBIN, A. B., GOMEZA, J., DOHLMAN, H. G., SONDEK, J.,
HARDEN, T. K., BOUVIER, M., LAPORTE, S. A., AOKI, J., FLEISCHMANN, B. K., MOHR, K.,
KONIG, G. M., TUTING, T. & KOSTENIS, E. 2015. The experimental power of FR900359 to
study Gq-regulated biological processes. *Nat Commun,* 6**,** 10156.

SHU, J., LIU, Z., JIN, L. & WANG, H. 2018. An RNAsequencing study identifies candidate genes for
angiotensin IIinduced cardiac remodeling. *Mol Med Rep,* 17**,** 1954-1962.
SMRCKA, A. V. 2008. G protein βγ subunits: central mediators of G protein-coupled receptor
signaling. *Cell Mol Life Sci,* 65.
SMRCKA, A. V., LEHMANN, D. M. & DESSAL, A. L. 2008. G protein βγ subunits as targets for
small molecule therapeutic development. *Comb Chem High Throughput Screen,* 11**,** 382-
95.
SPIEGELBERG, B. D. & HAMM, H. E. 2005. Gβγ binds histone deacetylase 5 (HDAC5) and inhibits
its transcriptional co-repression activity. *J Biol Chem,* 280**,** 41769-76.
STRATTON, M. S., LIN, C. Y., ANAND, P., TATMAN, P. D., FERGUSON, B. S., WICKERS, S. T.,
AMBARDEKAR, A. V., SUCHAROV, C. C., BRADNER, J. E., HALDAR, S. M. & MCKINSEY, T.
A. 2016. Signal-Dependent Recruitment of BRD4 to Cardiomyocyte Super-Enhancers Is
Suppressed by a MicroRNA. *Cell Rep,* 16**,** 1366-1378.
SUN, Y. & WEBER, K. T. 1996. Angiotensin converting enzyme and myofibroblasts during tissue
repair in the rat heart. *J Mol Cell Cardiol,* 28**,** 851-8.
TRAVERS, J. G., KAMAL, F. A., ROBBINS, J., YUTZEY, K. E. & BLAXALL, B. C. 2016. Cardiac Fibrosis:
The Fibroblast Awakens. *Circ Res,* 118**,** 1021-40.
TRAVERS, J. G., KAMAL, F. A., VALIENTE-ALANDI, I., NIEMAN, M. L., SARGENT, M. A., LORENZ, J.
N., MOLKENTIN, J. D. & BLAXALL, B. C. 2017. Pharmacological and Activated Fibroblast
Targeting of Gβγ-GRK2 After Myocardial Ischemia Attenuates Heart Failure Progression.
J Am Coll Cardiol, 70**,** 958-971.
WEBER, K. T. & DIEZ, J. 2016. Targeting the Cardiac Myofibroblast Secretome to Treat
Myocardial Fibrosis in Heart Failure. *Circ Heart Fail,* 9.
WEBER, K. T., SUN, Y., BHATTACHARYA, S. K., AHOKAS, R. A. & GERLING, I. C. 2013.
Myofibroblast-mediated mechanisms of pathological remodelling of the heart. *Nat Rev*
Cardiol, 10**,** 15-26.
YANG, J., TIAN, B. & BRASIER, A. R. 2017. Targeting Chromatin Remodeling in Inflammation and
Fibrosis. *Adv Protein Chem Struct Biol,* 107**,** 1-36.
YIM, Y. Y., BETKE, K. M., MCDONALD, W. H., GILSBACH, R., CHEN, Y., HYDE, K., WANG, Q., HEIN,
L. & HAMM, H. 2019. The in vivo specificity of synaptic Gβ and Gγ subunits to the
alpha2a adrenergic receptor at CNS synapses. *Sci Rep,* 9**,** 1718.
ZERBINO, D. R., ACHUTHAN, P., AKANNI, W., AMODE, M. R., BARRELL, D., BHAI, J., BILLIS, K.,
CUMMINS, C., GALL, A., GIRON, C. G., GIL, L., GORDON, L., HAGGERTY, L., HASKELL, E.,
HOURLIER, T., IZUOGU, O. G., JANACEK, S. H., JUETTEMANN, T., TO, J. K., LAIRD, M. R.,
LAVIDAS, I., LIU, Z., LOVELAND, J. E., MAUREL, T., MCLAREN, W., MOORE, B., MUDGE, J.,
MURPHY, D. N., NEWMAN, V., NUHN, M., OGEH, D., ONG, C. K., PARKER, A., PATRICIO,
M., RIAT, H. S., SCHUILENBURG, H., SHEPPARD, D., SPARROW, H., TAYLOR, K.,
THORMANN, A., VULLO, A., WALTS, B., ZADISSA, A., FRANKISH, A., HUNT, S. E.,
KOSTADIMA, M., LANGRIDGE, N., MARTIN, F. J., MUFFATO, M., PERRY, E., RUFFIER, M.,
STAINES, D. M., TREVANION, S. J., AKEN, B. L., CUNNINGHAM, F., YATES, A. & FLICEK, P.
2018. Ensembl 2018. *Nucleic Acids Res,* 46**,** D754-D761.
ZHANG, J. H., BARR, V. A., MO, Y., ROJKOVA, A. M., LIU, S. & SIMONDS, W. F. 2001. Nuclear
localization of G protein β 5 and regulator of G protein signaling 7 in neurons and brain.
J Biol Chem, 276**,** 10284-9.

ZHANG, Y., LIU, T., MEYER, C. A., EECKHOUTE, J., JOHNSON, D. S., BERNSTEIN, B. E., NUSBAUM,
C., MYERS, R. M., BROWN, M., LI, W. & LIU, X. S. 2008. Model-based analysis of ChIP-Seq
(MACS). *Genome Biol,* 9**,** R137.
ZHOU, Q., LI, T. & PRICE, D. H. 2012. RNA polymerase II elongation control. *Annu Rev Biochem,*
81**,** 119-43.
1104

Figure 1.

Figure 2.

Figure 3.

Figure 4.

A

	siRNA control	siRNA Gβ₁	siRNA Gβ₂	
DMEM	+ −	+ −	+ −	75 min treatment
1 μM AngII	− +	− +	− +	

IP: Gβ₁₋₄
Blot: Rpb1

IP: Gβ₁₋₄
Blot: Gβ₁₋₄

Total Lysate
Blot: Rpb1

Total Lysate
Blot: Gβ₁₋₄

B

Ratio of RNA Pol II immunoprecipitated with Gβ₁₋₄ (fold change over control)

*

siRNA control - DMEM
siRNA control - 1 μM AngII
siRNA Gβ₁ - DMEM
siRNA Gβ₁ - 1 μM AngII
siRNA Gβ₂ - DMEM
siRNA Gβ₂ - 1 μM AngII

C

Ang II Stimulation   Ang II Stimulation

Calcium Release (normalized to baseline siRNA control -Vehicle)

Time (s)

- siRNA control - Vehicle
- ▽ siRNA control - 1 μM AngII
- ■ siRNA Gβ₁ - Vehicle
- □ siRNA Gβ₁ - 1 μM AngII
- ▲ siRNA Gβ₂ - Vehicle
- □ siRNA Gβ₂ - 1 μM AngII

D

Calcium Release (Area under the curve)

*

siRNA control
siRNA Gβ₁
siRNA Gβ₂

Figure 5.

Figure 6.

A

1 μM AngII + siRNA Gβ₁   672   135   7   1 μM AngII + siRNA Control

B Flag-Gβ₁

Cluster 1   Cluster 2

— 75 min DMEM
— 75 min 1 μM Ang II

C RNAPII

Cluster 1   Cluster 2

— 75 min DMEM + siRNA Control
— 75 min AngII + siRNA Control
— 75 min DMEM + siRNA Gβ₁
— 75 min AngII + siRNA Gβ₁

D

GO ID	Term	p-value
GO:0071356	cellular response to tumor necrosis factor	3.3E-10
GO:0071347	cellular response to interleukin-1	6.2E-08
GO:0043065	positive regulation of apoptotic process	5.1E-08
GO:0044344	cellular response to fibroblast growth factor	2.2E-08

E

chr3:109,862 kb   109,866 kb   109,870 kb   109,874 kb

Flag-Gβ₁ DMEM [0-0.66]
Flag-Gβ₁ Ang II [0-0.66]
RNAPII DMEM + siRNA Control [0-1.78]
RNAPII Ang II + siRNA Control [0-1.78]
RNAPII DMEM + siRNA Gβ₁ [0-1.78]
RNAPII Ang II + siRNA Gβ₁ [0-1.78]

Thbs1

F

chr1:21,850 kb   21,851 kb   21,852 kb   21,853 kb   21,854 kb

Flag-Gβ₁ DMEM [0-1.11]
Flag-Gβ₁ Ang II [0-1.11]
RNAPII DMEM + siRNA Control [0-1.89]
RNAPII Ang II + siRNA Control [0-1.89]
RNAPII DMEM + siRNA Gβ₁ [0-1.89]
RNAPII Ang II + siRNA Gβ₁ [0-1.89]

Ctgf

Supplemental Figure 1.

Supplmental Figure 2.

A

	Cytosolic fraction		Nuclear fraction		Cytosolic fraction		Nuclear fraction		
DMSO	+	+	+	+	−	−	−	−	1 h pre-treatment
40μM Importazole	−	−	−	−	+	+	+	+	pre-treatment
DMEM	+	−	+	−	+	−	+	−	45 min treatment
1 mM carbachol	−	+	−	+	−	+	−	+	treatment

IP: Gβ$_{1-4}$ Blot: Rpb1

IP: Gβ$_{1-4}$ Blot: Gβ$_{1-4}$

Total Lysate Rpb1

Total Lysate Gβ$_{1-4}$

Total Lysate Nup62

Total Lysate Hsp90

B

Ratio of Rpb1 immunopreceipitated with Gβ$_{1-4}$ (fold change over control)

Vehicle treated — Cytosolic fraction / Nuclear fraction

Importazole treated — Cytosolic fraction / Nuclear fraction

*

- 45 min DMEM
- 45 min 1 mM Carbachol
- 45 min DMEM
- 45 min 1 mM Carbachol
- 45 min DMEM
- 45 min 1 mM Carbachol
- 45 min DMEM
- 45 min 1 mM Carbachol

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5

Supplemental Figure 6

Supplemental Figure 7

Supplemental Figure 8

Supplemental Figure 9

Cardiac Fibroblasts

$G\alpha q$

$PLC\beta$

Ca^{2+}

PKC

CaMKII

MEK

P-TEFb

$G\beta\gamma$-RNAPII
Interaction

HEK 293F cells

$G\alpha q$

$PLC\beta$

Ca^{2+}

Calcineurin

P-TEFb

$G\beta\gamma$-RNAPII
Interaction