Response of pneumococcus to changes in temperature and oxygen varies by

serotype, lineage, and site of isolation

Adrienn Tothpal[1,2*], Katherine Desobry[1], Anne L. Wyllie[1], Daniel M. Weinberger[1*]

[1]Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT 06510,

USA

[2]Institute of Medical Microbiology, Semmelweis University, Budapest 1089, Hungary

Running title: Response of pneumococcus to environment

The total number of words of the manuscript main text: 3028

The number of words of the abstract: 192

The number of main text figures: 6

The number of main text tables: 1

The number of supplementary figures: 5

Footnotes

Conflict of interest: AT and KD report no conflicts of interest. ALW has received consulting fees from Pfizer. DMW has received consulting fees from Pfizer, Merck, GSK, and Affinivax.

Funding: This work was supported by the National Institute of Health/National Institute of Allergy and Infectious Diseases [grant numbers R01AI123208, R56AI110449]. The funding agencies were not involved in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Presented in part: at ASM Microbe, New Orleans, 2017 and the 11th International Symposium on Pneumococci and Pneumococcal Diseases (ISPPD), Melbourne, 2018.

Corresponding authors: tothpal.adrienn@gmail.com, daniel.weinberger@yale.edu

Adrienn Tothpal: Mailing address: Institute of Medical Microbiology, Semmelweis University, Nagyvarad ter 4, Budapest, Hungary HU-1089. Phone: +36 1 459-1500. Fax: +36 1 459-1500

Daniel Martin Weinberger: Mailing address: Yale School of Public Health, 60 College Street, New Haven, CT 06510, USA. Phone: +1 203-737-6004. Fax: +1 203-737-1662

Abstract

Background: Pneumococcus, a commensal bacterium of the nasopharynx, has the potential to cause severe disease. Invasion of different body sites results in exposure to a range of environmental conditions, including different temperature and oxygen levels. The response to these variations could influence virulence and transmissibility.

Methods: We evaluated the effect of temperature and oxygen on the growth of 256 pneumococcal isolates representing 53 serotypes, recovered from healthy carriers and from invasive disease (IPD), conjunctivitis, and pneumonia patients. Strains were grown at a range of physiologically-relevant temperatures anaerobically or in ambient air with and without catalase and were monitored by reading the optical density.

Results: There was considerable variability between strains in response to changes in temperature and oxygen. Some of this variability was associated with serotype. Generally, carriage and IPD isolates grew to the maximal density at the temperature of the nasopharynx (~33C), but IPD isolates began growing earliest at internal body temperature. Growth curve characteristics correlated with epidemiological estimates of serotype-specific carriage prevalence and invasiveness.

Discussion: Environmental variability has differential effects on the growth of carriage and disease isolates and of different pneumococcal serotypes, which could influence virulence and transmissibility.

61 **Keywords:** pneumococcal infection, growth curves, temperature, carriage, anaerobic growth

Streptococcus pneumoniae (pneumococcus) is a commensal bacterium of the human nasopharynx. The nasopharynx is considered to be the reservoir and source of pneumococcal transmission between individuals [1]. Pneumococcus is an opportunistic pathogen that causes diseases ranging from mild but common illnesses like sinusitis and otitis media to serious illnesses such as pneumonia, sepsis and meningitis [2].

In different anatomical sites within the human host, pneumococci are exposed to variable temperature and oxygen levels. In the nasopharynx, considered its main niche, the average temperature is around 33°C, with some differences between children and adults [3–6]. The core body temperature, which would be encountered during invasion into tissues, is 37°C. The temperature in the lungs is constantly changing based on the temperature of inhaled air but is generally lower than 37°C [7]. During infection by pneumococci or during viral co-infection (such as influenza or RSV), both external and internal temperature increases [8–12]. Oxygen levels also vary within the host. In the nasopharynx, bacteria on top of the mucus layer are exposed to almost ambient air (20% O_2). Pneumococci in biofilms in the nasopharynx encounter lower levels of oxygen [13]. Entering the lower respiratory tract or the middle ear, pneumococci are exposed to micro-aerophilic conditions and to almost anaerobic conditions when present in blood and the cerebrospinal fluid (CSF) [14–16]. Likewise, mucus production during infection (i.e., due to influenza or RSV) can block the air passage and form micro-aerophilic (around 5% O_2) or even anaerobic microenvironments [15,16].

To understand how pneumococci successfully colonize the host and transition from nasopharyngeal colonization to severe disease, it is important to understand how different strains

respond to variations in the microenvironment. Pneumococcus is a diverse pathogen, with >90

serotypes (defined by the capsule polysaccharide) and has tremendous genetic variation that results

from recombination. This genetic variability influences the biology and epidemiology of the strains

[17]. Serotypes vary in their prevalence among healthy carriers and in the likelihood that they will

cause severe disease. Laboratory studies have evaluated variations in growth characteristics by

serotype and have identified a relationship between *in vitro* growth characteristics and prevalence

of serotypes in the nasopharynx of healthy children [18,19]. However, the effect of variation in

temperature and oxygen on the growth of different strains and serotypes has not been

systematically explored. The response to such environmental variation could influence the success

of a particular serotype in the nasopharynx or the likelihood of causing disease.

The aim of this study was to investigate how environmental variability in temperature and oxygen

influences the growth of different strains and serotypes of pneumococci. Using a diverse set of

clinical and nasopharyngeal isolates, as well as capsule-switch and capsule-knockout variants

generated in the lab, we quantified how the growth of pneumococcus *in vitro* varies under a range

of physiologically-relevant temperatures that represent colonization and invasive infections as well

in aerobic and anaerobic conditions. We evaluated the relationship between these *in vitro* growth

patterns and relevant serotype-specific epidemiological and biological variables.

Methods

Bacterial strains, culture media, and chemicals

Strains

Invasive pneumococcal disease (IPD) isolates were obtained from the isolate bank at the Centers

for Disease Control/Active Bacterial Core surveillance; carriage isolates were provided by Ron

Dagan (Ben-Gurion University, Israel), Adrienn Tothpal and Eszter Kovacs (Semmelweis

University, Hungary [20,21]) and Debby Bogaert and Anne Wyllie (UMC, Utrecht [22]) (**Table

1**). Capsule-switch variants generated on the TIGR4 genetic background and the serotype 6B

knockout strain were provided by Marc Lipsitch and generated as previously described [23].

Additional capsule-knockout strains were generated by replacement of the capsule biosynthesis

locus with the Sweet Janus cassette [24].

Culture media

Pneumococcal isolates were stored at -80 °C on Cryobeads (Cryobank, Copan Diagnostics,

Murrieta, CA). Strains were routinely grown at 37°C and 5% CO_2 overnight on tryptic soy agar

plates supplemented with 5% sheep blood (TSAII) (Thermo Fisher Scientific). Growth in broth

culture was performed in BHI (Becton, Dickinson, and Co., Sparks, MD) with and without catalase

(5000 units, Worthington Biochemical Corporation, Lakewood, NJ) for aerobic cultivation and

with Oxyrase® (Oxyrase, Inc., West Mansfield, OH) diluted 1:10 to create an anaerobic

environment.

Growth experiment

Strains were streaked onto TSAII plates and incubated at 37°C in a 5% CO2-enriched atmosphere overnight, then harvested into PBS to OD600 0.2 and diluted (6 fold) in BHI with or without catalase or Oxyrase. Growth was monitored in sterile flat-bottomed 96-well microtiter plates (BRAND GMBH, Wertheim, Germany) for 24 hours in a microplate reader (BioTek ELx808) with a built-in incubator, reading the optical density at 600 nm every 30 minutes, after 5s shaking (Gen5 program). Each strain was tested in all three oxygen conditions and across the full range of temperatures (30-39°C). An anaerobic control strain (*Bacteroides thetaiotaomicron*) was used to confirm the elimination of oxygen by Oxyrase.

Data analysis

Each growth curve was blanked by subtracting the OD600 reading at t=0 for that well. In instances where the t=30 minutes measurement was lower than the t=0 measurement due to measurement error at the first time point, the OD600 at t=30m was subtracted instead. For each growth curve, we estimated the length of the lag phase and the maximum growth rate using the groFit package in R [25]. For each curve we also extracted the maximum OD600 measurement. To quantify variations in growth characteristics by serotype, site of isolation, and environmental condition while adjusting for repeated measurements and strain-to-strain variations, we used linear mixed effects models (lme4 package in R) [26]. The outcome variable was maximum OD600, fixed effects variables included serotype, temperature (categorical), oxygen (aerobic+catalase, aerobic without catalase, anaerobic), site of isolation (categorical), and a random intercept for each isolate. Certain interactions among the fixed effects were also evaluated to test specific hypotheses (site of isolate*oxygen; oxygen*temperature; serotype*oxygen). The significance of these interactions

at different levels was evaluated using the interactionMeans function in the phia package in R [27].

Correlations of the effect of oxygen (ratio of maximum OD at a particular temperature for a

particular isolate grown anaerobically or aerobically with catalase) with previously described

epidemiological characteristics (invasiveness, carriage prevalence) were assessed using

Spearman's correlations [28]. Associations with the presence of specific capsule components were

assessed using Wilcoxon rank sum tests in R.

Principal components analysis and linkage to serotype-specific characteristics

We sought to evaluate the link between growth curve characteristics and previously published

serotype-specific characteristics (capsule structure, disease severity, invasiveness, pre-vaccine

carriage prevalence) [29–31]. Across all growth curves, isolates, conditions, and time points, there

was a large volume of data with many possible variables that could be generated for correlation

analyses (e.g., maximum OD600 achieved, OD600 at time point T). To avoid overfitting the data,

we first performed principal components analysis. For each of the possible

serotype/temperature/oxygen/isolation site combinations, we obtained the mean OD600 value at

each time point. We then created an S*C matrix where S represented the number of serotypes and

C represented the number of unique temperature/oxygen/isolation site/time point combinations.

Therefore, there was a variable for every tested environmental condition, measured at each time

point for isolates from different diseases sites, for each serotype. We also created variables for

maximum OD600 reached under the different environmental conditions. These analyses were

restricted to the carriage and invasive pneumococcal disease (IPD) isolates. With this matrix,

missing values occurred (i.e., no serotype 12F carriage isolates were tested, leading to missingness

in the summary values). Therefore, we performed a simple mean imputation of the missing values

(calculate mean of the non-missing values for a column and assign this value to the missing

values). The variables were then centered and scaled prior to use. Using this matrix with 1470

variables for 51 serotypes, we performed principal components analysis (princomp in R); Six of

the principal components (PCs) explained 50% of the variation in the data (PC1: 14%, PC2: 11%,

PC3: 9%, PC4: 6%, PC5: 6%, PC6: 6%). These 6 PCs were used in subsequent regression analyses.

In a series of linear regressions, we evaluated the association between these 6 PCs and pre-vaccine

carriage prevalence (composite of several studies, square root transformed), case-fatality rate,

complexity of the capsular polysaccharide (carbons/polysaccharide repeat) and invasiveness

(square root transformed). We also used logistic regression to evaluate the association between the

6 PCs and the presence of N-acetylated sugars in the capsule or the presence of GlcA and GalA

(both derived from same pathway).

Data availability

The raw data and R analysis scripts are available in a github repository

(https://github.com/weinbergerlab/GrowthVariation).

Results

Growth characteristics

In total, we performed more than 4900 growth curves on 256 different pneumococcal strains,

representing 53 different serotypes. We tested aerobic and anaerobic conditions ranging from 30-

39°C. There was a positive correlation between growth rate and maximum density achieved

(**Supplementary Figure 1**). Most strains reached their maximum density at low temperatures. The

length of the lag phase was shortest for carriage isolates at low temperatures (i.e., 33°C) and was

shortest for IPD isolates at 37°C. As a representative example, growth curves for serotype 19F

isolates from carriage, conjunctivitis, pneumonia, and IPD are shown in **Figure 1**. Growth curves

for   all   isolates   and   conditions   can   be   explored   interactively   at

https://weinbergerlab.shinyapps.io/ShinyGrowth_v2/.

The effect of temperature and oxygen varies based on site of isolation

The presence of oxygen had different effects on IPD and carriage isolates. On average, IPD isolates

grew to a significantly higher density under aerobic+catalase conditions compared with anaerobic

conditions (**Figure 2A**). This was true at both low and high temperatures (**Figure 2B, 2C**). In

contrast, carriage isolates grew similarly in aerobic+catalase and anaerobic conditions at high

temperatures but grew to a significantly higher density in aerobic+catalase at low temperatures.

Conjunctivitis and pneumonia isolates also grew similarly in anaerobic and aerobic+catalase

conditions, although fewer isolates were tested (**Figure 2**).

When grown in ambient air without catalase, pneumococcus typically produces toxic levels of

hydrogen peroxide. As expected, most IPD isolates hardly grew in ambient air due to this toxicity.

Unexpectedly, some IPD isolates carriage isolates grew moderately under these conditions (e.g.,

IPD isolates of serotypes 9N, 12F, 18A,23A, 33F IPD isolate; **Figure 5**). Moreover, carriage

isolates grew moderately well in ambient air without catalase, especially at higher temperatures

(35-39°C). (**Figure 2C, 3**).

Importance of oxygen varies by serotype

We next investigated whether the effect of oxygen on growth patterns varied by serotype (**Figure**

4, 5). For these analyses, we focused on comparing growth in aerobic+catalase versus anaerobic

conditions. Overall, certain serotypes (2, 4, 13, 23B, 35B) grew better under aerobic conditions

(with catalase), while other serotypes (6B, 8, 9N, 12F) did not show a difference between aerobic

and anaerobic growth (**Figure 4**). The differences between aerobic and anaerobic conditions were

more pronounced for the IPD isolates, but the direction of association was generally similar for

both IPD and carriage isolates.

The clinical isolates represented diverse genetic backgrounds. We attempted to evaluate the growth

characteristics of capsule-knockout strains as well as several capsule-switch variants. While the

results were ambiguous, they suggested that the effect of capsule production on growth was more

pronounced   during   anaerobic   growth   compared   with   aerobic   growth   with   catalase

(**Supplementary Figures 2-5**).

Relationship of growth characteristics to serotype-specific characteristics

Finally, we performed exploratory analyses to evaluate whether the characteristics of the growth

curves were related to serotype-specific epidemiological characteristics and biochemical

characteristics of the capsule. There was an association between the second principal component

(PC2, which explained 11% of the variability in the data) and pre-vaccine carriage prevalence

(p<0.001) and invasiveness (p=0.05) (**Figure 6**). We therefore evaluated which components of the

growth curves contributed most to PC2. PC2 was largely influenced by density at ~5 hours among

the carriage isolates at 33C and ~2-3 hours among the IPD isolates at 38°C. These time points

corresponded to periods when many of the isolates were in early log-phase growth but some were

still in stationary phase. PC2 was also influenced by density at later time points, which largely

(inversely) reflects density at early time points.

There was no association between PC1-6 and serotype-specific case fatality ratio, polysaccharide

complexity, or specific polysaccharide components. There were also no notable correlations

between the effect of oxygen (ratio of maximum OD600 in anaerobic vs aerobic growth with

catalase) and the invasiveness of a serotype, case-fatality of the serotype, carriage prevalence of

the serotype, or with any capsule components.

Discussion

We provide novel information about the effect of environmental conditions on the growth characteristics of pneumococcal strains. Testing a diverse set of isolates, pneumococci grew to the highest density under conditions that mimic the normal environment of the nasopharynx in terms of temperature and oxygen level. Important differences were observed between carriage and disease isolates and by serotype, possibly reflecting adaptation to the environment from which they were isolated. The hierarchy of strains growing at early time points correlated with several serotype-specific epidemiological and biological characteristics.

As the nasopharynx is the normal habitat of pneumococcus, we hypothesized that the nasopharyngeal temperature would be ideal for growth. Indeed, temperature played an important role in terms of the maximal density achieved and how quickly the isolates started growing (lag phase). Carriage isolates grew to the highest density and had the shortest lag at temperatures resembling those of the nasopharynx. This finding, along with recent work on the effect of lower temperatures on the immune response to pathogens in the upper respiratory tract [32], suggests that the environment of the nasopharynx is optimal for pneumococcal growth and survival. IPD isolates responded differently to environmental variations. While they also reached maximum density at low temperatures, the IPD isolates had the shortest lag phase at 37°C (i.e., core body temperature). Additionally, IPD isolates hardly grew in ambient air without catalase, while carriage isolates grew moderately well, especially at higher temperatures (**Figure 1-4**).

During the invasion process, when the pneumococcus transitions from the nasopharyngeal environment to the internal body environment, it has to adapt to many changes, including nutrient,

temperature and oxygen levels. Temperatures vary from the low-30° range in the upper respiratory

tract to 37°C in core body sites and even higher during fever. Likewise, oxygen levels can vary

considerably during infection. Increased mucus production (due to co-infection with viruses or

other pathogens in the upper respiratory tract) leads to lower oxygen levels and may generate a

local hypoxic environment [33–35] . The availability of oxygen is also decreased in pneumonia,

empyema, and otitis media. Oxygen levels in the uninflamed middle ear space, for example,

resemble that of venous blood, are less than a third that of the airway, and may be further reduced

by the presence of effusion [14,15,36]. These gradients of temperature and oxygen could influence

which strains succeed, and pneumococcus might adapt to these variations. We found that IPD

isolates, but not carriage isolates, started growing more quickly at body temperature (37°C) than

at lower temperature but reached a lower final density then the same isolates grown at lower

temperatures, and strains generally grew to a higher density with oxygen. However, some isolates

appeared to be less susceptible to these variations in oxygen and temperature, potentially

influencing the likelihood of a carriage episode leading to infection.

Some of the differences observed in growth phenotypes between carriage and disease isolates

could reflect opaque/transparent phase variation [37]. Opaque variants are generally isolated from

IPD and have increased capsule production and decreased production of certain surface proteins.

Phenotypically, the presence of oxygen accentuates differences in capsule production between

opaque and transparent variants [36]. This effect could be mediated via the pathways involved in

converting pyruvate to acetyl-CoA, an important biochemical precursor for capsule production for

many serotypes [38]. Variations in the use of this pathway between serotypes or the efficiency of

this pathway between lineages could influence some of the patterns that were observed in the

growth curves.

Serotype is a major determinant of pneumococcal epidemiology and biology. When comparing

growth in aerobic conditions (with catalase) with growth in anaerobic conditions, the benefit of

oxygen varied by serotype (**Figure 4, 5**). Serotypes 2, 4, 13, 23B, 35B and 38 grew better with

additional catalase than in anaerobic conditions, whereas serotypes 6B, 8, 9N, 12F, 22F and 42

grew similarly in both environments. These patterns did not correlate with the presence of N-

acetylated sugars in the capsule, which are derived from a pathway that would be influenced by

the presence of oxygen [38], nor did they correlate with invasiveness of the serotype. We tested a

relatively small number of isolates per serotype, so it is possible that non-capsular genetic

variations or differences in the culture history of the isolates could influence the observed

responses to oxygen.

This study had certain limitations. For the growth curves, we used BHI broth which is an artificial

growth medium that differs in nutrient composition from the host. We evaluated several minimal

media but found that growth was generally poor, making comparisons between strains difficult.

While we tested a large number of strains representing many serotypes, some serotypes were only

represented by a single isolate (i.e., 11B, 12F, 13). This could limit the generalizability of serotype-

specific findings in these instances, making it difficult to make inferences about whether variability

was due to serotype, site of isolation, or lineage effects. The strains used in this study were largely

a convenience sample from clinical studies. The genetic diversity of pneumococcus makes it

difficult to draw conclusions about the cause of differences between strains. The growth curves

with the capsule-knockout strains and capsule-switch variants suggests that the capsule itself could

influence these phenotypes. We did not perform any gene expression studies which could be highly

influenced by environmental conditions [39]. Further work could explore the genetic basis (both

capsular and non-capsular factors) for the differences in growth phenotypes between strains.

In conclusion, we demonstrate that the growth characteristics of pneumococcus are influenced by

environmental variations, that the effect of these variations depend on the site of isolation of the

bacterium, and that the optimal growth conditions for carriage isolates resemble the conditions of

the nasopharynx. Moreover, we demonstrate that the growth patterns among serotypes are

associated with carriage prevalence and other epidemiological and biological characteristics.

These results could help in understanding which of the serotypes has the greatest capacity to

emerge - in both carriage and disease - in the future.

Acknowledgements: The authors gratefully acknowledge Marc Lipsitch, Krzysztof Trzcinski and

Claudette Thompson for providing the TIGR4 and serotype 6B capsule-switch variants, the isolate

bank from the Active Bacterial Core surveillance (ABCs)/Emerging Infections Programs (EIP)

Network for providing the IPD isolates from the United States, Ron Dagan for providing the

isolates from Israel, and Eszter Kovacs and Orsolya Dobay for providing the isolates from

Hungary.

References

1. Bogaert D, Engelen MN, Timmers-Reker AJM, et al. Pneumococcal Carriage in Children in The Netherlands: a Molecular Epidemiological Study. J Clin Microbiol **2001**; 39(9):3316–20.

2. O'Brien KL, Wolfson LJ, Watt JP, et al. Burden of disease caused by Streptococcus pneumoniae in children younger than 5 years: global estimates. Lancet **2009**; 374(9693):893–902.

3. Geurkink N. Nasal anatomy, physiology, and function. J Allergy Clin Immunol **1983**; 72(2):123–8.

4. Jones N. The nose and paranasal sinuses physiology and anatomy. Adv Drug Deliv Rev **2001**; 51(1):5–19.

5. Keck T, Leiacker R, Heinrich A, Kühnemann S, Rettinger G. Humidity and temperature profile in the nasal cavity. Rhinology **2000**; 38(4):167–71.

6. Van Cauwenberge P, Sys L, De Belder T, Watelet J-B. Anatomy and physiology of the nose and the paranasal sinuses. Immunol Allergy Clin North Am **2004**; 24(1):1–17.

7. Webb P. Air Temperatures in Respiratory Tracts of Resting Subjects in Cold. J Appl Physiol **1951**; 4(5):378–82.

8. Ioannou S, Ebisch S, Aureli T, et al. The Autonomic Signature of Guilt in Children: A Thermal Infrared Imaging Study. PLoS ONE **2013**; 8(11).

9.   Keck T, Leiacker R, Riechelmann H, Rettinger G. Temperature Profile in the Nasal Cavity.

The Laryngoscope **2000**; 110(4):651–4.

10.   Lindemann J, Keck T, Scheithauer MO, Leiacker R, Wiesmiller K. Nasal mucosal

temperature in relation to nasal airflow as measured by rhinomanometry. Am J Rhinol

2007; 21(1):46–9.

11.   McFadden ER, Pichurko BM, Bowman HF, et al. Thermal mapping of the airways in

humans. J Appl Physiol **1985**; 58(2):564–70.

12.   Willatt DJ. Continuous infrared thermometry of the nasal mucosa. Rhinology **1993**;

31(2):63–7.

13.   Borriello G, Werner E, Roe F, Kim AM, Ehrlich GD, Stewart PS. Oxygen Limitation

Contributes to Antibiotic Tolerance of Pseudomonas aeruginosa in Biofilms. Antimicrob

Agents Chemother **2004**; 48(7):2659–64.

14.   Harell M, Mover-Lev H, Levy D, Sade J. Gas composition of the human nose and

nasopharyngeal space. Acta Otolaryngol **1996**; 116(1):82–4.

15.   Hergils L, Magnuson B. Middle ear gas composition in pathologic conditions: mass

spectrometry in otitis media with effusion and atelectasis. Ann Otol Rhinol Laryngol **1997**;

106(9):743–5.

16.   Yesilkaya H, Andisi VF, Andrew PW, Bijlsma JJ. Streptococcus pneumoniae and reactive

oxygen species: an unusual approach to living with radicals. Trends Microbiol **2013**;

21(4):187–95.

17.   Bentley SD, Aanensen DM, Mavroidi A, et al. Genetic analysis of the capsular biosynthetic

378   locus from all 90 pneumococcal serotypes. PLoS Genet **2006**; 2(3):e31.

379   18.   Bättig P, Hathaway LJ, Hofer S, Mühlemann K. Serotype-specific invasiveness and

380   colonization prevalence in Streptococcus pneumoniae correlate with the lag phase during in

381   vitro growth. Microbes Infect **2006**; 8(11):2612–7.

382   19.   Hathaway LJ, Brugger SD, Morand B, et al. Capsule type of Streptococcus pneumoniae

383   determines growth phenotype. PLoS Pathog **2012**; 8(3):e1002574.

384   20.   Tóthpál A, Kardos S, Laub K, et al. Radical serotype rearrangement of carried

385   pneumococci in the first 3 years after intensive vaccination started in Hungary. Eur J

386   Pediatr **2015**; 174(3):373–81.

387   21.   Tóthpál A, Laub K, Kardos S, et al. Epidemiological analysis of pneumococcal serotype

388   19A in healthy children following PCV7 vaccination. Epidemiol Infect **2016**; 144(7):1563–

389   73.

390   22.   Prevaes SMPJ, Winter-de Groot KM de, Janssens HM, et al. Development of the

391   Nasopharyngeal Microbiota in Infants with Cystic Fibrosis. Am J Respir Crit Care Med

392   **2015**; 193(5):504–15.

393   23.   Trzcinski K, Thompson CM, Lipsitch M. Construction of Otherwise Isogenic Serotype 6B,

394   7F, 14, and 19F Capsular Variants of Streptococcus pneumoniae Strain TIGR4. Appl

395   Environ Microbiol **2003**; 69(12):7364–70.

24. Li Y, Thompson CM, Lipsitch M. A modified Janus cassette (Sweet Janus) to improve

allelic replacement efficiency by high-stringency negative selection in Streptococcus

pneumoniae. PloS One **2014**; 9(6):e100510.

25. Matthias Kahm, Hasenbrink G, Lichtenberg-Frate H, Jost Ludwig, Kschischo M. grofit:

Fitting Biological Growth Curves with R. J Stat Softw **2010**; 33(7):1–21.

26. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using

lme4. J Stat Softw **2015**; 67(1).

27. Rosario-Martinez HD. phia: Post-Hoc Interaction Analysis. R package. **2015**. Available at:

https://CRAN.R-project.org/package=phia

28. Weinberger DM, Grant LR, Steiner CA, et al. Seasonal Drivers of Pneumococcal Disease

Incidence: Impact of Bacterial Carriage and Viral Activity. Clin Infect Dis Off Publ Infect

Dis Soc Am **2014**; 58(2):188–94.

29. Weinberger DM, Trzciński K, Lu YJ, et al. Pneumococcal capsular polysaccharide

structure predicts serotype prevalence. PLoS Pathog **2009**; 5(6):e1000476.

30. Weinberger DM, Grant LR, Weatherholtz RC, Warren JL, O'Brien KL, Hammitt LL.

Relating Pneumococcal Carriage Among Children to Disease Rates Among Adults Before

and After the Introduction of Conjugate Vaccines. Am J Epidemiol **2016**; 183(11):1055–62.

31. Weinberger DM, Harboe ZB, Sanders EAM, et al. Risk of death from pneumococcal

pneumonia is a stable serotype-associated property: a meta-analysis. Clin Infect Dis **2010**;

51(6):692–9.

32. Foxman EF, Storer JA, Fitzgerald ME, et al. Temperature-dependent innate defense against

the common cold virus limits viral replication at warm temperature in mouse airway cells.

Proc Natl Acad Sci U S A **2015**; 112(3):827–32.

33. Bakaletz LO. Immunopathogenesis of polymicrobial otitis media. J Leukoc Biol **2010**;

87(2):213–22.

34. Giebink GS, Ripley ML, Wright PF. Eustachian Tube Histopathology during Experimental

Influenza a Virus Infection in the Chinchilla. Ann Otol Rhinol Laryngol **1987**; 96(2):199–

206.

35. Sadé J, Luntz M, Levy D. Middle ear gas composition and middle ear aeration. Ann Otol

Rhinol Laryngol **1995**; 104(5):369–73.

36. Weiser JN, Bae D, Epino H, et al. Changes in Availability of Oxygen Accentuate

Differences in Capsular Polysaccharide Expression by Phenotypic Variants and Clinical

Isolates of Streptococcus pneumoniae. Infect Immun **2001**; 69(9):5430–9.

37. Weiser JN, Austrian R, Sreenivasan PK, Masure HR. Phase variation in pneumococcal

opacity: relationship between colonial morphology and nasopharyngeal colonization. Infect

Immun **1994**; 62(6):2582–9.

38. Echlin H, Frank MW, Iverson A, et al. Pyruvate Oxidase as a Critical Link between

Metabolism and Capsule Biosynthesis in Streptococcus pneumoniae. PLoS Pathog **2016**;

12(10).

39. Aprianto R, Slager J, Holsappel S, Veening JW. High-resolution analysis of the

pneumococcal transcriptome under a wide range of infection-relevant conditions. Nucleic

Acids Res **2018**. Available at: https://academic.oup.com/nar/advance-

article/doi/10.1093/nar/gky750/5079695

Figure legends

Figure 1. Growth patterns of serotype 19F strains isolated from different sites in the body. Growth was measured every 30 minutes over 24 hours at 30-39°C, with ambient air, ambient air with catalase, or under anaerobic conditions. Each line corresponds to a growth curve in a different condition—bluer temperatures correspond to colder conditions, orange colors correspond to warmer conditions. Each column corresponds to an isolate obtained from a different anatomical site.

Figure 2. Effect of oxygen on the maximum density achieved for carriage and clinical samples (A) across all temperatures, (B) at low temperature (30-35°C) and (C) at high temperature (37-39°C). Blue triangles represent aerobic growth with catalase, orange circles represent aerobic without catalase, green squares represent anaerobic growth. The y-axis shows the maximum density achieved for IPD, carriage, pneumonia, and conjunctivitis isolates, after adjusting for serotype and temperature. Mean+/- 95% confidence intervals.

Figure 3. The effect of temperature and oxygen on growth of (A) all isolates, (B) IPD isolate and (C) carriage isolates. The y-axis shows the maximum density at different temperature and oxygen conditions, after adjusting for serotype effects. Mean+/- 95% confidence intervals.

Figure 4. Relative difference in maximum density when grown in aerobic conditions with catalase versus anaerobic conditions. Positive values indicate that a higher density (as measured by optical density) was achieved under aerobic conditions. A value of 0 indicates no

difference. Mean+/- 95% confidence intervals. The reference in the regression is serotype 14, and the temperature was included as a covariate.

Figure 5. The effect of serotype and oxygen on growth of carriage isolates and IPD isolates. The y-axis shows the maximum density achieved for each serotype. Mean+/- 95% confidence intervals. Temperature was included as a covariate.

Figure 6. Correlation between the second principal component and (A) serotype-specific pre-vaccine carriage prevalence in a composite of settings and (B) serotype-specific log-invasiveness. Higher principal component scores correspond to higher density at an early time point. Both of the epidemiologic measures were estimated and described in previous studies. The size of the circle is proportional to the inverse variance of the invasiveness estimate [30].

Supplementary Figure 1. Correlation between maximum density achieved by an isolate and the growth rate across (A) carriage and (B) IPD isolates. Individual dots represent individual isolates. Lines represent LOESS fit. Strains with steep slopes reached a higher maximal OD.

Supplementary Figures 2-5. Growth curves for capsule-switch variants on the TIGR4 background and capsule knockout strains.

Table 1. Pneumococcal strains used in this study.

Source of isolate	Number of isolates	Country	Serotypes
Invasive pneumococcal disease (sepsis, meningitis)	40	US (CDC)	1, 2, 3, 4, 6A, 6B, 6C, 6D, 7C, 7F, 8, 9N, 9V, 10A, 10F, 11A, 11B, 12F, 13, 14, 15A, 15B, 15C, 15F, 17F, 18A, 18B, 18C, 19A, 19B, 20, 22F, 23F, 29, 31, 33F, 34, 35A, 35B, 37
	2	Hungary	3, 6B/D
Pneumonia	8	Hungary	3, 8, 10A, 15A, 19F, 35B, 43/45/38, NT
Conjunctivitis	16		3, 21, 31, 34, 42, 11A, 15A, 15B, 16F, 19A, 19F, 23A, NT
Carriage	52	Israel	6B, 14, 15B/C, 19A, 19F, 23F, NT
	87	Hungary	1, 3, 8, 21, 31, 34, 38, 6A, 9V, 10A, 11A, 15A, 15B/C, 16F, 18C, 19A, 19F, 22A, 22F, 23A, 23B, 23F, 24F, 28F, 35F
	51	Netherlands	3, 27, 10A, 11A, 15B/C, 16F, 19A, 19F, 22A, 22F, 23B, 33F, 35B, 35F, NT
Laboratory-generated genetic variants	4	Various	TIGR4 (cps-, 5, 14, 19F), 603 (6B, cps-), CDC-10A (10A, cps-), CDC-15B (15B, cps-)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary figure 1A

Supplementary figure 1B

Supplementary figure 2

Supplementary figure 3

Supplementary figure 4

Supplementary figure 5