A comprehensive data-driven model of cat primary visual cortex

Ján Antolík[1,2,3] [†*], Rémy Cagnol [1] [†], Tibor Rózsa[1], Cyril Monier[2,4], Yves Frégnac[2,4], Andrew P. Davison[2,4],

1 Faculty of Mathematics and Physics, Charles University, Malostranské nám. 25, 118 00, Prague 1, Czechia
2 Unit of Neuroscience, Information and Complexity (UNIC), CNRS FRE 3693, Gif-sur-Yvette, France
3 INSERM UMRI S 968; Sorbonne Université, UPMC Univ Paris 06, UMR S 968; CNRS, UMR 7210, Institut de la Vision, F-75012, Paris, France
4 Institut des neurosciences Paris-Saclay, Université Paris-Saclay, CNRS, 91400, Saclay, France

†These authors contributed equally to this work.
* antolikjan@gmail.com

1 Abstract

Knowledge integration based on the relationship between structure and function of the neural substrate is one of the main targets of neuroinformatics and data-driven computational modeling. However, the multiplicity of data sources, the diversity of benchmarks, the mixing of observables of different natures, and the necessity of a long-term, systematic approach make such a task challenging. Here we present a first snapshot of a long-term integrative modeling program designed to address this issue in the domain of the visual system: a comprehensive spiking model of cat primary visual cortex. The presented model satisfies an extensive range of anatomical, statistical and functional constraints under a wide range of visual input statistics. In the presence of physiological levels of tonic stochastic bombardment by spontaneous thalamic activity, the modelled cortical reverberations self-generate a sparse asynchronous ongoing activity that quantitatively matches a range of experimentally measured statistics. When integrating feed-forward drive elicited by a high diversity of visual contexts, the simulated network produces a realistic, quantitatively accurate interplay between visually evoked excitatory and inhibitory conductances; contrast-invariant orientation-tuning width; center surround interactions; and stimulus-dependent changes in the precision of the neural code. This integrative model offers insights into how the studied properties interact, contributing to a better understanding of visual cortical dynamics. It provides a basis for future development towards a comprehensive model of low-level perception.

2 Author summary

Computational modeling can integrate fragments of understanding generated by experimental neuroscience. However, most models considered only a few features of neural computation at a time, leading to either poorly constrained models with many parameters, or lack of expressiveness in over-simplified models. A solution is to develop detailed models, but constrain them with a broad range of anatomical and functional

data to prevent overfitting. This requires a long-term systematic approach. Here we present a first snapshot of such an integrative program: a large-scale spiking model of cat primary visual cortex, that is constrained by an extensive range of anatomical and functional features. Together with the associated modeling infrastructure, this study lays the groundwork for a broad integrative modeling program seeking an in-depth understanding of vision.

3 Introduction ¹

A key challenge in neuroscience is the consolidation of the multitude of experimental findings into a coherent characterization of cortical processing. Slow progress in such knowledge integration is evidenced by the surprisingly incomplete state of understanding even in well-explored cortical areas, such as the primary visual cortex (V1) [1].

For example, while V1 circuitry supports many different computations occurring concurrently, such as edge detection, depth processing, and contextual modulation, these have mostly been studied in isolation. Many computational studies have proposed mechanisms to explain V1 phenomena — one at a time — including layer-specific resting-state differences [2], contrast adaptation [3], or orientation tuning contrast invariance [4–8]. An alternative to such an unipotent model approach is to propose a common adaptive mechanism capable of explaining multiple V1 properties (e.g. [9,10]). However, only few concurrent phenomena have been actually demonstrated in such models (but see [11]), despite the potential these approaches hold for knowledge integration. The most comprehensive explanations of V1 function come from a few large-scale models combining well-established cortical mechanisms [12,13], but these studies still account for only a limited number of phenomena at a time (see section 5.3).

Failure to address the full spectrum of phenomena concurrently in a single model instance leads to under-constrained models, and proliferation of numerous alternative explanations for individual phenomena, while informing us little about how different V1 computations are multiplexed such that the same synapses participate in multiple simultaneous calculations. While further reductionist, hypothesis-led research is indisputably required, many of these questions require a systematic, integrative approach to progressively build a unified multi-scale theory of brain function [14,15]. A danger in detailed modeling is growth of free parameters, risking over-fitting and consequent lack of explanatory power. However, building and validating a model against a large number of diverse experimental studies adds strong constraints on parameters. Similarly, a multi-scale approach greatly increases the categories of experiments suitable as a source of model constraints [16].

In our view, the solution to these challenges is a sustained, collaborative effort to incorporate the full breadth of experimentally established constraints into a single pluripotent model of V1. While necessarily a long-term project, here we offer a basis for such an integrationist effort: a detailed, biologically realistic model of cat V1, validated against an unprecedented range of experimental measures, including intra- and extra-cellular functional recordings realized over the years in our laboratory. To support collaborative knowledge integration, the model and all experimental protocols are implemented in an easily extensible simulation framework [17], and all code and data are published online (*http://v1model.arkheia.org*).

The model comprises cortical layers 4 and 2/3, within a 5.0×5.0mm patch around the area centralis (Fig 1; see Methods). Afferent connections to layer 4 neurons are stochastically generated based on receptive-field templates parametrised by the neurons' retinotopic positions, and their location in an overlaid, pre-computed orientation map (Fig 2). Thalamocortical connectivity in Layer 4 follows a push-pull organization [5] (Fig 1E), while intra-cortical connectivity follows parameters derived from experimental

data [18–20] (Fig 1AB). The model has been probed by the most common stimulation paradigms used in early vision: drifting gratings (DGs), masked gratings of variable diameter, and naturalistic movies. Under background thalamic bombardment, a spontaneous cortical activity within a physiologically plausible resting-conductance regime emerges in the model. Under visual stimulation, the same model exhibits diverse patterns of interplay between evoked excitation and inhibition; stimulus-locked subthreshold variability; contrast-invariant orientation tuning; size tuning; stimulus-dependent firing precision; and a realistic distribution of Simple/Complex receptive fields.

The primary contribution of this study is that of knowledge integration: we advance our ability to capture the function of V1 within a single model, and provide insights into how the studied properties interact, contributing to a better understanding of cortical dynamics. We demonstrate for the first time in a model of visual system a self-consistent spontaneous state without unrealistic sources of variability, which allows us to reproduce stimulus dependent changes in trial-to-trial variability of V1 responses previously demonstrated experimentally [21]. To the best of our knowledge, we are the first to demonstrate how a model of V1 can achieve orientation tuning of firing rate, membrane-potential, and excitatory & inhibitory conductance simultaneously (in line with experimental evidence), which is an important constraint on the cortical dynamics driving generation of orientation tuning in V1. We demonstrate the utility of the proposed function-driven integrative approach and provide a basis for future development towards a comprehensive model of V1 and beyond, grounded in an open-science approach.

4 Results

Here we present a single model instance (i.e. a single set of equations with a single set of parameters was used for all simulations; only the visual stimuli changed) representing the cat early visual system, consisting of a retino-thalamic pathway feeding input in the form of spikes to layers 4 and 2/3 of a 5.0×5.0 mm patch of V1 (Fig 1,2) centered at 3 degrees of visual field eccentricity (see Methods).

4.1 Spontaneous activity

The only external source of variability in the model is a white noise current injection into lateral geniculate nucleus (LGN) cells which induces their spontaneous firing at about $17\,\mathrm{Hz}$ for ON cells and $8\,\mathrm{Hz}$ for OFF cells, in line with recordings from LGN in response to a gray blank stimulus at $50\,\mathrm{cd/m^2}$ [22]. This noise represents the combined effect of the retinal spontaneous discharge [23–27], and the intrinsic mechanisms generating ongoing activity in thalamus [28]. Due to the low proportion of synapses of thalamic origin in the model cortex [29], the variability of spontaneous and visually evoked cortical activity is largely driven by internal cortical dynamics.

In the spontaneous condition all modeled excitatory neuron populations fire irregularly (Fig 3A; Fig 4B: coefficient of variation of their inter-spike-intervals is above 0.9) and asynchronously (Fig 3A; Fig 4C: the mean cross-correlation of the spike counts is less than 0.01), with only occasional synchronized events, consistent with spontaneous firing patterns in awake animals [30, 31] and with the up-states in *in vivo* anesthetized preparations [30, 32]. The model exhibits higher spontaneous correlations between inhibitory than between excitatory neurons (Fig 3A; Fig 4C), which is consistent with the observation of higher synchrony among inhibitory neurons in macaque cortex during slow wave sleep [33]. Overall the spontaneous activity corresponds to the asynchronous

Fig 1. The model architecture. (A-B) Layer 2/3 and Layer 4 lateral connectivity. All cortical neuron types make local connections within their layer. Layer 2/3 excitatory neurons also make long-range functionally specific connections. For the sake of clarity A,B do not show the functional specificity of local connections and connection ranges are not to scale. (C) Extent of modeled visual field and example of receptive fields (RFs) of one ON and one OFF-center lateral geniculate nucleus (LGN) relay neuron. As indicated, the model is retinotopically organized. The extent of the modeled visual field is larger than the corresponding visuotopic area of modeled cortex in order to prevent clipping of LGN RFs. (D) Local connectivity scheme in Layer 2/3: connections are orientation- but not phase-specific, leading to predominantly Complex cell type RFs. Both neuron types receive narrow connections from Layer 4 excitatory neurons. (E) Local connectivity in Layer 4 follows a push-pull organization. (F) Afferent RFs of Layer 4 neurons are formed by sampling synapses from a probability distribution defined by a Gabor function overlaid on the ON and OFF LGN sheets, where positive parts of the Gabor function are overlaid on ON and negative on OFF-center sheets. The ON regions of RFs are shown in white, OFF regions in black.

Fig 2. The induction of orientation maps in the model. (A) The precomputed orientation map that serves as a template based on which orientation of the thalamo-cortical connectivity of Layer 4 neurons is determined. (B) The orientation assigned to neurons (each dot represents a neuron) in the two layers based on their position within the overlaid pre-computed orientation map. (C) The comparison between the orientation preference assigned to the neurons via the pre-computed orientation map (y axis) and the orientation preference measured using the sinusoidal grating protocol (x axis; see Section 6.4).

irregular state previously observed in balanced randomly connected network models [34–36]. ⁹²⁻⁹³

We observe higher mean spontaneous rates among inhibitory than excitatory neurons, in line with experimental evidence [37,38]. The spontaneous firing rates of the majority of excitatory neurons are low (<2 Hz, Fig 4AG), as observed experimentally [39,40]. In addition, higher spontaneous firing rates in Layer 2/3 than in Layer 4 have been reported in macaques [41], just like in the model. In both excitatory and inhibitory populations of Layer 2/3, the firing rates closely follow log-normal distributions (Fig 4G), a phenomenon previously shown in several cortical areas and species [42,43]. ⁹⁴⁻¹⁰⁰

The mean resting membrane potential (Vm) of neurons is close to -70 mV, as observed in *in vivo* cat V1 recordings [44]. As shown in Fig 3B and 4EF, the excitatory synaptic conductances recorded in model neurons during spontaneous activity (1.15 nS, averaged across all layers) are consistent with levels observed in cat V1 (~ 1 nS; layer origin not known) [44]. The inhibitory conductance during spontaneous activity was 3.56 nS when averaged across all model neurons, slightly lower than the value of 4.9 nS reported in cat [44]. These realistic levels of conductance are in stark contrast to those in most balanced random network models of spontaneous activity, in which the global synaptic conductance tends to be orders of magnitude higher [36,45]. Overall, the model exhibits a spontaneous state with balanced excitatory and inhibitory inputs well matched with cat V1 at both extra- and intracellular level, without the need for any unrealistic external sources of variability. This represents an advance over previous models of V1 and over dedicated models of spontaneous activity, and enables a self-consistent exploration of stimulus-driven changes in variability (see Sec 4.4). ¹⁰¹⁻¹¹⁴

Fig 3. Spontaneous activity in modeled cortical layers. (A) Raster plot of spiking activity recorded for 5 s of biological time in Layers 4 and 2/3 (from bottom to top; red: excitatory neurons, blue: inhibitory neurons). The relative number of spike trains shown corresponds to the relative number of neurons in the network (total of 1000 neurons shown). (B-E) the membrane potential (black) and the excitatory (red) and inhibitory (blue) synaptic conductances of randomly selected neurons. (B) excitatory neuron in Layer 2/3; (C) inhibitory neuron in Layer 2/3 (C); (D) excitatory neuron in Layer 4; (E) inhibitory neuron in Layer 4.

Fig 4. Statistics of spontaneous activity in the simulated network recorded for 40 seconds. (A–F) Six measures of spontaneous activity for excitatory (black bars) and inhibitory (white bars) neural populations in the two modeled cortical layers. Error bars report standard deviation. (A–C) Statistics of the spiking activity of the 4 modeled cortical populations based on spike trains recorded for 40 s. (A) Mean single-unit firing rates. (B) Irregularity of single-unit spike trains quantified by the coefficient of variation of the inter-spike intervals. (C) Synchrony of multi-unit spiking activity quantified as the mean correlation coefficient between the peri-stimulus time histogram (PSTH) (bin size 10 ms) of all pairs of recorded neurons in a given population. (D–F) Mean values of intracellular variables: membrane potential (D); excitatory (E) and inhibitory (F) synaptic conductances. (G) The distribution of firing rates (black points) in Layer 2/3 excitatory and inhibitory neurons is well fitted by a log-normal distribution with matching mean and variance (gray line; left-column). The log-normal distribution appears as a normal distribution on a semi-logarithmic scale. The data are better fit by a log-normal (gray line) than by an exponential (dashed line) distribution (right-column; log-log scale).

4.2 Responses to drifting sinusoidal gratings

Fig 5 shows the responses of representative excitatory and inhibitory neurons from both   116
model cortical layers to multiple trials of optimally- and cross-oriented sinusoidal   117
gratings of optimal spatial frequency, drifting at 2 Hz.

Fig 5. Responses of representative excitatory (A,C) and inhibitory (B,D) neurons from
Layer 4 (A,B) and Layer 2/3 (C,D) to 10 trials of DGs of optimal (left) and
cross-oriented (right) orientations. For all neurons, the top panel is a spike raster plot,
each line corresponding to a single trial. The middle panel shows the incoming
excitatory (red) and inhibitory (blue) synaptic conductances and the bottom panel
shows the membrane potential (thin lines: single trials; thick lines: mean). Stimulus
onset is at time zero.

The membrane potential of the Layer 4 excitatory cells follow the sinusoidal   119
modulation of the grating luminance — characteristic of Simple cells [44] — but   120
remains largely unmodulated and below threshold during the cross-oriented condition   121
(Fig 5A (bottom panel)). The sinusoidal shape of the membrane potential trace is the   122
result of the excitatory and inhibitory synaptic conductances being mutually in   123
anti-phase (Fig 5A middle panel), due to the push-pull connectivity in model Layer 4.   124
The trial-averaged excitatory and inhibitory synaptic conductances (mean value 1.66 nS   125
and 4.15 nS respectively) are consistent with observations in cat V1 [21, 44]. In layer 4   126
the spikes are generated in a periodic manner aligned to the upswings of the membrane   127
potential, while the cross-oriented grating does not elicit significant depolarization (Fig   128
5A (top panel)).   129

The spiking response of the inhibitory cells in Layer 4 (Fig 5B) is similar. The   130
smaller number of excitatory cortico-cortical synapses is compensated by the stronger   131
excitatory to inhibitory synaptic weights (see Methods), leading to comparable mean   132
input excitatory conductances (1.55 nS). However, the smaller number of inhibitory   133
cortico-cortical synapses yields lower mean input inhibitory conductances (2.65 nS),   134
resulting in a shift towards stronger relative excitation during visual stimulation, and   135
hence higher overall firing rates (see Section 4.2.1). Furthermore, due to the weaker   136
inhibition, the cross-oriented grating still elicits slight depolarization of the membrane   137
potential, leading to increased spiking activity, and resulting in broadening of the   138
orientation tuning curve, as further examined in Section 4.2.1.   139

Unlike the phasic response of Layer 4 cells, both excitatory and inhibitory cells in   140
Layer 2/3 show steady depolarization in response to an optimally orientated grating,   141
similar to non-linear Complex cells (Fig 5CD). The mean excitatory and inhibitory   142

synaptic conductances (2.48 nS and 14.01 nS respectively) are higher than in Layer 4 ₁₄₃
due to the higher number of recurrent connections (see Methods), and lie within the ₁₄₄
ranges observed experimentally [21, 44]. ₁₄₅

4.2.1 Orientation tuning of spike response ₁₄₆

We next examined the responses of the model to different orientations of the sinusoidal ₁₄₇
grating at two different contrasts. The mean tuning widths of excitatory and inhibitory ₁₄₈
neurons in Layer 4 and excitatory and inhibitory neurons in Layer 2/3, measured as ₁₄₉
half-width-at-half-height (HWHH), were 21.1°, 23.3°, 25.2° and 28.3° respectively (Fig ₁₅₀
6C), which is within the range of tuning widths in cat V1 reported by experimental ₁₅₁
studies [8, 46, 47]. Even though some sub-groups of inhibitory neurons have been shown ₁₅₂
to be broadly tuned to orientation [46], on average inhibitory neurons are well ₁₅₃
tuned [46, 47], just as in our model. Indeed, Nowak *et al.* [46] found that inhibitory ₁₅₄
layers are slightly more broadly tuned than excitatory ones (mean HWHH of regular ₁₅₅
spiking 23.4° vs. fast spiking neurons 31.9°). Our model follows the same trend (mean ₁₅₆
HWHH of excitatory neurons 23.4° vs. inhibitory neurons 26.0°). ₁₅₇

Fig 6. Orientation tuning properties of excitatory and inhibitory neurons in Layers 4
and 2/3. (A) Orientation tuning curves calculated as the mean firing rate of the neuron
at the given orientation. The large tuning curves on the left represent the mean of
centered orientation tuning curves across all the measured neurons of the given type.
The smaller tuning curves on the right correspond to randomly selected single neurons.
(B) Half-width-at-half-height (HWHH) at 100% vs 10% contrast. (C) The distribution
of HWHH measured at 100% contrast across the four modeled cortical populations. (D)
The distribution of relative unselective response amplitude (RURA) measured at 100%
contrast across the four modeled cortical populations.

Because the HWHH measure does not take into account the unselective portion of ₁₅₈
the response, cells that respond substantially above the spontaneous rate at all ₁₅₉
orientations, but where this unselective response is topped by a narrow peak, will yield ₁₆₀
low HWHH. To address this concern, Nowak *et al.* [46] also calculated the relative ₁₆₁

unselective response amplitude (RURA; see Methods), and showed that the RURA of ₁₆₂ inhibitory neurons is considerably broader than that of excitatory ones (mean for ₁₆₃ regularly spiking neurons 2.5% vs. fast spiking neurons neurons 18.1%). We have ₁₆₄ repeated this analysis, and found that the difference between the RURA of excitatory ₁₆₅ and inhibitory model neurons (Fig 6CD; -0.47% in excitatory vs 10.43% in inhibitory ₁₆₆ neurons) is indeed more pronounced than their difference in HWHH. ₁₆₇

Most excitatory and inhibitory neurons in the model cortical Layer 4 and 2/3 exhibit ₁₆₈ contrast invariance of orientation tuning width (Fig 6AB). On average we observe very ₁₆₉ minor broadening of the tuning curves at high contrast: the mean HWHH differences ₁₇₀ between low and high contrasts is 0.19° (excitatory, Layer 4), 0.75° (inhibitory, Layer 4), ₁₇₁ −0.05° (excitatory, Layer 2/3), and 1.06° (inhibitory, Layer 2/3). Such minor ₁₇₂ broadening has been observed experimentally in Simple cells in cat V1 (0.3°; layer ₁₇₃ origin and neural type not known [8]). ₁₇₄

Overall, the orientation tuning in our model is in good quantitative agreement with ₁₇₅ experimental data, with both excitatory and inhibitory neurons exhibiting sharp and ₁₇₆ contrast-invariant orientation tuning across all modeled layers, in contrast to many ₁₇₇ previous modeling studies that relied on untuned inhibition, or broad ₁₇₈ contrast-dependent inhibition [5, 6, 48]. See the Discussion section 5.4 for a mechanistic ₁₇₉ explanation of how these phenomena arise in this model in comparison to others. ₁₈₀

4.2.2 Orientation tuning of sub-threshold signals ₁₈₁

To better understand the mechanisms of orientation tuning in the model, we ₁₈₂ investigated the tuning of the membrane potential and the excitatory and inhibitory ₁₈₃ synaptic conductances. An essential mechanism responsible for orientation tuning in ₁₈₄ this model is the push-pull connectivity bias in Layer 4. In theory, a cell with a pure ₁₈₅ push-pull mechanism with perfectly balanced excitation and inhibition should exhibit ₁₈₆ perfectly linear (Simple cell) behavior, where orientation tuning is solely driven by the ₁₈₇ luminance modulation of the DG, and the mean membrane potential remains at rest at ₁₈₈ all orientations. To assess to what extent this idealized scheme is true in this model, we ₁₈₉ have plotted the orientation tuning curves of the mean (DC) and first harmonic (F1) ₁₉₀ components of the sub-threshold signals across the different neural types and layers in ₁₉₁ Fig 7. ₁₉₂

In Layer 4 the orientation tuning of the membrane potential is dominated by the F1 ₁₉₃ component, in line with the presence of strong push-pull mechanisms in this model layer ₁₉₄ (Fig 7 B). Layer 4 cells also have significant tuning of the F0 component (mean ₁₉₅ membrane potential), but of lesser magnitude (Fig 7 A). The F0 components of the ₁₉₆ membrane potential show noticeable broadening at higher contrast, consistent with ₁₉₇ experimental observations that, unlike the spiking response, the orientation tuning ₁₉₈ width of the mean of membrane potential is not contrast independent [8]. In Layer 2/3 ₁₉₉ the magnitude of the membrane potential components is reversed: the magnitude of the ₂₀₀ F0 component of the membrane potential is stronger than the F1 component. This is ₂₀₁ consistent with the lack of phase specific connectivity in Layer 2/3, and explains the ₂₀₂ predominance of Complex cells in this model layer (see Sec 4.2.3). ₂₀₃

To understand how the interplay between excitation and inhibition leads to the ₂₀₄ observed membrane potential tuning characteristics in Fig 7C-F, we plot the orientation ₂₀₅ tuning curves of the F0 and F1 components of excitatory (C,D) and inhibitory (E,F) ₂₀₆ synaptic conductances. In Layer 4, the most obvious difference is that, in comparison to ₂₀₇ the tuning of membrane potential, the F0 component of both excitatory and inhibitory ₂₀₈ conductances is not dominated by the F1 component. The weaker F0 component of the ₂₀₉ membrane potential in Layer 4 neurons is thus due to the cancellation between the ₂₁₀ excitatory and inhibitory F0 components, while the F1 component of the membrane ₂₁₁ potential orientation tuning remains strong, as cancellation between excitatory and ₂₁₂

Fig 7. Orientation tuning of membrane potential and excitatory and inhibitory conductances of excitatory and inhibitory neurons in Layer 4 and 2/3. Each row shows the average orientation tuning curves of all neurons of given neural type recorded in the given layer (see row labels on the left), with spontaneous level of each respective neural signal subtracted. Orientation tuning curves of: (A) F0 component of membrane potential; (B) F1 component of membrane potential; (C) F0 component of excitatory synaptic conductance; (D) F1 component of excitatory synaptic conductance; (E) F0 component of inhibitory synaptic conductance; (F) F1 component of inhibitory synaptic conductance. Low contrast: 10% (black line), high contrast: 100% (blue line).

inhibitory conductances does not occur due to the half-period phase shift between them   213
(see Fig 5A,B). In Layer 2/3 both F0 and F1 components of excitatory and inhibitory   214
conductances in both excitatory and inhibitory neurons are well tuned, but the F0   215
components dominate in magnitude the F1 components for both excitatory and   216
inhibitory conductances and for both neural types.   217

To the best of our knowledge, this is the first model where orientation tuning of both   218
F0 and F1 components of membrane potential and conductances is exhibited in all   219
layers and cell types, consistent with experimental observations [49, 50]. The contrast   220
invariance in orientation tuning arises as a complex interplay between excitation and   221
inhibition, differs between layers and neural types, and, in the model Layer 4, is   222
consistent with engagement of a push-pull mechanism [5]. However, additional   223
enhancement of activity is present due to the recurrent facilitatory dynamics among   224
neurons of similar orientation, which goes beyond the predictions of the push-pull   225
model, as evidenced by the presence of tuning in the F0 components of the membrane   226
potential of layer 4 neurons, and which accounts for the small residual broadening of   227
orientation tuning curve at low contrast.   228

4.2.3   Simple and Complex cell types   229

Next we examined the classification of the cells into Simple and Complex. The   230
modulation ratio (MR) of the peri-stimulus time histogram (PSTH) classified most   231
model Layer 4 neurons as Simple cells (see Fig 8A) and most Layer 2/3 neurons as   232
Complex cells (see Fig 8A), in-line with the experimentally observed laminar biases in   233
cat [51]. When pooled across the two layers the histogram of MRs forms the   234
characteristic experimentally observed bimodal distribution (see Fig 8A; [52]).   235

Fig 8. MRs of excitatory cells in the model. (A) Histogram of MRs calculated from the
PSTH over recorded excitatory cells in Layer 4 (first row), Layer 2/3 (second row) and
pooled across both layers. (B) as in (A) but MRs calculated from the membrane
potential after subtraction of background level. (C) As in A but the histograms are of
the absolute value of the F0 component of the membrane potential. (D) As in C but for
the F1 component of the membrane potential. (E) The relationship between the MR
calculated from PSTH and from membrane potential. In the pooled histogram plots
cells classified as Simple are marked white and as Complex are marked black.

Unlike for the PSTH, the MRs computed from the membrane potential (with the   236
resting potential subtracted) and pooled across the cortical layers form a unimodal   237
distribution (see Fig 8B), in line with experimental evidence in cat [52]. When the MRs   238
calculated from PSTH and membrane potential are plotted against each other (Fig 8E),   239

a characteristic hyperbolic relationship is revealed, in line with the observations of Priebe *et al.* [52].

We also analyzed the F0 and F1 components of the membrane potential individually. Both have a unimodal distribution as observed experimentally by Priebe *et al.* [52]. The range of values for the F1 component fits the experimental data well, as Priebe *et al.* [52] observes that most complex cells have an F1 component lower than 5 and that only few simple cells have an F1 component higher than 8. On the other hand, the distribution of the F0 component of membrane potential (Fig 8C,D) is narrower in both modeled layers than observed experimentally [52]. This could be due to intrinsic regularities present in the model, such as identical physiological parameters among all cells of the same cell type, or limited variability of the ratio of afferent and recurrent inputs to model Layer 4 neurons, or completely phase-non-specific connectivity from Layer 4 to 2/3. These artificial inherent regularities of the model could be exaggerating the differences between the two functional neural types.

4.3   Size tuning

Next we studied the response of model neurons to DGs confined to an aperture of increasing diameter, to assess the presence of surround suppression in the model. While most excitatory neurons in both cortical layers show surround suppression (Fig 9A-D), we observe a diversity of tuning patterns. An example of a Layer 4 cell with prototypical size tuning properties, exhibiting the expansion of facilitation diameter at low contrast [53, 54], is shown in the first row of Fig 9, whereas the second row of Fig 9 shows an example of a Layer 4 cell that exhibits only weak suppression. Neurons in Layer 2/3 also express classical size tuning effects, as exemplified in the third row of Fig 9.

The degree of the suppression varies widely in both cortical layers (Fig 9D). Overall the suppression is stronger in Layer 2/3 (mean suppression index (SI) at high contrast: 0.19 ± 0.004) than in Layer 4 (mean SI at high contrast: 0.10 ± 0.010). These values are on the lower end of the range of experimental observations in cat V1 (i.e. 0.16 [54], 0.35 [55], 0.44 [53] and 0.47 [56]; cells pooled across all cortical layers). This could come from the absence of higher visual areas in our model, which feedback connections have been shown to contribute positively to surround suppression in macaque V1 [57]. Stronger suppression in Layer 2/3 than in Layer 4 was observed in three studies [55, 56, 58] while [53] did not observe a statistically significant difference between the layers. Finally, Wang *et al.* [53] observed a decrease of suppression at low contrast, which is consistent with the results of the Layer 2/3 of our model (mean SI at low contrast in Layer 2/3: 0.12 ± 0.006; mean SI at low contrast in Layer 4: 0.10 ± 0.011).

At high contrast, we observe that the mean maximum facilitation diameters (MFD) for Layer 4 and Layer 2/3 neurons are respectively $2.30° \pm 0.06°$ and $2.00° \pm 0.03°$ (Fig 9G). It is difficult to precisely compare these numbers to experimental data, as RF size in cat V1, and hence MFD, grows with neurons' foveal eccentricity [59], but we are not aware of any study that reports eccentricity-dependent data on size tuning. Studies have reported a broad range of mean MFD values ($2.9°$ [60], $3.5°$ [58], $6°$ [54]). The pooled mean MFD we observe in our study is on the lower end of this range, which is consistent with the fact that all these studies recorded neurons at (on average) larger eccentricities than the $3°$ – at which our model is centered (see Methodology). We observe important trends in the model that are consistent with experimental data. First, we observe an increase of the MFD at lower contrast (Fig 9G; mean MFD at low contrast: $2.84° \pm 0.06$ Layer 4 and $2.63° \pm 0.04$ Layer 2/3). This change of 27% (pooled across layers) is in good agreement with Wang *et al.* [53] and Tailby *et al.* [54] who found increases of 33% and 36% respectively in cat V1. Second, we observe smaller MFD in Layer 2/3 than in Layer 4 in line with experimental observations [58]. Finally,

Fig 9. Size tuning in excitatory neurons. The first three rows show size tuning in three example model neurons. Top row is a typical Layer 4 cell with strong size tuning, second row shows an example of Layer 4 cells that does not exhibit size tuning, and the third row shows a typical Layer 2/3 cell. (A) The size tuning curve of trial-averaged spiking responses. Dots are measured responses, line is fitted size tuning curve. (B) The size tuning curve of trial-averaged mean membrane potential. (C) The size tuning curves of trial-averaged mean excitatory (red) and inhibitory (blue) synaptic conductances. Data shown for low and high contrast conditions: (A-B) low-contrast (10%) condition is black, high-contrast (100%) condition is blue; (C) low-contrast (10%) condition is un-saturated, high-contrast (100%) condition is saturated; (D-H) Scatterplots of size-tuning measures. Arrows mark means after pooling across layers. White (empty in H) dots correspond to individual Layer 4 neurons and black (full in H) dots correspond to Layer 2/3 neurons. (D) The suppression index of spike responses at low vs. high contrast. (E) The counter-suppression index of spike responses at low vs. high contrast. (F) The suppression index vs. the counter-suppression index at high contrast. (G) the maximum facilitation diameter at low vs. high contrast. (H) The suppression index of excitatory (red) and inhibitory (blue) synaptic conductances at low vs. high contrast.

we find that both excitatory and inhibitory conductances are size tuned in both simulated cortical layers (Fig 9H), in line with experimental evidence [61,62].

Many neurons do not exhibit monotonic suppression after reaching their classical RF size: after a certain diameter their responses partially recover from the suppression [53]. This phenomenon has been named counter-suppression. We observe this in a substantial subset of model neurons across both modeled layers (Fig 9E), as shown also in some of the single neurons examples (first row in Fig 9A). When quantifying the magnitude of this counter-suppression as the CSI index, we find that it is stronger at low contrast (mean CSI at low contrast: 0.05 ± 0.009 Layer 4 and 0.05 ± 0.005 Layer 2/3) than at high contrast (mean CSI at low contrast: 0.03 ± 0.006 Layer 4 and 0.02 ± 0.004 Layer 2/3), which is in line with Wang *et al.* [53]. It should be noted, however, that the limited area of the simulated cortical network likely prevents the full build-up of counter-suppression effects.

4.4 Trial-to-trial variability: response to natural images with simulated eye-movements

Next we probed the model with a stimulus consisting of a natural scene animated with simulated eye-movements, as described in the intra-cellular study of Baudot *et al.* [21]. As can be seen in Fig 10B, the natural image stimulus elicits a highly repeatable response in cat V1, both at the level of spikes and at the level of sub-threshold responses. In contrast, the response to presentation of a DG of medium contrast (30%) is only locked to the slow temporal frequency of the luminance modulation (Fig 10A). This stimulus dependent difference in response statistics is qualitatively consistent with the findings of Baudot *et al.* [21]. The presence of a strong short-term synaptic depression in the thalamo-cortical connections shapes the response in a stimulus-dependent way, as it is less triggered by the transient changes in the natural scene stimulus than by the sustained activation induced by the DG [63].

To further investigate the response precision and reliability of the model neurons we computed the cross-correlation between trials of the spiking and membrane potential responses. The reliability is given by the peak amplitude of the cross-correlation (a) at time zero, and the temporal precision by the standard deviation of the Gaussian fit (σ) [21]. As shown in Fig 10C,D, for the spiking response (top), both for model Layer 4 and Layer 2/3 neurons, the cross-correlation has a higher peak and is narrower for the natural-image-with-eye-movement (NI) stimulus (Layer 4: $a = 0.221$, $\sigma = 12$; Layer 2/3: $a = 0.081$, $\sigma = 15$), than for DGs at medium contrast (Layer 4: $a = 0.187$, $\sigma = 60$; Layer 2/3: $a = 0.037$, $\sigma = 60$), in line with the experimental results [21] (see Fig 10F).

For the membrane potential (Fig 10C,D bottom) the same relationship holds except that for Layer 4 neurons the peak (and thus the reliability) for gratings is slightly higher than for natural images (Layer 4 NI: $a = 0.441$, $\sigma = 18$; Layer 2/3 NI: $a = 0.064$, $\sigma = 15$; Layer 4 DG: $a = 0.63$, $\sigma = 99$; Layer 2/3 DG: $a = 0.025$, $\sigma = 71$), unlike the observations in [21] (Fig 10F), where the reliability is higher for the NI stimulus for both spikes and membrane potential.

Next, Baudot *et al.* [21] found that relative to the trial-to-trial variability of membrane potential during spontaneous activity, the variability of membrane potential during DG stimulation increases, while it decreases during NI stimulation (Fig 10J). The results of the same analysis in the model are shown in Fig 10(GH). In both model layers we observe an increase of stimulus-locked variance (calculated as the inverse of the stimulus-locked time-averaged standard deviation across trials, and expressed as a percentage relative to the value for ongoing activity) for DG ($91 \pm 1.05\%$ in Layer 4 and $88 \pm 0.2\%$ in Layer 2/3). For the NI condition, we observe a decrease of stimulus-locked variance in Layer 4, as well as a small increase in Layer 2/3 ($130 \pm 0.53\%$ in Layer 4 and

Fig 10. Trial to trial reliability of responses to a natural image with simulated eye movements (NI) and to drifting sinusoidal gratings (DG). (A) The response of a randomly sampled excitatory neuron in model Layer 4 to the DG stimulus (contrast at 30%). The top panel shows the spike raster histogram, the middle panel shows the incoming synaptic excitatory (red) and inhibitory (blue) conductances. The bottom panel shows the membrane potential. The thick lines show the mean of the analog signals over the 10 trials. (B) As (A), but for the NI stimulus. (C-F) Cross-correlation across trials for spikes (top) and subthreshold membrane potential activity (bottom), in response to DG and NI stimuli. The reliability is given by the peak amplitude at time zero, and the temporal precision by the standard deviation of the Gaussian fit to the cross-correlation. (C) Results averaged across all recorded excitatory neurons in model Layer 4. (D) Results averaged across all recorded excitatory neurons in model Layer 2/3. (E) Results averaged across all recorded excitatory neurons pooled across the two model layers. (F) The experimental data from [21]. Note that in that study two additional stimuli were also presented (dense noise : blue line; grating with simulated eye-movements : green line) that we have not used in this study. (G-I) The inverse of the stimulus-locked time-averaged standard deviation of membrane potential (1/SD) across trials, averaged across cells, relative to the value for ongoing activity. (G) Results averaged across all recorded excitatory neurons in model Layer 4. (H) Results averaged across all recorded excitatory neurons in model Layer 2/3. (I) Results averaged across all recorded excitatory neurons pooled across the two model layers. (J) Experimental results in cat [21].

97 $\pm$ 0.22% in Layer 2/3). However, note that in Baudot *et al.* [21] the results are ₃₄₁ pooled across neurons of all layers, and the same treatment of our simulation data yields ₃₄₂ a good quantitative match to the experimental data (Fig 10I). ₃₄₃

 The lack of direct external noise injected into cortical model neurons was crucial for ₃₄₄ achieving the results presented in this section. The trial-to-trial variability in the model ₃₄₅ is primarily shaped by the intra-cortical interactions during both spontaneous and ₃₄₆ stimulus driven activity. This is an advance over a previous model of a V1 column that ₃₄₇ explained the precision and reliability changes between the grating and naturalistic ₃₄₈ stimuli at the spiking level, but did not reproduce the stimulus-dependent variability ₃₄₉ effect at the level of membrane potential [63]. This was due to the inclusion of ₃₅₀ substantial random spiking input directly into cortical neurons, which induced ₃₅₁ substantial stimulus-independent variability of the membrane potential, overriding the ₃₅₂ internally generated variability components. These differences point out the importance ₃₅₃ of simultaneously capturing within a single model the cortical processing under multiple ₃₅₄ stimulation conditions and at different levels of signal integration, to be able to fully ₃₅₅ describe its operation. ₃₅₆

5   Discussion ₃₅₇

A coherent explanation for how the numerous V1 computations coexist within the ₃₅₈ underlying neural substrate is still lacking. We have approached this problem with a ₃₅₉ systematic, integrative, data-driven methodology, constructing a model of V1 firmly ₃₆₀ grounded in anatomical and physiological data. We show that a single set of plausible ₃₆₁ connectivity principles — (i) weakly orientation-biased thalamic input, (ii) local, weakly ₃₆₂ push-pull-biased intra-Layer 4 connections, and (iii) long-range weakly ₃₆₃ orientation-specific intra-Layer 2/3 connections — give rise to the most salient V1 ₃₆₄ computations. This single model instance provides insight into the underlying ₃₆₅ mechanisms of processing within the classical receptive field: orientation tuning of both ₃₆₆ spiking and subthreshold signals (Fig 6 and 7); and emergence of Simple and Complex ₃₆₇ cells (Fig 8). The same model gives an accurate account of computations requiring ₃₆₈ integration beyond the classical receptive field: size tuning of both spikes and synaptic ₃₆₉ conductances (Fig 9). Finally, beyond artificial stimuli, the model replicates the ₃₇₀ stimulus-dependent changes to the precision of spiking responses, and to the ₃₇₁ trial-to-trial variability of membrane potential (Fig 10) when stimulated by a ₃₇₂ naturalistic spatio-temporal stimulus. All these stimulus evoked properties are ₃₇₃ underpinned by a resting state that is in very good quantitative agreement with ₃₇₄ physiological data at multiple levels of integration (Fig 4). ₃₇₅

5.1   Integrative computational modelling approach ₃₇₆

To support model component reuse, and hence the long-term integrative modeling ₃₇₇ program proposed in this study, we built our model in the Mozaik [17] neural network ₃₇₈ modeling environment, optimized for efficient specification and reuse of model ₃₇₉ components, experimental and stimulation protocols and model analyses, and are ₃₈₀ making the resulting model code available under a liberal open-source license. Although ₃₈₁ neuroscience should ultimately converge upon a single model, there are many paths ₃₈₂ through the space of partial models to arrive there, and for the health of the field it is ₃₈₃ good to have several "competing" models, each of which will make different ₃₈₄ approximations [15]; nevertheless, all models should pass the same validation tests. We ₃₈₅ therefore also make available our library of integration tests via a dedicated data store ₃₈₆ (*http://v1model.arkheia.org*) implemented in the recent Arkheia framework [64], sharing ₃₈₇ model and virtual experiment specifications. The integration tests are structured so as ₃₈₈

to be independent of the model specification, and their ready-to-use implementation in the Mozaik environment is provided. Altogether, we provide here an advanced and ready-to-use platform for future long-term, systematic, incremental, integrative research on the early visual system of higher mammals and beyond.

5.2   Novelty and predictions

The central contribution of this study is the original convergence of the following key integrative principles: (i) the multitude of anatomical constraints respected, (ii) the multiplicity of neural signals (synaptic, single unit) and integration levels (conductance, neuronal, columns) validated, (iii) the multiplicity of the stimulation protocols tested, (iv) the diversity of functional phenomena explained, and (v) the multiplicity of spatial and temporal scales validated. While many of the properties demonstrated in this study have been previously modeled in isolation, to the best of our knowledge, the single V1 model instance presented here was validated with the broadest range of visual stimulation protocols to date (Table 1). As such, the present study represents a significant advance towards providing a comprehensive integrative treatment of the early visual system.

This model also provides a description of the asynchronous irregular resting regime that is quantitatively accurate at both supra- and sub-threshold levels, does not rely on *ad-hoc* noise input to cortex, and is compatible with evoked processing. This allows us to demonstrate, how the trial-to-trial variability of membrane potential in V1 neurons can increase during drifting grating stimulation and decrease during naturalistic movie stimulation [21].

This model also provides a mechanistic explanation of orientation tuning that is simultaneously consistent with all the following physiological constraints: (i) orientation tuning is sharp across cortical layers [47], (ii) inhibition is only moderately broader than excitation [47,65], and (iii) both DC and F1 components of spikes, membrane potential and conductances tend to be tuned to the cell's preferred orientation. Many past models have only explored the orientation tuning of spiking responses [66,67] and relied on either broad inhibition [5,48,68] or inhibition tuned to the orthogonal orientation [69,70], which is in contradiction with inhibitory conductances tuned to preferred orientation [49]. Few past models have been consistent with the constraint of sharply tuned responses of inhibitory cells and of inhibitory input tuned to the preferred orientation [71], and no modelling studies demonstrated the orientation tuning of both the DC and F1 components of all the sub-threshold signals.

The model offers a number of testable predictions about properties of cat V1:

1. Layer- and cell-type-specific differences in resting regime:

   (a) Higher inhibitory conductances in Layer 2/3 than in Layer 4.

   (b) Greater synchrony among the inhibitory neurons.

2. Layer-specific distributions of MRs of spiking and membrane potential responses (layer specific data is missing in cat).

3. Near contrast-invariance of inhibitory neurons.

4. Layer-specific differences in sub-threshold signal orientation tuning.

5. Layer-specific differences in conductance size tuning.

6. Important testable assumptions were made about the model parameters:

   (a) The sharpness of functional specificity of cortico-cortical connections.

	This study	Troyer 1998	Somers 1998	Wielaard series	Tononi series	Rangan series	Kremkow 2016	Chariker series
Spontaneous state without ad-hoc noise source	✓	✗	✗	✗	✓	✗	✗	✓
Realistic levels of cond. in the spontaneous state	✓						✓	
Log-normally distributed spont. firing rates in L2/3	✓							
Realistic mean rate in spontaneous state in L4	✓				✓			✓
Realistic mean rate in spontaneous state in L2/3	✓	○	○	○	✓	○	○	○
Spontaneous patterns of cortical activity					✓	✓		
Bimodal distribution of MR (spikes)	✓	✗		✓	✗	✓	✗	✓
Unimodal, monotonic distribution of MR (Vm)	✓			✓		✓		
Layer 4 orientation tuning width	✓	✓			○	✓	✓	✓
Layer 2/3 orientation tuning width	✓	○	○	○	○	○	○	○
Contrast invariant orientation tuning	✓	✓			○	✓	✓	
Conductance increase at onset & offset of flashed bar					✓			
Anti-phase cond. in Simple cells responding to DG	✓	✓	✗	✗	✗		✓	✗
Spatial frequency selectivity					✓			✓
Direction selectivity (DS)								✓
DS broadband across spatial frequencies								✓
DS broadband across temporal frequencies								✓
Precision and reliability of spike responses to NI	✓						✓	
Stimulus dependent trial-to-trial variance of Vm	✓						✗	
Gamma band synchronization						✓		✓
Size tuning	✓	○	✓	✓			○	
Contrast dependent surround-suppression	✓	○	✓	✓			○	
Contrast dependent shift in size tuning peak	✓	○	✓	✓			○	
Size-tuning of exc. and inh. conductances	✓	○		✓			○	
Size-tuning counter-suppression	✓	○					○	
Line-Motion illusion		○				✓	○	
Slow Wave Sleep mode with up & down states					✓			
Bi-modal distribution of Vm in sleep mode					✓			
Slow oscillation propagation in cortical space					✓			

Table 1. Validation of visual cortex models with respect to experimentally observed functional properties. A green tick means that the model gives a qualitatively good match to experimental findings. A red cross means that the model gives a response which is qualitatively different from experimental findings. A red open circle means that the model does not incorporate neural structures necessary for this feature to be meaningfully tested. Empty spaces mean that the model was not tested for the property, and it is not possible with certainty to conclude from the corresponding paper that the model does or does not match the corresponding experimental data.

(b) The excitatory-to-excitatory vs. excitatory-to-inhibitory pathway strength ratio.

5.3   Other integrative models of V1

In the following, we present a cross-comparison of the presented model with the most prominent integrative models of V1 to date. We restrict this comparison to models of cat and macaque V1, as the two species represent the two major animal models for vision in higher mammals, while their anatomical and functional similarity allows for straightforward comparison.

Somers *et al.* [12] investigated a topologically-organized model of V1, which incorporated excitatory connections that dominated at very short and long distances, and inhibitory ones that dominated at intermediate distances. Direct anatomical evidence for such an arragment of lateral connections is lacking [19]. All connection probabilities were biased towards co-oriented targets, but were independent of the phase preference of targets, which implies that the model was not able to replicate the anti-phase relationship between excitatory and inhibitory conductances [44,49,72]. The Somers *et al.* model was able to demonstrate the emergence of contrast-invariant orientation tuning, and contrast dependence of size tuning, but none of the other features of V1 processing considered here were examined.

Another similar pair of modeling studies was performed by Wielaard *et al.* [73,74]. These modeling studies did not consider long-range cortical interactions, nor functional specificity of intra-cortical interactions, implying that the models were not able to replicate the anti-phase of excitatory and inhibitory conductances [44,49,72]. Wielaard *et al.* demonstrate the emergence of both Simple and Complex cell types and show how a bimodal distribution of MRs at the level of spiking emerges despite unimodally distributed MRs of membrane potential. Furthermore, they show how such an only-locally-connected cortical model can explain the emergence of size tuning, quantitatively well matching experimental results. Interestingly, the authors do not demonstrate some of the more classical properties of V1 neurons, such as sharp mean orientation tuning (the limited data presented indicate a predominance of poorly tuned cells) or its contrast invariance. Similar to our model, Wielaard *et al.* demonstrate that the size tuning arises as a complex dynamic interplay between excitation and inhibition, whereby both conductances exhibit reduction at large stimulus sizes. Unfortunately Wielaard *et al.* do not assess the sub-threshold signals quantitatively, or characterize the background activity in the model. Finally, no naturalistic stimuli were tested in the model.

The Tononi group has also published a series of models focusing on the cat early visual system [75–78]. In addition to the LGN and cortical layers 4 and 2/3 considered in our model, the Tononi *et al.* model series also incorporated a model of sub-granular layers and a rudimentary model of a generic extra-striate area that represented the totality of feedback input from higher-level visual areas to V1. Tononi *et al.* aimed primarily at simulating the spontaneous activity during awake vs. slow wave sleep conditions. Accordingly, the model incorporated only rudimentary functionally specific circuitry, chiefly restricted to the classic Hubel & Wiesel type convergence of thalamic afferents onto cells in granular and infragranular layer that induced the most basic form of orientation selectivity. Only two orthogonal orientations assigned to two spatially separated cell groups were considered, and no other investigation of evoked activity was performed. On the other hand, the model demonstrated an impressive range of activity dynamics present in awake and slow wave sleep brain states including spontaneous patterns of cortical activity, slow wave sleep mode with up & down states, the slow oscillation propagation in cortical space, bi-modal distribution of membrane potential in sleep mode and gamma band synchronization in both sleep and awake modes.

One of the most comprehensive V1 modeling effort prior to the present study is a series of modeling studies [79–81] summarized in Rangan *et al.* [13], architecturally very similar to the Wielaard *et al.* models [73, 74]. While the models also focus on spontaneous regime dynamics, they rely on constant independent noisy spike input in the cortical neurons. In multiple related models the authors have demonstrated a range of V1 properties in both spontaneous and evoked states including a fluctuation-driven spontaneous activity regime characterized by low firing rates [79], emergence of sharply tuned Simple and Complex cell types with a bimodally distributed MR of spike responses and unimodally distributed MR of membrane potential [80], and approximate contrast invariance of orientation tuning. The study also demonstrated two features of V1 computation not examined in our model: the spontaneous activity dynamics that are correlated with the intrinsic orientation organization of cortex [79], and the spatio-temporal patterns of cortical activity in response to a line-motion illusion stimulus [81]. Unlike in the present study, neither the contextual modulation nor naturalistic stimulation were explored in these models. Most importantly, a detailed quantitative account of their sub-threshold behavior was not provided. Finally, while all the models summarized in Rangan *et al.* [13] share a common architecture, it is unclear whether all these studies used identical model parameterizations, and hence if all these features could be achieved within a single model instance.

Finally, recently Chariker *et al.* [48] have also made the case for a more comprehensive approach to the computational study of V1, presenting a model that is, similarly to the present study, firmly grounded in anatomical data. Their work is complementary to ours as they model layer 4cα of monkey V1, which receives sparse input from magnocellular cells. Their model displays both gamma oscillations [82] and realistic direction selectivity [83], which we did not investigate in our model. The authors do not probe the model with any stimulus other than the ubiquitous fullfield DG, and therefore do not investigate contextual modulation effects nor the response of their model to natural images. The characterization of the spontaneous state is missing. The model Simple cells also do not exhibit the anti-phase relationship between excitatory and inhibitory conductances. The authors do investigate contrast dependence of orientation tuning, but only by examining the ratio of the orthogonal response to the preferred response, a measure to which V1 neurons are naturally not contrast invariant to [84]. It is therefore unclear whether the neurons in their model display contrast-invariance of the HWHH measure instead. Finally, the analysis of the model properties at the sub-threshold level is limited.

5.4   Other models of V1

Models of orientation selectivity are typically categorized according to their circuitry ('feed-forward' vs 'recurrent'), the relative impact of such feed-forward vs recurrent contributions on the genesis of orientation selectivity, and the selectivity of the cortical inhibition. In 'feed-forward' models [5, 66, 85] orientation tuning is induced by a strong sharply tuned thalamo-cortical pathway, while intra-cortical connections have only a minor influence. In contrast, 'recurrent' models [86, 87] assume weak poorly tuned thalamic input but strong intra-cortical interactions. With respect to inhibition, the defining features are its orientation tuning, its presence or absence at orthogonal orientations, and its relationship to stimulus phase.

Although the opposition between thalamic-driven and cortical-recurrent computing schemes has long dominated the literature, this dichotomic categorization is highly oversimplified [7]. Following the anatomical and physiological evidence to determine the model architecture, as we have done here, results in a model that resides in between these standard classes. The thalamo-cortical connections in the model constitute a fraction of the synaptic input to Layer 4 ($\sim 10\%$), which would assign the model to the

recurrent category. On the otger hand, the dominating intra-cortical connections in Layer 4 are biased towards a push-pull organization, typically assumed in feed-forward models [5] (although, based on experimental evidence [20, 88], this functional bias is weaker than in previous models). Consequently, the inhibitory input to Layer 4 neurons is broad and present at orthogonal orientations. But the resulting orientation tuning of inhibitory neurons is sharp, comparable to that of excitatory ones, showing that even weak and poorly orientation-tuned afferent input and broad inhibitory input can lead to sharply tuned inhibitory neurons, in line with experiments [47, 49].

Furthermore, in classical feed-forward models the DC component of the feed-forward excitatory input is canceled-out by the DC component of the inhibition [5]. It is the F1 component of the membrane potential that induces the orientation tuning. But, in line with experimental evidence, our model still exhibits orientation tuning of the DC components of both membrane potential and excitatory and inhibitory conductances, suggestive of a boosting of the mean response at preferred orientations, typical of recurrent models. However, the push-pull interactions remain an important factor in the generation of contrast-invariant orientation tuning, as evidenced by the dominance of the F1 components in Simple cells. This shows that different strategies of generating orientation tuning can co-exist within the same neural circuit, facilitating the models' consistency with a broader range of physiological findings, hence exemplifying the benefits of the data-driven integrative approach in counteracting the fragmentation of knowledge.

Recently, a feed-forward explanation of contrast-invariant orientation tuning was proposed by Finn *et al.* [8], relying on two key mechanisms: an expansive non-linearity governing the relationship between membrane potential and the spike rate, and the contrast dependence of the thalamic input. However, although the formulation of the Finn *et al.* model is feed-forward, the assumed non-linearities may in the cortical substrate be implemented by recurrent dynamics. Furthermore, the Finn *et al.* model does not explain the fact that the modulation of the membrane potential is dominated by inhibition, nor the anti-phase relationship between inhibition and excitation [44, 49, 72].

Finally, a recent study from Arkhipov *et al.* [89] also followed a comprehensive, data-driven approach, employed a broad set of test protocols, and demonstrated orientation selectivity, amplification of thalamic inputs, presence of gamma oscillations, and log-normal distributions of firing rates. Unlike the present study it did not investigate more complex functional properties, such as size-tuning or trial-to-trial precision. The Arkhipov *et al.* model is complementary to the model presented here: cat V1 in the anesthetized state vs awake mice [89] (respective merits of the two preparations are discussed in section 5.5). The Arkhipov *et al.* study also incorporated neural morphology, although, they showed that a simplified, integrate-and-fire version of their model remained in good agreement with *in-vivo* data, justifying the level of detail chosen in the present study.

5.5 Modeling choices

We have chosen to focus our model on the cat visual system, but supplement the model with data from macaque whenever equivalent data from the cat is not available, as the closest available approximation. The most extensive body of data on V1 has been collected in cat, including several of the key datasets for this study, such as extensive parametric studies of visual RFs, the extents of inter-areal connectivity in V1 [19] and the relative strength of the different intra-areal pathways [18]. Furthermore, there is greater availability of in-vivo intracellular data, which are critical for constraining our model, from the anesthetized cat [44, 49, 72, 90–92]. Finally, cat area 17 can be equated with macaque area V1, based on criteria of relative position in the cortical mantle,

internal organization of visual field representations, and trans- and sub-cortical ₅₈₇ connections. ₅₈₈

The level of biological detail of the presented model implies a substantial set of ₅₈₉ model parameters that have to be determined. The process of parameter determination ₅₉₀ was divided into two stages. In stage one we first determined the parameter values or ₅₉₁ ranges for all model parameters from the experimental literature. Through this process ₅₉₂ we obtained constraints for an absolute majority of model parameters including: (i) ₅₉₃ physiological parameters of the neural model which were primarily based on intracellular ₅₉₄ patch clamp recordings in [44] (see section 6.1.1), (ii) spatial distributions of local ₅₉₅ connectivity between different cortical layers and neural types [93] (see section 6.1.4), ₅₉₆ (iii) functional specificity of layer 2/3 connectivity, based on [20] (see section 6.1.5), and ₅₉₇ (iv) synaptic delays, based on a set of paired recording studies (see section 6.1.7). ₅₉₈

Unfortunately, current experimental evidence is insufficient for precise determination ₅₉₉ of every single model parameter. In this study we hypothesize the existence of push-pull ₆₀₀ connectivity among Layer 4 neurons, as it is the only plausible explanation for the ₆₀₁ anti-phase relationship between excitatory and inhibitory conductances in Simple ₆₀₂ cells [21, 94] proposed to date (see section 6.1.5). As this circuitry has not been directly ₆₀₃ probed, its hypothetical parameters are unknown. Another poorly constrained ₆₀₄ parameter is the unitary synaptic strength between neurons. Existing experiments have ₆₀₅ reported a wide range of values, only indicative of the order-of-magnitude of the ₆₀₆ synaptic strength, and a systematic identification of the synaptic strength with respect ₆₀₇ to pre- and post-synaptic neuron type and layer membership is completely missing (see ₆₀₈ section 6.1.6). ₆₀₉

In stage two we fine-tuned the parameters that were not fully determined in stage ₆₁₀ one. The tuning was guided primarily by the goal of achieving stable dynamics and ₆₁₁ physiologically plausible mean values of intra and extra cellular signals in the ₆₁₂ spontaneous condition. In our experience, achieving physiological behavior in the ₆₁₃ spontaneous state secured physiological behavior also in the evoked conditions. While ₆₁₄ the model dynamics were robust to many of the parameters, we noticed that the model ₆₁₅ was particularly sensitive to parameters that governed the balance between excitation ₆₁₆ and inhibition, such as the synaptic strengths between different neural types. However, ₆₁₇ the broad range of stimulation protocols and functional features that the models is ₆₁₈ tested against ensures that this fitting process avoids major over-fitting that could skew ₆₁₉ the interpretation of the results. ₆₂₀

Having instantiated the model based on the parameterization determined in the ₆₂₁ process described above, we next validated the model against a broad set of ₆₂₂ experimental findings. Acknowledging the finite manpower resources, and the fact that ₆₂₃ the integrative program is inherently an incremental process, we identified a priority list ₆₂₄ of V1 phenomena to be addressed in this first snapshot. These validation tests were ₆₂₅ strategically selected to cover the most common stimulation paradigms (the stimuli ₆₂₆ represent a superset of visual stimulation protocols used in our own lab [21] and in the ₆₂₇ Allen Institute's Brain Observatory [95]) and a broad range of the most salient and well ₆₂₈ established functional features of early visual system. ₆₂₉

5.6   Model limitations ₆₃₀

The model makes five main simplifications: (i) the omission of layers 5 and 6 [96, 97], (ii) ₆₃₁ omission of the cortico-thalamic feedback pathway [98, 99], (iii) omission of the feedback ₆₃₂ from higher cortical areas [100, 101], (iv) reduction of neural dynamics to only two types ₆₃₃ (excitatory and inhibitory) [65, 102], and (v) the reduced cortical neuron density. The ₆₃₄ infra-granular layers have been omitted because they are much less understood and ₆₃₅ there is much less data available for constraining their model. Furthermore, they are ₆₃₆ believed to be largely involved in sub-cortical projections and feedback projection from ₆₃₇

higher level cortical areas [103], which are another two features not considered here (but see [104]). The cortico-cortical feedback, mediated partly by long-distance supra-granular connections, represents a particularly difficult challenge for modeling, as it in principle requires some form of treatment of all the numerous higher-cortical areas that send feedback to V1, while relying on a relatively sparse set of findings on the anatomy and physiology of these feedback structures. Due to recent methodological advances in selective targeting of different neural subtypes, the roles which the different inhibitory neural sub-types play in cortical processing are starting to be uncovered [105, 106]. This mapping out of the neural sub-type characteristics is, however, still in an early stage, and is largely confined to the mouse, without a clear understanding of how these findings will translate to cortical processing in higher mammals. Undoubtedly, continuing systematic exploration of the neural sub-type specificities, especially if successfully translated into higher mammalian models, would provide an invaluable trove of constraints to be incorporated into the model.

We model neurons as exponential integrate-and-fire units with conductance-based synapses [107]. This choice had several motivations. As we have demonstrated here, a number of important characteristics of V1 computations are manifested at the precision level of single spikes, and such quantities do not have a direct representation in firing-rate based models. We also show here how the dynamics of sub-threshold signals, including the excitatory and inhibitory synaptic conductances, provide an important means for constraining the model, which thus excludes usage of current-based synapse models. The particular choice of the exponential variant of the conductance-based integrate-and-fire scheme was motivated by our observations that the variable effective threshold in this neural model secures more stable asynchronous behavior than the simpler fixed-threshold variants. We chose not to pursue more detailed representations of single neuron dynamics, such as a full morphological representation or ion channel kinetics, for two main reasons: (i) the neuron counts necessitated by the cortical area considered in this study, and the length of the stimulation paradigms targeted in this study would make simulations with such detailed neural models nearly intractable on currently available computing resources, and (ii) lack of sufficient detailed reconstructions of neurons in cat V1 with associated intracellular functional data [108] and lack of detailed characterization of ion channel kinetics and ion channel dendritic distributions in cat.

Another limitation of our study is that we do not model the full spectrum of the RF properties present in V1. For example, we assume fixed size, spatial frequency and aspect ratio of the templates from which afferent connectivity (and thus 'afferent' RF) of Layer 4 V1 neurons is generated. We also assume that both the ON and OFF channels are of equal strength, even though recent electrophysiological investigations revealed the dominance of the OFF pathway and systematic variations of the ON-OFF pattern of RF following the orientation maps [109, 110]. All these structural parameters are known to have substantial biological variability [47, 111, 112]. They are the result of developmental processes, whether intrinsic or stimulus driven, which both have stochastic components, but more importantly likely generate features of cortical organization more broadly adapted to processing of visual stimuli that are not accounted for in our model (and presumably some or even most have not even been discovered yet). Unfortunately, quantitative estimates of these various sources of variability are not currently available. Our modeling paradigm could be used in the future to formulate predictions for these values by evaluating which additional sources of variability and of what magnitude could account for the experimental observations. An alternative intriguing possibility would be adoption of a hybrid scheme, whereby some of the variability due to the developmental processes would be directly simulated in dedicated adaptive models (e.g. see [10, 113]), and then transplanted into the detailed

spiking modeling paradigm for more thorough examination.

Ultimately, the main reason for omission of these important structural features was to restrict this initial study to a manageable (both computationally and conceptually) level of complexity. We would like to emphasize that even though we argue here for an integrative, comprehensive treatment of the neural systems under study, we acknowledge that this has to be an incremental process. All these omissions would be natural candidates for the next steps of the integrative research program initiated in this study.

6 Materials and Methods

The model's overall architecture was inspired by a previous rate-based model of V1 development [10] and a spiking model of a single cortical column [63]. The model is implemented, and all experiments and analysis defined in the Mozaik framework [17], which is freely accessible at *https://github.com/CSNG-MFF/mozaik*. The NEST simulator [114, version 3.4] was used as the underlying simulation engine.

6.1 V1 model

The cortical model corresponds to layers 4 and 2/3 of a 5.0×5.0 mm patch of cat primary visual cortex, and thus given the magnification factor of 1 at 3 degrees of visual field eccentricity [115], covers roughly 5.0×5.0 degrees of visual field. It contains 108150 neurons and ~ 155 million synapses. This represents a significant down-sampling ($\sim 10\%$) of the actual density of neurons present in the corresponding portion of cat cortex [116] and was chosen to make the simulations computationally feasible. The neurons are distributed in equal quantities between the simulated Layer 4 and Layer 2/3, which is consistent with anatomical findings showing that in cat primary visual cortex approximately the same number of neurons have cell bodies in these two cortical layers [116]. Each simulated cortical layer contains one population of excitatory neurons and one population of inhibitory neurons in the ratio 4:1 [117, 118].

We model both the feed-forward and recurrent V1 pathways; however, overall the model architecture is dominated by the intra-cortical connectivity, while thalamocortical synapses constitute about 14% of the synaptic input to Layer 4 cells (see Section 6.1.3), in line with experimental evidence [29]. The thalamic input reaches both excitatory and inhibitory neurons in Layer 4 (see Fig 1EF). In both cortical layers we implement short-range lateral connectivity between both excitatory and inhibitory neurons, and additionally in Layer 2/3 we also model long range excitatory connections onto other excitatory and inhibitory neurons [19, 20, 119] (see Fig 1AB). Layer 4 excitatory neurons send narrow projections to Layer 2/3 neurons (see Fig 1E).

6.1.1 Neuron model

All neurons are modeled as single-compartment exponential integrate-and-fire neurons (ExpIF; Eq. 1). The time course of the membrane potential $V(t)$ of all modeled neurons is governed by:

$$\tau_{\mathrm{m}}\frac{\mathrm{d}V}{\mathrm{d}t} = -(V - E_{\mathrm{L}}) + \Delta_T \exp\left(\frac{V - V_T}{\Delta_T}\right) + R_{\mathrm{m}}g_{\mathrm{exc}}(E_{\mathrm{exc}} - V) + R_{\mathrm{m}}g_{\mathrm{inh}}(E_{\mathrm{inh}} - V)$$

(1)

where g_{exc} and g_{inh} are the incoming excitatory and inhibitory synaptic conductances (see Section 6.1.6), and V_T is the threshold parameter of the exponential action potential generation mechanism. Spikes are registered when the membrane potential

crosses -40 mV, at which time the membrane potential is set to the reset value V_r of ₇₃₁ -60 mV. Each spike is followed by a refractory period during which the membrane ₇₃₂ potential is held at V_r. For excitatory neurons, E_L and V_T were respectively set to ₇₃₃ -80 mV and -57 mV; for inhibitory neurons, they were set to -78 mV and -58 mV. The ₇₃₄ membrane resistance in cat V1 in the absence of synaptic activity has been estimated to ₇₃₅ be on average $\sim 250\,\text{M}\Omega$ for excitatory neurons, and that of inhibitory neurons has been ₇₃₆ estimated to be higher [44]. To reflect these findings we set the membrane resistance of ₇₃₇ all cortical excitatory and inhibitory neurons R_m to $250\,\text{M}\Omega$ and $300\,\text{M}\Omega$ respectively. ₇₃₈ We set the membrane time constant τ_m of excitatory neurons to 8 ms, and of inhibitory ₇₃₉ neurons to 9 ms. The refractory period is 2 ms and 0.5 ms for excitatory and inhibitory ₇₄₀ neurons respectively. Overall these neural parameter differences between excitatory and ₇₄₁ inhibitory neurons reflect the experimentally observed greater excitability and higher ₇₄₂ maximum sustained firing rates of inhibitory neurons [120]. The excitatory and ₇₄₃ inhibitory reversal potentials E_{exc} and E_{inh} are set to 0 mV and -80 mV respectively in ₇₄₄ accordance with values observed experimentally in cat V1 [44]. The threshold slope ₇₄₅ factor Δ_T was set to 0.8 mV for all neurons based on fits to pyramidal neuron data [121]. ₇₄₆

6.1.2 Thalamo-cortical model pathway ₇₄₇

All neurons in the model Layer 4 receive connections from the model LGN (see Section ₇₄₈ 6.2). For each neuron, the spatial pattern of thalamo-cortical connectivity is determined ₇₄₉ by the sum of a Gabor distribution (Eq. 2), inducing the elementary RF properties in ₇₅₀ Layer 4 neurons [5, see Fig 1EF], and a Gaussian distribution. ₇₅₁

$$
\begin{aligned}
g(x, y, \lambda, \theta, \psi, \sigma, \gamma) &= \exp\left(\frac{x'^2 + y'^2\gamma^2}{2\sigma^2}\right)(G + \cos(2\pi x'\lambda + \psi)) \\
x' &= x\cos\theta + y\sin\theta \\
y' &= -x\sin\theta + y\cos\theta
\end{aligned}
\tag{2}
$$

where G is set at 0.085 and defines the relative weight of the Gaussian distribution with ₇₅₂ respect to the Gabor distribution. For individual neurons the orientation θ, phase ψ, ₇₅₃ size σ, frequency λ and aspect ratio γ of the distributions are selected as follows. To ₇₅₄ induce functional organization in the model, a pre-computed orientation map [122] ₇₅₅ corresponding to the 5.0×5.0 mm of simulated cortical area is overlaid onto the ₇₅₆ modeled cortical surface (see Fig 2A), thereby assigning each neuron an orientation ₇₅₇ preference θ (see Fig 2B). For the sake of simplicity, the phase ψ of the Gabor ₇₅₈ distribution is assigned randomly (but see Sec 5.6). The remaining parameters are set ₇₅₉ to constant values, matching the average of measurements in cat V1 RFs located in the ₇₆₀ para-foveal area. Taking into consideration the estimated mean RF diameter at ₇₆₁ para-foveal eccentricities [59], and assuming that the measured RF perimeter ₇₆₂ corresponds to the 3σ envelope of the Gabor function, we set the size σ parameter to ₇₆₃ 0.17 degrees of visual field. The spatial frequency λ was set to 0.8 Hz based on ₇₆₄ experimental measurements [123]. Finally, studies have produced wide range of ₇₆₅ estimates for the mean of the RF aspect ratio, ranging from 1.7 to 5 [124]. We have ₇₆₆ chosen to set the value of the γ parameter to 2.5, which is at the lower end of this range, ₇₆₇ as studies restricted to only layer 4 cells produced such lower estimates. ₇₆₈

6.1.3 Cortico-cortical connectivity ₇₆₉

The number of synaptic inputs per neuron in cat V1 has been estimated to be ₇₇₀ approximately 5800 [125]. While a substantial portion of these synapses come from ₇₇₁ outside V1, the exact number has not yet been established. A recent investigation ₇₇₂

by [93] found that even for a cortical section 800µm in diameter 76% of synapses come ₇₇₃ from outside of this region, while a rapid falloff of bouton density with radial distance ₇₇₄ from the soma has been demonstrated in cat V1 [20]. Furthermore, V1 receives direct ₇₇₅ input from a number of higher cortical areas and sub-cortical structures. Feedback from ₇₇₆ V2 alone accounts for as many as 6% of synapses in supra-granular layers of ₇₇₇ macaque [101, 126]. Altogether it is reasonable to extrapolate from these numbers that ₇₇₈ as many as 50% of the synapses in layers 4 and 2/3 originate outside of V1. ₇₇₉

Furthermore, cortical synapses have been shown to frequently fail to transmit ₇₈₀ arriving action potentials, with typically every other pre-synaptic action potential ₇₈₁ failing to evoke a post-synaptic potential [127, 128]. Because it would be ₇₈₂ computationally expensive to explicitly model the synaptic failures, to account for the ₇₈₃ resulting loss of synaptic drive, we have factored it into the number of simulated ₇₈₄ synapses per neuron. Thus, taking into account the average number of synaptic inputs ₇₈₅ (5800), and the estimates of the proportion of extra-areal input (50%) and of the failure ₇₈₆ rates of synaptic transmission (50%) we decided to model 1000 reliable synaptic inputs ₇₈₇ per modeled excitatory cell in Layer 4. In line with the observations from Binzegger *et* ₇₈₈ *al.* [18] (reported by Izhikevich and Edelman [129]) that Layer 2/3 has more numerous ₇₈₉ recurrent connections than Layer 4, each modeled excitatory cell in Layer 2/3 receives ₇₉₀ 2300 synaptic inputs. Inhibitory neurons receive 40% fewer synapses than their ₇₉₁ excitatory counterparts, to account for their smaller size, but otherwise synapses are ₇₉₂ formed proportionally to the two cell type densities. 22% of the Layer 2/3 neurons ₇₉₃ synapses originate from Layer 4 cells [129]. In addition Layer 4 cells from each excitatory ₇₉₄ and inhibitory population receive additional thalamo-cortical synapses from a uniform ₇₉₅ distribution with respective boundaries of [90, 190], and [112, 168], which lies within the ₇₉₆ range of experimental observations [29]. The synapses are drawn probabilistically with ₇₉₇ replacement (with functional and geometrical biases described below). However, ₇₉₈ because we allow formation of multiple synapses between neurons, the exact number of ₇₉₉ connected neurons and the effective unitary strength of these connections is variable. ₈₀₀

The geometry of the cortico-cortical connectivity is determined based on two main ₈₀₁ principles: (i) the connection probability falls off with increasing cortical distance ₈₀₂ between neurons [19, 20, 130] (see Fig 1AB), and (ii) connections have a functionally ₈₀₃ specific bias [20, 88]. The two principles are each expressed as a connection-probability ₈₀₄ density function, then multiplied and re-normalized to obtain the final connection ₈₀₅ probability profiles, from which the actual cortico-cortical synapses are drawn. The ₈₀₆ following two sections describe how the two probability density profiles of connectivity ₈₀₇ are obtained. ₈₀₈

Finally, apart from the connectivity directly derived from experimental data, we also ₈₀₉ modeled a direct feedback pathway from Layer 2/3 to Layer 4. Such direct connections ₈₁₀ from Layer 2/3 to Layer 4 are rare [18], however a strong feedback from Layer 2/3 ₈₁₁ reaching Layer 4 via layers 5 and 6 exists [18]. Since we found that closing of this strong ₈₁₂ cortico-cortical loop is important for correct expression of functional properties across ₈₁₃ the investigated layers, and because we do not explicitly model the sub-granular layers ₈₁₄ (see Section 6.1), we decided to instead model a direct Layer 2/3 to Layer 4 pathway. ₈₁₅ For the sake of simplicity, we have assumed it has the same geometry as the ₈₁₆ feed-forward Layer 4 to Layer 2/3 pathway (see following two sections), and considered ₈₁₇ it represents 20% of the synapses formed in Layer 4. ₈₁₈

6.1.4 Spatial extent of local intra-cortical connectivity ₈₁₉

The exact parameters of the spatial extent of the model local connectivity, with the ₈₂₀ exception of excitatory lateral connections in Layer 2/3, were established based on a ₈₂₁ re-analysis of data from cat [19]. Let M_{xyz}^{ij} be the probability of potential connectivity ₈₂₂ (Fig 7 in [19]), of a pre-synaptic neuron of type $i \in \{\text{exc}, \text{inh}\}$ at cortical depth x to ₈₂₃

	α				θ			
	L4 Exc	L4 Inh	L23 Exc	L23 Inh	L4 Exc	L4 Inh	L23 Exc	L23 Inh
L4 Exc	0.0139	0.0148	0.0174	0.0197	207.7	191.8	154.4	131.5
L4 Inh	0.0126	0.0119			237.5	256.4		
L23 Inh			0.0149	0.0150			189.5	188.61

Table 2. The parameters of hyperbolic profiles (Eq. 3) of potential connectivity derived from [19] used to generate the distance dependent profile of local connectivity in the model. The parameters for pathways not present in the model are not included in the table.

other post-synaptic neurons of type j at cortical depth y and lateral (radial) displacement z. We reduced the 3 dimensional matrix M^{ij}_{xyz} to a single spatial profile for each pair of layers and neural types (excitatory and inhibitory). This was done as follows. For each possible projection $L_{\mathrm{pre}}T_{\mathrm{pre}} \to L_{\mathrm{post}}T_{\mathrm{post}}$ where L corresponds to layers $L \in \{4, 2/3\}$ and T corresponds to neuron type $T \in \{\mathrm{exc}, \mathrm{inh}\}$ we did the following:

1. Select section of $M^{T_{\mathrm{pre}}T_{\mathrm{post}}}_{xyz}$ along the depth dimension corresponding to the pre-synaptic layer L_{pre} and post-synaptic layer L_{post}.

2. Average the selected section of M along the the dimensions corresponding to pre- and post-synaptic layer depth, resulting in a vector representing the average lateral profile of potential connectivity of a neuron residing in layer L_{pre} to neurons residing in layer L_{post}.

3. Normalize the resulting distance profile to obtain the probability density function.

To obtain a parametric representation of the distance connectivity profiles we fitted them with several probability distributions, including Gaussian, exponential and hyperbolic. We found that the best fits were obtained using the zero mean hyperbolic distribution:

$$pdf(x) = \exp\left(-\alpha\sqrt{\theta^2 + x^2}\right) \tag{3}$$

which was thus chosen as the parametric representation of the connectivity profiles. The resulting values of the parameters of the fitted hyperbolic distributions, for all combinations of pre- and post- synaptic layers and neuron types, which were used to generate the local connectivity distance dependent profiles in the model, can be found in Table 2.

This data [19] only reflects the local connectivity, due to its dependence on neural reconstruction in slices which cut off distal dendrites and axons further than 500μm from the cell body. In cat Layer 2/3 excitatory neurons send long-range axons up to several millimeters away [20, 119]. Following Buzás *et al.* [20] we model the lateral distribution of the Layer 2/3 excitatory connections as $G(\sigma_s) + \alpha G(\sigma_l)$, where G is a zero mean normal distribution, $\sigma_s = 270\,\mu$m and $\sigma_l = 1000\,\mu$m are the short and long-range space constants chosen in-line with Buzás *et al.* [20], and $\alpha = 4$ is the ratio between the short-range and long-range components. Furthermore, the long-range connections in Layer 2/3 have like-to-like bias [20]. How we incorporate this functional bias is described in the following section.

6.1.5 Functionally specific connectivity

The functionally specific connectivity in V1 poorly understood. Experimental studies have shown that local connections in cat are diffuse, without a strong orientation

bias [20], while long-range connections have a moderate bias towards iso-orientation ⁸⁵⁹ configurations [20, 92]. Recent more detailed studies of local connectivity in mice have ⁸⁶⁰ shown that local connections also have a weak bias towards connecting neurons with ⁸⁶¹ similar receptive field properties [88, 131]. Furthermore, the anti-phase relationship ⁸⁶² between excitatory and inhibitory synaptic conductances found in a majority of cat V1 ⁸⁶³ Simple cells [21, 94] can be interpreted as indirect evidence for inhibitory inputs to be ⁸⁶⁴ biased towards neurons with anti-correlated RFs (i.e. co-tuned but anti-phase; although ⁸⁶⁵ inhibitory conductance shows greater diversity of orientation tuning relative to ⁸⁶⁶ spike-based preference [39, 44, 90]). Overall, these anatomical and functional findings ⁸⁶⁷ point to a weak tendency of excitatory neurons towards connecting nearby neurons of ⁸⁶⁸ similar RF properties while this bias increases somewhat for more distant post-synaptic ⁸⁶⁹ neurons, and within Layer 4 Simple cells for inhibitory neurons to exhibit weak bias ⁸⁷⁰ towards connecting to neurons with anti-correlated RFs. ⁸⁷¹

Within Layer 4, the above findings point to the push-pull scheme of connectivity ⁸⁷² that has been demonstrated to explain a range of phenomena related to orientation ⁸⁷³ tuning [5] (see Fig 1E). To implement this connectivity in our model we performed the ⁸⁷⁴ following process. For each pair of Layer 4 neurons the correlation c between their ⁸⁷⁵ afferent RFs (i.e. RF defined only by the afferent inputs from thalamus) is calculated. ⁸⁷⁶ The connectivity likelihood for a given pair of neurons is then given by ⁸⁷⁷ $\frac{1}{\sigma\sqrt{2\pi}}e^{-(c-\mu)^2/2\sigma^2}$ where $\sigma = 1.3$, and $\mu = 1$ if the post-synaptic neuron is excitatory or ⁸⁷⁸ $\mu = -1$ if it is inhibitory. ⁸⁷⁹

In Layer 2/3 excitatory neurons send long-range connections spanning up to $6\,\text{mm}$ to ⁸⁸⁰ both excitatory and inhibitory neurons, preferentially targeting those with similar ⁸⁸¹ orientation preference [20]. To reflect this connectivity in the model we define the ⁸⁸² functional connectivity likelihood between pairs of neurons in Layer 2/3 as ⁸⁸³ $\frac{1}{\sigma\sqrt{2\pi}}e^{-(\Delta o)^2/2\sigma^2}$ where the Δo is the difference between the orientation preference of ⁸⁸⁴ the two neurons, and σ is set to 1.3. The connectivity likelihoods described above are ⁸⁸⁵ renormalized for each neuron to obtain probability density functions. Note that only the ⁸⁸⁶ long-range component of the Layer 2/3 model is multiplied by this functionally specific ⁸⁸⁷ bias, while the short-range component remains non-specific (see Section 6.1.4). Overall ⁸⁸⁸ this parameterization leads to only weak bias towards co-tuned connections in both ⁸⁸⁹ simulated cortical layers. ⁸⁹⁰

6.1.6 Synapses ⁸⁹¹

The relatively few studies that have examined in detail the strength of cortical synapses ⁸⁹² have found that unitary synaptic strengths are generally weak ($<2\,\text{nS}$), but broad ⁸⁹³ ranges have been reported [132, 133]. While the overall specificity of the synaptic ⁸⁹⁴ strength between pairs of neural types or layers remains unclear, a recent study in cat ⁸⁹⁵ V1 has shown that synapses from excitatory onto inhibitory cells are stronger than ⁸⁹⁶ those targeting excitatory neurons [132, 133]. Reflecting these insufficient experimental ⁸⁹⁷ constraints, the synaptic weights were selected to achieve an overall balance between ⁸⁹⁸ excitation and inhibition that supports reasonable levels of both spontaneous and ⁸⁹⁹ evoked activity, while being compatible with the limited physiological findings. ⁹⁰⁰ Specifically, we set the Layer 4 intra-laminar excitatory-to-excitatory synapses to ⁹⁰¹ $0.18\,\text{nS}$ and excitatory-to-inhibitory synapses to $0.22\,\text{nS}$, while inhibitory synapses are ⁹⁰² set to $1\,\text{nS}$. In Layer 2/3, the synaptic weights were set to the same value as Layer 4 ⁹⁰³ with the exception of the excitatory-to-inhibitory synaptic weights which were set to ⁹⁰⁴ $0.35\,\text{nS}$. The thalamo-cortical synapses were set to $1.2\,\text{nS}$, reflecting the findings that ⁹⁰⁵ these synapses tend to be larger [29] and more reliable [127] than their intra-laminar ⁹⁰⁶ counterparts. The weights of synapses formed by connections from Layer 4 to Layer 2/3 ⁹⁰⁷ were set to 1, while the weights of synapses formed by the feedback connection from ⁹⁰⁸

Layer 4 to Layer 2/3 were set to the same value as for the excitatory intra-laminar
synapses. Unitary synaptic inputs are modeled as transient conductance changes, with
exponential decay with time-constant $\tau_e = 1.5$ ms for excitatory synapses and
$\tau_i = 4.2$ ms for inhibitory synapses.

We have also modeled synaptic depression for thalamo-cortical, and excitatory
cortico-cortical synapses [134] using the model of [135]. For the thalamo-cortical
synapses we assume parameters corresponding to strong depression similar to [136]
and [63] (U=0.75, $\tau_{rec} = 125$ ms, $\tau_{psc} = 1.5$ ms and $\tau_{fac} = 0$ ms). For the cortico-cortical
excitatory synapses we assumed moderate depression (U=0.75, $\tau_{rec} = 30$ ms,
$\tau_{psc} = 1.5$ ms and $\tau_{fac} = 0$ ms), in line with [135], except for the feeedback synapses for
which we assumed weaker depression (U=0.75, $\tau_{rec} = 20$ ms, $\tau_{psc} = 1.5$ ms and
$\tau_{fac} = 0$ ms). The inhibitory neurons in our model correspond mostly to fast-spiking
basket cells, as they are the main source of inhibition in V1. Basket cells .2'3 form
depressing synapses with both excitatory [137] and inhibitory [138] neurons in the
neocortex of rodents, although this has not been yet quantified in cats to the best of our
knowledge. For Layer 4 inhibitory synapses, we decided to use parameters
corresponding to a stronger depression than for the cortico-cortical excitatory synapses
(U=0.75, $\tau_{rec} = 70$ ms, $\tau_{psc} = 4.2$ ms and $\tau_{fac} = 0$ ms), while for the inhibitory synapses
of Layer 2/3 we used the same recovery time constant as for the excitatory synapses
(U=0.75, $\tau_{rec} = 30$ ms, $\tau_{psc} = 4.2$ ms and $\tau_{fac} = 0$ ms).

6.1.7 Delays

We model two types of delays in the model circuitry. First, for all intra-cortical
connectivity a distance-dependent delay with propagation constant of
0.3 ms^{-1} [139–141] was used, which corresponds to the slow propagation of action
potentials along the intra-V1 (lateral) un-myelinated axons. The delays in the
feed-forward thalamo-cortical pathway are drawn from a uniform distribution within the
range $(1.4, 2.4)$ ms. Second, [142] have recently shown that delays of synaptic
transmission in cat V1 are dependent on both pre- and post-synaptic neural type, with
the notable feature of slow excitatory to excitatory and fast excitatory to inhibitory
transmission. Distance-dependent axonal propagation delay is unlikely to explain these
results as these experiments were performed in nearby neurons [142]. Thus, as suggested
by [142], we have included a constant additive factor in all synaptic delays, specifically
1.4 ms for excitatory to excitatory synapses, 0.5 ms for excitatory to inhibitory synapses,
1.0 ms for inhibitory to excitatory synapses and 1.4 ms for inhibitory to inhibitory
synapses, in line with the quantitative observations by [142]. We observed that the
addition of this neuron-type-dependent delay factor improved the stability of the
modeled cortical neural networks, reducing synchronous events during spontaneous
activity. We hypothesized that this is due to the ability of inhibition to respond faster
to any transient increase in activity in the network due to the shorter excitatory to
inhibitory delay.

6.2 Input model

The input model described in this section corresponds to the whole retino-thalamic
pathway. The cortical model corresponds to roughly $5.0 \times 5.0°$ of visual field (Fig 1CF).
To accommodate the full extents of the RFs of neurons at the edges of the model, the
LGN model corresponds to $6 \times 6°$ of visual field. In the same manner, to accommodate
the full extent of RFs of thalamic neurons the overall visual field from which the
thalamic model receives input corresponds to $11 \times 11°$.

We simulate the responses of the LGN neurons using the canonical center-surround
model of RF (Fig 1C). The centers of both ON and OFF LGN neurons RFs are

uniformly randomly distributed in the visual space, with density 100 neurons per square ₉₅₈ degree. Each LGN neuron has a spatiotemporal RF, with a difference-of-Gaussians ₉₅₉ spatial profile and a bi-phasic temporal profile defined by a difference of gamma ₉₆₀ functions [143]. Due to the relatively small region of visual space our model covers, we ₉₆₁ do not model the systematic changes in RF parameters with foveal eccentricity and thus ₉₆₂ assume that all ON and OFF LGN neurons have identical parameters. The exact ₉₆₃ spatial and temporal parameters have been fitted to match the observations in [143]. ₉₆₄

To obtain the spiking output of a given LGN neuron, the visual stimulus, sampled ₉₆₅ into 7 ms frames, is convolved with its spatiotemporal RF. In addition, saturation of the ₉₆₆ LGN responses with respect to local contrast and luminance is modeled [3, 22]. For ₉₆₇ simplicity, the local luminance ll is calculated as the mean of luminance values and local ₉₆₈ contrast lc as the standard deviation of the luminances within the RF of a given neuron. ₉₆₉ The response of the linear RF is separated into a DC (luminance) component r_l and a ₉₇₀ contrast component r_c. The saturation of the two components is modeled with two ₉₇₁ Naka-Rushton functions $\alpha_l \frac{r_l}{1.0+\beta_l ll}$ and $\alpha_c \frac{r_c}{1.0+\beta_c lc}$ respectively, where α is the gain and ₉₇₂ β is the saturation parameter of the corresponding component. The parameters α and β ₉₇₃ were empirically adjusted to obtain luminance and contrast response curves whose ₉₇₄ saturation point and level are within the ranges observed experimentally [3, 22]. ₉₇₅

The resulting luminance and contrast temporal traces are then summed and injected ₉₇₆ into integrate-and-fire neurons as a current, inducing stimulus dependent spiking ₉₇₇ responses. In addition to the stimulus-dependent drive, each LGN neuron is also ₉₇₈ injected with an independent white noise current. The magnitude and variance of this ₉₇₉ noise is such that ON and OFF LGN neurons fire at $\sim$ 17 and $\sim$ 8 spikes/s respectively ₉₈₀ in the spontaneous condition [22]. ₉₈₁

6.3 Stimulation protocols ₉₈₂

Each stimulation protocol consists of a series of visual stimuli which are interleaved with ₉₈₃ 150 ms of 50 cd/m^2 gray blank stimuli. The population of recorded neurons, with the ₉₈₄ exception of the size tuning protocol (see Section 6.3.2), was always restricted to the ₉₈₅ central circular patch (radius of 1 mm) of the model to avoid contamination by ₉₈₆ potential edge-effects due to the finite simulated cortical area. Because the recording of ₉₈₇ intracellular signals at high temporal resolution for such a long stimulus set as presented ₉₈₈ in this work is extremely memory consuming, we recorded the membrane potential and ₉₈₉ excitatory and inhibitory conductances only from a subset of 190 neurons confined to ₉₉₀ the central 0.2×0.2 mm patch of the simulated cortical space, and whose orientation ₉₉₁ preference was within 0.2 radians of horizontal. The durations of all visual stimuli are ₉₉₂ aligned with the 7 ms frame duration at which the retino-thalamic model operates (see ₉₉₃ Section 6.2). Within each experimental protocol, stimuli were presented in a random ₉₉₄ order. ₉₉₅

6.3.1 Drifting sinusoidal grating protocol ₉₉₆

The spatial and temporal frequency of the RFs of the modeled LGN neurons (see ₉₉₇ Section 6.1.2), and of the Gabor distribution template from which thalamo-cortical ₉₉₈ synapses were sampled, were identical for all simulated LGN and cortical neurons ₉₉₉ respectively. An important consequence of this simplification is that it allowed us to ₁₀₀₀ efficiently execute protocols requiring DGs. By employing a full-field stimulus with ₁₀₀₁ spatial frequency matching the Gabor template (0.8 Hz) and drifting at 2 Hz, we were in ₁₀₀₂ parallel stimulating all cortical neurons with a stimulus with optimal spatial frequency. ₁₀₀₃ We varied the orientation of the gratings between 0 and 180 degrees, in 8 equal steps. ₁₀₀₄ We used 3 values of contrast: 10%, 30% and 100%. Each grating was shown 10 times for ₁₀₀₅ 2002 ms. We recorded spikes from 2025 randomly selected neurons for this experiment. ₁₀₀₆

6.3.2 Size tuning protocol

Size tuning was measured using DGs of optimal spatial and temporal frequency (see
orientation tuning protocol for details), confined to an aperture of variable diameter.
The orientation of the gratings was horizontal (0°) and the position was in the center of
the simulated visual field. The diameter of the aperture was varying between 0° and
10° in 12 steps, with contrast of either 10% or 100%, each presentation lasting 2002 ms
and being repeated 10 times.

The center of the modeled cortical area was occupied by a horizontal orientation
domain (see Fig 2). During the size tuning protocol we recorded from neurons in this
central (horizontal) orientation domain and selected neurons whose RFs centers were
within 0.4° from the center of the simulated visual field and whose orientation
preference was within 0.25 radians from horizontal. This setup allowed us to
simultaneously determine the size tuning of the population of neurons reported here,
with precision comparable to experimental studies, while greatly reducing the
computational resources required for the simulation.

6.3.3 Natural images with simulated eye-movement protocol

We have replicated the natural image with simulated eye-movement (NI) protocol
introduced by Baudot *et al.* [21]. This protocol emulates the global retinal flow
experienced during the exploration of the natural environment by simulating the retinal
impact of visuomotor interaction. This is done by imposing shifts and drifts of a static
natural scene that reproduce the kinematics of a realistic ocular scanpath [21] recorded
in the intact behaving animal. We used the same image as in Baudot *et al.* [21], scaled
to match the size of the simulated visual space, and the same path. Presentation of the
resulting movie lasting 2002 ms was repeated 10 times. We recorded spikes from 2025
randomly selected neurons for this protocol.

6.4 Data analysis

Unless specified otherwise, all tuning curves were calculated as the trial averaged mean
firing rate during stimulation with the given stimulus and parameter value (e.g.
orientation of sinusoidal grating). Spontaneous activity level was not subtracted. All
analog signals were recorded at 1 ms resolution.

We calculated the *irregularity* of an individual neuron's spike train as the coefficient
of variation (CV) of its inter-spike-intervals (ISI):

$$CV_{ISI}^2 = \frac{\text{Var}[ISI]}{\langle ISI \rangle^2} \tag{4}$$

where Var corresponds to variance, and $\langle \rangle$ is the mean. To ensure accurate estimation of
the irregularity we exclude neurons that fired fewer than 10 spikes during the recording.
We consider the threshold for irregular firing to be $CV_{ISI} > 0.9$ [45]. We assess
synchrony in our network by calculating the average cross-correlation (CC) of spike
counts SC_i SC_j between disjoint pairs of all recorded neurons i, j.

$$CC[SC_i, SC_j] = \frac{\text{Cov}(SC_i, SC_j)}{\sqrt{\text{Var}(SC_i)\text{Var}(SC_j)}} \tag{5}$$

where Var is again variance and Cov is covariance of the spike count. The spike counts
were calculated by counting the number of spikes within 10 ms bins.
In order to assess orientation tuning, we calculated two complementary measures:
the half width at half height (HWHH) and relative unselective response amplitude

(RURA) [46]. To calculate HWHH we fitted the orientation tuning curves with a Gaussian function [144]:

$$R(\phi) = \beta + \alpha \exp\left(\frac{\phi - \phi_{\mathrm{pref}}}{2\sigma^2}\right) \tag{6}$$

where R is the spiking response of the given neuron to a sinusoidal grating with orientation ϕ, ϕ_{pref} is the preferred orientation of the given neuron, σ is the width of the tuning, β is the baseline activity and α a scale factor. Low responding neurons (less then 1 spike/s at optimal orientation) were excluded from the analysis, as reliable curve fitting was not possible with the amount of recorded data. Furthermore, neurons for which a reliable fit of the Gaussian curve was not possible ($MSE > 30\%$ of the tuning curve variance) were also excluded from this analysis. In total, a minority consisting of 11% of the neurons was excluded. HWHH was then calculated as $\sqrt{2\ln 2}\sigma$. RURA was calculated from the fitted Gaussian parameters according to the following equation [144]:

$$RURA = \frac{\beta}{\beta + \alpha} \tag{7}$$

The modulation ratio (MR) of spike responses was computed as the ratio of the fundamental Fourier component F1 and the DC component F0 of the neuron's peri-stimulus time histogram (PSTH) in response to a DG [145]. The PSTH was formed with 10 ms bins, and spontaneous activity was subtracted prior to MR calculation. Cells with MR greater than one are classified as Simple cells, cells with MR lower than one are classified as Complex cells. The MR of the membrane potential was calculated analogously as the V1/V0 component ratio of the membrane potential after subtraction of its resting level.

To quantify the parameters of size tuning curves, we first fitted the data with the Supplementary Difference of Gaussians (3G) model [53] defined as:

$$f(r) = K_{\mathrm{c}}G_{\mathrm{c}}(r) - K_{\mathrm{s}}G_{\mathrm{s}}(r) + K_{\mathrm{cs}}G_{\mathrm{cs}}(r) \tag{8}$$

where

$$G_x(d) = \left(\frac{2}{\sqrt{\pi}} \int_0^d e^{-(y/w_x)^2} dy\right)^2 \tag{9}$$

where d is the stimulus diameter, K_{c} is the strength of the center, K_{s} is the strength of the suppressive surround, K_{cs} is the strength of the counter-suppressive surround and w_x are the space constants of the corresponding terms. We then performed the following analysis on the fitted curves. We excluded low responding neurons (response of less then 2 spikes/s at the optimum size) from the analysis as well as neurons for which a reliable fit was not possible ($MSE > 30\%$), leading to 31% of the neurons being discarded. We defined the summation peak δ_{max} as the diameter where maximum response is achieved before suppression sets in (we also refer to this diameter as the maximum facilitation diameter (MFD)), suppression peak δ_{min} as the diameter where the minimum response is achieved across all aperture sizes greater than δ_{max}, and the counter-suppression peak δ_{cs} as the diameter where the maximum response is achieved across all aperture sizes greater than δ_{min}. This in turn allows us to define the peak summation response R_{max}, peak suppression response R_{min}, and peak counter-suppression response R_{cs} as the neuron's responses at the corresponding aperture sizes. We can then define the suppression index (SI) analogously to experimental studies [53] as:

$$SI = \frac{R_{\mathrm{max}} - R_{\mathrm{min}}}{R_{\mathrm{max}}} \tag{10}$$

and the counter-suppression index CSI as:

$$CSI = \frac{R_{\text{cs}} - R_{\min}}{R_{\max}} \tag{11}$$

The reliability and the precision of the responses were measured by fitting a Gaussian function to the mean cross-correlation across trials (Eq. 5) of the spiking responses, and of the sub-threshold membrane potential responses [21]. Spikes were removed from the membrane potential prior to the analysis, by replacing the membrane potential within a 5 ms window centered on each spike with the signal interpolated between the levels of membrane potential before and after the window. The reliability was then defined as the CC peak amplitude at time zero, and the temporal precision by the standard deviation of the Gaussian fit.

Quantitative values in this article are expressed as mean ± SEM.

7   Acknowledgments

This work was supported by the Centre National de la Recherche Scientifique, Paris-Saclay IDEX (NeuroSaclay); the French National Research Agency (Complex-V1); the European Union's Seventh Framework Program under grant agreements 269921 (BrainScaleS) and 604102 (Human Brain Project); through the project Improvement of internationalization in the field of research and development (`CZ.02.2.69/0.0/0.0/17_050/0008466`) at Charles University and through institutional funding from Charles University (projects PRIMUS/20/MED/006 and GAUK/4089/2022).

References

1. Olshausen BA. 20 years of learning about vision: Questions answered, questions unanswered, and questions not yet asked. In: 20 Years of Computational Neuroscience. Springer; 2013. p. 243–270. Available from: `http://link.springer.com/10.1007/978-1-4614-1424-7http://link.springer.com/10.1007/978-1-4614-1424-7{_}12`.

2. Potjans TC, Diesmann M. The cell-type specific cortical microcircuit: Relating structure and activity in a full-scale spiking network model. Cerebral Cortex. 2014;24(3):785–806. doi:10.1093/cercor/bhs358.

3. Bonin V, Mante V, Carandini M. The suppressive field of neurons in lateral geniculate nucleus. Journal of Neuroscience. 2005;25(47):10844–10856. doi:10.1523/JNEUROSCI.3562-05.2005.

4. Carandini M, Heeger DJ. Summation and division by neurons in primate visual cortex. Science. 1994;264(5163):1333–1336. doi:10.1126/science.8191289.

5. Troyer TW, Krukowski aE, Priebe NJ, Miller KD. Contrast-invariant orientation tuning in cat visual cortex: thalamocortical input tuning and correlation-based intracortical connectivity. The Journal of Neuroscience. 1998;18(15):5908–5927.

6. Lauritzen TZ, Miller KD. Different roles for simple-cell and complex-cell inhibition in V1. Journal of Neuroscience. 2003;23(32):10201–10213. doi:23/32/10201 [pii].

7. Teich AF, Qian N. Comparison among some models of orientation selectivity. Journal of Neurophysiology. 2006;96(1):404–419. doi:10.1152/jn.00015.2005.

8. Finn IM, Priebe NJ, Ferster D. The emergence of contrast-invariant orientation tuning in simple cells of cat visual cortex. Neuron. 2007;54(1):137–52. doi:10.1016/j.neuron.2007.02.029.

9. Olshausen BA, Field DJ. Emergence of simple-cell receptive field properties by learning a sparse code for natural images. Nature. 1996;381(6583):607–609. doi:10.1038/381607a0.

10. Antolík J, Bednar JAJA. Development of Maps of Simple and Complex Cells in the Primary Visual Cortex. Frontiers in Computational Neuroscience. 2011;5(April):1–19. doi:10.3389/fncom.2011.00017.

11. Zhu M, Rozell CJ. Visual Nonclassical Receptive Field Effects Emerge from Sparse Coding in a Dynamical System. PLoS Computational Biology. 2013;9(8):1–15. doi:10.1371/journal.pcbi.1003191.

12. Somers DC, Todorov EV, Siapas AG, Toth LJ, Kim DS, Sur M. A local circuit approach to understanding integration of long-range inputs in primary visual cortex. Cerebral Cortex. 1998;8(3):204–217. doi:10.1093/cercor/8.3.204.

13. Rangan A, Tao L, Kovacic G, Cai D. Multiscale modeling of the primary visual cortex. IEEE Engineering in Medicine and Biology Magazine. 2009;28(3):19–24. doi:10.1109/MEMB.2009.932803.

14. Sejnowski TJ, Koch C, Churchland PS. Computational Neuroscience. Science. 1988;241(4871):1299–1306.

15. Einevoll GT, Destexhe A, Diesmann M, Grün S, Jirsa V, de Kamps M, et al.. The Scientific Case for Brain Simulations; 2019. Available from: https://www.sciencedirect.com/science/article/abs/pii/S0896627319302909?via{%}3Dihub.

16. Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, et al. Reconstruction and Simulation of Neocortical Microcircuitry. Cell. 2015;163(2):456–492. doi:10.1016/j.cell.2015.09.029.

17. Antolík J, Davison AP. Integrated workflows for spiking neuronal network simulations. Frontiers in Neuroinformatics. 2013;7(December):1–15. doi:10.3389/fninf.2013.00034.

18. Binzegger T, Douglas RJ, Martin KAC. A quantitative map of the circuit of cat primary visual cortex. Journal of Neuroscience. 2004;24(39):8441–8453. doi:10.1523/JNEUROSCI.1400-04.2004.

19. Stepanyants A, Hirsch JA, Martinez LM, Kisvárday ZF, Ferecskó AS, Chklovskii DB. Local potential connectivity in cat primary visual cortex. Cerebral Cortex. 2008;18(1):13–28. doi:10.1093/cercor/bhm027.

20. Buzás P, Kovács K, Ferecskó AS, Budd JML, Eysel UT, Kisvárday ZF. Model-based analysis of excitatory lateral connections in the visual cortex. Journal of Comparative Neurology. 2006;499(6):861–881. doi:10.1002/cne.21134.

21. Baudot P, Levy M, Marre O, Monier C, Pananceau M, Frégnac Y. Animation of natural scene by virtual eye-movements evokes high precision and low noise in V1 neurons. Frontiers in Neural Circuits. 2013;7(DEC):206. doi:10.3389/fncir.2013.00206.

22. Papaioannou J, White A. Maintained activity of lateral geniculate nucleus neurons as a function of background luminance. Experimental Neurology. 1972;34(3):558–566. doi:10.1016/0014-4886(72)90050-7.

23. Cavaggioni A. The dark-discharge of the eye in the unrestrained cat. Pflügers Archiv European Journal of Physiology. 1968;304(1):75–80. doi:10.1007/BF00586720.

24. Levine MW. The distribution of the intervals between neural impulses in the maintained discharges of retinal ganglion cells. Biological Cybernetics. 1991;65(6):459–467. doi:10.1007/BF00204659.

25. Field GD, Rieke F. Mechanisms regulating variability of the single photon responses of mammalian rod photoreceptors. Neuron. 2002;35(4):733–747. doi:10.1016/S0896-6273(02)00822-X.

26. Croner LJ, Purpura K, Kaplan E. Response variability in retinal ganglion cells of primates. Proceedings of the National Academy of Sciences of the United States of America. 1993;90(17):8128–8130. doi:10.1073/pnas.90.17.8128.

27. Salari V, Scholkmann F, Bokkon I, Shahbazi F, Tuszynski J. The Physical Mechanism for Retinal Discrete Dark Noise: Thermal Activation or Cellular Ultraweak Photon Emission? PLoS ONE. 2016;11(3):1–16. doi:10.1371/journal.pone.0148336.

28. Rogala J, Waleszczyk WJ, Wróbel A, Wójcik DK. Effect of cortex inactivation on spontaneous activity of cells in perigeniculate and dorsal lateral geniculate nuclei. BMC Neuroscience. 2013;14(S1):P418. doi:10.1186/1471-2202-14-s1-p418.

29. Da Costa NM, Martin KAC. How thalamus connects to spiny stellate cells in the cat's visual cortex. Journal of Neuroscience. 2011;31(8):2925–2937. doi:10.1523/JNEUROSCI.5961-10.2011.

30. Esposito F, Mulert C, Goebel R. Combined distributed source and single-trial EEG-fMRI modeling: Application to effortful decision making processes. NeuroImage. 2009;47(1):112–121. doi:10.1016/j.neuroimage.2009.03.074.

31. Lee AK, Manns ID, Sakmann B, Brecht M. Whole-Cell Recordings in Freely Moving Rats. Neuron. 2006;51(4):399–407. doi:10.1016/j.neuron.2006.07.004.

32. Destexhe A, Rudolph-Lilith M. Neuronal noise. Springer Series in Computational Neuroscience. Springer US; 2012. Available from: http://link.springer.com/10.1007/978-0-387-79020-6.

33. Le Van Quyen M, Muller LE, Telenczuk B, Halgren E, Cash S, Hatsopoulos NG, et al. High-frequency oscillations in human and monkey neocortex during the wake–sleep cycle. Proceedings of the National Academy of Sciences. 2016;113(33):9363–9368. doi:10.1073/pnas.1523583113.

34. Van Vreeswijk C, Sompolinsky H. Chaos in neuronal networks with balanced excitatory and inhibitory activity. Science. 1996;274(5293):1724–1726. doi:10.1126/science.274.5293.1724.

35. Brunel N. Dynamics of networks of randomly connected excitatory and inhibitory spiking neurons. Journal of Physiology-Paris. 2000;94(5-6):445–463. doi:10.1016/S0928-4257(00)01084-6.

36. Vogels TP, Abbott LF. Signal propagation and logic gating in networks of integrate-and-fire neurons. Journal of Neuroscience. 2005;25(46):10786–10795. doi:10.1523/JNEUROSCI.3508-05.2005.

37. Bringuier V, Frégnac Y, Baranyi A, Debanne D, Shulz DE. Synaptic origin and stimulus dependency of neuronal oscillatory activity in the primary visual cortex of the cat. Journal of Physiology. 1997;500(3):751–774. doi:10.1113/jphysiol.1997.sp022056.

38. Swadlow HA. Fast-spike Interneurons and Feedforward Inhibition in Awake Sensory Neocortex. Cerebral Cortex. 2003;13(1):25–32. doi:10.1093/cercor/13.1.25.

39. Monier C, Chavane F, Baudot P, Graham LJ, Frégnac Y. Orientation and direction selectivity of synaptic inputs in visual cortical neurons: a diversity of combinations produces spike tuning. Neuron. 2003;37(4):663–80.

40. Bardy C, Huang JY, Wang C, FitzGibbon T, Dreher B. 'Simplification' of responses of complex cells in cat striate cortex: Suppressive surrounds and 'feedback' inactivation. Journal of Physiology. 2006;574(3):731–750. doi:10.1113/jphysiol.2006.110320.

41. Kim T, Freeman RD. Direction selectivity of neurons in the visual cortex is non-linear and lamina-dependent. European Journal of Neuroscience. 2016;43(10):1389–1399. doi:10.1111/ejn.13223.

42. Hromadka T, DeWeese MR, Zador AM. Sparse Representation of Sounds in the Unanesthetized Auditory Cortex. PLoS biology. 2008;6(1):124–137.

43. Buzsáki G, Mizuseki K. The log-dynamic brain: how skewed distributions affect network operations. Nature Reviews Neuroscience. 2014;15(4):264–278. doi:10.1038/nrn3687.

44. Monier C, Fournier J, Frégnac Y. In vitro and in vivo measures of evoked excitatory and inhibitory conductance dynamics in sensory cortices. Journal of Neuroscience Methods. 2008;169(2):323–365. doi:10.1016/j.jneumeth.2007.11.008.

45. Kumar A, Schrader S, Aertsen A, Rotter S. The high-conductance state of cortical networks. Neural Computation. 2008;20(1):1–43. doi:10.1162/neco.2008.20.1.1.

46. Nowak LG, Sanchez-Vives MV, McCormick DA. Lack of orientation and direction selectivity in a subgroup of fast-spiking inhibitory interneurons: Cellular and synaptic mechanisms and comparison with other electrophysiological cell types. Cerebral Cortex. 2008;18(5):1058–1078. doi:10.1093/cercor/bhm137.

47. Cardin JA, Palmer LA, Contreras D. Stimulus feature selectivity in excitatory and inhibitory neurons in primary visual cortex. Journal of Neuroscience. 2007;27(39):10333–10344. doi:10.1523/JNEUROSCI.1692-07.2007.

48. Chariker L, Shapley R, Young LS. Orientation selectivity from very sparse LGN inputs in a comprehensive model of macaque V1 cortex. Journal of Neuroscience. 2016;36(49):12368–12384. doi:10.1523/JNEUROSCI.2603-16.2016.

49. Anderson JS, Carandini M, Ferster D. Orientation tuning of input conductance, excitation, and inhibition in cat primary visual cortex. Journal of Neurophysiology. 2000;84(2):909–926. doi:10.1152/jn.2000.84.2.909.

50. Schummers J, Mariño J, Sur M. Synaptic integration by V1 neurons depends on location within the orientation map. Neuron. 2002;36(5):969–978. doi:10.1016/S0896-6273(02)01012-7.

51. Hirsch JA, Martinez LM. Laminar processing in the visual cortical column. Current Opinion in Neurobiology. 2006;16(4):377–384. doi:10.1016/j.conb.2006.06.014.

52. Priebe NJ, Mechler F, Carandini M, Ferster D. The contribution of spike threshold to the dichotomy of cortical simple and complex cells. Nature Neuroscience. 2004;7(10):1113–1122. doi:10.1038/nn1310.

53. Wang C, Bardy C, Huang JY, FitzGibbon T, Dreher B. Constrast dependence of center and surround integration in primary visual cortex of the cat. Journal of Vision. 2009;9(1):20.1–15. doi:10.1167/9.1.20.

54. Tailby C, Solomon SG, Peirce JW, Metha AB. Two expressions of "surround suppression" in V1 that arise independent of cortical mechanisms of suppression. Visual Neuroscience. 2007;24(1):99–109. doi:10.1017/S0952523807070022.

55. Okamoto M, Naito T, Sadakane O, Osaki H, Sato H. Surround suppression sharpens orientation tuning in the cat primary visual cortex. European Journal of Neuroscience. 2009;29(5):1035–1046. doi:10.1111/j.1460-9568.2009.06645.x.

56. Akasaki T, Sato H, Yoshimura Y, Ozeki H, Shimegi S. Suppressive effects of receptive field surround on neuronal activity in the cat primary visual cortex. Neurosci Res. 2002;43(3):207–20. doi:10.1016/S0168-0102(02)00038-X.

57. Nurminen L, Merlin S, Bijanzadeh M, Federer F, Angelucci A. Top-down feedback controls spatial summation and response amplitude in primate visual cortex. Nature Communications. 2018;9(1). doi:10.1038/s41467-018-04500-5.

58. Jones HE, Andolina IM, Oakely NM, Murphy PC, Sillito AM. Spatial summation in lateral geniculate nucleus and visual cortex. Experimental Brain Research. 2000;135(2):279–284. doi:10.1007/s002210000574.

59. Wilson JR, Sherman SM. Receptive-field characteristics of neurons in cat striate cortex: Changes with visual field eccentricity. Journal of Neurophysiology. 1976;39(3):512–533. doi:10.1152/jn.1976.39.3.512.

60. Sengpiel F, Sen A, Blakemore C. Characteristics of surround inhibition in cat area 17. Experimental Brain Research. 1997;116(2):216–228. doi:10.1007/PL00005751.

61. Anderson JS, Lampl I, Gillespie DC, Ferster D. Membrane potential and conductance changes underlying length tuning of cells in cat primary visual cortex. Journal of Neuroscience. 2001;21(6):2104–2112.

62. Ozeki H, Finn IM, Schaffer ES, Miller KD, Ferster D. Inhibitory Stabilization of the Cortical Network Underlies Visual Surround Suppression. Neuron. 2009;62(4):578–592. doi:10.1016/j.neuron.2009.03.028.

63. Kremkow J, Perrinet LU, Monier C, Alonso JM, Aertsen A, Frégnac Y, et al. Push-pull receptive field organization and synaptic depression: Mechanisms for reliably encoding naturalistic stimuli in V1. Frontiers in Neural Circuits. 2016;10(MAY). doi:10.3389/fncir.2016.00037.

64. Antolík J, Davison APA. Arkheia: Data Management and Communication for Open Computational Neuroscience. Frontiers in Neuroinformatics. 2018;12:6. doi:10.3389/fninf.2018.00006.

65. Nowak LG, Azouz R, Sanchez-Vives MV, Gray CM, McCormick Da. Electrophysiological classes of cat primary visual cortical neurons in vivo as revealed by quantitative analyses. Journal of Neurophysiology. 2003;89(3):1541–1566. doi:10.1152/jn.00580.2002.

66. Hubel DH, Wiesel TN. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. The Journal of Physiology. 1962;160(1):106–154. doi:10.1113/jphysiol.1962.sp006837.

67. Pugh M, Ringach D, Shapley R, Shelley M. Computational Modeling of Orientation Tuning Dynamics in V1 Neurons. J Comp Neuroscience. 2000;8:143–159.

68. Somers DC, Nelson SB, Sur M. An emergent model of orientation selectivity in cat visual cortical simple cells. Journal of Neuroscience. 1995;15(8):5448–5465. doi:10.1523/jneurosci.15-08-05448.1995.

69. Sillito AM, Kemp JA, Milson JA, Berardi N. A re-evaluation of the mechanisms underlying simple cell orientation selectivity. Brain Research. 1980;194:517–520.

70. Creutzfeldt OD, Kuhnt U, Benevento LA. An intracellular analysis of visual cortical neurones to moving stimuli: Responses in a co-operative neuronal network. Experimental Brain Research. 1974;21(3):251–274. doi:10.1007/BF00235746.

71. Rubin DB, VanHooser SD, Miller KD. The stabilized supralinear network: A unifying circuit motif underlying multi-input integration in sensory cortex. Neuron. 2015;85(2):402–417. doi:10.1016/j.neuroN.2014.12.026.

72. Hirsch Ja, Alonso JM, Reid RC, Martinez LM. Synaptic Integration in Striate Cortical Simple Cells. Journal of Neuroscience. 1998;18(22):9517–9528.

73. Wielaard J, Sajda P. Circuitry and the classification of simple and complex cells in V1. Journal of Neurophysiology. 2006;96(5):2739–2749. doi:10.1152/jn.00346.2006.

74. Wielaard J, Sajda P. Extraclassical receptive field phenomena and short-range connectivity in V1. Cerebral Cortex. 2006;16(11):1531–1545. doi:10.1093/cercor/bhj090.

75. Hill S, Tononi G. Modeling sleep and wakefulness in the thalamocortical system. Journal of neurophysiology. 2005;93(3):1671–98. doi:10.1152/jn.00915.2004.

76. Lumer ED, Edelman GM, Tononi G. Neural dynamics in a model of the thalamocortical system. I. Layers, loops and the emergence of fast synchronous rhythms. Cerebral Cortex. 1997;7:207–27.

77. Lumer ED, Edelman GM, Tononi G. Neural dynamics in a model of the thalamocortical system . II . The role of neural synchrony tested through perturbations of spike timing. Cerebral Cortex. 1997;7:228–236.

78. Esser SK, Hill SL, Tononi G. Sleep homeostasis and cortical synchronization: I. Modeling the effects of synaptic strength on sleep slow waves. Sleep. 2007;30(12):1617–1630. doi:10.1093/sleep/30.12.1617.

79. Cai D, Rangan AV, McLaughlin DW. Architectural and synaptic mechanisms underlying coherent spontaneous activity in V1. Proceedings of the National Academy of Sciences of the United States of America. 2005;102(16):5868–5873. doi:10.1073/pnas.0501913102.

80. Tao L, Cai D, McLaughlin DW, Shelley MJ, Shapley R. Orientation selectivity in visual cortex by fluctuation-controlled criticality. Proceedings of the National Academy of Sciences of the United States of America. 2006;103(34):12911–12916. doi:10.1073/pnas.0605415103.

81. Rangan AV, Cai D, McLaughlin DW. Modeling the spatiotemporal cortical activity associated with the line-motion illusion in primary visual cortex. Proceedings of the National Academy of Sciences of the United States of America. 2005;102(52):18793–18800. doi:10.1073/pnas.0509481102.

82. Chariker L, Shapley R, Young LS. Rhythm and synchrony in a cortical network model. Journal of Neuroscience. 2018;38(40):8621–8634. doi:10.1523/JNEUROSCI.0675-18.2018.

83. Chariker L, Shapley R, Hawken M, Young LS. A Computational Model of Direction Selectivity in Macaque V1 Cortex Based on Dynamic Differences between On and Off Pathways. Journal of Neuroscience. 2022;42(16):3365–3380. doi:10.1523/JNEUROSCI.2145-21.2022.

84. Chariker L, Shapley R, Young LS. Contrast response in a comprehensive network model of macaque V1. Journal of Vision. 2020;20(4):1–19. doi:10.1167/jov.20.4.16.

85. Ferster D, Miller KD. Neural Mechanisms of Orientation Selectivity in the Visual Cortex. Annual Review of Neuroscience. 2000;23(1):441–471. doi:10.1146/annurev.neuro.23.1.441.

86. Ben-Yishai R, Lev Bar-Or R, Sompolinsky H. Theory of orientation tuning in visual cortex. Proceedings of the National Academy of Sciences of the United States of America. 1995;92(9):3844–3848. doi:10.1073/pnas.92.9.3844.

87. Douglas R, Koch C, Mahowald M, Martin K, Suarez H. Recurrent excitation in neocortical circuits. Science. 1995;269(5226):981–985. doi:10.1126/science.7638624.

88. Ko H, Hofer SB, Pichler B, Buchanan KA, Sjöström PJ, Mrsic-Flogel TD. Functional specificity of local synaptic connections in neocortical networks. Nature. 2011;473(7345):87–91. doi:10.1038/nature09880.

89. Arkhipov A, Gouwens NW, Billeh YN, Gratiy S, Iyer R, Wei Z, et al. Visual physiology of the layer 4 cortical circuit in silico. PLOS Computational Biology. 2018;14(11):e1006535. doi:10.1371/journal.pcbi.1006535.

90. Fournier J, Monier C, Levy M, Marre O, Sari K, Kisvarday ZF, et al. Hidden Complexity of Synaptic Receptive Fields in Cat V1. Journal of Neuroscience. 2014;34(16):5515–5528. doi:10.1523/JNEUROSCI.0474-13.2014.

91. Frégnac Y, Bathellier B. Cortical Correlates of Low-Level Perception: From Neural Circuits to Percepts. Neuron. 2015;88:110–126. doi:10.1016/j.neuron.2015.09.041.

92. Gerard-Mercier F, Carelli PV, Pananceau M, Troncoso XG, Frégnac Y. Synaptic correlates of low-level perception in V1. Journal of Neuroscience. 2016;36(14):3925–3942. doi:10.1523/JNEUROSCI.4492-15.2016.

93. Stepanyants A, Martinez LM, Ferecsko AS, Kisvarday ZF. The fractions of short- and long-range connections in the visual cortex. Proceedings of the National Academy of Sciences. 2009;106(9):3555–3560. doi:10.1073/pnas.0810390106.

94. Martinez LM, Wang Q, Reid RC, Pillai C, Alonso JM, Sommer FT, et al. Receptive field structure varies with layer in the primary visual cortex. Nature Neuroscience. 2005;8(3):372–379. doi:10.1038/nn1404.

95. Allen Brain Observatory contributors. Allen Institute for Brain Science: Allen Brain Observatory; 2016. http://observatory.brain-map.org/visualcoding.

96. Thomson AM. Neocortical layer 6, a review. Frontiers in neuroanatomy. 2010;4(March):13. doi:10.3389/fnana.2010.00013.

97. Thomson AM. Interlaminar Connections in the Neocortex. Cerebral Cortex. 2003;13(1):5–14. doi:10.1093/cercor/13.1.5.

98. Wang W, Jones HE, Andolina IM, Salt TE, Sillito AM. Functional alignment of feedback effects from visual cortex to thalamus. Nature Neuroscience. 2006;9(10):1330–1336. doi:10.1038/nn1768.

99. Sillito AM, Jones HE, Gerstein GL, West DC. Feature-linked synchronization of thalamic relay cell firing induced by feedback from the visual cortex. Nature. 1994;369(6480):479–482. doi:10.1038/369479a0.

100. Angelucci A, Bullier J. Reaching beyond the classical receptive field of V1 neurons: Horizontal or feedback axons? Journal of Physiology Paris. 2003;97(2-3):141–154. doi:10.1016/j.jphysparis.2003.09.001.

101. Budd JML. Extrastriate feedback to primary visual cortex in primates: A quantitative analysis of connectivity. Proceedings of the Royal Society B: Biological Sciences. 1998;265(1400):1037–1044. doi:10.1098/rspb.1998.0396.

102. Xu X, Callaway EM. Laminar specificity of functional input to distinct types of inhibitory cortical neurons. Journal of Neuroscience. 2009;29(1):70–85. doi:10.1523/JNEUROSCI.4104-08.2009.

103. Markov NT, Vezoli J, Chameau P, Falchier A, Quilodran R, Huissoud C, et al. Anatomy of hierarchy: Feedforward and feedback pathways in macaque visual cortex. Journal of Comparative Neurology. 2014;522(1):225–259. doi:10.1002/cne.23458.

104. Guarino DG, Davison AP, Frégnac Y, Antolík J. The cortico-thalamic loop attunes competitive lateral interactions across retinotopic and orientation preference maps. BioRxiv. 2022;.

105. Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M. Inhibition of inhibition in visual cortex: The logic of connections between molecularly distinct interneurons. Nature Neuroscience. 2013;16(8):1068–1076. doi:10.1038/nn.3446.

106. Harris KD, Mrsic-Flogel TD. Cortical connectivity and sensory coding; 2013. Available from: `http://www.nature.com/doifinder/10.1038/nature12654`.

107. Fourcaud-Trocmé N, Hansel D, van Vreeswijk C, Brunel N. How spike generation mechanisms determine the neuronal response to fluctuating inputs. Journal of Neuroscience. 2003;23(37):11628–40.

108. Ascoli GA, Donohue DE, Halavi M. NeuroMorpho.Org: A Central Resource for Neuronal Morphologies. Journal of Neuroscience. 2007;27(35):9247–9251. doi:10.1523/JNEUROSCI.2055-07.2007.

109. Kremkow J, Jin J, Komban SJ, Wang Y, Lashgari R, Li X, et al. Neuronal nonlinearity explains greater visual spatial resolution for darks than lights. Proceedings of the National Academy of Sciences of the United States of America. 2014;111(8):3170–5. doi:10.1073/pnas.1310442111.

110. Wang Y, Jin J, Kremkow J, Lashgari R, Komban SJ, Alonso JM. Columnar organization of spatial phase in visual cortex. Nature Neuroscience. 2014;18(1):97–103. doi:10.1038/nn.3878.

111. Jones JP, Palmer LA. The two-dimensional spatial structure of simple receptive fields in cat striate cortex. Journal of Neurophysiology. 1987;58(6):1187–1211. doi:10.1152/jn.1987.58.6.1187.

112. Ringach DL. Mapping receptive fields in primary visual cortex. Journal of Physiology. 2004;558(3):717–728. doi:10.1113/jphysiol.2004.065771.

113. Kayser AS, Miller KD. Opponent Inhibition: A Developmental Model of Layer 4 of the Neocortical Circuit. Neuron. 2002;33(1):131–142. doi:10.1016/S0896-6273(01)00570-0.

114. Sinha A, de Schepper R, Pronold J, Mitchell J, Mørk H, Nagendra Babu P, et al.. NEST 3.4; 2023. Available from: `https://doi.org/10.5281/zenodo.6867800`.

115. Tusa RJ, Palmer La, Rosenquist aC. The retinotopic organization of area 17 (striate cortex) in the cat. The Journal of comparative neurology. 1978;177(2):213–35. doi:10.1002/cne.901770204.

116. Beaulieu C, Colonnier M. Number of neurons in individual laminae of areas 3B, 4γ, and $6a\alpha$ of the cat cerebral cortex: A comparison with major visual areas. Journal of Comparative Neurology. 1989;279(2):228–234. doi:10.1002/cne.902790206.

117. Beaulieu C, Kisvarday Z, Somogyi P, Cynader M, Cowey A. Quantitative distribution of gaba-immunopositive and -immunonegative neurons and synapses in the monkey striate cortex (area 17). Cerebral Cortex. 1992;2(4):295–309. doi:10.1093/cercor/2.4.295.

118. Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C. Interneurons of the neocortical inhibitory system. Nature Reviews Neuroscience. 2004;5(10):793–807. doi:10.1038/nrn1519.

119. Angelucci A, Levitt JB, Walton EJS, Hupé JM, Bullier J, Lund JS. Circuits for local and global signal integration in primary visual cortex. Journal of Neuroscience. 2002;22(19):8633–8646. doi:10.1523/jneurosci.22-19-08633.2002.

120. McCormick DA, Connors BW, Lighthall JW, Prince DA. Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. Journal of Neurophysiology. 1985;54(4):782–806. doi:10.1152/jn.1985.54.4.782.

121. Naud R, Marcille N, Clopath C, Gerstner W. Firing patterns in the adaptive exponential integrate-and-fire model. Biological Cybernetics. 2008;99(4-5):335–347. doi:10.1007/s00422-008-0264-7.

122. Antolík J. Rapid long-range disynaptic inhibition explains the formation of cortical orientation maps. Frontiers in Neural Circuits. 2017;11(March):1–15. doi:10.3389/fncir.2017.00021.

123. Movshon JA, Thompson ID, Tolhurst DJ. Spatial and temporal contrast sensitivity of neurones in areas 17 and 18 of the cat's visual cortex. The Journal of Physiology. 1978;283(1):101–120. doi:10.1113/jphysiol.1978.sp012490.

124. Payne B, Peters A. The Cat Primary Visual Cortex. Elsevier Science; 2001. Available from: `https://books.google.cz/books?id=tBlZA2NSQWsC`.

125. Beaulieu C, Colonnier M. A laminar analysis of the number of round asymmetrical and flat symmetrical synapses on spines, dendritic trunks, and cell bodies in area 17 of the cat. Journal of Comparative Neurology. 1985;231(2):180–189. doi:10.1002/cne.902310206.

126. Johnson RR, Burkhalter A. Microcircuitry of forward and feedback connections within rat visual cortex. Journal of Comparative Neurology. 1996;368(3):383–398. doi:10.1002/(SICI)1096-9861(19960506)368:3¡383::AID-CNE5¿3.0.CO;2-1.

127. Stratford KJ, Tarczy-Hornoch K, Martin KAC, Bannister NJ, Jack JJB. Excitatory synaptic inputs to spiny stellate cells in cat visual cortex. Nature. 1996;382(6588):258–261. doi:10.1038/382258a0.

128. Allen C, Stevens CF. An evaluation of causes for unreliability of synaptic transmission. Proceedings of the National Academy of Sciences of the United States of America. 1994;91(October):10380–10383. doi:10.1073/pnas.91.22.10380.

129. Izhikevich EM, Edelman GM. Large-scale model of mammalian thalamocortical systems. Proceedings of the National Academy of Sciences of the United States of America. 2008;105(9):3593–3598. doi:10.1073/pnas.0712231105.

130. Budd JM, Kisvárday ZF. Local lateral connectivity of inhibitory clutch cells in layer 4 of cat visual cortex (area 17). Experimental Brain Research. 2001;140(2):245–250. doi:10.1007/s002210100817.

131. Lee WCA, Bonin V, Reed M, Graham BJ, Hood G, Glattfelder K, et al. Anatomy and function of an excitatory network in the visual cortex. Nature. 2016;532(7599):370–374. doi:10.1038/nature17192.

132. Hoffmann JHO, Meyer HS, Schmitt AC, Straehle J, Weitbrecht T, Sakmann B, et al. Synaptic conductance estimates of the connection between local inhibitor interneurons and pyramidal neurons in layer 2/3 of a cortical column. Cerebral Cortex. 2015;25(11):4415–4429. doi:10.1093/cercor/bhv039.

133. Cruikshank SJ, Lewis TJ, Connors BW. Synaptic basis for intense thalamocortical activation of feedforward inhibitory cells in neocortex. Nature Neuroscience. 2007;10(4):462–468. doi:10.1038/nn1861.

134. Abbott LF. Synaptic Depression and Cortical Gain Control. Science. 1997;275(5297):221–224. doi:10.1126/science.275.5297.221.

135. Markram H, Wang Y, Tsodyks M. Differential signaling via the same axon of neocortical pyramidal neurons. Proceedings of the National Academy of Sciences of the United States of America. 1998;95(9):5323–5328. doi:10.1073/pnas.95.9.5323.

136. Banitt Y, Martin KAC, Segev I. A biologically realistic model of contrast invariant orientation tuning by thalamocortical synaptic depression. Journal of Neuroscience. 2007;27(38):10230–10239. doi:10.1523/JNEUROSCI.1640-07.2007.

137. Beierlein M, Gibson JR, Connors BW. Two Dynamically Distinct Inhibitory Networks in Layer 4 of the Neocortex. Journal of Neurophysiology. 2003;90(5):2987–3000. doi:10.1152/jn.00283.2003.

138. Ma Y, Hu H, Agmon A. Short-term plasticity of unitary inhibitory-to-inhibitory synapses depends on the presynaptic interneuron subtype. Journal of Neuroscience. 2012;32(3):983–988. doi:10.1523/JNEUROSCI.5007-11.2012.

139. Bringuier V, Chavane F, Glaeser L, Frégnac Y. Horizontal propagation of visual activity in the synaptic integration field of area 17 neurons. Science. 1999;283(5402):695–699. doi:10.1126/science.283.5402.695.

140. Frégnac Y. Reading Out the Synaptic Echoes of Low-Level Perception in V1. In: Fusiello A, Murino V, Cucchiara R, editors. Lecture Notes in Computer Science 1. vol. 7583 of Lecture Notes in Computer Science. Springer; 2012. p. 486–495. Available from: `http://dx.doi.org/10.1007/978-3-642-33863-2{_}50http://link.springer.com/10.1007/978-3-642-33863-2{_}50`.

141. Chavane F, Sharon D, Jancke D, Marre O, Frégnac Y, Grinvald A. Lateral spread of orientation selectivity in V1 is controlled by intracortical cooperativity. Frontiers in Systems Neuroscience. 2011;5(FEBRUARY 2011):1–26. doi:10.3389/fnsys.2011.00004.

142. Ohana O, Portner H, Martin KAC. Fast Recruitment of Recurrent Inhibition in the Cat Visual Cortex. PLoS ONE. 2012;7(7):e40601. doi:10.1371/journal.pone.0040601.

143. Cai D, DeAngelis GC, Freeman RD. Spatiotemporal receptive field organization in the lateral geniculate nucleus of cats and kittens. Journal of Neurophysiology. 1997;78(2):1045–1061. doi:10.1152/jn.1997.78.2.1045.

144. Alitto HJ, Usrey WM. Influence of contrast on orientation and temporal frequency tuning in ferret primary visual cortex. Journal of Neurophysiology. 2004;91(6):2797–2808. doi:10.1152/jn.00943.2003.

145. Skottun BC, De Valois RL, Grosof DH, Movshon JA, Albrecht DG, Bonds AB. Classifying simple and complex cells on the basis of response modulation. Vision Research. 1991;31(7-8):1078–1086. doi:10.1016/0042-6989(91)90033-2.