Quantitative proteomics reveals key roles for post-transcriptional gene regulation in the

molecular pathology of FSHD

Sujatha Jagannathan[1,2,3,#,*], Yuko Ogata[4], Philip R. Gafken[4], Stephen J. Tapscott[3*], and Robert

K. Bradley[1,2*]

[1]Computational Biology Program, Public Health Sciences Division, Fred Hutchinson Cancer

Research Center, Seattle, WA 98109, USA

[2]Basic Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA

[3]Human Biology Division, Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA

[4]Proteomics Shared Resource, Fred Hutchinson Cancer Research Center, Seattle, WA 98109,

USA

[#]Current affiliation: Department of Biochemistry and Molecular Genetics & RNA Bioscience

Initiative, University of Colorado Denver School of Medicine, Aurora, CO 80045, USA

*Corresponding authors

Email: rbradley@fredhutch.org; sujatha.jagannathan@ucdenver.edu; stapscot@fredhutch.org

Abstract

DUX4 is a transcription factor whose misexpression in skeletal muscle causes facioscapulohumeral muscular dystrophy (FSHD). While DUX4's transcriptional activity has been extensively characterized, the DUX4-induced proteome remains undescribed. Here, we report concurrent measurement of RNA and protein levels in DUX4-expressing cells via RNA-seq and quantitative mass spectrometry. DUX4 transcriptional targets were robustly translated, confirming the likely clinical relevance of proposed FSHD biomarkers. However, a multitude of mRNAs and proteins exhibited discordant expression changes upon DUX4 expression. Our dataset revealed unexpected proteomic, but not transcriptomic, dysregulation of diverse molecular pathways, including Golgi apparatus fragmentation, as well as extensive post-transcriptional buffering of stress response genes. Key components of RNA degradation machineries, including UPF1, UPF3B, and XRN1, exhibited suppressed protein, but not mRNA, levels, explaining the build-up of aberrant RNAs that characterizes DUX4-expressing cells. Our results provide a resource for the FSHD community and illustrate the importance of post-transcriptional processes to DUX4-induced pathology.

Introduction

Facioscapulohumeral muscular dystrophy is caused by the inappropriate expression of an early embryonic transcriptional activator, DUX4, in adult muscle, leading to cell death (1, 2). Decades of work have generated a detailed parts list of the genes and pathways affected by DUX4 that may underlie FSHD pathophysiology (3-10). However, an integrated model for how those DUX4-induced changes lead to disease has remained elusive (11-13). As transient and pulsatile expression of DUX4 is sufficient induce pathology and cell death (14), it is critical that we understand the cellular events and pathways set in motion by DUX4 that lead to eventual cell death in order to develop effective therapeutics for FSHD.

DUX4 induces changes in the expression of hundreds of genes that impact dozens of highly interconnected pathways (3-10), making a cause-and-effect relationship between dysregulated gene expression and FSHD pathology difficult to discern. Because DUX4 is a strong transcriptional activator, most studies of DUX4 activity have focused on measuring gene expression at the transcript level (3, 14, 15), thereby implicitly assuming that the transcriptome accurately represents the cellular proteome in DUX4-expressing cells. While this is a reasonable assumption, it is well known that RNA and protein levels are not always concordant and that post-transcriptional regulation can results in divergent RNA and protein levels (16). While a few proteomics studies have been conducted on FSHD muscle biopsies, these early studies lack the depth necessary to make meaningful comparisons with the DUX4-induced transcriptome (17-19). Furthermore, given our recent discovery that DUX4 induces proteolysis of a key RNA binding protein, UPF1 (7), we hypothesized that conducting paired measurements of RNA and protein levels might be particularly important in order to identify altered post-transcriptional gene regulation in DUX4-expressing cells. Hence, we set out to generate reliable RNA- and protein-level measurements of DUX4-induced gene expression and thereby elucidate the extent of post-transcriptional gene dysregulation in DUX4-expressing cells.

We used our previously established and validated cell culture models of DUX4 expression

(20) to conduct RNA-seq and Stable Isotope Labeling with Amino acids in Cell culture (SILAC)

coupled with quantitative mass spectrometry (21). The resulting data enabled us to measure

DUX4-induced alterations in the transcriptome as well as protein levels for ~4,000 genes with

high confidence. Comparing the transcript-level and protein-level alterations revealed three

distinct patterns of expression for different subsets of genes: 1) Concordant changes in

expression at the RNA and protein level for many transcriptional targets of DUX4; 2) Post-

transcriptional buffering of the expression of many genes, especially of those involved in stress

response pathways; and 3) Discordant gene expression changes at the RNA versus protein levels

for many genes, particularly those involved in RNA surveillance. Together, these findings highlight

the importance of measuring the expressed proteome in order to fully understand DUX4 biology

and the FSHD disease process.

Results

Determining protein-level alterations in DUX4-expressing cells via quantitative mass spectrometry.

In order to measure DUX4-induced changes to the cellular proteome, we conducted SILAC-based mass spectrometry in two independent DUX4 expression systems (**Figure 1A).** We previously showed that DUX4 expressed via a lentiviral vector versus an inducible transgene integrated into the genome of a myoblast cell line both yield comparable gene expression profiles that accurately capture the transcriptome of FSHD cells (20). Here, we used both of these expression systems in order to internally corroborate our results, thereby ensuring that our proteomic data were robust with respect to choice of model system.

We first established the efficiency of SILAC labeling and determined appropriate cell culture conditions and experimental timing in order to ensure near-complete labeling of proteins with heavy isotopic Arginine and Lysine. To this end, we cultured immortalized MB135 myoblasts in SILAC labeling media for a week and subjected the total protein from labeled cells, as well as a 1:1 mix of labeled to unlabeled cells, to mass spectrometry. Using the resulting mass spectra, we calculated labeling efficiency in two ways. First, we quantified the relative abundance of heavy and light spectra for several individual peptides and found that it showed a >95% labeling efficiency (a representative example shown in **Figure S1A**). Next, we analyzed the proteome-wide distribution of log-transformed heavy to light ratios of peptides from both the heavy-only (**Figure S1B**) and heavy:light mixed samples (**Figure S1C**). The distribution of heavy-to-light ratios was strongly skewed to the right in the heavy-only samples, indicating robust labeling with the heavy amino acids. In contrast, the heavy:light mixed sample yielded a distribution centered roughly around zero, as expected. Based on these data, we proceeded with the labeling conditions and increased the labeling time even further (to three weeks) in order to achieve maximal SILAC labeling of the MB135 myoblasts.

MB135 myoblasts were adapted to light or heavy SILAC media for 3 weeks were

transduced with lentivirus carrying DUX4 (vDUX4) or GFP (vGFP) expression constructs.

Samples were collected 24- and 36-hours post-transduction. In an independent experiment,

MB135 cells carrying a doxycycline-inducible DUX4 transgene (iDUX4; (20)) were adapted to

SILAC media for 3 weeks and DUX4 expression was induced with 1µg/ml of doxycycline for 14

hours in two replicates carrying heavy and light SILAC labels. Paired controls with no treatment

were also collected with both heavy and light labels. Total protein from cells expressing DUX4

were mixed with an equal amount of total protein from cells without DUX4 expression containing

the opposite SILAC label to generate samples that were then subjected to mass spectrometry.

Peptide-spectrum matches (PSMs) with quantified heavy to light ratios were subject to

thorough screening for quality (e.g., filtering out single-peak spectra and spectra without unique

mapping; **Figure 1A**; see *Materials & Methods* for further details; Supplemental Data Set 1). A

histogram of log-transformed DUX4 to Control abundance ratios of peptides mapping to a DUX4

target gene, ZSCAN4, from both vDUX4 and iDUX4 datasets showed highly skewed $\log_2$

(DUX4/Control) ratio, consistent with significant upregulation of the protein upon DUX4

expression (**Figure 1B-C**). In contrast, plotting the $\log_2$ (DUX4/Control) ratio of all individual

peptides mapping to the housekeeping gene RPL15 showed that the ratio is centered around

zero (**Figure 1B-C**), as would be expected for a gene with no differential expression upon DUX4

induction. These example plots illustrate the strong agreement between the expected and

observed protein fold change values determined by SILAC mass spectrometry. Moreover, out of

the 65 genes identified by Yao *et al* (22) as potential FSHD biomarkers based on transcriptome

analysis of FSHD patient samples, 8 were quantified in the vDUX4 proteomics study and 25 were

quantified in the iDUX4 proteomics study and both show high induction at the protein level (**Figure

1D, E**). Note that the lower number of peptides (and hence quantified proteins) from the vDUX4

sample indicates the lower depth of this dataset, and yet yields fold changes that are highly

consistent with the iDUX4 dataset with a higher depth.

Assessing concordance of RNA and protein expression fold change.

Next, using the iDUX4 dataset, we performed peptide to protein summarization via median

heavy/light ratios of all the peptides mapping to a certain protein in both replicates to obtain gene-

level $\log_2$ (DUX4/Control) ratios (**Figure 1A and S1D-E**; Supplemental Data Set 2). After filtering

out genes that were only observed in one of the two label-swap replicates, we obtained

quantitative proteomics information for 4005 genes, 3961 of which also had a corresponding RNA-

seq measurement (**Figure 2A;** RNA-seq data previously reported in (20)). The lower number of

genes quantified via proteomics compared to RNA-seq is expected as proteomics is known to

have lower sensitivity compared to RNA-seq.

To qualitatively compare the RNA and protein expression level changes upon DUX4

expression, we assessed the overlap of genes with an expression change of 4-fold or above.

Among genes that are upregulated (> 2 log2 fold change), the concordance between RNA and

protein was roughly 40-50%, whereas similarly downregulated genes show very little concordance

(**Figure 2A**). To obtain a more quantitative measure of concordance, we generated a scatter plot

of the RNA versus protein fold change for the 3961 genes (**Figure 2B**). We found a reasonable

level of correlation between these values with a Pearson's correlation coefficient, r, of 0.51 (p-

value < 2.2e-16).

To assess whether similar pathways were affected at the RNA versus protein level, we

conducted Gene Ontology (GO) analysis for genes up- or down-regulated at the RNA and protein

(**Figure 2C**). Surprisingly, we observed that the pathways that are affected at the RNA versus

protein levels are quite distinct. While the transcript level changes occur in genes involved in

transcription and mRNA processing, protein-level changes impact pathways including humoral

immune response, proteolysis and exocytosis. As the exocytosis pathway has not been implicated

in any of the previous DUX4 gene expression studies, we sought to examine this phenomenon

further by imaging the Golgi apparatus, which is the source of exocytotic vesicles in the cell (23).

We found that DUX4 expressing cells showed severe fragmentation of Golgi apparatus, which

could be an indicator of a perturbation in the cellular secretory pathways (24) (**Figure 2D;**

quantified in **Figure 2E**).

Taken together, these results demonstrate that analyzing the protein measurements may

give us insights that are not discernable in the transcriptome fold change analysis performed in

earlier studies.

Post-transcriptional buffering of stress response genes may exacerbate DUX4 toxicity.

Though many of the genes induced at the transcript level are largely also induced at the

protein-level, a subset of genes showed no change in their protein level despite their transcripts

were up- or down-regulated to a significant degree (678 genes, shaded blue in **Figure 3A**),

indicating post-transcriptional buffering of the protein levels. Most notably, several housekeeping

genes that respond to protein folding stress or dsRNA-induced stress showed transcriptional

upregulation with minimal protein-level upregulation (**Figure 3B**).

Given that both unfolded protein and dsRNA induced stresses converge in

phosphorylation of eIF2α and lead to translation inhibition (25), we asked if the timing of

transcription of various stress-response genes coincide with translational downregulation. We

found that HSPA5, a prominent marker of the unfolded protein response pathway (26), shows

transcriptional upregulation at a time period that temporally coincides with eIF2α phosphorylation

and reduced incorporation of [^{35}S]-labeled methionine, a proxy for bulk translation efficiency

(**Figure 3 C-E**). These data demonstrate that translation inhibition caused by various cellular

stresses and the resulting post-transcriptional buffering prevents DUX4-expressing cells from

mounting a robust stress response.

Post-transcriptional modulation of RNA quality control pathway by DUX4.

Next, we focused our analysis on the subset of genes that showed significant changes at

the protein level with either no change in their transcript abundance or a change in the opposite

direction (198 genes shown as 'gold' circles in **Figure 4A**). Pathway analysis did not reveal any

significant trends among these genes. So instead, we decided to focus on one of the pathways

that we have previously shown to be post-transcriptionally modulated – namely, the nonsense-

mediated RNA decay (NMD) pathway (7).

A diagram showing RNA- vs protein-level changes to various components of this pathway

demonstrates substantial post-transcriptional regulation in this pathway (**Figure 4B**). Many of

these genes, including UPF1, UPF2, UPF3B and XRN1, showed downregulation at the protein

level. The downregulation of XRN1 is of particular interest as it is the 5'-3' exonuclease that

degrades NMD targets upon cleavage by the endonuclease, SMG6 (27). Moreover, SMG6 too is

downregulated to a log2 fold change of -4.7, though it is only detected via a single peptide and

hence was filtered out of our analysis. Thus, DUX4-induced NMD inhibition appears to be a result

of post-transcriptional downregulation of multiple key players of the NMD pathway, which explains

the severity of NMD inhibition in DUX4-expressing cells.

Post-transcriptional downregulation of a gene can be achieved via two means: reduced

translation or increased protein degradation. We have previously shown that DUX4 induces

proteasome-mediated degradation of UPF1 (7). Hence, we asked if DUX4 affects known

components and regulators of the ubiquitin proteasome. A scatterplot of all ubiquitin proteasome

regulators shows a change in the expression of several such genes, one or more of which may

underlie the rapid degradation of UPF1 (**Figure 4C**). Further studies are needed to understand

the precise molecular mechanism behind this regulatory pathway and its downstream

consequences. In summary, we propose that the post-transcriptional gene regulation plays a

critical role in inhibiting NMD and perturbing the proteostasis in DUX4-expressing cells and may

thus underlie key aspects of FSHD pathology (**Figure 4D**).

Finally, in order to enable researchers and patients in the FSHD community to access the

data generated in this study, we developed a web tool for easy visualization of these data

(Screenshot shown in **Figure S2**). This tool can be freely accessed at the following URL:

https://dynamicrna.shinyapps.io/dataviz/.

Discussion

Most of the highly induced DUX4 transcriptional targets are germline and early embryonic genes that are normally never expressed in adult muscle (3). So, it is possible that despite being expressed at the transcript level, such genes may be translated poorly and/or be degraded rapidly upon translation due to the lack of cell-type chaperones or other factors. Here, we used quantitative mass spectrometry in two different cell culture models of FSHD to demonstrate that DUX4-induced transcripts are efficiently translated into stable proteins in muscle cells. As we previously demonstrated that our vDUX4 and iDUX4 systems accurately capture the transcriptional program of FSHD cells (20), it is reasonable to assume that DUX4-induced transcripts are similarly translated into stable proteins in FSHD muscle. However, further work is required to definitively confirm this. Similarly, our data strongly motivate focused investigation of these DUX4-induced proteins to test their utility as potential FSHD biomarkers in relevant clinical samples (22).

Next, we asked if the changes in the DUX4 proteome are largely reflective of the changes to the transcriptome. We found that this is not the case. While there is a positive correlation between these measurements (Pearson's correlation coefficient of 0.51), hundreds of genes deviate from this trend. GO analysis of the most differentially expressed genes at the transcript versus protein levels revealed RNA splicing and processing as being the prominent categories impacted at the transcript level, while protein-level changes impacted an entirely different set of pathways. We take these results as indication that the transcriptome-level analysis paints an incomplete picture of DUX4 biology that should be complemented with proteome-level analysis to thoroughly understand how DUX4 misexpression causes FSHD.

We next pursued the various mechanisms by which protein-level changes deviate from the corresponding transcriptomic changes. We found that genes induced by dsRNA and unfolded protein stress are transcriptionally induced, but translationally buffered as a result of the

translational repression that accompanies activation of these stress-response pathways.

Motivated by this result, we hypothesize that DUX4-expressing cells are unable to mount a robust

stress response despite inducing the transcripts necessary to alleviate stress. We also show that

multiple proteins in RNA surveillance pathways, including UPF1 and XRN1, are downregulated

at the protein level, which may explain the drastic reduction in RNA quality control capacity that

we observed in DUX4-expressing cells. From the proteomics data, we identified many genes

involved in the ubiquitin proteasome pathway whose mis-regulation could alter protein stability.

These genes may serve as a starting point for further investigation of how DUX4, which is best

known as a transcriptional activator, induces widespread alterations in protein turnover.

Materials and Methods

Accession codes. Mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE partner repository (Vizcaino et al., 2013) with the

dataset identifier PXD010221 (Reviewer account details: Username: reviewer37110@ebi.ac.uk;

Password: aMrE4BW0). The RNA-seq data are available through the NCBI SRA database under

accession number GSE85461 (20).

Cell culture and SILAC labeling. Proliferating human myoblasts were cultured in F10 medium

(Gibco/Life Technologies) supplemented with 20% fetal bovine serum (Thermo Scientific), 10ng

bFGF (Life Technologies), 1μM dexamethasone (Sigma) and 50U/50μg penicillin/streptomycin

(Life Technologies). Cells were labeled in SILAC media containing heavy lysine (Lys8) and

arginine (Arg10) for 3 weeks before DUX4 induction experiments. To induce DUX4 expression in

the MB135 iDUX4 cells, 1μg/ml of doxycycline was added for 8 or 14 hours, as indicated. For viral

DUX4 expression, MB135 cells were transduced with lentivirus carrying DUX4 coding sequence

under a hPGK promoter in the presence of polybrene.

Gel slice digestion. Total RNA and protein were extracted from whole cells using TRIzol[RT]

reagent (Ambion) following the manufacturer's instructions. 50 μg of total protein was subjected

to SDS PAGE using a 4-15% bis-TRIS gel. The gel was stained using GelCode blue (Pierce) as

per manufacturer's instructions, destained overnight in ultrapure water and the entire lane

containing protein was cut into 16 fractions using a GelCutter (Gel company Inc.). Individual gel

slices in 1.5 mL tubes (Eppendorf) were consecutively washed with water and incubated with

25mM ammonium bicarbonate, 50% acetonitrile for 2 hrs. The gel pieces were dehydrated with

acetonitrile. And the dried gel slices were reduced by covering them with 10 mM dithiothreitol in

100 mM ammonium bicarbonate and heating them at 56^{o}C for 45 min. The solution was removed

and discarded. The gel slices were alkylated by covering them with a solution of 50 mM

iodoacetamide in 100 mM ammonium bicarbonate and incubating in the dark at ambient

temperature for 30 min. The solution was removed and discarded. The gel slices were dehydrated

with acetonitrile, then washed with 100 mM ammonium bicarbonate 100 mM for 10 min. The

solution was removed, discarded and the gel slices were dehydrated once again with acetonitrile.

After removing acetonitrile, the gel slices were then hydrated with 5 ng/uL sequencing grade

trypsin (Promega) in 50 mM ammonium bicarbonate and digested overnight at 37^{o}C on an orbital

shaker. Following digestion, the supernatants were collected, and the gel slices were washed with

0.1% trifluoroacetic acid, and after 30 min an equal volume of acetonitrile was added followed by

washing for an additional 1 hour. The original digestion supernatant and the wash for a single

sample were combined into a single tube and dried by vacuum centrifugation. The digestion

products were desalted using Ziptips (Millipore) per the manufacturer's instructions, eluted with

70% acetonitrile/0.1% trifluoroacetic acid, and dried by vacuum centrifugation.

Mass spectrometry. The desalted material was resuspended in 20 µL of 2% acetonitrile in 0.1%

formic acid, and 18 µL was analyzed using one of two LC/ESI MS/MS configurations. The first

configuration consisted of an Easy-nLC II (Thermo Scientific) coupled to a Orbitrap Elite ETD

(Thermo Scientific) mass spectrometer using a trap-column configuration as described (28). A

trap of 100 µm × 20 mm packed with Magic C_{18}AQ (5-µm, 200 Å resin; Michrom Bioresources)

packing material was used for in-line desalting and a column of 75 µm × 250 mm packed with

C_{18}AQ (5-µm, 100 Å resin; Michrom Bioresources) was used for analytical peptide separations.

Chromatographic separations were carried out using a 60-minute gradient from 5% to 35%

solvent B (solvent A: 0.1% formic acid, solvent B: 0.1% formic acid in acetonitrile) at a flowrate of

300 nL/min. The analytical column temperature was maintained at 40℃. The Orbitrap Elite

instrument was operated in the data-dependent mode, switching automatically between MS

survey scans in the Orbitrap (AGC target value 1E6, resolution 240,000, and maximum injection

time 250 ms) and collision induced dissociation (CID) MS/MS spectra acquisition in the linear ion

trap (AGC target value of 10,000 and injection time 100 ms). The twenty most intense precursor

ions from the OrbiTrap full scan were each consecutively selected for fragmentation by CID in the

linear ion trap using a normalized collision energy of 35%. Ions of +2 and +3 charge states were

selected for MS/MS and selected ions were dynamically excluded for 30 seconds. The second

configuration consisted of an Easy nanoLC 1000 (Thermo Scientific) HPLC connected to an

OrbiTrap Fusion (Thermo Scientific) mass spectrometer. In-line chromatographic separations (no

trap column) were carried out using a 75 μm × 400 mm column packed with Magic $C_{18}AQ$ (5-μm,

100 Å resin; Michrom Bioresources) packing material at a flowrate of 300 nL/min.

Chromatographic elution consisted of a 90-minute gradient from 3% to 27% B and the column

temperature was maintained at 40℃. The OrbiTrap Fusion was operated in the 2 second "top

speed" data dependent acquisition mode with MS survey scans in the OrbiTrap at least every 2

seconds (AGC target value 4E5, resolution 120,000, and maximum injection time of 50 ms).

Quadrupole isolation was set to 1.6 FWHM and higher energy collision dissociation (HCD) was

used for fragmentation at a collision energy of 28% and MS/MS detection was carried out in the

linear ion trap set at rapid scan speed (injection time of 250 ms and AGC target of 10E2).

Positively charged ions from 2 to 6 were selected for MS/MS and selected ions were dynamically

excluded for 30 seconds.

Data analysis and statistical methods. Qualitative and quantitative data analysis were

performed using Proteome Discoverer 2.1 (Thermo Scientific). The data were searched against

a human UniProt database (downloaded 11-04-16) that was appended with protein sequences

from the common Repository of Adventitious Proteins (cRAP; www.thegpm.org/crap/) an *in silico*

translation products of noncanonical transcript isoforms stabilized due to NMD inhibition. In

downstream analyses, peptides that only mapped to NMD targets were not considered any further

in the current study and will be pursued in a future investigation. Searches were conducted with

the trypsin enzyme specificity. The precursor ion tolerance was set to 10 ppm and the fragment

ion tolerance was set to 0.6 Da. Variable modifications were set for oxidation on methionine

(+15.995 Da), carbamidomethyl (+57.021 Da) on cysteine, and acetylation (+42.010 Da) on the

N-terminus of proteins. Heavy SILAC amino acids for lysine (+8.014 Da) and arginine (10.008)

were also accounted for in the analysis as variable modifications. All search results were

evaluated by Percolator (29) for false discovery rate (FDR) evaluation of the identified peptides.

Peptide identifications were filtered to a peptide FDR of 1%.

The peptide-spectrum matches (PSMs) and the corresponding quantification data were

then exported as a tab-delimited text file from Proteome Discoverer 2.1, and all downstream data

analysis was conducted using the R statistical programming language. The Custom R script

(**Supplementary File 3**) to reproduce the analyses described above and figures in this manuscript

is deposited in github, available at the following URL: https://github.com/sjaganna/2018-

jagannathan_et_al. Briefly, we filtered PSMs to remove the spectra carrying the following quality

flags assigned by Proteome Discoverer – "Inconsistently labeled", "Indistinguishable channels",

"No quantitative values", "Not unique", "Redundant spectra", "Single peak spectra", and "Excluded

by method" – all of which represent poor quality spectra. Next, we removed peptides that failed

to map to a gene, and peptides mapping to genes in the cRAP database or to uncharacterized

genes. We then assigned the median peptide fold change within an experiment as the

summarized protein fold change value.

Upon comparing the gene-level fold changes obtained from four different samples, we

noticed a strong correlation between the two viral replicates (Pearson's r = 0.81), as well as the

data from iDUX4 replicate 1 and both vDUX4 datasets (Pearson's r = 0.79 and 0.77). The iDUX4

replicate 2, in which we swapped the isotopic labels used for the control and DUX4 samples,

however, showed poor correlation with the three other samples due to a small set of proteins (103

proteins; 2.6% of total) that did not label efficiently. Proteins that do not label efficiently result in

apparent anticorrelation between label-swap samples. This issue, wherein a small subset of

proteins does not label efficiently, is a well-known problem in SILAC experiments (30). Rather

than ignore this subset of proteins, which would require the imposition of potentially arbitrary

thresholds, we instead followed the standard SILAC analysis procedure of taking the median

value of H/L ratio across the two iDUX4 replicates as the final iDUX4 H/L ratio (30, 31). This

procedure preserves the H/L ratio for proteins that labeled efficiently (the vast majority of the

proteome) while bringing the H/L ratio for proteins that did not label efficiently to approximately 1

(corresponding to no measurable change). We then calculated *p*-values for the fold change

calculated for the combined iDUX4 samples using a bootstrap approach. For a protein with n

peptides whose H/L ratios were used to calculate protein fold change, we calculated the

probability that n peptides chosen randomly out of all measured peptides would yield a similar

fold change by chance. To this end, we generated a null distribution for each protein as follows.

First, the same number of H/L ratios as the number of peptides quantified for that protein were

randomly chosen from the set of all calculated peptide-level H/L ratios (n=229,558). With these

randomly chosen H/L ratios, a fold change was calculated by computing the median. This process

was repeated 1,000 times to generate the null distribution of H/L ratios for that protein. Using this

null distribution, a *p*-value was calculated using a two-tailed Z test.

Gene Ontology analysis. GO analysis was performed via the Overrepresentation Enrichment

Analysis method using WebGestalt server (pmid: 15980575; www.webgestalt.org). One of the

GO categories identified as enriched in this analysis was skin development, which we

subsequently removed from Figure 2C as many of the genes that contributed to this GO category

were extracellular proteins including keratins that could be environmental contaminants.

Immunofluorescence microscopy. Cells were permeabilized with PBS containing 0.1% Triton

X-100 for 5 minutes at room temperature and rinsed thrice in PBS. Primary antibody against

GM130 (Bethyl Laboratories, Cat # A303-402A-T) was diluted 1:200 in PBS and incubated for 1

hour at room temperature. After three washes in PBS, secondary anti-Rabbit TRITC (Jackson

ImmunoResearch, cat # 711-025-152) diluted 1:400 was added and incubated for 45 minutes at

room temperature. Cells were washed thrice in PBS with the nuclear counterstain DAPI included

in the final wash. Images were collected on a Cytation 5 multimode reader (BioTek) and analyzed

using GenPrime software (BioTek).

Figure legends

Figure 1. Quantitative mass spectrometry of DUX4-expressing cells.

A) Schematic of experimental set up and the subsequent data analysis steps to measure protein

fold change in cells expressing vDUX4 or iDUX4.

B) Histogram of normalized, log2-transformed DUX4/Control ratios for all peptides mapping to

ZSCAN4, a DUX4 target gene (top panels), and RPL15, a housekeeping gene (bottom panels),

following 24 and 36 hours of vDUX4 expression.

C) Histogram of normalized, log2-transformed DUX4/Control ratios for all peptides mapping to

ZSCAN4, a DUX4 target gene (top panels), and RPL15, a housekeeping gene (bottom panels),

in the two label-swap replicates of iDUX4 expression.

D) Box plot of normalized, log2-transformed DUX4/Control ratio for peptides corresponding to the

FSHD biomarker transcripts identified by Yao et al (2015) in the vDUX4 dataset. Each dot refers

to an individual peptide that was quantified and the color represents the time point of vDUX4

expression (24 hours in blue and 36 hours in purple).

E) Box plot of normalized, log2-transformed DUX4/Control ratio for peptides corresponding to the

FSHD biomarker transcripts identified by Yao et al (2015) in vDUX4 dataset. Each dot refers to

an individual peptide that was quantified and the color denotes replicate 1 (red) or replicate 2

(purple) of iDUX4 expression.

Figure 2. Concordant changes in RNA and protein abundance of several DUX4

transcriptional targets.

A) Table showing number of genes for which we could quantify fold changes in RNA and protein

levels in the iDUX4 datasets.

B) Scatter plot of $\log_2$ fold change in RNA levels (DUX4/control) versus $\log_2$ fold change in protein

levels (DUX4/control) of all genes.

C) Gene Ontology analysis of genes that are up- or down-regulated > 4-fold at the RNA level or

protein level. Multiple testing correction done via Benjamini-Hochberg procedure.

D) Fluorescence micrographs of cells with and without DUX4 induction for 14h, stained with DAPI

(blue), anti-GM130 (Golgi apparatus; in green) and phalloidin-555 (actin; in red).

E) Quantification of percent of cells with DUX4-positive nuclei and fragmented Golgi apparatus.

Error bars represent standard deviation.

Figure 3. Extensive post-transcriptional buffering of stress response genes in DUX4-

expressing cells.

A) Scatter plot of log2 fold change of RNA levels (DUX4/control) versus log2 fold change of protein

levels (DUX4/control). Genes that show an upregulation at the RNA level, but no significant

change at the protein level fall in the blue rectangle.

B) RNA and protein fold change across the different datasets for representative genes that show

post-transcriptional buffering.

C) Changes in RNA levels of the chaperone HSPA5 over time of expression of vDUX4 expression

(or control) as measured by qRT-PCR. The region shaded in blue represents the region where

vDUX4 sample significantly deviates from Control due to induction of unfolded protein response.

D) Levels of phosphorylated eIF2α in control versus DUX4-expressing cells over an expression

time course as detected by immunoblots.

E) Percentage of [^{35}S]-Methionine incorporation (representing translation efficiency) in DUX4-

expressing cells normalized over control cells during a time course of DUX4 expression.

Figure 4. DUX4 induces post-transcriptional gene regulation.

A) Scatter plot of log2 fold change in RNA levels (DUX4/control) versus log2 fold change in

protein levels (DUX4/control). Genes regulated post-transcriptionally result in drastic changes

in protein level without major changes in RNA levels and are highlighted in gold.

B) Schematic representation of RNA- and protein-level changes for the genes involved in mRNA

surveillance. Colors represent a heat map of actual fold changes in RNA levels (top) and in

protein levels (bottom). Proteins with fewer than two quantified peptides are outlined by dotted

lines.

C) Scatter plot of log2 fold change in RNA levels (DUX4/control) versus log2 fold change in

protein levels (DUX4/control) for genes in the ubiquitin proteasome pathway.

D) Model for DUX4-induced post-transcriptional gene regulation.

Supplementary Figures.
438

Figure S1. Demonstration of peptide to protein summarization for two candidate genes.

A) The mass spectra for a representative peptide from a heavy-labeled sample.

B) Histogram of log2 transformed heavy to light ratio of peptides from cells labeled with heavy

amino acids for 1 week.

C) Histogram of log2 transformed heavy to light ratio of peptides from an equal mixture of

unlabeled cells with cells labeled with heavy amino acids for 1 week.

D) Histogram shows the DUX4/Control ratios (log2) for all peptides in the iDUX4 dataset (top

panel), peptides mapping to RPL15 (middle panel) or ZSCAN4 gene (bottom panel)

E) Box and whisker plot showing the median log2 DUX4/control ratio for RPL15 and ZSCAN4.

Figure S2. Tool for easy access of the data generated in this study

Screenshot of the Shiny web server showing a sample analysis of the RNA and protein-level data

for a DUX4 transcriptional target, ZSCAN4.

Supplementary Data

Supplementary data 1: PSM-level data (deposited to Dryad; Link: https://datadryad.org/review?doi=doi:10.5061/dryad.ck06k75)

Supplementary data 2: Gene level data for RNA and protein fold change

Supplementary file 3: R markdown document with the code needed to process mass spectrometry data and generate figures.

Acknowledgements

The authors wish to acknowledge Heidi Dvinge for helpful discussions on proteomics data analysis. We also thank current and past members of the Bradley and Tapscott laboratories for comments and suggestions that have improved this body of work over time. This research was supported by the NINDS P01NS069539 (RKB and SJT), and the FSH Society FSHS-22014-01 (SJ). RKB is a Scholar of The Leukemia & Lymphoma Society. The Fred Hutchinson Cancer Research Center Proteomics Facility is supported by a National Institutes of Health Cancer Center Support Grant (P30 CA015704). The OrbiTrap Fusion mass spectrometer used in this research was purchased with a grant from the M.J. Murdock Charitable Trust.

476 **Competing interests**
477
478 No competing interests declared.

References

1. Tawil R, van der Maarel SM, & Tapscott SJ (2014) Facioscapulohumeral dystrophy: the path to consensus on pathophysiology. *Skelet Muscle* 4:12.

2. Lemmers RJ*, et al.* (2010) A unifying genetic model for facioscapulohumeral muscular dystrophy. *Science* 329(5999):1650-1653.

3. Geng LN*, et al.* (2012) DUX4 activates germline genes, retroelements, and immune mediators: implications for facioscapulohumeral dystrophy. *Dev Cell* 22(1):38-51.

4. Block GJ*, et al.* (2013) Wnt/beta-catenin signaling suppresses DUX4 expression and prevents apoptosis of FSHD muscle cells. *Hum Mol Genet* 22(23):4661-4672.

5. Young JM*, et al.* (2013) DUX4 binding to retroelements creates promoters that are active in FSHD muscle and testis. *PLoS Genet* 9(11):e1003947.

6. Banerji CR*, et al.* (2015) beta-Catenin is central to DUX4-driven network rewiring in facioscapulohumeral muscular dystrophy. *J R Soc Interface* 12(102):20140797.

7. Feng Q*, et al.* (2015) A feedback loop between nonsense-mediated decay and the retrogene DUX4 in facioscapulohumeral muscular dystrophy. *Elife* 4.

8. Homma S, Beermann ML, Boyce FM, & Miller JB (2015) Expression of FSHD-related DUX4-FL alters proteostasis and induces TDP-43 aggregation. *Ann Clin Transl Neurol* 2(2):151-166.

9. Dmitriev P*, et al.* (2016) DUX4-induced constitutive DNA damage and oxidative stress contribute to aberrant differentiation of myoblasts from FSHD patients. *Free Radic Biol Med* 99:244-258.

10. Shadle SC*, et al.* (2017) DUX4-induced dsRNA and MYC mRNA stabilization activate apoptotic pathways in human cell models of facioscapulohumeral dystrophy. *PLoS Genet* 13(3):e1006658.

11. Campbell AE, Belleville A, Resnick R, Shadle SC, & Tapscott SJ (2018) Facioscapulohumeral dystrophy: Activating an early embryonic transcriptional program in human skeletal muscle. *Hum Mol Genet*.

12. Lek A, Rahimov F, Jones PL, & Kunkel LM (2015) Emerging preclinical animal models for FSHD. *Trends Mol Med* 21(5):295-306.

13. Tassin A*, et al.* (2013) DUX4 expression in FSHD muscle cells: how could such a rare protein cause a myopathy? *J Cell Mol Med* 17(1):76-89.

14. Rickard AM, Petek LM, & Miller DG (2015) Endogenous DUX4 expression in FSHD myotubes is sufficient to cause cell death and disrupts RNA splicing and cell migration pathways. *Hum Mol Genet* 24(20):5901-5914.

15. Knopp P*, et al.* (2016) DUX4 induces a transcriptome more characteristic of a less-differentiated cell state and inhibits myogenesis. *J Cell Sci* 129(20):3816-3831.

16. Schwanhausser B*, et al.* (2011) Global quantification of mammalian gene expression control. *Nature* 473(7347):337-342.

17. Tassin A*, et al.* (2012) FSHD myotubes with different phenotypes exhibit distinct proteomes. *PLoS One* 7(12):e51865.

18. Celegato B*, et al.* (2006) Parallel protein and transcript profiles of FSHD patient muscles correlate to the D4Z4 arrangement and reveal a common impairment of slow to fast fibre differentiation and a general deregulation of MyoD-dependent genes. *Proteomics* 6(19):5303-5321.

19. Laoudj-Chenivesse D*, et al.* (2005) Increased levels of adenine nucleotide translocator 1 protein and response to oxidative stress are early events in facioscapulohumeral muscular dystrophy muscle. *J Mol Med (Berl)* 83(3):216-224.

20. Jagannathan S*, et al.* (2016) Model systems of DUX4 expression recapitulate the transcriptional profile of FSHD cells. *Hum Mol Genet* 25(20):4419-4431.

21. Harsha HC, Molina H, & Pandey A (2008) Quantitative proteomics using stable isotope
labeling with amino acids in cell culture. *Nat Protoc* 3(3):505-516.
22. Yao Z*, et al.* (2014) DUX4-induced gene expression is the major molecular signature in
FSHD skeletal muscle. *Hum Mol Genet* 23(20):5342-5352.
23. Rodriguez-Boulan E & Musch A (2005) Protein sorting in the Golgi complex: shifting
paradigms. *Biochim Biophys Acta* 1744(3):455-464.
24. Bexiga MG & Simpson JC (2013) Human diseases associated with form and function of
the Golgi complex. *Int J Mol Sci* 14(9):18670-18681.
25. Claudio N, Dalet A, Gatti E, & Pierre P (2013) Mapping the crossroads of immune
activation and cellular stress response pathways. *EMBO J* 32(9):1214-1224.
26. Oslowski CM & Urano F (2011) Measuring ER stress and the unfolded protein response
using mammalian tissue culture system. *Methods Enzymol* 490:71-92.
27. Palacios IM (2013) Nonsense-mediated mRNA decay: from mechanistic insights to
impacts on human health. *Brief Funct Genomics* 12(1):25-36.
28. Licklider LJ, Thoreen CC, Peng J, & Gygi SP (2002) Automation of nanoscale
microcapillary liquid chromatography-tandem mass spectrometry with a vented column.
Anal Chem 74(13):3076-3083.
29. Kall L, Canterbury JD, Weston J, Noble WS, & MacCoss MJ (2007) Semi-supervised
learning for peptide identification from shotgun proteomics datasets. *Nat Methods*
4(11):923-925.
30. Park SS*, et al.* (2012) Effective correction of experimental errors in quantitative proteomics
using stable isotope labeling by amino acids in cell culture (SILAC). *J Proteomics*
75(12):3720-3732.
31. Ong SE & Mann M (2006) A practical recipe for stable isotope labeling by amino acids in
cell culture (SILAC). *Nat Protoc* 1(6):2650-2660.
555

Figure 1

Figure 2

A

Criteria	Gene number		
	RNA	Protein	Both
Measured	21184	4005	3961
log2 fold change > 2	107	132	54
log2 fold change < -2	36	76	3

B

r = 0.51 (p-value < 2.2e-16)

C

D

E

Figure 3

A

n = 678

Protein log2 Fold change

RNA log2 Fold change

B

HSP90AA1

iDUX4: RNA
iDUX4: Protein
vDUX4: RNA
vDUX4: Protein (24h)
vDUX4: Protein (36h)

DUX4/Control ratio (log2)

HSPA5

DUX4/Control ratio (log2)

DDX5

iDUX4: RNA
iDUX4: Protein
vDUX4: RNA
vDUX4: Protein (24h)
vDUX4: Protein (36h)

DUX4/Control ratio (log2)

DHX9

DUX4/Control ratio (log2)

C

HSPA5 mRNA

Normalized fold change

UPR

Control
DUX4

D

Phosphorylated eIF2a

Control

DUX4

0 8 12 16 18 20 22 24 28 32 36

E

S35-Met incorporation (DUX4/Ctrl)

Time (in hours)

DUX4 expression

Figure 4

Figure S1

A

Peptide: VEIIANDQGNR

H = 8.25E6

L = 1.33E5

Relative Abundance

m/z

B

Frequency

log2 (heavy/light)

C

Frequency

log2 (heavy/light)

D

All genes

Peptide Number

RPL15

Peptide Number

ZSCAN4

Peptide Number

Normalized log2(DUX4/Control)

E

log2 (DUX4/Ctrl)

RPL15   ZSCAN4

Figure S2