The key protein of endosomal mRNP transport Rrm4

binds translational landmark sites of cargo mRNAs

Lilli Olgeiser[1], Carl Haag[1], Susan Boerner[2], Jernej Ule[3,4], Anke Busch[5], Janine Koepke[6],

Julian König[5], Michael Feldbrügge[1, $] and Kathi Zarnack[2, $]

$ shared corresponding authorship

ORCID IDs: M. Feldbrügge 0000-0003-0046-983X and K. Zarnack 0000-0003-3527-3378

[1] Heinrich Heine University Düsseldorf, Institute for Microbiology, Cluster of Excellence on Plant Sciences, 40204 Düsseldorf, Germany
[2] Buchmann Institute for Molecular Life Sciences (BMLS), Goethe University Frankfurt, Max-von-Laue-Str. 15, 60438 Frankfurt am Main, Germany
[3] The Francis Crick Institute, Midland Road 1, Kings Cross, London NW1 1AT, UK
[4] Department of Molecular Neuroscience, UCL Institute of Neurology, Queen Square, London, WC1N3BG, UK
[5] Institute of Molecular Biology gGmbH, Ackermannweg 4, 55128 Mainz, Germany
[6] Medical Clinic II (Molecular Pneumology), Justus Liebig University of Gießen, Excellence Cluster Cardio-Pulmonary System, Aulweg 130, 35392 Gießen, Germany

Running title: Endosomal mRNP transport

Key words: endosome / fungus / iCLIP / microtubule / RNA-binding protein / RNA transport

[*] **Corresponding authors:**

Dr. Michael Feldbrügge

Institute for Microbiology, Cluster of Excellence on Plant Sciences

Heinrich Heine University Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf, Germany

Phone: +49 (211) 81-15475

feldbrue@hhu.de

Dr. Kathi Zarnack

Buchmann Institute for Molecular Life Sciences (BMLS)

Goethe University Frankfurt, Max-von-Laue-Str. 15, 60438 Frankfurt am Main, Germany

Phone: + 49 (69) 798-42506

kathi.zarnack@bmls.de

Abstract

RNA-binding proteins (RBPs) determine spatiotemporal gene expression by mediating active

transport and local translation of cargo mRNAs. Here, we cast a transcriptome-wide view on

the transported mRNAs and cognate RBP binding sites during endosomal messenger

ribonucleoprotein (mRNP) transport in *Ustilago maydis*. Using individual-nucleotide

resolution UV crosslinking and immunoprecipitation (iCLIP), we compare the key transport

RBP Rrm4 and the newly identified endosomal mRNP component Grp1 that is crucial to

coordinate hyphal growth. Both RBPs bind predominantly in the 3´ untranslated region of

thousands of shared cargo mRNAs, often in close proximity. Intriguingly, Rrm4 precisely

binds at stop codons, which constitute landmark sites of translation, suggesting an intimate

connection of mRNA transport and translation. Towards uncovering the code of recognition,

we identify UAUG as specific binding motif of Rrm4 that is bound by its third RRM domain.

Altogether, we provide first insights into the positional organisation of co-localising RBPs on

individual cargo mRNAs.

Introduction

All eukaryotic cells must accurately regulate the expression of proteins in time and space. To

this end, many mRNAs accumulate at specific subcellular sites, and their local translation is

exactly timed [1-3]. mRNA localisation is achieved most commonly by active motor-

dependent transport along the cytoskeleton. Functional transport units are messenger

ribonucleoprotein complexes (mRNPs), consisting of various RNA-binding proteins (RBPs),

accessory proteins and cargo mRNAs. Key factors are RBPs that recognise localisation

elements (LEs) within mRNAs. For movement, the RBPs either interact with motors directly

or are connected via linker proteins [1, 4].

We discovered co-transport of mRNPs on the cytoplasmic surface of early endosomes as

a novel translocation mechanism of cargo mRNAs during hyphal growth in fungi [5, 6]. These

endosomes shuttle along microtubules by the concerted action of plus-end directed kinesin

and minus-end directed dynein [7, 8]. They serve as multipurpose platforms functioning not

only during endocytic recycling but also during long-distance transport of whole organelles

such as peroxisomes [9-12].

Endosomal mRNA transport was uncovered analysing the RBP Rrm4 in the dimorphic

phytopathogenic fungus *Ustilago maydis* (Fig EV1A) [13, 14]. Loss of Rrm4 has no effects

on the yeast form of the fungus. However, the absence of Rrm4 causes characteristic defects

in unipolar growth when switching to the hyphal form: the fraction of bipolarly growing

hyphae increases and the insertion of basal septa is delayed [6, 15]. In line with endosomal

mRNA transport, Rrm4 binds mRNAs and shuttles on early endosomes along microtubules *in*

vivo [5, 16]. Using the poly(A)-binding protein Pab1 as an mRNA marker revealed that loss

of Rrm4 abolishes this transport, resulting in a gradient of mRNAs declining from the nucleus

towards the cell periphery [17]. Thus, one function of Rrm4 might be the general distribution

of mRNAs within hyphae [17].

Initial CLIP experiments with Rrm4 identified target mRNAs encoding chitinase Cts1

and septin Cdc3, among others [17, 18]. The subcellular localisation of both proteins was

Rrm4-dependent: Loss of Rrm4 strongly reduced the secretion of the chitinase Cts1.

Moreover, shuttling of the Cdc3 protein on early endosomes was abolished, and the gradient

of septin filaments at the growth pole of hyphae was no longer formed [6]. Since *cdc3* mRNA

and its encoded protein are found together with ribosomes on the same shuttling endosomes,

we hypothesised that endosome-coupled translation of *cdc3* mRNA during long-distance

transport is critical for the efficient formation of septin filaments at the growth pole [6]. This

was supported by demonstrating that all four septin-encoding mRNAs are present on

endosomes and that septin proteins assemble into heteromeric complexes on the cytoplasmic

face of endosomes during long-distance transport [19]. Thus, Rrm4-dependent mRNA

transport regulates the specific localisation of the corresponding translation products. To

understand this complex process at the transcriptome-wide level, we present herein an *in vivo*

snapshot of RNA binding sites of endosomal RBPs on cargo mRNAs at single-nucleotide

resolution.

Results

Loss of the glycine/glutamine-rich protein Grp1 affects hyphal growth

In order to identify additional protein components involved in endosomal mRNA transport,

we performed pilot affinity tag purification using Rrm4 as bait. We identified the potential

RBP glycine-rich protein 1 (Grp1; UMAG_02412), which carries an N-terminal RNA

recognition motif (RRM) domain followed by a short C-terminal region rich in glycine and

glutamine (GQ-rich; Fig 1A, Fig EV1B-D). The protein was similar to other small RRM

proteins, such as human CIRBP or RBM3 and plant RBG7 (*At*GRP7), all previously

described as global stress regulators (Fig 1A) [20, 21].

For functional analysis, we generated deletion mutants in laboratory strain AB33. In this

strain the master transcription factor controlling hyphal growth is under control of an

inducible promoter. Thus, hyphal growth can be elicited synchronously by changing the

nitrogen source in the medium. The corresponding hyphae grow like wild type by tip

expansion at the apical pole, while the nucleus is positioned in the centre and septa are

inserted in regular intervals at the basal pole (Fig EV 1A) [22]. In the yeast form of AB33, we

observed that loss of Grp1 resulted in slower proliferation as well as increased cell size (Fig

EV1E-G). At lower temperatures, growth of the *grp1Δ* strain was affected even more strongly

and it exhibited an altered colony morphology (Fig EV1H). This was consistent with a

potential function in cold stress response, similar to the plant and human orthologues [20, 21].

Furthermore, colony growth of the *grp1Δ* strain was strongly reduced upon treatment with

inhibitors of cell wall biosynthesis, such as Calcofluor White (CFW) or Congo Red (CR) [23,

24]. Hence, loss of Grp1 might cause defects in cell wall formation (Fig EV1I).

Studying hyphal growth revealed that, unlike observed in *rrm4Δ* strains, loss of Grp1 did

not cause an increased amount of bipolar cells as it is characteristic for defects in microtubule-

dependent transport (see *rrm4Δ* hyphae for comparison; Fig 1B-C) [12]. On the contrary,

under optimal growth conditions hyphae were significantly longer (Fig 1B-C), and the length

of empty sections at the basal pole was increased (Fig 1D; Fig EV1J). Hence, the coordination

of hyphal growth may be disturbed in the absence of Grp1. In order to further investigate this,

we stressed hyphae three hours post induction (h.p.i.) of hyphal growth with the cell wall

inhibitor CFW. In comparison to wild type hyphae, we observed a strongly increased number

of *grp1Δ* hyphae with abnormal shapes (86%), indicating that cell wall integrity might be

affected (Fig 1E-F).

In summary, loss of Grp1 affects both yeast-like and hyphal growth. During the latter,

Grp1 seems to be crucial for the correct coordination of cell wall expansion, which becomes

particularly apparent during stress conditions.

Grp1 is a novel component of endosomal mRNA transport

To analyse the subcellular localisation of Grp1, we generated AB33 strains expressing Grp1

fused at its C-terminus to Gfp by homologous recombination. The functional Grp1-Gfp

version accumulated in the cytoplasm as well as in the nucleus of hyphae. In comparison, the

poly(A)-binding protein Pab1-Gfp was absent from the nucleus, suggesting that this

localisation pattern is specific for Grp1 (Fig 2A). Importantly, a subpopulation of Grp1-Gfp

moved bi-directionally in the cytoplasm with a velocity comparable to Rrm4-Gfp and Pab1-

Gfp, which are known to shuttle on early endosomes (Fig 2A-B; Supplemental Video 1).

To test whether Grp1 shuttles on Rrm4-positive endosomes, we performed dynamic co-

localisation studies using dual-view technology [25]. We generated AB33 strains co-

expressing Grp1-Gfp and Rrm4 fused C-terminally to the red fluorescent protein Tag-Rfp

(tRfp) [26]. For comparison, we used a strain expressing Pab1 fused to the red fluorescent

protein mCherry [17, 27]. Analysing hyphae 6 h.p.i. 99% of processive Grp1-Gfp signals co-

migrated with Rrm4-tRfp, revealing extensive co-localisation of both proteins in shuttling

units (Fig 2C-D; Supplemental Video 2). Consistently, 97% of processive Grp1-Gfp signals

co-migrated with Pab1-mCherry, indicating that Grp1, like Pab1, was present on Rrm4-

positive endosomes (Fig 2C-D; Supplemental Videos 3-4). Thus, Grp1 appears to be a novel

component of endosomal mRNPs that might already be recruited to transport mRNPs in the

nucleus.

The endosomal localisation of Grp1 depends on Rrm4

To investigate whether Grp1 has an influence on the shuttling of Rrm4-positive endosomes,

we studied Rrm4 movement in *grp1Δ* strains. Loss of Grp1 altered neither processive Rrm4-

Gfp movement nor the velocity of the respective endosomes (Fig 2E-F; Supplemental Video

5). Vice versa, studying Grp1-Gfp movement in the absence of Rrm4 revealed that its

endosomal localisation depended on Rrm4 (Fig 2G). Importantly, similar to Pab1-Gfp, a

gradient of Grp1-Gfp was formed in *rrm4Δ* hyphae, with a decreasing signal intensity towards

the growing apex (Fig 2H-I) [17]. Similar to Pab1, which is expected to associate with almost

all poly(A) tails of mRNAs [28], Grp1 might therefore be distributed in association with

many mRNAs (see below).

To test whether Grp1-Gfp binds to endosomes in an mRNA-dependent manner, we

generated AB33 strains expressing Rrm4^{mR123}-tRfp. This Rrm4 variant carried point

mutations in the RNP1 regions of RRM domains 1-3 causing a reduced RNA binding activity

and loss of function of Rrm4 [16]. In dual-view experiments we observed that Grp1-Gfp like

Pab1-Gfp no longer shuttled in the presence of Rrm4^{mR123}-tRfp (Fig 2J). Thus, the localisation

of Grp1 depends on the presence of functional Rrm4, more precisely on its capability to bind

RNA. In summary, we identified Grp1 as a novel component of endosomal mRNPs whose

shuttling on Rrm4-positive endosomes depended on Rrm4 and mRNA.

Rrm4 and Grp1 share thousands of target transcripts

In order to learn more about the function of the two endosomal mRNP components Rrm4 and

Grp1 during hyphal growth, we performed a comparative transcriptome-wide analysis of their

RNA binding behaviour using individual-nucleotide resolution UV crosslinking and

immunoprecipitation (iCLIP) [29]. For application with fungal RBPs, we had to modify a

number of steps in the iCLIP protocol (Fig EV2A-D; see Materials and methods) [30]. One

major challenge was the high RNase and protease activity in fungal cell extracts that resulted

in a low yield of crosslinked protein-RNA complexes and short mRNA fragments. The most

critical changes to the protocol came with the fast processing of crosslinked material and the

identification of the optimal UV-C irradiation dose (Fig EV2B).

139 Using the improved protocol, we found that Rrm4-Gfp and Grp1-Gfp displayed

140 substantial crosslinking to RNA *in vivo* (compared to Gfp control; Fig 3A). As expected, the

141 RNA signal was dependent on UV-C irradiation and sensitive to RNase I digestion. Upon

142 iCLIP library preparation, we obtained more than 100 million sequencing reads,

143 corresponding to 4.7 x 10^6 and 14.8 x 10^6 crosslink events for Rrm4 and Grp1, respectively

144 (Fig EV3A-B). Reproducibility between two replicate experiments was high for both proteins,

145 demonstrating the quality of the obtained data set (Pearson correlation coefficient > 0.96, p

146 value < 2.22e-16; Fig EV3C).

147 Consistent with the abundance of both proteins, the crosslink events accumulated into

148 thousands of clusters that spread across major parts of the transcriptome (Fig EV3A). In order

149 to focus on the most prominent sites, we used the crosslink frequency within each cluster

150 relative to the background signal within the same transcript to determine the 25% most

151 prominent binding sites for Rrm4 and Grp1 ('signal-over-background'; see Materials and

152 methods). This procedure identified a total of 6,412 binding sites for Rrm4 and 6,478 binding

153 sites for Grp1, residing in 3,262 and 3,165 target transcripts, respectively (Fig 3B). This

154 represented a substantial fraction of the about 6,700 protein-encoding genes in the *U. maydis*

155 genome [31]. Extensive endosomal transport of mRNA is consistent with a role in evenly

156 distributing mRNAs throughout hyphae (see Discussion).

157 Comparing Rrm4 and Grp1 revealed a large overlap of 2,114 target transcripts that were

158 conjointly bound by both proteins (Fig 3B-C, Dataset EV1; see below). In this shared target

159 set, we observed an enrichment for functional categories like mitochondrion, vesicle transport

160 and cytoskeleton (Fig 3D). Moreover, we found several known Rrm4 target transcripts,

161 including for instance all four *septin* mRNAs (Fig 3C, 3E). Binding sites of Rrm4 and Grp1 in

162 the *septin* mRNAs were almost exclusively located in the 3´ untranslated region (UTR),

163 consistent with the hypothesis that these mRNAs are transported in a translationally active

state (see Discussion) [19]. *cts1* mRNA, another known target of the Rrm4 transport

machinery [18], also carried binding sites of both RBPs in the 3´ UTR (Fig EV3D).

To assess the function of target mRNAs that are specifically recognised by only one of

the two RBPs, we applied more stringent criteria to define 280 and 520 transcripts that were

uniquely bound by Rrm4 and Grp1, respectively (Fig EV3E, Dataset EV2-3). While the

Rrm4-unique set displayed no clear trend, the Grp1-unique set showed an enrichment for

mRNAs encoding nuclear proteins that were involved in transcriptional regulation and

chromatin remodelling (Fig 3D). Although these mRNAs were expressed and bound by Grp1,

they were most likely not transported by the Rrm4 machinery, which we presume would result

in perinuclear localisation. This could facilitate an efficient nuclear import of the translation

products, as being described in mammalian cells for transcription factors like c-Myc or for

metallothionin [32, 33].

In summary, the comparative iCLIP approach revealed that Grp1 and Rrm4 conjointly

bind thousands of shared target mRNAs. These offer a comprehensive view on the full

spectrum of cargo mRNAs transported by the endosomal mRNP transport machinery in *U.*

maydis.

Rrm4 binds to functionally important sites of target transcripts

Studying the distribution of binding sites in different transcript regions revealed that Rrm4

and Grp1 preferentially bound in the 3´ UTR (Fig 4A). Within this region, both proteins

frequently bound in close proximity, with 51% of Rrm4 binding sites directly overlapping

with a Grp1 binding site (compared to only 5% in the open reading frame, ORF; Fig 4B-C).

Thus, the cargo mRNAs of the transport mRNPs are often conjointly recognised by both

RBPs in the 3´ UTR.

In contrast to Grp1 that was almost exclusively attached to the 3´ UTR, Rrm4 bound a

substantial fraction of target mRNAs within the ORF (1,315 mRNAs with 1,783 ORF binding

sites; Fig 4A, 4C). Taking a closer look at the binding pattern of Rrm4 along ORFs, we

observed binding sites at the start and stop codons of a subset of target mRNAs, reflected in

increased crosslinking of Rrm4 at these sites (Fig 4D). Whereas only two transcripts

harboured a Grp1 binding site at the start codon, Rrm4 binding sites overlapped the start

codon in 47 target mRNAs, like in the transcript encoding the translation initiation factor Sui1

(UMAG_02665; Fig EV3D; Dataset EV4). Of note, the *rrm4* mRNA itself exhibited Rrm4

binding sites around the start codon, hinting at a potential autoregulation (Fig 4E).

Even more prominently than at start codons, we observed a strong accumulation of Rrm4

binding sites at the stop codons of multiple target transcripts (291 cases; Fig 4D-E; Dataset

EV5). These included, for example, both subunits of cytoplasmic dynein (Dyn1 and Dyn2)

[34]. Furthermore, the stop codon-bound targets were significantly enriched for mRNAs

encoding mitochondrial proteins, including for instance the majority of nucleus-encoded

subunits and accessory components of the F_oF_1-ATPase (Fig EV4A-C). Through binding at

the stop codon, Rrm4 might influence translation termination for these targets. This

hypothesis was supported by revisiting data from a previous proteomics approach analysing

the membrane-associated proteome of wild type and *rrm4Δ* hyphae, in which the amount of

the membrane-associated component Atp4 was threefold reduced in *rrm4Δ* hyphae compared

to wild type (Fig EV4D) [18].

In essence, the high-resolution mapping of binding sites for the two endosomal RBPs

Rrm4 and Grp1 revealed (i) that both proteins conjointly bind in the 3´ UTR, and (ii) that

Rrm4 additionally recognises binding sites in the ORF and at start and stop codons.

Rrm4 specifically recognises the motif UAUG via its third RRM

In order to address the RNA sequence specificity of both RBPs, we used the motif analysis

algorithm DREME [35] to search for sequence motifs in a 30-nt window around the RBP

binding sites. This analysis retrieved UAUG as the most significantly enriched sequence motif

at Rrm4 binding sites (Fig 5A). Analysing the relative positioning of the motif showed that

more than one third of all Rrm4 binding sites harboured a UAUG motif precisely at the centre

of the binding site (2,201 out of 6,412; Fig 5B). The motif did not accumulate at Grp1 binding

sites, supporting the notion that the motif was specifically recognised by Rrm4. In order to

estimate the relative strength of Rrm4 binding, we calculated the 'signal-over-background'

(SOB), i. e the ratio of crosslink events within the binding site over the background of

crosslink events in the surrounding sequence (see Materials and methods). The background

binding served as a proxy for the abundance of underlying transcript. Since the SOB

procedure did not correct for UV crosslinking biases and similar confounding factors,

comparisons between binding sites were only performed at a global scale. We observed that

the Rrm4 binding sites with UAUG showed stronger relative binding than those lacking the

motif (Fig 5C), suggesting a tight interaction of Rrm4 with the UAUG-associated binding

sites.

A similar sequence analysis of the Grp1 binding sites initially suggested the sequence

UGUA as a potential recognition motif (Fig EV5A). However, the same motif also frequently

occurred at Rrm4 binding sites and showed no clear positioning relative to the Grp1 binding

sites (Fig EV5B), making it questionable whether it was directly involved in the RNA

recognition of Grp1. We therefore did not pursue this motif further.

In order to independently test whether Rrm4 specifically recognises the sequence motif

UAUG, we applied the yeast three-hybrid assay (Fig 5D). We previously used this approach to

successfully identify SELEX-derived RNA aptamers that were recognised by the third RRM

domain of Rrm4 *in vivo* (RRM3) [36]. Intriguingly, two RNA aptamers, SELEX-A1 and

SELEX-B1, contained the UAUG motif. We chose SELEX-A1 (Fig 5E) [36] to mutate the

UAUG motif and tested RNA binding using the yeast three-hybrid assay. In contrast to the

initial SELEX-A1 aptamer, the mutant version was no longer recognised by Rrm4 (Fig 5F;

Fig EV5C). Consistent with earlier results, mutating the third RRM domain of Rrm4 gave

similar results in the context of the initial SELEX-A1 aptamer [36]. Thus, our computational

and experimental analyses indicate that Rrm4 specifically recognises the sequence motif

UAUG via its third RRM domain.

Interestingly, Rrm4 binding sites in the whole ORF region showed a strong enrichment

for the Rrm4 motif UAUG (Fig 5G), such that 61% of all ORF binding sites harboured

UAUG. Since the start codon AUG is contained within this motif, the vast majority of start

codon-associated binding sites exhibited the UAUG motif (88%, Fig EV5D). However, the

frequency of Rrm4 binding was not increased compared to UAUG motifs in the surrounding

sequence, indicating that Rrm4 does not show a particular preference for UAUG motifs in the

context of translational start sites (Fig EV5E). Nevertheless, the Rrm4 recognition motif

intrinsically overlaps with the start codon, suggesting a potential link to translation regulation.

Binding of Rrm4 at the start codon might interfere, for example, with translational initiation

of the bound target mRNAs.

In contrast to the UAUG prevalence in the ORF, stop codons seem to be recognised

differently (UGA overlaps with UAUG i.e. UA<u>UGA</u>; but UAA was the most common stop

codon, 55%, Fig 5H). Since UAUG-containing binding sites showed particularly strong Rrm4

binding (Fig 3C; as example, see *cdc12* in Fig 3E), Rrm4 appears to exhibit a tight association

with the ORF via its third RRM domain. Uniting these observations, we hypothesised that

Rrm4 simultaneously recognised multiple regions of the same cargo mRNAs. In line with this

notion, we found that transcripts with a Rrm4 binding site in the 3´ UTR were significantly

enriched for a second Rrm4 binding site in the ORF (663 out of 1,703 transcripts with at least

two Rrm4 binding sites; p value < 2.22e-16, Fisher's exact test). In 69% of these cases, the

ORF binding site harboured UAUG, and in 56%, the 3´ UTR binding site of Rrm4 overlapped

with a Grp1 binding site. Taken together, these observations would be consistent with a model

that Rrm4 binds with its RRM domains RRM1 and/or RRM2 close to Grp1 in the 3´ UTR and

via its third RRM domain to a UAUG-containing binding site in the ORF (Fig 6).

Discussion

At present, a small number of high-throughput studies provide a global view on transported and localised mRNAs. In oocytes and embryos from fruit fly, transcriptome-wide RNA *in situ* hybridisation approaches revealed that the majority of transcripts exhibit a defined localisation pattern [37, 38]. In neuronal cells, the localised transcriptome and proteome have also been compiled [39, 40]. However, it is still unclear (i) how these cargo mRNAs reach their destination, (ii) which RBPs mediate their transport, and (iii) what are the precise interaction sites within the target mRNAs. Here, we applied iCLIP to study the newly identified endosomal RBP Grp1 and the key transport RBP Rrm4 during endosomal mRNP transport in *U. maydis.* To the best of our knowledge, this is the first detailed iCLIP analysis of mRNA transport.

The GQ-rich RNA-binding protein Grp1 is a novel component of endosomal mRNPs

The small GQ-rich protein Grp1 shares similarity with other glycine-rich RBPs from humans and plants. A characteristic feature of this conserved class of RBPs is an N-terminal RRM domain followed by a glycine-rich low-complexity sequence that is typical for intrinsically disordered regions (IDRs). In RBPs, IDRs can mediate the assembly of membrane-less organelles through phase transitions [41], and this assembly is important, for example, during splicing [42]. Interestingly, IDRs have also been implicated in the formation of RNA granules for neuronal RNA transport [43, 44].

Small glycine-rich RBPs function in a wide variety of biological processes. The plant protein *At*GRP7, for example, is involved in cold stress adaption, osmotic stress response, circadian rhythm and plant immunity [21, 45, 46]. Human CIRBP regulates telomerase activity and the human cold shock protein RBM3 is involved in translational reprogramming during the cooling response of neuronal cells [20, 47, 48]. Globally, these proteins might

function as RNA chaperones that prevent the formation of aberrant RNA secondary structures

under stress conditions [45, 49].

In this study, we observe that loss of the fungal orthologue Grp1 causes aberrant

alterations of the hyphal growth programme. Under optimal conditions *grp1Δ* hyphae grow

significantly longer, suggesting an unusual acceleration of cell wall formation. Moreover, cell

wall stress revealed clear abnormal morphologies in comparison to wild type. Consistent with

a role in hyphal growth, Grp1 shuttles on Rrm4-positive endosomes that are the main

transport units for long-distance transport of mRNAs in hyphae. Moreover, Grp1 and Rrm4

conjointly bind in the 3´ UTRs of thousands of target mRNAs (Fig 6A, see below). We

therefore propose that the potential RNA chaperone Grp1 most likely constitutes an accessory

component of endosomal mRNPs. Its function could be particularly important under

suboptimal conditions. Alternatively, Grp1 might regulate stability and/or translation of

mRNAs encoding proteins involved in hyphal growth independent of endosomal mRNA

transport.

It is of note that the RRM protein Hrb27C (Hrp48) from fruit fly with a related domain

architecture containing two N-terminal RRMs followed by a C-terminal GQ-rich region was

found as an mRNP component during transport of *oskar* and *gurken* mRNAs [50-52]. Hence,

the presence of small glycine-rich RBPs in transport mRNPs might be preserved across

organisms.

Endosomal RBPs recognise a broad spectrum of cargo mRNAs

In order to obtain a comprehensive view on the *in vivo* mRNA targets of an RBP, UV

crosslinking techniques are currently the method of choice [53-55]. Here, we applied iCLIP to

study fungal mRNA transport. A strength of our approach was the use of strains expressing

functional Gfp-tagged versions of Grp1 and Rrm4 using homologous recombination to avoid

overexpression artefacts. For application in fungi, we had to improve several steps [30], of

which optimising the dose and duration of UV-C irradiation was most critical. Thereby, we

were able to obtain a transcriptome-wide view of the cargo mRNAs, including their

interaction sites with cognate RBPs at single nucleotide resolution.

Comparing two distinct RBPs present in endosomal mRNPs enabled us to disentangle

the precise binding behaviour of the two co-localising RBPs. We identified more than 2,000

shared target transcripts, covering a substantial amount of the approximately 6,700 annotated

protein-coding genes (http://pedant.helmholtz-muenchen.de/index.jsp; ORF update

2016.11.08) [31]. The broad target spectrum of endosomal mRNA transport fits with earlier

observations that Rrm4 transported all mRNAs under investigation, albeit with different

processivity of transport [17]. Moreover, loss of Rrm4 impairs the global mRNA distribution

in hyphae, indicated by a disturbed subcellular distribution of the poly(A)-binding protein

Pab1 [17]. Thus, one function of the endosomal mRNA transport machinery might be the

equal distribution of mRNPs to supply all parts of the hyphae with mRNAs. This might be

particularly important for those parts that are distant from the mRNA-synthesising nucleus.

Such a universal mRNA transport mode resembles the "sushi belt model" in neuronal cells, in

which the shuttling of mRNPs by active transport along microtubules is thought to distribute

mRNAs throughout the dendrite to serve synapses that are in demand of mRNAs [56].

Furthermore, since many mRNAs in *U. maydis* appear to be transported in a translationally

active state (see below), this mRNA distributer function would also disclose the mechanism of

how ribosomes are transported by endosomes, as observed previously [57].

Notably, we found that most nuclear-encoded subunits and accessory components of

mitochondrial F_0F_1 ATPase are targets of endosomal mRNP transport in *U. maydis*.

Consistent with the idea that a precise spatiotemporal regulation of translation might be

important for efficient mitochondrial protein import, we observed that the abundance of Atp4

is reduced in *rrm4Δ* hyphae [18]. These results agree with previous findings that the 3´ UTR

of *ATP2* mRNA from *S. cerevisiae* is important for efficient mitochondrial uptake of mature

Atp2p [58]. A close link between RNA biology and mitochondrial protein import is thus an

emerging theme in mitochondrial biology [59-63].

Distinct binding patterns of Rrm4 may allow orchestration of mRNA transport and

translation

The positional information obtained by single nucleotide resolution was essential to uncover

the precise binding behaviour of the involved RBPs. In the majority of cases, Rrm4 binds

together with Grp1 in the 3´ UTR (1,700 transcripts; Fig 6A). The vicinity of these binding

sites to the poly(A) tail fits the previous observation that the endosomal adaptor protein Upa1

interacts with both Rrm4 and Pab1 [64]. While translating ribosomes would potentially

remove RBPs from the ORF [65], such binding in the 3´ UTR, as seen e.g. on all four *septin*

mRNAs, would allow simultaneous translation and transport of mRNAs. Consistently, we

have recently provided evidence that endosome-coupled translation of *septin* mRNAs

mediates endosomal assembly and transport of heteromeric septin complexes [6, 19]. Since

transport of translationally active mRNAs has recently been observed in neurons [66], this

mode of transport might be more widespread than currently anticipated.

Our transcriptome-wide RNA binding maps illustrate an intriguing binding pattern of

Rrm4 at translational landmark sites, indicating an intimate connection of endosomal mRNP

transport and translation in *U. maydis* (Fig 6A): First, in over 1,000 target transcripts Rrm4

binds within the ORF, indicating that it may be involved in stalling translational elongation.

Notably, a similar mechanism was suggested in neurons, in which a subset of mRNAs are

translationally stalled during transport [67]. Similar to Rrm4, the neuronal RBP FMR1 binds

its translationally stalled target mRNAs preferably in the coding sequence [68]. Second, in

about 50 and 300 target transcripts, respectively, Rrm4 precisely binds at the start and stop

codons, suggesting modulation of translation initiation and termination. At start codons and

within the ORF, the Rrm4 binding sites frequently harboured UAUG. This motif is recognised

by the third RRM domain of Rrm4. In accordance with its ELAV-type domain organisation,

we therefore propose that Rrm4 binds UAUG-containing binding sites via its third RRM to

influence translation (Fig 6B), while the two tandem RRMs (RRM1/2) bind the target

mRNAs in the 3´ UTR, possibly together with Grp1 (Fig 6B). In a previous study, we

observed that mutations in RRM1 and RRM3 led to strongly reduced overall RNA binding of

Rrm4, although mutations in RRM3 did not show a mutant phenotype with respect to hyphal

growth [16]. In contrast, mutations in RRM1 affected hyphal growth strongly [16],

accompanied by reduced endosomal shuttling of Pab1 as well as *cdc3* mRNA [6]. Therefore,

the potential translational regulation during endosomal mRNP transport mediated by RRM3

may be an additive and used for fine-tuning. The tandem RRM domains RRM1 and RRM2 of

Rrm4 might mediate the recognition of target mRNAs for transport, possibly via binding in

the 3´ UTR.

In essence, our comprehensive transcriptome-wide view on endosomal mRNA transport

revealed the precise positional deposition of co-localising RBPs during transport. The key

RBP Rrm4 exhibits a particular binding behaviour by recognising distinct landmarks of

translation in target mRNAs. Thereby, translation and transport might be intimately coupled

and precisely synchronised for the specific expression needs of each target transcript.

Materials and methods

Plasmids, strains and growth conditions

Cloning was done using *E. coli* K-12 derivate Top10 (Life Technologies, Carlsbad, CA,

USA). All strains were constructed by the transformation of cells with linearised plasmids,

and homologous integration events were verified by Southern blot analysis [69]. Genomic

DNA of wild type strain UM521 (*a1b1*) served as template for PCR amplifications unless

noted otherwise. Proteins were tagged with eGfp (enhanced green fluorescent protein;

Clontech, Mountain View, CA, USA), Tag-Rfp or mCherry [26, 70]. Plasmid sequences are

available upon request. The accession numbers of *U. maydis* genes used in this study are:

rrm4 (UMAG_10836), *grp1* (UMAG_02412), *pab1* (UMAG_03494). Detailed information is

supplied in Appendix Tables S1-S3. The conditions used for cultivation of *U. maydis* are

described elsewhere [69]. For the induction of filamentous growth of lab strain AB33 and

derivates, cultures were grown to an OD_{600} of 0.5 in complete medium supplemented with 1%

glucose (CM-glc) and shifted to nitrate minimal medium supplemented with 1% glc (NM-

glc). The growth curves of *U. maydis* strains were recorded by cultivating strains in CM-glc at

28°C and measuring the OD_{600} every 2 h.

Multiple sequence alignments

Orthologous proteins were identified using BLAST P (https://blast.ncbi.nlm.nih.gov/blast/).

Clustal W and GeneDoc 2.6 were used for multiple amino acid sequence alignment and

graphical representation, respectively [71].

Preliminary affinity purification experiments using Rrm4-GfpTT as bait

Preliminary purification experiments were performed using a strain expressing Rrm4 fused C-

terminally to Gfp and tandem affinity purification (tap) tag [16]. We analysed the eluted

fractions by mass spectrometry either as complex mixture in solution (Fig EV1C) or by

cutting individual bands from an SDS-PAGE gel (Fig EV1D). In both cases, the protein

purification followed a published protocol (CLIP) [16]. Both experiments were considered to

be pilot studies and were only performed once. In case of the in-solution digest, the protein

samples were precipitated with 20% trichloroacetic acid (1:2, v/v) and washed five times with

cold acetone. Samples were digested with sequencing-grade modified trypsin (Promega), and

the resulting peptides were separated into fractions by nanoLC (PepMap100 C-18 RP

nanocolumn and UltiMate 3000 liquid chromatography system, Dionex). Fractions were

spotted with matrix solution on a MALDI plate and subsequently analysed by MALDI-TOF

MS (4800 Proteomics Analyser, AB Sciex). Mass spectrometry data were searched against the

MUMDB (Munich *Ustilago Maydis* Data Base; 6785 entries; IBIS Institute of Bioinformatics

and Systems, German Research Center for Environmental Health) [31] using Mascot

embedded into GPS Explorer software (AB Sciex). The total peptide score is the sum of all

peptide scores corresponding to the predicted proteins, excluding the scores of duplicate

matches. The best peptide score is the best score from all identified peptides corresponding to

the predicted proteins.

Applying as filter a total peptide score of larger than 46, we identified 23 proteins as potential

interaction partners of Rrm4 (Fig EV1C). Among these proteins were interesting candidates

that were identified before in independent approaches: Chitinase Cts1 and ribosomal protein

Rps19 were found in a differential proteomics study, comparing wild type and *rrm4Δ* hyphae

[18]. Upa2 is a PAM2 motif-containing protein that was identified as potential interaction

partner of Rrm4 in a bioinformatics approach [64]. Grp1 (UMAG_02412) was chosen for

further analysis, since it contained an RNA recognition motif (RRM) for RNA binding.

In the gel-based approach, the calmodulin beads were washed three times and incubated in

SDS-loading buffer for 8 min at 90°C. The supernatant was subjected to SDS-PAGE (10%

polyacryl amide). Proteins were staining with Coomassie Blue. After excision of selected

bands samples were destained and dried under vacuum. The dried material was suspended in

30 µl 10 mM ammonium bicarbonate (pH 8.3) containing 0.6 µg of trypsin (Promega) and

10% acetonitrile. After incubation at room temperature for 12 h, the trypsin digest was

extracted and analysed as described above. We were able to verify Rrm4-GfpTT (TEV

cleavage product 117 kDa) and Grp1 (18 kDa; Fig EV1D) in selected bands matching their

expected gel position after electrophoresis.

Colony morphology, temperature stress and cell wall stress

For growth on solid media, cell suspensions (OD_{600} of 0.5) were inoculated onto respective

plates at 28°C. Colony morphology was tested on CM (1% glc) medium. Temperature stress

was analysed on CM (1% glc) medium, incubated at 28°C, 20°C or 16°C. Cell-wall stress was

analysed by supplementing CM (1% glc) medium with 50 µM Calcofluor White (Sigma-

Aldrich, Taufkirchen, Germany) or 40 µg/ml Congo Red (Sigma-Aldrich). All plates were

kept for at least 48 h at the required temperature. The set-up used to acquire images was

described before [64]. To analyse cell wall stress in hyphae, 2.5 µM Calcofluor White was

added to cultures 3 hours post induction (h.p.i) and incubated for an additional 2 h.

Microscopy and image processing

The microscope set-ups and dual-colour imaging were used as described before [5, 25, 72].

Gfp and tagRfp or mCherry fluorescence was simultaneously detected using a two-channel

imager (DV2, Photometrics, Tucson, AZ, USA). All images and videos were acquired and

analysed using Metamorph (Versions 7.7.0.0 and 7.7.4.0; Molecular Devices, Sunnyvale, CA,

USA).

Cell length was assessed by measuring the length of single cells from pole-to-pole. For

hyphae, empty sections were not included in the measurements. The length of empty sections

was assessed by measuring the distance from septum to septum of the first empty section at

the distal pole of hyphae. Cell wall defects induced by Calcoflour White were quantified by

manually scoring for the presence of abnormal cell wall shapes.

For the analysis of co-localisation and velocity of moving signals the acquired videos

were converted to kymographs using Metamorph. Co-localisation was assessed by

quantifying kymographs acquired by dual-colour imaging. Changes in direction were counted

as individual signals. Processive signals (distance travelled > 5 μm) were counted manually.

Velocity was only measured for processive signals (movement > 5 μm). For all

quantifications, at least three independent experiments were analysed. Statistical analysis was

performed using Prism5 (Graphpad, La Jolla, CA, USA).

The gradient of Grp1-Gfp, Pab1-Gfp and Gfp in the presence and absence of Rrm4 was

quantified by measuring the fluorescence intensity in a previously specified region of interest

(ROI; ROI1 in vicinity to the nucleus and ROI2 at the hyphal tip). The fluorescence close to

the nucleus was then set in relation to the signal intensity at the tip. Statistical analysis was

performed using Prism5 (Graphpad, La Jolla, CA, USA).

iCLIP experiments

The iCLIP protocol was modified from [30] and the original application in *U. maydis* [17]. (i)

The initial two-step tap tag purification [17] was switched to single-step purification using

strains expressing C-terminal Gfp fusions and Gfp-trap immunoprecipitation (GFP-

Trap®_MA, ChromoTek, Martinsried, Germany) [73]. (ii) For UV-C crosslinking, cells were

irradiated continuously in a single session. (iii) The addition of DNase and RNase I was

omitted from the samples used for iCLIP library preparation. Due to the high RNase activity

in fungal cells, the addition of external RNases was unnecessary, therefore RNase I (10

Units/sample; 16 minutes, 37°C; Thermofisher Scientific, Darmstadt, Germany) was only

added in the control experiments (Fig 3A).

All iCLIP experiments were performed with 150 ml cultures grown to an $OD_{600} = 0.5$ in CM-

medium (supplemented with 1% glc) and then shifted to NM-medium (supplemented with 1%

glc) to induce hyphal growth. After 6 h, each culture was split into 3 x 50 ml, harvested by

centrifugation at 6,280 g for 10 min at 4°C, and the cell pellets were resuspended in 5 ml ice-

cold PBS. For UV-C crosslinking, cells were irradiated once with 200 mJ/cm^2 at 254 nm

(Biolink UV-Crosslinker, Vilber-Lourmat, Eberhardzell, Germany) as a thin layer in a square

petri dish (10 cm^2), pooled in a 50 ml tube, and harvested by centrifugation at 6,280 g for 10

min at 4°C. The cell pellet was resuspended in 6 ml lysis buffer (50 mM Tris-HCl, pH 7.4,

100 mM NaCl, 1% Nonidet P-40, 0.1% SDS, 0.5% sodium deoxycholate) supplemented with

inhibitors (per 10 ml lysis buffer): 2× Complete protease inhibitor EDTA-free (Roche

Diagnostics, Mannheim, Germany), 1 mM reduced DTT (GERBU, Heidelberg, Germany), 5

mM Benzamidine (Sigma), 1 mM PMSF (Sigma), 0.75 µg/µl heparin (Sigma), 5.25 ng/µl

pepstatin A (Sigma) and 15 µl SUPERase-In (20 U/µl; Thermo Fisher Scientific). The

solution was split into three times 2 ml for cell lysis and immunoprecipitation. Cell lysis was

performed in a Retsch ball mill (2 ml cell suspension/grinding jar, 2 grinding balls, d=12 mm;

MM400; Retsch, Haan, Germany) 3 x 10 min at 30 Hz while keeping samples frozen using

liquid nitrogen. All of the following steps up to the 5´ labelling of RNA were performed at

4°C. The cell lysates were pooled in a 15 ml tube, split in precooled 1.5 ml tubes and cleared

by centrifugation at 16,200 g for 15 min at 4°C. The protein-RNA complexes were

immunoprecipitated from each strain by using a total of 60 µl GFP-Trap magnetic agarose

beads (Chromotek; GFP-Trap®_MA) [73]. Before immunoprecipitation, the 60 µl beads were

added to a 2 ml tube and pre-washed three times with 500 µl ice-cold lysis buffer (without

inhibitors) and afterwards split into three times 20 µl beads in 2 ml tubes. For

immunoprecipitation, 2 ml cleared cell lysate were added to 20 µl beads and the tubes were

incubated rotating for 1 h at 4°C. Next, the beads were combined in a 2 ml tube and washed

three times with 900 µl high-salt wash buffer (50 mM Tris-HCl, pH 7.4, 1 M NaCl, 1 mM

EDTA, 1% Igepal CA-630, 0.1% SDS, 0.5% sodium deoxycholate) and four times with 900

µl PNK wash buffer (20 mM Tris-HCl, pH 7.4, 10 mM $MgCl_2$, 0.2% Tween-20). For each

subsequent enzymatic reaction, the 60 µl beads were split into three times 20 µl beads in 2 ml

tubes and re-pooled for the washing steps. 3´ end RNA dephosphorylation, L3 adapter

ligation, 5´ end phosphorylation, SDS-PAGE and nitrocellulose transfer were performed as

described [30] with minor changes implemented. For 5´ end phosphorylation, 50% of each

sample was radioactively labelled with ^{32}P-γ-ATP for 10 min at 37°C and the labelled beads

were washed once with PNK wash buffer before they were combined with the unlabelled

beads. The beads were diluted in 80 µl NuPAGE LDS loading buffer (Thermo Fisher

Scientific) with 0.1 M DTT added. Samples were heated at 70°C for 5 min and split among

two adjacent wells on a 4-12% NuPAGE Bis-Tris gel (Thermo Fisher Scientific). The protein-

RNA complexes were separated at 180 V for 70 min in MOPS running buffer with NuPAGE

reducing agent added (Thermo Fisher Scientific) and transferred onto a nitrocellulose

membrane. For Western blotting, Gfp was detected using monoclonal α-GFP primary

antibodies (clones 7.1 and 13.1; Sigma) and a mouse IgG HRP conjugate (H+L; Promega,

Madison, WI, USA) as secondary antibody. Peroxidase activity was determined using the

AceGlow blotting detection system (Peqlab, Erlangen, Germany).

Labelled RNA was detected by exposure of X-ray films ranging from 2 h - overnight at

-80°C. cDNA library preparation was performed as described before [30]. To avoid over-

amplification of the cDNA library, the optimal number of PCR cycles producing PCR

products of around 150 nt (cDNA insert = 20-30 nt; L3 adapter, RT-primer and P3/P5 Solexa

primers = 128 nt) was tested for each protein in every experiment (PCR cycler PTC-200, MJ

Research, St. Bruno, Quebec, Canada; Fig EV2D). For all Grp1 and Gfp replicates, 18 PCR

cycles were determined to be optimal, while the optimal PCR cycle numbers for the Rrm4

replicates were determined to be 18 and 22 PCR cycles. The iCLIP libraries were multiplexed

and sequenced on an Illumina HiSeq 2500 (San Diego, CA, USA; 51-nt reads, single-end),

yielding a total of 118 million reads.

iCLIP data processing

All bioinformatics analyses are based on the *U. maydis* 521 genome sequence (original

PEDANT database name p3_t237631_Ust_maydi_v2GB) and the associated gene annotation

(version p3_t237631_Ust_maydi_v2GB.gff3; both downloaded from

ftp://ftpmips.gsf.de/fungi/Ustilaginaceae/Ustilago_maydis_521/). We extended all genes by

300 nt on either side to include potential 5´ and 3´ UTR regions which are currently not

annotated in the *U. maydis* genome. For manual annotation of transcript ends, RNASeq data

(AB33 hyphae, 6 h.p.i.) were used, and transcript ends were defined at the position where

read coverage dropped below 10.

Basic quality checks were applied to all sequenced reads using FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Afterwards, iCLIP reads were

filtered based on sequencing quality (Phred score) in the barcode region, keeping only reads

with at most one position with a Phred score < 20 in the experimental barcode (positions 4 to

7) and without any position with a Phred score < 17 in the random barcode (positions 1 to 3

and 8 to 9). The reads were then de-multiplexed based on the experimental barcode at

positions 4 to 7 using Flexbar (version 2.4, GitHub, San Francisco, CA, USA) without

allowing mismatches [74]. The following analysis steps were applied to all individual

samples: remaining adapter sequences were trimmed from the 3´ end of the reads using

Flexbar (version 2.4) allowing one mismatch in 10 nt, requiring a minimal overlap of 1 nt

between read and adapter as well as removing all reads with a remaining length of less than

24 nt (including the 9-nt barcode). The first 9-nt of each read containing the barcode were

trimmed off and added to the read name in the fastq file.

Filtered and trimmed reads were mapped to the *U. maydis* genome and its gene annotation

using STAR (version 2.5.1b, GitHub) [75], allowing up to two mismatches and without soft-

clipping on the 5´ end of the reads. Only uniquely mapped reads were kept for further

analysis.

After mapping and filtering, duplicate reads were marked using the *dedup* function from

bamUtil (version 1.0.7; https://github.com/statgen/bamUtil) and removed if carrying an

identical random barcode, and hence representing technical duplicates. The nucleotide

position upstream of each aligned read was considered as the 'crosslink nucleotide', with each

read counted as individual 'crosslink event'. The total number of crosslink events for the

different iCLIP libraries can be found in Fig EV3A. To assess the reproducibility between

biological replicates (Fig EV3C), we counted the number of crosslink events within each

gene.

We then proceeded to define putative RBP 'binding sites', i.e. sites that show a

significant enrichment of crosslink events compared to the surrounding sequence. Since most

RBPs contact multiple positions in the RNA, the crosslink events in a binding site commonly

spread across more than one nucleotide. For the identification of such windows with

significantly increased crosslink event frequency, peak calling was performed using ASPeak

that accounts for the different expression levels of the underlying transcripts [76]. Peak

calling was performed on merged replicates for each RBP to maximise the signal. In order to

obtain binding sites of a uniform width, we further processed the predicted peaks [77]. We

first determined the summit of each peak (i.e. the position with highest number of crosslink

events within the peak; if two or more positions showed the same count, the most upstream

position was taken) and then defined a binding site window of 9-nt that was centred around

the summit position. Overlapping peaks were merged and newly centred on the summit of the

combined window as described above. Finally, we filtered the binding sites as follows: (i) To

account for reproducibility, we required each binding site to be detected by at least 5 crosslink

events from each biological replicate. (ii) We further removed all Rrm4 and Grp1 binding

sites that overlapped by at least 1 nt with any of 88 reproducible Gfp binding sites. To define

the set of target genes that were exclusively bound by Rrm4 and showed no evidence of Grp1

binding ('Rrm4-unique targets'), we subtracted the intermediate list of binding sites at this

point for Grp1 from the final Rrm4 binding sites (after SOB filtering, see below), and vice

versa.

For the remaining analysis, we included an additional processing step to focus on the

top 25% of binding sites. To determine this set, we used the ratio of crosslink events within

the binding site over the background crosslink events in the surrounding sequence (termed

'signal-over-background', SOB) as a proxy of binding site strength: Crosslink events from

both replicates were summed up within each binding site and divided by the gene-specific

background to obtain an SOB value for each binding site. The background for each gene was

defined as the number of crosslink events outside the binding sites divided by the number of

nucleotides in the gene that harbour such background crosslink events. We note that while this

procedure alleviates the impact of transcript-level differences, it does not correct for UV

crosslinking bias and similar confounding factors, and should therefore only be used as a

proxy for bulk comparisons of binding sites. Only binding sites within the top 25% of the

SOB distribution for each RBP were taken into consideration for further analyses.

This SOB filtering procedure yielded a total of 6,412 and 6,478 binding sites for Rrm4

and Grp1, respectively (Fig 3B, Fig EV3). The binding sites corresponded to 3,262 and 3,165

target transcripts for Rrm4 and Grp1, respectively. When assigning genomic nucleotides and

RBP binding sites to distinct transcript regions (Fig 3D), we applied the following hierarchy

to resolve overlapping annotation: 3´ UTR > 5´ UTR > exon > intron. Binding sites of Rrm4

and Grp1 were considered as overlapping if at least 1 nt was shared between the 9-nt binding

sites.

To study the distribution of binding sites of Rrm4 and Grp1 in 3´ UTRs (Fig 4C), we

manually annotated transcript ends. This was necessary, because the *U. maydis* gene

annotation only includes ORF regions of genes. To this end, RNASeq data from AB33 hyphae

(6 h.p.i.) were visually inspected and 3´ ends were defined due to strong reduction in RNASeq

coverage.

Enriched sequence motifs around RBP binding sites were identified using DREME [35;

parameter –norc to search the coding strand only) to analyse a 30-nt window around all

binding site summits compared to shuffled nucleotides. Based on the sequence profile and the

UAUG content around binding sites (Fig 5A-B), we counted a binding site as UAUG-

containing if it harboured the motif with the last 5 nt of the 9-nt binding sites. In order to

assess the frequency of Rrm4 binding sites at UAUG motifs at start codons and in other

regions (Fig EV5E), we used all transcripts that carry Rrm4 binding sites anywhere within the

transcript and are hence sufficiently expressed to be detected in our iCLIP analysis.

Analysing functional categories of cellular components on identified targets was

performed using the FunCat annotation scheme (http://mips.gsf.de/funcatDB/; version 2.1,

reference genome: p3_t237631_Ust_maydi_v2GB) [78]. Categories were filtered during

analysis for enrichment by setting the p-value threshold to <0.05.

RNASeq library preparation and data processing

RNA was extracted from AB33 hyphae 6 h.p.i. using the RNeasy Mini kit following the

manufacturer's instructions for preparation of total RNA from yeast (Qiagen, Hilden,

Germany). To this end, AB33 hyphae were opened in a Retsch ball mill (3 balls, d=4 mm ;

MM400; Retsch, Haan, Germany) 4 times for 5 min at 30 Hz while keeping samples frozen

using liquid nitrogen. The resulting cell powder was resolved in 450 µl RLT buffer (+β-

mercaptoethanol) and centrifuged at 13,000 rpm for 2 min at 4°C. The supernatant was

transferred to a new reaction tube, mixed with 1 volume 70% EtOH and then added to the

RNeasy spin column. All following processing steps were performed according to

manufacturer's instructions. TruSeq RNA Library Prep kit v2 (Illumina, San Diego, CA,

USA) was used for cDNA library generation. The cDNA libraries were sequenced using the

HiSeq 2000 platform (Illumina) with 151-nt single-end reads.

Basic quality checks were applied to all sequenced reads using FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Afterwards, RNASeq reads

were trimmed based on sequencing quality (Phred score) using Flexbar (version 3.0.3) [74].

Specifically, adapter sequences were removed (TruSeq Universal Adapter), and reads were

trimmed at the first position with a Phred score < 20 and removed if the remaining read length

was less than 20 nt. The trimmed reads were mapped to the *U. maydis* genome and its gene

annotation using STAR (version 2.5.3a) [75], allowing up to five mismatches with soft-

clipping. Uniquely mapped reads were kept for further analysis.

Yeast three-hybrid analysis

Yeast three-hybrid experiments were performed as described previously [36, 79]. To test the

interaction with Rrm4, the plasmids (Appendix Table S4) encoding the RNA aptamers

SELEX-A1[36] or mutated SELEX-A1 (mUAUG; this work) were cotransformed in strain

L40 coat with the corresponding plasmids encoding for Rrm4 or mutated variants [36, 80].

Transformed cells were incubated on SC -ura -leu plates (2-3 d at 28°C) before single clones

were selected. Interaction was assayed as growth on selection medium SC -his +1 mM 3-AT

(3-amino-1,2,4-triazole; Sigma-Aldrich; 3 d at 28°C). For the serial dilution assays, single

clones were grown in SC -ura -leu medium to a starting $OD_{600} = 0,5$ and sequentially diluted

1:5 in water. The dilutions were then spotted on control (SC -ura -leu) and selection (SC -his

+1 mM 3-AT) plates and incubated at 28°C.

Data Availability

The iCLIP and RNASeq dataset are available from GEO under the accession numbers

GSE109557 and GSE109560, respectively. The associated SuperSeries is GSE109561.

Acknowledgements

We thank Dr. J. Kahnt for pilot Tap tag experiments and Dr T. Pohlmann for initial work on

Grp1 fusion proteins. We acknowledge R.F.X. Sutandy for iCLIP support, M. Brüggemann for

RNASeq data analysis, F. Finkernagel for bioinformatics and Dr. M. Seiler for critical reading

of the manuscript. We are grateful to U. Gengenbacher and S. Esch for excellent technical

assistance and members of the IMB Genomics core facility for technical assistance. The work

was funded by grants from the Deutsche Forschungsgemeinschaft to MF (FE 448/8-1;

FOR2333-TP03 FE448/10-1; CEPLAS EXC1028) and KZ (FOR2333-TP10 ZA881/1-1) and

JK (SPP1935 KO4566/2-1) and JU (European Research Council; 617837-Translate).

Author contributions

LO, CH, JU, JKö, MF and ZK designed this study and analysed the data. LO and CH

performed experiments to characterise Grp1 and comparative iCLIP analysis. JKoe performed

preliminary affinity purification experiments. SB, AB, and ZK performed all computational

iCLIP data analyses. MF and ZK drafted and revised the manuscript with input from all co-

authors. MF and ZK directed the project.

Competing interests

The authors declare that they have no competing interests.

References

1.   Holt CE, Bullock SL (2009) Subcellular mRNA localization in animal cells and why it

matters. *Science* **326**: 1212-1216

2.   Martin KC, Ephrussi A (2009) mRNA localization: gene expression in the spatial

dimension. *Cell* **136**: 719-730

3.   Eliscovich C, Singer RH (2017) RNP transport in cell biology: the long and winding

road. *Current opinion in cell biology* **45**: 38-46

4.   Mofatteh M, Bullock SL (2017) SnapShot: Subcellular mRNA localization. *Cell* **169**:

178.e1

5.   Baumann S, Pohlmann T, Jungbluth M, Brachmann A, Feldbrügge M (2012) Kinesin-3

and dynein mediate microtubule-dependent co-transport of mRNPs and endosomes. *J*

Cell Sci **125**: 2740-2752

6.   Baumann S, König J, Koepke J, Feldbrügge M (2014) Endosomal transport of septin

mRNA and protein indicates local translation on endosomes and is required for correct

septin filamentation. *EMBO Rep* **15**: 94-102

7.   Egan MJ, McClintock MA, Reck-Peterson SL (2012) Microtubule-based transport in

filamentous fungi. *Curr Opin Microbiol* **15**: 637-645

8.   Steinberg G (2014) Endocytosis and early endosome motility in filamentous fungi. *Curr*

Opin Microbiol **20**: 10-18

9.   Guimaraes SC, Schuster M, Bielska E, Dagdas G, Kilaru S, Meadows BR, Schrader M,

Steinberg G (2015) Peroxisomes, lipid droplets, and endoplasmic reticulum "hitchhike"

on motile early endosomes. *J Cell Biol* **211**: 945-954

10.   Salogiannis J, Egan MJ, Reck-Peterson SL (2016) Peroxisomes move by hitchhiking on

early endosomes using the novel linker protein PxdA. *The Journal of cell biology* **212**:

289-296

11.   Salogiannis J, Reck-Peterson SL (2016) Hitchhiking: a non-canonical mode of

709   microtubule-based transport. *Trends in cell biology* **27**: 141-150

710   12.   Haag C, Pohlmann T, Feldbrugge M (2017) The ESCRT regulator Did2 maintains the

711   balance between long-distance endosomal transport and endocytic trafficking. *PLoS*

712   *Genet* **13**: e1006734

713   13.   Haag C, Steuten B, Feldbrügge M (2015) Membrane-coupled mRNA trafficking in

714   fungi. *Annu Rev Microbiol* **69**: 265-281

715   14.   Vollmeister E, Feldbrügge M (2010) Posttranscriptional control of growth and

716   development in *Ustilago maydis*. *Curr Opin Microbio* **13**: 693-699

717   15.   Becht P, Vollmeister E, Feldbrügge M (2005) Role for RNA-binding proteins implicated

718   in pathogenic development of *Ustilago maydis*. *Euk Cell* **4**: 121-133

719   16.   Becht P, König J, Feldbrügge M (2006) The RNA-binding protein Rrm4 is essential for

720   polarity in *Ustilago maydis* and shuttles along microtubules. *J Cell Sci* **119**: 4964-4973

721   17.   König J, Baumann S, Koepke J, Pohlmann T, Zarnack K, Feldbrügge M (2009) The

722   fungal RNA-binding protein Rrm4 mediates long-distance transport of *ubi1* and *rho3*

723   mRNAs. *EMBO J* **28**: 1855-1866

724   18.   Koepke J, Kaffarnik F, Haag C, Zarnack K, Luscombe NM, König J, Ule J, Kellner R,

725   Begerow D, Feldbrügge M (2011) The RNA-binding protein Rrm4 is essential for

726   efficient secretion of endochitinase Cts1. *Mol Cell Proteom* **10**: M111.011213 1-15

727   19.   Zander S, Baumann S, Weidtkamp-Peters S, Feldbrügge M (2016) Endosomal assembly

728   and transport of heteromeric septin complexes promote septin cytoskeleton formation. *J*

729   *Cell Sci* **129**: 2778-2792

730   20.   Zhu X, Buhrer C, Wellmann S (2016) Cold-inducible proteins CIRP and RBM3, a

731   unique couple with activities far beyond the cold. *Cellular and molecular life sciences :*

732   *CMLS* **73**: 3839-3859

21.   Kang H, Park SJ, Kwak KJ (2013) Plant RNA chaperones in stress response. *Trends*

Plant Sci **18**: 100-106

22.   Brachmann A, Weinzierl G, Kämper J, Kahmann R (2001) Identification of genes in the

bW/bE regulatory cascade in *Ustilago maydis*. *Mol Microbiol* **42**: 1047-1063

23.   Imai K, Noda Y, Adachi H, Yoda K (2005) A novel endoplasmic reticulum membrane

protein Rcr1 regulates chitin deposition in the cell wall of Saccharomyces cerevisiae. *J*

Biol Chem **280**: 8275-84

24.   Ram AF, Klis FM (2006) Identification of fungal cell wall mutants using susceptibility

assays based on Calcofluor white and Congo red. *Nat Protoc* **1**: 2253-2256

25.   Baumann S, Takeshita N, Grün N, Fischer R, Feldbrügge M (2015) *Live cell imaging of*

endosomal trafficking in fungi. In Methods in Mol Biol: Membrane trafficking, Tang BL

(ed) pp 347-363. New York: Springer

26.   Merzlyak EM, Goedhart J, Shcherbo D, Bulina ME, Shcheglov AS, Fradkov AF,

Gaintzeva A, Lukyanov KA, Lukyanov S, Gadella TW*, et al.* (2007) Bright monomeric

red fluorescent protein with an extended fluorescence lifetime. *Nature methods* **4**: 555-

557

27.   Campbell RE, Tour O, Palmer AE, Steinbach PA, Baird GS, Zacharias DA, Tsien RY

(2002) A monomeric red fluorescent protein. *Proc Natl Acad Sci U S A* **99**: 7877-7882

28.   Hogan DJ, Riordan DP, Gerber AP, Herschlag D, Brown PO (2008) Diverse RNA-

binding proteins interact with functionally related sets of RNAs, suggesting an

extensive regulatory system. *PLoS Biol* **6**: e255

29.   König J, Zarnack K, Rot G, Curk T, Kayikci M, Zupan B, Turner DJ, Luscombe NM,

Ule J (2010) iCLIP reveals the function of hnRNP particles in splicing at individual

nucleotide resolution. *Nat Struct Mol Biol* **17**: 909-915

30.   Huppertz I, Attig J, D'Ambrogio A, Easton LE, Sibley CR, Sugimoto Y, Tajnik M,

758   König J, Ule J (2014) iCLIP: protein-RNA interactions at nucleotide resolution.

759   *Methods* **65**: 274-287

760   31.   Kämper J, Kahmann R, Bölker M, Ma LJ, Brefort T, Saville BJ, Banuett F, Kronstad

761   JW, Gold SE, Müller O*, et al.* (2006) Insights from the genome of the biotrophic fungal

762   plant pathogen *Ustilago maydis*. *Nature* **444**: 97-101

763   32.   Chabanon H, Mickleburgh I, Burtle B, Pedder C, Hesketh J (2005) An AU-rich stem-

764   loop structure is a critical feature of the perinuclear localization signal of c-myc mRNA.

765   *Biochem J* **392**: 475-483

766   33.   Levadoux M, Mahon C, Beattie JH, Wallace HM, Hesketh JE (1999) Nuclear import of

767   metallothionein requires its mRNA to be associated with the perinuclear cytoskeleton.

768   *The Journal of biological chemistry* **274**: 34961-34966

769   34.   Straube A, Enard W, Berner A, Wedlich-Söldner R, Kahmann R, Steinberg G (2001) A

770   split motor domain in a cytoplasmic dynein. *EMBO J* **20**: 5091-5100

771   35.   Bailey TL (2011) DREME: motif discovery in transcription factor ChIP-seq data.

772   *Bioinformatics* **27**: 1653-1659

773   36.   König J, Julius C, Baumann S, Homann M, Göringer HU, Feldbrügge M (2007)

774   Combining SELEX and yeast three-hybrid system for *in vivo* selection and

775   classification of RNA aptamers. *RNA* **13**: 614-622

776   37.   Lécuyer E, Yoshida H, Parthasarathy N, Alm C, Babak T, Cerovina T, Hughes TR,

777   Tomancak P, Krause HM (2007) Global analysis of mRNA localization reveals a

778   prominent role in organizing cellular architecture and function. *Cell* **131**: 174-187

779   38.   Jambor H, Surendranath V, Kalinka AT, Mejstrik P, Saalfeld S, Tomancak P (2015)

780   Systematic imaging reveals features and changing localization of mRNAs in Drosophila

781   development. *eLife* **4**

39.   Cajigas IJ, Tushev G, Will TJ, tom Dieck S, Fuerst N, Schuman EM (2012) The local

transcriptome in the synaptic neuropil revealed by deep sequencing and high-resolution

imaging. *Neuron* **74**: 453-466

40.   Rangaraju V, Tom Dieck S, Schuman EM (2017) Local translation in neuronal

compartments: how local is local? *EMBO Rep* **18**: 693-711

41.   Calabretta S, Richard S (2015) Emerging roles of disordered sequences in RNA-binding

proteins. *Trends Biochem Sci* **40**: 662-672

42.   Ying Y, Wang XJ, Vuong CK, Lin CH, Damianov A, Black DL (2017) Splicing

activation by Rbfox requires self-aggregation through Its tyrosine-rich domain. *Cell*

170: 312-323 e10

43.   Jung H, Yoon BC, Holt CE (2012) Axonal mRNA localization and local protein

synthesis in nervous system assembly, maintenance and repair. *Nat Rev Neurosci* **13**:

308-324

44.   Kato M, Han TW, Xie S, Shi K, Du X, Wu LC, Mirzaei H, Goldsmith EJ, Longgood J,

Pei J*, et al.* (2012) Cell-free formation of RNA granules: low complexity sequence

domains form dynamic fibers within hydrogels. *Cell* **149**: 753-767

45.   Ciuzan O, Hancock J, Pamfil D, Wilson I, Ladomery M (2015) The evolutionarily

conserved multifunctional glycine-rich RNA-binding proteins play key roles in

development and stress adaptation. *Physiol Plant* **153**: 1-11

46.   Meyer K, Koster T, Nolte C, Weinholdt C, Lewinski M, Grosse I, Staiger D (2017)

Adaptation of iCLIP to plants determines the binding landscape of the clock-regulated

RNA-binding protein AtGRP7. *Genome Biol* **18**: 204

47.   Peretti D, Bastide A, Radford H, Verity N, Molloy C, Martin MG, Moreno JA, Steinert

JR, Smith T, Dinsdale D*, et al.* (2015) RBM3 mediates structural plasticity and

protective effects of cooling in neurodegeneration. *Nature* **518**: 236-239

48.   Bastide A, Peretti D, Knight JR, Grosso S, Spriggs RV, Pichon X, Sbarrato T, Roobol A,

Roobol J, Vito D*, et al.* (2017) RTN3 is a novel cold-induced protein and mediates

neuroprotective effects of RBM3. *Curr Biol* **27**: 638-650

49.   Kim JS, Park SJ, Kwak KJ, Kim YO, Kim JY, Song J, Jang B, Jung CH, Kang H (2007)

Cold shock domain proteins and glycine-rich RNA-binding proteins from *Arabidopsis*

thaliana can promote the cold adaptation process in *Escherichia coli*. *Nucleic Acids Res*

35: 506-516

50.   Yano T, Lopez de Quinto S, Matsui Y, Shevchenko A, Shevchenko A, Ephrussi A (2004)

Hrp48, a *Drosophila* hnRNPA/B homolog, binds and regulates translation of oskar

mRNA. *Dev Cell* **6**: 637-648

51.   Huynh JR, Munro TP, Smith-Litiere K, Lepesant JA, St Johnston D (2004) The

Drosophila hnRNPA/B homolog, Hrp48, is specifically required for a distinct step in

osk mRNA localization. *Dev Cell* **6**: 625-635

52.   Goodrich JS, Clouse KN, Schupbach T (2004) Hrb27C, Sqd and Otu cooperatively

regulate *gurken* RNA localization and mediate nurse cell chromosome dispersion in

Drosophila oogenesis. *Development* **131**: 1949-1958

53.   König J, Zarnack K, Luscombe NM, Ule J (2012) Protein-RNA interactions: new

genomic technologies and perspectives. *Nat Rev Genet* **13**: 77-83

54.   Van Nostrand EL, Pratt GA, Shishkin AA, Gelboin-Burkhart C, Fang MY,

Sundararaman B, Blue SM, Nguyen TB, Surka C, Elkins K*, et al.* (2016) Robust

transcriptome-wide discovery of RNA-binding protein binding sites with enhanced

CLIP (eCLIP). *Nature methods* **13**: 508-514

55.   Zarnegar BJ, Flynn RA, Shen Y, Do BT, Chang HY, Khavari PA (2016) irCLIP platform

for efficient characterization of protein-RNA interactions. *Nature methods* **13**: 489-492

56.   Doyle M, Kiebler MA (2011) Mechanisms of dendritic mRNA transport and its role in

synaptic tagging. *EMBO J* **30**: 3540-3552

57.   Higuchi Y, Ashwin P, Roger Y, Steinberg G (2014) Early endosome motility spatially

834   organizes polysome distribution. *J Cell Biol* **204**: 343-357

835   58.   Margeot A, Blugeon C, Sylvestre J, Vialette S, Jacq C, Corral-Debrinski M (2002) In

836   *Saccharomyces cerevisiae*, *ATP2* mRNA sorting to the vicinity of mitochondria is

837   essential for respiratory function. *EMBO J* **21**: 6893-6904

838   59.   Gehrke S, Wu Z, Klinkenberg M, Sun Y, Auburger G, Guo S, Lu B (2015) PINK1 and

839   Parkin control localized translation of respiratory chain component mRNAs on

840   mitochondria outer membrane. *Cell Metab* **21**: 95-108

841   60.   Gold VA, Chroscicki P, Bragoszewski P, Chacinska A (2017) Visualization of cytosolic

842   ribosomes on the surface of mitochondria by electron cryo-tomography. *EMBO Rep* **18**:

843   1786-1800

844   61.   Lesnik C, Golani-Armon A, Arava Y (2015) Localized translation near the

845   mitochondrial outer membrane: An update. *RNA Biol* **12**: 801-809

846   62.   Zhang Y, Chen Y, Gucek M, Xu H (2016) The mitochondrial outer membrane protein

847   MDI promotes local protein synthesis and mtDNA replication. *EMBO J* **35**: 1045-1057

848   63.   Sen A, Cox RT (2016) Clueless is a conserved ribonucleoprotein that binds the

849   ribosome at the mitochondrial outer membrane. *Biol Open* **5**: 195-203

850   64.   Pohlmann T, Baumann S, Haag C, Albrecht M, Feldbrügge M (2015) A FYVE zinc

851   finger domain protein specifically links mRNA transport to endosome trafficking. *eLife*

852   **4**: 10.7554/eLife.06041

853   65.   Halstead JM, Lionnet T, Wilbertz JH, Wippich F, Ephrussi A, Singer RH, Chao JA

854   (2015) Translation. An RNA biosensor for imaging the first round of translation from

855   single cells to living animals. *Science* **347**: 1367-1671

856   66.   Wu B, Eliscovich C, Yoon YJ, Singer RH (2016) Translation dynamics of single

857   mRNAs in live cells and neurons. *Science* **352**: 1430-1435

858   67.   Graber TE, Hebert-Seropian S, Khoutorsky A, David A, Yewdell JW, Lacaille JC,

859   Sossin WS (2013) Reactivation of stalled polyribosomes in synaptic plasticity. *Proc*

860   *Natl Acad Sci U S A* **110**: 16205-16210

861   68.   Darnell JC, Van Driesche SJ, Zhang C, Hung KY, Mele A, Fraser CE, Stone EF, Chen

862   C, Fak JJ, Chi SW*, et al.* (2011) FMRP stalls ribosomal translocation on mRNAs linked

863   to synaptic function and autism. *Cell* **146**: 247-261

864   69.   Brachmann A, König J, Julius C, Feldbrügge M (2004) A reverse genetic approach for

865   generating gene replacement mutants in *Ustilago maydis*. *Mol Gen Genom* **272**: 216-

866   226

867   70.   Shaner NC, Campbell RE, Steinbach PA, Giepmans BN, Palmer AE, Tsien RY (2004)

868   Improved monomeric red, orange and yellow fluorescent proteins derived from

869   *Discosoma* sp. red fluorescent protein. *Nat Biotechnol* **22**: 1567-1572

870   71.   Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H,

871   Valentin F, Wallace IM, Wilm A, Lopez R*, et al.* (2007) Clustal W and Clustal X version

872   2.0. *Bioinformatics* **23**: 2947-2948

873   72.   Baumann S, Zander S, Weidtkamp-Peters S, Feldbrügge M (2016) *Live cell imaging of*

874   *septin dynamics in Ustilago maydis*. In Septins, Gladfelter AS (ed) pp 143-149. Elsevier

875   Inc.

876   73.   Rothbauer U, Zolghadr K, Muyldermans S, Schepers A, Cardoso MC, Leonhardt H

877   (2008) A versatile nanotrap for biochemical and functional studies with fluorescent

878   fusion proteins. *Mol Cell Proteomics* **7**: 282-289

879   74.   Dodt M, Roehr JT, Ahmed R, Dieterich C (2012) FLEXBAR-flexible barcode and

880   adapter processing for next-generation sequencing platforms. *Biology* **1**: 895-905

881   75.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,

882   Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15-

883   21

884   76.   Kucukural A, Ozadam H, Singh G, Moore MJ, Cenik C (2013) ASPeak: an abundance

885   sensitive peak detection algorithm for RIP-Seq. *Bioinformatics* **29**: 2485-2486

886   77.   Sutandy FXR, Ebersberger S, Huang L, Busch A, Bach M, Kang HS, Fallmann J,

887   Maticzka D, Backofen R, Stadler PF*, et al.* (2018) *In vitro* iCLIP-based modeling

888   uncovers how the splicing factor U2AF2 relies on regulation by cofactors. *Genome Res*

889   **28**: 699-713

890   78.   Ruepp A, Zollner A, Maier D, Albermann K, Hani J, Mokrejs M, Tetko I, Guldener U,

891   Mannhaupt G, Münsterkötter M*, et al.* (2004) The FunCat, a functional annotation

892   scheme for systematic classification of proteins from whole genomes. *Nucleic Acids Res*

893   **32**: 5539-5545

894   79.   Vollmeister E, Haag C, Zarnack K, Baumann S, König J, Mannhaupt G, Feldbrügge M

895   (2009) Tandem KH domains of Khd4 recognize AUACCC and are essential for

896   regulation of morphology as well as pathogenicity in *Ustilago maydis*. *RNA* **15**: 2206-

897   2218

898   80.   SenGupta DJ, Zhang B, Kraemer B, Pochart P, Fields S, Wickens M (1996) A three-

899   hybrid system to detect RNA-protein interactions *in vivo*. *Proc Natl Acad Sci U S A* **93**:

900   8496-8501

901   81.   Kozak M (2005) Regulation of translation via mRNA structure in prokaryotes and

902   eukaryotes. *Gene* **361**: 13-37

Figure Legends

Fig 1. Grp1 is important for hyphal growth under suboptimal conditions.

(**A**) Sequence alignment of glycine-rich proteins (Fig EV1B). *Um*Grp1 from *U. maydis*

(UMAG_02412), *At*GRP7 (RBG7) from *A. thaliana* (NC_003071.7), *Hs*RBM3 and *Hs*CIRBP

from *H. sapiens* (NC_000023.11 and NC_000019.10, respectively). Amino acid positions

within RRM that are identical in at least three proteins are highlighted in green (boxes

indicate RNA contact regions RNP1 and RNP2). Glycine and arginine/glutamine residues in

the glycine-rich region are labelled in red and blue, respectively.

(**B**) Hyphae of AB33 derivatives (6 h.p.i.). Growth direction and basal septa are marked by

arrows and asterisks, respectively (scale bar, 10 µm).

(**C**) Hyphal length over time. Black and grey dots represent hyphae with and without septa,

respectively. Shown are merged data (> 200 hyphae per strain) from three independent

experiments, overlaid with the mean of means, red line, and standard error of the mean

(s.e.m.). Significance was assessed using paired two-tailed Student's t-test on the mean hyphal

lengths from the replicate experiments, followed by multiple testing correction (Benjamini-

Hochberg). Significant p-values ($p < 0.05$) for comparison against wild type within each time

point are indicated above.

(**D**) Length of empty sections (see Fig EV1J). Merged data from three independent

experiments are shown together, overlaid with mean of means, red line, and s.e.m. (total

hyphae analysed: *wt*, 250; *grp1Δ*, 190; *grp1-gfp*, 332; difference in means between *wt* and

grp1-gfp was statistically not significant, ns, p= 0.95; paired two-tailed Student's t-test on the

mean lengths from the replicate experiments (n=3).

(**E**) Differential interference contrast (DIC, top) and fluorescence images (bottom) of AB33

hyphae (5 h.p.i.) stressed at 3 h.p.i. with cell wall inhibitor CFW (2.5 µM). Arrowheads

indicate aberrant cell wall deformation (scale bar, 10 µm).

(**F**) Percentage of hyphae with normal cell walls with (stressed) and without (unstressed)

CFW treatment (data points represent percentages of three independent experiments, n=3;

mean, dark grey lines, and standard error of the mean, s.e.m., > 100 hyphae analysed per

experiment; paired two-tailed Student's t-test on the means).

Fig 2. Grp1 shuttles on Rrm4-positive endosomes throughout hyphae.

(**A**) Micrographs (DIC and inverted fluorescence image; scale bar, 10 µm) and corresponding

kymographs of AB33 hyphae (6 h.p.i.) expressing Grp1-Gfp, Rrm4-Gfp or Pab1-Gfp (arrow

length on the left and bottom indicate time and distance, 10 s and 10 µm, respectively). To

visualise directed movement of signals (distance over time) within a series of images,

kymographs were generated by plotting the position of signals along a defined path (x-axis)

for each frame of the corresponding video (y-axis). Bidirectional movement is visible as

diagonal lines (yellow arrowheads; N, nucleus; Supplemental Video 1). For an example image

of a complete AB33 hypha, see Fig EV1A.

(**B**) Average velocity of fluorescent signals per hypha for strains shown in A (movement of

tracks with > 5 µm were scored as processive). Data points represent averages from three

independent experiments (n=3), with mean, red line, and s.e.m.. At least 20 signals/hyphae

were analysed out of 12 hyphae per experiment (ns; p=0.18 and p=0.23) using an paired two-

tailed Student's t-test on the means.

(**C**) Kymographs of hyphae of AB33 derivatives (6 h.p.i.) expressing pairs of red and green

fluorescent proteins as indicated. Fluorescence signals were detected simultaneously using

dual-view technology (arrow length as in A). Processive co-localising signals are marked by

yellow arrowheads (Supplemental Videos 2-4).

(**D**) Percentage of red fluorescent signals exhibiting co-localisation with the green fluorescent

signal for strains shown in (C). Data points represent observed co-localisation in three

independent experiments, mean, dark grey line, and s.e.m. (n=3, 11 hyphae each; paired two-

tailed Student's t-test on the means; ns; p=0.63 and p=0.5).

(**E**) Kymographs comparing hyphae of AB33 derivatives (6 h.p.i.) expressing Rrm4-Gfp in

the wild type (left) or *grp1Δ* strains (right) (processive signals marked by yellow arrowheads;

arrow length on the left and bottom indicate time and distance, 10 s and 10 µm, respectively;

Supplemental Video 5).

(**F**) Average velocity of fluorescent signals per hyphae for strains shown in (E) (movement of

tracks with > 5 µm were scored as processive). Data points represent averages from three

independent experiments (n=3) with mean, black line, and s.e.m.. At least 20 signals/hypha

were analysed out of at least 10 hyphae per experiment (ns; p=0.27 and p=0.4) using an paired

two-tailed Student's t-test on the means.

(**G**) Kymographs comparing hyphae (6 h.p.i.) expressing Grp1-Gfp or Pab1-Gfp in wild type

(left) with *rrm4Δ* strains (right; processive signals marked by yellow arrowheads; arrow

length as in A).

(**H**) Hyphal tips (4 h.p.i.) of AB33 derivatives expressing Grp1-Gfp, Pab1-Gfp or Gfp alone

comparing wild type (top) with *rrm4Δ* strains (bottom). Fluorescence micrographs in false

colours (black/blue to red/white, low to high intensities, respectively; scale bar, 10 µm; ROI1

and ROI2-labelled circles exemplarily indicate regions-of-interest analysed in E).

(**I**) Ratio of signal intensities in strains shown in (H) comparing Gfp fluorescence at the tip

(ROI1) and in close vicinity to the nucleus (ROI2; see Materials and methods). Bars represent

mean and s.e.m. (*wt*: n=160; *rrm4Δ*: n=152), Pab1G (n=126; *rrm4Δ*: n=170), Gfp (n=152;

rrm4Δ: n=210), unpaired two-tailed Student's t-test.

(**J**) Kymographs of hyphae of AB33 derivatives (6 h.p.i.) expressing pairs of red and green

fluorescent proteins as indicated (arrow length as in A; Supplemental Videos 6-7).

Fluorescence signals were detected simultaneously using dual-view technology. Processive

co-localising signals are marked by yellow arrowheads. Note that processive movement is

completely lost in the lower panels. Only static signals, visualised as vertical lines, are

remaining.

Fig 3. Rrm4 and Grp1 bind to thousands of target transcripts.

(**A**) Autoradiograph and Western blot analyses for representative iCLIP experiments with

Rrm4-Gfp, Grp1-Gfp and Gfp. Upper part: Upon radioactive labelling of co-purified RNAs,

the protein-RNA complexes were size-separated in a denaturing polyacrylamide gel. Protein-

RNA complexes are visible as smear above the size of the protein (Rrm4-Gfp, 112 kDa; Grp1-

Gfp, 45 kDa; Gfp, 27 kDa; indicated by arrowheads on the right). Samples with and without

UV-C irradiation and RNase I (see Material and methods) are shown. Lower part:

corresponding Western blot analysis using α-Gfp antibody (arrowheads and asterisks indicate

expected protein sizes and putative degradation products, respectively).

(**B**) Summary of binding sites and target transcripts of Rrm4 and Grp1 (top). Venn diagram

(below) illustrates the overlap of Rrm4 and Grp1 target transcripts.

(**C**) iCLIP data for Rrm4 and Grp1 on *cdc3* (UMAG_10503; crosslink events per nucleotide

from two experimental replicates [light grey/light blue] and merged data [grey/blue] from

AB33 filaments, 6 h.p.i.). Track below the merged iCLIP data show binding sites for each

protein (bs, red). Note that crosslink events in the 5´ UTR and first exon of *cdc3* were not

assigned as binding sites due to low reproducibility between replicates (see Materials and

methods). RNASeq coverage from wild type AB33 filaments (6 h.p.i.) is shown for

comparison. Gene model with exon/intron structure below was extended by 300 nt on either

side to account for 5´ and 3´ UTRs (green).

(**D**) Functional categories of cellular components (FunCat annotation scheme, 78) for proteins

encoded by target transcripts that are shared between Rrm4 and Grp1 (left) or unique to Grp1

(right). P values for the enrichment of the listed category terms are depicted by colour (see

scale below).

(**E**) iCLIP data (crosslink events from merged replicates) for Rrm4 (blue) and Grp1 (grey) as

well as RNASeq coverage on selected target transcripts (*cdc10*, UMAG_10644; *cdc11*,

UMAG_03449; *cdc12*, UMAG_03599). Enlarged regions (indicated by boxes) of the 3´ UTR

(green) are shown on the right. Datasets and visualisation as in (C). Binding sites (bs) are

shown (red; orange indicates overlap with UAUG).

1012

Fig 4. Rrm4 binds target transcripts at the start and stop codon.

(**A**) Distribution of binding sites within different transcript regions: 5´ UTR, 3´ UTR, ORF

and intron. Percentage and absolute number of binding sites are given for each category. On

the left, a transcriptome-wide distribution of nucleotides per transcript region is shown for

comparison.

(**B**) Percentage of Rrm4 binding sites (bs) overlapping with Grp1 bs (by at least 1 nt) within

shared target transcripts, shown for all bs and separated into transcript regions. The total

number of binding sites per category is indicated on top.

(**C**) Positional maps of Rrm4 (top) and Grp1 (bottom) bs relative to the stop codon (position

0). Binding sites in ORFs and 3´ UTR are given in red and orange, respectively. 234 target

transcripts were randomly selected carrying an Rrm4 bs in the 3´ UTR (with > 100 Rrm4

crosslink events; out of 1,715 Rrm4/Grp1 shared targets with Rrm4 bs in 3´ UTR; Dataset

EV6). Transcripts were ordered by decreasing 3´ UTR length (see Materials and methods).

(**D**) Metaprofiles of Rrm4 (top) and Grp1 (bottom) crosslink events relative to the start and

stop codon (position 0). Note that crosslink events are substantially more frequent towards

ORF ends, reflected in different y-axis scales.

(**E**) Genome browser views of Rrm4 and Grp1 iCLIP events as well as RNASeq data of *sui1*

(UMAG _02665) and *dyn2* (UMAG_04372). Visualisation as in Fig 3C.

1031

Fig 5. Rrm4 recognises UAUG *in vivo*.

(**A**) Logo representation of the most enriched sequence motif at Rrm4 binding sites. At each

position, the height of the stack is proportional to the information content, while the relative

height of each nucleotide within the stack represents its relative frequency at this position.

(**B**) Frequency of the Rrm4 motif UAUG around Rrm4 and Grp1 binding sites. Shown is the

percentage of binding sites that harbour an UAUG starting at a given position in an 81-nt

window around the binding site summit.

(**C**) Box plot comparing relative binding strength at Rrm4 binding sites (signal-over-

background, SOB; see Materials and methods) with or without UAUG (n=2,201 and n=4,211,

respectively; unpaired Student's t-test). Box limits represent quartiles, centre lines denote 50th

percentiles, and whiskers extend to most extreme values within 1.5x interquartile range.

(**D**) Schematic representation of the yeast three-hybrid system: *lexA* operator (*lexA op*)

sequences are bound by the LexA-MS2 coat protein (CP) hybrid (grey), recruiting the MS2-

SELEX-RNA hybrid (black and red, respectively) to the promoter region of the *HIS3* reporter

gene. Transcription is activated by binding of the third hybrid AD-Rrm4-Gfp (green) carrying

a Gal4 activation domain (AD).

(**E**) RNA structure prediction of aptamer SELEX-A1 with UAUG (red) or the mutated version

mUAUG containing UCUC(A) (mutated bases in green).

(**F**) Colony growth on control and selection plates of yeast cells expressing protein and RNA

hybrids indicated on the right. RNA binding is scored by growth on selection plates (SC -his

+3-AT, 3-amino-1,2,4-triazole). mRRMx, Rrm4 variants harbouring mutations in RRM1, 2, 3

or 1 and 2.

(**G**) Percentage of Rrm4 bs containing the motif UAUG, shown for all bs and separated into

transcript regions. Total number of binding sites is indicated on top.

(**H**) Relative contribution of the three stop codon variants to all (left) and Rrm4-bound stop

codons (left). Although opal stop codons (UGA) fit best with a UAUG-containing binding site

(UA<u>UGA</u>; present at 32 out of 63 bound UGA stop codons, 51%), they are depleted from

Rrm4-bound stop codons.

1060

Fig 6. Model of Rrm4/Grp1-mediated endosomal mRNA transport

(**A**) Schematic drawing of target transcript with Rrm4-bound regions on top (5´ cap structure,

blue circle; 5´ and 3´ UTR, light blue; ORF, dark blue; poly(A) tail, AAAA). Different

categories of Rrm4 target transcripts were defined according to the presence of Rrm4 binding

sites at the start codon, within the ORF, at the stop codon and in the 3´ UTR. Approximate

number of target transcripts and selected examples are given for each category. About 900 of

1,300 target transcripts with an Rrm4 binding site in the ORF harbour an UAUG motif within

the ORF binding site. Potential RRM domains of Rrm4 and Grp1 that may mediate RNA

binding in the different transcript regions are given in green and magenta, respectively.

(**B**) Simplified model proposing the spatial arrangement of endosomal RBPs with bound

target transcripts. The three RRM domains of Rrm4 are schematically displayed and labelled

by numbers (FYVE zinc finger domain; PI3P, phosphatidylinositol 3-phosphate; M,

MademoiseLLE domain; P and PL, PAM2 and PAM2-like sequence, respectively; further

details, see text). Note that complete endosomal mRNPs will consist of additional mRNA and

protein components forming more complex, higher-order structures.

1076

Expanded View Figure Legends

Figure EV1. Loss of Grp1 causes defects in cell growth.

(**A**) Hyphal form (6 hours post induction, h.p.i.) of laboratory strain AB33 expressing a Gfp-

tagged protein with nuclear localisation signal to stain the nucleus (N; λN-NLS-Gfp, phage

protein λN fused to triple Gfp, containing a nuclear localisation signal; inverted fluorescence

image shown; scale bar, 10 μm). Hyphae expand at the apical pole (arrow) and insert septa

(asterisks) at the basal pole in regular time intervals resulting in the formation of empty

sections.

(**B**) Schematic representation of the domain architecture of four small glycine-rich proteins

(RRM, RNA recognition motif, green; GQ/R, glycine-rich region with arginine or glutamine,

red). *Um*Grp1 from *U. maydis* (UMAG_02412), *At*GRP7 from *Arabidopsis thaliana* (RBG7;

NC_003071.7), *Hs*RBM3 and *Hs*CIRBP from *Homo sapiens* (NC_000023.11 and

NC_000019.10, respectively). Number of amino acids indicated on the right.

(**C**) Results of preliminary affinity purification experiments using Rrm4-GfpTT as bait (see

Materials and methods). Proteins with a functional link to Rrm4 are marked in red (this study)

[18, 64]. Peptide count: number of identified peptides corresponding to predicted protein;

total peptide score: sum of all peptide scores corresponding to predicted protein, excluding the

scores of duplicate matches; best peptide score: best score from all identified peptides

corresponding to predicted protein. Note that the difference between total peptide score and

best peptide score is a correction of the software depending on how many possible predicted

candidates match to the identified peptide mass.

(**D**) Tandem affinity purification using Rrm4-GfpTT as bait. Protein bands were stained with

Coomassie Blue after SDS-PAGE. Proteins in boxed areas were identified as Rrm4 and Grp1

(size of marker proteins in kDa on the right).

(**E**) Growth curve of indicated AB33 derivatives growing in liquid culture. Data points

represent averages from three independent experiments (n=3). Error bars show s.d. (**F**)

Differential interference contrast (DIC) images of AB33 derivatives as yeast-like budding

cells (scale bar, 10 µm).

(**G**) Length of budding cells (shown are merged data from three independent experiments,

n=3; > 100 cells per strain were analysed, *wt*, 269; *rrm4Δ*, 122 (only two independent

experiments); *grp1Δ*, 263; *grp1G*, 318), overlaid with the mean of means, red line, and s.e.m.;

paired two-tailed Student's t-test on the mean cell lengths from the replicate experiments.

(**H**) Colonies of indicated AB33 strains grown in the yeast form incubated at different

temperatures (28°C for 1 d or 16°C and 21°C for 5 d).

(**I**) Colonies of indicated AB33 strains grown in the yeast form. Incubated plates contained

cell wall inhibitors (CM, complete medium for 1 d; CFW, 50 µM Calcofluor White for 4 d;

CR, 57.4 µM Congo Red for 4 d).

(**J**) Fluorescence images of the basal pole of hyphae of AB33 derivatives (6 h.p.i.). Septa

(asterisks) were stained with CFW. White bars indicate exemplary length measurements of

empty sections shown in Fig 1D (scale bar, 10 µm).

1117

Figure EV2 Improving the iCLIP protocol for fungal RBPs.

(**A**) Grp1-Gfp/RNA complexes were size-separated on denaturing PAGE after UV-C

irradiation and transferred to a nitrocellulose membrane (left). RNA was radioactively

labelled, and protein-RNA complexes with covalently linked RNAs of different sizes were

visible as smear above the expected molecular weight of the Grp1-Gfp protein (45 kDa;

marked by arrowhead). RNA of four different regions of the membrane (A to D indicated on

the right) were isolated from the membrane and size separated on a denaturing gel (6%)

(right; nucleotide size marker on the left, bp).

(**B**) Autoradiographs showing Rrm4-Gfp, Grp1-Gfp and Gfp in complex with RNA after UV-

C irradiation at 0, 160, 320, 480 and 640 mJ/cm^2. Corresponding Western blots using anti-Gfp

are shown below. Arrowheads indicate the expected molecular weight of the proteins (Rrm4-

Gfp, 112 kDa; Grp1-Gfp, 45 kDa; Gfp, 27 kDa). After each irradiation step, the cells were

mixed. Note that increased UV-C irradiation in combination with slow processing due to long

time intervals was particularly harmful for the Rrm4 protein, which was completely degraded

after four minutes of UV-C irradiation. Putative degradation products are marked by asterisks.

(**C**) Autoradiographs showing Rrm4-Gfp, Grp1-Gfp and Gfp in complex with RNA after

single UV-C irradiation at 0, 100, 200, 300 or 400 mJ/cm². This time, mixing breaks were

omitted and cells were harvested as quickly as possible. Corresponding Western blots are

shown below. Labelling as above. We chose 200 mJ/cm^2 as optimal UV-C irradiation dose,

since the amount of unspecific Gfp-RNA complexes increased at higher doses. Arrowheads

and asterisks as in (B).

(**D**) Amplification of the Rrm4, Grp1 and Gfp derived cDNA libraries with different numbers

of PCR cycles (between 18 and 24; ctrl, control without template cDNA). The PCR products

were separated on a native gel (6%) and stained with SYBR green I (nucleotide size marker

on the left, bp). The size of the cDNA insert together with the adapters (cDNA insert = 20-30

nt; L3 adapter, RT-primer and P3/P5 Solexa primers = 128 nt) is expected to be ~ 150-160 nt

after amplification.

1145

Figure EV3. Comparative iCLIP procedure results in high-quality dataset.

(**A**) Summary of the iCLIP libraries including initial number of sequencing reads, uniquely

mapped reads, crosslink events (xlinks) for both replicates for Rrm4, Grp1 and Gfp. In

addition, sum of crosslink events as well as resulting binding sites and target transcripts are

given for merged replicates. Gfp binding sites were only filtered for reproducibility but not for

relative signal intensity (SOB; see Material and methods).

(**B**) Stacked bar chart showing percentage of reads mapping to a unique, multiple (multiple

mapping) or no location (unmapped) in the *U. maydis* genome for Rrm4, Grp1 and Gfp.

(**C**) Scatter plot comparing number of crosslink events per gene from two independent

replicate experiments for Rrm4, Grp1 and Gfp (PCC, Pearson correlation coefficient).

(**D**) Genome browser views of Rrm4 and Grp1 iCLIP events as well as RNASeq data of *cts1*

(UMAG_10419) and *rrm4* (UMAG_10836). Visualisation as in Fig 3C.

(**E**) Venn diagram to identify target transcripts that are uniquely bound by Rrm4 (left) or Grp1

(right). Unique target transcripts (numbers given in bold) are selected only if they show no

evidence of binding by the other RBP (considering all binding sites before SOB filtering, see

Materials and methods).

1162

Figure EV4. Accumulation of crosslink events at stop codons of mRNAs encoding

subunits of the mitochondrial F_OF_1-ATPase.

(**A**) Heatmap of crosslink events of Grp1 (left) and Rrm4 (right) in a window around the stop

codons (position 0 = first position of 3´ UTR) of mRNAs encoding subunits of the

mitochondrial F_OF_1-ATPase (nomenclature and gene identifiers for *U. maydis* on the right).

Crosslink events per nucleotide are represented by a colour scale (right).

(**B, C**) iCLIP data of Rrm4 and Grp1 as well as RNASeq data across selected mRNAs of the

F_1 subcomplex (**B**) and F_O subcomplex (**C**) that carry an Rrm4 binding site precisely at the

stop codon. Visualisation as in Fig 3C.

(**D**) Cy2 image of a representative differential gel electrophoresis (DIGE) analysis comparing

membrane-associated proteins of wild type (blue, Cy5-labelled) and *rrm4Δ* hyphae (red, Cy3-

labelled; size marker on the left, pH range at the top; 6 h.p.i.). Protein variants exhibiting at

least 2.5-fold differences in protein amounts are indicated by numbered arrowheads (taken

from earlier study) [18]. The area marked with a yellow rectangle is enlarged on the right. The

corresponding standardised logarithmic protein abundances for the Atp4 spot obtained from

three biological replicates are given on the right (internal standard set to 0).

Figure EV5. Control experiments for the yeast three-hybrid analysis.

(**A**) Logo representation of the most enriched sequence motif at Grp1 binding sites. At each

position, the height of the stack is proportional to the information content, while the relative

height of each nucleotide within the stack represents its relative frequency at this position.

(**B**) Frequency of UGUA around Rrm4 and Grp1 binding sites. Shown is the percentage of

binding sites that harbour an UGUA starting at a given position in an 81-nt window around

the binding site summit. Representation as in Fig 5B.

(**C**) Colony growth on control and selection plates of yeast cells expressing protein and RNA

hybrids indicated on the right. RNA binding is scored by growth on selection plates (SC –his

+3-AT, 3-amino-1,2,4-triazole). This control experiment demonstrates that growth on

selection plates (see Fig 5F) depends the presence of Rrm4 variant and cognate hybrid RNA.

mRRMx, Rrm4 variants harbouring mutations in RRM 1, 2, 3 or 1 and 2.

(**D**) Relative occurrence of NAUG sequence context for all (left) and Rrm4-bound (right) start

codons. CAUG fits to the Kozak sequence in eukaryotes [81]. The fraction of start codons

coinciding with the Rrm4 recognition motif UAUG is shown in blue. This sequence context

was strongly enriched among the Rrm4-bound target mRNAs, whereas it comprises only 8%

of all annotated start codons in the *U. maydis* genome.

(**E**) UAUG motifs at start codons were not more frequently bound than UAUG motifs in the

surrounding sequence (see Materials and methods). Out of a total of 14,748 UAUG motifs in

expressed transcripts 15.2% are bound by Rrm4. Similarly, 14.5% of the 282 UAUG motifs

directly at start codons are bound by Rrm4. This is only marginally more than at 215 and 483

UAUG motifs within 100 nt upstream and downstream of the start codons, out of which

13.0% and 11.0% are bound, respectively.

A

```
                                    |──────────────────────────────RRM──────────────────────────────|  |
Grp1   MAA-----KVYVGNLSWNTTDDSLAHAFSTYGQLTDYIVMKDRETGRSRGFGFVTFATQAEADAAIAALNEQELDGRIRVNMANSRPAG
GRP7   MASGDVEYRCFVGGLAWATDDRALETAFAQYGDVIDSKIIMDRETQRSRGFGFVTFKDEKAMKDAIEGMNGQDLDGRSITVNEAQSRGSG
RBM3   MSSE--EGKLFVGGLNFNTDEQALEDHFSSFGPISEVVVVKDRETQRSRGFGFVTFTNPEHASVAMRAMNGESLDGRQIRVDHAGKSARG
CIRBP  MASD--EGKLFVGGLSFDTNEQSLEQVFSKYGQISEVVVVKDRETQRSRGFGFVTFENIDDAKDAMMAMNGKSLDGRIRVDQAGKSSDN
                RNP2                                         RNP1
```

```
        |────────────────────────────GQ/R-rich────────────────────────────|
Grp1   GMGGGYGGVTGQYGANAYGAQGGYGGYGGQPGGFQQPGGFQQGGYPQQGGYGGYQQPGFQPQQGGYGAPQQGYGAPQQGGYGGYNGQSQ
GRP7   GGGGHRGGGGGGYRSGGGGGYSGGGGSYGGGGGRREGGGGYSGGGGGYSSRGGGGGSYGGGRREGGGGYGGGEGGGYGGSGGGGGW---
RBM3   TRGGGFGAHGRGRSYSRGGGDQGYGSGRYYDSRPGGYGYGYGRSRDYNGRNQGGYDRYSGGNYRDNYDN-------------------
CIRBP  RSRGYRGGSAGGRGFFRGGRGRGRGFSRGGGDRGYGGNRFESRSGGYGGSRDYYSSRSQGGYSDRSSGGSYRDSYDSYATHNE------
```

B

wt

grp1Δ

rrm4Δ

grp1-gfp

C

length of hyphae (μm)

- with septa
- without septa

2 h.p.i.   4 h.p.i.   6 h.p.i.

wt grp1Δ rrm4Δ grp1-gfp

p=0.02   p=0.01   p=0.02

D

empty sections (μm)

ns

p=0.02

wt grp1Δ grp1-gfp

E

grp1-gfp   *grp1Δ*

stressed   stressed

F

normal hyphae (%)

- unstressed
- stressed

p=0.007

grp1-gfp grp1Δ rrm4-gfp rrm4Δ

B

	RRM	GQ/R-rich	
*Um*Grp1			175
*At*GRP7			176
*Hs*RBM3			157
*Hs*CIRBP			172

C

Nr.	UMAG	Annotation	MW[a]	PC[b]	Total PS[c]	Best PS[d]
1	10836	Rrm4	85	24	3086,8	252,7
2	11918	conserved hypothetical protein	95	4	582,2	196,6
3	10419	chitinase Cts1	55	5	513,1	130,3
4	06140	related to programmed cell death protein (calcium-binding protein)	34	4	457,7	182,1
5	05365	probable protein phosphatase 2A regulatory B subunit	53	3	322,4	124,2
6	10526	probable SSB2 - heat shock protein of HSP70 family, cytosolic	68	2	144,6	98,8
7	04080	probable DED1 - ATP-dependent RNA helicase	71	1	87,1	89,6
8	05158	related to Nucleoporin nup189 (SonB)	229	1	76,1	80,1
9	10350	Upa2	232	2	75,2	47,3
10	03791	Ums2heat shock 70 kd protein 2	70	1	71,4	73,9
11	02412	Grp1, related to CIRBP - glycine-rich RNA-binding protein	18	1	68,8	71,3
12	11551	probable 40S small subunit ribosomal protein S19	18	1	66,9	66,9
13	04619	related to HSE1, binds ubiquitin, mediates endosomal protein sorting	66	1	61,7	64,3
14	11211	Cutinase	51	1	59,3	63,3
15	01227	hypothetical protein	88	2	57,3	40,4
16	11719	probable ribosomal protein S4.e, cytosolic	30	1	56,9	59,4
17	06404	probable thioredoxin peroxidase	25	1	54,0	54,0
18	10163	probable 60S large subunit ribosomal protein L20	20	1	53,2	55,7
19	03738	probable ATP dependent RNA helicase	250	2	51,7	32,2
20	01077	putative protein	76	1	50,9	55,9
21	02578	related to Het-c heterokaryon incompatibility protein	108	2	50,0	31,9
22	01434	Fer3 siderophore peptide synthetase, ferrichromeA biosynthesis	528	2	46,8	31,3
23	01702	conserved hypothetical protein	20	1	46,2	46,2

[a] molecular weight [kDa]; [b] peptide count; [c] total peptide score; [d] best peptide score

A

Grp1

B

position relative to
binding site summit (nt)

C

D

start codons
n=6,734

Rrm4 at start
n=47

■ UAUG ■ CAUG ■ AAUG ■ GAUG

E