Alcohol consumption and mate choice in UK Biobank: comparing

observational and Mendelian randomization estimates

Authors: Laurence J Howe*, [1, 3] Daniel J Lawson, [1] Neil M Davies, [1] Beate St.

Pourcain, [2] Sarah J Lewis, [1] George Davey Smith, [1, #] and Gibran Hemani [1, #]

[1] Medical Research Council Integrative Epidemiology Unit, Population Health
Sciences, University of Bristol, United Kingdom

[2] Max Planck Institute for Psycholinguistics, Nijmegen, Netherlands

[3] Institute of Cardiovascular Science, University College London, United Kingdom

* Correspondence to lh14833@bristol.ac.uk

# These authors contributed equally

Abstract

Alcohol use is correlated within spouse-pairs, but it is difficult to disentangle

the effects of alcohol consumption on mate-selection from social factors or

cohabitation leading to spouses becoming more similar over time. We hypothesised

that genetic variants related to alcohol consumption may, via their effect on alcohol

behaviour, influence mate selection.

Therefore, in a sample of over 47,000 spouse-pairs in the UK Biobank we

utilised a well-characterised alcohol related variant, rs1229984 in *ADH1B*, as a

genetic proxy for alcohol use. We compared the phenotypic concordance between

spouses for self-reported alcohol use with the association between an individual's

self-reported alcohol use and their partner's rs1229984 genotype using Mendelian

randomization. This was followed up by an exploration of the spousal genotypic

concordance for the variant and an analysis determining if relationship length may be

related to spousal alcohol behaviour similarities.

We found strong evidence that both an individual's self-reported alcohol

consumption and rs1229984 genotype are associated with their partner's self-

reported alcohol use. The Mendelian randomization analysis found that each unit

increase in an individual's weekly alcohol consumption increased their partner's

alcohol consumption by 0.26 units (95% C.I. 0.15, 0.38; $P=1.10 \times 10^{-5}$). Furthermore,

the rs1229984 genotype was concordant within spouse-pairs, suggesting that some

spousal concordance for alcohol consumption existed prior to cohabitation. Although

the SNP is strongly associated with ancestry, our results suggest that this

concordance is unlikely to be explained by population stratification. Overall, our

findings suggest that alcohol behaviour directly influences mate selection.

Introduction

Human mate choice is highly non-random; spouse-pairs are generally more

phenotypically similar than would be expected by chance [1-6]. Previous studies

suggest that alcohol related phenotypes, ranging from consumption to alcohol

dependence, are highly correlated within spouse-pairs [7-13]. However, the extent to

which the spousal concordance is due to the effect of alcohol behaviour on mate

selection (assortative mating) is currently unclear. Indeed, the spousal concordance

may be related to assortment on other social and environmental factors (social

homogamy) or be a consequence of an individual's partner influencing their alcohol

behaviour after the individuals have paired up (partner interaction effects) or even

relate to spousal similarities influencing relationship length (relationship dissolution)

[11-13]. The mechanism explaining spousal concordance for alcohol consumption could

have important implications relating to human social and reproductive behaviour.

Figure 1 illustrates possible explanations for spousal concordance on alcohol

consumption.

One biological mechanism that partially explains the phenotypic concordance

between spouse-pairs is that they are on average more genetically similar across the

genome than non-spouse-pairs [14]. Genotypes implicated in the aetiology of height,

education, blood pressure and several chronic diseases have been shown to be

correlated within spouse-pairs [15-18]. It is not known whether genetic variants

implicated in alcohol metabolism, via their effect on alcohol behaviour, contribute to

mate selection.

Alcohol behaviour has been shown to be highly heritable with estimates of

30-50% for alcohol use disorders [19 20] and a common variant heritability of 13% for

self-reported alcohol consumption [21]; Genome-wide Association Studies (GWAS)

have identified more than 15 loci implicated in either the aetiology of alcohol

dependence [22-26] or alcohol consumption volume [21 24 27-29]. Notably, genetic variants

in the Alcohol Dehydrogenase (*ADH*) and Aldehyde Dehydrogenase (*ALDH*) gene

families are associated with differences in alcohol consumption [30]. For example,

ADH1B is involved in the production of enzymes that oxidise alcohol and so

individuals with certain alleles may find alcohol consumption unpleasant, resulting in

lower intake. Similarly, a genetic variant in *ALDH2*, rare in non-east Asian

populations, is associated with a "flush reaction" to alcohol [31 32].

Alcohol consumption-related genetic variants can be useful to determine the

most likely explanation for the spousal phenotypic concordance for alcohol use, by

analogy with Mendelian randomization studies [33 34]. Genetic variants for alcohol

consumption are in theory less susceptible to confounding from socioeconomic and

behavioural factors than measured alcohol consumption so can be used to rule out

the possibility that social homogamy is driving the spousal phenotypic concordance

[33 35]. The timing of the effects of alcohol consumption can be discerned by evaluating

the spousal genotypic concordance for alcohol use-related variants. Genotypic

concordance would imply that an effect exists prior to pairing, suggesting that some

degree of the spousal phenotypic concordance is attributable to assortative mating

(**Figure 2**).

In this study we aimed to explore spousal similarities for alcohol consumption

using observational and genetic data. First, we estimated the association of an

individual's self-reported alcohol use with the self-reported alcohol use of their

partner. Second, we used a Mendelian randomization framework to estimate the

effect of an individual's alcohol use on their spouse's alcohol use. Here, we used

their partner's rs1229984 genotype, a missense mutation in *ADH1B* strongly

associated with alcohol consumption as an instrumental variable for self-reported

alcohol consumption. Third, we estimated the association of rs1229984 genotype

between spouses, to evaluate the timing of possible causal effects, and investigate

the possibility of bias from population stratification. Fourth, using the mean age of

each couple as a proxy for relationship duration, we determined if there was an

association between longer relationships and more similar spousal alcohol

behaviour. As a positive control, to demonstrate the validity of derived spouse pairs

and the usage of a Mendelian randomization framework, we also analysed height,

known to be correlated between spouses, using similar methods.

107   **Materials and Methods**

Study participants

UK Biobank

UK Biobank is a large-scale cohort study, including 502,655 participants aged

between 40-69 years. Study participants were recruited from 22 recruitment centres

across the United Kingdom between 2006 and 2010 [36] [37]. For the purposes of our

analyses, we restricted the dataset to a subset of 463,827 individuals of recent

European descent with available genotype data, with individuals of non-European

descent removed based on a k-means cluster analysis on the first 4 genetic principal

components [38]. The different subsets of UK Biobank utilised in our analyses are

illustrated in **Supplementary Figure 1**.

Spouse-pair subsample

Spouse information is not explicitly available, therefore we used similar

methods to previous studies [15-17] to identify spouse-pairs in the UK Biobank. Starting

with the European subsample described above, household sharing information was

used to extract pairs of individuals who (a) report living with their spouse (6141-0.0),

(b) report the same length of time living in the house (699-0.0), (c) report the same

number of occupants in the household (709-0.0), (d) report the same number of

vehicles (728-0.0), (e) report the same accommodation type and rental status (670-

0.0, 680-0.0), (f) have identical home coordinates (rounded to the nearest km)

(20074-0.0, 20075-0.0), (g) are registered with the same UK Biobank recruitment

centre (54-0.0) and (h) both have available genotype data. If more than two

individuals shared identical information across all variables, these individuals were

excluded from analysis. At this stage, we identified 52,471 potential spouse-pairs.

We excluded 4,866 potential couples who were the same sex (9.3% of the

sample), as unconfirmed same sex pairs may be more likely to be false positives.

Although sexual orientation data was collected in UK Biobank, access is restricted

for privacy/ethical reasons. To reduce the possibility that identified spouse-pairs are

in fact related or non-related familial, non-spouse pairs; we removed 3 pairs

reporting the same age of death for both parents (1807-0.0, 3526-0.0). Then we

constructed a genetic relationship matrix (GRM) amongst derived pairs and removed

53 pairs with estimated relatedness (IBD >0.1). To construct the GRM; we used a

pool of 78,341 markers which were derived by LD pruning (50KB, steps of 5 KB,

r2<0.1) 1,440,616 SNPs from the HapMap3 reference panel [39] using the 1000

Genomes CEU genotype data [40] as a reference panel. The final-sample included

47,549 spouse-pairs.

Non-spouse-pair samples

For secondary analyses requiring data from unrelated individuals, we derived

a sample of individuals of European descent and a more restrictive sample believed

to be of white British descent. Starting with the UK Biobank subset of 463,827

individuals of recent European descent, we removed 78,540 related individuals

(relevant methodology has been described previously [38]) to generate the European

sample and using lists provided by UK Biobank, further restricted this sample to

337,114 individuals identifying as being of "white British" descent.

Height and educational attainment

At baseline, the height (cm) of UK Biobank participants was measured using a

Seca 202 device at the assessment centre (ID: 50-0.0). Measured height was used

as a positive control for the application of a Mendelian randomization framework in

the context of assortative mating.

Educational attainment as characterised by years in full-time education was

defined as in a previous publication [41]. Individuals born outside England, Scotland or

Wales were removed because of schooling system differences, participants with a

college or university degree were classified with a leaving age of 21 years and

participants who self-reported leaving school when younger than 15 years were

classified with a leaving age of 15. Educational attainment was included as a

covariate in phenotypic analyses of spousal alcohol behaviour similarities as a

possible confounder.

Self-reported alcohol variables

At baseline, study participants completed a questionnaire. Participants were

asked to describe their current drinking status (never, previous, current, prefer not to

say) (ID: 20117-0.0) and estimate their current alcohol intake frequency (daily or

almost daily, three or four times a week, once or twice a week, one to three times a

month, special occasions only, never, prefer not to say) (ID: 1558-0.0). Individuals

reporting a current intake frequency of at least "once or twice a week" were asked to

estimate their average weekly intake of a range of different alcoholic beverages (red

wine, white wine, champagne, beer, cider, spirits, fortified wine) (ID: 1568-0.0, 1578-

0.0, 1588-0.0, 1598-0.0, 1608-0.0).

From these variables, we derived three measures: ever or never consumed

alcohol (current or former against never), a binary measure of current drinking for

self-reported current drinkers (three or more times a week against less than three

times a week) and an average intake of alcoholic units per week, derived by

combining the self-reported estimated intakes of the different alcoholic beverages

consumptions across the five drink types, as in a previous study [21]. The

questionnaire used the following measurement units for each of the five alcoholic

drink types: measures for spirits, glasses for wines and pints for beer/cider which

were estimated to be equivalent to 1, 2 and 2.5 units respectively. Individuals

reporting current intake frequency of "one to three times a month", "special

occasions only" or "never" (for whom this phenotype was not collected), were

assumed to have a weekly alcohol consumption volume of 0. More information on

alcohol variables used in this study is contained in **Supplementary Table 1**.

Genotyping

488,377 UK Biobank study participants were assayed using two similar

genotyping arrays, the UK BiLEVE Axiom™ Array by Affymetrix1 (N= 49,950) and

the closely-related UK Biobank Axiom™ Array (N= 438,427). Directly genotyped

variants were pre-phased using SHAPEIT3 [42] and then imputed using Impute4 using

the UK10K [43], Haplotype Reference Consortium [44] and 1000 Genomes Phase 3 [40]

reference panels. Post-imputation, data were available on approximately ~96 million

genetic variants.

Statistical analysis

Utilising genetic variation to disentangle spousal correlations

In general, the effects of genetic variation on a phenotype can be assumed to

be via the variant's effect on intermediary observable or unobservable phenotypes.

In the context of assortative mating, it is unlikely that individuals would assort based

directly on genotype but rather on an observed phenotype influenced by genetic

factors. Assuming that a phenotype is influenced by genetic factors G and individuals

assort on the phenotype such that the phenotypic correlation between spouses is

equal to C, then expected correlations between an index individual's G and their

partner's phenotype and G induced by assortment can be shown to be a function of

the heritability of the phenotype and the spousal phenotypic correlation C

(**Supplementary Methods**). This implies that estimates of assortative mating

utilising genetic data are likely to be attenuated compared to the true value of

phenotypic assortment, unless genetic factors completely explain variation in the

phenotype of interest.

However, there are notable advantages of applying genetic approaches such

as Mendelian randomization and genetic correlation analyses to the context of

assortative mating for mechanistic understanding. In conventional Mendelian

randomization studies [33][34], genetic variants are used as proxies for a measured

exposure to evaluate potential causal relationships between an exposure and an

outcome (e.g. LDL cholesterol and coronary heart disease [45]). Genetic proxies may

be more reliable than the measured exposure because of the reduced potential for

confounding and reverse causation.

In the context of Mendelian randomization across spouses, the premise is

largely similar; the exposure is an individual's phenotype (e.g. alcohol consumption),

proxied by a genetic instrument, and the outcome is their partner's phenotype (e.g.

alcohol consumption). A Mendelian randomization approach can evaluate a direct

effect of an individual's alcohol consumption on the alcohol consumption of their

partner as opposed to effects of social homogamy. A direct effect captured by a

Mendelian randomization framework could capture; individuals being likely to select

a mate with similar behaviour (assortative mating), an individual's alcohol

consumption influencing their partner's during the relationship (partner interaction

effects) or more similar couples staying together for longer (relationship dissolution).

Interpretation can be nuanced, as for example, it seems unlikely that an individual's

height could influence the height of their partner, but partner interaction effects are

highly plausible for alcohol behaviour.

Similarly, estimating the genotypic concordance between-spouses for variants

relating to a trait of interest can be used to improve mechanistic understanding. The

interpretation of genotypic concordance is comparable to that of Mendelian

randomization across spouses with two important distinctions. First, genotypic

concordance will not capture partner interaction effects as germline DNA is fixed for

both spouses prior to assortment. Second, concordance induced by assortment will

be further attenuated compared to a Mendelian randomization approach.

Phenotypic spousal concordance for height

To verify the validity of the derived spouse-pair sample, we evaluated the

spousal phenotypic concordance for height. Previous studies have found strong

evidence of spousal concordance for height, so comparable results would be

consistent with derived spouses being genuine. The spousal phenotypic

concordance was estimated using a linear regression of an individual's height

against the height of their partner, adjusting for sex. With one unique phenotype

pairing within couples (male spouse height/ female spouse height), each individual in

the data-set was included only once as either the reference individual or their

partner.

Mendelian randomization: Genetically influenced height and measured height of

partner

We validated the application of a Mendelian randomization approach to

assortative mating using height as a positive control; genotypes influencing height

have previously demonstrated to be highly correlated between spouse-pairs [15]. As a

measure of genetically influenced height, we started with 382 independent SNPs,

generated using LD clumping ($r^2 < 0.001$) in MR-Base [46], from a recent Genome-wide

Association Study (GWAS) of adult height in Europeans [47].

For the purposes of the Mendelian randomization analysis, we restricted

analyses to spouse-pairs with complete measured height data and genotype data.

First, we estimated the association between 378 SNPs (4 SNPs were unavailable in

the QC version of the data-set) and height in the same individual, using the spouse-

pair sample with sex included as a covariate. Second, we estimated the association

between the 378 SNPs and spousal height. PLINK [48] was used to estimate the SNP-

phenotype associations also including sex as a covariate. We then estimated the

effect of a 1 cm increase in an individual's height on their partner's height using the

TwoSampleMR R package [46] and the internally derived weights described above.

The fixed-effects Inverse-Variance Weighted (IVW) method was used as the primary

analysis. Cochran's Q test and the I^2 statistic were used to test for heterogeneity in

the fixed-effects IVW [49]. MR Egger [50] was used to test for directional pleiotropy. The

weighted median [51] and mode [52] were used to test the consistency of the effect

estimate. With two unique pairings between genotype and phenotype in each couple

(male spouse genotype/ female spouse height and the converse), each individual in

the data-set was included twice as both the reference individual and as the partner.

Spousal genetic concordance for height

To evaluate spousal genotypic concordance for height, we evaluated the

association between height genetic risk scores (GRS) across spouse-pairs. Height

GRS were constructed using previously described height loci in PLINK [48]. The cross-

spouse association was estimated using linear regression of an individual's GRS

against the GRS of their partner. With one unique genotype pairing within couples

(male spouse genotype/female spouse genotype), each individual in the dataset was

included only once as either the reference individual or their partner.

Phenotypic spousal concordance for self-reported alcohol behaviour

To evaluate the phenotypic concordance on alcohol use we compared self-

reported alcohol behaviour between spouses. We estimated the spousal

concordance for the two binary measures (ever or never consumed alcohol, three or

more times a week) using a logistic regression of the relevant variable for an

individual against the relevant variable for their partner, adjusting for sex, age and

partner's age. In addition, we included recruitment centre, height and education (of

both spouses) in the model as potential confounders. Similarly, linear regression was

used to estimate the spousal-concordance for continuous weekly alcohol

consumption volume, adjusting for the same covariates. Spouse-pairs with any

missing phenotype data, or where one or more spouses reported their weekly

alcohol consumption volume to be more than five standard deviations away from the

mean (calculated using the sample of individuals with non-zero weekly drinking)

were removed from relevant analyses. With one unique phenotype pairing within

couples (male alcohol variable/ female alcohol variable), each individual in the data-

set was included only once as either the reference individual or their partner.

Mendelian randomization: Genetically influenced alcohol consumption volume and

self-reported alcohol consumption of partner

We then applied the Mendelian randomization framework to investigate if an

individual's genotype at rs1229984 in *ADH1B* affects the self-reported alcohol

consumption volume of their partner. Given the rarity of individuals homozygous for

the minor allele in European populations, the MAF is 2.9% in the 1000 Genomes

CEU population [40], we first determined whether an additive or a dominant model (as

used in previous studies [45 53]) was most appropriate for the SNP by comparing the

association of genotype at rs1229984 with self-reported weekly alcohol consumption

in the European and British samples. We found strong evidence to suggest that the

SNP has an additive effect on alcohol consumption (**Supplementary Table 2**) and

assumed this model in all relevant analyses.

For the Mendelian randomization analysis, we restricted analysis to spouse-

pairs where both members had genotype data, and one or more members had self-

reported alcohol consumption volume. First, we estimated the association of the

rs1229984 genotype with alcohol consumption in the same individual after adjusting

for sex, age, centre and the first 10 principal components of the reference individual.

Second, we estimated the association between rs1229984 and spousal alcohol

consumption after adjusting for sex, age (of both spouses), centre and the first 10

principal components of both spouses. PLINK [48] was used to estimate the SNP-

phenotype associations. We then estimated the effect of a 1 unit increase in an

individual's weekly alcohol consumption volume on the same variable in their

partner. The Wald ratio estimate was obtained using mr_wald_ratio function in the

TwoSample MR R package [46] using internally derived weights. Sensitivity analyses

were limited due to the use of a single genetic instrument. With two unique pairings

between genotype and phenotype in each couple (male alcohol variable/ female

genotype and the converse), each individual in the data-set was included twice as

both the reference individual and as the partner.

Spousal genotypic concordance for rs1229984 genotype

We then investigated properties of the rs122984 variant in the UK Biobank

that may be relevant to assortative mating. Starting with the UK Biobank subset of

463,827 individuals of recent European descent, we removed 78,540 related

individuals (relevant methodology has been described previously [38]) and tested

Hardy-Weinberg Equilibrium (HWE) in the resulting sample of 385,287 individuals.

To evaluate the possibility of population stratification, we investigated the association

of both the SNP and self-reported alcohol consumption with genetic principal

components and birth coordinates. As a sensitivity analysis, we also restricted the

sample to a more homogeneous sample of British individuals, provided by the UK

Biobank, and repeated analyses.

We then estimated the genotypic concordance between derived spouse-pairs

for rs1229984 genotype using linear regression. As a sensitivity analysis, we then

investigated the possibility that spousal-concordance for rs1229984 was driven by

fine-scale assortative mating due to geography, which is itself associated with

genetic variation within the UK [54][55]. For this, we restricted the sample to include only

28,653 spouse-pairs born within 100 miles of each other. To test the validity of this

sensitivity analysis, we explored whether birth or genetic differences (as determined

by principal components) between spouses are associated with alcohol behaviour or

rs122984 genotype differences in the restricted and full spouse-pair samples. The

spouse-pairs were then stratified into the 22 different UK Biobank recruitment

centres and logistic regression analyses were re-run to estimate the spousal-

concordance of the *ADH1B* genotype by centre. With one unique genotype pairing

within couples (male genotype/female genotype), each individual in the dataset was

included only once as either the reference individual or their partner. Geographical

patterns of heterogeneity across the different UK Biobank recruitment centres would

provide evidence of population stratification.

Relationship duration and spousal alcohol behaviour

Relationship length may influence spousal similarities for alcohol behaviour

because spouses become more similar over time or because pairs with similar

alcohol behaviour tend to have longer relationships. To explore these possibilities,

we investigated the association between relationship length and alcohol behaviour

and rs122984 genotype similarities. Without available data on relationship length, we

used the mean age of each couple as a proxy and evaluated associations using a

linear regression of mean couple age against spousal difference in weekly alcohol

consumption and rs1229984 genotype. Analyses were adjusted for the sex of

reference individual.

A list of derived spouse-pairs has been returned to UK Biobank. For details please

contact access@ukbiobank.ac.uk.

Results

Spousal concordance for height

Phenotypic concordance for height

Measured height was strongly concordant between spouse-pairs. In a sample

of 47,377 spouse-pairs, a 1 unit increase in an individual's height was associated

with a 0.24-unit increase (95% C.I. 0.23, 0.25, $P<10^{-16}$) in their partner's height. This

result is consistent with previous findings [56][57], validating the derived spouse pairs.

Mendelian randomization framework: Genetically influenced height and height of

partner

The application of Mendelian randomization to spousal height was consistent

with the previous evidence for assortative mating on height. Across 47,377 spouse-

pairs, a 1 cm increase in an individual's height was associated with a 0.19 cm

increase in their partner's height (95% C.I. 0.18, 0.21; $P<10^{-16}$), distinctly smaller

than the phenotype estimate (Z-test for difference of means: $P=8.3\times10^{-8}$). The I^2

statistic (2.9%) and Cochran's Q test (P=0.64) suggested consistent effects across

SNPs, and estimates were consistent across the weighted median, weighted modal

and MR-Egger estimators with the MR-Egger intercept test finding no strong

evidence for directional pleiotropy (**Table 1**).

Table 1: Mendelian randomization estimates for the effect of a 1 cm increase in
height on partner's height

Test	Interpretation	Estimate (95% C.I.)	P-value
Phenotypic association for comparison	N/A	0.24 (0.23, 0.25)	$<10^{-16}$
Inverse variance weighted	Primary causal estimate[1]	0.19 (0.18, 0.21)	$<10^{-16}$
Heterogeneity of Inverse variance weighted	Balanced pleiotropy	I^2=3.6%	0.68
MR-Egger	Intercept test for directional pleiotropy[2]	0.001 (-0.006, 0.008)	0.75
	Regression estimate[1]	0.19 (0.15, 0.21)	$<10^{-16}$
Weighted median	Consistency[1]	0.18 (0.15, 0.21)	$<10^{-16}$
Weighted mode	Consistency[1]	0.17 (-0.23, 0.57)	0.41

1 Units: mm change in partner's height per 1-unit increase in individual's height

2 Units: Average pleiotropic effect of a height genetic variant on partner's height

Genotypic concordance for height

Similarly, the genotypic concordance analysis for height was strongly

concordant with previous findings; we found strong evidence that spouses have

similar genotypes at height influencing loci. Each 1 S.D. increase in an individual's

height GRS was associated with a 0.024 S.D. increase in their partner's GRS (95%

C.I. 0.015, 0.033; P=1.96×10^{-7}).

397 **Spousal concordance for self-reported alcohol behaviour**

398 *Phenotypic spousal concordance*

399 The majority of derived spouse-pairs had complete data for relevant self-

400 reported alcohol behaviour phenotypes. Strong evidence was found for phenotypic

401 concordance between spouse-pairs for all self-reported alcohol variables. Amongst

402 45,066 spouse-pairs, an individual self-reporting as a never-drinker was associated

403 with increased odds (OR 13.03, 95% C.I., 10.98, 15.44 $P<10^{-16}$) of their partner self-

404 reporting as a never-drinker. Similarly, when restricting to 40,723 pairs who both

405 reported being current-drinkers, an individual drinking three or more times a week

406 had increased odds (OR 6.24, 95% C.I., 5.95, 6.54 $P<10^{-16}$) of their partner also

407 drinking three or more times a week.

408 For self-reported alcohol consumption volume; 44,886 spouse-pairs had

409 either complete phenotype data or reported their consumption frequency as less

410 than weekly (in which case their weekly volume was assumed to be 0). After

411 removing 189 pairs with outlying values (>5 S.D from the mean) from one or more

412 members, the final sample included 47,321 spouse-pairs. In this sample, each unit

413 increase in an individual's weekly alcohol consumption volume was associated with

414 a 0.37-unit increase (95% C.I. 0.36, 0.38 $P<10^{-16}$) in the same variable in their

415 partner.

416 *Mendelian randomization: Genetically influenced alcohol consumption and self-*

417 *reported alcohol behaviour of partner*

418 To evaluate the degree to which an individual's alcohol consumption is

419 affected by their partner's genetically influenced alcohol consumption, we used a

420 sample of 47,321 spouse-pairs with available data on weekly alcohol consumption.

421 In this sample, each additional copy of the *ADH1B* major allele was associated with

an increased weekly alcohol consumption of 3.98 units a week (95% C.I. 3.51, 4.43;

P<10^{-16}) in the same individual. Each additional copy of the major allele was

associated with an increased weekly alcohol consumption of 1.04 units a week (95%

C.I. 0.58, 1.51; P=1.09x10^{-5}) in the reference individual's partner. After scaling the

estimate using a Wald estimator; a 1 unit increase in an individual's alcohol

consumption led to having partner's with alcohol consumption 0.26 units higher than

baseline (95% C.I. 0.15, 0.38; P=1.10x10^{-5}). This effect is slightly lower than the

phenotypic estimate of 0.37 units (95% C.I. 0.36, 0.38) although confidence intervals

overlap (Z-test for difference of means: P=0.064).

Characteristics of rs1229984 in the UK Biobank

In the sample of 385,287 individuals of recent European descent, the MAF of

rs1229984 was 2.8% and very strong evidence was found for the SNP violating

HWE (Chi^2 = 275, P <10^{-16}) due to fewer heterozygotes compared to expectation

(expected=20,972, observed=20,194). However, when restricting to the sample of

337,114 individuals of British descent, the MAF of rs1229984 was 2.2% and there

was little evidence of the SNP violating HWE (Chi^2 = 2.0, P=0.16) and there were

more heterozygotes compared to expected (expected= 14,506 observed=14,743)

(**Supplementary Table 3**). Evidence was found of allele frequency differences for

rs1229984 between the two samples (Chi^2=445, P<10^{-16}) suggesting that population

substructure differences may explain the HWE results.

The SNP was found to be strongly associated with both genetic principal

components and birth coordinates in both samples. In the less restrictive European

sample, each additional major allele of rs1229984 was associated with being born

24.6 miles farther north (95% C.I. 22.2, 27.0) and 13.3 miles farther west (95% C.I.

12.1, 14.5). The SNP was similarly associated with principal components and birth

coordinates in the sample of British descent although there were differences in effect

estimates between the two samples (**Supplementary Table 4**). We also found

strong evidence that self-reported alcohol consumption is strongly associated with

birth coordinates and principal components in both samples concordant directionally

with the SNP associations (**Supplementary Table 5**).

Genotypic concordance

Amongst 47,549 spouse-pairs, strong concordance was observed for the

genotype of rs1229984. Each additional copy of the major rs1229984 allele was

associated with an increased number of major alleles in their partner (Beta 0.019;

95% C.I. 0.010, 0.028; P=5.0×10^{-5}).

As a sensitivity analysis, we restricted the sample to 28,653 spouse-pairs

born within 100 miles of each other and stratified spouse-pairs by the 22 different UK

Biobank recruitment centres. In this sample, we did not find strong evidence that

birth location differences were associated with similarities in alcohol behaviour or

rs1229984 genotype, contrasting with clear evidence of associations in the full

spouse-sample. However, we did find evidence that genomic principal component

differences were associated with spousal similarities for these variables, likely

reflecting the fine-scale population structure of UK Biobank (**Supplementary Table**

6). Of the 22 centres, 2 centres were omitted from the meta-analysis because the

limited sample sizes led to convergence issues in regression. A fixed-effects meta-

analysis was then used to estimate the spousal-concordance across the remaining

20 centres and 28,615 spouse-pairs. Evidence was found of spousal concordance

for rs1229984 (Beta 0.016; 95% C.I. 0.004, 0.028; P=0.011), consistent with the

previous analysis. Cochran's Q test for heterogeneity across the betas suggested no

strong evidence for heterogeneity (P= 0.34) across the different centres (**Table 2**).

Table 2: Meta-analysis of spousal-concordance for rs1229984 across the UK
Biobank recruitment centres

Recruitment Centre	Number of spouse-pairs born within 100 km of each other	Beta (95% C.I.)
Stockport	9	N/A[1]
Manchester	662	0.024 (-0.088, 0.0675)
Oxford	669	-0.010 (-0.088, 0.067)
Cardiff	930	0.022 (-0.043, 0.088)
Glasgow	1046	0.072 (0.019, 0.125)
Edinburgh	611	-0.047 (-0.166, 0.070)
Stoke	1215	-0.012 (-0.075, 0.051)
Reading	1352	0.003 (-0.055, 0.060)
Bury	2244	0.012 (-0.031, 0.055)
Newcastle	2976	-0.025 (-0.064, 0.013)
Leeds	2563	0.041 (0.001, 0.081)
Bristol	2117	0.015 (-0.030, 0.060)
St Bartholomew's Hospital	122	-0.073 (-0.220, 0.074)
Nottingham	2342	0.025 (-0.017, 0.066)
Sheffield	2260	0.037 (-0.009, 0.082)
Liverpool	2632	0.023 (-0.020, 0.066)
Middlesbrough	1477	0.002 (-0.050, 0.053)
Hounslow	838	0.073 (-0.000, 0.147)
Croydon	1034	0.044 (-0.027, 0.115)
Birmingham	1440	-0.019 (-0.068, 0.031)
Swansea	85	-0.068 (-0.283, 0.146)
Wrexham	29	N/A[1]
Combined (Fixed effects)	**28,615**	**0.016 (0.004, 0.028) P=0.011**

475
    1 Linear regression estimates did not converge due to limited sample sizes, these studies were excluded from the meta-
analysis.

Relationship length and spousal alcohol behaviour similarities

We did not find strong evidence that increased mean couple age, used as a

proxy for relationship length, was associated with more concordant spousal alcohol

behaviour. Per 1-year increase in couple mean age, spousal differences in terms of

weekly alcohol units consumed were 0.017 smaller (95% C.I. -0.040, 0.007, P=0.16).

In terms of genotypic differences at rs1229984, we found weak evidence that older

couples were more dissimilar at the locus. Per 1-year increase in couple mean age,

spousal allelic differences at rs1229984 were 0.0004 larger (95% C.I. 0.0000,

0.0009; P=0.035).

487   **Discussion**

In this study, we used a large sample of derived spouse-pairs in a UK-based

cohort to demonstrate that an individual's self-reported alcohol use and their

genotype for an alcohol implicated variant, rs1229984 in *ADH1B*, are associated with

their partner's self-reported alcohol use. Furthermore, we showed that the genotype

of the variant is concordant within spouse-pairs. There are several possible

explanations for our findings. First, that rs1229984 influences alcohol behaviour,

which has a downstream effect on mate selection. Second, that a participant's

alcohol use is influenced by their partner's alcohol use. Third, spouse-pairs with

more similar alcohol behaviour were more likely to remain in a relationship, and so

be present in our study sample. Fourth, that given the strong association of the SNP

with both genetic principal components and birth coordinates, the spousal

concordance is related to factors influencing social homogamy, independent of

alcohol behaviour, such as place of birth, ancestry or socio-economic status. Indeed,

the allele frequency of rs1229984 was found to deviate between European and

British subsets of the UK Biobank.

However, we presented evidence suggesting that a substantial proportion of

the spousal concordance is likely to be explained by the biological effects of the

variant on alcohol consumption in the index individual. Firstly, we have tested the

association between a causal SNP for alcohol consumption, and not the measured

consumption itself, thereby avoiding any post-birth confounding factors suggesting

that alcohol use has a direct effect on spousal alcohol use. Secondly, because

rs1229984 is concordant between spouses, there must be some degree of

assortment on alcohol consumption prior to cohabitation. Furthermore, we found little

evidence to suggest that the mean age of each spouse-pair, used as a proxy for

relationship length, was associated with alcohol behaviour similarities. These

findings suggest that the spousal concordance is unlikely to be due to relationship

dissolution after the age of 40. Thirdly, we accounted for possible effects of ancestral

factors, which could have induced confounding, by including principal components

as covariates in the Mendelian randomization analysis. Additionally, as a sensitivity

analysis, we conducted a within centre sensitivity analysis excluding spouse-pairs

born more than 100 miles apart, finding a consistent effect estimate.

The strong evidence for spousal-concordance on the variant has implications

for conventional Mendelian randomization studies (i.e. estimating the causal effect of

an exposure on an outcome) [33] which use the SNP as a genetic proxy for alcohol

intake [45]. Assortative mating could lead to a violation of the Mendelian randomization

assumption, that the genetic instrument for the exposure is not strongly associated

with confounders of the exposure-outcome relationship. If both genetic and

environmental factors affect alcohol consumption, then assortative mating on alcohol

consumption could contribute to associations between genetic and environmental

factors in the offspring, with the strength of association dependent on the degree of

assortative mating [58].

Interestingly, the minor allele of rs1229984 (i.e. associated with lower alcohol

consumption) has been previously found to be positively associated with years in

education [45] and socio-economic related variables, such as the Townsend

deprivation index and number of vehicles in household [59] [60]. Each copy of the minor

allele was associated with an additional 0.023 (95% C.I. 0.012 to 0.034, P=0.00005)

years of education and a 0.016 S.D. (95% C.I. -0.001 to 0.033, P=0.06) increase in

intelligence [61] [62]. These associations may be down-stream causal effects of alcohol

consumption, which implies that some of the spousal concordance for alcohol

consumption could be explained by assortative mating on educational attainment [15]

or alternatively these associations may reflect maternal genotype and intrauterine

effects [63]. Over time, assortative mating on alcohol consumption may further

strengthen the associations between rs1229984 and socio-economic related

variables [58]. Of further interest is that the variant has previously been shown to be

under selection [64] suggesting that the variant has historically had a substantial effect

on reproductive fitness and may partially explain the violation of HWE observed

across Europeans in our analyses.

The analyses in this study extended previous work on the concordance

between spouse-pairs for alcohol behaviour [7-12] by comparing the phenotypic

concordance with analyses utilising a genetic variant strongly associated with alcohol

consumption. A major strength of this study is the use of distinct methods with

different non-overlapping limitations, allowing for improved inference by triangulating

the results from the different methods [65]. First, we evaluated the spousal phenotypic

concordance for self-reported alcohol consumption, second we investigated the

effect of an individual's rs1229984 genotype on the alcohol consumption of their

spouse using Mendelian randomization, third we demonstrated spousal genotypic

concordance for rs1229984 and fourth we explored whether older couples have

more similar alcohol behaviour. The use of the UK Biobank data-set was a

considerable strength for these analyses because of the low frequency of the

rs1229984 minor allele; the large scale of the UK Biobank allowed for the

identification of thousands of genotyped spouse-pairs. A further strength of these

analyses is that we have demonstrated the utility of a Mendelian randomization

framework for application to assortative mating by applying it to height and alcohol

use. Indeed, the evidence for differences between the observational and Mendelian

randomization estimates for spousal height suggest that the observational estimate

may be inflated by confounding factors although differences could also be related to

the attenuated effects of phenotypic assortment on genetic associations. A similar

approach using polygenic risk scores has previously demonstrated assortative

mating on educational attainment [18]. However, the use of Mendelian randomization

has a notable advantage over polygenic approaches because of the possibility of

using various sensitivity analyses to test for heterogeneity and consistency of the

effect estimate [50-52].

There are several limitations of this study. First, although spouse-pairs were

identified using similar methods to previous studies [15-17], the identified spouse-pairs

have not been confirmed. However, the phenotypic spousal concordance estimate

for height found in this study is highly concordant with previous estimates [56],

consistent with derived couples being genuine. Second, despite follow-up analyses,

it is difficult to definitively prove that the spousal concordance is a direct result of

assortative mating on alcohol consumption. Assortment independent of alcohol use,

potentially relating to ancestral or geographical factors, cannot be completely ruled

out and down-stream pleiotropic effects of the variant may influence mate selection.

Third, the use of a single genetic instrument in the Mendelian randomization

analysis, limited the use of sensitivity analyses [50-52] and meant it is not possible to

infer similar associations for other alcohol-implicated variants. Fourth, selection into

the UK Biobank, particularly with regards to participation of spouse-pairs is a

potential source of bias [66]. Fifth, it is unclear whether the mean age of each couple is

a suitable proxy for relationship length, which limits conclusions regarding the

possibilities of partner interactions and relationship dissolution. Indeed, patterns of

assortment on alcohol behaviour changing over time would confound the use of this

proxy. Finally, it is difficult to extrapolate the results of this study in the UK Biobank to

non-European populations. This is because of potential contextual influences; for

example, in some East Asian populations, males are much more likely to consume

alcohol than females[67,68]. Indeed, even within the UK, there may be regional

variation that we were unable to detect in this study. Additionally, there is some

evidence that the effect of genetic contributors to alcohol varies across different

populations [29].

To conclude, our results suggest that there is non-random mating on

rs1229984 in *ADH1B*, likely related to the effect of the variant on alcohol behaviour.

These results suggest that alcohol use influences mate selection and argue for a

more nuanced approach to considering social and cultural factors when examining

causality in epidemiological studies. Further research investigating other alcohol-

implicated variants, and other societies and ethnicities, and assortment on other

phenotypes, would strengthen these conclusions.

602   **Figure titles and descriptions**

Figure 1 Possible explanations for spousal concordance on alcohol use.

(A) Assortative mating. Alcohol behaviour influences mate selection; individuals are

more likely to select a mate with similar alcohol consumption.

(B) Social homogamy or confounding. An unknown confounder influence mate

selection independent of alcohol behaviour. For example, ancestry or socio-

economic status may influence both alcohol use and mate choice.

(C) Partner interaction effects. As spouse-pairs cohabitate their alcohol behaviour

becomes more similar over time.

(D) Relationship dissolution. Spouse-pairs with more similar alcohol behaviour are

more likely to remain in a relationship and be recruited into UK Biobank or similarly,

are more likely to participate in the study together.

Figure 2 Interpretations of phenotypic concordance, Mendelian randomization and

genotypic concordance analyses between-spouses.

(A) Phenotypic concordance. Spousal concordance for alcohol use could be

explained by a direct effect of an individual's alcohol consumption on their partner's

alcohol consumption (assortative mating, partner interaction effects or relationship

dissolution) or confounding factors such as assortment on social factors (social

homogamy) leading to spousal correlation for alcohol use.

(B) Mendelian randomization framework. An association between an individual's

alcohol influencing genotype and their spouse's alcohol use would suggest that the

spousal concordance is explained by a direct effect of alcohol consumption. Genetic

variants are unlikely to be associated with socio-economic confounders suggesting

that social homogamy is unlikely. Spousal phenotype/genotype associations induced

by assortment are dependent on the heritability of the trait (see Supplementary

Methods).

(C) Genotypic concordance. Genotypic concordance for alcohol related genetic

variants would suggest that some degree of the spousal concordance is explained

by assortative mating or relationship dissolution. Partner interaction effects cannot

lead to genotypic concordance because genotypes are fixed from birth. Spousal

genotypic concordance induced by assortment is dependent on the trait heritability

(see Supplementary Methods).

645   **Acknowledgements**

LJH was a Medical Research Council funded PhD student at the University of Bristol

and is now funded by the British Heart Foundation and University College London.

NMD, SJL and GDS work in the Medical Research Council Integrative Epidemiology

Unit at the University of Bristol (MC_UU_00011/1) which is supported by the Medical

Research Council and the University of Bristol. NMD is supported by the Economics

and Social Research Council (ESRC) via a Future Research Leaders grant

[ES/N000757/1]. DJL [WT104125MA] and GH [208806/|/17/|] are both supported by

the Wellcome Trust. UK Biobank received ethical approval from the Research Ethics

Committee (11/NW/0383). This research was approved as part of application 15825

(PI: Dr Philip Haycock).

Conflicts of interest

Neil Davies reports a grant for research unrelated to this work from the Global

Research Awards for Nicotine Dependence (GRAND), an independent grant making

body funded by Pfizer.

Contributions

LJH, GDS, GH and DJL formulated the project outline and analysis plan. LJH

performed all statistical analyses and drafted the first manuscript draft under

supervision from GDS, GH, SJL, BSP and NMD. All authors contributed to

interpretation of results and writing of the final manuscript.

References

1. Vandenberg SG. Assortative mating, or who marries whom? *Behavior Genetics*
1972;2(2-3):127-57.
2. Buss DM. Human mate selection: Opposites are sometimes said to attract, but in
fact we are likely to marry someone who is similar to us in almost every
variable. *American Scientist* 1985;73(1):47-51.
3. Mare RD. Five decades of educational assortative mating. *American Sociological*
Review 1991:15-32.
4. Silventoinen K, Kaprio J, Lahelma E, et al. Assortative mating by body height and
BMI: Finnish twins and their spouses. *American Journal of Human Biology*
2003;15(5):620-27.

5. Krueger RF, Moffitt TE, Caspi A, et al. Assortative mating for antisocial behavior:
Developmental and methodological implications. *Behavior Genetics*
1998;28(3):173-86.
6. Watson D, Klohnen EC, Casillas A, et al. Match makers and deal breakers:
Analyses of assortative mating in newlywed couples. *Journal of Personality*
2004;72(5):1029-68.
7. Agrawal A, Heath AC, Grant JD, et al. Assortative mating for cigarette smoking
and for alcohol consumption in female Australian twins and their spouses.
Behavior Genetics 2006;36(4):553-66.
8. Hall RL, Hesselbrock VM, Stabenau JR. Familial distribution of alcohol use: II.
Assortative mating of alcoholic probands. *Behavior Genetics* 1983;13(4):373-
82.
9. Hall RL, Hesselbrock VM, Stabenau JR. Familial distribution of alcohol use: I.
Assortative mating in the parents of alcoholics. *Behavior Genetics*
1983;13(4):361-72.
10. McLeod JD. Spouse concordance for alcohol dependence and heavy drinking:
Evidence from a community sample. *Alcoholism: Clinical and Experimental*
Research 1993;17(6):1146-55.
11. Reynolds CA, Barlow T, Pedersen NL. Alcohol, tobacco and caffeine use:
spouse similarity processes. *Behavior Genetics* 2006;36(2):201.
12. Grant JD, Heath AC, Bucholz KK, et al. Spousal concordance for alcohol
dependence: evidence for assortative mating or spousal interaction effects?
Alcoholism: Clinical and Experimental Research 2007;31(5):717-28.
13. Ask H, Rognmo K, Torvik FA, et al. Non-random mating and convergence over
time for alcohol consumption, smoking, and exercise: the Nord-Trøndelag
Health Study. *Behavior Genetics* 2012;42(3):354-65.
14. Domingue BW, Fletcher J, Conley D, et al. Genetic and educational assortative
mating among US adults. *Proceedings of the National Academy of Sciences*
2014;111(22):7996-8000.
15. Robinson MR, Kleinman A, Graff M, et al. Genetic evidence of assortative mating
in humans. *Nature Human Behaviour* 2017;1:0016.
16. Rawlik K, Canela-Xandri O, Tenesa A. Indirect assortative mating for human
disease and longevity. *bioRxiv* 2017:185207.
17. Tenesa A, Rawlik K, Navarro P, et al. Genetic determination of height-mediated
mate choice. *Genome Biology* 2016;16(1):269.
18. Hugh-Jones D, Verweij KJ, Pourcain BS, et al. Assortative mating on educational
attainment leads to genetic spousal resemblance for polygenic scores.
Intelligence 2016;59:103-08.
19. Verhulst B, Neale MC, Kendler KS. The heritability of alcohol use disorders: a
meta-analysis of twin and adoption studies. *Psychological Medicine*
2015;45(5):1061-72.
20. Walters GD. The heritability of alcohol abuse and dependence: a meta-analysis
of behavior genetic research. *The American Journal of Drug and Alcohol*
Abuse 2002;28(3):557-84.
21. Clarke T-K, Adams MJ, Davies G, et al. Genome-wide association study of
alcohol consumption and genetic overlap with other health-related traits in UK
Biobank (N= 112 117). *Molecular Psychiatry* 2017;22(10):1376.
22. Gelernter J, Kranzler H, Sherva R, et al. Genome-wide association study of
alcohol dependence: significant findings in African-and European-Americans
including novel risk loci. *Molecular Psychiatry* 2014;19(1):41.

23. Park BL, Kim JW, Cheong HS, et al. Extended genetic effects of ADH cluster
genes on the risk of alcohol dependence: from GWAS to replication. *Human*
Genetics 2013;132(6):657-68.
24. Bierut LJ, Goate AM, Breslau N, et al. ADH1B is associated with alcohol
dependence and alcohol consumption in populations of European and African
ancestry. *Molecular Psychiatry* 2012;17(4):445.
25. Bierut LJ, Agrawal A, Bucholz KK, et al. A genome-wide association study of
alcohol dependence. *Proceedings of the National Academy of Sciences*
2010;107(11):5082-87.
26. Treutlein J, Cichon S, Ridinger M, et al. Genome-wide association study of
alcohol dependence. *Archives of General Psychiatry* 2009;66(7):773-84.
27. Schumann G, Coin LJ, Lourdusamy A, et al. Genome-wide association and
genetic functional studies identify autism susceptibility candidate 2 gene
(AUTS2) in the regulation of alcohol consumption. *Proceedings of the*
National Academy of Sciences 2011;108(17):7119-24.
28. Schumann G, Liu C, O'Reilly P, et al. KLB is associated with alcohol drinking,
and its gene product β-Klotho is necessary for FGF21 regulation of alcohol
preference. *Proceedings of the National Academy of Sciences*
2016;113(50):14372-77.
29. Jorgenson E, Thai KK, Hoffmann TJ, et al. Genetic contributors to variation in
alcohol consumption vary by race/ethnicity in a large multi-ethnic genome-
wide association study. *Molecular Psychiatry* 2017;22(9):1359.
30. Edenberg HJ, McClintick JN. Alcohol Dehydrogenases, Aldehyde
Dehydrogenases, and Alcohol Use Disorders: A Critical Review. *Alcoholism:*
Clinical and Experimental Research 2018;42(12):2281-97.
31. Thomasson HR, Edenberg HJ, Crabb DW, et al. Alcohol and aldehyde
dehydrogenase genotypes and alcoholism in Chinese men. *American Journal*
of Human Genetics 1991;48(4):677.
32. Luczak SE, Glatt SJ, Wall TJ. Meta-analyses of ALDH2 and ADH1B with alcohol
dependence in Asians: American Psychological Association, 2006.
33. Davey Smith G, Ebrahim S. 'Mendelian randomization': can genetic
epidemiology contribute to understanding environmental determinants of
disease? *International Journal of Epidemiology* 2003;32(1):1-22.
34. Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal
inference in epidemiological studies. *Human molecular genetics*
2014;23(R1):R89-R98.
35. Davey Smith G, Lawlor DA, Harbord R, et al. Clustered environments and
randomized genes: a fundamental distinction between conventional and
genetic epidemiology. *PLoS Medicine* 2007;4(12):e352.
36. Sudlow C, Gallacher J, Allen N, et al. UK Biobank: an open access resource for
identifying the causes of a wide range of complex diseases of middle and old
age. *PLoS Medicine* 2015;12(3):e1001779.
37. Bycroft C, Freeman C, Petkova D, et al. The UK Biobank resource with deep
phenotyping and genomic data. *Nature* 2018;562(7726):203.
38. Mitchell RE, Hemani G, Dudding T, et al. UK Biobank Genetic Data: MRC-IEU
Quality Control, version 1, 13/11/2017 2017 [
39. International HapMap 3 Consortium. Integrating common and rare genetic
variation in diverse human populations. *Nature* 2010;467(7311):52-58.
40. Genomes Project Consortium. A global reference for human genetic variation.
Nature 2015;526(7571):68-74.

41. Mountjoy E, Davies NM, Plotnikov D, et al. Education and myopia: assessing the
direction of causality by mendelian randomisation. *BMJ* 2018;361:k2022.
42. O'Connell J, Sharp K, Shrine N, et al. Haplotype estimation for biobank-scale
data sets. *Nature Genetics* 2016;48(7):817-20.
43. Consortium UK. The UK10K project identifies rare variants in health and disease.
Nature 2015;526(7571):82-90.
44. McCarthy S, Das S, Kretzschmar W, et al. A reference panel of 64,976
haplotypes for genotype imputation. *Nature Genetics* 2016;48(10):1279.
45. Holmes MV, Dale CE, Zuccolo L, et al. Association between alcohol and
cardiovascular disease: Mendelian randomisation analysis based on
individual participant data. *BMJ* 2014;349:g4164.
46. Hemani G, Zheng J, Elsworth B, et al. The MR-Base platform supports
systematic causal inference across the human phenome. *eLife*
2018;7:e34408.
47. Wood AR, Esko T, Yang J, et al. Defining the role of common variation in the
genomic and biological architecture of adult human height. *Nature Genetics*
2014;46(11):1173.
48. Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome
association and population-based linkage analyses. *The American Journal of*
Human Genetics 2007;81(3):559-75.
49. Greco M, Del F, Minelli C, et al. Detecting pleiotropy in Mendelian randomisation
studies with summary data and a continuous outcome. *Statistics in Medicine*
2015;34(21):2926-40.
50. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid
instruments: effect estimation and bias detection through Egger regression.
International Journal of Epidemiology 2015;44(2):512-25.
51. Bowden J, Davey Smith G, Haycock PC, et al. Consistent estimation in
Mendelian randomization with some invalid instruments using a weighted
median estimator. *Genetic Epidemiology* 2016;40(4):304-14.
52. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data
Mendelian randomization via the zero modal pleiotropy assumption.
International Journal of Epidemiology 2017;46(6):1985-98.
53. Zuccolo L, Lewis SJ, Davey Smith G, et al. Prenatal alcohol exposure and
offspring cognition and school performance. A 'Mendelian
randomization'natural experiment. *International Journal of Epidemiology*
2013;42(5):1358-70.
54. Leslie S, Winney B, Hellenthal G, et al. The fine-scale genetic structure of the
British population. *Nature* 2015;519(7543):309.
55. Haworth S, Mitchell R, Corbin L, et al. Common genetic variants and health
outcomes appear geographically structured in the UK Biobank sample: Old
concerns returning and their implications. *bioRxiv* 2018:294876.
56. Price RA, Vandenberg SG. Spouse similarity in American and Swedish couples.
Behavior Genetics 1980;10(1):59-71.
57. Pearson K, Lee A. On the laws of inheritance in man: I. Inheritance of physical
characters. *Biometrika* 1903;2(4):357-462.
58. Hartwig FP, Davies NM, Davey Smith G. Bias in Mendelian randomization due to
assortative mating. *Genetic epidemiology* 2018
59. Canela-Xandri O, Rawlik K, Tenesa A. An atlas of genetic associations in UK
Biobank. *Nature Genetics* 2018;50(11):1593.

60. Howe LJ, Sharp GC, Hemani G, et al. Prenatal alcohol exposure and facial
828   morphology in a UK cohort. *Drug and Alcohol Dependence* 2019
829   61. Lee JJ, Wedow R, Okbay A, et al. Gene discovery and polygenic prediction from
830   a genome-wide association study of educational attainment in 1.1 million
831   individuals. *Nature Genetics* 2018;50(8):1112.
832   62. Hill W, Marioni R, Maghzian O, et al. A combined analysis of genetically
833   correlated traits identifies 187 loci and a role for neurogenesis and myelination
834   in intelligence. *Molecular Psychiatry* 2018:1.
835   63. Hemani G, Bowden J, Haycock PC, et al. Automating Mendelian randomization
836   through machine learning to construct a putative causal map of the human
837   phenome. *bioRxiv* 2017:173682.
838   64. Galinsky KJ, Bhatia G, Loh P-R, et al. Fast principal-component analysis reveals
839   convergent evolution of ADH1B in Europe and East Asia. *The American
840   Journal of Human Genetics* 2016;98(3):456-72.
841   65. Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology.
842   *International Journal of Epidemiology* 2016;45(6):1866-86.
843   66. Munafò MR, Tilling K, Taylor AE, et al. Collider scope: when selection bias can
844   substantially influence observed associations. *International Journal of
845   Epidemiology* 2017;47(1):226-35.
846   67. Spiller W, Slichter D, Bowden J, et al. Detecting and correcting for bias in
847   Mendelian randomization analyses using Gene-by-Environment interactions.
848   *International Journal of Epidemiology* 2018:187849.
849   68. Chen L, Davey Smith G, Harbord RM, et al. Alcohol intake and blood pressure: a
850   systematic review implementing a Mendelian randomization approach. *PLoS
851   Medicine* 2008;5(3):e52.