On simulating cold stunned turtle strandings on Cape Cod

Liu Xiaojian[1]¶, James Manning[2]¶*, Robert Prescott[3], Zou Huimin[4], Mark Faherty[3]

[1] School of Remote Sensing and Information Engineering,

Wuhan University , Wuhan, China

[2] National Oceanic Atmospheric Administrations's Northeast Fisheries Science

Center, Woods Hole, Massachusetts, United States of America

[3] Massachusetts Audubon's Wellfleet Bay Wildlife Sanctuary, Wellfleet,

Massachusetts 02663 United States of America

[4] Shandong Agricultural University, Taian, China

*Corresponding author, E-mail: james.manning@noaa.gov

¶ LX and JM are Joint Senior Authors

Abstract

Kemp's ridley turtles were on the verge of extinction in the 1960s. While they have slowly recovered, they are still endangered. In the last few years, the number of strandings on Cape Cod Massachusetts beaches has increased by nearly an order of magnitude relative to preceding decades. This study uses a combination of ocean observations and models to investigate the causes and transport of cold-stunned animals in Cape Cod Bay. The Finite Volume Community Ocean Model (FVCOM) is used here as an example of a well-respected high resolution model. First, using data from satellite-tracked drifters and local temperature moorings, the FVCOM ocean model was validated. The model was used to simulate the ocean currents in the Cape Cod Bay during periods of anomalous strandings and to explain stranding locations as observed by volunteers. Nearby observations and model estimates of water temperature and wind are used to document the forcing mechanisms associated with strandings at particular times and places. Finally, the origin of the stranding turtles was estimated by backtracking through modelled flow fields.

Key words: Kemp's ridley turtles; FVCOM model; ocean current; particle tracking

Introduction

Kemp's ridley is the smallest and the most endangered sea turtle in the world and shows little sign of recovery [1,2]. Their perilous situation is attributed primarily to the over-harvesting of their eggs during the last century. In 1970, it was listed by the U.S. Fish and Wildlife Service [3] as "endangered throughout its range" and it has received Federal protection under the U.S. Endangered Species Act of 1973 ever since [2].

During the past 18 years, Padre Island National Seashore (PAIS) has been using

satellite telemetry to understand migratory patterns and habitat of adult Kemp's

ridleys [4,5,6]. Kemp's ridley adults are primarily found in the Gulf of Mexico [7-8].

Juveniles are also found along the eastern seaboard of the United States as far north as

Nova Scotia, Canada [9]. The adult Kemp's ridley turtles are mainly active in coastal

waters less than 165 feet deep, and the smallest juveniles Kemp's ridley are found in

shallow waters, often foraging in less than 3 feet of water [10]. Young sea turtles have

ventured up the Northeast Coast, leaving warm seas to feed on crabs and other prey.

Sea turtles on the East Coast of the US swim to warmer waters to spend the

winter. For example, turtles in the vicinity of Virginia, tend to migrate south in late

fall each year [11,12,13]. However, even in Florida, significant mortalities have been

reported [14] as a result of being caught in some water bodies/estuaries. In laboratory

experiments, Schwartz [15] demonstrated that Kemp's ridleys become inactive at

temperatures below 10^{o}C, although they are somewhat more resistant than other

species. If a sea turtle becomes cold stunned there is little chance of survival without

assistance. Morreale, et al. [16] studied cold stunned stranding events in the late 1980s

on Long Island, NY. Burke et al [17] specifically addressed the causes of these events

and found one year, in particular, when fewer strandings occurred. They attributed

this inter-annual variability to slight changes in the prevalent wind direction at the

critical time of year when the water temperature fell below 10^{o}C. As documented by

Still et al, 2005, when Kemp's ridley's inside Cape Cod Bay are late in heading south

and the water temperature gets below 10.4^o, they become "cold stunned" [18].

Kemp ridleys in Cape Cod Bay are unable to move south due to the presence of

the Cape Cod landmass. These turtles, typically 2 to 3 years old, are sized between a

dinner plate and a serving platter [19]. They are so small that we can not easily use the

satellite telemetry devices typically used to track adult turtles in Gulf of Mexico to

track their position. In recent years, a large number of cold stunned Kemp's ridley

turtles have been stranding on the shores of Cape Cod Bay on the eastern coast of the

United States in late Fall. These turtles are extremely weak, often wash ashore, and

need to be rescued by dozens of trained volunteers each fall. Since 1979, Wellfleet

Bay Wildlife Sanctuary staff and volunteers have patrolled the beaches of Cape Cod,

on the lookout for cold-stunned turtles, which are rapidly transported to the New

England Aquarium for evaluation and rehabilitation. In the last few years, the number

of strandings has increased by nearly an order of magnitude relative to the strandings

in the preceding decades.

This paper explores the transport of cold stunned Kemp's ridleys using the Finite

Volume Community Ocean Model (FVCOM), a numerical ocean model well-suited

to simulating coastal ocean processes. First, we attempt to validate FVCOM by using

drifter data and mooring data from the region. Secondly, we look at the number of sea

turtles stranded in various towns along the shore of Cape Cod Bay from 2012 to 2015

and try to explain these distributions based on FVCOM-derived simulated particle

tracks, wind records, and observations of water temperature.

Methods

FVCOM model

FVCOM is a prognostic, unstructured-grid, Finite-Volume, free-surface, three-

dimensional (3-D) primitive equations Community Ocean Model developed originally

by Chen et al [20]. FVCOM is solved numerically by a second-order accurate discrete

flux calculation in the integral form of the governing equations over an unstructured

triangular grid. This approach combines the best features of finite-element methods

(grid flexibility) and finite-difference methods (numerical efficiency and code

simplicity) and provides a much better numerical representation of both local and

global momentum, mass, salt, heat, and tracer conservation. The ability of FVCOM to

accurately solve scalar conservation equations in addition to the topological flexibility

provided by unstructured meshes and the simplicity of the coding structure has made

FVCOM ideally suited for many coastal and interdisciplinary scientific applications

[21].

FVCOM model has been run on several grids spanning the Northeast Continental

Shelf of the US and is still evolving. The third generation grid (GOM3) was used in

this experiment covering Gulf of Maine and Georges Bank waters and hourly flow

fields are available from 1978 to 2015. The GOM3 grid was developed primarily for

the prediction of coastal ocean dynamics, with horizontal grid resolution ranging from

20 meters to 15 kilometers and includes tides. The vertical domain was divided into

45 s-coordinate levels [22]. We also experimented on finer-resolution "MassBay"

grid that had hourly flow fields available from 2011-2014.

Drifter observations

The surface drifters used in this study track the top meter of the water column

[23]. This type is typically referred to as the "Davis-style" or "CODE" drifter first

developed by Russ Davis for the Coastal Ocean Dynamics Experiment in the early

1980s (Fig 1) [24]. Our version of this drifter typically has a metal-frame and four

fabric sails measuring 91 cm long and 48 cm wide. These units were deployed in the

vicinity of Cape Cod from 1978 to 2010 with tracks archived for public access. More

than a thousand of these student-built drifters have been deployed off the New

England coast in the past few decades [25].

Fig 1. Schematic of standard surface drifter from Poulain [24] (left) and photo of

surface drifter used in this report being recovered after stranding in Barnstable

MA (right).

Moored temperature observations

There are at least three different sets of moored ocean temperature observations

available in the Northeast US. The Northeastern Regional Association of Coastal

Ocean Observing System (NERACOOS) typically has several mooring sites in the

Gulf of Maine [26]. In 2017, they added a mooring (CDIP) inside Cape Cod Bay. The

NERACOOS Mooring A location is shown in Fig 2. The Environmental Monitors on

Lobster Trap Project (eMOLT) [27] comprises another collection of moorings with

18+ years of hourly bottom temperatures from dozens of locations off the New

England coast include several within Cape Cod Bay. Finally, the Mass Division of

Marine Fisheries (DMF) maintains several diver-serviced bottom temperature times

series at a set of shipwrecks off the Massachusetts coast.

Fig 2. Locations of moored observations used in this report including two

NERACOOS moorings (blue triangles), five eMOLT moorings (red circles)

and four Massachusetts Division of Marine of Fisheries moorings (green

squares).

On using models and observations

Forward tracks through modeled flow compared with drifter tracks

Surface drifter tracks from 2012 to 2015 are used to quantify the difference

between FVCOM model currents and observed Lagrangian currents. We limited

drifter data to those with sample intervals less than 2.5 hours. This resulted in 55

drifter tracks in Cape Cod Bay that had a total of 1365 days. We also compared the

model track to the drifter track by restarting the comparison every 3 days for a total of

454 comparisons. When the drifter trajectory is compared with the model trajectory,

the "separation distance" and "distance ratio" are often used as two measures of

accuracy. They are defined as the distance between the drifter point and the modeled

track point at the same time and the ratio of separation distance to the drifter's total

distance traveled, respectively.

Backtracking turtle trajectories

We would like to estimate the origin of stranded turtles based on backtracking

particles through numerical models. However, we first need to get the simulated turtle

off the beach in order to begin moving it backward through the water. The process of

selecting a "short distance off the beach" is not easy. We experimented with multiple

methods (simple to complex) to solve this problem. For the purpose of this study, we

chose the simplest method which involves a) finding the nearest point on the model-

based coastline to the stranding location, b) calculating the line from the specified

stranding to that point, and c) then extending the line X km out to sea. After

experimenting with several values, we chose to set "X =1.5 km". The backward

tracking is conducted by moving the simulated turtle with speed and direction as

prescribed by the model at the nearest grid nodes with hourly time steps. The tracking

is terminated when the estimated water temperature exceeds the cold-stunned level.

For reasons discussed below, we chose this to be 10.5 ^{0}C.

Binned averaged current and wind to help explain strandings

The model data used is the hourly output of the FVCOM model as described in section 2.1 above. Even before looking at particle tracks, we first averaged the observed current, modeled current, and wind stress on 0.05 ° bins. The "observed current" is derived from averaging the first differences of the near-hourly drifter fixes within each bin. The "modeled current" in each grid cell is the average velocities of all nodes within each bin whenever drifters were present in the bin. While technically different types of current velocity (Lagrangian vs Eulerian), we assume they are the same. After removing the tidal effects by sequentially averaging over the solar (24 hour) and lunar (24.841) periods, we calculated the average surface current of each bin. The wind stress was derived from the same weather model that drives the FVCOM ocean model.

Turtle stranding data

The Kemp's ridley turtle data used in this experiment was provided by Mass Audubon's Wellfleet Bay Wildlife Sanctuary. These data include information on date and location of stranding for each cold-stunned Kemp's ridley Sea turtle from 1991 to 2015. Hundreds of trained volunteers have been participating in this project since 1978. Beach walkers cover the coastline each fall to document strandings and report their findings.

Sea-surface temperature observations

In addition to the modeled sea-surface temperature (SST) estimates available hourly, we also have observations of SST from satellite. While clear images are only available a few times per month in this region, they provide another snapshot

180 realization of the conditions in Cape Cod Bay. For the purpose of this report, we

181 accessed the raw data prepared by the University of Delaware and posted via the

182 MARACOOS.org website.

Results

Kemp's ridley strandings on Cape Cod

Fig 3 shows the number of Kemp's ridleys stranded on the shores of Cape Cod

Bay from 1991 to 2015 (with some years 1994-1998 not available). As shown, the

number of strandings vary from year to year but have increased dramatically in recent

years.

Fig 3. 1991-2015 Cape Cod Bay turtle strandings.

Turtles were recovered in several towns each year between 2012 and 2015 (Figs

4 and 5). The towns with the largest numbers of strandings between 2012-2015 were

located on the mid to outer Cape (Table 1). Among them, each town reported

strandings in more than one year of the study period, and strandings were reported

every year in Wellfleet and Brewster. While most stranding events occur on the Outer

Cape, some occur Mid Cape. Our hypothesis is that year-to-year changes of stranding

locations are primarily due to small changes in the surface current.

Fig 4. Total turtle strandings per town from 2012-2015. Most of the "Outer

Cape" towns (green dots) had >400 strandings while some of the "Mid Cape"

towns (purple) had ~200.

Fig 5. Chart of the number of strandings by town between 2012 - 2015 showing

most in the first three Outer Cape towns.

Table 1. The top four towns with the largest number of strandings from 2012 to 2015

Years	1st	2nd	3rd	4th
2012	Dennis	Barnstable	Brewster	Wellfleet
2013	Brewster	Eastham	Dennis	Wellfleet
2014	Eastham	Truro	Wellfleet	Brewster
2015	Wellfleet	Truro	Eastham	Brewster

Verifying FVCOM model

After describing the distribution of strandings in different years and towns, we

now turn to verifying the FVCOM model before using it to simulate turtle advection.

Before any particle tracking is conducted, we first examine the bin-averaged current

and wind. The current averaging is conducted for both the drifter observations and the

model output and shown in Fig 6.

Fig 6. Surface current derived from all drifters (a) and FVCOM model (b),

number of drifter observations prior to 2018 (c), and contoured bathymetry

in meters (d). It is clear from panel c that there are relatively few drifter

observations inside Cape Cod Bay compared to other parts of Mass Bay and

the ocean side of Cape Cod.

If we derive the mean field from drifters whenever there was GOM3 model data

available (<2015), we get the vector plot Fig 6a and compare it to the model-derived

field in Fig 6b. In a mean-flow sense, the model does well to estimate the flow of the

coastal waters in this region. While there are some areas of discrepancies in, for

example, the middle of the bay, most of the region is resolved by the model. When we

calculate the overall mean differences (observed-mean), we find values of X and Y

for the eastward and northward flows, respectively. The variability of the flow is less

resolved with the mean standard deviations of these means being 0.07m/s and

0.09m/s, respectively. Given additional drifter deployments in Cape Cod Bay

providing more observations of current (Fig 6c) especially in recent years (2016-

2017), we derived a new mean field in Fig 6d showing very small subtidal mean flow

field.

We next compare observed and modeled drifter tracks directly. The results are

shown in Fig 7. The 3-day "separation distances" (defined earlier) are mainly

distributed around 20 km, the minimum value is close to 0km and the maximum is 80

km or more; while the "distance ratios" are distributed around 0.2, the minimum value

is close to 0 and the maximum value is above 1.0. Based on these comparisons, we

estimate that the FVCOM model can be used to simulate particle pathways with an

uncertainty of 200 m per kilometer of trackline.

Fig 7. The 3-day separation distance and the distance ratio between observed

and modeled drifter track.

Taking 2012 and 2013 as an example, we compared the surface temperature

monitored by a mooring and the temperature simulated by the nearest FVCOM model

grid cell (less than 1km away). As shown in Fig 8a and Fig 8b and Table 2, the

surface temperature monitored by NERACOOS mooring A is close (<0.4 $^{\circ}$C) from

the temperature simulated by FVCOM. We also compared the bottom temperature

monitored by a nearby (again <1km from model grid node) eMOLT station and the

temperature simulated by the FVCOM model, as shown in Fig 8c and Fig 8d.

FVCOM bottom temperatures were less accurate near the bottom but still within 0.6

$^{\circ}$C of the observed.

Fig 8. Observed (NERACOOS Mooring A) vs modeled (FVCOM) surface (2

meters) temperatures in 2012 (a) and 2013 (b). Observed (eMOLT RM04) vs

modelled (FVCOM) bottom temperatures in 2012 (c) and 2013 (d).

Table 2. Observed minus modeled temperatures where NERACOOS "Mooring A" is
observed at the surface and eMOLT "RM04" is observed at the bottom.
250

Time	index	Surface temperature	Bottom temperature
		NERACOOS-FVCOM	eMOLT-FVCOM
2012	mean	-0.385	-0.540
	std	0.334	0.517
2013	mean	-0.220	-0.600
	std	0.267	0.500
2014	mean	-0.087	NA
	std	0.270	NA

Examining binned current and wind

Despite recent efforts to survey the bay from both ships and planes, we do not

know the exact location of the cold stunned turtles in the bay or in the water column.

However, considering that the cold-stunned turtles are not capable of swimming, we

assume they generally follow the surface currents with some effects by the wind and

waves. It is expected that more skillful predictions of stranding positions may be

made using observed ocean surface currents and wind stress to numerically advect

water parcels forward. Fig 9 shows the number of turtles stranded on the shores of

Cape Cod Bay per day in late 2014 with most occurring the week of November 21st.

Fig 10 maps the mean current and the mean wind stress from November 18 – 23,

2014, and the distribution of the stranded sea turtles for these six days.

Fig 9. The number of turtles stranded on Cape Cod Bay beaches in Nov-Dec 2014.

Fig 10. November 18-23, 2014 mean wind (black arrows) with November 3, 2014 sea surface temperature (a), Mass Bay grid simulation of mean ocean surface currents (black arrows) during that time overlaid on surface current variance ellipses (pink) (b), and distribution of turtle strandings by town with November 19, 2014 sea surface temperature (c) , and mean wind for the entire month of November 2014 with December 26, 2014 sea surface temperature (d). The legends are posted on the land mast on the western side of the charts. The largest red dot in panel c denotes 260 turtles stranded in Eastham during the period 18-23 November 2014.

From November 18, 2014 to November 23, 2014, as depicted on the turtle stranding map (Fig 10c), Truro, Wellfleet, and Eastham had the most strandings (253, 225, and 260, respectively), and their positions are consistent with the oceanic currents and winds shown by the simulated current and wind maps (Fig 10a and 10b), distributed in the area where currents and wind stress are most intense, and in their direction. It can be seen that the combination of ocean current and wind calculated by FVCOM model can be used to explain the general distribution of strandings. Comparing the mean wind for one week in mid-November 2014 (Fig 10a) to the mean wind for the entire month of November 2014 (Fig 10d), we see the intensity of the mid-November period explains the unusually high rate of strandings on the Outer Cape. The intensity of these events is best described in the form of time series plots below. In the future turtle rescue work, FVCOM 3 day forecast model can be used to simulate stranding events and help volunteer beach walkers focus their efforts.

As previous studies have found, when the water temperature of the Cape Cod

Bay is less than 50F (10℃), the metabolism of Kemp's ridley turtle will slow to the

point where they are unable to move and are at the mercy of the wind, waves, and

currents [18]. As shown in Fig 11, we set up 10 temperature detection points and 10

wind stress monitoring points across Cape Cod Bay and Mass Bay to extract the

temperature and wind stress simulated by the model. In the following plots (Figs 12-

14), we examine the relationship between wind stress and strandings especially when

the ocean temperature is below 10.5 ^{o}C.

Fig 11. Ten modeled wind stress points (green dots) and ten modeled surface

temperature points (green).

Fig 12. 2012: Top panel shows wind stress (Pa) direction shown in green vectors

(north up) and magnitude in purple. Bottom panel shows Mid-Cape strandings

(blue bars) the Outer-Cape strandings (red bars), modeled surface temperature

(green line), and observed bottom temperature (blueline) in 2012.

Fig 13. 2013: Top panel: Wind stress with direction shown in green vectors

(north up) and magnitude in purple. Bottom panel: Mid-Cape (blue bars) and

Outer-Cape (red bars) strandings, modeled surface temperature (green line),

and observed bottom temperature (blueline) in 2013.

Fig 14. 2014: Top panel: Wind stress with direction shown in green vectors

(north up) and magnitude in purple. Bottom panel: Mid-Cape strandings (blue

bars), Outer-Cape strandings (red bars), modeled surface temperature (green

line), and observed bottom temperature (blueline) in 2014.

Figs 12-14 show the relationship between the wind stress (top panels), and the

number of strandings in "Outer Cape" and "Mid-Cape" (bottom panels) towns in

2012, 2013, and 2014, respectively. The average of the surface temperature simulated

by all the FVCOM model points (specific model sites shown in Fig 11) and the

average of the bottom temperature observed in Cape Cod Bay (specific observation

sites shown in Fig 2) are also plotted in the bottom panels. The wind stress vector is

the average simulated by FVCOM model points (sites shown in Fig 11). When the

temperature is less than 10.5 $^{\circ}$C and the wind stress from a particular direction is

greater than ~0.4 Pa, the strandings generally increase but both conditions must be

met. In the case of 2012 (Fig 12), for example, the strong wind in early November

resulted in near-zero strandings since the water temperature was still above 11°C. In

the case of 2013 (Fig 13), there was a pair of events with strong northwesterly wind

events at the end of November that coincided with a drop in temperatures resulting in

peaks in strandings around 26 November and 1 December. Note the strong winds

(>0.4Pa) in mid-December 2013 resulted in very few strandings presumably due to

the turtles having fled or already stranded by that time. In the case of 2014 (Fig 14),

the wind in mid-to-late November was not strong but it was persistently from the west

over many days with a drop in both surface and bottom temperature of a few degrees

which evidently resulted in historically high amounts of strandings.

In order to quantify the relationship between wind stress and the number of

strandings, we calculated the correlation between wind stress and the number of

strandings when the temperature was below 10.5°C. As noted above, it is not only the

strength of the wind that may be important but the persistence of the wind over time.

For this reason, we chose to calculate the "3-day sum of the windstress components" .

Fig 15 (left hand panels) shows the correlation between the number of turtle

strandings in "Outer Cape" towns and the 3-day sum of the **eastward** wind stress in

2012, 2013, 2014 and mean respectively. Fig 15 (right hand panels) show the

correlation between the number of turtles strandings in "Mid Cape" towns and the

southward wind stress in 2012, 2013, 2014, and mean respectively. In most cases, the

correlation is greater than 90% and exceeds 95% in several cases. Wind stress and the

number of strandings per three days show a linear correlation.

Fig 15. Linear correlations between eastward (left) and southward (right)

components of the wind stress and number of strandings (summed over 3 days)

in 2012 , 2013 ,2014 and mean (2012-2014) for Outer Cape (left) and Mid Cape

(right) towns.

In all of the above cases (2012-2014), it is clear that water temperature is

important as well as the wind so it is necessary to understand the change of ocean

temperature before estimating the origin of stranded turtles. Time series of surface

temperature are extracted from 10 FVCOM grid nodes across Cape Cod Bay (Fig 16).

The temperature across the 10 nodes is similar, dropping to 10.7 $^\circ$C on November 17

in both years. As is shown in this figure, there is very little spatial variability in the

surface temperature across the bay and any one monitoring site could potential work

as a marker of change. If one site has cold stun conditions, they all do.

Fig 16. Simulation of surface temperature for Cape Cod Bay and its adjacent

waters in 2012 (a) and 2013 (b). The locations a-j are plotted in Fig 11.

We backtracked turtles for 2012 and 2013[1]. Using the hourly currents from

FVCOM model, we selected the 167 turtle strandings in these years who's simulated

trajectory stayed off the beach. In other words, we focus on only those 167 cases

where the backtracking generated origins within the bay (Fig 17). We also looked at

the estimate of temperature along their track (Fig 18). In these time series we see that

a) many of the stranded turtles were exposed to <10.5 $^\circ$C water in mid-to-late

November and b) there were more December stranding events in 2012.

Fig 17. The backtracked positions of the turtles in 2012 and 2013. Red dots

show landing locations in November and December. Blue and green dots

show estimated locations of back-tracked turtles that stranded in November

and December, respectively, when they first encountered <10.5^{0}C. water.

This assumes passive (non-swimming) turtles were only transported by the

current when the modeled surface temperature is less than 10.5 ^{0}C. The blue

---

[11] Note: Since there were evidently issues with 2014 model hindcast, we chose to focus only on the previous two years in this analysis.

line is one example of backtracking a turtle that landed just south of Boston

Harbor in December. According to the model, that turtle was first cold-

stunned off Gloucester, MA.

We might ask where were the turtles that stranded in November located when

they were first exposed to <10.5 $^{\circ}$C surface waters? Our results show that turtles

stranded in November 2012 and 2013 may be cold stunned at a variety of locations

(blue dots in Fig 17) primarily in Cape Cod Bay but some in Massachusetts Bay.

Turtles stranded in December 2012 and 2013 (green dots in Fig 17) may originate

even further away, some as far as northern Massachusetts Bay. After first being

exposed to <10.5 $^{\circ}$C surface waters, the turtles are evidently in temperatures that vary

considerably but gradually decrease by a few degrees at most before stranding (Fig

18).

Fig 18. Modelled temperature of backtracked surface particles in 2012 (top

panel) and 2013 (bottom panel).

Discussion

Although the state-of-the-art coastal ocean models like FVCOM can help

simulate cold stunned turtle pathways, there are a variety of issues to consider: First,

uncertainty in the numerically simulated drifter tracks result in differences of up to

several kilometers per day relative to the observed satellite-tracked drifter tracks.

Obviously, there is significant uncertainty in the backtracking results shown in Fig 17.

The 200-meter-per-kilometer difference in modeled vs observed trajectory shown

earlier grows over long distances and time so that the numerical circulation models

and particle tracking routines will need to be improved considerably before these

particle tracking tools can be used confidently and operationally.

Simulated trajectories might also differ from actual trajectories due wind and

waves influencing cold stun turtle trajectories. There is a growing body of literature

on surface wave effects on the transport of surface particles. Much of this research

comes from the coast guard search and rescue operations [28]. Methods have been

devised to add the effects of multiples forces including direct wind effect, breaking

waves [29], and Stokes drift [30] and these will need to be considered in future work.

Additional uncertainties are introduced with the tracking methodology near the

boundary of the model grid (i.e. in shallow waters near the shoreline). We need to

further explore algorithms for forward and backward particle tracking near the coast.

Hence, it will be necessary to improve the accuracy of both the circulation models

like FVCOM with more data assimilation but we also need to improve our particle

tracking models with both the effects of wind and waves and processes near the coast.

Despite these limitations, models like FVCOM are improving every year

particularly with increasing grid resolution, incorporation of more realistic

bathymetry, and more observations to assimilate. In order to examine the effect of

different grid resolution, for example, we conducted a sensitivity study comparing

particle tracks on the "GOM3" vs "MassBay" grid (Fig 19). It can be seen here that

the particles released on the GOM3 grid stranded on the Cape Cod beaches faster than

those released on the MassBay. While the final stranding locations would likely be

similar, the particles on the latter grid were not yet stranded at the end of the week-

long simulation.

Fig 19. Comparing particle tracks on the GOM3 (left) and MassBay (right) grid

during the week of 18-25 Nov 2014.

Still more observations are needed to monitor in-situ conditions throughout Cape

Cod Bay including more realizations of the deeper currents not depicted by the

standard surface drifters used here. There is uncertainty in both the deeper currents

and the depth at which the turtles generally reside during this critical time in the fall.

As the frequency of extreme weather events is apparently on the increase [31],

the chances of turtles being suddenly surprised by wind and temperature changes will

increase. As more tropical species migrate north with warming conditions, will we see

a larger sea turtle population in Cape Cod Bay in the future and will that result in

larger strandings?

Conclusion

With some specified degree of uncertainty, the ocean current and temperatures

simulated by the FVCOM ocean model is similar to that observed by satellite-tracked

drifters and moored observations. The model can help predict the strandings of the

Kemp's ridley turtle on the shores of Cape Cod and also be used to backtrack their

source. In the forward prediction runs, the FVCOM model helped explain the

anomalously large stranding event that occurred on Outer Cape Cod between

November 18 − 23, 2014. In general, the geographic distribution of standings is

consistent with the model simulation of ocean current and wind. Given these

findings, coastal ocean models such as FVCOM can help explain the general

locations of strandings and may help volunteers to rescue turtles in the future.

In the backtracking experiments, the FVCOM model was used to estimate the

origination of stranded Kemp's ridley turtles in 2012 and 2013, showing with a

specified degree of uncertainty that most turtles stranded in November were originally

located in a variety of locations in both Cape Cod Bay and Massachusetts Bay when

they were first exposed to cold-stun temperatures (<10.5 ^{o}C). The turtles stranded in

December may have originated well outside Cape Cod Bay just a few weeks earlier.

There are still many uncertainties as to when the turtles are cold-stunned but a

simple regression of different components of the wind stress (with water temperatures

below the 10.5^{o}C threshold) vs number of strandings indicate that values greater than

0.4 Pa will induce more stranding events (r^2=95). As expected, strong winds out of

the west will result in Outer Cape strandings and strong winds out of the north will

result in Mid-Cape strandings.

Acknowledgements

Thanks to all those who have helped with this analysis, the many turtle rescue

volunteers, the staff at Mass Audubon's Wellfleet Bay Wildlife Sanctuary, Dr. Vitalli

Sheremet for help in coding, an anonymous NEFSC reviewer, and Dr. Dai Honglei

for long term encouragement and support. The Massachusetts Division of Marine

Resources provided some of their moored temperature observations from Cape Cod

Bay and the UMASS Dartmouth FVCOM modeling group provided on-line access to

their simulations.

References

1. Plotkin, PT. National Marine Fisheries Service and U. S. Fish and Wildlife
Service Status Reviews for Sea Turtles Listed under the Endangered Species
Act of 1973. National Marine Fisheries Service, Silver Spring, Maryland, 2009.
109-116.

2. National Marine Fisheries Service, U.S. Fish and Wildlife Service, and
Secretaría de Medio Ambiente y Recursos Naturales . Bi-national Recovery
Plan for the Kemp's ridley Sea Turtle (Lepidochelys kempii). Second Revision.
2011. National Marine Fisheries Service, Silver Spring, Maryland, USA.

3. U.S. Fish and Wildlife Service. List of endangered foreign fish and wildlife.
Federal Register 1970;35:18319−18322.

4. McClellan, CM, Read AJ, Cluse WM, and Godfrey MH. Using telemetry to
mitigate the bycatch of long-lived marine vertebrates. Ecological Applications
2009;19(6):1660-1671.

5. Putman, NF and Mansfield KL. 2015. Direct evidence of swimming
demonstrates active dispersal in the sea turtle "lost years." Current Biology
25:1-7.

6. Shaver, DJ, Hart KM, Fujisaki I, Rubio C, Sartain AR, Pena J, et al Foraging
area fidelity for Kemp's ridleys in the Gulf of Mexico. Ecology and Evolution
2013;3(7):2002-2012.

7. Shaver, DJ, Schroeder BA, Byles RA, Burchfield PM, Peña J, Márquez R, et
al. Movements and home ranges of adult male Kemp's ridley sea turtles
(Lepidochelys kempii) in the Gulf of Mexico investigated by satellite telemetry.
Chelonian Conserv. Biol. 2005;4:817−827.

8. Shaver, DJ, Rubio C. Post-nesting movement of wild and head-started Kemp's

ridley sea turtles (Lepidochelys kempii) in the Gulf of Mexico. Endanger.

Species Res. 2008;4:43–55.

9. Bleakney, JS. Four records of the Atlantic ridley turtle, Lepidochelys kempi,

from Nova Scotia. Copeia 1955;2:137.

10. Kemp's ridley Sea Turtle. Lepidochelys kempi. URL: http://www.docin.com/p-

984318341.html.

11. Bianca SS, David MK, Marjorie A.M., Susan GB, Katherine LM, James PM.

Consequences of drift and carcass decomposition for estimating sea turtle

mortality hotspots. 2017;Ecological Indicators,84:391-396.

12. Keinath, JA, Musick JA, Byles RA. 1987. Aspects of the biology of Virginia's

sea turtles: 1979-1986. Virginia J. Sci. 38(4):329-336.

13. Byles, RA 1988. Behavior and ecology of sea turtles from Chesapeake Bay,

Virginia. Ph.D. dissertation, College of William and Mary, Virginia Institute of

Marine Science, Gloucester, VA.

14. Blair EW, Ehrhart LM. Hypothermic Stunning and Mortality of Marine Turtles

in the Indian River Lagoon System, Florida. American Society of Ichthyologists

and Herpetologists. 1989;3:696-703.

15. Schwartz FJ. Behavioral and tolerance responses to cold water temperatures by

three species of seaturtles (Reptilia,Chelonidae) in North Carolina. Fla Mar Res

Publ.1978;33:16-18 .

16. Morreale SJ, Meylan AB, Sadove SS, Standora EA, et al. Annual Occurrence

and Winter Mortality of Marine Turtles in New York Waters. Journal of

Herpetology, 1992;26(3):301–308. JSTOR, www.jstor.org/stable/1564885.

17. Burke VJ, Standora EA, Morreale SJ. Factors Affecting Strandings of Cold-

Stunned Juvenile Kemp's ridley and Loggerhead Sea Turtles in Long Island,

New York. Copeia, 1991;4:1136–1138. www.jstor.org/stable/1446115.9.

18. Still BM, Griffin CR, Prescott R. Climatic and oceanographic factors affecting

daily patterns of juvenile sea turtle cold-stunning in Cape Cod Bay,

Massachusetts. Chelonian Conserv Biol. 2005; 4: 883–89 0

19. Gorman J. A Surge of Stranded Turtles. The New York Times. URL:

https://www.nytimes.com/2014/12/13/science/a-cape-cod-mystery-hundreds-of-

sea-turtles-stranded-on-beaches.html. 2014.

20. Chen C, Liu H., Beardsley RC. An unstructured, finite-volume, three-

dimensional, primitive equation ocean model: application to coastal ocean and

estuaries. J. Atm.&Oceanic Tech., 2003a;20: 159-186.

21. Chen C, Beardsley RC, Cowles G, Qi J, Lai Z, Gao G, et al. An Unstructured

Grid, Finite-Volume Coastal Ocean Model FVCOM User Manual. 2011.

SMAST/UMASSD-11-1101.

22. The unstructured Grid Finite Volume Community Ocean Model (FVCOM).

2014. URL: http://fvcom.smast.umassd.edu/2014/01/17/1-research-gulf-model-

domain/

23. Lynch DR, Greenberg DA, Bilgili A , McGillicuddy DJ, Manning JP,

Aretxabaleta AL . Particles in the Coastal Ocean Theory and Applications.

Cambridge University Press, America. 2015, pp. 20-21

24. Poulain P. Drifter observations of surface circulation in the Adriatic Sea

between December 1994 and March 1996. Journal of Marine Systems, 1999;20:

231-253.

25. Manning JP, McGillicuddy DJ, Pettigrew N, Churchill JH, Incze L. Drifter
Observations of Maine Coastal Current Drift. Continental Shelf   Res.
2009;doi:10.1016/j.csr.2008.12.008.

26. The GoMOOS Moored Buoy Design. URL:
http://gyre.umeoce.maine.edu/GoMoos/oceans2003.pdf.

27. Manning JP, Pelletier E. Environmental Monitors on Lobster Traps: longterm
temperature observations from New England's bottom waters. Jour. of
Operational Oceanography. 2009;2:25-33.

28. Breivik Ø, Allen AA, Maisondieu C and Roth JC. Wind-induced drift of
objects at sea: the leeway field method. 2011 doi:10.1016/j.apor.2011.01.005

29. Carniel S, Warner JC, Chiggiato J, and Sclavo M. Investigating the impact of
surface wave breaking on modeling the trajectories of drifters in the Northern
Adriatic Sea during a wind-storm event. Ocean Modelling. 2009;30:225-239.

30. Röhrs J, Christensen, KH, Hole, LR, et al. Observation-based evaluation of
surface wave effects on currents and trajectory forecasts. Ocean Dynamics
2012;62: 1519-1533.

31. Vose, RS. and Coauthors. Monitoring and understanding changes in extremes:
Extratropical storms, winds, and waves. *Bull. Amer. Meteor. Soc.* 2014;**95**:377–
386. doi:https://doi.org/10.1175/BAMS-D-12-00162.1.

CODE SURFACE DRIFTER

8.9 cm

Flotation line

100.33 cm

63.5 cm

26.67 cm

100.97 cm

11.43 cm

Weight = 8.4 Kg

2012 wind stickplot

a

2012 temperture vs strandings

Nov 16 2012

bottom tempe
surface tempe
#strandings/d
#strandings/d

Nov 12 2012   Nov 19 2012   Nov 26 2012   Dec 03 2012   Dec 10 2012   Dec 17 2012   Dec

b

2013 wind stickplot

a

2013 temperture vs strandings

- bottom temp
- surface temp
- #strandings/d
- #strandings/d

Nov 13 2013

Nov 11 2013   Nov 18 2013   Nov 25 2013   Dec 02 2013   Dec 09 2013   Dec 16 2013   Dec 23

b

2014 wind stickplot

a

2014 temperture vs strandings

Legend:
- bottom tempe...
- surface tempe...
- #strandings/da...
- #strandings/da...

Nov 15 2014

Nov 10 2014   Nov 17 2014   Nov 24 2014   Dec 01 2014   Dec 08 2014   Dec 15 2014   Dec 22 201

b

Eastward wind stress vs strandings on Outer Cape towns

2012
$r^2=0.94$

2013
$r^2=0.98$

2014
$r^2=0.75$

mean(2012-2014)
$r^2=0.56$

sum of eastward wind stress per 3 days (pa)

Northward wind stress vs strandings on

2012
$r^2=-0.99$

2013
$r^2=-0.95$

2014
$r^2=-0.96$

mean(2

sum of northward wind stress per 3

2012

2013

11/25   12/4   12/12   12/20   12/29 11/1   11/9   11/17   11/25

a

b

the temperature along the trajectory

Legend:
- ● Temperature after stran...
- ● Pre-freezing temperatur...

2012

2013