High dispersal levels and lake warming are emergent drivers of cyanobacterial community assembly over the Anthropocene in peri-Alpine lakes

Authors:

Marie-Eve Monchamp me.monchamp@gmail.com, ORCIDiD: 0000-0001-5361-5162 [1,2];

Piet Spaak spaak@eawag.ch ORCIDiD: 0000-0003-0631-373X[1,2];

Francesco Pomati francesco.pomati@eawag.ch, ORCIDiD: 0000-0002-9746-3735 [1,2,#]

[1] Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department of Aquatic Ecology, 8600 Dübendorf, Switzerland;

[2] Swiss Federal Institute of Technology (ETH) Zürich, Institute of Integrative Biology, 8092 Zürich, Switzerland.

 # Corresponding author address:

Eawag, Department of Aquatic Ecology

Überlandstrasse 133, 8600 Dübendorf, Switzerland.

Email: francesco.pomati@eawag.ch

Phone: +41 58 765 5410

Short title: Cyanobacterial community assembly patterns

Keywords: Beta diversity, community phylogenetic structure, stochastic assembly, environmental forcing, meta-community, peri-Alpine lakes, eutrophication, climate change

Disentangling the relative importance of deterministic and stochastic processes in shaping

natural communities is central to ecology. Studies on community assembly over broad

temporal and spatial scales in aquatic microorganisms are scarce. Here, we used 16S rDNA

sequence data from lake sediments to test for community assembly patterns in

cyanobacterial phylogenies across ten European peri-Alpine lakes and over a century of

eutrophication and climate warming. We studied phylogenetic similarity in cyanobacterial

assemblages over spatial and temporal distance, and environmental gradients, comparing

detected patterns with theoretical expectations from deterministic and stochastic processes.

We found limited evidence for deviation of lake communities from a random assembly model

and no significant effects of geographic distance on phylogenetic similarity, suggesting no

dispersal limitation and high levels of stochastic assembly. We did not detect a significant

effect of phosphorus and nitrogen levels on deviation of community phylogenies from

random. We found however a significant decay of phylogenetic similarity for non-random

communities over a gradient of air temperature and water column stability. We show how

phylogenetic data from sedimentary archives can improve our understanding of microbial

community assembly processes, and support previous evidence that climate warming has

been the strongest environmental driver of cyanobacterial community assembly over the

Anthropocene.

Introduction

Understanding the mechanisms that determine changes in the structure and composition of natural communities over large spatial and temporal scales is critical, given the impacts that human activities have on biodiversity and ecosystem functions [1]. The relative importance of stochastic and deterministic processes driving community assembly might vary over space and time: dispersal, demography, ecological interactions and evolutionary processes can all influence the structure of natural communities across scales [2-7]. It is an on-going challenge to understand how anthropogenic environmental changes influence ecological and evolutionary mechanisms determining community assembly, particularly in aquatic microbes whose dispersal appears to have no boundaries [8,9].

Assembly studies focusing on ecological mechanisms in lake cyanobacterial communities have been scarce due to a lack of data at the appropriated spatial and temporal scale, despite the importance that these organisms have reached over the past decades for freshwater ecosystem functioning and services [10]. Over the last century, the frequency and severity of cyanobacteria-dominated communities have increased in lakes and reservoirs worldwide despite remediation measures applied at the regional and international scale [11,12]. Cyanobacterial blooms are often dominated by toxic species, and there is a global concern that environmental changes are promoting the geographic expansion of some potentially harmful taxa [13,14], due to a combined effect of increasing temperature and nutrient loads [11,15,16]. Toxic species such as *Dolichospermum lemmermannii* and *Planktothrix rubescens* have indeed widened their geographic distribution, supporting the idea that some harmful cyanobacteria are spreading across temperate lakes [16]. The role of geographic dispersal (where distance limits the establishment of new taxa) relative to turnover of taxa driven by environmental gradients has not been explicitly explored in the assembly of these globally important microorganisms.

In this study, we analysed cyanobacterial community composition data spanning over a hundred years and across ten lakes. We used 16S rDNA sequences from sediment cores of

European peri-Alpine lakes (Fig. S1) that underwent directional environmental change

characterised by climate warming and eutrophication [16]. Our previous work has explored the

patterns of long term change in alpha and beta diversity in lake cyanobacterial communities,

showing a homogenization of assemblage composition at the regional scale [16]. The aim of

this study was to test for emergent deterministic (environment-driven) and stochastic

(dispersal-driven) patterns in the phylogenetic structure of cyanobacterial assemblages

across these different lakes of the same region [2,17].

We used a null-model that accounted for temporal changes in the size of the species pool to

simulate random assembly. We then tested for deviation from random patterns as

phylogenetic clustering and overdispersion: the tendency for taxa to co-occur with larger or

smaller expectancy, respectively, than predicted by the null-model (Fig. 1A) [18-22]. In most

cases, dispersal-driven assembly would generate random taxa co-occurrence patterns, while

environmental drivers would lead to deviation from random assembly [19,21,23,24]. There can be

interactions among assembly mechanisms that generate exceptions to these predictions [25,26].

We however expect that comparison of phylogenetic structures to null-model simulations,

combined with the patterns of community phylogenetic similarity across lakes and spatial or

ecological distance, will allow us testing for deterministic and stochastic signatures in

cyanobacterial community assembly.

Specifically, when dispersal of cyanobacterial taxa among lakes is not limited (Fig. 1B), we

expect that phylogenetic community similarity will decrease over an environmental gradient,

while no change is expected when the system is driven only by dispersal (Fig. 1C) [2]. If there

are barriers to dispersal of cyanobacteria, we predict differences in similarities among lake

communities that are only dependent on the geographic distance (Fig. 1B), and no effects

driven by ecological gradients (Fig. 1C) [2]. The environmental-driven decrease in phylogenetic

community similarity will not be influenced by dispersal limitation (Fig. 1B), and will vary

deterministically as a consequence of the gradient itself (Fig. 1C) [2]. This is because we

expect that, under environment-driven assembly, the turnover of taxa along the ecological

gradient will determine community structure in each lake. Here, we investigated whether

cyanobacterial community phylogenetic structures within and across lakes over time

matched these expectations from assembly processes, and what patterns dominate.

Results

Community phylogenetic structure. We calculated a standardised effect sizes (SES) of

mean-pairwise distances (MPD) within each local community based on the comparison of the

observed MPD values with the values a randomly assembled community (Methods). Based

on the SES_{MPD} values, a large number of cyanobacterial communities did not show a

phylogenetic structure that significantly differed from the null (random) expectation (Fig. 2). In

fact, only two communities were significantly phylogenetically clustered, whereas eighteen

out of seventy-six communities showed significant phylogenetic overdispersion, with SES_{MPD}

values outside the 90% confidence interval of the null model simulation. Although the

remaining forty-six communities analysed did not show significant signal of non-randomness,

most (especially since the 1980s) of the SES_{MPD} values were negative, suggesting a

tendency towards overdispersion. The samples that displayed significant non-random

phylogenetic structure and their corresponding MPD and SES_{MPD} values are listed in Table

S2 of the Supplementary Materials.

Distance-decay relationships. We estimated the phylogenetic similarity (beta-diversity)

across all pairs of communities reconstructed from the sedimentary archives of the ten lakes

at each time-period and investigated the role of geographical and temporal distance (Fig. 3).

Our analysis did not reveal an increase in phylogenetic beta-diversity with geographic

distance (Fig. 3A), suggesting no dispersal limitation of cyanobacteria at the regional (peri-

Alpine) scale. On the contrary, we observed for all lakes a clear decay of phylogenetic

similarity along the temporal gradient representing the history of each lake (Fig. 3B), with the

exception of lakes Geneva and Annecy (the latter due to insufficient data points) (Fig. S2).

Community similarity over environmental gradients. All non-random samples identified in

Fig. 2 using the SES$_{MPD}$ metric were used to investigate the role of chemical (TP and NO$_3^-$-N)

and physical (air temperature and water column stability) drivers in explaining cyanobacterial

community deviation from a random assembly. We found no evidence for a role of nutrients

(TP and NO$_3^-$-N) in explaining non-random community structure (Fig. 4). On the other hand,

OLS regression showed a significant decrease in community similarity along with both air

temperature (p = 0.0003, adjusted R^2 = 0.057) and SSI gradients (p = 0.019, adjusted R^2 =

0.058) (Fig. 4). This suggests that communities in lakes characterized by similar physical

characteristics related to lake water temperature are more similar in cyanobacterial

community composition compared to lakes that display greater differences in temperature

and stratification.

Discussion

Most of the cyanobacterial communities obtained in this study from sedimentary archives did

not show significant deviation from a random phylogenetic structure, suggesting a strong

signal of stochastic (dispersal-driven) community assembly in lake cyanobacteria. Our

previous work has shown that DNA-based reconstructions of cyanobacterial communities are

robust [16,27], therefore the observed patterns are unlikely to be driven by biases in

sedimentary DNA-based community reconstructions. On the contrary, it has been noted that

a strong effect of stochastic relative to deterministic processes is expected in the community

assembly of microalgae, particularly in highly productive (eutrophic) environments [2]. The

decay in phylogenetic similarity over time coupled with the lack of a geographic distance-

decay relationship across lake communities (Fig. 3) suggest temporally dynamic

communities with no limitation to dispersal at the regional (peri-Alpine) scale. In a recent

study on genetic divergence among populations of a marine diatom, no significant

relationship was found between genetic and geographic distances at regional and global

scales [9]. Most reports about microbial dispersal so far did no show clear evidence for

geographic distance-decay patterns at the local (0 - 100 km) and regional (101 - 5,000 km)

scales [28]. The scale of distances in our study was not suited to capture dissimilarity changes

among cyanobacterial communities along very large geographical distances (e.g.

continental), where an effect of geographical isolation might emerge [29]. Nevertheless, our

research suggests that cyanobacterial communities present weak dispersal limitation among

lakes of the same region, even around and across barriers such as the European Alpine

mountain range.

Previous work has shown that communities of cyanobacteria have become more

homogeneous in terms of composition across peri-Alpine lakes over the last decades, in

favour of a few clades of bloom-forming and potentially toxic taxa [14,16]. This could result in an

increase of phylogenetic clustering over time, if the traits under selection by environmental

changes are phylogenetically conserved [19]. Our data show that the phylogenetic structure of

assemblages is, in most cases, not clustered. Only two out of seventy-six tested

communities displayed significant clustering, which strongly suggests that the environmental

driving forces of cyanobacterial assembly in the lakes are not acting on phylogenetically

conserved traits. Coloniality and buoyancy regulation are in fact multiphyletic traits that have

been associated to the invading taxa, which belong to different cyanobacterial phylogenetic

lineages within the orders *Chroococcales*, *Nostocales* and *Oscillatoriales* [10,11,16]. The more

prevalent signal of overdispersion in the community phylogenies might support this

hypothesis of environmental selection for convergent traits.

Gradients of NO_3^--N and TP concentrations across lakes did not significantly explain

deviation from random assembly in the investigated cyanobacterial communities (Fig. 4). It is

important to note that most of the lakes investigated here classify as meso-eutrophic to

eutrophic [30] (see average P and NO_3^--N concentrations reported by [16]). Significant patterns of

phylogenetic dissimilarity might emerge across communities characterized by a broader

nutrient gradient, but this remains to be tested by surveys or experimentally. Rather, the

difference in physical conditions, such as temperature and strength of the water column

stratification, seemed in our study to explain a significant proportion of the observed variance

in phylogenetic relatedness among lake cyanobacterial communities. The increasing trend in

air temperatures, which has accelerated since the 1980s across the peri-Alpine region, has

caused modifications in thermal characteristics of lakes, e.g., via changes in the duration and

strength of the water column stratification [16,31], which in turn affects recirculation and

availability of nutrients for phytoplankton growth [10,22,32]. This effect has been amply

documented and has favoured, as mentioned above, buoyant cyanobacterial forms that are

able to control their vertical position in the water column to reach optimal nutrient and light

conditions [14,16,22,31,32]. Our findings therefore support previous evidence and suggest that

climate warming is the strongest environmental driver of the assembly of lake cyanobacterial

communities [33], and might select for specific traits such as, for example, buoyancy regulation.

In conclusion, this is the first study to our knowledge that explicitly tests for deterministic and

stochastic assembly patterns in cyanobacterial communities across regional scales and over

the Anthropocene. We successfully reconstructed the phylogenetic diversity in seventy-six

cyanobacterial communities over a period of ca. 100 years using DNA from sediment cores

collected in ten lakes. Our study shows that both stochastic (dispersal-driven) and

deterministic (environmental-driven) processes are important in assembling cyanobacterial

communities across lakes of the European peri-Alpine region. Cultural eutrophication and

climate change are the most notable environmental factors favouring cyanobacterial growth,

but the deterministic processes governing community assembly appeared in our study to be

more significantly driven by lake warming. Our results confirm previous evidence [16,33] and

expand our understanding of cyanobacterial community assembly processes. Knowledge

about the relative importance of (potentially controllable) environmental drivers and (likely

uncontrollable) dispersal of organisms in shaping the structure of cyanobacterial

assemblages is important for the management of aquatic ecosystems whose services are

threatened by an increasing prevalence of potentially toxic taxa.

Materials and methods

Data collection

We used high-resolution 16S rDNA sequence data from ten European peri-Alpine lakes,

spanning between early 1900s and 2016, to estimate phylogenetic diversity of cyanobacterial

communities [16]. Briefly, sediment cores were collected in ten lakes between 2013 and 2016

using a gravity corer, and layers were dated by varve counting and, in most cases, with

radionuclide (Pb^{210}, Cs^{237}) measurements [16,27]. Based on the sediment age models, sediment

sub-samples were collected at various depths in cores from each lake to capture the

cyanobacterial community composition over the last ~100 years. DNA was extracted from

bulk sediments in a clean laboratory facility, and the DNA extract were used for PCR and

high-throughput sequencing of the cyanobacterial 16S rRNA gene (Table S1) on a MiSeq

Illumina platform as previously described [16,27].

The clean, primmer-trimmed sequences were clustered in operational taxonomic units

(OTUs) with a 97% threshold of sequence similarity in QIIME [34] using the UPARSE workflow

[35]. PyNast [34] and the Greengenes microbial sequence database [36] were used for sequence

alignment, and FastTree [37] was used to estimate a phylogeny based on maximum-likelihood

containing all OTUs found in the lakes. OTUs were taxonomically assigned with a confidence

threshold of 85% and the ones assigned to phyla other than photosynthetic cyanobacteria

were removed from the dataset. Each sample was rarefied to 2,744 sequences

(cyanobacteria only) prior to phylogenetic analyses.

The physical (air temperature) and chemical (nitrogen [NO_3^--N], total phosphorus [TP]) data

were collected over several decades consist of several decades of monitoring of the ten

lakes [16] and the annual maximal Schmidt Stability Index (SSI; the maximal strength of the

water column stratification) was derived from water temperature and hypsometry data [16,38].

Euclidean distances for each environmental variable were calculated among lakes to derive

environmental gradients used in the linear models in the R package 'vegan' version 2.4.4 [39].

Phylogenetic analyses

To derive the phylogenetic structure of each community, we quantified the mean pairwise

phylogenetic distance (MPD) [40] using *mpd* and *ses.mpd* in the package 'picante' version

1.6.2 for R [41] and used null-model simulations of random assembly that account for temporal

changes in the size of the species pool [21]. We calculated a standardised effect size of MPD

(SES_{MPD}) within each local community subtree based on the comparison of the observed

MPD values with the values in the random distribution using 999 randomisations of the

species at the tip of the phylogenetic tree, while species richness was maintained [41-43]:

SES_{MPD} = Mean($MPD_{Observed}$ − $MPD_{Randomized}$)/SD($MPD_{Randomized}$). The SES_{MPD} values were

multiplied by −1 to be equivalent to the net relatedness index (NRI) [40].

We derived the Unifrac phylogenetic distance [44] between all pairs of samples using the *dist*

function in the Bioconductor package 'phyloseq' [45] based on the OTU table and the fasta files

from amplicon sequencing. The geographic distance-decay relationship was measured on

binned communities each representing a period of one decade (from the 1930s to the 2010s;

decades 1900s, 1910s and 1920s were excluded due to insufficient number of samples). The

binning was done to remove the factor time from the analysis, as it would introduce a bias

when comparing multiple samples from single lakes over time. The temporal distance-decay

pattern of phylogenetic similarity was studied by plotting log-transformed Unifrac distances

against natural log-transformed time distances (years) for each lake in the dataset.

Significance of the distance-decay relationship at each decade was tested using Mantel tests

in the R package 'ade4' with a significance threshold of $p \leq 0.05$. To test whether the

environment (physical and chemical parameters) play a role in driving community assembly,

we used the samples that were identified as phylogenetically non-random (i.e., those which

SES_{MPD} values were outside the 90% confidence interval of the null model simulation) in

linear ordinary least square (OLS) regressions where physical and chemical lake data were

the explanatory variables.

Acknowledgments

255   This work was supported by the Swiss Enlargement Contribution, project IZERZ0 – 142165

256   'CyanoArchive' to P.S. in the framework of the Romanian–Swiss Research Programme.

Authors contributions

M-E.M., F.P. and P.S. designed the study. M-E.M. collected data and performed data analysis. M-E.M. and F.P. wrote the manuscript, which was commented and edited by P.S.

Conflict of Interest

The authors declare no competing financial interests.

References

Cardinale, B. J. *et al.* Biodiversity loss and its impact on humanity. *Nature* **486**, 59-67, doi:10.1038/nature11148 (2012).

Chase, J. M. Stochastic community assembly causes higher biodiversity in more productive environments. *Science (New York, N.Y.)* **328**, 1388-1391, doi:10.1126/science.1187820 (2010).

Vergnon, R., Van Nes, E. H. & Scheffer, M. Emergent neutrality leads to multimodal species abundance distributions. *Nature Communications* **3**, 663-666, doi:10.1038/ncomms1663 (2012).

Siepielski, A. M., Hung, K.-L., Bein, E. E. B. & McPeek, M. a. Experimental evidence for neutral community dynamics governing an insect assemblage. *Ecology* **91**, 847-857 (2010).

Segura, A. M. *et al.* Emergent neutrality drives phytoplankton species coexistence. *Proceedings of The Royal Society of London B* **278**, 2355-2361, doi:10.1098/rspb.2010.2464 (2011).

Wagner, C. E., Harmon, L. J. & Seehausen, O. Ecological opportunity and sexual selection together predict adaptive radiation. *Nature* **487**, 366-369, doi:10.1038/nature11144 (2012).

Hillerislambers, J., Adler, P. B., Harpole, W. S., Levine, J. M. & Mayfield, M. M. Rethinking Community Assembly through the Lens of Coexistence Theory. *Annu. Rev. Ecol. Evol. Syst* **43**, 227-248, doi:10.1146/annurev-ecolsys-110411-160411 (2012).

Patterson, D. J. Seeing the big picture on microbe distribution. *Science* **325**, 1506-1507, doi:10.1126/science.1179690 (2009).

Whittaker, K. A. & Rynearson, T. A. Evidence for environmental and ecological selection in a microbe with no geographic limits to gene flow. *Proceedings of the National Academy of Sciences* **114**, 2651-2656, doi:10.1073/pnas.1612346114 (2017).

290   10   Huisman, J. *et al.* Cyanobacterial blooms. *Nature Reviews Microbiology,* online ahead
291   of print-online ahead of print, doi:10.1038/s41579-018-0040-1 (2018).

292   11   Carey, C. C., Ibelings, B. W., Hoffmann, E. P., Hamilton, D. P. & Brookes, J. D. Eco-
293   physiological adaptations that favour freshwater cyanobacteria in a changing climate.
294   *Water Research* **46**, 1394-1407, doi:10.1016/j.watres.2011.12.016 (2012).

295   12   Taranu, Z. E., Zurawell, R. W., Pick, F. & Gregory-Eaves, I. Predicting cyanobacterial
296   dynamics in the face of global change: the importance of scale and environmental
297   context. *Global Change Biology* **18**, 3477-3490, doi:10.1111/gcb.12015 (2012).

298   13   Sinha, R. *et al.* Increased incidence of Cylindrospermopsis raciborskii in temperate
299   zones - Is climate change responsible? *Water Research* **46**, 1408-1419,
300   doi:10.1016/j.watres.2011.12.019 (2012).

301   14   Salmaso, N., Capelli, C., Shams, S. & Cerasino, L. Expansion of bloom-forming
302   Dolichospermum lemmermannii (Nostocales, Cyanobacteria) to the deep lakes south
303   of the Alps: colonization patterns, driving forces and implications for water use.
304   *Harmful Algae* **50**, 76-87, doi:10.1016/j.hal.2015.09.008 (2015).

305   15   Rigosi, A., Carey, C. C., Ibelings, B. W. & Brookes, J. D. The interaction between
306   climate warming and eutrophication to promote cyanobacteria is dependent on
307   trophic state and varies among taxa. *Limnology and Oceanogaphy* **59**, 99-114,
308   doi:10.4319/lo.2014.59.01.0099 (2014).

309   16   Monchamp, M.-E. *et al.* Homogenization of lake cyanobacterial communities over a
310   century of climate change and eutrophication. *Nature Ecology & Evolution* **2**, 317–
311   324, doi:10.1038/s41559-017-0407-0 (2018).

312   17   Ruokolainen, L., Ranta, E., Kaitala, V. & Fowler, M. S. When can we distinguish
313   between neutral and non-neutral processes in community dynamics under ecological
314   drift? *Ecology Letters* **12**, 909-919, doi:10.1111/j.1461-0248.2009.01346.x (2009).

315   18   Gotelli, N. J. & McCabe, D. J. Species co-occurrence: A meta-analysis of J. M.
316   Diamond's assembly rules model. *Ecology* **83**, 2091-2096, doi:10.1890/0012-
317   9658(2002)083[2091:SCOAMA]2.0.CO;2 (2002).

318   19   Emerson, B. C. & Gillespie, R. G. Phylogenetic analysis of community assembly and
319   structure over space and time. *Trends in Ecology and Evolution* **23**, 619-630,
320   doi:10.1016/j.tree.2008.07.005 (2008).

321   20   Cadotte, M., Albert, C. H. & Walker, S. C. The ecology of differences: assessing
322   community assembly with trait and evolutionary distances. *Ecology letters* **16**, 1234-
323   1244, doi:10.1111/ele.12161 (2013).

324   21   Kathleen Lyons, S. *et al.* Holocene shifts in the assembly of plant and animal
325   communities implicate human impacts. *Nature* **529**, 80-83, doi:10.1038/nature16447
326   (2016).

22 Pomati, F., Matthews, B., Seehausen, O. & Ibelings, B. W. Eutrophication and climate
warming alter spatial (depth) co-occurrence patterns of lake phytoplankton
assemblages. *Hydrobiologia* **787**, 375-385, doi:10.1007/s10750-016-2981-6 (2017).
23 Cavender-Bares, J., Kozak, K. H., Fine, P. V. A. & Kembel, S. W. The merging of
community ecology and phylogenetic biology. *Ecology Letters* **12**, 693-715,
doi:10.1111/j.1461-0248.2009.01314.x (2009).
24 Kraft, N. J. B. *et al.* Disentangling the drivers of β diversity along latitudinal and
elevational gradients. *Science* **333**, 1755-1758, doi:10.1126/science.1208584 (2011).
25 Mayfield, M. M. & Levine, J. M. Opposing effects of competitive exclusion on the
phylogenetic structure of communities. *Ecology letters* **13**, 1085-1093,
doi:10.1111/j.1461-0248.2010.01509.x (2010).
26 Bell, T., Newman, J. A., Silverman, B. W., Turner, S. L. & Lilley, A. K. The contribution
of species richness and composition to bacterial services. *Nature* **436**, 1157-1160,
doi:10.1038/nature03891 (2005).
27 Monchamp, M.-E., Walser, J.-C., Pomati, F. & Spaak, P. Sedimentary DNA reveals
cyanobacteria community diversity over 200 years in two peri-alpine lakes. *Applied*
and Environmental Microbiology **82**, 6472-6482, doi:10.1128/AEM.02174-16 (2016).
28 Hanson, C. A., Fuhrman, J. A., Horner-Devine, M. C. & Martiny, J. B. H. Beyond
biogeographic patterns: processes shaping the microbial landscape. *Nature reviews.*
Microbiology **10**, 497-506, doi:10.1038/nrmicro2795 (2012).
29 Papke, R. T., Ramsing, N. B., Bateson, M. M. & Ward, D. M. Geographical isolation in
hot spring cyanobacteria. *Environmental Microbiology* **5**, 650-659, doi:10.1046/j.1462-
2920.2003.00460.x (2003).
30 Oecd. *Eutrophication of waters. Monitoring, assessment and control. Final report,*
OECD cooperative programme on monitoring of inland waters (eutrophication
control). (Environment Directorate, OECD, 1982).
31 Posch, T., Köster, O., Salcher, M. M. & Pernthaler, J. Harmful filamentous
cyanobacteria favoured by reduced water turnover with lake warming. *Nature Climate*
Change **2**, 809-813, doi:10.1038/nclimate1581 (2012).
32 Yankova, Y., Neuenschwander, S., Köster, O. & Posch, T. Abrupt stop of deep water
turnover with lake warming: Drastic consequences for algal primary producers.
Scientific Reports **7**, 13770-13770, doi:10.1038/s41598-017-13159-9 (2017).
33 Mantzouki, E. *et al.* Temperature Effects Explain Continental Scale Distribution of
Cyanobacterial Toxins. *Toxins (Basel)* **10**, doi:10.3390/toxins10040156 (2018).
34 Caporaso, J. G. *et al.* PyNAST: A flexible tool for aligning sequences to a template
alignment. *Bioinformatics* **26**, 266-267, doi:10.1093/bioinformatics/btp636 (2010).

35 Edgar, R. C. UPARSE: highly accurate OTU sequences from microbial amplicon
reads. *Nature methods* **10**, 996-998, doi:10.1038/nmeth.2604 (2013).
36 McDonald, D. *et al.* An improved Greengenes taxonomy with explicit ranks for
ecological and evolutionary analyses of bacteria and archaea. *The ISME Journal* **6**,
610-618, doi:10.1038/ismej.2011.139 (2012).
37 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 - Approximately maximum-
likelihood trees for large alignments. *PLoS ONE* **5**, e9490-e9490,
doi:10.1371/journal.pone.0009490 (2010).
38 Schmidt, W. Über die Temperatur- und Stabilitätsverhältnisse von Seen. *Geografiska*
Annaler **10**, 145-177 (1928).
39 Oksanen, J. *et al.* vegan: Community Ecology Package. R package version 2.0-10.
http://CRAN.R-project.org/package=vegan (2013).
40 Webb, C. O. Exploring the phylogenetic structure of ecological communities: An
example for rain forest trees. *The American Naturalist* **156**, 145-155,
doi:10.1086/303378 (2000).
41 Swenson, N. G. *Functional and phylogenetic ecology in R.* (Springer, 2014).
42 Erickson, D. L. *et al.* Comparative evolutionary diversity and phylogenetic structure
across multiple forest dynamics plots: a mega-phylogeny approach. *Frontiers in*
genetics **5**, 358-358, doi:10.3389/fgene.2014.00358 (2014).
43 Hardy, O. J. & Senterre, B. Characterizing the phylogenetic structure of communities
by an additive partitioning of phylogenetic diversity. *Journal of Ecology* **95**, 493-506,
doi:10.1111/j.1365-2745.2007.01222.x (2007).
44 Lozupone, C. & Knight, R. UniFrac: a new phylogenetic method for comparing
microbial communities. *Applied and environmental microbiology* **71**, 8228-8235,
doi:10.1128/AEM.71.12.8228-8235.2005 (2005).
45 McMurdie, P. J. & Holmes, S. Phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS ONE* **8**,
doi:10.1371/journal.pone.0061217 (2013).

 Figure legends

Figure 1. Schematic description of the theoretical expectations for phylogenetic community

assembly within and between communities, based on stochastic (dispersal-driven) and

deterministic (environment-driven) processes. **(A)** Taxa associations are analysed using their

mean pairwise phylogenetic distance (MPD) for each date of each time-series and by

comparing it to expected patterns from null-model simulations of random assembly (grey

box): clustering and overdispersion (above and below the random expectation, respectively)

signal communities that are composed of species phylogenetically closer or further apart

than expected by chance, respectively, as a sign of deterministic processes. **(B)** Predicted

patterns in phylogenetic community similarity depending on limitation (black line) and no

limitation (grey line) in taxa dispersal among sites. **(C)** Predicted change in phylogenetic

similarity for completely stochastic (grey) and deterministic (black) models of community

assembly along an ecological environmental gradient (e.g. lake physics and chemistry).

Figure 2. Time series of the standardised effect size of mean pairwise phylogenetic

distances (SES$_{MPD}$) values multiplied by −1 for the equivalent of the NRI index [40], calculated

for each local community compared with null model simulations. The samples outside the

central area delimited by the dashed lines show significant community structure (above or

below the 95% [grey lines] and 90% [red lines] confidence intervals of the null-model

simulation). Negative values signal phylogenetic overdispersion, whereas positive values

signal phylogenetic clustering.

Figure 3. Distance-decay in cyanobacterial communities. A) Natural log-transformed

phylogenetic (Unifrac) similarity between all pairwise cyanobacterial communities at each

decade between the 1900s to the 2010s plotted against a gradient of natural log-transformed

geographic distances between lakes. A null distance signifies that the pairwise phylogenetic

dissimilarity was calculated between samples of the same lake at a given period. The vertical

dashed lines mark the distances of 50 km and 130 km for reference (the absolute pairwise

distances separating all lakes are summarized in Table S2 of Supplementary Information). **B)**

Natural log-transformed Unifrac distances plotted against the natural log-transformed

temporal gradient (years) for each lake (colour coding).

Figure 4. Cyanobacterial beta-diversity over environmental gradients. Phylogenetic

(Unifrac) similarity between samples are plotted against environmental distances (euclidean)

in **A)** TP (n = 104) and **B)** NO_3^--N (n = 77) concentrations, **C)** air temperature (n = 209), and

D) maximal annual Schmidt Stability Index (SSI) (n = 65). Only significant regressions ($p \leq$

0.05) are shown. The number of samples in each analysis depends on the availability of lake

monitoring data (details on the time-series of monitoring data are found in [16]). Samples used

in the regression analysis are the non-random communities (i.e., showing significant

phylogenetic structure based on SES_{MPD} deviation from the null model expectation).