Analysis of the gut and gill microbiome of resistant and susceptible lines of

rainbow trout (*Oncorhynchus mykiss*)

Ryan M. Brown[1], Gregory D. Wiens[2], Irene Salinas[1]

[1] University of New Mexico, Department of Biology, Center for Evolutionary

and Theoretical Immunology (CETI), Albuquerque, New Mexico, USA

[2] National Center for Cool and Cold Water Aquaculture, Agriculture Research

Service, United States Department of Agriculture Kearneysville, West Virginia,

USA

<u>Abstract</u>

Commensal microorganisms present at mucosal surfaces play a vital role in

protecting the host organism from bacterial infection. There are multiple factors

that contribute to selecting for the microbiome, key of which are host genetics.

Flavobacterium psychrophillum, the causative agent of Bacterial Cold Water Disease

in salmonids, accounts for acute losses in wild and farmed Rainbow Trout

(Oncorhynchus mykiss). The U.S. National Center for Cool and Cold Water

Aquaculture has used family-based selective breeding to generate a line of rainbow

trout with enhanced resistance to *F. psychrophilum.* The goal of this study is to

determine whether selective breeding impacts the gut and gill microbiome of the *F.*

psychrophilum-resistant as compared to a background matched susceptible trout

line. Mid-gut and gill samples were collected from juvenile fish (mean bwt 118g)

and microbial diversity assessed by 16S rDNA amplicon sequencing. Results indicate

that alpha diversity was significantly higher in the mid-gut of the susceptible line

compared to the resistant line, while no significant differences in alpha diversity

were observed in the gills. *Mycoplasma sp.* was the dominant taxon in the mid-gut of

both groups, although it was present at a decreased abundance in the susceptible

line. We also observed an increased abundance of taxa that could potentially be

pathogenic in the susceptible line, including *Brevinema sp.* and Enterobacteriaceae

members. Within the gills, both lines exhibited similar microbial profiles, with

Candidatus Branchiomonas being the dominant taxon. Together, these results

suggest that selectively bred *Flavobacterium psychrophillum*-resistant trout may

harness a more resilient gut microbiome, attributing to the disease resistant

phenotype, providing a framework for future experiments.

Keywords:

Rainbow trout; Aquaculture; *Flavobacterium psychrophilum*; Bacterial Cold Water

Disease; ARS-Fp-R; ARS-Fp-S; Microbiome; Disease resistance

1. Introduction

The microbiome has well established roles in pathogen exclusion and host

immunity, including systemic and mucosal innate and adaptive immune responses

and development of the immune system [1–3]. There is strong evidence to support a

role of host genetics in the selection of the gut microbiome in humans and other

mammals [4–6], although this has not been well characterized in fish. The host

microbial composition is also shaped by other factors including environment, diet,

and disease [7–10]. To begin to disentangle the contribution of host genetics and

environmental factors shaping the fish microbiome, here we utilize a rainbow trout

model in which two genetic lines of rainbow trout have been established by

selective breeding that differ in susceptibility to a common environmental gram

negative pathogen, *Flavobacterium psychrophilum*.

Flavobacterium psychrophilum is the causative agent of Bacterial Cold Water Disease

(BCWD), which is a major concern in the United States aquaculture industry

affecting a range of cold-water fish species, including the commercially relevant

rainbow trout *(Oncorhynchus mykiss)*. *F. psychrophillum* is a mucosal pathogen that

typically infects the skin and gills of fish [11]. Symptoms of BCWD in developed fish

include necrosis of the caudal region, skin lesions, eroded fin tips, and loss of

appetite. *F. psychrophilum* has a more pronounced effect on young fry, a condition

referred to as rainbow trout fry syndrome. Rainbow trout fry syndrome is

responsible for acute losses in trout farms worldwide, as the associated mortality

rate is reported to be greater than 50% [12]. BCWD is becoming an increasingly

difficult disease to treat, as *F. psychrophilum* strains have developed resistance to

several commonly used antibiotics [13–15], and there is currently no commercially

available licensed vaccine.

The National Center for Cool and Cold Water Aquaculture (NCCWA) utilized

family-based selective breeding to develop two distinctive genetic lines of rainbow

trout that confer enhanced resistance (ARS-Fp-R), or susceptibility (ARS-Fp-S) to

the pathogen *F. psychrophilum* [16]. Enhanced resistance to *F. psychrophilum*-

induced mortality in the ARS-Fp-R line has been described, both in the laboratory

setting and on trout farms [17,18]. Previous studies have begun to investigate

possible host mechanisms that attribute to enhanced resistance. For instance, a

strong correlation between resistance to *F. psychrophilum* and increased spleen size

has been described, although this relation does not appear to translate to other

common fish pathogens, such as *Yersinia ruckeri* [19]. Additionally, whole-body

transcriptome analysis has identified numerous acute phase proteins and

inflammatory cytokines that are differentially expressed in each line following

challenge with *F. psychrophilum* [20]. Further work is needed to better characterize

the mechanism(s) by which enhanced resistance is achieved in the ARS-Fp-R line.

In this paper, we begin to investigate the hypothesis that the higher innate disease

resistance of the ARS-Fp-R line is due to differences in the microbial composition at

host mucosal surfaces. Using 16S rDNA amplicon sequencing, we determine and

compare the mid-gut and gill microbiomes of rainbow trout lines selectively bred

for *F. psychrophilum* resistance and susceptibility.

2. Materials and Methods

2.1. Animals and sampling

The Institutional Animal Care and Use Committee (Leetown, WV) reviewed and

approved all animal husbandry practices and disease challenge protocols per

standards set forth in the USDA, ARS Policies and Procedures 130.4.v.3 titled

'Institutional Animal Care and Use 84 Committee'. Fish used in these experiments

were from the 2017 Year Class and maintained as specific pathogen free as

determined by biannual testing as previously described [17]. A total of 33 single

sire-dam matings contributed to the pool of ARS-Fp-R line fish and 31 matings

contributed to the pool of ARS-Fp-S line. The disease resistant phenotype of each

genetic line was evaluated at two time-points, 75 and 276 days post-hatch. Fish

were challenged with *F. psychrophilum* strain CSF259-93 and survival recorded over

21 days as previously described [19]. Mean fish body weight at the first evaluation

was 1.9 g and a total of 120 ARS-Fp-R and 119 ARS-Fp-S fish (n=3 tanks per line)

were challenged by intraperitoneal injection with a dose of 1.4E+07 CFU g^{-1} in a

total volume of 50 μL using a 26g needle fitted onto an Eppendorf repeating pipette.

Mean body weight at the second evaluation was 194 g and a total of 70 ARS-Fp-R

and 70 ARS-Fp-S fish (n=2 tanks per line) were challenged by intramuscular

injection with a dose of 3.5E+06 CFU g^{-1} body weight in a total volume of 50 μL using

a 26g needle fitted onto an Eppendorf repeating pipette. The fish utilized in the

second experiment were part of a larger study evaluating experimental vaccination

and these fish had been sham vaccinated with PBS 35 days prior to challenge. In

both challenges, *F. psychrophilum* was isolated from mortalities and confirmed by

PCR genotyping.

At the time of microbiome sampling, fish were reared under two different tank

conditions as described in Supplemental data file 1. Ten fish from each tank system

per genetic line were euthanized using 200 mg L^{-1} MS222 for 5 minutes. Each fish

was photographed, weighted, gill tissue sampled, intestinal mid-gut sampled, and

spleen weighed within 30 minutes of euthanasia. Sample were placed in SLB [21]

on ice and then moved to storage at -80° C. Instruments were cleaned between each

fish and gloves changed between tank groups. Three control tubes containing SLB

alone were included as negative controls.

2.2. DNA Extraction, 16S rDNA PCR Amplification, and Sequencing

Whole genomic DNA was extracted from skin and gill samples by first lysing tissue

samples using sterile 3 mm tungsten beads (Qiagen) and Qiagen TissueLyser II.

Next, using the cetyltrimethylammonium bromide method as previously described

[21], DNA was isolated and suspended in 50 μL RNase and DNase free molecular

biology grade water. DNA concentration and purity was then assessed using a

Nanodrop ND 1000 (Thermo Scientific).

Bacterial DNA was then replicated by PCR using Illumina adapter fused primers

targeting the V1-V3 region of the prokaryotic 16S rDNA gene. The primer sequences

were as follows: 28F 5'-GAGTTTGATCNTGGCTCAG-3' and 519R

5'GTNTTACNGCGGCKGCTG-3' (where N = any nucleotide, and K = T or G). DNA

samples were diluted 1:10 or 1:100, and Quantabio 5PRIME HotMasterMix was

used. 16S amplicons were generated using the following conditions: 94° C for 90s;

33 cycles of 94° C for 30s, 52° C for 30s, 72° C for 90s; and a final extension of 72° C

for 7 min A positive control of a verified 16S V1-V3 amplicon, and a negative control

of molecular biology grade water was included in every PCR reaction. In addition,

we included a mock community positive control with each sequencing run, which

consisted of equal DNA amounts of 7 different bacterial isolates previously cloned

and a negative control that consisted of SLB handled in the same way as the rest of

the tubes during sampling to which no tissue was added. Amplicons were purified

using the Axygen AxyPrep Mag PCR Clean-up Kit (Thermo Scientific), and eluted into

30 μL molecular biology grade water. Unique oligonucleotide barcodes were ligated

to the 5' and 3' ends of each sample, as well as the Nextera adaptor sequences, using

the Nextera XT Index Kit v2 set A (Illumina). DNA concentrations were quantified

using a Qubit, and normalized to a concentration of 200 ng/uL for DNA library

pooling. Pooled samples were cleaned once more using the Axygen PCR clean-up kit

before being sent off for sequencing. Sequencing was performed on the Illumina

Miseq platform using the MiSeq Reagent Kit v3 (600 cycle) at the Clinical and

Translational Sciences Center at the University of New Mexico Health Sciences

Center.

2.3. Data Analysis and statistics

Differences in survival between genetic lines were determined using the product

limit method of Kaplan and Meier and calculations were performed using GraphPad

v4.0 software. Log-rank (Mantel-Cox) test was used to compare survival curves.

Initial sequence data was analyzed using the latest version of Quantitative Insights

Into Microbial Ecology 2 (Qiime2 v2018.6) [22]. Demultiplexed sequence reads

were quality filtered using DADA2 [23]. Samples were then rarefied to a sampling

depth of 12,603 sequences per sample for mid-gut reads, and 2020 sequences per

sample for gill reads. Amplicon sequence variants (ASVs) were picked by aligning to

the latest Silva 16S database (version 132). Core diversity metrics were analyzed,

including number of ASVs and Shannon's diversity index for alpha diversity, and

PERMANOVA for beta diversity. Nonmetric multidimensional scaling and

generation of heat maps were performed in RStudio [24] using the phyloseq

package [25]. Random forest modeling was performed in Qiime2. Differential

abundance testing was performed in Qiiime2 using ANCOM [26], as well as in

RStudio using the DESeq2 package [27]. For all statistical analyses, fish were split

into groups based on 'Treatment' (ARS-Fp-R, ARS-Fp-S) or 'Tank Treatment'.

3. Results

3.1. Phenotype Confirmation of Disease Resistance/Susceptibility

The relative phenotype of the two genetic lines was evaluated at time-points either

before or after microbiome sampling. At both time points, the survival of the ARS-

Fp-R genetic line was significantly higher ($P<0.001$) than the ARS-Fp-S line and

consistent with estimated mid-point breeding values. In the first evaluation, a total

of 3/120 (3%) ARS-Fp-R line fish died compared to 82/119 (69%) ARS-Fp-S line

fish. In the second evaluation, 2/70 (3%) ARS-Fp-R fish died, while 58/70 (83%)

ARS-Fp-S line fish died within the 21-day challenge period.

3.2. High throughput sequencing analysis

A total number of 3,598,038 raw reads were obtained from all mid-gut samples.

After merging paired ends, quality filtering, and removal of chimeric reads with

DADA2, as well as filtering out non-specific trout genomic reads, a total of 1,413,104

reads remained, with a mean of 35,328 reads per sample. Samples were rarified to a

sample depth of 12,031, which excluded two ARS-Fp-R samples and two ARS-Fp-S

samples. The sample size after rarefaction was n=18 for both the ARS-Fp-R and ARS-

Fp-S lines.

Gill sample sequencing produced a total of 4,646,971 raw reads. Quality filtering in

DADA2 and removal of non-specific reads retained 333,281 reads, with a mean of

13,331 reads per sample. Samples were rarified to 2010 reads. The sample size after

rarefaction was n=11 for ARS-Fp-R and n=14 for ARS-Fp-S.

3.3. Resistant and susceptible lines display significant differences in alpha

diversity in the mid-gut, but not the gills

Comparison of alpha diversity metrics obtained from the mid-gut showed

significantly lower measures of gut microbial community richness (Observed ASVs),

as well as Shannon's diversity index in the ARS-Fp-R line compared to the ARS-Fp-S

line (Fig 1A and 1B). There were a total of 15 ASVs reported in the mid-gut of the

resistant line, and 29 in the mid-gut of the susceptible line. In the gills, there were no

significant differences in alpha diversity between lines with a total of 57 ASVs found

in the gills of the resistant line, and 50 in the susceptible line (Fig 1C and 1D).

Fig 1. Comparison of alpha diversity metrics for the mid-gut and gill microbiome of ARS-Fp-R

and ARS-Fp-S trout.

(A) Total number of observed ASVs in the mid-gut. (B) Shannon's diversity index in the mid-gut. (C)

Total number of observed ASVs in the gills. (D) Shannon's diversity index in the gills. ** indicates

statistically significant differences p<0.01.

3.4. Beta diversity analysis suggests possible tank effect on the gut

microbiome

We assessed the microbial diversity between different treatments, as well as

between tanks by performing Nonmetric Multidimensional Scaling (NMDS) using

the Bray Curtis distance metric. This ordination showed a discrete grouping of the

two tanks containing the ARS-Fp-S line, while the two tanks containing the ARS-Fp-

R line were more tightly clustered (Fig 2A). PERMANOVA analysis [28] identified

"Treatment" (P value = 0.02) and "Tank" (P value = 0.048 for ARS-Fp-R, P value =

0.001 for ARS-Fp-S) as significant determinants of the mid-gut microbial community

composition. In the gills, NMDS ordination showed a similar pattern to that found in

the gut, where fish from tank 26 of the susceptible line clustered tightly together

while fish from tank 12 showed greater variability. Meanwhile, there was no clear

separation between individuals held in separate tanks of the resistant line (Fig 2B).

PERMANOVA identified only tank housing within the ARS-Fp-S line (P value =

0.004) as a significant factor in determining gill microbial communities.

Fig 2. NMDS ordination plots.

NMDS ordination performed using Bray Curtis distance matrix of the (A) mid-gut and (B) gill

microbiome of ARS-Fp-R and ARS-Fp-S trout.

3.5. Gut microbial community composition

A total of eight different phyla were identified in the mid-gut across both lines,

although only four of these were represented over 1%, including Tenericutes,

Spirochaetes, Proteobacteria, and Firmicutes. Tenericutes composed the vast

majority of the mid-gut microbiome of both lines, constituting 81% of the total

microbial diversity in the susceptible line and 89% in the resistant line (Fig 3A).

At the genus level, all Tenericutes reads were identified as *Mycoplasma sp.* The

genus *Brevinema sp.* showed greater abundance in the ARS-Fp-S (8.5%) compared

to the ARS-Fp-R line (4.8%). Similarly, members of the family Enterobacteriaceae

had greater abundance in the ARS-Fp-S line than the ARS-Fp-R line (2.2% and 0%,

respectively); however, these differences were not significant. Differential

abundance testing with ANCOM revealed three genera that were differentially

abundant in the mid-gut of both trout lines; including *Hydrotalea sp., Paenibacillus*

sp., and *Variovorax sp.* This was replicated by differential abundance testing using

DESeq2, an R package originally developed for differential expression analysis in

RNA-seq data, but has been proposed to be used in microbiome studies as well

[29,30].

The relative distribution of ASVs at the genus level was notably different between

tanks of the same line (Fig 3B). Potential opportunistic taxa such as *Brevinema sp.*

and Ambiguous Enterobacteriaceae were present in tanks 11 and 12, but were not

identified in either tank 25 or 26. This trend is further shown in a heatmap

representing the top 25 ASVs observed in each sample (Fig 4). Two of the tanks that

were in close proximity to one another displayed similar microbial profiles, despite

holding different lines. This was not the case for tanks 25 and 26, as tank 26

displayed a microbial profile different from all other tanks since it contained ASVs

representative of Enterobacterieaceae, *Hydrotalea sp.*, *Paenibacillus sp,* and

Variovorax sp.

Fig 3. Relative microbial composition in the gut of each line.

(A) Relative abundance of ASVs at the phylum level for each line. (B) Relative abundance of ASVs for

each tank the two lines were divided into.

Fig 4. Heatmap representing the top 25 ASVs present in the mid-gut of ARS-Fp-R and ARS-Fp-S

trout.

Each column represents one individual. Each row represents one ASV.

3.6. Gill microbial community composition

A total of nine different phyla were present in the trout gills across both lines. Five

of these were represented at abundance greater than 1%, including Proteobacteria,

Tenericutes, Spirochaetes, Bacteriodetes, and Firmicutes. Proteobacteria was the

most abundant phylum in both groups, representing 85% of all bacterial diversity in

the ARS-Fp-S line and 95% in the ARS-Fp-R line (Fig 5A). At the genus level, most

Proteobacteria reads were identified as *Candidatus Branchiomonas*. This taxon

constituted 74% of all diversity in the gills of the susceptible line and 85% in the

resistant line. We identified trace amounts of *Flavobacterium sp.* in both lines, as this

taxon constituted 0.16 % of the ARS-Fp-S line and 1.1% of the ARS-Fp-R line. No

ASVs were differentially abundant in the gill microbial community of both groups

through ANCOM or DEseq2 analyses.

We observed discernable differences in the microbial community composition of

fish of the same line housed in different tanks (Fig 5B). For example, *Candidatus*

Branchiomonas was present at levels above 95% in tanks 25 (ARS-Fp-R) and 26

(ARS-Fp-S), whereas it constituted 71% of tank 11 (ARS-Fp-R) and 20% of tank 12

(ARS-Fp-S). The aforementioned potential opportunistic pathogens *Brevinema sp.*

was only identified in tank 12. A heatmap of the top 30 ASVs in each sample (Fig 6)

shows signatures of *Brevinema sp.* in fish from tank 12, as well as a reduced

abundance of *Candidatus Branchiomonas* in this tank compared to all others.

Fig 5. Microbial community composition of the gills of ARS-Fp-R and ARS-Fp-S trout.

(A) Relative abundance of ASVs at the phylum level for each line. (B) Relative abundance of ASVs for

tank the two lines were divided into.

Fig 6. Heatmap representing the top 30 ASVs represented in the gill of ARS-Fp-R and ARS-Fp-S

trout.

Each column represents one individual fish. Each row represents one ASV.

3.7. Random Forest Modeling

Random forest modeling [31] was performed in order to see if a machine learning

module could accurately predict treatment group as well as spleen index based on

microbial community composition. In the mid-gut, this model was able to accurately

predict 100% of FPR samples, but only 50% of FPS samples (Fig 7). Table 1 shows

features at the genus level that were rendered as being important in classifying

treatment group. Due to the lower sampling size in the gills, we did not include

random forest analyses of the gill samples. Finally, we also performed a random

forest regressor to attempt to predict spleen index based on gut microbial

composition, and found that there was no correlation between these two variables.

Table 1: Features at the genus level that were identified by random forest modeling as being

important in the classification of treatment group

Feature	Importance
Ralstonia	0.194781898
Mycoplasma	0.190003235
Variovorax	0.129277426
Brevinema	0.098010406
Paenibacillus	0.093643145
Hydrotalea	0.084976654
Ambiguous Desulfovibrionaceae	0.08063865
Streprococcus	0.034762947
Staphylococcus	0.027645825
Ambiguous Gammaproteobacteria	0.024251868
Shewanella	0.023781255
Ambiguous Ruminococcaceae	0.014827708
Exiguobacterium	0.002446602
Ambiguous Moraxellaceae	0.000952381

Fig 7. Confusion matrix for random forest modeling in the gut.

This heatmap represents how frequently samples within each line were correctly classified. The

correct label is represented on the y-axis and the predicted label is represented on the x-axis, with

the prediction frequency shown as a gradient.

4. Discussion

Commensal microbes have co-evolved with their eukaryotic counterparts, forming

an intricate relationship that benefits both parties involved. Several studies have

revealed that host genetics influences gut microbiota composition in a variety of

species, including humans and rodents [4,5], chickens [32], and *Drosophila* [33].

However, other factors such as host diet and environmental factors are also deeply

intertwined and deeply shape host microbial communities [7–9].

Teleost fish live in symbiosis with complex microbial communities that inhabit

every mucosal barrier (gut, gills, skin and nose) [34]. Fish microbial community

composition is influenced by age [35], tissue site [36], diet [37–39], stress [40] and

pathogen infection [41]. However, few studies have investigated the impact host

genetics have on shaping teleost microbiomes. A study on brook charr identified

three quantitative trait loci associated with abundance of commensal strains in the

skin [42]. Another study in Atlantic salmon found significant differences in the skin

and gut microbial composition amongst distinct wild populations that are not likely

attributed to environmental conditions alone [43]. These observations suggest that

host genetics play an important role in teleost microbiome assembly, although more

work is needed to better understand this relation.

Farmed fish are susceptible to many pathogens that threaten the sustainability of

the finfish farming industry. Among the most prominent bacterial diseases, BCWD is

particularly problematic in rainbow trout. The development of two genetic trout

lines with different susceptibilities to BCWD agent, *F. psychrophilum*, offers an

excellent platform to understand how host genetics may shape fish microbial

communities. The current study suggests that host genetics influence the microbial

community composition in the mid-gut, since the mid-gut bacterial community of

the susceptible line was significantly more diverse than that of the resistant line. In

agreement with previous studies, the mid-gut communities of both lines were

dominated by *Mycoplasma sp.* This taxon appears to be highly abundant in the

gastrointestinal microbiome of all salmonid species studied, including Atlantic

salmon [44–46], rainbow trout [36,47] and Chinook salmon [48]. Interestingly, we

identified a lower abundance of *Mycoplasma sp.* in the susceptible line, suggesting

that disease susceptibility may be associated with decreased *Mycoplasma sp.* levels

in the gut. In support, a recent study investigated the intestinal microbiome of

offshore farmed Chinook salmon [48], finding that abundance of potential

pathogenic *Vibrio sp.* appeared to be inversely correlated with the presence of

Mycoplasma sp. Mycoplasma sp. are characterized by their uniquely small genome

and lack of a cell wall, which makes culture-based approaches to studying this

bacterium difficult to achieve. Considering the widespread presence and abundance

of *Mycoplasma sp.* in salmonid gastrointestinal tract across a wide range of

geographical locations, including both lab-reared fish and wild-caught, it appears as

if there have been strong evolutionary forces that have enabled *Mycoplasma sp.* to

thrive in this microenvironment. Future studies are needed to determine the nature

of this relationship and the ability of *Mycoplasma sp.* to prevent pathogen

colonization in the gastrointestinal tract of salmonids.

Potential opportunistic pathogens were observed within the mid-gut of the ARS-Fp-

S line at a greater abundance than in the ARS-Fp-R line. Expansions of opportunistic

taxa have been previously described in other fish microbiome studies including in

the skin of Atlantic salmon experimentally infected with salmonid alphavirus (Reid,

2017) and in Atlantic salmon experimentally infected with the parasite

Lepeophtheirus salmonis [41]. Specifically, we found *Brevinema sp.* and ambiguous

members of the family Enterobacteriaceae in the susceptible line. *Brevinema sp.* was

further identified as *Brevinema andersonii* at the species level in the Silva database,

which is a pathogenic taxa that has been isolated in rodents [49]. However, upon

sequence search using BLAST, they were most closely identified as an uncultured

Spirochaeta sequence derived from an aquatic environment. Spirochaetes members

have been described to cause disease in other marine species [50], and there are

multiple well-studied members that are human pathogens [51]. Deeper taxonomic

characterization would be necessary to determine if this ASV is in fact a pathogen in

rainbow trout. Meanwhile, Enterobacteriaceae reads were most closely identified as

Hafnia paralvei, which is associated with infections in a variety of animals, including

fish [52]. Although the relative abundance of these taxa was small, these results

suggest that ARS-Fp-R trout may possess a more resilient mid-gut microbiome, as

compared to ARS-Fp-S. It is important to note that these fish were reared on spring

water and sampled only at one timepoint. Additional studies are needed to examine

whether these trends are observed in other environments and stages of

development.

Differences in the microbial community composition between both trout lines were

less pronounced in the gills compared to the gut, although it should be noted that

the sample size was smaller in this tissue due to difficulties amplifying the 16S rDNA

region in certain samples. Nonetheless, alpha diversity metrics were similar in each

line. Interestingly, *Candidatus Branchiomonas* was the dominant taxon in both lines.

This is a known pathogen that has been shown to cause gill epitheliocystis in

Atlantic salmon [53], although it has not been previously described in rainbow

trout. Fish from both lines were visually healthy, suggesting that *Candidatus*

Branchiomonas may be a common member of the trout gill microbiome in certain

environments. Further studies should evaluate which factors favor the colonization

of *Candidatus Branchiomonas* in trout gills.

Flavobacterium sp. was identified in the gills of both lines, although sequence search

using BLAST did not yield species level taxonomic resolution. Although this taxon

was present at low abundance in both lines, it was surprising that we detected

higher *Flavobacterium sp.* abundance in the gill microbiome of the resistant line

compared to the susceptible line. Considering that the resistant and susceptible

lines maintained the survival phenotype following the challenge experiment, our

results indicate that susceptibility to *F. psychrophilum* infection is not due to

increased abundance of this pathogen as a member of the indigenous microbial

community. Disease susceptibility may be instead due to the greater ability of this

pathogen to displace the microbial communities in the susceptible line compared to

the resistant line.

One of the interesting aspects of the present study was the identification of

differences in the microbial composition between tanks of the same trout genetic

lines. All tanks were supplied water from the same source, and proper water quality

was maintained throughout the experiment. This tank effect complicates the ability

to discern between genetic lines, although there still appears to be notable

differences when comparing each line, particularly in the gut as discussed earlier.

There were similarities between two of the neighboring tanks that housed different

lines, in tanks 11 and 12. However, this did not occur in the other two neighboring

tanks, as the susceptible trout from tank 26 displayed an expansion of pathogenic

taxa compared to tank 25, which housed resistant trout.

Work in zebrafish has shown that interhost dispersal can actually outweigh genetic

factors that contribute to microbiome assembly [54]. In order to assess whether

host genetics contribute to host microbiome assembly, all external factors that

potentially contribute to this network must also be accounted for. In the present

study, diet was consistent across fish of all tanks, and environmental factors such as

water quality and temperature were also consistently maintained. Still, differences

amongst tanks were present that add noise to the between-group comparisons. This

finding highlights the importance of adequate experimental design in fish

microbiome studies as well as the fact that different factors differentially shape

microbial assembly at different tissue sites (i.e. gut versus gills).

5. Conclusions

In conclusion, the present study reveals differences in the microbial composition of

the gut but not the gills of two rainbow trout lines with differential susceptibility to

F. psychrophilum infection. Disease susceptibility was associated with a more

diverse gut microbiome and the presence of potentially pathogenic taxa although

important tank effects were also detected. Thus, selective breeding programs may

not only select for host genetic factors but also, as a consequence, unique microbial

assemblies, which in turn, may render the host more or less resilient to pathogen

invasion or infection.

Acknowledgements

We thank Kurt Schwalm and Dr. Darrell Dinwiddie for handling the Illumina MiSeq

runs. We thank Dr. Timothy Leeds for breeding the ARS-Fp-R and ARS-Fp-S genetic

lines and Travis Moreland for fish rearing and Jeremy Everson for assistance with

fish sampling. This work was supported in part by Agricultural Research Service

Project 1930-32000-006. Mention of trade names or commercial products in this

publication is solely for the purpose of providing specific information and does not

imply recommendation or endorsement by the U.S Department of Agriculture. USDA

is an equal opportunity employer.

Data Availability

All datasets obtained have been deposited at NCBI BioProject and are publicly

available under BioProject ID number PRJNA488363.

References

[1]   C.A. Thaiss, N. Zmora, M. Levy, E. Elinav, The microbiome and innate immunity, Nature. 535 (2016) 65–74. doi:10.1038/nature18847.

[2]   N. Shi, N. Li, X. Duan, H. Niu, Interaction between the gut microbiome and mucosal immune system, Mil. Med. Res. 4 (2017). doi:10.1186/s40779-017-0122-9.

[3]   C.G. Buffie, E.G. Pamer, Microbiota-mediated colonization resistance against intestinal pathogens, Nat. Rev. Immunol. 13 (2013) 790–801. doi:10.1038/nri3535.

[4]   M.J. Bonder, A. Kurilshikov, E.F. Tigchelaar, Z. Mujagic, F. Imhann, A.V. Vila, P. Deelen, T. Vatanen, M. Schirmer, S.P. Smeekens, D. V. Zhernakova, S.A. Jankipersadsing, M. Jaeger, M. Oosting, M.C. Cenit, A.A.M. Masclee, M.A. Swertz, Y. Li, V. Kumar, L. Joosten, H. Harmsen, R.K. Weersma, L. Franke, M.H. Hofker, R.J. Xavier, D. Jonkers, M.G. Netea, C. Wijmenga, J. Fu, A. Zhernakova, The effect of host genetics on the gut microbiome, Nat. Genet. 48 (2016) 1407–1412.

doi:10.1038/ng.3663.

[5]   A. Spor, O. Koren, R. Ley, Unravelling the effects of the environment and host

genotype on the gut microbiome, Nat. Rev. Microbiol. 9 (2011) 279–290.

doi:10.1038/nrmicro2540.

[6]   M. Fulde, F. Sommer, B. Chassaing, K. van Vorst, A. Dupont, M. Hensel, M. Basic,

R. Klopfleisch, P. Rosenstiel, A. Bleich, F. Bäckhed, A.T. Gewirtz, M.W. Hornef,

Neonatal selection by Toll-like receptor 5 influences long-term gut microbiota

composition, Nature. 560 (2018) 489–493. doi:10.1038/s41586-018-0395-5.

[7]   L.A. David, C.F. Maurice, R.N. Carmody, D.B. Gootenberg, J.E. Button, B.E. Wolfe,

A. V. Ling, A.S. Devlin, Y. Varma, M.A. Fischbach, S.B. Biddinger, R.J. Dutton, P.J.

Turnbaugh, Diet rapidly and reproducibly alters the human gut microbiome,

Nature. 505 (2014) 559–563. doi:10.1038/nature12820.

[8]   D. Rothschild, O. Weissbrod, E. Barkan, A. Kurilshikov, T. Korem, D. Zeevi, P.I.

Costea, A. Godneva, I.N. Kalka, N. Bar, S. Shilo, D. Lador, A.V. Vila, N. Zmora, M.

Pevsner-Fischer, D. Israeli, N. Kosower, G. Malka, B.C. Wolf, T. Avnit-Sagi, M.

Lotan-Pompan, A. Weinberger, Z. Halpern, S. Carmi, J. Fu, C. Wijmenga, A.

Zhernakova, E. Elinav, E. Segal, Environment dominates over host genetics in

shaping human gut microbiota, Nature. 555 (2018) 210–215.

doi:10.1038/nature25973.

[9]   R.K. Singh, H.-W. Chang, D. Yan, K.M. Lee, D. Ucmak, K. Wong, M. Abrouk, B.

Farahnik, M. Nakamura, T.H. Zhu, T. Bhutani, W. Liao, Influence of diet on the

gut microbiome and implications for human health, J. Transl. Med. 15 (2017)

73. doi:10.1186/s12967-017-1175-y.

[10]   and V.B.Y. Schreiner, Andrew B., John Y. Kao, The gut microbiome in health

493   and in disease, Curr. Opin. Gastroenterol. 31 (2015) 69–75.

494   doi:10.1097/MOG.0000000000000139.The.

495   [11]   A. Nematollahi, A. Decostere, F. Pasmans, F. Haesebrouck, Flavobacterium

496   psychrophilum infections in salmonid fish, J. Fish Dis. 26 (2003) 563–574.

497   doi:10.1046/j.1365-2761.2003.00488.x.

498   [12]   C.E. Starliper, Bacterial coldwater disease of fishes caused by Flavobacterium

499   psychrophilum, J. Adv. Res. 2 (2011) 97–108. doi:10.1016/j.jare.2010.04.001.

500   [13]   M.S. Bruun, A.S. Schmidt, L. Madsen, I. Dalsgaard, Antimicrobial resistance

501   patterns in Danish isolates of Flavobacterium psychrophilum, Aquaculture.

502   187 (2000) 201–212. doi:10.1016/S0044-8486(00)00310-0.

503   [14]   M.S. Bruun, L. Madsen, I. Dalsgaard, Efficiency of oxytetracycline treatment in

504   rainbow trout experimentally infected with Flavobacterium psychrophilum

505   strains having different in vitro antibiotic susceptibilities, Aquaculture. 215

506   (2003) 11–20. doi:10.1016/S0044-8486(01)00897-3.

507   [15]   C. Kum, S. Kirkan, S. Sekkin, F. Akar, M. Boyacioglu, Comparison of in vitro

508   antimicrobial susceptibility in Flavobacterium psychrophilum isolated from

509   rainbow trout fry, J. Aquat. Anim. Health. 20 (2008) 245–251.

510   doi:10.1577/H07-040.1.

511   [16]   J.T. Silverstein, R.L. Vallejo, Y. Palti, T.D. Leeds, C.E. Rexroad, T.J. Welch, G.D.

512   Wiens, V. Ducrocq, Rainbow trout resistance to bacterial cold-water disease is

513   moderately heritable and is not adversely correlated with growth, J. Anim. Sci.

514   87 (2009) 860–867. doi:10.2527/jas.2008-1157.

515   [17]   G.D. Wiens, S.E. LaPatra, T.J. Welch, J.P. Evenhuis, C.E. Rexroad, T.D. Leeds, On-

516   farm performance of rainbow trout (Oncorhynchus mykiss) selectively bred

517   for resistance to bacterial cold water disease: Effect of rearing environment

518   on survival phenotype, Aquaculture. 388–391 (2013) 128–136.

519   doi:10.1016/j.aquaculture.2013.01.018.

520   [18]   G.D. Wiens, Y. Palti, T.D. Leeds, Three generations of selective breeding

521   improved rainbow trout (Oncorhynchus mykiss) disease resistance against

522   natural challenge with Flavobacterium psychrophilum during early life-stage

523   rearing, Aquaculture. 497 (2018) 414–421.

524   [19]   S. Hadidi, G.W. Glenney, T.J. Welch, J.T. Silverstein, G.D. Wiens, Spleen Size

525   Predicts Resistance of Rainbow Trout to Flavobacterium psychrophilum

526   Challenge, J. Immunol. 180 (2008) 4156–4165.

527   doi:10.4049/jimmunol.180.6.4156.

528   [20]   D. Marancik, G. Gao, B. Paneru, H. Ma, A.G. Hernandez, M. Salem, J. Yao, Y. Palti,

529   G.D. Wiens, Whole-body transcriptome of selectively bred, resistant-, control-,

530   and susceptible-line rainbow trout following experimental challenge with

531   Flavobacterium psychrophilum, Front. Genet. 5 (2015).

532   doi:10.3389/fgene.2014.00453.

533   [21]   K.R. Mitchell, C.D. Takacs-Vesbach, A comparison of methods for total

534   community DNA preservation and extraction from various thermal

535   environments, J. Ind. Microbiol. Biotechnol. 35 (2008) 1139–1147.

536   doi:10.1007/s10295-008-0393-y.

[22]   J.G. Caporaso, J. Kuczynski, J. Stombaugh, K. Bittinger, F.D. Bushman, E.K.

Costello, N. Fierer, A.G. P⬜a, J.K. Goodrich, J.I. Gordon, G.A. Huttley, S.T. Kelley,

D. Knights, J.E. Koenig, R.E. Ley, C.A. Lozupone, D. McDonald, B.D. Muegge, M.

Pirrung, J. Reeder, J.R. Sevinsky, P.J. Turnbaugh, W.A. Walters, J. Widmann, T.

Yatsunenko, J. Zaneveld, R. Knight, QIIME allows analysis of high-throughput

community sequencing data, Nat. Methods. 7 (2010) 335–336.

doi:10.1038/nmeth.f.303.

[23] B.J. Callahan, P.J. McMurdie, M.J. Rosen, A.W. Han, A.J.A. Johnson, S.P. Holmes,

DADA2: High-resolution sample inference from Illumina amplicon data, Nat.

Methods. 13 (2016) 581–583. doi:10.1038/nmeth.3869.

[24] R.C. Team, R: A language and environment for statistical computing, (2018).

https://www.r-project.org/.

[25] P.J. McMurdie, S. Holmes, Phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data, PLoS One. 8 (2013).

doi:10.1371/journal.pone.0061217.

[26] S. Mandal, W. Van Treuren, R.A. White, M. Eggesbø, R. Knight, S.D. Peddada,

Analysis of composition of microbiomes: a novel method for studying

microbial composition., Microb. Ecol. Health Dis. 26 (2015) 27663.

doi:10.3402/mehd.v26.27663.

[27] M.I. Love, W. Huber, S. Anders, Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2, Genome Biol. 15 (2014).

doi:10.1186/s13059-014-0550-8.

[28] M.J. Anderson, PERMANOVA Permutational multivariate analysis of variance,

Austral Ecol. (2005) 1–24. doi:10.1139/cjfas-58-3-626.

[29] P.J. McMurdie, S. Holmes, Waste Not, Want Not: Why Rarefying Microbiome

Data Is Inadmissible, PLoS Comput. Biol. 10 (2014).

doi:10.1371/journal.pcbi.1003531.

[30] S. Weiss, Z.Z. Xu, S. Peddada, A. Amir, K. Bittinger, A. Gonzalez, C. Lozupone,

J.R. Zaneveld, Y. Vázquez-Baeza, A. Birmingham, E.R. Hyde, R. Knight,

Normalization and microbial differential abundance strategies depend upon

data characteristics, Microbiome. 5 (2017). doi:10.1186/s40168-017-0237-y.

[31] a Liaw, M. Wiener, Classification and Regression by randomForest, R News. 2

(2002) 18–22. doi:10.1177/154405910408300516.

[32] L. Zhao, G. Wang, P. Siegel, C. He, H. Wang, W. Zhao, Z. Zhai, F. Tian, J. Zhao, H.

Zhang, Z. Sun, W. Chen, Y. Zhang, H. Meng, Quantitative genetic background of

the host influences gut microbiomes in chickens, Sci. Rep. 3 (2013).

doi:10.1038/srep01163.

[33] J.M. Chaston, A.J. Dobson, P.D. Newell, A.E. Douglas, Host genetic control of the

microbiota mediates the Drosophila nutritional phenotype, Appl. Environ.

Microbiol. 82 (2016) 671–679. doi:10.1128/AEM.03301-15.

[34] C. Kelly, I. Salinas, Under pressure: Interactions between commensal

microbiota and the teleost immune system, Front. Immunol. 8 (2017).

doi:10.3389/fimmu.2017.00559.

[35] P. Smith, D. Willemsen, M. Popkes, F. Metge, E. Gandiwa, M. Reichard, D.R.

Valenzano, Regulation of life span by the gut microbiota in the short-lived

african turquoise killifish, Elife. 6 (2017). doi:10.7554/eLife.27014.

[36] L. Lowrey, D.C. Woodhams, L. Tacchi, I. Salinas, Topographical mapping of the

rainbow trout (Oncorhynchus mykiss) microbiome reveals a diverse bacterial

community with antifungal properties in the skin, Appl. Environ. Microbiol. 81

(2015) 6915–6925. doi:10.1128/AEM.01826-15.

[37]   H.I. Reid, J.W. Treasurer, B. Adam, T.H. Birkbeck, Analysis of bacterial

populations in the gut of developing cod larvae and identification of Vibrio

logei, Vibrio anguillarum and Vibrio splendidus as pathogens of cod larvae,

Aquaculture. 288 (2009) 36–43. doi:10.1016/j.aquaculture.2008.11.022.

[38]   S. Miyake, D.K. Ngugi, U. Stingl, Diet strongly influences the gut microbiota of

surgeonfishes, Mol. Ecol. 24 (2015) 656–672. doi:10.1111/mec.13050.

[39]   C. Neuman, E. Hatje, K.Z. Zarkasi, R. Smullen, J.P. Bowman, M. Katouli, The

effect of diet and environmental temperature on the faecal microbiota of

farmed Tasmanian Atlantic Salmon (Salmo salar L.), Aquac. Res. 47 (2016)

660–672. doi:10.1111/are.12522.

[40]   L. Tacchi, L. Lowrey, R. Musharrafieh, K. Crossey, E.T. Larragoite, I. Salinas,

Effects of transportation stress and addition of salt to transport water on the

skin mucosal homeostasis of rainbow trout (Oncorhynchus mykiss),

Aquaculture. 435 (2015) 120–127. doi:10.1016/j.aquaculture.2014.09.027.

[41]   M.S. Llewellyn, S. Leadbeater, C. Garcia, F.E. Sylvain, M. Custodio, K.P. Ang, F.

Powell, G.R. Carvalho, S. Creer, J. Elliot, N. Derome, Parasitism perturbs the

mucosal microbiome of Atlantic Salmon, Sci. Rep. 7 (2017).

doi:10.1038/srep43465.

[42]   S. Boutin, C. Sauvage, L. Bernatchez, C. Audet, N. Derome, Inter individual

variations of the fish skin microbiota: Host genetics basis of mutualism?, PLoS

One. 9 (2014). doi:10.1371/journal.pone.0102649.

[43]   T.M. Uren Webster, S. Consuegra, M. Hitchings, C. Garcia de Leaniz, Inter-

population variation in the Atlantic salmon microbiome reflects

environmental and genetic diversity, Appl. Environ. Microbiol. (2018)

AEM.00691-18. doi:10.1128/AEM.00691-18.

[44]   C.E. Dehler, C.J. Secombes, S.A.M. Martin, Seawater transfer alters the

intestinal microbiota profiles of Atlantic salmon (Salmo salar L.), Sci. Rep. 7

(2017). doi:10.1038/s41598-017-13249-8.

[45]   M.S. Llewellyn, P. McGinnity, M. Dionne, J. Letourneau, F. Thonier, G.R.

Carvalho, S. Creer, N. Derome, The biogeography of the atlantic salmon (Salmo

salar) gut microbiome, Isme J. 10 (2015) 1–5. doi:10.1038/ismej.2015.189.

[46]   W.E. Holben, P. Williams, M. Saarinen, L.K. S??rkilahti, J.H.A. Apajalahti,

Phylogenetic analysis of intestinal microflora indicates a novel Mycoplasma

phylotype in farmed and wild salmon, Microb. Ecol. 44 (2002) 175–185.

doi:10.1007/s00248-002-1011-6.

[47]   P.P. Lyons, J.F. Turnbull, K.A. Dawson, M. Crumlish, Phylogenetic and

functional characterization of the distal intestinal microbiome of rainbow

trout Oncorhynchus mykiss from both farm and aquarium settings, J. Appl.

Microbiol. 122 (2017) 347–363. doi:10.1111/jam.13347.

[48]   M. Ciric, D. Waite, J. Draper, J.B. Jones, Characterisation of gut microbiota of

farmed Chinook salmon using metabarcoding, bioRxiv. (2018) 288761.

doi:10.1101/288761.

[49]   D.L. Defosse, R.C. Johnson, B.J. Paster, F.E. Dewhirst, G.J. Fraser, Brevinema

andersonii gen. nov., sp. nov., an infectious spirochete isolated from the short-

tailed shrew (Blarina brevicauda) and the white-footed mouse (Peromyscus

leucopus)., Int. J. Syst. Bacteriol. 45 (1995) 78–84. doi:10.1099/00207713-45-

1-78.

[50] T. Matsuyama, M. Yasuike, A. Fujiwara, Y. Nakamura, T. Takano, T. Takeuchi,

N. Satoh, Y. Adachi, Y. Tsuchihashi, H. Aoki, K. Odawara, S. Iwanaga, J. Kurita, T.

Kamaishi, C. Nakayasu, A Spirochaete is suggested as the causative agent of

Akoya oyster disease by metagenomic analysis, PLoS One. 12 (2017).

doi:10.1371/journal.pone.0182280.

[51] J.J. Halperin, A Tale of Two Spirochetes: Lyme Disease and Syphilis, Neurol.

Clin. 28 (2010) 277–291. doi:10.1016/j.ncl.2009.09.009.

[52] G. Huys, M. Cnockaert, S.L. Abbott, J.M. Janda, P. Vandamme, Hafnia paralvei

sp. nov., formerly known as Hafnia alvei hybridization group 2, Int. J. Syst.

Evol. Microbiol. 60 (2010) 1725–1728. doi:10.1099/ijs.0.018606-0.

[53] E.R. Toenshoff, A. Kvellestad, S.O. Mitchell, T. Steinum, K. Falk, D.J. Colquhoun,

M. Horn, A novel betaproteobacterial agent of gill epitheliocystis in seawater

farmed Atlantic salmon (Salmo salar), PLoS One. 7 (2012).

doi:10.1371/journal.pone.0032696.

[54] A.R. Burns, E. Miller, M. Agarwal, A.S. Rolig, K. Milligan-Myhre, S. Seredick, K.

Guillemin, B.J.M. Bohannan, Interhost dispersal alters microbiome assembly

and can overwhelm host innate immunity in an experimental zebrafish model,

Proc. Natl. Acad. Sci. (2017) 201702511. doi:10.1073/pnas.1702511114.