Tissue-expression profiles unveils the gene interaction of hepatopancreas, eyestalk, and ovary in precocious female Chinese mitten crab, *Eriocheir sinensis*

Xiaowen Chen[1, 2, 3, a], Jun Wang[1, 2, 3, a], Xin Hou[1, 2, 3], Wucheng Yue[1, 2, 3], Shu Huang[1, 2, 3], Chenghui Wang[1, 2, 3, *]

[1] Key Laboratory of Freshwater Aquatic Genetic Resources, Ministry of Agriculture, Shanghai, 201306, China.

[2] National Demonstration Center for Experimental Fisheries Science Education (Shanghai Ocean University), Shanghai, 201306, China.

[3] Shanghai Engineering Research Center of Aquaculture, Shanghai, 201306, China

Running title：Transcriptome of precocious crab.

[a] These authors contributed equally to this study.

*Corresponding author: Chenghui Wang

Address: 999, Hucheng huan Road, Lingang New City, Shanghai, 201306, China

Email address: wangch@shou.edu.cn   Tel / Fax: +86 21 6190 0439

Abstract

Sexual precocity is a serious and common biological phenomenon in animal species. Large amount of precocity individuals was identified in Chinese mitten crab, *Eriocheir sinensis*, which caused huge economical loss every year. However, the underlying genetic basis of precocity in *E. sinensis* is still lack. In this study, histology observation, comparative transcriptome was conducted among different stages of precocious one-year old and normal two-year old *E. sinensis*, tissue-expression profiles of ovary, hepatopancreas, and eyestalk tissues were presented and compared. Genes associated with lipid metabolic process, lipid transport, vitelline membrane formation, vitelline synthesis and neuropeptide hormone related genes were upregulated in the ovary, hepatopancreas and eyestalk of precocious *E. sinensis*. Our results indicated eyestalk involved in neuroendocrine system providing neuropeptide hormone that may induce vitellogenesis in hepatopancreas and further stimulate ovary development. Hepatopancreas is a site for energy storage, vitellogenin synthesis and may assist to induce oogenesis through lipid transport in precocious *E. sinensis*. The genetic basis of precocity in *E. sinensis* is an integrated gene regulatory network of eyestalk, hepatopancreas, and ovary tissues. Our study provides effective convenient phenotype measurement method for identification of potential precocious *E. sinensis* detection, and valuable genetic resources and novel insights into the research of molecular mechanism of precocity in *E. sinensis*.

Keywords: transcriptome; ovary development; genetic network

Introduction

Sexual precocity refers to the reproductive system (gonad) that abnormally develop into sexual mature stage in advance during puberty, which is a natural phenomenon in most animal species, even in human [1-3]. Growth and development retardation, increased illness rate, and other associated physiological defects are believed to be the consequences of sexual precocity [1, 4]. Sexual precocity is a complex physiological process that induced by extrinsic environment factors and intrinsic genetic factors [2, 5]. The early development of gonads is considered to be the molecular response to environment factors, such as hormone, nutrition, temperature, disease and so on [5]. The scientific communities are struggled with the genetic mechanism of sexual precocity; however, the underlying mechanism is still elusive.

In vertebrate, the gonad development is believed to be regulated by hypothalamus-pituitary-gonad axis (HPG), however, in invertebrate, the regulation of reproductive system is vague [6, 7]. Chinese mitten crab *Eriocheir sinensis*, an economical crustacean widely cultured in China is also suffered from severe precocious problems [8]. Huge economic losses in the *E. sinensis* aquaculture industry is caused by substantial proportion of precocious *E. sinensis* individuals every year [2, 9]. Environment factors such as temperature, salinity, light, and stocking density are believed to induce sexual precocity in *E. sinensis*, however, the intrinsic molecular feedback to the stimulation of environment factors is largely unknown in *E. sinensis* [10-12].

A similar X-organ-sinus gland complex neuroendocrine system existed in eyestalk is believed to play analogous roles as HPG axis in crustaceans [7, 13, 14]. Gonad inhibiting hormone (*GIH*), molt-inhibiting hormone (*MIH*), crustacean hyperglycaemic hormone (*CHH*), neuropeptide F (*NPF*) genes/neuropeptides expressed and synthesized in eyestalk are believed to play essential roles in regulating gonad development of *E. sinensis* [15]. Meanwhile, hepatopancreas is an essential organ for energy metabolism, providing essential energy source for gonad development of *E. sinensis* [16]. Studies have also indicated exogenous vitellogenin is synthesized in hepatopancreas and transferred to ovary during vitellogenesis process in *E.sinensis* [17]. However, how the environment factors stimulate and activate the gonad early development and what is the specific biological function of eyestalk and hepatopancreas in regulating gonad development was largely unknown in precocious *E. sinensis*.

E. sinensis is a catadromous species and the life cycle is generally two years. The mating and spawning occurs during winter in brackish water and the fertilized eggs develop into larvae in spring, then the larva will migrate to freshwater rivers/lakes and spend nearly two years with nearly 20-times molting before they get sexual mature [8, 18]. Generally, the gonad development of *E. sinensis* initiates at the second year. As for precocious *E. sinensis*, the gonad starts to develop and get completely sexual mature in the first year [8]. Molting and growth is terminated in sexual mature precocious *E. sinensis* and these individuals are usually discarded due to their unworthiness in the aquaculture [9]. The most direct and accurate way to identify precocious *E. sinensis* is through histology observation which is quite inconvenient during aquaculture for farmers. During the aquaculture process, precocious female *E.*

sinensis individuals are always identified based on the shape of abdominal sternite by

experienced farmers (**Figure 1A**) [15, 19]. The female *E. sinensis* gets sexual mature

when the abdominal sternite completely covers the whole abdomen (Figure 1A).

Previous studies have indicated the ratio of abdominal sternite length is a candidate

phenotype character to discriminate the precocious level in female *E. sinensis* [19]

(Figure 1A). However, researches link the abdominal sternite length to ovary

development stages are limited.

In this study, potential precocious female *E. sinensis* were firstly collected based on

phenotype characters (ratio of abdominal sternite length) and then confirmed by

histology observation. Comparative transcriptome was conducted on eyestalk,

hepatopancreas, and ovary tissues of precocious female *E. sinensis* from different

ovary development stages to 1) provide a convenient way to identify precocious

E.sinensis in aquaculture, 2) reveal the tissue-expression profiles in precocious

female *E. sinensis* in chronological order, 3) identify potential candidate

genes/pathways involved in early ovary development, and 4) provide novel insights

into the genetic network of eyestalk and hepatopancreas in regulating ovary

development in *E. sinensis*.

Materials and methods

Animal sampling and ethics

This study was approved by the Institutional Animal Care and Use Committee

(IACUS) of Shanghai Ocean University (Shanghai, China). Sampling procedures

complied with the guideline of IACUS on the care and use of animals for scientific

purposes. All the *E. sinensis* individuals in this study were collected from the Aquatic

Animal Germplasm Station of Shanghai Ocean University (Shanghai, China). In this

study, potential precocious female *E. sinensis* individuals were firstly collected

according to the ratio of abdominal sternite length (B5-B5'/C5-C5') and then

confirmed by histology observation (**Figure 1**).

Measurement of phenotype characters and histology observation

For the collected *E. sinensis* individuals, phenotype characters such as body weight,

ovary weight and hepatopancreas weight were measured, and abdominal sternite

length were also recorded according to previous research [19]. Hepatopancreas index

(HSI), gonad index (GI), and ratio of abdominal sternite length were calculated

according to the following formula.

Hepatopancreas index = (wet hepatopancreas weight/wet body weight) $\times$100%

Gonad index= (Wet gonad weight/wet body weight) $\times$100%

Rate of abdominal sternite length= (B5-B5')/(C5-C5') (Figure 1A)

Ovary tissues from potential precocious *E. sinensis* individuals were fixed using

Bouin's fixative (Sangon Biotech, China) at room temperature for 24h. Then the

ovary tissue-slices were prepared and stained according to hematoxylin-eosin (HE)

staining method. The tissue-slices were observed under a DM500 microscope system

(LEIKA, Germany) and Image Analysis Software Toup View.

RNA isolation and transcriptome sequencing

According to the histology observation results, precocious *E. sinensis* individuals with

ovary developmental stages in major growth stage I, major growth stage II, and sexual

mature stage were collected with three biological replicates in each group. Meanwhile,

three normal developed two-year old female *E. sinensis* individuals (sexual mature

stage) were also sampled as control group. All the collected *E. sinensis* were

anaesthetized on ice and eyestalk, hepatopancreas, and ovary tissues were quickly

collected and immediately stored in liquid nitrogen before RNA extraction. Total

RNA was extracted from each collected *E. sinensis* individual with RNAiso Reagent

(Takara, China) according to the manufacturer's instructions. The RNA integrity and

quantity were examined using agarose gel electrophoresis and an Agilent 2100

Bioanalyzer (Agilent, Shanghai, China), respectively. A total of 5 μg RNA with an

RNA integrity number (RIN) exceeding 8.0 was used for RNA-seq library

construction using the Truseq™ RNA sample Prep Kit for Illumina (Illumina, USA).

These indexed libraries were sequenced on an Illumina Hiseq™4000, with 150 bp

pair-end reads produced.

Differential expression and GO enrichment analysis

After sequencing, raw sequencing reads were firstly trimmed using Trimmomatic

software[20]. And then, clean reads were mapped to our previously assembled

reference transcriptome assembly (NCBI TSA accession number: GGQO00000000)

using Bowtie 1.0.0 [21]. Gene abundance, the TPM (transcripts per million transcripts)

value was measured using the RSEM 1.3.0 software [22]. The resulting data matrix

with expression value (TPM) for all the samples was generated and used as input data

for differential expression analysis. Then the differentially expressed genes (DEGs)

was measured by DESeq2 software using $P < 0.001$ for the false discovery rate (FDR)

and a fold change $> 2^2$ [23]. After normalizing the DEG TPM values using log2 and

mean centered, cluster analysis was performed using the hierarchical cluster method

based on the euclidean distance using heatmap module in R. GO enrichment analysis

of the DEGs was conducted using TopGO software with an adjusted P value < 0.001

[24]. Pearson correlation was calculated and plotted by corrplot package in R.

qRT-PCR validation

Quantitative real-time PCR (qRT-PCR) was carried out to validate the DEGs

identified in this study. Eight DEGs in ovary, hepatopancreas, and eyestalk were

chosen for qRT-PCR assays. PCR primers were designed according to our previous

reference transcriptome assembly (**Table S1**). In this study, three reference genes

ubiquitin conjugating enzyme (Ube), beta-actin (β-actin), and *ribosomal S27 fusion*

protein (S27) were selected to normalize the gene expression level. qRT-PCR was

conducted using SYBR Green Premix Ex Taq (Takara, China) in a QIAxcel real-time

PCR system (Qiagen, German). A standard curve was firstly generated to assess

amplification accuracy, and primers with an amplification efficiency between 95%

and 105%, and Pearson correlation (R^2) >0.98 were chosen for following qRT-PCR

experiments. Three biological and three technical replicates were chosen for each

selected DEG. The relative expression was estimated using the $2^{-\Delta\Delta Ct}$ method with

normal developed sex mature *E. sinensis* individuals as a calibration control [25].

Relative expression results were presented as the fold-change relative to normal

developed sex mature *E. sinensis* individuals. Statistical significance ($P < 0.05$) was

determined using one-way ANOVA tests under SPSS 25.0.

Data archive

Sequencing reads are available at NCBI SRA database (SRR7777398, SRR7777399,

SRR7777400,   SRR7777401,   SRR7777402,   SRR7777403,   SRR7777404,

SRR7777405, SRR7777406, SRR7777407, SRR7777408, SRR7777409)

Results

Phenotype characters measurement and ovary histology observation

According to the ratio of abdominal sternite length and the histological observation

from collected potential precocious *E. sinensis* individuals, four groups of *E. sinensis*

with different ovary developmental stage were clearly identified. Group I, the ratio

was less than 0.7 and no clearly ovary tissue was discovered; Group II, the ratio was

ranged from 0.71-0.85, the ovary developmental stage was in major growth stage I;

Group III, the ratio was 0.86-1.0 and the ovary developmental stage was in major

growth stage II; Group IV, the ratio was greater than 1.0 and the ovary stage was

completely mature with clearly oocytes (Table 1, Figure 1B). The hepatopancreas

index decreased from 10.45% to 5.40% and the gonad index increased from 0 to 7.59%

accompanied with the abdominal sternite length ratio increased from 0.70 to 1.00.

(**Table 1, Figure 1**).

Tissue gene expression profiles in precocious *E. sinensis*

Regarding ovary tissue, a total of 957 DEGs were identified among different groups

of precocious *E. sinensis*. Three clusters were defined based on the hierarchical

clustering results revealing different expression patterns in the ovary of precocious *E.*

sinensis. GO enrichment analysis indicated that genes in cluster 1 including

sodium-dependent nutrient amino acid transporter 1 (*NAAT1*), MFS-type transporter

(*SLC18B1*), solute carrier family 13 member 3 (*SLC13A3*), solute carrier family 10

member 6 (*SLC10A6*), monocarboxylate transporter 12 (*SLC16A12*),

glucose-6-phosphate exchanger (*SLC37A4*), low-density lipoprotein receptor 1

(*LDLR-A*), nose resistant to fluoxetine protein 6 (*NRF-6*) genes associated with

transmembrane transport for amino acid, creatine, glucose, lipid transport, and

estradiol 17-beta-dehydrogenase 8 (*HSD17B8*) gene associated with estrogen

biosynthetic process were highly expressed in major growth stage II group (**Figure**

2A, 2B Cluster1). Genes associated with oogenesis, border follicle cell migration,

steroid metabolic process and lipid transport were highly expressed in completely sex

mature precocious group enriched in cluster 2, including myosin heavy chain,

non-muscle (*ZIP*), dynamin (*SHI*), Ets DNA-binding protein pokkuri (*AOP*), Protein

catecholamines up (*CATSUP*), very low-density lipoprotein receptor (*VLDLR*),

sulfotransferase 1A1 (*SULTLA1*), sortilin-related receptor (*SORL1*), low-density

lipoprotein receptor-related protein (*LRP*) (**Figure 2A, 2B Cluster 2**). Genes enriched

in cluster 3 such as Delta (24)-sterol reductase (*DHCR24*), dehydrogenase/reductase

SDR family member 11 (*DHRS111*), NPC intracellular cholesterol transporter 1

(*NPC1*) were involved in cholesterol metabolic process, steroid biosynthesis were

highly expressed in major growth stage I (**Figure 2A, 2B, Cluster 3**).

Compared sex mature precocious ovary with normal two-year-old sex mature ovary,

only 11 DEGs were identified, among them up-regulated genes like Neuroparsin-A

(*NPAB*), Lipase 3 (*LIP3*) in precocious ovary were associated with neuropeptide

hormone activity and lipid catabolic process (Table S2, Figure 3).

Considering hepatopancreas tissue, a total of 806 DEGs were identified among

different groups of precocious hepatopancreas. Two clusters were defined based on

the hierarchical clustering results. GO enrichment indicated that genes such as

ATP-binding cassette sub-family A member 3 (*ABCA3*), sodium/bile acid

cotransporter (*SLC10A1*), ileal sodium/bile acid cotransporter (*SLC10A2*), vitelline

membrane outer layer protein 1 (*VMO1*), vitellogenin (*VG*), beta-hexosaminidase

subunit beta (*HEXB*), hemolymph juvenile hormone binding protein (*JHBP*) in cluster

1 associated with lipid transport, vitelline membrane formation, oogenesis were

highly expressed in sex mature precocious group (**Figure 2C, 2D Cluster 1, Figure**

3). Genes such as glycerol-3-phosphate acyltransferase 3 (*GPAT3*), bile salt-activated

lipase (*CEL*), lysosomal acid lipase (*LIPA*), pancreatic triacylglycerol lipase (*PNLIP*)

in cluster 2 were associated with lipid metabolic process (**Figure 2C, 2D Cluster 2**).

Compare normal two-year-old sexual mature hepatopancreas with sexual mature

precocious hepatopancreas, 372 DEGs were identified, Genes up-regulated in

completely sexual mature precocious hepatopancreas like *VMO1*, *VG* were enriched

in vitellogenin synthesis, vitelline membrane formation, lipid transport biological

process (Table S2, Figure 3).

For eyestalk tissue, a total of 1,081 DEGs were identified among different groups of

precocious eyestalks. Two clusters were defined based on the hierarchical clustering

results revealing different expression patterns. GO enrichment analysis indicated

genes such as neuropeptide F (*NPF*), *MIH, CHH*, vasotocin-neurophysin VT 1 (*VT1*),

glycoprotein hormone beta-5 (*GPHB5*), Cytochrome P450 18a1 (*CYP18A1*),

corticotropin-releasing factor-binding protein (*CRHBP*) in cluster 1 highly expressed

in sexual mature precocious eyestalk were associated with neuropeptide hormone

activity, fatty acid biosynthetic process (**Figure 2E, 2F Cluster1, Figure 3**). Genes

such as cuticle protein CP498, cuticle protein AM1159, insulin-like growth

factor-binding protein-related protein 1 (*IGFBP*), platelet-derived growth factor

receptor alpha (*PDGFRA*), protein 60A (*GBB*) in cluster 2 were involved in chitin

metabolic process, growth factor activity (**Figure 2E, 2F Cluster 2**).

Between normal two-year-old sexual mature eyestalk and sexual mature precocious

eyestalk, 449 DEGs were identified, up-regulated genes like *VT1*, *NPF*,

Prohormone-3 (*PROH3*), helicostatins (*helicostatins*), pro-neuropeptide Y (*NPY*) in

completely sexual mature precocious eyestalk were associated with neuropeptide

hormone activity, neuropeptide signaling pathway (Table S2, Figure 3).

DEGs related to neuropeptide hormone activity and lipid transport

Gene expression values from a total of 15 DEGs (GO:0005184, neuropeptide

hormone activity) and 12 DEGs (GO:0006869, lipid transport) from the studied

individuals were extracted. 12 out of 15 DEGs related to neuropeptide hormone

activity were identified in eyestalk tissue and most of the DEGs (*MIH*, *GPHB5*, *NPA*,

CHH, *VT1*, *NPF, RPCH*, *PDH1*, *CCAP*, *NPY*) were upregulated in precocious

eyestalk than in normal sex mature eyestalk. 11 out of 12 DEGs related to lipid

transport were identified in hepatopancreas tissue, and most of the DEGs (*SLC10A1*,

Apolipophorin, *VG*, *LDLR-A*, *SLC10A2*) were upregulated in precocious

hepatopancreas than in normal sex mature hepatopancreas (Table S3).

The correlation coefficient adjacency matrix indicated that DEGs in eyestalk

annotated as neuropeptide hormone activity such as *NPY*, *RPCH* (SP|Q23757),

GHBP5, *NPF*, *ORCKA* (SP|Q9NL83_ORCKA), *CCAP*, Helicostatins

(SP|Q44314_ALLP) were positively correlated with DEGs in hepatopancreas

annotated as lipid transport such as *VG* (SP|Q6RG02), *NPC2* (SP|P61917), *SLC10A1*

(SP|Q14973), *SLC10A2* (SP|Q28727) (*P*<0.05) (Figure 4A, red shade area).

Meanwhile, DEGs in hepatopancreas annotated as lipid transport such as *VG*

(SP|Q6RG02), *NPC2* were positively correlated with DEGs in ovary annotated as

oogenesis, border follicle cell migration, steroid metabolic process, lipid transport

(Figure 4B, yellow shade area) (*P*<0.05). However, most DEGs in eyestalk annotated

as neuropeptide hormone activity were not correlated with DEGs in ovary, only

SLC10A3 (SP|P09131) showed positively correlation with Helicostatins, *RPCH*,

GHBP5, and *CCAP* genes (**Figure S1**).

Discussion

Sexual precocity is a complex biological process with many genes/pathways involved

in specific organs to induce gonad early development [1, 5]. In this study, we utilized

the ratio of abdomen sternite length to discriminate ovary developmental stages and

believed it is a convenient method for the detection of potential precocious *E. sinensis*

in advance. *E. sinensis* with the ratio of abdomen sternite length above 0.70 during the

first year are potential precocious *E. sinensis* that should be abandoned in the

aquaculture. Compared with the expression profiles of ovary between two-year old

and one-year old sex mature *E. sinensis*, only 11 DEGs were identified and the tissue

histology observation showed normal function of precocious ovary which indicated

that precocious *E. sinensis* are functional and able for spawning [2]. *Neuroparsin-A*

(NPA) is believed to be involved in the regulation of insect reproduction through

inhibit the effects of juvenile hormone was highly expressed in precocious *E. sinensis*

indicating that early ovary development may due to the abnormal expression of *NPA*

caused by environment factors [26]. However, comprehensive functional studies need

to be conducted to reveal the molecular mechanism. Our results also provided novel

insights into the genetic interaction of eyestalk and hepatopancreas on ovary early

development of precocious *E. sinensis*.

Significant different gene expression profiles were identified in hepatopancreas and

eyestalk between normal and precocious *E. sinensis*, which indicating the important

function of hepatopancreas and eyestalk in regulating ovary development.

Hepatopancreas is an essential organ for energy storage and metabolism of

crustaceans, providing required energy for the growth and development [16, 27]. It is

also a site for the synthesis and metabolism of certain steroid hormones required by

crustaceans, which plays essential roles in vitellogenesis process [17, 28]. In this

study, DEGs associated with lipid metabolic process were upregulated in major

growth stage I, stage II during vitellogenesis process in precocious *E. sinensis* (Figure

2B, cluster1), indicating the initiation of ovary development depend on the lipid

metabolic in hepatopancreas, which may provide energy and steroid hormones for

early ovary development [29, 30].

Previous studies also pointed out that nutriment like sugar/lipid will be absorbed and

accumulated in hepatopancreas, and excessive nutrition will be transferred to gonad

continuously which will induce gonad early development [31]. Interestingly, DEGs

associated with lipid transport were also upregulated in precocious *E. sinensis*

compared with normal *E. sinensis* in both hepatopancreas and ovary tissues (Figure

2B, Cluster 2, Figure 3). These up-regulated lipid transport related genes may provide

the genetic basis for the lipid transfer from hepatopancreas to ovary [30-32]. The

hepatopancreas index measured in this study decreased with the increased gonad

index also supported that hepatopancreas may transfer required energy/lipid to ovary

(**Figure 1**). Meanwhile, vitellogenin (VG) is the key factor component in

vitellogenesis in *E. sinensis*, which include endogenous VG and exogenous VG.

Endogenous VG is believed to be synthesized by oocyte, while exogenous VG is

believed to be synthesized in hepatopancreas and transferred to ovary [33]. In this

study, *VG* genes were significantly up-regulated in hepatopancreas of precocious *E.*

sinensis, indicating hepatopancreas is the main vitellogenin synthesis site in *E.*

sinensis and all the VG protein required for vitellogenesis originated from

hepatopancreas. Meanwhile, *VG* genes expression in hepatopancreas was positively

correlated with oogenesis, lipid transport DEGs in ovary indicating excessive VG

expression in hepatopancreas may stimulate the ovary development in precocious *E.*

sinensis. *VMO1* which is associated with vitelline membrane formation was also

up-regulated in precocious *E. sinensis*, indicating hepatopancreas may also participate

in the vitelline membrane formation process.

Our results confirmed the essential roles of hepatopancreas in regulating ovary

development, energy and steroid hormone required for oogenesis; vitellogenin

synthesis and vitelline membrane formation for vitellogenesis were fulfilled in

hepatopancreas. The intrinsic genetic factors of sexual precocity in *E. sinensis* at some

content due to the abnormal expression of the above-mentioned candidate DEGs in

hepatopancreas.

It is believed that X-organ-sinus gland complex system existed in eyestalk is an

important neuroendocrine system in crustaceans [7]. Previous study indicated by

regulating the endocrine system of shrimp and crab, the growth rate and the mature

time will be improved, and eyestalk ablation (ESA) is a way to stimulate gonad

development and ovulation [34]. Consistent with previous hypothesis, more

neuropeptide hormone like *NPF*, *NPY*, Prohormone-3 were identified as up-regulated

in precocious eyestalk in this study indicating the essential regulatory mechanism of

eyestalk in ovary development. *NPF* and *NPY* belong to NPY family are neuropeptide

hormone that accelerates ovarian maturation in female *Schistocerca gregaria* [35],

and plays a role in the regulation of visceromotor functions during egg laying [36].

CHH, *MIH* have long been considered to inhibit crustacean molting, and after *E.*

sinensis finish their last reproductive molting like insects, they will stop molting and

initiate their gonad development [37, 38]. Therefore, extremely highly expressed

CHH, *MIH* in eyestalk may inhibit the molting and induce precocity. More

interestingly, DEGs in eyestalk such as *NPY*, *RPCH*, Helicostatins, *GHBP5*, *NPF*,

ORCKA, *CCAP* were positively correlated with *VG* genes expression in

hepatopancreas which indicate these neuropeptide hormone genes may target

hepatopancreas and induce *VG* expression, however, further functional experiments

need to be conducted to confirm the hypothesis (Figure 4B). All the upregulated genes

in precocious eyestalk indicated the neuropeptide hormone synthesized in eyestalk

may stimulate ovary development like HPG axis. Our results pointed out that eyestalk

is indeed essential organ for gonad development and may play essential roles in

regulating vitellogenesis in precocious *E. sinensis*. However, the interplay of eyestalk,

hepatopancreas, and ovary requires more further functional researches.

Sexual precocity is a serious situation in the aquaculture of *E. sinensis*, scientific

researchers and farmers were struggled with the solutions and genetic mechanism of

sexual precocity. In this study, there were few expression differences identified

between precocious and normal sexual mature ovary, and the early development of

ovary may be affected by abnormal developed eyestalk and hepatopancreas. Several

stimulation factors such as high temperature, salt, stock density, nutrition may induce

the metabolic disorder of genes associated with neuropeptide hormone, steroid

hormone as identified in this study, leading to the abundant expression and

accumulation of VG in hepatopancreas and further initiating the ovary development.

However, comprehensive functional studies should be conducted to further reveal the

genetic mechanism of sexual precocity, especially the regulatory mechanism for

eyestalk and hepatopancreas. Our study provides valuable genetic resources for the

research of sexual precocity in *E. sinensis* in future. Meanwhile, the effective

convenient phenotype measurement method and related candidate DEGs identified in

this study will provide guidance for the detection of precocious *E. sinensis* in

aquaculture.

Acknowledgements

This work was funded by Agriculture Research System of Shanghai, China (Grant No.

201804), Shanghai Agriculture Applied Technology Development Program, China

(Grant No. G2017-02-08-00-10-F00076), Shanghai Science and Technology

Committee Programs, China (No.16391905300; No. 13DZ2251800), Leading

Agricultural Talents in Shanghai Project, China (Grant No. D-8004-16-0217).

Conflict of interest

The authors declare no competing interests.

References

1. Parent A-S, Teilmann G, Juul A, Skakkebaek NE, Toppari J, Bourguignon J-P. The Timing of
Normal Puberty and the Age Limits of Sexual Precocity: Variations around the World, Secular
Trends, and Changes after Migration. Endocrine Reviews 2003; 24:668-693.
2. Chang G, Wu X, Cheng Y, Zeng C, Yu Z. Reproductive performance, offspring quality,
proximate and fatty acid composition of normal and precocious Chinese mitten crab *Eriocheir*
sinensis. Aquaculture 2017; 469:137-143.
3. Takada L, Barbero MMD, Oliveira HN, de Camargo GMF, Fernandes Júnior GA,
Aspilcueta-Borquis RR, Souza FRP, Boligon AA, Melo TP, Regatieri IC, Feitosa FLB,
Fonseca LFS, et al. Genomic association for sexual precocity in beef heifers using
pre-selection of genes and haplotype reconstruction. PLOS ONE 2018; 13:e0190197.
4. Guiry A, Flynn D, Hubert S, O'Keeffe AM, LeProvost O, White SL, Forde PF, Davoren P,
Houeix B, Smith TJ, Cotter D, Wilkins NP, et al. Testes and brain gene expression in
precocious male and adult maturing Atlantic salmon (Salmo salar). BMC Genomics 2010;
11:1-18.

5. Abreu AP, Macedo DB, Brito VN, Kaiser UB, Latronico AC. A new pathway in the control of
the initiation of puberty: the MKRN3 gene. Journal of Molecular Endocrinology 2015;
54:R131-R139.
6. Bao C, Yang Y, Huang H, Ye H. Neuropeptides in the cerebral ganglia of the mud crab, *Scylla*
paramamosain: transcriptomic analysis and expression profiles during vitellogenesis.
Scientific Reports 2015; 5:17055.
7. Nagaraju GP. Reproductive regulators in decapod crustaceans: an overview. J Exp Biol 2011;
214:3-16.
8. Wang W, Wang C, Ma X. Ecological Aquaculture of Chinese mitten crab (In Chinese). Beijing,
China: China Agriculture Press; 2013.
9. Xu Y, Li X, Deng Y, Lu Q, Yang Y, Pan J, Ge J, Xu Z. Comparative transcriptome sequencing
of the hepatopancreas reveals differentially expressed genes in the precocious juvenile
Chinese mitten crab, *Eriocheir sinensis* (Crustacea: Decapoda). Aquaculture Research 2017;
48:3645-3656.
10. Li X, Li Z, Liu J, Silva SS. Advances in precocity research of the Chinese mitten crab
Eriocheir sinensis. Aquaculture International 2010; 19:251-267.
11. Li X, Dong S, Lei Y, Li Y. The effect of stocking density of Chinese mitten crab *Eriocheir*
sinensis on rice and crab seed yields in rice–crab culture systems. Aquaculture 2007;
273:487-493.
12. Wu X, Wang Z, Cheng YX, Zeng C, Yang X, Lu J. Effects of dietary phospholipids and highly
unsaturated fatty acids on the precocity, survival, growth and hepatic lipid composition of
juvenile Chinese mitten crab, *Eriocheir sinensis* (H. Milne-Edwards). Aquaculture Research
2011; 42:457-468.
13. Keller R. Crustacean neuropeptides: Structures, functions and comparative aspects.
Experientia 1992; 48:439-448.
14. Böcking D, Dircksen H, Keller R. The Crustacean Neuropeptides of the CHH/MIH/GIH
Family: Structures and Biological Activities. In: Wiese K (ed.) The Crustacean Nervous
System. Berlin, Heidelberg: Springer Berlin Heidelberg; 2002: 84-97.
15. Ci Y. Preliminary research on the molecular mechanism of precocious Chinese mitten crab,
Eriocheir sinensis (In Chinese) Shanghai Ocean University; 2015. Master.
16. Huang S, Wang J, Yue W, Chen J, Gaughan S, Lu W, Lu G, Wang C. Transcriptomic variation
of hepatopancreas reveals the energy metabolism and biological processes associated with
molting in Chinese mitten crab, *Eriocheir sinensis*. Scientific Reports 2015; 5:14015.
17. Wang W, Wu X, Liu Z, Zheng H, Cheng Y. Insights into Hepatopancreatic Functions for
Nutrition Metabolism and Ovarian Development in the Crab *Portunus trituberculatus*: Gene
Discovery in the Comparative Transcriptome of Different Hepatopancreas Stages. PLoS One
2014; 9:e84921.
18. Wang J, Xu P, Zhou G, Li X, Lu Q, Liu X, Zhou J, Wang C. Genetic Improvement and
Breeding Practices for Chinese Mitten Crab, *Eriocheir sinensis*. Journal of the World
Aquaculture Society 2018.
19. Li C, Wang C, Li S. Discriminant analysis on preconciousness of Chinese mitten crab,
Eriocheir sinensis (In Chinese). Fisheries Science & Technology Information 1998:25-28.
20. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina Sequence
Data. Bioinformatics 2014.

21.　Langmead B, Trapnell C, Pop M, Salzberg S. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. Genome Biology 2009; 10:R25.

22.　Li B, Dewey C. RSEM: accurate transcript quantification from RNA-Seq data with or without
a reference genome. BMC Bioinformatics 2011; 12:323.

23.　Anders S, Huber W. Differential expression analysis for sequence count data. Genome
Biology 2010; 11:R106.

24.　A A, J R. topGO: Enrichment analysis for Gene Ontology. R package version 2.22.0. 2010.

25.　Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time
Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. Methods 2001; 25:402-408.

26.　Badisco L, Claeys I, Van Loy T, Van Hiel M, Franssens V, Simonet G, Broeck JV.
Neuroparsins, a family of conserved arthropod neuropeptides. General and Comparative
Endocrinology 2007; 153:64-71.

27.　Wang L, Yan B, Liu N, Li Y, Wang Q. Effects of cadmium on glutathione synthesis in
hepatopancreas of freshwater crab, *Sinopotamon yangtsekiense*. Chemosphere 2008; 74:51-56.

28.　Uawisetwathana U, Leelatanawit R, Klanchui A, Prommoon J, Klinbunga S, Karoonuthaisiri
N. Insights into Eyestalk Ablation Mechanism to Induce Ovarian Maturation in the Black
Tiger Shrimp. PLOS ONE 2011; 6:e24427.

29.　Du N, Lai W, Chen P, Cheng Y, Nan C. Studies on vitellogenesis of *Eriocheir sinensis* (In
Chinese). Acta Zool Sin 1999; 45:88-92.

30.　Wang C, Li S, Li C, Zhao J. Observation and analysis on appearing difference of precocious
crab of Yangtze population and Liaohe population cultured in ponds (In Chinese). Journal of
Lake Sciences 2001:57-62.

31.　Teng W, Cheng Y, Wu X, Yang X, Bian W, Lu Q, Wang W. A comparative study on biological
index changes concerned with gonad development between two population of the Chinese
mitten crab (*Eriocheir sinensis*): Rhine and Yangtze. Journal of Shanghai Fisheries University
2008:65-71.

32.　Li S, Wang C, Zhao N. Studies on gonad development rule of lake stocked mitten crab of
Yangtze population (In Chinese). Acta Hydrobiologica Sinica 2001:350-357.

33.　Wahli W. Evolution and expression of vitellogenin genes. Trends in Genetics 1988; 4:227-232.

34.　Wang X, Zhang Z, Wang Y, Jia X, Li P. Research progress on regulation mechanism of ovary
development in shrimps and crabs. Biotechnology Bulletin 2013:29-35.

35.　Schoofs L, Clynen E, Cerstiaens A, Baggerman G, Wei Z, Vercammen T, Nachman R, De
Loof A, Tanaka S. Newly discovered functions for some myotropic neuropeptides in locusts．
Peptides 2001; 22:219-227.

36.　Rajpara SM, Garcia PD, Roberts R, Eliassen JC, Owens DF, Maltby D, Myers RM, Mayeri E.
Identification and molecular cloning of a neuropeptide Y homolog that produces prolonged
inhibition in aplysia neurons. Neuron 1992; 9:505-513.

37.　Chan S-M, Gu P-L, Chu K-H, Tobe SS. Crustacean neuropeptide genes of the CHH/MIH/GIH
family: implications from molecular studies. General and Comparative Endocrinology 2003;
134:214-219.

38.　Huiyang H, Chunru F, Xuelei C, Jie G, Xiaoshuai H, Haihui Y. Molt．inhibiting hormone
(MIH) gene from the green mud crab *Scylla paramamosain* and its expression during the
molting and ovarian cycle. Aquaculture Research 2015; 46:2665-2675.

Table 1. Information on phenotype characters and histology observation of

precocious *E.sinensis*.

Rate of abdominal sternite length	No. of individuals	Average weight	Average hepatopancreas index	Average gonad index	Ovary developmental stage
<0.70	6	5.62±0.72g	10.45±1.56%	0	Not applicable
0.71-0.85	9	13.72±1.23g	7.16±1.52%	0.16±0.09%	Major growth stage I
0.86-0.99	9	19.47±4.01g	6.03±1.70%	0.34±0.10%	Major growth stage II
≥1.0	6	40.90±1.68g	5.40±0.98%	7.59±1.07%	Mature stage

Figure 1. Phenotype characters, histology observation, and hepatopancreas and gonad

indexes among different ovary developmental stages of precocious *E.sinensis*. A.

Shape of abdominal sternite of female *E.sinensis*. B5-B5' indicated the outside length

of the fifth abdominal sternite, C5-C5' indicated the inside length of the fifth

abdominal sternite. B. Different shapes of abdominal sternite corresponding to

different ovary developmental stages in female *E.sinensis*. C. Average hepatopancreas

and gonad index among different groups of precocious *E.sinensis*. HIS: hepapancreas

index, GI: gonad index.

Figure 2. Expression patterns of differentially expressed genes in precocious ovary,

hepatopancreas and eyestalk tissues. A. Heatmap of differentially expressed genes in

ovary. B. Gene expression plot for the three defined clusters. C. Heatmap of

differentially expressed genes in hepatopancreas. D. Gene expression plot for the two

defined clusters. E. Heatmap of differentially expressed genes in eyestalk. F. Gene

expression plot for the two defined clusters.

P1O, P1HP, P1E indicated precocious ovary, hepatopancreas, and eyestalk with ovary

in major growth stage I; P2O, P2HP, P2E indicated precocious ovary, hepatopancreas,

and eyestalk with ovary in major growth stage II; PMO, PMHP, PME indicated

precocious ovary, hepatopancreas, and eyestalk with ovary in sex mature stage.

Figure 3. Expression profiles of eight differential expressed genes from RNA-Seq

(orange) and qRT-PCR (blue) at different developmental stages of the ovary.

P1O, P1HP, P1E indicated precocious ovary, hepatopancreas, and eyestalk with ovary

in major growth stage I; P2O, P2HP, P2E indicated precocious ovary, hepatopancreas,

and eyestalk with ovary in major growth stage II; PMO, PMHP, PME indicated

precocious ovary, hepatopancreas, and eyestalk with ovary in sex mature stage.

Figure 4. Pearson correlation of gene expression levels of DEGs between tissues

(ovary, hepatopancreas, eyestalk). A. Pearson correlation of gene expression levels of

DEGs between hepatopancreas and eyestalk. Red shade indicated the positively

correlated genes in hepatopancreas and eyestalk ($P<0.05$). B. Pearson correlation of

gene expression levels of DEGs between hepatopancreas and ovary ($P<0.05$).

"X" symbol indicated P value >0.05 for the Pearson correlation.