RAPID AND REPEATED LOCAL ADAPTATION TO CLIMATE IN AN

INVASIVE PLANT

Lotte A. van Boheemen[1*], Daniel Z. Atwater[2], Kathryn A. Hodgins[1]

[1]School of Biological Sciences, Monash University, Clayton, VIC 3800, Australia

[2]Department of Biology, Earlham College, Richmond, IN 47374, USA

*Corresponding author, la.vanboheemen@gmail.com

SUMMARY

• Biological invasions provide opportunities to study evolutionary processes
occurring over contemporary timescales. To explore the speed and repeatability of
adaptation, we examined the divergence of life-history traits to climate, using
latitude as a proxy, in the native North American and introduced European and
Australian ranges of the annual plant *Ambrosia artemisiifolia*.

• We explored niche changes following introductions using climate niche dynamic
models. In a common garden, we examined trait divergence by growing seeds
collected across three ranges with highly distinct demographic histories.
Heterozygosity-fitness associations were used to explore effect of invasion history
on potential success. We accounted for non-adaptive population differentiation
using 11,598 SNPs.

• We revealed a centroid shift to warmer, wetter climates in the introduced ranges.
We identified repeated latitudinal divergence in life-history traits, with European
and Australian populations positioned at either end of the native clines.

• Our data indicate rapid and repeated adaptation to local climates despite the recent
introductions and a bottleneck limiting genetic variation in Australia. Centroid
shifts in the introduced ranges suggest adaptation to more productive
environments, potentially contributing to trait divergence between the ranges.

KEYWORDS

Invasion, trait evolution, climate niche dynamics, local adaptation, latitudinal clines,
climate adaptation, heterozygosity fitness correlations

INTRODUCTION

During biological invasions species commonly spread over large and climatically diverse geographic areas. In doing so, they often re-establish within climatic niches found in their native ranges, or flourish in new environments (Sax & Brown, 2000; Allendorf & Lundquist, 2003; Atwater *et al.*, 2018). Although plasticity and broad ecological tolerance may facilitate success of invaders in many cases (e.g. Geng *et al.*, 2007; Zhang *et al.*, 2010), a growing number of empirical examples suggest that rapid adaptation to local conditions can also enable the establishment and spread of these species in the face of new selective dynamics (Huey *et al.*, 2000; Lachmuth *et al.*, 2011; Colautti & Barrett, 2013; Chown *et al.*, 2014; Turner *et al.*, 2014; Oduor *et al.*, 2016). As such, invasions provide an opportunity to study contemporary adaptive processes, which is key in an era of rapid, human-induced, environmental change.

Comparisons of clinal patterns in life history traits within native and introduced ranges provide an important opportunity to observe if rapid and repeated adaptation can occur in response to similar climatic gradients (Moran & Alexander, 2014). However, many factors could impact trait evolution of native and invasive populations evolving in response to similar climatic gradients resulting in divergent outcomes. Mating system and demographic events, such as bottlenecks, genetic drift and admixture could differentially affect native and invasives' adaptive capacity or influence the route by which adaptive evolution proceeds (Lee, 2002; Facon *et al.*, 2006; Prentis *et al.*, 2008; Rius & Darling, 2014; Estoup *et al.*, 2016). Also the probability of observing trait clines in the introduced range would be low if the strength of climate mediated selection is weak and/or the invasion is recent. By contrast, the invasion history could create patterns of trait variation (e.g., through climate matching (Maron *et al.*, 2004)) mimicking adaptive population divergence (Colautti *et al.*, 2009; Colautti & Lau, 2015). Shifts in the biotic environment could lead divergence of multiple traits linked by trade-offs (Blossey & Notzold, 1995). Dissection of these various mechanisms is required to advance our understanding of the role of rapid evolution in invasive spread (Keller *et al.*, 2009; Bonhomme *et al.*, 2010; Lachmuth *et al.*, 2011; Agrawal *et al.*, 2015; Cristescu, 2015; Dlugosch *et al.*, 2015a). More generally, as many single species have invaded several distinct regions of the globe, comparisons of the native range to multiple successful introductions could illuminate if and when traits evolve in parallel.

Across environmental gradients, trade-offs among life history traits can shape

adaptive trait divergence in response to local conditions and impact the evolutionary

trajectory of trait combinations in invasive populations (Etterson & Shaw, 2001;

Griffith & Watson, 2005; Colautti *et al.*, 2010; Hodgins & Rieseberg, 2011; Colautti

& Barrett, 2013). For example, reductions in season-length at higher latitudes are

frequently reported to select for early flowering at the cost of diminished plant size

(Colautti *et al.*, 2010; Li *et al.*, 2014). Coordinated shifts in life-history traits along

latitudinal gradients within ranges have been documented in several invasive plants

(Dlugosch & Parker, 2008b; Hodgins & Rieseberg, 2011; Colautti & Barrett, 2013).

Latitudinal patterns in plant size could have important evolutionary

consequences for other plant traits. Variation in plant size can influence optimal

resource allocation to male and female sex function (Charnov, 1982; De Jong &

Klinkhamer, 1989; Klinkhamer *et al.*, 1997). In wind-pollinated plants, height can

affect fitness returns directly through more effective pollen dispersal in taller plants

(Burd & Allen, 1988; Klinkhamer *et al.*, 1997), as well as indirectly through increased

availability of resources (Lloyd & Bawa, 1984; De Jong & Klinkhamer, 1989;

Klinkhamer *et al.*, 1997; Zhang, 2006). Outcrossing wind-pollinated plants are

predicted to adaptively change sex allocation to be more male-biased with increase in

size (Lloyd, 1984; De Jong & Klinkhamer, 1989; de Jong & Klinkhamer, 1994;

Klinkhamer *et al.*, 1997), as local seed dispersal should lead to saturating female gain

curves (Lloyd & Bawa, 1984; Sakai & Sakai, 2003), yet more linear male function

gain curves are expected (Klinkhamer *et al.*, 1997; Friedman & Barrett, 2009).

Latitudinal clines in height could therefore be expected to lead to adaptive shifts in

sex allocation. However, studies investigating the evolution of sex allocation patterns

over wide geographic ranges are rare (Guo *et al.*, 2010; Barrett & Hough, 2012).

We examine the repeatability and divergence of important life-history traits in

the native North American and introduced European and Australian ranges of

Ambrosia artemisiifolia. We raised seeds collected at 77 locations from broad climatic

scales in a common garden and accounted for non-adaptive genetic differentiation

using 11,598 genotype-by-sequencing SNPs, as neutral processes could impact trait

variation. We investigate four specific questions: *1) Do native and introduced*

populations occur in similar climates niches? As climate is likely important in

governing trait variation in this species, we examined climatic niche shifts following

introduction to assess how traits might be predicted to diverge within and among the

ranges. *2) Is there evidence for rapid parallel adaptation to latitude?* Repeatable trait
clines for each range along latitudinal gradients, highly associated with many aspects
of climate, would provide strong evidence that rapid adaptation to similar selective
environments has occurred. We additionally explore coordinated shifts in traits
potentially linked by trade-offs. *3) Is there evidence for trait differentiation between*
native and introduced ranges? By examining patterns across multiple introduced
ranges, we can explore if novel recipient communities generated trait divergence
during introduction, or if adaptation to local climates dominates patterns of trait
variation. 4*) Is there a correlation between heterozygosity and fitness related traits?*
Significant correlations would reveal if demographic changes such as bottlenecks,
admixture and inbreeding have likely impacted the evolution of traits during this
species' extensive range expansion. Such correlations are predicted at the individual
and population level in regions that have recently expanded their range, including
those that have undergone admixture (Peischl & Excoffier, 2015).

MATERIAL AND METHODS

Study species

Ambrosia artemisiifolia is a wind pollinated, outcrossing, hermaphrodictic
annual, which has aggressively spread from its native North America to many regions
worldwide (Laaidi *et al.*, 2003; Smith *et al.*, 2013). The first records documenting the
invasion are in central France around 1850 (Chauvel *et al.*, 2006). Multiple
introductions from distinct native sources ensued to both east and west Europe,
resulting in levels of genetic variation equivalent to those found in the native range
(Chun *et al.*, 2010; Gladieux *et al.*, 2010; Gaudeul *et al.*, 2011; van Boheemen *et al.*,
2017). Genetic analysis suggests the Australian populations originate from a
subsequent single introduction event around 80 years ago, derived from the European
introduction, although the exact source is unknown (van Boheemen *et al.*, 2017).
Range expansion likely occurred both north- and southward following this south-
Queensland introduction (Palmer & McFadyen, 2012; van Boheemen *et al.*, 2017)

Latitudinal clines in phenology have been observed within the native North
American range and the introduced ranges of Europe (Chun *et al.*, 2011; Leiblein-
Wild & Tackenberg, 2014) and China (Li *et al.*, 2014), with earlier reproduction and
greater relative investment in reproductive biomass in high-latitudinal compared to
low-latitudinal populations (Chun *et al.*, 2011; Hodgins & Rieseberg, 2011; Li *et al.*,

2014). The wind-spread pollen is a major cause of hayfever worldwide and its

medical treatment costs millions of dollars each year (Taramarcaz *et al.*, 2005),

providing considerable incentive to understand the factors impacting pollen

production in this species.

Climate niche dynamics

To estimate the climatic niche occupied by *A. artemisiifolia* in its native North

American, introduced Eurasian and Australasian ranges, we used ordination-based

species distribution models. Models were taken from a larger study of 835 species

(Atwater *et al.*, 2018), where the methods are described in detail. Briefly, occurrence

data were collected from the Global Biodiversity Information Facility and plotted in

2-dimensional climate space based on rotated component variables (RCA1:

temperature and RCA2: precipitation) of the 19 WorldClim variables (Hijmans *et al.*,

2005) representing annual trends, seasonality and means in temperature or

precipitation (Table S1, Supplementary Material). We selected these variables as they

are commonly used in studies on species distribution and local adaptation including

for studies of *A. artemisiifolia* (e.g. Leiblein-Wild & Tackenberg, 2014; Sun *et al.*,

2017). Geographic and climatic sampling bias was removed to produce estimates of

occurrence probability in each set of climatic variables, for each of the three ranges

(North American, Eurasian, and Australasian). We used Monte Carlo resampling ($n =$

120) to compare observed niche dynamics to those expected using randomly

resampled occurrence points (Atwater *et al.*, 2018). We used Schoener's *D* (Schoener,

1968) to estimate niche overlap between the native range and each introduced range,

and we estimated niche stability, expansion (species occupying climates in their

introduced range that are unoccupied in the native range), and unfilling (species

occupying climates in the native range that are unoccupied in the introduced

range)(Guisan *et al.*, 2014). Finally, we tested whether the location of the niche

centroid differed between native and introduced ranges.

Data collection

To investigate local environments, we described climatic differences between

27 populations in the native range of North America, 32 populations in the introduced

European range, and 18 populations in the introduced Australian range (Fig. S1,

Supplementary Material). We used the 19 WorldClim variables and added a

geographic dimension to the data by including altitude, latitude and longitude, as these variables are shown to be important in *A. artemisiifolia* growth and fitness (Chun *et al.*, 2011; Chapman *et al.*, 2014), with the latter affecting season length and photoperiod (Ziska *et al.*, 2011). To explore associations between climatic and geographic variables in the sampled populations, we applied a principal component analysis (PCA). We opted to present latitudinal trait clines only, as *i)* the primary principle component was highly correlated with latitude (Fig. S2); *ii)* precipitation and temperature variables were strongly associated with latitude (Spearman's ρ^2=0.326-0.417 (Table S2); *iii)* exploratory analyses revealed associations between each trait and many climatic and geographic variable were highly similar to trait~latitude trends (results not shown); and *iv)* is shown to affect *A. artemisiifolia* season length and photoperiod (Ziska *et al.*, 2011).

To assess the potential for adaptive differentiation along latitudinal gradients, we measured trait variation in a common garden of raised seeds collected at broad geographical scales across the three ranges. This is a traditional approach to detect genetic differences among populations (e.g. Bossdorf *et al.*, 2005; Colautti *et al.*, 2009; Hodgins & Rieseberg, 2011; Savolainen *et al.*, 2013; de Villemereuil *et al.*, 2016). We collected seeds in 2013-2014 and randomly selected an average of 12 maternal families with 20 seeds per family from each population (for a full description of data collection methods, see Supplementary Methods). Following a 6-week stratification at 4°C (Willemsen, 1975), we placed seeds in a 30°C germination chamber with 12h light/dark cycle. Two weeks after germination, we planted a randomly selected seedling from each maternal line in a random order into 100ml kwikpot trays with Debco Seed Raising Superior Germinating Mix. We top-watered all plants and artificially manipulated daylight following the light cycle at 47.3°N (median latitude over all sampling locations). One month later, we performed a second transplant (hereafter day 0) to 0.7L pots with Debco mix and 1.5ml slow-release fertilizer (Osmocote Pro, eight to nine months). We examined variation in life-history traits including phenology, growth, and vegetative and reproductive allocation (Table 1).

To assess neutral genetic differentiation underlying trait variation resulting from non-adaptive evolutionary processes, we extracted DNA from leaf tissue of 861 individuals and performed double-digest genotype-by-sequencing library preparation (see Supplementary Methods). We aligned and filtered raw sequences following van

Boheemen *et al.* (2017). Briefly, SNPs were aligned using BWA-mem (Li & Durbin,
2009) to a draft reference genome for *A. artemisiifolia* (van Boheemen *et al.*, 2017).
We called variants with GATK HaplotypeCaller and filtered SNPs using GATK hard-
filtering recommendations (McKenna *et al.*, 2010; Van der Auwera *et al.*, 2013). We
identified a total of 11,598 polymorphic biallelic SNPs with 50% SNP call rate. We
inferred population genetic structure and calculated individual and population level q-
scores for the most likely number of clusters K (=2) with the Bayesian clustering
method STRUCTURE v2.3.4 (Pritchard *et al.*, 2000). We used these STRUCTURE q-
values as a measure of genetic and, therefore, trait differentiation resulting from non-
adaptive (neutral) mechanisms.

To explore the impact of heterozygosity on fitness related traits we calculated
heterozygosity (H_O) as the proportion of heterozygous loci out of the total number of
called genotypes for each individual. Introductions could diminish or increase
heterozygosity, which in turn could inhibit or stimulate invasion (e.g. inbreeding due
to genetic drift or heterosis following admixture).

Statistical analyses

We conducted all statistical analyses in R v3.4.3 (R Core Team, 2017). We
improved normality and reduced heteroscedacity of the data by square root or log-
transforming traits where appropriate. We tested all univariate linear mixed models
using the lme4 package (Bates *et al.*, 2014). To account for demographic history in
patterns of trait divergence, we included STRUCTURE q-scores (individual or
population mean as appropriate) as a random effect in all univariate mixed models.
We calculated significance of fixed effects using type III Wald F tests with Kenward-
Roger's approximation of denominator degrees of freedom and step-wise removed
non-significant ($p>0.05$) fixed effects using the lmerTest package (Kuznetsova *et al.*,
2015), starting with the highest order interactions. To reduce false discovery of
associations due to the number of traits being tested, we 'fdr' corrected p-values
(hereafter q-values) of the combined final models using the p.adjust function, further
reducing models when applicable.

To explore patterns of latitudinal trait divergence among ranges, potentially
indicative of local adaptation, we tested population mean trait responses to range
(native North America, introduced Europe and Australia), latitude, their interaction
and latitude[2] (to account for non-linear trait responses) in multi- and univariate

models ((M)ANCOVA). In these analyses, latitude values are not randomly

distributed among ranges due to the geographic distribution of *A. artemisiifolia*,

suggesting a violation of independence (Miller & Chapman, 2001). However, the

values of the covariate (latitude) are observational and not manipulated by the

independent variables (range) and the (M)ANCOVA assumptions are thus not

violated (Keppel, 1991). We increased the power of the multivariate analysis

(Scheiner, 2001) by removing highly correlated traits (Spearman's $\rho^2 > 0.6$, Table S3,

Supplementary Material) and calculated the approximate F-statistics and Wilks' λ

(multivariate F-value) to measure the strength of the associations.

To explore differences in latitudinal trait clines between ranges as revealed in

the ANCOVAs, we tested for significant two-way interactions between range and

latitude which is indicative of non-parallel trait~latitude slopes among the ranges. To

further dissect the extent of trait divergence and its dependence on latitude, we

compared model estimates of traits in the introduced ranges to the native estimates at

the minimum and maximum observed latitude in each range, where applicable (EU_{min},

NA_{max}, NA_{min} and AU_{max}) (Fig. S4, Supplementary Material). We tested overall

pairwise range differences in trait values for significant range effects, when higher

order interactions involving range were not significant. We explored the highest order

significant interactions, using χ^2 tests with Holm p-value correction using the phia

package (De Rosario-Martinez, 2013.).

We verified the presence of the well-described plant height-flowering time

trade-off and examined associated patterns in other reproductive traits (dichogamy

and sex allocation) within ranges. We tested linear relations between individual trait

values of control treated plants in mixed models, with plant height, range and their

interaction as fixed effects. In addition to individual STRUCTURE q-values, we

added population as random effect. A significant interaction between range and height

indicated a differential slope between the focal reproductive trait and plant height

between ranges.

To investigate geographical patterns in genetic diversity of populations, we

tested the effect of range, latitude, their interaction and $latitude^2$ on population mean

H_O. To explore the effect of heterozygosity on selected traits, we included population

mean H_O, latitude, their interactions with range and $latitude^2$ as fixed effects. To

reduce false positive association due to multiple testing, we only tested the response

of growth (plant height and biomass) and fitness (total reproductive biomass and

average seed weight) related traits. As we identified signatures of repeated local

adaptation in floral sex allocation (Results), we also tested the response of this trait.

RESULTS

Climate niche dynamics

Niche overlap (D) was significantly lower than expected between the North

American native and Eurasian invasive range ($P<0.001$) although the native and

Australasian range did not have significantly lower overlap than expected ($P=0.425$).

However, niche stability was low between the native and both invasive ranges

($P<0.001$). Climatic niche unfilling and expansion were not significantly different

than the null model, except that especially low expansion was found in the Eurasian

population ($P=0.017$), meaning that while the niche shifted, the species did not tend to

occupy completely novel climates in its Eurasian range. In both introduced ranges the

niche centroid shifted significantly towards hotter, wetter climates ($P<0.001$; Fig. 1).

We note that this climate analysis compares the entire Eurasian and Australasian

ranges with the native North American range. For our common garden analysis we

focused only on specific latitudinal transects in Europe and Australia to examine how

traits have evolved along latitudinal clines during invasion, and did not include any

Asian populations. We include this more general analysis of climate niche to assess

how climate shifts might contribute to trait divergence among the ranges generally,

although our sampling for the common garden was more limited.

Climate of sampled populations

Climatic variables across all populations in the native North American,

introduced European and introduced Australian ranges could be effectively

summarized by the first two principle components (PC) in the PCA (Fig. S2, Table

S2, Supplementary Material), which together explained 70.06% of between-

population variation. Here, PC1 was strongly associated with latitude, temperature

and seasonality, whereas PC2 was mostly precipitation related (Fig S2, Table S2).

The climate experienced by Australian populations was distinct from the North

American and European ranges (Fig. S2, Table S2), with higher annual, winter and

summer temperatures, and with lower seasonal variation. Moreover, given the sub-

tropical location of Australian populations, the sampled populations experienced

higher precipitation during the growing season (Fig. S2).

Repeatability in trait clines among ranges

Traits across all North American, European and Australian ranges were well

summarized by the PCA, where the first two PCs explained 83.1% of all variation

(Fig. S3, Supplementary Material). The main PCs were associated with each trait to a

similar extent so no trait syndromes were apparent. Traits measured in Australian

plants were generally distinct from the other populations although there was some

overlap in multivariate space among the ranges (Fig. S3, Table S3). Multivariate trait

analyses revealed a significant two-way interaction between latitude and range

($F_{14,120}$=1.796, p=0.047, Wilks' λ=0.684) (Table S4) suggesting latitudinal trait clines

exist, but do not have the same relationship within ranges for all traits. Further

dissection of these patterns in univariate analyses revealed maximum growth rate,

flowering onset, dichogamy, average seed weight, total reproductive biomass and

specific leaf area (SLA) displayed similar latitudinal clines among ranges, indicated

by a significant latitude effect but an absence of a higher-order interaction (Fig. 2,

Table 2). We identified range differences in latitudinal trait clines for maximum

height, total biomass, floral sex allocation and relative reproductive biomass, as

indicated by significant range:latitude interactions. However, all of these slopes were

significantly different from zero and were in the same direction as the native North

American patterns (Fig. 2, Table S5a). These results are consistent with local

adaptation along a latitudinal gradient in all three ranges.

At higher latitudes, plants were shorter, weighed less, reached lower

maximum growth rates and flowered earlier. Flowering onset extremes were 14-133

days after transplant (population means for EU20 and AU13, Fig. 2). In all ranges,

dichogamy (the temporal separation of pollen dispersal and emergence of receptive

stigma within an individual plant) was prevalent. Protogyny (emergence of stigma

prior to pollen release) predominated at higher latitudinal populations, with receptive

stigmas being visible up to 40 days before any pollen was released within the same

plant (EU20)(Fig. 2). Conversely, protandry prevailed at latitudes below 40° from the

equator, with pollen release occurring up to 14 days before stigmas became receptive

(NC)(Fig. 2). Floral sex allocation followed a similar trend across ranges, with a slight

male function bias at lower latitudes, shifting towards an extreme female function bias at high-latitude populations (Fig. 2). The biggest seeds were found at mid-latitudinal populations (38.6°N in KY), with seeds decreasing in size towards the high and low latitudes (Fig. 2, Table 2). Total reproductive biomass also showed a similar curved relationship, with the combined weight of racemes (male floral sex function) and seeds (female floral sex function) being up to three times as high at mid-latitudinal populations compared to high-latitudinal plants. In contrast, the relative reproductive biomass increased with latitude. Within each range, plants from lower latitudes had lower SLA (Fig. 2).

Trait divergence between ranges

While latitudinal trait clines were repeatable for many traits as described above, we identified shifts in trait values at comparable latitudes as revealed by significant range effects (Table 2). Maximum growth rates were highest in Europe and lowest in Australia (Fig. 2, Table 3a). European plants also flowered later than North American and Australian plants at similar latitudes. The temporal separation between pollen release and the appearance of receptive stigma (dichogamy) was greater in the native North America compared to Europe (Fig. 2, Table 3a). European seeds were heavier and plants had higher total reproductive biomass than those measured in the other ranges. At any given latitude, Australian leaves had higher SLA compared to the native range, with lowest SLA in European populations (Fig. 1, Table 3a).

Dissection of range differences in latitudinal trait clines (maximum plant height, total biomass, floral sex allocation and relative reproductive biomass) revealed most significant interactions between range and latitude were prompted by clinal differences between the introduced European and native North American populations (Table S5, Supplementary Material). For these traits, the discrepancy between North American and European trait values increased with increasing latitudes, such that at high-latitude populations, European plants were taller, heavier and less female-biased in floral sex allocation (Table 3b). Moreover, Australian plants found closest to the equator were significantly shorter than native North American expectations (Table 3b).

Trade-offs between life-history traits

We tested for the presence of a trade-off between plant height and flowering

time and investigated associated patterns in dichogamy and floral sex allocation and

height. As expected, taller plants started flowering later in both the native and the

introduced European range, though this pattern was not significant in introduced

Australian populations (Fig. 3, Table 4, Table S6, Supplementary Material).

Correspondingly, we observed protogyny and a large female-biased sex allocation in

short plants versus protandry with a slight male bias in tall plants. These dichogamy

associations with height were not significant in Australia (Fig. 3, Table 4 & S6).

However, it is possible that the height of some large Australian plants might have

been truncated due to greenhouse conditions.

Associations between heterozygosity and life-history traits

To identify geographic patterns in observed heterozygosity (H_O), we tested the

effect of range, latitude and their interaction on H_O. We found no latitudinal patterns

in H_O varying within ranges (range:latitude, χ^2_1=3.811, p=0.149) or among all ranges

(latitude, χ^2_1=0.000, p=0.986). We did identify variable between ranges (range,

(χ^2_1=6.446, p=0.040), resulting from significantly lower H_O in Australia compared to

native North America (χ^2_1=6.446; p=0.033). When accounting for latitude and

population genetic structure, we found a significant interaction effect between mean

population H_O and range on total biomass (Fig. 4, Table 5). Pairwise range

comparisons in post-hoc tests revealed a higher H_O that was associated with heavier

Australian plants (Table S7, Supplementary Material). Moreover, we found that mean

population H_O was positively correlated with seed size in all ranges (Fig. 4, Table 5).

We found no effect of individual genomic heterozygosity on plant height, phenology,

dichogamy, total or relative reproductive investment and floral sex allocation (Table

5).

DISCUSSION

We show genetically based differentiation along multiple latitudinal clines in

all examined traits including plant size, growth, reproductive investment, phenology,

dichogamy, SLA, and sex allocation. Remarkably, the clinal patterns apparent in the

native range evolved repeatedly within both introduced ranges over the course of only

100-150 years and despite limited neutral genetic variation in the introduced

Australian range. These patterns are consistent with rapid adaptation, as we accounted

for neutral genetic differentiation. Moreover, low effects of maternal environment are

expected (Hodgins & Rieseberg, 2011) and the introduction history of this species

(van Boheemen *et al.*, 2017) reveals climate-matching (Maron *et al.*, 2004) is

unlikely. The adaptive trait divergence at similar latitudes, together with a centroid

shift to warmer and wetter climates in the introduced ranges, could suggest invasive

populations have adapted to more productive environments following introduction.

The observed rapid evolution has implications for the evolutionary potential of this

species and further range expansion following climate change. Furthermore, the

divergence of reproductive traits such as flowering time, sex allocation and seed size

during recent range expansion should impact the production of allergenic pollen as

well as the abundance and dispersal of seed that could impact spread.

Climate niche shifts

Higher resource levels, such as increased water availability, are a known

contributor to invasion in many plant species (Blumenthal, 2006; Dlugosch *et al.*,

2015b). Increased resource availability may occur through a shift in the fundamental

or realized niche during invasion. The latter can result from reductions in competition,

perhaps reflecting the presence of a vacant niche in the introduced range (e.g.

Dlugosch *et al.*, 2015b). Climate niche dynamics analysis reveals higher *A.*

artemisiifolia abundance in warmer and wetter climates in the introduced ranges

compared to the native range. It is possible that this centroid shift reflects an historic

effect where colonization of warmer and wetter environments occurred earlier, or

perhaps by genotypes pre-adapted to these climates (*but see* van Boheemen *et al.*,

2017). Alternatively, the shift might reflect changes in biotic interactions leading to

greater abundance of this species in high resource environments or differences in the

availability of these climates in the introduced regions. Evolutionary processes that

allow introduced species to colonize warmer and wetter environments than those

occupied by native plants could also cause centroid shifts. This evolutionary

interpretation is supported as Australian populations follow trait trajectories parallel

to, but extending beyond, those of the native range.

Repeated latitudinal clines

Our common garden experiments using samples collected across multiple

similar latitudinal gradients, reveal that local adaptation can happen quickly and

predictably, with repeated evolution of native clines in both of the introduced ranges.

Latitudinal clines in phenology and size are a common feature of many

geographically widespread plant species (e.g. Colautti *et al.*, 2010; Li *et al.*, 2014),

with *A. artemisiifolia* flowering shown to be driven by the association between season

length and latitude (Ziska *et al.*, 2011). Short season lengths at high latitudes can

select for earlier flowering (Bradshaw & Holzapfel, 2008; Colautti & Barrett, 2013),

while the evolution of delayed flowering at low latitudes reflects the trade-off

between size and the timing of reproductive maturity, where fitness is maximized by

flowering later at a large size (Colautti *et al.*, 2010; Colautti & Barrett, 2013). This

correlation between plant size and flowering time has been reported for *A.*

artemisiifolia (Hodgins & Rieseberg, 2011; Leiblein-Wild & Tackenberg, 2014;

Scalone *et al.*, 2016) and our results are consistent with rapid genetic differentiation in

plant size, growth rates and phenology in response to latitude-associated selection

pressures such as season length.

We exposed repeated patterns of genetic differentiation in sex allocation

strategy over similar latitudinal clines, consistent with rapid adaptation to local

climate. Plants sourced from higher latitudes displayed female-biased sex-allocation

and protogyny, with more balanced floral sex allocation and a decrease in the

temporal separation of male and female function towards the equator. Previous studies

on *A. artemisiifolia* showed plasticity for sex-allocation and dichogamy in relation to

plant size (Paquin & Aarssen, 2004; Friedman & Barrett, 2009; Friedman & Barrett,

2011) and ample genetic variation for evolution to act on (Friedman & Barrett, 2011).

The patterns observed this study are consistent with theory on sex allocation in

outcrossing wind-pollinated hermaphrodites with local seed dispersal. Spatial

heterogeneity has been observed in animal pollinated plants, where small, resource

limited individuals often allocate more resources to male function (Korpelainen,

1998; Guo *et al.*, 2010). However, these studies are on wild populations and cannot

separate environmental and genetic effects in allocation patterns along resource

gradients. Our current findings from common garden experiments are therefore novel

in identifying genetic differentiation among populations in sex allocation over spatial

gradients.

Trait divergence among the ranges

Many hypotheses aim to explain the success of invasive species, including the

evolution of increased competitive ability (EICA) through escape from native

herbivores (Blossey & Notzold, 1995). Moreover, if trade-offs between performance

and abiotic stress tolerance occur, greater resources could facilitate the evolution of

more competitive phenotypes in introduced ranges (Grime, 1977; Bossdorf *et al.*,

2005; He *et al.*, 2010; Dlugosch *et al.*, 2015b). Our reported trait shifts in European

population compared to natives at equivalent latitudes indeed suggest an increase in

competitive ability, commonly measured as elevated plant growth and reproductive

effort (Felker□Quinn *et al.*, 2013). As no evidence for EICA is found in Europe

(Hodgins & Rieseberg, 2011), these observations might reflect the warmer and wetter

European climate (Fig. 1). Notably, although Europe was identified as the

introduction source for Australian populations (van Boheemen *et al.*, 2017), traits

measured within each range were highly dissimilar. Most of the sampled Australian

populations extended beyond absolute latitudes of the other populations and were

situated in warmer, less seasonal climates (Fig. S2). These factors might explain

Australian trait variation beyond values observed in source populations.

Heterozygosity and invasion

Genetic drift within small founding populations and on the invasion front can

lead to reduced genetic diversity, potentially impacting additive genetic variation

(Dlugosch & Parker, 2008a; Excoffier & Ray, 2008; Peischl *et al.*, 2013; Bock *et al.*,

2015). In *A. artemisiifolia*, Australian populations were bottlenecked and potentially

subjected to high genetic drift, whereas multiple introductions into Europe from

distinct native sources has implicated admixture as a driver of invasion success (van

Boheemen *et al.*, 2017). We found the biomass of Australian, but not European, plants

was indeed associated with heterozygosity, providing only partial support for the

fitness benefits of heterozygosity during invasion (Peischl & Excoffier, 2015).

Admixture and heterosis are unlikely to be main drivers of invasiveness in Europe, as

we found no relationship between population level heterozygosity and any trait other

than seed size. Indeed, in most experimentally admixed European and native *A.*

artemisiifolia crosses heterosis was absent (Hahn & Rieseberg, 2017). These

observations suggest demographic processes had very limited consequences (negative

or positive) on the fitness of these introduced populations. However, local adaptation

of life history traits such as plant size across broad environmental gradients may mask

heterozygosity-fitness correlations. Future studies could address this question by
examining the association in single populations (e.g. Conte *et al.*, 2017).

In plants, reduced seed size is one trait that could aid dispersal and might
therefore be expected to evolve during range expansion (Bartle *et al.*, 2013; Huang *et*
al., 2015). Spatial sorting for dispersal traits at the expansion front has been well
documented in other invasions, such as the cane toads (Estoup *et al.*, 2004; Phillips *et*
al., 2006). In Europe, spatial sorting for increased dispersal, and therefore smaller
seeds, could have occurred at the range edge during expansion northwards. However,
this mechanism would only explain the seed size decline in low-latitudinal
populations in Australia, where range expansion likely occurred both north- and
southward (Palmer & McFadyen, 2012; van Boheemen *et al.*, 2017). The association
between seed size and mean population heterozygosity we identified in all three
ranges could be expected when small seeds aid dispersal, as founder effects should
also reduce heterozygosity during colonization. Though additional factors likely shape
seed size evolution, our findings suggest seed size divergence could represent an
important difference in life-history strategies between ranges. Moreover, we observed
patterns indicating a relationship between genomic and ecological dynamics
potentially linked to range expansion and colonization.

Conclusion

Invasive species often exhibit rapid adaptation despite facing novel selective
pressures (Lachmuth *et al.*, 2011; Colautti & Barrett, 2013; Chown *et al.*, 2014;
Turner *et al.*, 2014). Moreover, the success of invasives is considered paradoxical as
strong demographic changes are predicted to enhance inbreeding and reduce genetic
variation and, consequently, evolutionary potential (Allendorf & Lundquist, 2003).
We investigated these apparent contradictions in a comprehensive study. We
compared the native range with multiple introduced ranges with highly distinct
demographic histories, characterized similarities and shifts in climatic niches, tested
adaptation in a large number of life-history traits and explored heterozygosity-fitness
associations while accounting for non-adaptive population differentiation. We found
strong evidence for parallel adaptation in all three ranges. This study therefore
emphasizes that although introduction dynamics can affect genetic diversity
(Dlugosch & Parker, 2008a) the adaptive potential of those traits might not be
constrained to a similar extent (Dlugosch *et al.*, 2015a).

ACKNOWLEDGEMENTS

We thank J. Stephens and A. Wetherhill for sample collection, S. Bou-Assi, M. Kourtidou, J. Taylor, G. Boinnard, T. Freeman and E. Barnett for greenhouse assistance, S. Bou-Assi and K. Nurkowski for genomic analyses and T. Connallon for manuscript suggestions. A Monash University Dean's International Postgraduate Research Scholarship was provided to LAB, a Monash University Start-up Grant to KAH.

AUTHOR CONTRIBUTIONS

LAB and KH designed the project, with data collection and analyses carried out by LAB and refined by KH. DZA developed and carried out the niche distribution modelling. All authors discussed the results and contributed to the MS writing.

DATA ACCESSBILITY

Sequence data are available at the National Center for Biotechnology Information (NCBI) Sequence Read Archive under Bioproject PRJNA449949.

558   **Table 1**. Traits included in this study.

Trait	Description
Max. height	Maximum measured height
Total biomass	Above- and belowground dry biomass
Max. growth rate	Sensu Chuine *et al.* (2001)
Flowering onset	First recorded day of flowering (number of days after second transplant); first day of pollen release (male function) or receptive female function
Dichogamy	First recorded day of pollen release - first recorded day of receptive female function (a positive value is protogyny, a negative value is protandry)
Floral sex allocation (female/male)	Female (seeds) / male (raceme) dry weight (a value >1 is higher biomass allocation to female function)
Weight per seed	Dry weight per seed in milligrams, averaged over 20 seeds (where available)
Total reproductive biomass	Female (seeds) and male (raceme) dry weight
Relative reproductive biomass	Total reproductive biomass / total plant biomass
Specific leaf area	Leaf area of fully expanded fresh leaf/leaf dry weight

560 **Table 2.** *Ambrosia artemisiifolia* population mean trait responses to range, latitude, their interaction and latitude[2], with population q-values as random effects. We reported Wald type III F-values, with Kenward-Roger degrees of freedom as subscript and symbols specifying significance (fdr corrected q) of effect.

	Range	Latitude	Latitude[2]	Range:Latitude
Max. height	$3.886_{2,70.57}$*	$55.963_{1,71.9}$***	$0.581_{1,70.97}$(ns)	$4.068_{2,70.35}$*
Total biomass	$3.344_{2,69.25}$#	$3.227_{1,70.98}$#	$5.214_{1,70.97}$*	$3.667_{2,69.14}$*
Max. growth rate	$6.405_{2,71.98}$**	$21.658_{1,70.76}$***	$2.025_{1,73}$(ns)	$1.101_{2,69.14}$(ns)
Flowering onset	$6.862_{2,71.978}$**	$158.301_{1,70.763}$***	$1.102_{1,73}$(ns)	$3.43_{2,69.14}$#
Dichogamy	$10.521_{1,73}$**	$8.163_{2,71.18}$**	$6.805_{1,72.99}$*	$3.157_{2,69.14}$#
Floral sex allocation (female/male)	$6.354_{2,69.82}$**	$6.048_{1,66.76}$*	$7.949_{1,66.27}$*	$7.094_{2,69.77}$**
Weight per seed	$4.156_{2,66.13}$*	$25.758_{1,72.41}$***	$23.704_{1,72.88}$***	$0.18_{2,67.29}$(ns)
Total reproductive biomass	$27.478_{1,73}$***	$6.856_{2,71.18}$**	$20.509_{1,72.99}$***	$3.427_{2,69.14}$#
Relative reproductive biomass	$6.436_{2,70.57}$**	$22.839_{1,71.9}$***	$2.997_{1,70.97}$(ns)	$6.422_{2,70.35}$**
Specific leaf area	$7.626_{2,67.39}$**	$17.167_{1,66.45}$***	$1.807_{1,63.65}$(ns)	$3.003_{2,65.14}$#

ns q>0.1; # q<0.1; * q<0.05, ** q<0.01; *** q<0.001

Table 3. Range differences of *A. artemisiifolia* population mean traits at comparable

latitudes for significant (q<0.05) range effects (a, Table 2) and trait differences

between ranges at minimum (min) and maximum (max) latitudes (Figure S4,

Supplementary Material) for significant range:latitude interactions by comparing trait

values at (b, Table 2).

a)	North America - Europe	North America - Australia	Europe - Australia
Max. growth rate	12.328_1**	4.264_1*	1.278_1**
Flowering onset	16.465_1***	0.123_1(ns)	13.269_1***
Dichogamy	18.8_1***	0.077_1(ns)	4.87_1#
Weight per seed	7.44_1*	2.059_1(ns)	6.002_1*
Total reproductive biomass	15.228_1***	0.811_1(ns)	6.516_1*
Specific leaf area	2.734_1#	14.936_1***	14.736_1***

)	North America-Europe	North America-Australia	Europe-Australia
Max. growth rate	13.269_1***	4.178_1*	11.606_1**
Flowering onset	16.465_1***	0.123_1(ns)	13.269_1***
Dichogamy	18.8_1***	0.077_1(ns)	4.87_1#
Weight per seed	7.44_1*	2.059_1(ns)	6.002_1*
Total reproductive biomass	15.228_1***	0.811_1(ns)	6.516_1*
Specific leaf area	2.734_1#	14.936_1***	14.736_1***

b) Trait	North America (NA) - Europe (EU)		North America (NA) - Australia (AU)	
	EU_{min}	NA_{max}	NA_{min}	AU_{max}
Max. height	8.679_1**	25.236_1***	4.818_1#	0.039_1(ns)
Total biomass	0.339_1(ns)	15.572_1***	0.564_1(ns)	1.307_1(ns)
Floral sex allocation (female/male)	0.012_1(ns)	26.417_1***	1.586_1(ns)	0.096_1(ns)
Relative reproductive biomass	4.289_1(ns)	7.557_1*	0.065_1(ns)	0.017_1(ns)

579 ns q>0.1; # q<0.1; * q<0.05, ** q<0.01; *** q<0.001

580 **Table 4**. Flowering time and sex function allocation responses to maximum plant
581 height (Height), range and their interaction, with individual q-values and population
582 as random effects. We reported Wald type III F-values test, Kenward-Roger degrees
583 of freedom as subscript and symbols specifying significance of effect.
584

Trait	Range	Height	Range:Height
Flowering onset	$20.658_{2,606.14}$***	$105.082_{1,766.54}$***	$14.767_{2,671.02}$***
Dichogamy	$6.503_{510.44}$**	$51.235_{1,670.96}$***	$10.621_{2,545.18}$***
Floral sex allocation (female/male)	$17.451_{2,508.47}$***	$138.084_{1,659.31}$***	$16.963_{2,508.47}$***

585 ns q>0.1; # q<0.1; * q<0.05, ** q<0.01; *** q<0.001

586 **Table 5.** Trait responses (population mean) to range, latitude, latitude^2, heterozygosity (H$_O$) and interactions, with population q-values as random

587 effects. We reported Wald type III F-values test values, Kenward-Roger degrees of freedom as subscript and symbols specifying significance of

588 effect.

	H$_O$	Range	Latitude	Latitude^2	H$_O$:Range	Range:Latitude
Max. height	$0.007_{1,70.75}$(ns)	$3.435_{2,69.52}$#	$51.007_{1,70.95}$***	$0.198_{1,69.96}$(ns)	$0.095_{2,59.46}$(ns)	$3.699_{2,69.23}$#
Total biomass	**$8.455_{1,53.099}$***	$4.942_{2,65.157}$*	$108.362_{1,68.316}$***	$0.699_{1,69.812}$(ns)	**$3.947_{2,64.456}$***	$2.055_{2,66.152}$(ns)
Total reproductive biomass	$0.263_{1,71.8}$(ns)	$7.123_{2,70.12}$**	$19.116_{1,71.91}$***	$24.912_{1,71.96}$***	$0.274_{2,59.46}$(ns)	$1.721_{2,68.2}$(ns)
Weight per seed	**$5.858_{1,38.45}$***	$3.271_{2,47.3}$#	$23.516_{1,71.93}$***	$20.776_{1,71.84}$***	$0.069_{2,25.95}$(ns)	$0.332_{2,55.71}$(ns)
Floral sex allocation (female/male)	$0.12_{1,69.63}$(ns)	$6.619_{2,68.56}$**	$6.229_{1,69.87}$*	$7.55_{1,69.84}$*	$0.096_{2,59.46}$(ns)	$7.287_{2,68.47}$**

590 ns q>0.1; # q<0.1; * q<0.05, ** q<0.01; *** q<0.001

REFERENCES

Agrawal AA, Hastings AP, Bradburd GS, Woods EC, Züst T, Harvey JA, Bukovinszky T. 2015. Evolution of plant growth and defense in a continental introduction. *The American Naturalist* **186**: E1-E15.

Allendorf FW, Lundquist LL. 2003. Introduction: population biology, evolution, and control of invasive species. *Conservation Biology* **17**: 24-30.

Atwater DZ, Ervine C, Barney JN. 2018. Climatic niche shifts are common in introduced plants. *Nature Ecology & Evolution* **2**: 34.

Barrett SC, Hough J. 2012. Sexual dimorphism in flowering plants. *Journal of Experimental Botany*: 67-82.

Bartle K, Moles AT, Bonser SP. 2013. No evidence for rapid evolution of seed dispersal ability in range edge populations of the invasive species *Senecio madagascariensis*. *Austral Ecology* **38**: 915-920.

Bates D, Maechler M, Bolker B, Walker S. 2014. lme4: Linear mixed-effects models using Eigen and S4. *R package version* **1**: 1-23.

Blossey B, Notzold R. 1995. Evolution of increased competitive ability in invasive nonindigenous plants: a hypothesis. *Journal of Ecology* **83**: 887-889.

Blumenthal DM. 2006. Interactions between resource availability and enemy release in plant invasion. *Ecology Letters* **9**: 887-895.

Bock DG, Caseys C, Cousens RD, Hahn MA, Heredia SM, Hübner S, Turner KG, Whitney KD, Rieseberg L. 2015. What we still don't know about invasion genetics. *Molecular Ecology* **24**: 2277-2297.

Bonhomme M, Chevalet C, Servin B, Boitard S, Abdallah JM, Blott S, San Cristobal M. 2010. Detecting selection in population trees: the Lewontin and Krakauer test extended. *Genetics*.

Bossdorf O, Auge H, Lafuma L, Rogers WE, Siemann E, Prati D. 2005. Phenotypic and genetic differentiation between native and introduced plant populations. *Oecologia* **144**: 1-11.

Bradshaw WE, Holzapfel CM. 2008. Genetic response to rapid climate change: it's seasonal timing that matters. *Molecular Ecology* **17**: 157-166.

Burd M, Allen T. 1988. Sexual allocation strategy in wind-pollinated plants. *Evolution* **42**: 403-407.

Chapman DS, Haynes T, Beal S, Essl F, Bullock JM. 2014. Phenology predicts the native and invasive range limits of common ragweed. *Global Change Biology* **20**: 192-202.

Charnov EL. 1982. The theory of sex allocation. *Monographs in Population Biology* **18**: 1-355.

Chauvel B, Dessaint F, Cardinal-Legrand C, Bretagnolle F. 2006. The historical spread of *Ambrosia artemisiifolia* L. in France from herbarium records. *Journal of Biogeography* **33**: 665-673.

Chown SL, Hodgins KA, Griffin PC, Oakeshott JG, Byrne M, Hoffmann AA. 2014. Biological invasions, climate change and genomics. *Evolutionary Applications* **8**: 23-46.

Chuine I, Aitken SN, Ying CC. 2001. Temperature thresholds of shoot elongation in provenances of *Pinus contorta*. *Canadian Journal of Forest Research* **31**: 1444-1455.

Chun YJ, Fumanal B, Laitung B, Bretagnolle F. 2010. Gene flow and population admixture as the primary post-invasion processes in common ragweed (*Ambrosia artemisiifolia*) populations in France. *New Phytologist* **185**: 1100-1107.

Chun YJ, V LEC, Bretagnolle F. 2011. Adaptive divergence for a fitness-related trait among invasive *Ambrosia artemisiifolia* populations in France. *Molecular Ecology* **20**: 1378-1388.

Colautti RI, Barrett SC. 2013. Rapid adaptation to climate facilitates range expansion of an invasive plant. *Science* **342**: 364-366.

Colautti RI, Eckert CG, Barrett SC. 2010. Evolutionary constraints on adaptive evolution during range expansion in an invasive plant. *Proceedings of the Royal Society B: Biological Sciences* **277**: 1799-1806.

Colautti RI, Lau JA. 2015. Contemporary evolution during invasion: evidence for differentiation, natural selection, and local adaptation. *Molecular Ecology* **24**: 1999-2017.

Colautti RI, Maron JL, Barrett SCH. 2009. Common garden comparisons of native and introduced plant populations: latitudinal clines can obscure evolutionary inferences. *Evolutionary Applications* **2**: 187-199.

Conte GL, Hodgins KA, Yeaman S, Degner JC, Aitken SN, Rieseberg LH, Whitlock MC. 2017. Bioinformatically predicted deleterious mutations reveal

complementation in the interior spruce hybrid complex. *BMC Genomics* **18**:

970.

Cristescu ME. 2015. Genetic reconstructions of invasion history. *Molecular Ecology*

24: 2212-2225.

De Jong T, Klinkhamer P. 1989. Size-dependency of sex-allocation in

hermaphroditic, monocarpic plants. *Functional Ecology*: 201-206.

de Jong TJ, Klinkhamer PG. 1994. Plant size and reproductive success through

female and male function. *Journal of Ecology*: 399-402.

De Rosario-Martinez H 2013. Phia: post-hoc interaction analysis. *R package*.

de Villemereuil P, Gaggiotti OE, Mouterde M, Till-Bottraud I. 2016. Common

garden experiments in the genomic era: new perspectives and opportunities.

Heredity **116**: 249.

Dlugosch KM, Anderson SR, Braasch J, Cang FA, Gillette HD. 2015a. The devil

is in the details: genetic variation in introduced populations and its

contributions to invasion. *Molecular Ecology* **24**: 2095-2111.

Dlugosch KM, Cang FA, Barker BS, Andonian K, Swope SM, Rieseberg LH.

2015b. Evolution of invasiveness through increased resource use in a vacant

niche. *Nature Plants* **1**.

Dlugosch KM, Parker IM. 2008a. Founding events in species invasions: genetic

variation, adaptive evolution, and the role of multiple introductions. *Molecular*

Ecology **17**: 431-449.

Dlugosch KM, Parker IM. 2008b. Invading populations of an ornamental shrub

show rapid life history evolution despite genetic bottlenecks. *Ecology Letters*

11: 701-709.

Estoup A, Beaumont M, Sennedot F, Moritz C, Cornuet J-M. 2004. Genetic

analysis of complex demographic scenarios: spatially expanding populations

of the cane toad, *Bufo marinus*. *Evolution* **58**: 2021-2036.

Estoup A, Ravigné V, Hufbauer R, Vitalis R, Gautier M, Facon B. 2016. Is There

a Genetic Paradox of Biological Invasion? *Annual Review of Ecology,*

Evolution, and Systematics **47**: 51-72.

Etterson JR, Shaw RG. 2001. Constraint to adaptive evolution in response to global

warming. *Science* **294**: 151-154.

Excoffier L, Ray N. 2008. Surfing during population expansions promotes genetic

revolutions and structuration. *Trends in Ecology & Evolution* **23**: 347-351.

Facon B, Genton BJ, Shykoff J, Jarne P, Estoup A, David P. 2006. A general eco-
evolutionary framework for understanding bioinvasions. *Trends in Ecology &*
Evolution **21**: 130-135.

Felker-Quinn E, Schweitzer JA, Bailey JK. 2013. Meta-analysis reveals evolution
in invasive plant species but little support for Evolution of Increased
Competitive Ability (EICA). *Ecology and Evolution* **3**: 739-751.

Friedman J, Barrett SC. 2009. Wind of change: new insights on the ecology and
evolution of pollination and mating in wind-pollinated plants. *Annals of*
Botany **103**: 1515-1527.

Friedman J, Barrett SC. 2011. Genetic and environmental control of temporal and
size-dependent sex allocation in a wind-pollinated plant. *Evolution* **65**: 2061-
2074.

Gaudeul M, Giraud T, Kiss L, Shykoff JA. 2011. Nuclear and chloroplast
microsatellites show multiple introductions in the worldwide invasion history
of common ragweed, *Ambrosia artemisiifolia*. *PLoS One* **6**: e17658.

Geng Y-P, Pan X-Y, Xu C-Y, Zhang W-J, Li B, Chen J-K, Lu B-R, Song Z-P.
2007. Phenotypic plasticity rather than locally adapted ecotypes allows the
invasive alligator weed to colonize a wide range of habitats. *Biological*
Invasions **9**: 245-256.

Gladieux P, Giraud T, Kiss L, Genton BJ, Jonot O, Shykoff JA. 2010. Distinct
invasion sources of common ragweed (*Ambrosia artemisiifolia*) in Eastern and
Western Europe. *Biological Invasions* **13**: 933-944.

Griffith TM, Watson MA. 2005. Is evolution necessary for range expansion?
Manipulating reproductive timing of a weedy annual transplanted beyond its
range. *The American Naturalist* **167**: 153-164.

Grime JP. 1977. Evidence for the existence of three primary strategies in plants and
its relevance to ecological and evolutionary theory. *The American Naturalist*
111: 1169-1194.

Guisan A, Petitpierre B, Broennimann O, Daehler C, Kueffer C. 2014. Unifying
niche shift studies: insights from biological invasions. *Trends in Ecology &*
Evolution **29**: 260-269.

Guo H, Mazer SJ, Du G. 2010. Geographic variation in primary sex allocation per
flower within and among 12 species of *Pedicularis* (Orobanchaceae):

proportional male investment increases with elevation. *American Journal of Botany* **97**: 1334-1341.

Hahn MA, Rieseberg LH. 2017. Genetic admixture and heterosis may enhance the invasiveness of common ragweed. *Evolutionary Applications* **10**: 241-250.

He W-M, Thelen GC, Ridenour WM, Callaway RM. 2010. Is there a risk to living large? Large size correlates with reduced growth when stressed for knapweed populations. *Biological Invasions* **12**: 3591-3598.

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**: 1965-1978.

Hodgins KA, Rieseberg L. 2011. Genetic differentiation in life-history traits of introduced and native common ragweed (*Ambrosia artemisiifolia*) populations. *Journal of Evolutionary Biology* **24**: 2731-2749.

Huang F, Peng S, Chen B, Liao H, Huang Q, Lin Z, Liu G. 2015. Rapid evolution of dispersal-related traits during range expansion of an invasive vine *Mikania micrantha*. *Oikos* **124**: 1023-1030.

Huey RB, Gilchrist GW, Carlson ML, Berrigan D, Serra Ls. 2000. Rapid evolution of a geographic cline in size in an introduced fly. *Science* **287**: 308-309.

Keller SR, Sowell DR, Neiman M, Wolfe LM, Taylor DR. 2009. Adaptation and colonization history affect the evolution of clines in two introduced species. *New Phytologist* **183**: 678-690.

Keppel G. 1991. *Design and analysis: A researcher's handbook*: Prentice-Hall, Inc.

Klinkhamer PG, De Jong TJ, Metz H. 1997. Sex and size in cosexual plants. *Trends in Ecology & Evolution* **12**: 260-265.

Korpelainen H. 1998. Labile sex expression in plants. *Biological Reviews* **73**: 157-180.

Kuznetsova A, Brockhoff PB, Christensen RHB. 2015. Package 'lmerTest'. *R package version* **2**.

Laaidi M, Laaidi K, Besancenot J-P, Thibaunod M. 2003. Ragweed in France: an invasive plant and its allergenic pollen. *Annals of Allergy, Asthma & Immunology* **91**: 195-201.

Lachmuth S, Durka W, Schurr FM. 2011. Differentiation of reproductive and
competitive ability in the invaded range of *Senecio inaequidens*: the role of
genetic Allee effects, adaptive and nonadaptive evolution. *New Phytologist*
192: 529-541.

Lee CE. 2002. Evolutionary genetics of invasive species. *Trends in Ecology and*
Evolution **17**: 386-391.

Leiblein-Wild MC, Tackenberg O. 2014. Phenotypic variation of 38 European
Ambrosia artemisiifolia populations measured in a common garden
experiment. *Biological Invasions* **16**: 2003-2015.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**: 1754-1760.

Li X-M, She D-Y, Zhang D-Y, Liao W-J. 2014. Life history trait differentiation and
local adaptation in invasive populations of *Ambrosia artemisiifolia* in China.
Oecologia **177**: 669-677.

Lloyd D. 1984. Gender allocations in outcrossing cosexual plants.

Lloyd DG, Bawa KS. 1984. Modification of the gender of seed plants in varying
conditions. *Evolutionary Biology* **17**: 255-338.

Maron JL, Vilà M, Bommarco R, Elmendorf S, Beardsley P. 2004. Rapid
evolution of an invasive plant. *Ecological Monographs* **74**: 261-280.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A,
Garimella K, Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome
Analysis Toolkit: a MapReduce framework for analyzing next-generation
DNA sequencing data. *Genome Research* **20**: 1297-1303.

Miller GA, Chapman JP. 2001. Misunderstanding analysis of covariance. *Journal of*
Abnormal Psychology **110**: 40.

Moran EV, Alexander JM. 2014. Evolutionary responses to global change: lessons
from invasive species. *Ecology Letters* **17**: 637-649.

Oduor AM, Leimu R, Kleunen M. 2016. Invasive plant species are locally adapted
just as frequently and at least as strongly as native plant species. *Journal of*
Ecology **104**: 957-968.

Palmer B, McFadyen RC 2012. *Ambrosia artemisiifolia* L. - annual ragweed.
Biological control of weeds in Australia. Collingwood: CSIRO, 52-59.

Paquin V, Aarssen LW. 2004. Allometric gender allocation in *Ambrosia*
artemisiifolia (Asteraceae) has adaptive plasticity. *American Journal of*
Botany **91**: 430-438.

Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. 2013. On the accumulation of
deleterious mutations during range expansions. *Molecular Ecology* **22**: 5972-
5982.

Peischl S, Excoffier L. 2015. Expansion load: recessive mutations and the role of
standing genetic variation. *Molecular Ecology* **24**: 2084-2094.

Phillips BL, Brown GP, Webb JK, Shine R. 2006. Invasion and the evolution of
speed in toads. *Nature* **439**: 803-803.

Prentis PJ, Wilson JR, Dormontt EE, Richardson DM, Lowe AJ. 2008. Adaptive
evolution in invasive species. *Trends in Plant Sciences* **13**: 288-294.

Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure
using multilocus genotype data. *Genetics* **155**: 945-959.

R Core Team 2017. R: a language and environment for statistical computing.

Rius M, Darling JA. 2014. How important is intraspecific genetic admixture to the
success of colonising populations? *Trends in Ecology & Evolution* **29**: 233-
242.

Sakai A, Sakai S. 2003. Size-dependent ESS sex allocation in wind-pollinated
cosexual plants: fecundity vs. stature effects. *Journal of Theoretical Biology*
222: 283-295.

Savolainen O, Lascoux M, Merila J. 2013. Ecological genomics of local adaptation.
Nature Review Genetics **14**: 807-820.

Sax DF, Brown JH. 2000. The paradox of invasion. *Global Ecology and*
Biogeography **9**: 363-371.

Scalone R, Lemke A, Štefanić E, Kolseth A-K, Rašić S, Andersson L. 2016.
Phenological variation in *Ambrosia artemisiifolia* L. facilitates near future
establishment at northern latitudes. *PLoS One* **11**: e0166510.

Scheiner SM 2001. Multiple response variables and multi-species interactions.
Design and analysis of ecological experiments. New York: Chapman & Hall,
99-133.

Schoener TW. 1968. The Anolis lizards of Bimini: resource partitioning in a complex
fauna. *Ecology* **49**: 704-726.

Smith M, Cecchi L, Skjøth C, Karrer G, Šikoparija B. 2013. Common ragweed: a
threat to environmental health in Europe. *Environment International* **61**: 115-
126.

Sun Y, Brönnimann O, Roderick GK, Poltavsky A, Lommen ST, Müller-Schärer
H. 2017. Climatic suitability ranking of biological control candidates: a
biogeographic approach for ragweed management in Europe. *Ecosphere* **8**:
e01731.

Taramarcaz P, Lambelet C, Clot B, Keimer C, Hauser C. 2005. Ragweed
(*Ambrosia*) progression and its health risks: will Switzerland resist this
invasion? *Swiss Medical Weekly* **135**: 538-548.

Turner KG, Hufbauer RA, Rieseberg LH. 2014. Rapid evolution of an invasive
weed. *New Phytologist* **202**: 309-321.

van Boheemen LA, Lombaert E, Nurkowski KA, Gauffre B, Rieseberg LH,
Hodgins KA. 2017. Multiple introductions, admixture and bridgehead
invasion characterize the introduction history of *Ambrosia artemisiifolia* in
Europe and Australia. *Molecular Ecology* **26**: 5421-5434.

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-
Moonshine A, Jordan T, Shakir K, Roazen D, Thibault J. 2013. From
FastQ data to high-confidence variant calls: the genome analysis toolkit best
practices pipeline. *Current protocols in bioinformatics* **43**: 11.10. 11-11.10.
33.

Willemsen RW. 1975. Dormancy and germination of common ragweed seeds in the
field. *American Journal of Botany* **62**: 639-643.

Zhang D-Y 2006. Evolutionarily stable reproductive investment and sex allocation in
plants. In: Harder LD, Barrett RD eds. *Ecology and evolution of flowers*.
Oxford: Oxford University Press, 41-60.

Zhang YY, Zhang DY, Barrett SC. 2010. Genetic uniformity characterizes the
invasive spread of water hyacinth (*Eichhornia crassipes*), a clonal aquatic
plant. *Molecular Ecology* **19**: 1774-1786.

Ziska L, Knowlton K, Rogers C, Dalan D, Tierney N, Elder MA, Filley W,
Shropshire J, Ford LB, Hedberg C. 2011. Recent warming by latitude

associated with increased length of ragweed pollen season in central North

America. *Proceedings of the National Academy of Sciences* **108**: 4248-4251.

THE FOLLOWING SUPPORTING INFORMATION IS AVAILABLE FOR

THIS ARTICLE:

Fig. S1 Sampling locations

Fig. S2 Principle component analysis of climatic and geographic variables

Fig. S3 Principle component analysis of traits

Fig. S4 Graphical representation of performed trait comparisons at range edges

Table S1 List of climatic and geographic variables

Table S2 Correlation and principle component loadings of climatic and geographic

variables

Table S3 Trait correlation and principle component loadings

Table S4 MANCOVA testing the effect of range treatment and latitude on selected

traits

Table S5 Trait-latitude associations within ranges and range differences

Table S6 Flowering onset, dichogamy and floral sex allocation associations with

height within ranges and range differences

Table S7 Range differences of heterozygosity-biomass associations

Methods S1 Detailed methods for common garden set-up and data collection

Notes S1 Australian trait divergence

FIGURE LEGENDS

Figure 1. Climatic niche models of the native North American population (A) and

introduced Australasian (B) and Eurasian (C) populations. Each panel shows the

climate space occupied in the respective range, with a PCA variable corresponding to

temperature on the x-axis and a PCA variable corresponding to precipitation on the y-

axis. Color indicates occurrence probability in a given climate (red: high occurrence,

green: medium occurrence, blue: low occurrence, grey: no occurrence). Open circles

plot climates of the North American source localities. Closed circles plot the climates

of the respective introduced range. On panels B and C, the dashed line encloses the

climates shared by both the native North American and respective introduced range.

Figure 2. Traits responses (population means) to absolute latitude in the native North
American (blue triangles), introduced European (green squares) and Australian (red
circles) ranges, with model predictions and 95% shaded confidence intervals from
step-wise reduced models (Table 2).

Figure 3. Flowering time, dichogamy and floral sex allocation responses to maximum
plant height (individual values) in the native North American (NA, blue triangles),
introduced European (EU, green squares) and Australian (AU, red circles) ranges,
with model predictions and 95% shaded confidence intervals from step-wise reduced
models (Table 4). Differences in slopes are indicated by letters and are significantly
different from zero unless otherwise indicated (ns)(Table S5, supporting information).

Figure 4. Total biomass and weight per seed response to heterozygosity (population
means) in the native North American (blue triangles), introduced European (green
squares) and Australian (red circles) ranges, with model predictions and 95% shaded
confidence intervals from step-wise reduced models (Table 5). Slopes of predicted
lines are significantly different from zero, unless otherwise indicated as (ns).

Fig. 1

Fig. 2

Fig. 3

Fig. 4