Title: Temporal pattern separation in hippocampal neurons through multiplexed neural codes

Short Title: Neural codes supporting pattern separation in hippocampus

Authors: Antoine D. Madar[1,2], Laura A. Ewell[1,3], Mathew V. Jones[1]

Authors affiliations
[1]Department of Neuroscience, University of Wisconsin-Madison, WI 53705, USA
[2]Department of Neurobiology, Neuroscience Institute, University of Chicago, IL 60637, USA
[3]Institute of Experimental Epileptology and Cognition Research, University of Bonn, Germany

Corresponding authors
Antoine D. Madar (madar@wisc.edu, madar@uchicago.edu)

Author contributions
Conceptualization: MVJ, ADM. Data curation: ADM. Formal analysis: ADM, MVJ. Funding acquisition: MVJ, ADM. Investigation/data collection: ADM, LAE. Methodology: MVJ, ADM, LAE. Project administration: MVJ, ADM. Resources: N/A. Software: ADM, MVJ. Supervision: MVJ. Validation: MVJ, LAE, ADM. Visualization: ADM, MVJ. Writing – original draft: ADM. Writing – review & editing: MVJ, ADM, LAE.

Acknowledgements
We thank Jesse Pfammatter and Kyrie Sellnow for their help in collecting the data.

Funding
This work was supported by the University of Wisconsin Institute for Clinical and Translational Research (M.V.J.; NIH/NCATS UL1TR000427) and Lily's Fund for Epilepsy Research (A.D.M.; 2015 fellow).

The authors declare no competing interests in relation to the present work.

Abstract:

Pattern separation is a central concept in current theories of episodic memory: this computation is thought to support our ability to avoid confusion between similar memories by transforming similar cortical input patterns of neural activity into dissimilar output patterns before their long-term storage in the hippocampus. Because there are many ways one can define patterns of neuronal activity and the similarity between them, pattern separation could in theory be achieved through multiple coding strategies. Using our recently developed assay that evaluates pattern separation in isolated tissue by controlling and recording the input and output spike trains of single hippocampal neurons, we explored neural codes through which pattern separation is performed by systematic testing of different similarity metrics and various time resolutions. We discovered that granule cells, the projection neurons of the dentate gyrus, can exhibit both pattern separation and its opposite computation, pattern convergence, depending on the neural code considered and the statistical structure of the input patterns. Pattern separation is favored when inputs are highly similar, and is achieved through spike time reorganization at short time scales (< 100 ms) as well as through variations in firing rate and burstiness at longer time scales. These multiplexed forms of pattern separation are network phenomena, notably controlled by GABAergic inhibition, that involve many celltypes with input-output transformations that participate in pattern separation to different extent and with complementary neural codes: a rate code for dentate fast-spiking interneurons, a burstiness code for hilar mossy cells and a synchrony code at long time scales for CA3 pyramidal cells. Therefore, the isolated hippocampal circuit itself is capable of performing temporal pattern separation using multiplexed coding strategies that might be essential to optimally disambiguate multimodal mnemonic representations.

Author Summary:

Pattern separation (the process of disambiguating incoming patterns of neuronal activity) is a central concept in all current theories of episodic memory, as it is hypothesized to support our ability to avoid confusion between similar memories. For the last thirty years, pattern separation has been attributed to the dentate gyrus of the hippocampus, but this has been hard to test experimentally. Moreover, because it is unclear how to define activity patterns in the brain, such a computation could be achieved in many different ways. Here, we demonstrate that pattern separation is performed by hippocampal networks (dentate gyrus and CA3) through a variety of neural codes. By systematically testing different definitions of what it means for spike trains to be similar (using a range of time scales and various standard and innovative metrics that assume different views of the neural code), we assessed how the input-output transformation of multiple hippocampal celltypes relate to pattern separation and found that different celltypes favor complementary coding strategies. This might help storing rich but concise and unambiguous representations of complex events. Finally, we provide the first experimental evidence of the importance of inhibitory signals in mediating pattern separation, and identify through which coding strategies.

Introduction:

Understanding what computations hippocampal neurons perform and how they support episodic memory is a longstanding goal in neuroscience. The hippocampus is notably critical for the discrimination of memories that are similar in content [1]. This cognitive function has long been hypothesized to be supported by a neural process called pattern separation, which is generally thought to be implemented in the dentate gyrus (DG) [2-4]. Pattern separation is

defined as the transformation of a non-simultaneous set of similar input patterns of neuronal

activity into less similar output patterns [5], and is theorized to happen in the hippocampus

before encoding a memory trace in order to avoid confusion between similar memories [6].

Despite a long history of research on the subject, mostly in silico [7], it is still unclear how the

activity of single neurons underlies this computation.

Generally speaking, single-neuron computations have been investigated experimentally

using two broad strategies: 1) by determining how the spiking of individual neurons, recorded in

vivo, is tuned to different parameters of the behavior or environment [8] or 2) by relating the

neuronal infra or suprathreshold membrane potential responses, generally recorded in vitro, to

their synaptic inputs.

The first strategy has led to the discovery that the principal cells of the hippocampus fire

preferentially in specific locations of an environment called place fields [9]. Upon sufficient

modification of the environment, the place field(s) of a given neuron remap by relocating or

changing their firing rate (in extreme cases, disappearing) [10, 11]. These different forms of

remapping have often been used as a proxy for pattern separation, yielding conflicting

interpretations on the role of DG and CA3 [12-14]. However, neurons remapping between two

similar environments is not sufficient evidence that those neurons participate in pattern

separation because place field remapping is assessed without knowledge of the direct synaptic

inputs: it is possible that separated representations are inherited from upstream networks. Thus,

despite extensive recordings of large populations of several hippocampal celltypes during

behavior [11, 14], it remains unknown how each celltype participates in pattern separation.

In contrast to the first strategy, the second one explicitly takes into account the synaptic

inputs of recorded neurons. Such an approach has been used to study dendritic integration [15],

short [16, 17] or long-term plasticity [18, 19] and the relationship between excitatory inputs and

the output spiking probability [19-23], but most of these studies were limited to simple input

activity patterns (i.e. rhythmic series of spikes or bursts of spikes) and simple output patterns (i.e.

a single spike). Some have used more complex sequences of spatially distributed synaptic inputs

[24-26], but the resulting input patterns were still spikes regularly spaced in time. Yet, neuronal

spike trains recorded in vivo are not so regular but instead approximately follow a Poisson

distribution [27] or a more bursty distribution [28, 29], and what neurons integrate is thus a

barrage of irregular synaptic inputs [30, 31]. To infer the single-neuron suprathreshold input-

output relationship in more naturalistic regimes, some have thus recorded the spiking response to

intracellular injections of current waveforms resulting from either white noise [32] or from the

simulated random discharges of a large number of presynaptic neurons [27, 30]. But this only

gives insight on the operations performed by a neuron isolated from its network. To understand

the role of the network, rare studies have directly stimulated afferents with naturalistic spike

trains to characterize short-term [33-35] or long-term [35, 36] synaptic dynamics of hippocampal

neurons. An even smaller number of studies have aimed at characterizing the suprathreshold

input-output function of a neuron in response to external Poisson spike trains [37, 38], and such

experiments have determined that this function was different when evaluated with simpler input

patterns [39], confirming the necessity to assess neuronal computations with complex,

naturalistic input patterns.

Surprisingly, because none of the above studies have quantified and systematically

varied the similarity between input patterns, none have directly addressed how the input-output

transformation in single hippocampal neurons relates to pattern separation. To answer this

question, we developed a new paradigm [40], in mouse brain slices, where afferent axons of the

DG are stimulated with complex input spike trains of varying similarity while the output spike

trains from a single neuron are recorded. Thus, in contrast to in vivo studies, we considered input

and output temporal patterns instead of spatial ones. With this paradigm, we recently

demonstrated that the suprathreshold responses of the principal neurons of the DG and CA3 were

strongly decorrelated compared to their inputs, whereas DG interneurons exhibited less temporal

decorrelation [40].

Importantly, our previous investigation only considered one type of neural code (binwise

synchrony, as measured by the Pearson's correlation coefficient), even though temporal pattern

separation could, in principle, be achieved in a variety of ways depending on the features

considered relevant in a spike train. For example, if the only feature carrying information in a

spike train was its firing rate, a pattern separator would convert a series of input spike trains with

similar rates into a series of output spike trains with dissimilar rates. Alternatively, the firing rate

could be irrelevant and two output trains with the same number of spikes could be considered

separated if their spiketimes are desynchronized. Pattern separation is thus a group of potential

computations, each computation corresponding to a specific neural code, or, in other words, to a

specific definition of "similarity". By testing a wide range of similarity metrics and time scales,

the present work aims to determine through which neural codes temporal pattern separation is

performed in the hippocampus.

Results:

Temporal pattern separation via decorrelation, orthogonalization and scaling

In Madar et al. (2018) [40], we demonstrated for the first time that single GCs exhibit

high levels of pattern decorrelation thanks to a novel pattern separation assay in acute brain

slices. Our general paradigm has three steps (**Fig 1, Methods - Experiments**): 1) ensembles of

stimulus patterns (simulating trains of action potentials) are generated, with known degrees of

similarity to each other. These sets of input spike trains are then fed into the DG by stimulating

the lateral perforant path. 2) The simultaneous response of a single neuron is recorded in whole-

cell current-clamp. 3) The similarity between the output spike trains is compared to the similarity

between the input patterns, revealing the degree of separation or convergence.

We first reanalyzed a dataset, presented in Madar et al. (2018) [40], of GC recordings

performed in response to input sets constituted of 10 Hz Poisson trains (**Fig 1B**). A pairwise

similarity analysis (**Fig 1E**), with the Pearson's correlation coefficient R as a measure of

similarity, confirmed our finding that the output spike trains of GCs are significantly less

correlated than their inputs at both low and high input correlations (**Fig 1F**). As detailed in

Madar et al. (2018) [40], even the repetition of the same input train leads to decorrelated output

trains, due to the probabilistic nature of synaptic transmission.

We used R as a similarity measure between spike trains because it is easy to implement

and is commonly used to quantify the similarity between neural activity patterns, both in

computational [41] and experimental studies [12, 13, 42, 43]. However, the original Hebb-Marr

framework theorized pattern separation as the orthogonalization of the input patterns [5, 44]. As

a result, the terms "decorrelation" (corresponding to output patterns, viewed as variables, with a

lower Pearson's correlation coefficient than their inputs) and "orthogonalization" (corresponding

to output patterns, viewed as vectors, that are closer to a right angle than inputs) are often

conflated in the literature, even though they are not mathematically equivalent and have a

nonlinear relationship to each other (**Fig 2A-C, S1**. See **Methods – Similarity metrics** and

Supplements - 1). For instance, pairs of spike trains can be uncorrelated ($R = 0$) without being

orthogonal, or can be orthogonal without being uncorrelated (**Fig 2A-C** and **S1**). To determine

whether output spike trains of GCs are truly orthogonalized, we explicitly considered spike trains

as vectors of spike-counts and computed the normalized dot product (NDP, i.e. the cosine of the

angle between two vectors) between pairs of spike trains to assess their similarity (**Fig 2A, C**.

See **Methods – Similarity metrics** and **Supplements - 1**). For every recording set, NDP_{output}

was lower than NDP_{input}, indicating that the angle between output spike trains was closer to a

right angle (i.e., closer to orthogonal) than their inputs (**Fig 2D-E**).

Vectors can differ by their angle, but also by their norm. In other words, even if neurons

fire in the same time bins (relative to the start of each sweep), the number of spikes per bin can

be different, as quantified by the ratio between their norms (scaling factor, SF) (**Fig 2A, C.** See

Methods – Similarity metrics and **Supplements - 1**). Our results show that for input patterns

highly similar in terms of SF, SF_{output} is slightly but significantly lower than SF_{input} (**Fig 2F**). This

indicates that variations in the binwise firing rate of single GCs in response to similar inputs is a

potential, but weak, mechanism of pattern separation at the 10 ms time scale.

As a whole, these results confirm that input spike trains are decorrelated in the DG at the

level of single GCs, and demonstrate that, at the 10 ms time scale, GCs exhibit temporal pattern

separation mediated by high levels of orthogonalization and weak levels of scaling.

Pattern separation of spike trains occurs through multiplexed neural codes

Trains of action potentials can be described in many different ways: their overall firing

rate, the timing of their spikes or the number of bursts of spikes, among other statistics (from

now on referred to as *spike train features*). Measuring the similarity between two spike trains is

not a trivial problem, if only because it is unclear what spike train features are relevant to the

brain and what counts as different or similar. In principle, pattern separation could be achieved

by the variation of any spike train feature (e.g. if the relevant feature is the total number of spikes

in two second bins, pattern separation will be achieved if this number varies more in the set of

output spike trains than in the input set). In other words, many different forms of pattern

separation could be performed depending on the neural code considered. In our analyses, the

neural code (i.e. the set of assumptions on which spike train features are considered relevant and

what definition of similarity is used) is determined by the choice of similarity metric and time

resolution (see **Supplements – 1**). In order to better characterize which neural code(s) are used to

perform temporal pattern separation in the hippocampus, we thus tested multiple metrics and

time scales. This approach also allowed us to ask whether several coding strategies could be

multiplexed (i.e. simultaneously relevant over different time scales) to achieve pattern

separation.

The three measures of similarity we have used above (R, NDP and SF) all carry different

assumptions about the neural code (**Table 1**): R and NDP are mostly, but differently, sensitive to

the binwise synchrony whereas SF evaluates variations in spike number (**Fig 2A-C** and see

Supplements - 1). On the other hand, these metrics resemble each other in that they require

binning spike trains in time windows of a prespecified duration (τ_w). The time scales that are

meaningful for the brain are uncertain so we assessed the separation of spike trains for different

τ_w. Indeed, different durations of τ_w assume a different window to read the information contained

in a spike train. Our analysis shows that pattern separation, measured through R or NDP, is more

pronounced at short time scales (e.g. 5 ms) than at longer ones (e.g. 100 ms) (**Fig 3A-B**). In

contrast, in the case of high input similarity, pattern separation through scaling is rather weak at

short time scales but gets stronger at longer ones (**Fig 3C**). This demonstrates that multiplexed

coding allows temporal pattern separation to be carried out by the DG for a large range of time

scales.

In addition, because a long stream of research suggests that spike trains can carry

information directly through the timing of individual spikes [45-47], we also assessed the

similarity between spike trains using SPIKE, a binless metric purely based on spike times [48],

thus assuming a neural code completely different from the binned metrics above (**Table 1,**

Methods – Similarity metrics and **Supplements**). Our results show that input spike trains with

similar spike times relative to their sweep start (defined here as spike trains with a high degree of

synchrony, see **Methods – Similarity metrics**), are transformed into significantly less

synchronous outputs, thereby demonstrating that temporal pattern separation in single GCs can

occur through spike timing modifications (**Fig 3D**). This is consistent with the high levels of

decorrelation and orthogonalization at very short time scales (**Fig 3A-B**) and confirms that

temporal pattern separation can be achieved through multiple coding strategies.

Table 1. Similarity metrics assumption on the neural code

	R	NDP	SF	SPIKE
Binned (spike count code)	X	X	X	
Binless (spike time code)				X
Preserved time relationship between time points or bins				X
"raw" (not baseline-subtracted) spike count vectors		X	X	
Assumption of linear relationship between spike trains	X	X	X	
Assumption of Euclidean space	X	X	X	
Excludes common periods of silence		X		
Mostly sensitive to binwise synchrony	X	X		
Sensitive to spike time synchrony				X
Sensitive to differences in firing rate or burstiness between spike trains			X	X

Single GC computations depend on the statistical structure of their input patterns

Results in **Fig 2F** and **3C** show that the similarity of GCs output spike trains in terms of SF is almost flat, suggesting it might be independent of the input similarity, hence leading to significant pattern convergence in certain conditions of low input similarity and short time scales. However, those experiments were based on responses to ~10 Hz Poisson input spike trains that did not differ much in terms of SF. To further characterize single GC computations, we designed two new input sets spanning a wider range of SF values (**Fig 4A-B**). From its definition, it was apparent that, unlike R and NDP, SF is very sensitive to both differences of overall firing rate and differences in burstiness between spike trains (**Fig S2**). Input set A was thus made of highly correlated Poisson spike trains with different firing rates (7-31.5 Hz), and input set B was made of uncorrelated spike trains with constant firing rate but different levels of

burstiness (see **Methods - Experiments**). By recording the spiking output of single GCs in

response to these input sets we could thus test how GCs transform input patterns with vastly

different statistical structures, and whether they can still perform pattern separation in those

conditions. Our results indicate that for both input sets GCs can perform weak levels of pattern

separation via scaling for highly similar inputs (**Fig 4C**), consistent with our experiments using

10 Hz Poisson trains (see Fig 2F and 3C). However, although not independent of input similarity,

the similarity of GCs output spike trains (in terms of SF) is comprised in a relatively narrow

band, yielding high levels of pattern convergence for relatively dissimilar input spike trains (**Fig

4C**). In other words, when assuming a neural code based on the norm of spike count vectors,

GCs outputs are maintained relatively similar to each other, even when their corresponding input

patterns are very different. This is true from the millisecond to the second time scale, but pattern

convergence decreases and separation increases with longer time scales (**Fig 4C**), consistent with

previous experiments (see Fig 3C).

Although SF makes mathematical sense when measuring the similarity between vectors,

it is difficult to interpret in terms of basic spike train features, in part because it is sensitive to

both firing rate and burstiness (see **Supplements – 1** for details). We thus aimed to gain a more

intuitive view of how those spike train features vary to result in pattern separation or

convergence. For that, we calculated the mean firing rate of each spike train and designed two

new indicators providing complementary information on the burstiness of individual spike trains:

compactness and occupancy (**Fig 5A**). Briefly, compactness is anticorrelated to the number of

time bins occupied by at least one spike (the less occupied bins, the more compact a spike train

is), whereas occupancy is the average number of spikes per occupied bin (**Methods – Firing

rate and burstiness codes** and **Supplements - 2**).

First, we asked how GCs transform their inputs in terms of firing rate and burstiness. For

all time scales, GCs have generally equal or higher compactness than their inputs but maintain a

constant low range of occupancy (**Fig 5B left**). This combination suggests that, using **Table S1**,

GC responses are mostly a sparser version of their inputs, with a decrease in the number of

spikes per burst when inputs are bursty. Accordingly, the overall firing rate of GC output spike

trains is generally lower than their inputs (**Fig 5B right**), which is consistent with the known

sparsity of GC activity in vivo and fits the view of the DG acting as a filter of cortical activity

[49].

We next asked whether temporal pattern separation is performed through variation of the

firing rate between GCs output spike trains, or through variations of compactness or occupancy,

or a combination of those three. Our results indicate that pattern separation, but also pattern

convergence, can be exhibited for all three types of codes, depending on certain conditions (**Fig**

6). First, separation is favored at longer time scales (for burstiness codes) and high input

similarity (for burstiness and rate codes) (**Fig 6A**). Second, the direction of the computation

(separation or convergence) strongly depends on the statistical structure of the input patterns (**Fig**

6B): 1) In response to Poisson inputs with very similar firing rates (P10Hz), GCs exhibit low but

significant levels of pattern separation through small variations of compactness, occupancy and

firing rate, which become more relevant with larger time scales. Pattern separation measured

through scaling (see Fig 2F and 3C) was thus due to the variability in all three spike train

features combined. 2) In response to Poisson inputs with varying firing rates (PΔFR, input set A)

GCs switch, as the time scale is increased, from pattern convergence to pattern separation

through compactness variations, while increasing their levels of convergence in terms of

occupancy. Accordingly, their output spike trains have more similar firing rates than their inputs

(this is because GCs maintain a narrow range of firing rates regardless of the rate of their inputs, as shown in Fig 5B). 3) In response to inputs with a constant firing rate but varying levels of burstiness (B 10.5 Hz, input set B), GCs show low but significant pattern convergence in terms of compactness and occupancy. Pattern convergence in terms of occupancy progressively decreases as the time scale increases. This translates into significant pattern separation through variations of the firing rate (note that, because spike trains were 2 s long, pattern separation via FR variations is equivalent to pattern separation via occupancy variations in a single 2 s time scale). Interestingly, this analysis demonstrates that GCs can exhibit both separation and convergence simultaneously through different neural codes: for instance, at the 250 ms time scale, the compactness of GCs output spike trains vary more than the compactness in input set A, whereas GCs output spike trains occupancy vary less (**Fig 6B, Table 2**).

Table 2 related to Fig 6B

One sample t-test (difference from 0)

p value < 0.05 in red.

Input set	$\tau_w = 10$ ms	$\tau_w = 100$ ms	$\tau_w = 250$ ms	Firing rate
	Compactness			
P10Hz	< 0.0001	< 0.0001	< 0.0001	
PΔFR	0.0004	0.1576	0.0023	
B10.5Hz	0.0135	0.0213	0.0298	
	Occupancy			**Firing rate**
P10Hz	< 0.0001	0.0004	< 0.0001	< 0.0001
PΔFR	0.0095	< 0.0001	< 0.0001	0.00016
B10.5Hz	0.0014	0.0003	0.00048	0.0244

Inhibition enforces temporal pattern separation in GCs.

GCs are known to receive strong feedforward and feedback inhibition from a variety of

GABAergic interneurons [50, 51] that impact GCs spiking output [21, 52, 53]. Furthermore,

theoretical work has long suggested a role for interneuron interactions with GCs in mediating

pattern separation [4, 54, 55], but experimental evidence is lacking. To test the influence of fast

inhibitory transmission on the complex computations of GCs characterized above, we recorded

GCs responses to 10 Hz Poisson trains before and after application of a nonsaturating

concentration of Gabazine, a $GABA_A$ receptor antagonist (**Fig 7A**, see **Methods - Experiments**).

Under these conditions, IPSC amplitude was reduced by ~30% (not shown), which led to a slight

but visible increase in firing rate, in part due to a higher propensity to fire short bursts of action

potentials riding a single EPSP (**Fig 7A-B**). This pharmacological manipulation led to lower

levels of temporal pattern separation as measured by R, NDP, but also SF (**Fig 7C**). This was

true at all time scales (**Fig 7D**), with the importance of inhibition for pattern separation even

increasing at longer time scales. Interestingly, the impact on pattern separation at larger time

scale was not only for SF, whose relevance increases with larger τ_w (see Fig 3C) but also for R

and NDP, whose relevance normally decreases with larger τ_w (see Fig 3A-B).

The average R_{output} was well correlated to the average firing rate of a recording set (linear

regression: $R^2 = 94\%$, $p < 0.0001$, $n = 14$ data points from 7 GCs), suggesting that the inhibitory

control of sparsity in GCs firing helps temporal pattern separation as measured with R.

Interestingly, in a previous report we did not detect such a strong relationship between R and the

firing rate levels of GCs [40]. Moreover, R is mathematically independent of proportional

increases of pairwise firing rate levels and not very sensitive to non-proportional ones (see

Supplements - 1 and **Fig S1C**). In the present experiment, the correlation between R and the

firing rate is thus likely to arise from physiological reasons that conjointly affect both the

sparsity and the binwise synchrony of output spike trains rather than from the mathematical

assumptions behind R.

In contrast to R and NDP, the mathematical definition of SF makes it sensitive to

pairwise firing rate levels (i.e. SF systematically increases for pairs of spike trains with higher

firing rates. See **Supplements - 1** and **Fig S1C**). As seen before, SF is also designed to be

sensitive to differences in firing rate and burstiness of individual spike trains (e.g. Fig 5 and S2).

Therefore, the effect of gabazine on decreasing pattern separation as measured by SF (**Fig 7C-D**)

could be due: 1) to a global increase in firing rates in all the output spike trains, 2) to variations

of firing rate or burstiness between output spike trains, or 3) to a combination of all of the above.

The direct assessment of pattern separation in terms of firing rate or burstiness indicates that

inhibition actually does not affect temporal pattern separation through rate or compactness codes

and favors pattern convergence through occupancy codes (**Fig 7E**). We can thus conclude that

the effect of inhibition on pattern separation via scaling (SF) is mostly due to the global firing

rate increase under gabazine.

Overall, we provide here the first experimental evidence that fast inhibitory transmission

in the DG controls levels of temporal pattern separation and convergence through different

multiplexed neural codes.

Different hippocampal celltypes support pattern separation through different neural codes

GCs are embedded in a network of multiple celltypes all participating in generating the

output of the DG. For instance, computational models of the DG have suggested that different

populations of inhibitory and excitatory interneurons could modulate certain forms of pattern

separation [43, 54]. To develop a comprehensive understanding of the computations performed

by the DG, and hippocampal networks in general, it is important to characterize the input-output

transformation of a diverse set of celltypes. We recorded from dentate fast-spiking GABAergic

interneurons (FSs), glutamatergic hilar mossy cells (HMCs) and CA3 pyramidal cells (PCs) in

response to 10 Hz (for GC, FS, HMC) or 30 Hz (for PC and GC) Poisson input trains with

varying levels of correlation (**Fig 8A** and **Methods – Experiments**).

Previously, we found that temporal pattern separation, measured using R, was not

specific to GCs but that GCs exhibit the highest levels of decorrelation among tested dentate

neurons, and that CA3 PCs express even more temporal decorrelation than GCs [40]. Here we

ask whether this is still the case when considering other neural codes based on different

similarity metrics. A pairwise similarity analysis and a comparison of the pattern separation

function (the linear regression model describing S_{output} as a function of S_{input}, with S representing

a given similarity metric) across celltypes shows that pattern orthogonalization (NDP) and

scaling (SF) levels significantly differ at most time scales (**Fig 8B-D, Table 3**).

For NDP, we notice a progression from FSs to HMCs to GCs to CA3 PCs, with GCs

performing the highest levels of separation among DG neurons, and CA3 PCs surpassing GCs,

especially with their ability to orthogonalize inputs already relatively dissimilar (**Fig 8B**). The

orthogonalization function of GCs, HMCs and CA3 PCs diverge as the time scale is increased

(**Fig 8C**).

For SF, scaling functions do not depend much on the input similarity, except for CA3

PCs. Mostly based on the scaling functions intercept, we notice two groups (**Fig 8D**): FSs and

GCs exhibit low levels of pattern separation through scaling, in contrast to HMCs and CA3 PCs

(and their GC controls) that produce spike trains with lower similarities in terms of SF. Note that

among celltypes tested with 10 Hz Poisson trains, HMCs are the best at pattern separation

through scaling, not GCs.

Table 3 related to Fig 8C-D

Post-hoc t-tests p-values with correction for multiple comparisons when necessary (Tukey-

Kramer) comparing slope or intercept estimates (from ANCOVA) between celltypes. Note that

CA3+gzn and GC+gzn are only compared with each other. P-values < 0.05 in red.

Compared Celltypes		$\tau_w = 10$ ms	$\tau_w = 50$ ms	$\tau_w = 100$ ms NDP	$\tau_w = 250$ ms	$\tau_w = 1000$ ms
GC	FS	slope: 0.3197, intercept: 0.0205	slope: 0.2177, intercept: 0.0028	slope: 0.7884, intercept: 0.2455	slope: 0.6601, intercept: 0.3798	slope: 0.8479, intercept:0.8847
GC	HMC	slope: 0.0038, intercept: <0.0001	slope: 0.0013, intercept: 0.0009	slope: 0.0140, intercept: 0.0177	slope: 0.0758, intercept: 0.0818	slope: <0.0001, intercept: <0.0001
FS	HMC	slope: 0.0069, intercept: 0.9758	slope: 0.8737, intercept: 0.4558	slope: 0.6957, intercept: 0.9756	slope: 0.9990, intercept: 0.8882	slope: 0.0001, intercept: < 0.0001
GC+gzn	CA3+gzn	slope: 0.8228, intercept: 0.0110	slope: 0.0920, intercept: 0.0084	slope: 0.0004 , intercept: <0.0001	slope: <0.0001, intercept: <0.0001	slope: <0.0001, intercept: <0.0001
Compared Celltypes				**SF**		
GC	FS	slope: 0.7487, intercept: 0.6242	slope: 0.7583, intercept: 0.6209	slope: 0.9279, intercept:0.9838	slope: 0.7575, intercept: 0.9719	slope: 0.5867 , intercept: 0.5271
GC	HMC	slope: 0.4701, intercept: 0.1568	slope: 0.1848, intercept: 0.0138	slope: 0.0749, intercept: 0.0069	slope: 0.0028, intercept: <0.0001	slope: 0.970, intercept: 0.0052
FS	HMC	slope: 0.3924, intercept: 0.1585	slope: 0.2396, intercept: 0.0477	slope: 0.6326, intercept: 0.2795	slope: 0.8893, intercept: 0.5867	slope: 0.3993, intercept: 0.2901
GC+gzn	CA3+gzn	slope: 0.1092, intercept: 0.1247	slope: 0.0969, intercept: 0.1231	slope: 0.1436, intercept: 0.1573	slope: 0.2810, intercept: 0.3182	slope: 0.1605, intercept: 0.1942

In contrast to GCs, HMCs and FSs tend to fire bursts of spikes (although the

characteristics of bursts differ between the two celltypes), which affects, at least partially, their

levels of pattern decorrelation [40]. An analysis of the distribution of inter-spike-intervals in all

recorded celltypes confirmed that HMCs and FSs are bursty (as they diverge from a Poisson

process) but not to the same extent (**Fig 9A,** and see **Methods - Firing rate and burstiness**

codes). It also revealed that, although neither CA3 PCs nor their GC controls (recorded under

gabazine and 30 Hz inputs) displayed bursts elicited by a single input pulse [40], they can be

considered bursty in the sense that their output spikes tend to be clustered together in time in a

nonrandom fashion (**Fig 9A**). We thus asked whether these celltypes with diverse bursty

behaviors could perform pattern separation through compactness, occupancy or firing rate codes.

Our analysis shows indeed that these features vary across the output spike trains of hippocampal

neurons, but in different ways for each celltype (**Fig 9B, Table 4**): for example, at long time

scales FSs and HMCs achieve pattern separation mostly through variations of occupancy and

firing rate, whereas CA3 PCs achieve pattern separation mostly through variations of

compactness.

Overall, we demonstrate here that not all hippocampal celltypes exhibit the same

computations, but that they all can perform temporal pattern separation to different degrees

through different neural codes and over different time scales. This suggests that different types of

neurons can serve different, complementary, computations that allow the isolated hippocampal

network to perform temporal pattern separation using multiplexed coding strategies.

Table 4 related to Fig 9B

Posthoc t-test after Kruskal-Wallis one-way ANOVA, with Tukey-Kramer correction for

multiple comparisons. CA3+gzn and GC+gzn are only compared with each other. P values <

0.05 in red.

Compared Celltypes		$\tau_w = 10$ ms	$\tau_w = 100$ ms	$\tau_w = 250$ ms	
		Compactness			
GC	FS	<0.0001	0.4368	0.0339	
GC	HMC	0.0064	0.9214	0.7015	
FS	HMC	0.6255	0.437	0.0326	
GC+gzn	CA3+gzn	0.0728	0.0112	0.3864	
		Occupancy			**FR**
GC	FS	< 0.0001	< 0.0001	< 0.0001	<0.0001
GC	HMC	0.0044	< 0.0001	0.1911	0.067
FS	HMC	0.6787	0.7675	0.0467	0.1138
GC+gzn	CA3+gzn	0.1732	0.8045	0.4212	0.0592

Discussion

In summary, we report that the hippocampus can perform temporal pattern separation or

convergence at the level of single neurons through different multiplexed neural codes. Pattern

separation can be achieved by varying many different spike train features in the ensemble of

output spike trains, like the spike times, the firing rate, the number of bursts or the number of

spikes in a burst. We found that GCs, the output neurons of the DG, mostly use a

"desynchronization" strategy leading to an orthogonalization of their output spike trains,

especially at short time scales. When considering a neural code focused on firing rates or

burstiness, GC output varies little, leading to pattern convergence or separation depending on the

degree of similarity of the inputs. Other hippocampal neurons perform different computations

through complementary combinations of various neural codes (mostly rate for FSs, burstiness for

HMCs and synchrony at long time scales for CA3 PCs), and their network interactions constrain

the ability of the hippocampus to perform temporal pattern separation, notably through inhibition

of GC activity. Given that, during behavior, hippocampal neurons have a mixed selectivity to

different aspects of the environment and cognitive task contingencies [56-58], the use of

multiplexed codes is likely critical to encode rich representations efficiently [59]. Therefore, the

multiplexed forms of pattern separation we have characterized here might be essential to

optimally disambiguate between similar multidimensional memories.

The hippocampus performs multiplexed forms of pattern separation

Overall, our study highlights that measuring neuronal computations implies a certain

vision of the neural code, and that brain networks or single neurons perform many different

computations depending on the assumed coding scheme.

The "neural code" is a pervasive but vague concept that refers to the set of features of

neural activity that convey information, for example to represent our experiences and memories.

The first difficulty in breaking this code comes from the infinitely large number of potential

features and combinations of features that could be relevant in the brain. Since the early work of

Adrian [60], the most common view is that neurons transmit information about their stimulus

through their firing rate, i.e. the number of output spikes averaged over a certain time window.

This rate code is often contrasted to time codes, but this is a false dichotomy [45]: there is an

infinite number of possible rate codes, as there is an infinite number of time windows over which

the number of spikes could be integrated (are spikes counted by reader neurons over

milliseconds, seconds or minutes?). In addition to spike counts, information can also be carried

by the timing of single spikes [45, 46, 61], bursts [62, 63] or even more complex spike patterns

[61, 64-66]. Even when focusing on time or burstiness codes, many subfeatures could be

considered words in the elusive language of the brain: for example, the information could be

encoded in the specific spike times [67] or the interspike intervals [64, 68], the relative or exact

latency or synchrony with regard to other spike trains [67] or the relative timing of individual

spikes with regard to a reference network oscillation [69], the timing or frequency of bursts [62,

63], the spike times within a burst or the spike counts per burst [63, 70].

Pattern separation has long been thought to cover several potential computations

depending on the neural code considered [5]. Indeed, past investigations did not always define

brain activity patterns the same way, nor did they choose the same similarity metrics and time

resolution, thus assuming different forms of pattern separation. When the hypothesis of pattern

separation was first formulated, activity patterns were defined as binary vectors representing

populations of ON or OFF neurons, without considering a time dimension [4, 6, 44]. As a result,

modeling work that followed often excluded the dynamics of neural activity, and used the

number of common active neurons between two population patterns as their measure of

similarity [43, 54, 71]. In addition to this purely spatial population code, others have defined

activity patterns as maps of firing rates averaged over minutes, or as population matrices of those

maps [12, 14], measuring pattern separation using R [72] or NDP [73] as their similarity metric.

Only two modeling studies have investigated pattern separation by directly considering spike

trains at a time scale less than a minute: one considered an input pattern as a static population of

ON/OFF neurons, an output pattern as a population vector of firing rates averaged over 500 ms

and the similarity between two output patterns was based on the sum of firing rate ratios [71]; the

second considered an input pattern as an ensemble of 30 ms spike trains (0 or 1 spike per input

channel, i.e. akin to a population of ON/OFF neurons but with the time of an input spike carrying

some information due to the possibility of temporal summation) and an output pattern as an

ensemble of 200 ms spike trains, using R with a τ_w of 20 ms to measure the similarity between

spatiotemporal patterns [41].

Past computational studies suggest, together, that the DG could perform pattern

separation through different codes, but our investigation is the first to systematically explore and

compare pattern separation levels based on diverse coding strategies. It is also the first focusing

on time codes. Thanks to our slice-based pattern separation assay, we could control input

patterns while recording the output of the DG, thus allowing us to experimentally test pattern

separation in the DG for the first time [5, 40]. We considered activity patterns (for both input and

output) to be two-second long spike trains, and used multiple definitions of spike train similarity

at time scales going from submillisecond to second resolution. We first measured spike count

correlations with R and NDP, two common metrics with different sets of assumptions on the

neural code (e.g. NDP does not consider common periods of silence as correlated; see

Supplements), showing that the DG performs decorrelation and orthogonalization of its outputs

at subsecond time scales at the level of single GCs (**Fig 3A-B**). Because this phenomenon is

stronger at millisecond time scales, and that R and NDP are mostly sensitive to binwise

synchrony, we measured spike train similarity with SPIKE, assuming a synchrony code purely

based on spike times instead of spike counts [48]. This confirmed that GC outputs are separated

through "desynchronization" of their spike times from sweep to sweep (**Fig 3D**). Moreover, we

designed a new simple similarity metric, SF (which complements NDP in the description of the

similarity between two spike count vectors), as well as two complementary measures of spike

train "burstiness" (compactness and occupancy). These revealed that, in addition to

desynchronization strategies, GCs can exhibit pattern separation through firing rate or burstiness

codes, and that depending on the time scale, the neural code considered and the input statistical

structure, different levels of pattern separation or convergence are performed (**Fig 6**).

In conclusion, our work demonstrates that single neuron pattern separation can be

achieved in isolated hippocampal tissue through multiplexed coding, which is a way of

simultaneously representing several features of a stimulus or, in our case, simultaneously

performing different computations, through different temporal scales and spike train features [59,

63, 74]. Our work illustrates the importance of considering multiple neural codes when studying

neuronal computations for three reasons: 1) The fact that conceptually different metrics lead to

converging results bolsters our conclusion that temporal pattern separation occurs within the DG

and CA3 at the level of single neurons; 2) a single neuron can exhibit either pattern convergence

or separation depending on the coding scheme; 3) a given neural code can be more relevant for

pattern separation/convergence in certain celltypes than others.

Our finding that DG can perform pattern convergence in some conditions is particularly

intriguing. If pattern separation is conceptualized to support mnemonic discrimination, then

pattern convergence would support generalization, another elusive mnemonic phenomenon [75].

In the future, experiments correlating mnemonic discrimination and generalization with the

levels of pattern separation or convergence through various potential codes will help pinpoint

which computations are actually used in the brain to support episodic memory.

Network mechanisms supporting hippocampal computations

Our work, in showing that spiking was difficult to elicit in GCs and that their responses

were sparser than their inputs (**Fig 5**) is in line with past research [20, 23]. It also resonates with

the long-standing idea that the DG acts as the gate of the hippocampus, filtering out incoming

cortical activity before it reaches CA3 to prevent overexcitation [49]. Figuring out the

computations of the DG is equivalent to reverse-engineering the transfer function defining the

DG filter. Some have suggested that the DG is a low-pass [22, 76] or a ~10 Hz band-pass filter

[21]. Our results confirm that the output frequency of GCs is constrained in a narrow band (**Fig**

5B) and go one step further by providing the first demonstration that the filtering properties of

the DG allow it to perform different computations including pattern separation in response to

complex naturalistic inputs (**Fig 6**).

The question is then: how are these filtering properties implemented? The physiological

mechanisms underlying DG computations in general are understudied and remain a mystery. In

Madar et al. (2018) [40], we provided a proof-of-principle that the short-term dynamics of

synaptic transmission at the perforant path-GC synapses could explain the levels of temporal

pattern decorrelation in the DG. But GCs also receive feedforward and feedback inhibition and

excitation from multiple interneurons that also interact between each other [51, 77, 78]. Indeed,

our finding that partial block of $GABA_A$-mediated neurotransmission impairs temporal pattern

separation in the DG (**Fig 7**) strongly suggests that the computations we investigated are network

phenomena. Interactions between multiple celltypes are likely involved, and it is thus critical to

characterize the suprathreshold input-output transformation of each celltype to parse out their

computational role. In the present work, we shed light on the computations of dentate FSs (also

known as basket cells providing fast feedforward and feedback perisomatic inhibition to GCs

[79]), HMCs (glutamatergic interneurons providing feedback monosynaptic excitation and

disynaptic inhibition to GCs mostly outside of the lamellar plane [77]), and CA3 PCs, the output

neurons of the hippocampal subnetwork directly downstream of the DG in the trisynaptic circuit,

but which can also provide indirect feedback to the DG via collaterals targeting hilar neurons

[77]. We discovered that, in contrast to GCs (**Fig 8-9**, and see [40]): 1) FSs generally exhibit low

levels of pattern separation through binwise desynchronization (R, NDP) or scaling (SF, likely

due to high firing rates), but high levels of separation through simple variation of local firing

rates from sweep to sweep, especially at the 100 ms time scale and higher; 2) HMCs exhibit

intermediate to low levels of binwise desynchronization up to the 500 ms time scale. However,

contrarily to GCs and FSs, they are able to achieve high levels of pattern separation through

scaling, thanks to a variable bursting response and low overall firing rates; 3) CA3 PCs exhibit

high levels of pattern separation through all tested neural codes, but significantly more than their

GC controls through binwise desynchronization only, especially at long time scales, suggesting

that CA3 might complement and amplify the separation inherited from the DG, possibly to make

pattern completion more efficient [40]. In the future, computational modelling combining those

results with knowledge of the synaptic dynamics between these populations of neurons will help

dissect the role played by each element of the network in hippocampal computations.

Our study especially calls for more work on the computational role of inhibitory synaptic

dynamics as it reveals the first experimental evidence that fast GABAergic transmission favors

pattern separation through multiple neural codes (binwise desynchronization and scaling) and

impairs it through another code (occupancy variations at short time scales) (**Fig 7**). Past research

has long suspected the importance of inhibition to hippocampal computations. Computational

models of the DG implementing perisomatic feedforward inhibition (provided by FSs) in the

form of a "k-winners take all" rule controlling the number of GCs activated by a given input

pattern have stressed the importance of such inhibition on pattern separation through a

population code [4, 54, 55]. The seminal model of Myers and Scharfman (2009) also suggested

that certain hilar interneurons, providing feedforward and feedback inhibition on the distal

dendrites of GCs, could impair population pattern separation through their influence on other

GABAergic interneurons (see also Danielson et al. 2017). Experimentally, a recent behavioral

study on mice showed that normal levels of tonic inhibition in GCs are critical for mnemonic

discrimination [80]. Our results now demonstrate that inhibition is also critical for computations

that could support this episodic memory function. Future experiments will need to clarify the

exact role of different interneuron populations as well as different types of inhibition (e.g.

feedforward vs feedback or somatic vs dendritic). Such work will especially be critical to further

understand CA3 computations and how they differ from those of DG, because our findings on

CA3 are based on recordings with partially reduced inhibition. On one hand, the enhancement of

pattern separation we observed in CA3 could be due to abnormal pharmacological conditions, on

the other hand, the necessity to lower inhibition to get CA3 PCs to spike could instead indicate

that CA3 must be disinhibited during episodic memory formation.

Methods:

Herein, we revisited experiments performed in Madar et al. (2018) [40] and expanded on

them by using additional types of input patterns, new pharmacological conditions, and by

analyzing the previous and new datasets with multiple similarity metrics assuming different

neural codes.

Experiments

All experiments were performed in accordance with the National Institute of Health

guidelines outlined in the National Research Council Guide for the Care and Use of Laboratory

Animal (2011) and regularly monitored and approved by the University of Wisconsin

Institutional Animal Care and Use Committee.

Briefly, we performed experiments on C57Bl6 male mice (Harlan/Envigo) following the

general paradigm described in Madar et al. (2018) [40]. Horizontal acute slices of a mouse brain

hemisphere containing the hippocampus were prepared in cutting solution (CS; two version were

used. CS#1/CS#2, in mM: 83/80 NaCl, 26/24 NaHCO$_3$, 2.5/2.5 KCl, 1/1.25 NaH2PO$_4$, 0.5/0.5

CaCl$_2$, 3.3/4 MgCl2, 22/25 D-Glucose, 72/75 Sucrose, 0/1 Na-L-Ascorbate, 0/3 Na-Pyruvate,

bubbled with 95% O$_2$ and 5% CO$_2$) and recorded at 33-35°C in regular artificial cerebrospinal

fluid (aCSF, in mM: 125 NaCl, 25 NaHCO3, 2.5 KCl, 1.25 NaH2PO4, 2 CaCl2, 1 MgCl2, and

25 D-Glucose). Two second input trains of varying similarity were delivered to neural afferents

of DG or CA3 neurons via bipolar electrical stimulation through a glass theta pipette, while

recording the membrane potential of a single responsive patched neuron in whole-cell current-

clamp mode. For the intracellular solution (IS), we used two different recipes that yielded similar

electrophysiological behaviors and results (IS#1/IS#2 in mM: 140/135 K-gluconate, 0/5 KCl,

10/0.1 EGTA, 10/10 HEPES, 20/20 Na-phosphocreatine, 2/2 Mg$_2$ATP, 0.3/0.3 NaGTP, 0/0.25

CaCl2, adjusted to pH 7.3 and 310 mOsm with KOH and H$_2$O). The location of the stimulation

(>100um away from dendrites of the recorded neuron) and its intensity (0.1-1 mA) was

determined to yield ~50% probability of spiking after an input pulse (range: ~20-80%). A

stimulation protocol consisted of five or ten trains (i.e. a given input set), interleaved (with 5 s of

pause between each train) and repeated ten or five times, respectively, such that fifty output

traces were recorded (**Fig 1C**). The amount of pattern separation or convergence effectuated for

a given recording was assessed by pairwise similarity analysis (defined below), displayed as

what we refer to as a "pattern separation graph": the pairwise input similarity across all pairs of

input trains (excluding self-comparisons) was compared to the average similarity of all pairs of

output spike trains coming from the corresponding pairs of input trains (excluding comparisons

between output spike trains coming from the same input train) (**Fig 1D-E**).

Figs 1-3 (GCs) and **Figs 8-9** (all celltypes) are based on recordings from GCs, fast-

spiking interneurons (FSs), hilar mossy cells (HMCs) and CA3 pyramidal cells (CA3 PCs)

reanalyzed from Madar et al. (2018) [40]. They were performed on slices (CS#1 for GCs, CS#2

for other celltypes; IS#1 for all) from young mice (p15-25), using 11 input sets consisting of five

Poisson spike trains with a mean rate of 10 Hz (GCs, FSs, HMCs) or 30 Hz (CA3 PCs and their

GC controls). GCs (10 Hz), FSs and HMCs were recorded in regular aCSF. Because it was very

difficult to promote firing of CA3 PCs in aCSF, CA3 PCs and their GCs control were recorded in

aCSF with 100 nM gabazine (gzn, SR-95531) to decrease inhibitory postsynaptic currents

(IPSC) amplitude by ~30% and allow CA3 PCs to occasionally escape inhibition and fire action

potentials. Note that the GCs recorded in gabazine as controls for the CA3 PC experiments could

not be directly compared to GC recordings in regular aCSF because input sets with different

mean rates were used.

To determine the role of inhibition on pattern separation in GCs, we performed additional

recordings of single GCs in slices from young mice (p22-32, 6 animals) (**Fig 7**). After cutting

(CS#2), slices were moved to a storing chamber containing 100% CS, at 37°C for 30 minutes

after dissection. As in all other experiments, the storing chamber was then placed at room

temperature and slices left there for at least 30 min. Slices were then transferred to the recording

chamber and perfused (5 ml/min) with oxygenated regular aCSF at 33-35°C. Patch-clamp

recordings of GCs were performed using IS#1. An experiment consisted of recording the same

GC under the same input set of five 10 Hz Poisson trains (average R_{input} = 0.76 at 10 ms, shown

in **Fig 1B**) before and after addition of 100 nM gabazine to the flow of regular aCSF. We waited

at least five minutes after the solution switch, to allow for equilibrium to be reached. The holding

potential (~-70mV), the location of the stimulation electrode, the stimulus current intensity and

all other parameters were unchanged between the two recordings (e.g. under gabazine, the

spiking probability was not retargeted to ~50%, unlike the gabazine experiments using 30 Hz

inputs mentioned above). The intrinsic properties of recorded GCs (n = 7) were (mean ± SEM):

1) Before gzn: resting membrane potential V_{rest} = -76.5 ± 2.5 mV, input resistance R_i = 194 ± 30

MΩ and membrane capacitance C_m = 20 ± 1.4 pF; 2) After gzn: V_{rest} = -74.4 ± 2 mV, R_i = 184 ±

29 MΩ and C_m = 17 ± 1.6 pF. R_i was not significantly affected and V_{rest} and C_m were only

slightly decreased after treatment (Wilcoxon signed rank test: V_{rest} p = 0.09; R_i p = 0.44; C_m p =

0.03). The slight drift of C_m was likely due to the small increase in access resistance occurring

over time (before: 6 ± 0.8 MΩ, after: 7.3 ± 1.4 MΩ; p = 0.06).

To explore the impact of a variety of input pattern statistical structures on single neuron

computations, we designed two new input sets comprised of 10 different two seconds spike

trains (**Fig 4-6**). Trains in "input set A" follow a Poisson (P) distribution and have an average

Pearson's correlation of 75% at the 10 ms time scale, but each train has a different mean firing

rate (PΔFR, **Fig 4A**). Trains in "input set B" do not follow a Poisson distribution: they all have

21 spikes, making them all 10.5 Hz overall, but are more or less bursty (B) (i.e. for each spike

train the same number of spikes was distributed in a different number of time bins. Occupied

time bins were selected randomly – following a uniform distribution – among all the possible

time bins) (B 10.5 Hz, **Fig 4B**). For these experiments, we used slices from two adult mice (p145

and p153) injected with saline at p46 as controls for a separate set of experiments (not described

here). Slices were prepared as described above (CS#2) and patch-clamp recordings were

performed in regular aCSF using IS#2. The intrinsic properties of recorded GCs were: 1) Input

set A (n = 5): V_{rest} = -80.4 ± 2.9 mV, R_i = 155 ± 19 MΩ and C_m = 23 ± 2 pF; 2) Input set B (n =

3): V_{rest} = -78.0 ± 5.5 mV, R_i = 178 ± 14 MΩ and C_m = 24 ± 3 pF.

All chemicals and drugs were purchased from Sigma-Aldrich (USA).

Input spike trains

As explained above, three types of input sets were used, defined by the statistical

structure (P for Poisson, B for Bursty) and the firing rate of their spike trains (constant or varying

across trains): 1) P10Hz (**Fig 1-3, 7-9**), 2) PΔFR (input set A in **Fig 4-6**), 3) B10.5Hz (input set B

in **Fig 4-6**).

Input sets with Poisson spike trains (P10Hz and PΔFR) were generated using two

different algorithms allowing to specify the number of spike trains in an input set (5 or 10,

respectively), the firing rate of each spike train, and the average Pearson's correlation coefficient

across all spike trains (computed with τ_w = 10ms). First, through iterative modifications of a

random Poisson spike train, we generated five different P10Hz input sets of 5 trains with average

correlation R_{input} = 0.88, 0.84, 0.74, 0.65, 0.56. Six other P10Hz input sets of 5 trains (R_{input} =

1.00, 0.95, 0.76, 0.48, 0.26, 0.11) and one PΔFR set of 10 trains (R_{input} = 0.75) were generated

using the Matlab toolbox provided by Macke and colleagues [81], which uses a more efficient

and mathematically principled algorithm that randomly samples from a multivariate Poisson

distribution with preset rate (FR) and covariance matrix. For both algorithms, although pairwise

correlations were constrained around the specified mean R_{input} (e.g. at τ_w = 10 ms, on average

across all P10Hz input sets, the relative standard error of pairwise R_{input} values is 4% of the mean

R_{input}), some variability remained (see **Fig 1E right**) which allowed to test a wide array of

pairwise input correlations (**Fig 1F**). Because this variability may be larger when measuring

spike train similarity with different metrics than R, or at τ_w values that were not used to design

input sets, we reported pairwise similarity values (as opposed to the average similarity across all

trains) throughout the article.

To design an input set with 10 spike trains spanning a range of burstiness (B10.5Hz), we

simply constrained the number of spikes to be equal in all trains and specified the number of bins

with at least one spike for each train (bursty trains have a lower number of occupied bins).

Spikes were assigned to randomly selected bins and to random times within those bins.

Similarity metrics

We assessed the similarity between spike trains in four ways, spanning a range of

different assumptions on the neural code: 1) with the Pearson's correlation coefficient (R), 2)

with the normalized dot product (NDP), 3) with the scaling factor (SF) and 4) with a distance

metric called SPIKE specifically designed to assess the dissimilarity between two spike trains

[48]. In addition, we used measures of differences between spike trains that assume different

neural codes, related to the one assumed by SF but easier to interpret in terms of basic spike train

features (see **Firing rate and burstiness codes** and **Dispersion metrics** sections below).

For R, NDP and SF, two spike trains X and Y of the same duration were divided into N

time bins of duration τ_w, with X_i and Y_i the respective numbers of spikes in bin i for each spike

train. In contrast, the SPIKE metric is binless and based on the spike times of each spike train.

For all similarity measures, empty spike trains (i.e. sweeps during which no spikes were evoked)

were excluded.

R (Pearson's correlation coefficient) can take values between 1 (perfectly correlated, i.e.

identical) and -1 (anticorrelated) with 0 indicating X and Y are uncorrelated. It was computed

with the following equation, where *cov* is the covariance (scalar), σ is the standard deviation and

$\bar{X}$ and $\bar{Y}$ are the means of X and Y:

$$R = \frac{cov(X,Y)}{\sigma_X . \sigma_Y} = \frac{\sum_{i=1}^{N}(X_i - \bar{X})(Y_i - \bar{Y})}{\sqrt{\sum_{i=1}^{N}(X_i - \bar{X})^2} \sqrt{\sum_{i=1}^{N}(Y_i - \bar{Y})^2}} \qquad (eq\ 1)$$

702   NDP (Normalized Dot Product) is the cosine of the angle θ between the two vectors: it is

703   0 when they are perfectly orthogonal, 1 when they are collinear (see **Fig 2A, C**). NDP was

704   computed as the dot product of X and Y divided by the product of their norms, with the

705   following equation (where X_i and Y_i are the coordinates of X and Y):

$$NDP = \cos(\theta) = \frac{\sum_{i=1}^{N} X_i Y_i}{\sqrt{\sum_{i=1}^{N} X_i^2} \sqrt{\sum_{i=1}^{N} Y_i^2}} \qquad (eq\ 2)$$

708   SF (Scaling Factor) quantifies the difference of length between the two vectors X and Y.

709   We have defined it as the ratio between the norms of each vector, the smaller norm always

710   divided by the bigger one to have SF values ranging from 0 to 1. SF = 1 means X and Y are

711   identical in terms of binwise spike number. The closer to 0 SF is, the more dissimilar are X and

712   Y (see **Fig 2A, C**). SF was computed with the following equation (where $0 < ||X|| \leq ||Y||$):

$$SF = \frac{||X||}{||Y||} = \frac{\sqrt{\sum_{i=1}^{N} X_i^2}}{\sqrt{\sum_{i=1}^{N} Y_i^2}} \qquad (eq\ 3)$$

NDP and SF focus on complementary features of a system of two vectors (angle vs

norms), and thus, together they are sufficient to fully describe the similarity between two vectors

in a Euclidean space.

The binless SPIKE similarity metric was computed from lists of spike times using the

Matlab toolbox provided at **www.fi.isc.cnr.it/users/thomas.kreuz/sourcecode.html**. The provided

software computes the SPIKE-distance between two spike trains (called D(t) in our study and

S(t) in eqn. 19 of the original paper) [48] and we derived the SPIKE similarity as: SPIKE =

$1 - \frac{1}{T}\int_0^T D(t)dt$, where T is the duration of the spike train. Because D(t) ranges from 0 to 1,

SPIKE is thus also between 0 and 1, like NDP and SF. When SPIKE is equal to 1, spike trains

have exactly the same spike times, i.e. they are synchronous (n.b. in our experiments, spike trains

were not simultaneously recorded, but we use "synchronous" in the sense of spike trains aligned

to the start of each sweep). Note that SPIKE has a large dynamic range (i.e. sensitivity over large

differences of spiketimes), and, as a result, realistic spike trains like in our input sets rarely have

a SPIKE similarity lower than 0.45 [48, 82] (see **Fig 3D**).

The ranges of possible values for the metrics we used are listed below, where closer to 1

always means more similar (although what similar means is different for each metric):

$-1 \leq R \leq 1$
$0 \leq NDP \leq 1$
$0 < SF \leq 1$
$0 \leq \text{SPIKE-similarity} \leq 1$; but ≥ 0.45 in most cases

Each metric assumes a specific neural code. For instance, R, NDP and SF all consider

that the basic informative feature in spike trains is the number of spikes in a time bin (i.e. a spike

count code), but R and NDP suppose that spike trains are similar when the spike counts increase

or decrease similarly in the same time bins (with empty bins not counting for NDP), thus

assuming a binwise synchrony code, whereas SF assumes a code where it is the similarity in

overall burstiness or firing rate that is relevant (see **Fig 4A-B**), not the binwise synchrony (see

Fig 2A-C). SPIKE, on the other hand, considers a spike time synchrony code. Details on the

neural codes assumed by each metric, how they differ from each other and how they are

impacted by global increases in firing rate can be found in the **Supplements** (part 1). The main

assumptions and properties of each similarity metric on the neural code are summarized in **Table**

1. Note that we do not consider one metric to be better than the other, or better than alternative

metrics not used in this study. Each metric can be used as a tool to analyze different aspects of

the neural code, with a set of pros, cons and assumptions that complement other methods.

Firing rate and burstiness codes

R, NDP, SF and SPIKE are considering different aspects of spike trains and thus assume

different neural codes. However, it can be difficult to interpret those similarity measures in terms

of common properties of spike trains like the firing rate or the burstiness. To get a better sense of

the neural code assumed by each similarity metric, and to directly assess whether pattern

separation was performed through easily interpretable strategies like variation of the firing rate

or variation of the burstiness, we measured such properties in each spike train.

The firing rate is simply the number of spikes divided by the duration of a sweep (2

seconds). Measuring burstiness is less straightforward as there is currently no consensus on the

definition of a burst of action potentials [83-85]. We used three measures, detailed below.

Spike trains are often considered bursty when they diverge from a Poisson process [28,

85]. To determine the tendency of a neuron to fire bursts, we computed the burstiness of a given

output set as the Kullback-Leibler divergence (D_{KL}) of its interspike interval (ISI) frequency

distribution M (normalized to the median ISI) from the normalized ISI frequency distribution P

of a Poisson process. In practice, output sets from a 10 Hz input set were compared to the ISI

distribution from all combined 10 Hz input sets, whereas input sets from 30 Hz input set were

compared to the combined distribution of all 30 Hz input sets.

$$Burstiness = D_{KL}(M||P) = \sum_j M(j) \log\left(\frac{M(j)}{P(j)}\right) \qquad (eq\ 4)$$

where M(j) or P(j) is the value of M or P in the jth bin of the distribution. Normalized ISI

frequency distributions were discretized in 250 bins going from 0 to 50, and D_{KL} was computed

using the KLDIV matlab function (v 1.0.0.0), written by David Fass, available at:

https://www.mathworks.com/matlabcentral/fileexchange/13089-kldiv.

See **Supplements - 2** for details on why D_{KL} is a suitable measure of cell burstiness.

To assess burstiness in single spike trains, the D_{KL} measure being not well-suited for this

(see **Supplements - 2**), we designed two simple nonparametric measures that we called

compactness and *occupancy*. Instead of explicitly defining "burstiness", these measures are based

on the binning of spike trains, offering the advantage of being directly comparable to the neural

codes assumed by binned similarity metrics like R, NDP and SF (**Fig S2**). A spike train is thus

seen as a vector of spike counts, and a time bin (or vector coordinate) is considered a window in

which spikes can be clustered. Thus, instead of counting the number of detected bursts and the

number of spikes in a burst, our two measures provide information on the number of occupied

bins (Compactness) and the number of spikes in occupied bins (Occupancy). These measures of

burstiness should be viewed as complementary to previous definitions, not as replacements.

The compactness of a spike train was computed as follows:

$$Compactness = 1 - proportion\ of\ occupied\ bins$$

$$= 1 - \frac{number\ of\ bins\ with\ at\ least\ 1\ spike}{total\ number\ of\ bins} \qquad (eq\ 5)$$

Compactness thus runs between 0 (all bins are occupied by at least one spike) and 1 (no spikes:

this case was excluded so the true maximum, corresponding to any spike train with all its spikes

clustered in just one bin, was $1 - \frac{1}{total\ number\ of\ bins}$). Values closer to 1 mean very few bins

are occupied: either many spikes are clustered in a small number of bins or the measured spike

train has very few spikes.

The occupancy of a spike train is the average number of spikes per occupied bin:

$$Occupancy = \frac{number\ of\ spikes}{number\ of\ occupied\ bins} \qquad (eq\ 6)$$

Because we excluded empty spike trains, the occupancy can theoretically go from 1 to infinity.

Note that the lowest value (i.e. 1 spike per occupied bin) means that all spikes are distributed in

different bins (or that there is just 1 spike).

Compactness and occupancy are both dependent on the firing rate of a given spike train,

but not in the same way (see **Supplements - 2** for details). They are complementary measures

that, together, allow determining whether two spike trains with different firing rates are also

different in terms of burstiness (**Table S1**). When spike trains have the same firing rate (e.g.

input set B 10.5 Hz in **Fig 4B**, which were all spike trains with 21 spikes each but were

constructed by specifying different numbers of occupied bins), compactness and occupancy are

redundant (**Fig S2C-D**) and, for both measures, higher values mean higher burstiness.

Dispersion metrics

To assess the amount of pattern separation performed by variation of a given spike train

feature like the firing rate, compactness or occupancy, we measured the difference in this feature

between two spike trains. In the same spirit as for similarity metrics, we could then compute the

difference between all pairs of trains in an input set (excluding self-comparisons) and compare to

the difference of all corresponding pairs of output spike trains, (excluding comparisons between

output spike trains associated with the same input spike train). This would yield matrices of

pairwise differences organized in the same fashion as the similarity matrix displayed in **Fig 1E**,

which could then be converted to pattern separation graphs showing the average pairwise output

absolute difference as a function of the pairwise input difference (**Fig 6A**). Note that the mean

absolute difference (MAD) between all spike trains of a given set (either input or output)

corresponds to the Gini mean difference [86], a common measure of dispersion of a sample akin

to the standard deviation (**Fig 5A**). The difference in MAD between an input set and its output

set indicates whether pattern separation or convergence was achieved through variation of the

measured spike train feature (**Fig 6B**).

Software and statistics

Data analysis was performed using MATLAB (Mathworks, Natick, MA, USA). Sample sizes were chosen based on the literature and estimations of the variance and effect size from preliminary data. The one-sample Kolmogorov-Smirnov test was used to verify the normality of data distributions. Parametric or non-parametric statistical tests were appropriately used to assess significance (p-value < 0.05). Assumptions on equal variances between groups were avoided when necessary. All tests on means or medians were two-tailed. To determine whether distributions of similarity values [S_{input}, S_{output}] (S standing for any similarity metric) were significantly different at a given time scale (**Fig 7, 8**), we performed an analysis of the covariance (ANCOVA) using separate lines regression models, with S_{input} as a continuous predictor and Treatment (**Fig 7**) or Celltype (**Fig 8**) as a categorical predictor with 2 or 5 levels respectively (the *aoctool* function in Matlab and a custom-written code yielded the same results). The 95% confidence interval around the slope and intercept of a given linear model was determined from the *regress* function (**Fig 8**).

Data and code availability

Data and code used in this manuscript are freely available:

- Data: https://www.ebi.ac.uk/biostudies/studies/S-BSST219

- Code: https://github.com/antoinemadar/PatSepSpikeTrains

Fig 1. Temporal pattern decorrelation at the level of single dentate granule cells.
(**A**) Histology of the DG in a horizontal slice (Cresyl violet/Nissl staining; scale bar: 250 μm), overlaid with a schematic of the experimental setup: a theta pipette in the ML is used to focally stimulate the perforant path (input) while a responding GC is recorded via whole-cell patch-clamp (output). GCL: granule cell layer, H: hilus, ML: molecular layer. Celltype color code: green for granule cell (GC); red for fast spiking interneuron (FS); orange for hilar mossy cell (HMC); black for CA3 pyramidal cell. Solid lines represent dendrites and dashed lines axons

(B) Examples of input sets. *Top*: each input set is constituted of five different trains of electrical
pulses following a Poisson distribution and an average rate of 10 Hz. *Bottom*: correlation
coefficient matrix for each input set, each square is the Pearson's correlation coefficient (R)
between two input trains considered as vectors of spike counts with a binning window (τ_w) of 10
ms. R_{input} is the average of coefficients, diagonal excluded. We used 11 such input sets (R_{input} =
0.11 to 1).
(C) Current-clamp recordings of the membrane potential of a single GC in response to a single
input set (average R_{input} = 0.76). Each input set (five input trains) is repeated ten times, yielding a
recording set of fifty output voltage traces.
(D) Sorted input and output rasters from the recording set in C (each parent input train has ten
children output trains forming an output subset, with matching colors). PW means pairwise.
(E) *Left*: Corresponding 55x55 correlation coefficients matrix using a τ_w of 10 ms. Each small
grey square represents the correlation coefficient between two spike trains. The average of
coefficients from children output trains corresponding to one of the ten pairs of input trains
(identified by color-coded squares), was computed to yield ten pairwise R_{output} coefficients. This
excludes comparisons between outputs generated from the same parent input. *Right*: mean across
multiple cells, following the same color-code as displayed in the matrix on the left.
(F) Pattern separation graph showing the mean $\pm$ SEM pairwise R_{output} across 102 recordings
(from 28 GCs), for 11 different input sets (i.e. 110 pairwise R_{input}). Values under the identity line
(dashed) demonstrate that output spike trains were less similar to each other than inputs, and thus
that pattern separation was performed. Blue indicates significant difference from the identity line
(one-sample t-test on the difference pairwise R_{input} - pairwise R_{output}, $p < 0.05$). The solid grey
line is a parabolic fit to the 102 recording sets.
[Fig adapted from Madar et al. (2018) [40] (fig. 1, 2, 5)]

871
Fig 2. Orthogonalization of input spike trains is a strong component of temporal pattern separation by single granule cells.
(**A-C**) Six hypothetical cases of pairs of spike trains and their associated Pearson's correlation coefficients (R), normalized dot products (NDP, cosine of the angle between two vectors) and scaling factors (SF, ratio of the norms), showing that the three metrics assume different neural codes and are not equivalent. (**A**) Synthetic spike trains (X and Y pairs) divided into six bins, with the corresponding number of spikes per bin. (**B**) R between each pair of X and Y describes the linear regression between the number of spikes in the bins of X versus the corresponding bins in Y (jitter was added to make all points visible). (**C**) Geometric view of vectors X and Y, where each bin is a dimension of a 6-dimensional space, and the number of spikes in a bin is the coordinate along this dimension. NDP measures how far from orthogonal two spike trains are and SF measures how different their binwise firing rates are. (**A-C**) NDP and R are sensitive to whether spike trains have spikes in the same bins or not (row 1), whereas SF is sensitive to differences in spike counts per bin (row 2), although neither NDP nor R is purely about "bin synchrony" (row 3). Note that these examples provide the intuition that orthogonal vectors (NDP = 0) necessarily correspond to a negative correlation between the spike trains (row 1 and 5) but that anticorrelated spike

trains ($R < 0$) are not necessarily orthogonal (row 6), and that orthogonal spike trains are not necessarily
perfectly anticorrelated as in row 1 because R, unlike NDP, considers common silent bins as correlated
(row 5). See also **Fig S1.**
(D) Vector representation of experimental data from one recording set, showing the average similarity
between a set of input spike trains (dashed line and green angle, $R_{input} = 0.76$) and the average similarity
between the fifty corresponding output spike trains excluding comparisons between outputs from same
parent input (solid line, purple angle). The angles are derived from the NDP whereas the lengths of each
vector express differences in binwise firing rates (SF). Here, outputs are more orthogonal (closer to 90˚)
than their inputs with little difference in scaling.
(E-F) Pattern separation graphs showing the pairwise output similarity as a function of the pairwise input
similarity, as measured by the NDP or SF (mean+/-SEM) across 102 recording sets (28 GCs) for 11
different input sets. As in Fig 1F, blue indicates a significant difference from the identity line (one-sample
t-test, $p < 0.05$). The solid grey line is a parabolic (E) or linear (F) fit to the 102 recording sets. **(E)**
Outputs are closer to orthogonality (NDP = 0) than their respective inputs, demonstrating strong levels of
pattern separation through orthogonalization at the 10 ms time scale. **(F)** At the 10 ms time scale, pattern
separation by scaling is present for inputs very similar in terms of SF (SF_{input} close to 1), but weak, at least
over the span of tested SF_{input} values.
904

Fig 3. Single granule cells exhibit pattern separation on millisecond to second time scales using different codes.

(**A-C**) Pattern separation/convergence assessed at different time scales and using different measures of similarity (S). *Top*: pairwise S_{output} (average across recording sets) as a function of pairwise S_{input}, measured with different binning windows τ_w. Solid curves are linear fits. Each color corresponds to a different τ_w ranging from 10 ms to 500 ms. *Bottom*: Effect of τ_w on the effective decorrelation ($S_{input} - S_{output}$), interpolated from the linear regressions. Note that as τ_w increases, pattern separation through decorrelation (R) or orthogonalization (NDP) becomes weaker while it becomes stronger through scaling (SF). (**D**) Similarity between spike trains is here assessed with the binless SPIKE metric, directly using spike times. *Left*: example of two input spike trains associated with two output spike trains from a GC recording set, and the corresponding distances D(t) between spike trains. D(t) can then be integrated over time to give a single value D. *Middle:* example of 55x55 matrix of SPIKE similarity (1-D) between all spike trains of an example recording set. 0 means that spikes of two trains never happened close in time, and 1 that they were perfectly synchronous. The output SPIKE similarity ($SPIKE_{output}$) is defined as the average similarity excluding comparisons between spike train from the same parent input train (i.e. average of the 10 pairwise $SPIKE_{output}$). *Right*: $SPIKE_{output}$ of the same 102 GC recordings as in Fig 1-3, as a function of $SPIKE_{input}$, fitted with a parabola (red line). Most data points are below the dashed identity line indicating that output spike trains are less similar than inputs. The average $SPIKE_{input}$-$SPIKE_{output}$ is significantly above 0 for all input sets except the two most dissimilar ($SPIKE_{input}$ = 0.74, 0.78) (one-sample T-tests, p < 0.05).

Fig 4. Pattern convergence via scaling in single granule cells.

(A-B) Two new input sets were designed to explore single GCs computations on inputs with a wide range of pairwise similarity as measured by SF. Both input sets have ten input trains of 2 s, and were repeated five times during the whole-cell patch-clamp recording of a single GC, yielding 50 output spike trains.

(A) *Left*: Input set A was constituted of spike trains following a Poisson distribution, each with a different firing rate (FR), increased from 7 to 31.5 Hz, but with an average R_{input} constrained around 0.75 ($\tau_w = 10$ ms). The ten input trains were thus well correlated but varied in their firing rate. *Middle*: matrix of SF coefficients for each pair of trains in input set A, showing a wide range of values unlike the 11 input sets with 10 Hz trains used in previous experiments. *Right*: SF values are indeed sensitive and anti-correlated to differences in overall firing rate between two spike trains (data points correspond to pairs of trains).

(B) Because SF is dependent on how differently clustered spikes are in two different trains, SF can vary even when the overall firing rate is the same. *Left*: Input set B was constituted of spike trains with 21 spikes (FR = 10.5 Hz) that were clustered in more or less bins, leading to trains with varying burstiness

and thus a wide range of SF values (*Middle*). R was not constrained but ended up close to 0 for all pairs.
Right: SF values are indeed sensitive and anticorrelated to differences in occupancy (average number of
spikes per occupied bins), a direct measure of burstiness when FR is constant across trains (data points
correspond to pairs of trains).
(C) Pattern separation graphs showing the pairwise output similarity as a function of the pairwise input
similarity, as measured by SF, across 5 GCs responses to input set A and 3 GCs responses to input set B
(Both input sets tested GC responses over 45 pairwise R_{input}). *Left and Middle*: Mean+/-SEM. Blue
indicates a significant difference from the identity line (one-sample t-test, $p < 0.05$). The solid grey line is
a linear fit to the 5x45 (*Left*) or 3x45 (*Middle*) data points. Both experiments confirm that, at the 10 ms
time scale, GCs exhibit pattern separation through scaling for highly similar input trains (SF close to 1) as
in Fig 2F, but show that GCs can exhibit pattern convergence for more dissimilar inputs (SF < 0.7). *Right*:
Mean across GC recordings for both experiments combined, measured at different time scales. It confirms
that SF_{output} decreases when using larger binning windows, leading to more pattern separation for high
input similarity and lower pattern convergence for lower input similarities.

Fig 5. Granule cells input-output transformations in terms of firing rate and burstiness.
(**A**) Graphs shown correspond to the same GC in response to either input set A or B. Data points
correspond to individual spike trains, (black empty circles for inputs: 10 trains per set; green filled circles
for GC outputs: 50 trains per set). For each spike train, we measured its overall FR, binwise compactness
and binwise occupancy, all spike train features providing information on sparseness and burstiness (see
Methods). Error bars correspond to two different measures of dispersion. Black: mean +/- SD; red: min-
max center of gravity ± the mean absolute difference (self-comparisons and comparisons between output
trains with the same parent input train were excluded). Comparing mean values of a given spike train
feature between inputs and outputs shows the direction of the transformation from inputs to outputs.
Comparing the dispersion between inputs and outputs shows whether pattern separation (larger output
dispersion) or convergence (smaller output dispersion) was achieved through a neural code focused on the
measured spike train feature.
(**B**) For each output spike train, the output spike train measures of compactness, occupancy or overall FR
is plotted against the measure of its parent input train (input set A or B). For all time scales, GCs have
generally higher compactness than their inputs, but maintain a low and narrow range of occupancy
regardless of their input statistics, showing that GCs are sparser (see **Table S1**). Consistently, GCs
maintain a lower FR than their inputs.

Fig 6. Depending on inputs statistical structure, granule cells exhibit pattern separation or convergence via burstiness and firing rate codes.

(**A**) Pattern separation graphs showing the average pairwise output absolute difference across GC recordings as a function of the pairwise input absolute difference, for the binwise compactness (*Left*), binwise occupancy (*Middle*) and overall FR (*Right*). Insets zoom in on the smallest values of the input-axis. Experiments using input set A and B were combined. Note that because we plot the absolute difference between two spike trains (i.e. the distance between two data points in Fig 5A) and not a measure of similarity, pattern separation corresponds to points above the identity line (dashed) and pattern convergence to points below. Error bars are SEM (not displayed in middle graph for readability). *Right*: input set A recordings (PΔFR) in black, input set B (B 10.5 Hz) in green.

(**B**) The difference in dispersion (Dispersion$_{output}$ − Dispersion$_{input}$; dispersion is the mean absolute difference, i.e. red bars in Fig 5A) of binwise compactness, binwise occupancy or overall FR for each GC recording set from three experiments using different types of input sets: Poisson trains with 10 Hz FR and varying R (P10Hz: 102 recordings, 28 GCs), Poisson trains with R ≈ 0.75 and varying FR (PΔFR, i.e. input set A: 5 GCs), trains with 10 Hz FR and varying burstiness (B 10.5 Hz, i.e. input set B: 3 GCs). Positive values correspond to pattern separation through variations of the given spike train property (compactness, occupancy or FR) and negative values to pattern convergence. Asterisks denote significant difference from 0, i.e. either pattern separation or convergence (one-sample t-test, p < 0.05. See **Table 2**).

Fig 7. Inhibition controls levels of pattern separation in granule cells through different neural codes

(**A**) Single GCs were recorded twice in response to the same input set (P10Hz, R = 0.76 at τ_w = 10 ms, shown in Fig 1B): First in normal aCSF, then under conditions of partial inhibitory block (aCSF + 100 nM gabazine, gzn), without changing any other parameter (i.e. same stimulation location and intensity).

(**B**) Partial disinhibition of GCs led to higher FR and a higher propensity to fire small bursts of spikes riding on the same EPSP, as illustrated in A and evidenced by an increased p(Burst) (probability of firing more than one spike between two input pulses). FR and p(Burst) were computed for each recording set (n = 7 GCs). Paired t-test (FR and p(Burst) respectively): p = 0.02, p = 0.04.

(**C**) Pattern separation graphs showing the pairwise output similarity as a function of the pairwise input similarity, as measured by R, NDP or SF at the 10 ms time scale (mean+/-SEM) across recordings in 7 GCs. Data points below the identity line (dashed) correspond to pattern separation. Partial block of inhibition in GCs led to a significant decrease in pattern separation through all similarity metrics (ANCOVA with separate-lines model fitting 70 data points per treatment group: p < 0.0001). Solid lines are the linear fits used for the ANCOVA.

(**D**) Cumulative frequency distributions of the distance of (pairwise S_{input}, pairwise S_{output}) data points to the identity line in pattern separation graphs as in C (insets show medians). Positive values of the x-axis correspond to pattern separation, and negative values to pattern convergence. Red curves (after gzn) are all shifted to the left of their black counterparts (before gzn), showing a decrease in pattern separation at all time scales. The shift is more pronounced at larger time scales for R and NDP, at smaller time scales for SF. ANCOVA on (pw S_{input}, pw S_{output}) data points with separate-lines model: p < 0.0001 for τ_w up to 1000 ms for R and NDP, p < 0.02 for SF.

(**E**) Average levels of pattern separation via compactness, occupancy or FR codes were measured for each recording set (as in Fig 6B). Surprisingly, partial block of inhibition did not significantly change the variations in FR or binwise compactness of output spike trains, but it led to less pattern convergence or even pattern separation through variations of occupancy (paired t-test: p = 0.008, 0.024, 0.031 for τ_w = 10, 20 and 50 ms respectively. No significant difference between treatments for larger τ_w or other spike train features at any τ_w.

1020
Fig 8. Levels of pattern separation through orthogonalization and scaling differ between hippocampal celltypes.

(A) Temporal pattern separation assays were performed by single whole-cell current-clamp recordings from GCs (green, same experiments as Fig 1-3), FSs (red), HMCs (orange) and CA3 PCs (black). For GCs, FSs and HMCs, input sets with Poisson trains of 10 Hz FR (e.g. Fig 1B) were delivered by stimulating in the outer molecular layer (OML), and recordings done in regular aCSF. For CA3 PCs those stimulus parameters rarely if ever elicited spiking, therefore input sets were constituted of Poisson trains with a 30 Hz FR delivered in the GCL, and recordings were performed under partial block of inhibition (100 nM Gabazine). Control experiments (teal) were performed in GCs under the same pharmacological conditions using 30 Hz input sets, but the stimulations were delivered in the OML. The membrane potential was maintained around -70 mV (between -70 and -60 mV for CA3 PCs). Note: although all celltypes and treatments are displayed on the same graphs for concision (B-D and Fig 9), CA3 PCs should only be directly compared to their corresponding GC controls.

(B) *Top*: Pattern separation graphs showing the pairwise output similarity as a function of the pairwise input similarity, as measured by NDP or SF at the 50 ms time scale (mean+/-SEM: 28 GCs, 3-13 recordings per input set; 4 FSs, 4 recordings per input set; 11 HMCs, 5-7 recordings per input set; 14 CA3 PC + gzn, 6-9 recordings per input set; 13 control GCs + gzn, with 11 recordings per input set). Same color code as in A. Data points below the identity line (dashed) correspond to pattern separation. All celltypes performed pattern separation through both orthogonalization and scaling but with different levels. *Bottom*: We performed ANCOVAs with separate-lines model for all five celltypes (e.g. solid lines shown in the top graphs) for 11 different time scales (5-1000 ms). Dashed horizontal line represents

significance level at 0.05: orthogonalization and scaling levels differ between celltypes at most time
scales.
(C-D) Slope and intercept of the linear models (e.g. solid lines in B, top) used to fit [S_{input}, S_{output}] data
points for a given celltype and time scale. Shaded patches correspond to the 95% confidence interval of
the parameter estimate. Significant differences between celltypes in pattern separation functions (with
Tukey-Kramer correction for multiple comparisons) are detailed in **Table 3**.
1048

1049

1050
Fig 9. Levels of pattern separation and convergence through burstiness and firing rate
codes differ between hippocampal celltypes.
(**A**) Assessment of burstiness in different celltypes (binless analysis).
Left: Cumulative frequency distributions of interspike intervals (ISI) observed in the output spike trains of
different hippocampal celltypes and their associated input trains. ISI of all recordings of a given celltype
or input set type were pooled. The ISIs were normalized to their median for a given celltype or input type,
such that their cumulative distribution would cross at frequency = 0.5, and that all exponential cumulative
distributions characteristic of a Poisson process (as both type of input sets are) would be super-imposed or
very close to each other. Thus, GCs are visually close to an exponential distribution, suggesting their
output is close to a Poisson process, i.e. not bursty. Distributions falling away from their corresponding
input distribution suggest that spikes are organized in a non-Poisson manner, suggestive of a bursty
behavior. In this sense, HMCs, CA3 PCs and their GC controls seem to exhibit a similar bursty behavior,
and FSs an even more bursty one.
Right: Quantification of the burstiness of each recording set (102 for GCs, 20 for FSs, 18 for HMCs, 15
for CA3 PCs+gzn and 22 for GCs+gzn). Burstiness was assessed as the Kullback-Leibler divergence from

P to M, with M the frequency distribution of normalized ISI in an output set (50 output trains), and P the
frequency distribution of normalized ISI in the associated input set type (either Poisson 10Hz or Poisson
30 Hz input sets were combined, as in A). The higher the divergence, the burstier a cell is. This analysis
confirms that GCs outputs are close to a Poisson process, and that all other celltypes and conditions are
significantly burstier (Kruskal-Wallis one-way ANOVA: $p < 0.0001$; post-hoc t-tests with Tukey-Kramer
correction for multiple comparisons: $p < 0.01$ for all GC (10Hz) to other celltypes comparisons. CA3 PCs
and GC controls (30Hz) are not significantly different ($p = 0.99$) nor FSs and HMCs ($p = 0.64$)). Box
plots: central mark and edges are the median, the 25th and the 75th percentiles, respectively. Whiskers are
most extreme data points not considered outliers (red +).
(B) Average levels of pattern separation via binwise compactness, binwise occupancy or FR codes were
measured for each recording set of every celltype, as in Fig 6B (negative values indicate pattern
convergence). Levels were different between celltypes (Kruskal-Wallis one-way ANOVA: $p < 0.0001$ at
all time scales; post-hoc t-tests with Tukey-Kramer correction for multiple comparisons in **Table 4**).

1081
1082   References
1083
1.   Kirwan CB, Hartshorn A, Stark SM, Goodrich-Hunsaker NJ, Hopkins RO, Stark CE. Pattern
separation deficits following damage to the hippocampus. Neuropsychologia. 2012; 50(10): 2408-14.
doi: 10.1016/j.neuropsychologia.2012.06.011.
2.   Treves A, Tashiro A, Witter MP, Moser EI. What is the mammalian dentate gyrus good for?
Neuroscience. 2008; 154(4): 1155-72. doi: 10.1016/j.neuroscience.2008.04.073.
3.   Berron D, Schutze H, Maass A, Cardenas-Blanco A, Kuijf HJ, Kumaran D, et al. Strong Evidence for
Pattern Separation in Human Dentate Gyrus. J Neurosci. 2016; 36(29): 7569-79. doi:
10.1523/jneurosci.0518-16.2016.
4.   O'Reilly RC, McClelland JL. Hippocampal conjunctive encoding, storage, and recall: avoiding a
trade-off. Hippocampus. 1994; 4(6): 661-82. doi: 10.1002/hipo.450040605.
5.   Santoro A. Reassessing pattern separation in the dentate gyrus. Front Behav Neurosci. 2013; 7:
96. doi: 10.3389/fnbeh.2013.00096.
6.   Rolls ET. A computational theory of episodic memory formation in the hippocampus. Behav
Brain Res. 2010; 215(2): 180-96. doi: 10.1016/j.bbr.2010.03.027.
7.   Chavlis S, Poirazi P. Pattern separation in the hippocampus through the eyes of computational
modeling. Synapse. 2017. doi: 10.1002/syn.21972.
8.   Lisman J, Buzsaki G, Eichenbaum H, Nadel L, Ranganath C, Redish AD. Viewpoints: how the
hippocampus contributes to memory, navigation and cognition. Nat Neurosci. 2017; 20(11): 1434-47.
doi: 10.1038/nn.4661.
9.   O'Keefe J. Place units in the hippocampus of the freely moving rat. Exp Neurol. 1976; 51(1): 78-
109. doi.
10.   Colgin LL, Moser EI, Moser MB. Understanding memory through hippocampal remapping.
Trends Neurosci. 2008; 31(9): 469-77. doi: 10.1016/j.tins.2008.06.008.
11.   Hainmueller T, Bartos M. Parallel emergence of stable and dynamic memory engrams in the
hippocampus. Nature. 2018. doi: 10.1038/s41586-018-0191-2.
12.   Leutgeb JK, Leutgeb S, Moser MB, Moser EI. Pattern separation in the dentate gyrus and CA3 of
the hippocampus. Science. 2007; 315(5814): 961-6. doi: 10.1126/science.1135801.
13.   Neunuebel JP, Knierim JJ. CA3 retrieves coherent representations from degraded input: direct
evidence for CA3 pattern completion and dentate gyrus pattern separation. Neuron. 2014; 81(2): 416-
27. doi: 10.1016/j.neuron.2013.11.017.
14.   Nakazawa K. Dentate Mossy Cell and Pattern Separation. Neuron. 2017; 93(3): 465-7. doi.
15.   Krueppel R, Remy S, Beck H. Dendritic integration in hippocampal dentate granule cells. Neuron.
2011; 71(3): 512-28. doi: 10.1016/j.neuron.2011.05.043.
16.   Hayakawa H, Samura T, Kamijo TC, Sakai Y, Aihara T. Spatial information enhanced by non-
spatial information in hippocampal granule cells. Cogn Neurodyn. 2015; 9(1): 1-12. doi: 10.1007/s11571-
014-9309-x.
17.   Liu YC, Cheng JK, Lien CC. Rapid dynamic changes of dendritic inhibition in the dentate gyrus by
presynaptic activity patterns. J Neurosci. 2014; 34(4): 1344-57. doi: 10.1523/JNEUROSCI.2566-13.2014.
18.   Bliss TV, Lomo T. Long-lasting potentiation of synaptic transmission in the dentate area of the
anaesthetized rabbit following stimulation of the perforant path. J Physiol. 1973; 232(2): 331-56. doi.
19.   Lopez-Rojas J, Heine M, Kreutz MR. Plasticity of intrinsic excitability in mature granule cells of
the dentate gyrus. Sci Rep. 2016; 6: 21615. doi: 10.1038/srep21615.
20.   Mongiat LA, Esposito MS, Lombardi G, Schinder AF. Reliable activation of immature neurons in
the adult hippocampus. PLoS One. 2009; 4(4): e5320. doi: 10.1371/journal.pone.0005320.

21.   Ewell LA, Jones MV. Frequency-tuned distribution of inhibition in the dentate gyrus. J Neurosci.
2010; 30(38): 12597-607. doi: 10.1523/jneurosci.1854-10.2010.
22.   Pardi MB, Ogando MB, Schinder AF, Marin-Burgin A. Differential inhibition onto developing and
mature granule cells generates high-frequency filters with variable gain. Elife. 2015; 4: e08764. doi:
10.7554/eLife.08764.
23.   Dieni CV, Panichi R, Aimone JB, Kuo CT, Wadiche JI, Overstreet-Wadiche L. Low excitatory
innervation balances high intrinsic excitability of immature dentate neurons. Nat Commun. 2016; 7:
11313. doi: 10.1038/ncomms11313.
24.   Kamijo TC, Hayakawa H, Fukushima Y, Kubota Y, Isomura Y, Tsukada M, et al. Input integration
around the dendritic branches in hippocampal dentate granule cells. Cogn Neurodyn. 2014; 8(4): 267-76.
doi: 10.1007/s11571-014-9280-6.
25.   Zylberberg J, Hyde RA, Strowbridge BW. Dynamics of robust pattern separability in the
hippocampal dentate gyrus. Hippocampus. 2016; 26(5): 623-32. doi: 10.1002/hipo.22546.
26.   Hyde RA, Strowbridge BW. Mnemonic representations of transient stimuli and temporal
sequences in the rodent hippocampus in vitro. Nat Neurosci. 2012; 15(10): 1430-8. doi:
10.1038/nn.3208.
27.   Fellous JM, Rudolph M, Destexhe A, Sejnowski TJ. Synaptic background noise controls the
input/output characteristics of single cells in an in vitro model of in vivo activity. Neuroscience. 2003;
122(3): 811-29. doi.
28.   Legendy CR, Salcman M. Bursts and recurrences of bursts in the spike trains of spontaneously
active striate cortex neurons. J Neurophysiol. 1985; 53(4): 926-39. doi: 10.1152/jn.1985.53.4.926.
29.   Weyand TG, Boudreaux M, Guido W. Burst and tonic response modes in thalamic neurons
during sleep and wakefulness. J Neurophysiol. 2001; 85(3): 1107-18. doi: 10.1152/jn.2001.85.3.1107.
30.   Destexhe A, Rudolph M, Pare D. The high-conductance state of neocortical neurons in vivo. Nat
Rev Neurosci. 2003; 4(9): 739-51. doi: 10.1038/nrn1198.
31.   Pernía-Andrade Alejandro J, Jonas P. Theta-Gamma-Modulated Synaptic Currents in
Hippocampal Granule Cells In Vivo Define a Mechanism for Network Oscillations. Neuron. 2014; 81(1):
140-52. doi: 10.1016/j.neuron.2013.09.046.
32.   Cook EP, Guest JA, Liang Y, Masse NY, Colbert CM. Dendrite-to-soma input/output function of
continuous time-varying signals in hippocampal CA1 pyramidal neurons. J Neurophysiol. 2007; 98(5):
2943-55. doi: 10.1152/jn.00414.2007.
33.   Dobrunz LE, Stevens CF. Response of hippocampal synapses to natural stimulation patterns.
Neuron. 1999; 22(1): 157-66. doi.
34.   Fuhrmann G, Segev I, Markram H, Tsodyks M. Coding of temporal information by activity-
dependent synapses. J Neurophysiol. 2002; 87(1): 140-8. doi.
35.   Gundlfinger A, Breustedt J, Sullivan D, Schmitz D. Natural spike trains trigger short- and long-
lasting dynamics at hippocampal mossy fiber synapses in rodents. PLoS One. 2010; 5(4): e9961. doi:
10.1371/journal.pone.0009961.
36.   Mistry R, Dennis S, Frerking M, Mellor JR. Dentate gyrus granule cell firing patterns can induce
mossy fiber long-term potentiation in vitro. Hippocampus. 2011; 21(11): 1157-68. doi:
10.1002/hipo.20815.
37.   Berger TW, Eriksson JL, Ciarolla DA, Sclabassi RJ. Nonlinear systems analysis of the hippocampal
perforant path-dentate projection. II. Effects of random impulse train stimulation. J Neurophysiol. 1988;
60(3): 1076-94. doi.
38.   Xie X, Song D, Wang Z, Marmarelis VZ, Berger TW. Interaction of short-term neuronal plasticity
and synaptic plasticity revealed by nonlinear systems analysis in dentate granule cells. Conf Proc IEEE
Eng Med Biol Soc. 2006; 1: 5543-6. doi: 10.1109/iembs.2006.259706.

39.   Berger TW, Eriksson JL, Ciarolla DA, Sclabassi RJ. Nonlinear systems analysis of the hippocampal
perforant path-dentate projection. III. Comparison of random train and paired impulse stimulation. J
Neurophysiol. 1988; 60(3): 1095-109. doi: 10.1152/jn.1988.60.3.1095.
40.   Madar AD, Ewell LA, Jones MV. Pattern separation of spiketrains in hippocampal neurons; 2018.
Preprint. Available from: bioRxiv: 10.1101/107706.
41.   Yim MY, Hanuschkin A, Wolfart J. Intrinsic rescaling of granule cells restores pattern separation
ability of a dentate gyrus network model during epileptic hyperexcitability. Hippocampus. 2014. doi:
10.1002/hipo.22373.
42.   Senzai Y, Buzsáki G. Physiological properties and behavioral correlates of hippocampal granule
cells and mossy cells. Neuron. 2017; 93(3): 691-704. e5. doi.
43.   Danielson NB, Turi GF, Ladow M, Chavlis S, Petrantonakis PC, Poirazi P, et al. In Vivo Imaging of
Dentate Gyrus Mossy Cells in Behaving Mice. Neuron. 2017; 93(3): 552-9. e4. doi.
44.   McNaughton BL, Nadel L. Hebb-Marr networks and the neurobiological representation of action
in space. Neuroscience and connectionist theory. 1990: 1-63. doi.
45.   Rieke F, Warland W, de Ruyter van Steveninck R, Bialek W. Spikes: Exploring the Neural Code.
THE MIT PRESS: A Bradford Book; 1999.
46.   VanRullen R, Guyonneau R, Thorpe SJ. Spike times make sense. Trends Neurosci. 2005; 28(1): 1-
4. doi: 10.1016/j.tins.2004.10.010.
47.   Gutig R. To spike, or when to spike? Curr Opin Neurobiol. 2014; 25: 134-9. doi:
10.1016/j.conb.2014.01.004.
48.   Kreuz T, Chicharro D, Houghton C, Andrzejak RG, Mormann F. Monitoring spike train synchrony.
J Neurophysiol. 2013; 109(5): 1457-72. doi: 10.1152/jn.00873.2012.
49.   Dengler CG, Coulter DA. Normal and epilepsy-associated pathologic function of the dentate
gyrus. Prog Brain Res. 2016; 226: 155-78. doi: 10.1016/bs.pbr.2016.04.005.
50.   Hosp JA, Struber M, Yanagawa Y, Obata K, Vida I, Jonas P, et al. Morpho-physiological criteria
divide dentate gyrus interneurons into classes. Hippocampus. 2014; 24(2): 189-203. doi:
10.1002/hipo.22214.
51.   Ramaswamy S. Exciting times for inhibition: GABAergic synaptic transmission in dentate gyrus
interneuron networks. Front Neural Circuits. 2015; 9: 13. doi: 10.3389/fncir.2015.00013.
52.   Lee CT, Kao MH, Hou WH, Wei YT, Chen CL, Lien CC. Causal Evidence for the Role of Specific
GABAergic Interneuron Types in Entorhinal Recruitment of Dentate Granule Cells. Sci Rep. 2016; 6:
36885. doi: 10.1038/srep36885.
53.   Hsu TT, Lee CT, Tai MH, Lien CC. Differential Recruitment of Dentate Gyrus Interneuron Types by
Commissural Versus Perforant Pathways. Cereb Cortex. 2016; 26(6): 2715-27. doi:
10.1093/cercor/bhv127.
54.   Myers CE, Scharfman HE. A role for hilar cells in pattern separation in the dentate gyrus: a
computational approach. Hippocampus. 2009; 19(4): 321-37. doi: 10.1002/hipo.20516.
55.   Temprana SG, Mongiat LA, Yang SM, Trinchero MF, Alvarez DD, Kropff E, et al. Delayed coupling
to feedback inhibition during a critical period for the integration of adult-born granule cells. Neuron.
2015; 85(1): 116-30. doi: 10.1016/j.neuron.2014.11.023.
56.   McKenzie S, Frank AJ, Kinsky NR, Porter B, Riviere PD, Eichenbaum H. Hippocampal
Representation of Related and Opposing Memories Develop within Distinct, Hierarchically Organized
Neural Schemas. Neuron. 2014; 83(1): 202-15. doi: 10.1016/j.neuron.2014.05.019.
57.   Eichenbaum H. On the integration of space, time, and memory. Neuron. 2017; 95(5): 1007-18.
doi.
58.   Lin L, Osan R, Shoham S, Jin W, Zuo W, Tsien JZ. Identification of network-level coding units for
real-time representation of episodic experiences in the hippocampus. Proc Natl Acad Sci U S A. 2005;
102(17): 6125-30. doi: 10.1073/pnas.0408233102.

59.   Panzeri S, Brunel N, Logothetis NK, Kayser C. Sensory neural codes using multiplexed temporal
scales. Trends in neurosciences. 2010; 33(3): 111-20. doi.
60.   Adrian ED. The basis of sensation. Christophers; London, 22 Berners Steeet, W. 1; 1928.
61.   Osborne LC, Palmer SE, Lisberger SG, Bialek W. The neural basis for combinatorial coding in a
cortical population response. J Neurosci. 2008; 28(50): 13522-31. doi: 10.1523/jneurosci.4390-08.2008.
62.   Kepecs A, Lisman J. Information encoding and computation with spikes and bursts. Network.
2003; 14(1): 103-18. doi.
63.   Mease RA, Kuner T, Fairhall AL, Groh A. Multiplexed Spike Coding and Adaptation in the
Thalamus. Cell Rep. 2017; 19(6): 1130-40. doi: 10.1016/j.celrep.2017.04.050.
64.   Lundstrom BN, Fairhall AL. Decoding stimulus variance from a distributional neural code of
interspike intervals. J Neurosci. 2006; 26(35): 9030-7. doi: 10.1523/jneurosci.0225-06.2006.
65.   Oram M, Wiener M, Lestienne R, Richmond B. Stochastic nature of precisely timed spike
patterns in visual system neuronal responses. Journal of neurophysiology. 1999; 81(6): 3021-33. doi.
66.   Dettner A, Munzberg S, Tchumatchenko T. Temporal pairwise spike correlations fully capture
single-neuron information. Nat Commun. 2016; 7: 13805. doi: 10.1038/ncomms13805.
67.   Thorpe S, Delorme A, Van Rullen R. Spike-based strategies for rapid processing. Neural Netw.
2001; 14(6-7): 715-25. doi.
68.   Chacron MJ, Longtin A, Maler L. Negative interspike interval correlations increase the neuronal
capacity for encoding time-dependent stimuli. J Neurosci. 2001; 21(14): 5328-43. doi.
69.   Lisman JE, Jensen O. The theta-gamma neural code. Neuron. 2013; 77(6): 1002-16. doi:
10.1016/j.neuron.2013.03.007.
70.   Elijah DH, Samengo I, Montemurro MA. Thalamic neuron models encode stimulus information
by burst-size modulation. Front Comput Neurosci. 2015; 9: 113. doi: 10.3389/fncom.2015.00113.
71.   Chavlis S, Petrantonakis PC, Poirazi P. Dendrites of dentate gyrus granule cells contribute to
pattern separation by controlling sparsity. Hippocampus. 2017. doi.
72.   Neher T, Cheng S, Wiskott L. Memory storage fidelity in the hippocampal circuit: the role of
subregions and input statistics. PLoS Comput Biol. 2015; 11(5): e1004250. doi:
10.1371/journal.pcbi.1004250.
73.   Aimone JB, Wiles J, Gage FH. Computational influence of adult neurogenesis on memory
encoding. Neuron. 2009; 61(2): 187-202. doi: 10.1016/j.neuron.2008.11.026.
74.   Ratté S, Hong S, De Schutter E, Prescott SA. Impact of neuronal properties on network coding:
roles of spike initiation dynamics and robust synchrony transfer. Neuron. 2013; 78(5): 758-72. doi.
75.   Hersman S, Rodriguez Barrera V, Fanselow M. Assigning Function to Adult-Born Neurons: A
Theoretical Framework for Characterizing Neural Manipulation of Learning. Front Syst Neurosci. 2015; 9:
182. doi: 10.3389/fnsys.2015.00182.
76.   Scullin CS, Partridge LD. Modulation by pregnenolone sulfate of filtering properties in the
hippocampal trisynaptic circuit. Hippocampus. 2012; 22(11): 2184-98. doi: 10.1002/hipo.22038.
77.   Scharfman HE. The enigmatic mossy cell of the dentate gyrus. Nat Rev Neurosci. 2016; 17(9):
562-75. doi: 10.1038/nrn.2016.87.
78.   Amaral DG, Scharfman HE, Lavenex P. The dentate gyrus: fundamental neuroanatomical
organization (dentate gyrus for dummies). 2007; 163: 3-790. doi: 10.1016/s0079-6123(07)63001-5.
79.   Hu H, Gan J, Jonas P. Interneurons. Fast-spiking, parvalbumin(+) GABAergic interneurons: from
cellular design to microcircuit function. Science. 2014; 345(6196): 1255263. doi:
10.1126/science.1255263.
80.   Engin E, Zarnowska ED, Benke D, Tsvetkov E, Sigal M, Keist R, et al. Tonic Inhibitory Control of
Dentate Gyrus Granule Cells by alpha5-Containing GABAA Receptors Reduces Memory Interference. J
Neurosci. 2015; 35(40): 13698-712. doi: 10.1523/jneurosci.1370-15.2015.

81.   Macke JH, Berens P, Ecker AS, Tolias AS, Bethge M. Generating spike trains with specified
correlation coefficients. Neural Comput. 2009; 21(2): 397-423. doi: 10.1162/neco.2008.02-08-713.
82.   Satuvuori E, Kreuz T. Which spike train distance is most suitable for distinguishing rate and
temporal coding? Journal of neuroscience methods. 2018; 299: 22-33. doi.
83.   Robin K, Maurice N, Degos B, Deniau JM, Martinerie J, Pezard L. Assessment of bursting activity
and interspike intervals variability: a case study for methodological comparison. J Neurosci Methods.
2009; 179(1): 142-9. doi: 10.1016/j.jneumeth.2009.01.020.
84.   Lobb C. Abnormal Bursting as a Pathophysiological Mechanism in Parkinson's Disease. Basal
Ganglia. 2014; 3(4): 187-95. doi: 10.1016/j.baga.2013.11.002.
85.   Cotterill E, Charlesworth P, Thomas CW, Paulsen O, Eglen SJ. A comparison of computational
methods for detecting bursts in neuronal spike trains and their application to human stem cell-derived
neuronal networks. J Neurophysiol. 2016; 116(2): 306-21. doi: 10.1152/jn.00093.2016.
86.   Yitzhaki S. Gini's mean difference: A superior measure of variability for non-normal distributions.
Metron. 2003; 61(2): 285-316. doi.
1284
1285

Supplements

3   This document provides supplementary Figures, tables and a discussion on how to

4   interpret the similarity and burstiness metrics used in the article "Temporal pattern separation in

5   hippocampal neurons through multiplexed neural codes".

7 – 1 – How to interpret similarity metrics

Measuring the similarity between two spike trains is a difficult problem [1, 2], and many

methods have been designed to this effect, each with a set of pros, cons and assumptions [2-6].

We could not be exhaustive but, in order to span a large range of potential neural codes, we used

four different similarity metrics: R (Pearson's correlation coefficient), NDP (Normalized Dot

product), SF (Scaling Factor) and SPIKE (1 - SPIKE distance). (Note: 'metrics' is used with a

loose sense throughout the paper). Their pros, cons and assumptions are detailed in this section

and summarized in **Table 1** of the main text.

16   Binned vs binless metrics

We used two different kinds of similarity metrics: 1) binned metrics (R, NDP, SF)

consider a spike train as a list X of spike counts X_i (with X_i the number of spikes in time bin i of

a given duration τ_w) and thus assume a neural code where spike counts are the basic informative

feature; 2) a binless metric SPIKE that considers a spike train as a list of spike times and thus

assumes a neural code where the time relative to the start of the given spike train is the basic

informative feature.

Although R has already been used to measure the similarity between spike trains [2, 7],

none of the binned metrics we used were designed with spike trains in mind and thus have

properties that might not be well suited for the study of spike trains. For instance, R, NDP and

SF have the drawback of considering all X_i (i.e. binned spike counts) as independent

observations, which may not be a realistic assumption. Moreover, they assume spike trains are

related linearly, or that they belong to a Euclidean space. Binless metrics like SPIKE avoid these

limitations [8].

Another limitation of binned metrics as implemented here (and this is also true for

compactness and occupancy) is that they are subject to edge effects: small shifts of the bins can

include or exclude spikes, which would lead to differences in the similarity measure. However,

the fact that marginally different binsizes (e.g. 5 ms vs 10 ms) lead to similar results (in other

words, similarity measures as a function of τ_w does not show spurious values: see Fig 3A-C, 6A,

7D, 8) suggests that edge effects do not impair our conclusions.

Differences between binned metrics

<u>R vs NDP and SF: epistemological differences</u>

The Pearson's correlation coefficient R was designed by early statisticians as a tool to

quantify the strength and direction of a linear regression [9]. R thus considers two spike trains X

and Y as variables, with X_i and Y_i the observations, and assesses, across all i, whether X_i and Y_i

tend to vary in a similar fashion (**Fig 2B**).

On the other hand, NDP and SF are metrics coming from the fields of Euclidean

geometry and linear algebra. In contrast to R, they explicitly consider spike trains as vectors,

with each time bin i corresponding to a coordinate, or dimension, in an N-dimensional space (N

being the total number of bins). NDP measures the angle between two vectors, and SF is based

on vector norms (**Fig 2C**).

R, NDP and SF are not equivalent

Although they have been designed for different purposes, both R [7] and NDP [10, 11]

have been used to measure the similarity between neuronal activity patterns, including spike

trains. In this context, NDP has sometimes loosely been called a measure of correlation [10, 12].

What difference is there between R and NDP then, especially in terms of the assumed neural

code? A comparison of equation 1 and 2 (**Methods**) reveals that R is actually an NDP, but of

baseline-subtracted vectors. In other words, R does not consider spike trains as raw spike count

vectors but as vectors of deviation from their respective mean spike count. This ensures that R is

an optimal estimator of correlation [9] but it has the inconvenience of not conserving the angular

relationship between X and Y, making NDP necessary if one wants to evaluate pattern separation

as an orthogonalization process. Moreover, using baseline-subtracted vectors also means that R

considers common silent periods (i.e. $X_i = Y_i = 0$) as correlated, which is an assumption on the

neural code that has been considered unrealistic by some [2]. In contrast, NDP excludes those

empty bins and thus assumes a neural code where common periods of inactivity are irrelevant.

Comparing equations 1, 2 and 3 (**Methods**), the relationships between R, NDP and SF

appear not trivial and not linear. To further evaluate the relationship between R, NDP and SF, we

compared the values of each similarity metrics for all pairs of a set of simulated spike trains. This

set contained the $3^6 = 729$ possible spike count vectors made of six bins, with each bin containing

0, 1 or 2 spikes (some examples are displayed in **Fig 2A**). This analysis confirmed that the three

metrics are not equivalent and provides an intuition on what each metric represents (**Fig S1A**).

Interestingly, even though R, NDP and SF are not linearly related in principle, our

measures of R and NDP between experimentally recorded spike trains have a near linear

relationship, as opposed to R and SF (**Fig S1B**). This correlation is likely due to our choice of

input spike trains as well as to biological constraints (e.g. refractory period) forcing the temporal

structure of output spike trains to vary only along certain dimensions. It is thus reflective of the

actual computations performed in the DG and shows that although R and NDP carry different

sets of assumptions about the neural code, spike trains recorded in GCs vary mostly along

properties that both R and NDP are sensitive to.

R and NDP vs SF: sensitivity to individual firing rate differences and dependency on

pairwise firing rate levels

To interpret R, NDP and SF and their different assumed neural codes, it is important to

understand their relationship with the firing rate. Studying the impact of firing rate on similarity

metrics can however mean different things.

First, a similarity metric can be sensitive to differences of firing rate between two spike

trains X and Y. For instance, if $X = kY$ (with k an integer > 1), that means that X and Y have

spikes in exactly the same bins (i.e. X and Y have the same temporal structure) but that Y has a

higher firing rate. In such a case, both R and NDP would consider X and Y as perfectly similar

(see example #2 in **Fig 2A-C**). In contrast, SF is designed to extract $1/k$ and thus strongly picks

up on differences of firing rate. On the other hand, if X and Y have the same firing rate but that

their spikes are in different bins (different temporal structure), R and NDP will consider those

spike trains completely dissimilar, and SF will see them as perfectly similar (see example #1 in

Fig 2A-C). Other cases, where X and Y have spikes in the same time bins but only some bins

have a different number of spikes (e.g. example #3 in **Fig 2A-C**) reveal that R and NDP are not

totally insensitive to differences of firing rate when those are not proportional, but overall these

two metrics are mostly sensitive to binwise synchrony, unlike SF (**Fig S2**).

Second, similarity metrics can be influenced by the overall firing rate level of a pair of

spike train (for instance measured as the geometric mean of the firing rate of X and Y). Whether

an ideal similarity metric should be independent of the firing rate in such a way is debatable, and

finding one is not trivial [2, 13], yet it is generally seen as an undesirable property [2] because it

makes the comparison of similarity values between two pairs of spike trains more challenging to

interpret [1]. Indeed, for spike trains X and Y with random spike times, the probability of spikes

occurring close in time will intuitively be higher if X and Y have higher rates [14], even though

their temporal structure would be conserved.

Because R is often used to measure spike train correlations [1], its relationship to

pairwise firing rate levels has already been studied. It was for instance demonstrated that, when

X and Y are two Poisson spike trains with the same rate λ, the normalization factor and the use

of variables centered on their mean (which is, in the case of spike trains, a linear function of the

firing rate) makes R mathematically independent of λ [14]. However, in the general case, R has

still a complex relationship with the firing rates of both spike trains [13].

To get a better sense of this relationship for R, and study the effect of pairwise firing rate

levels on NDP and SF as well, we considered two cases: 1) proportional firing rate increases, and

2) additive firing rate increases.

In the first case, we considered two spike trains A and B such that A = kX and B = kY

(with k an integer > 1). Some metrics aiming to measure the similarity between spike train in

terms of temporal structure, like the correlation index, have been shown to result in higher values

for the (A, B) pair compared to the (X, Y) pair [2]. However, in this case, R, NDP and SF values

would be the same for both pairs (A, B) and (X, Y): thanks to the denominator in their respective

formula, all three are insensitive to proportional increases in firing rates.

In the second case, we used the set of 729 6-bin spike count vectors described above and

generated new sets of related vectors with higher firing rates, such that for each spike train X of

the original set there was a vector A in the new set with $A_i = X_i + k$ when $X_i > 0$, and $A_i = X_i$

otherwise. In other words, the rate was additively (not proportionally) increased while preserving

the temporal structure of the initial spike trains and maintaining constant the difference in the

binwise number of spikes between spike trains (e.g. for k = 5, if X = [010101] and Y = [020202],

then A = [060606] and B = [070707]. For both pairs, the difference is [010101].). Our analysis

shows that such an increase in the firing rate can affect all three metrics, but in different ways

(**Fig S1C**): NDP and R can increase or decrease, and on average do not change much, whereas

SF systematically increases. A larger increase of the firing rate leads to more variability of the

effect size for NDP and R whereas the effect monotonically increases for SF.

SF is thus clearly dependent on pairwise firing rate levels, and R and NDP can be

affected somewhat as well. Thus, to directly assess the variability of specific spike train features

across a set of spike trains (i.e. individual firing rate, compactness and occupancy) without

contaminating interpretations with consideration of pairwise firing rate levels (especially in the

case of SF), we used additional metrics detailed in **Methods – dispersion metrics**.

<u>Temporal resolution of binned metrics</u>

As binned metrics, R, NDP and SF are dependent on the binsize, i.e. a specified temporal

resolution τ_w. In other words, in equations 1 to 3, X_i and Y_i are functions of τ_w. Temporal

resolution logically influences the value of metrics measuring similarity in terms of the temporal

structure of spike trains, such as R [1] or NDP. However it also influences SF: even though SF is

sensitive to differences in firing rate, it is designed to be sensitive to differences in binwise firing

rate, which makes it also sensitive to differences in burstiness even when the firing rate is

constant (e.g. **Fig 4B** and **Fig S2**).

In **Fig 3**, we evaluated the influence of τ_w on our results by varying it between 5 ms and

1000 ms. Note that because our spike trains are two-seconds long, using $\tau_w = 2000$ ms would

mean the spike trains can be seen as variables with only one observation, or as unidimensional

vectors (i.e., scalars) that can only vary by their norm (i.e., firing rate). In this case, R is

meaningless, NDP is necessarily 1, which indicates collinearity and SF would directly assess the

difference in total spike number (i.e. overall firing rate) between the two spike trains.

152
Fig S1 (related to Fig 2). **R, NDP and SF are not equivalent.**
(**A**) Relationships between these 3 similarity metrics (R, NDP and SF) computed between the 3^6 possible
spike count vectors with six bins and that can have 0, 1 or 2 spikes per bin (All pairs combinations =
499,500 data points).
(**B**) Relationships between these three similarity metrics for the 124,950 pairs of spike trains from the 102
experimental GC recording sets. Green lines correspond to a linear regression (R^2 and p-value in each
panel). Note that although R and NDP are well correlated in our experimental data ($R^2 > 0.95$), there is not
a linear relationship between R and NDP in theory.
(**C**) Impact of non-proportional firing rate increase on similarity values for all three metrics. For each 6-
bin spike count vector X (same as in A), a vector A was generated such that for each bin i with at least
one spike, $A_i = X_i + k$, with k = 1, 10 or 100, and $A_i = X_i$ when $X_i = 0$ (e.g. for k = 1, if X = [010101] and
Y = [020202], then A = [020202] and B = [030303]. For both pairs, the difference is [010101]). NDP, SF
and R values were computed for all pairs in each set of vectors, and the values of the initial set (x-axis)
was compared to the values of the X+k set (y-axis). Data points away from the identity line (red)
demonstrate an effect of the firing rate increase.

168
Fig S2 (related to Fig 4)**. Influence of firing rate and burstiness differences on binned**
similarity metrics.
Data points correspond to single pairs of input trains from input set A (**A-B**) or input set B (**C-D**). The
similarity between two spike trains (R, NDP or SF), their difference of compactness (proportion of
occupied bins), and their difference in occupancy (average number of spikes per occupied bins) were
measured using binning windows of 10 ms (**A, C**) or 100 ms (**B, D**). When FR is constant across spike
trains, compactness and occupancy differences are direct measures of burstiness differences, given a
certain time scale. As intuited from Fig 2A-C, SF is very sensitive to variations in FR, compactness and
occupancy between spike trains, whereas R and NDP are only mildly influenced at best (and mostly at
larger time scales), by such variations.

2 – Burstiness metrics

Cell burstiness vs spike train burstiness

To assess the tendency of a neuron to fire bursts, we computed the Kullback-Leibler

divergence $D_{KL}(M\|P)$ of the ISI distribution in a set of output spike trains from the ISI

distribution of a Poisson process. $D_{KL}(M\|P)$ is an asymmetric, information theory-based measure

of the inefficiency of the assumption that the distribution of ISI M is a distribution P (Poisson, in

our case), and thus can be interpreted as the divergence of one probability distribution from

another. Note that a train with regularly spaced spikes would diverge from Poissonness without

being bursty, but that natural spike trains are rarely regular, and accordingly none of our

recordings showed a regular ISI distribution, suggesting that $D_{KL}(M\|P)$ is a good indirect

indicator of burstiness.

The D_{KL} approach above has the advantage of being independent of the firing rate, but

has two limitations. First it is parametric, based on the nonconsensual opinion that a Poisson

process is the least bursty of processes [15]. Second, because it is based on the comparison of ISI

distributions, it requires a large sample of ISIs to be accurate and thus cannot be used to measure

the burstiness of a single two second spike train.

In order to assess the burstiness of single spike trains, we designed two binned metrics:

compactness and occupancy. We used binned metrics to avoid the difficult problem of burst

detection that generally requires some arbitrary threshold to define what is considered a burst

[16]. Unlike D_{KL} these metrics are not independent of the firing rate, but together they allow

distinguishing between burstiness and firing rate (see below).

Compactness and occupancy: complementary relationships to the firing rate allow

disambiguating between burstiness and lack of sparseness

It is unclear whether the concept of burstiness should be independent or not of the firing rate (for instance, should we consider a spike train with 100 spikes clustered in the same 10 ms bin as bursty as a spike train with just 10 spikes in a single bin?), but, as explained above for similarity metrics, a dependency of burstiness metrics on the firing rate can make interpretations more challenging. Both compactness and occupancy are dependent on the firing rate, but in different ways, which resolves the ambiguity: 1) Compactness does not provide information on the firing rate per se, and is only related to the firing rate in the sense that, in the case of Poisson-like spiking, higher rates lead to a higher chance of occupying a large number of bins. Thus a spike train with just one spike will be considered as compact as a spike train with a large number of spikes all clustered into one bin. In contrast, the occupancy is directly related to the number of spikes and thus to the firing rate of a given spike train (Indeed, when considering a time bin as long as the full duration of a spike train, i.e. two seconds in our case, the occupancy is exactly proportional to the firing rate.). Occupancy would thus distinguish between a sparse spike train with just one spike (Occupancy = 1) and a spike train with 10 spikes clustered in one bin (Occupancy = 10). 2) On the other hand, spike trains with different firing rates can have the same occupancy (e.g. if all spikes are distributed in different bins, Occupancy = 1 regardless of the number of spikes) but would be differentiated by their level of compactness (100 distributed spikes would be considered much less compact than a spike train with just one spike). Therefore, when assessing pairs of spike trains with different firing rates, compactness and occupancy provide complementary information that allows determining whether one spike train is unambiguously burstier or simply has a higher firing rate (see **Table S1**).

Table S1. (related to Fig 7) **How to interpret compactness and occupancy in terms of burstiness and sparseness when comparing two spike trains?**

The table shows what it means when *compactness* (C) is increased, stays constant or decreases while *occupancy* (O) increases, stays constant or decreases. We distinguish the cases when the mean firing rate (FR) is constant (FR absolute difference = 0) or different (FR abs diff > 0). The second spike train is either unambiguously burstier (B+) or less bursty (B-), has a greater FR (F+) or is sparser (F-), or a combination of those (i.e. B+F+ or B-F-). In each case, we provide an example (4-dimensional vectors of spike counts) when the conditions can coexist (NA means not applicable). Notice that the resulting table of examples is centrally symmetric.

C \ O		↗	—	↘
↗	FR abs diff = 0	**B+** 0101 → 0200	NA	NA
	FR abs diff > 0	**B+F+** 0111 → 0501	**F-** 0202 → 0200	**F-** 0222 → 0101
—	FR abs diff = 0	NA	2020 → 0202	NA
	FR abs diff > 0	**F+** 0101 → 0202	NA	**F-** 0202 → 0101
↘	FR abs diff = 0	NA	NA	**F-** 0200 → 0101
	FR abs diff > 0	**F+** 0101 → 0222	**F+** 0200 → 0202	**B-F-** 0501 → 0111

236   References
237

1.   Cohen MR, Kohn A. Measuring and interpreting neuronal correlations. Nature neuroscience.
2011; 14(7): 811. doi.
2.   Cutts CS, Eglen SJ. Detecting pairwise correlations in spike trains: an objective comparison of
methods and application to the study of retinal waves. J Neurosci. 2014; 34(43): 14288-303. doi:
10.1523/jneurosci.2767-14.2014.
3.   Lyttle D, Fellous JM. A new similarity measure for spike trains: sensitivity to bursts and periods
of inhibition. J Neurosci Methods. 2011; 199(2): 296-309. doi: 10.1016/j.jneumeth.2011.05.005.
4.   Kreuz T. Measures of spike train synchrony. Scholarpedia [Internet]. 2011; 6(10). Available from:
http://www.scholarpedia.org/article/Measures_of_spike_train_synchrony.
5.   Rusu CV, Florian RV. A new class of metrics for spike trains. Neural Computation. 2014; 26(2):
306-48. doi.
6.   Satuvuori E, Kreuz T. Which spike train distance is most suitable for distinguishing rate and
temporal coding? Journal of neuroscience methods. 2018; 299: 22-33. doi.
7.   Yim MY, Hanuschkin A, Wolfart J. Intrinsic rescaling of granule cells restores pattern separation
ability of a dentate gyrus network model during epileptic hyperexcitability. Hippocampus. 2014. doi:
10.1002/hipo.22373.
8.   Aronov D, Victor JD. Non-Euclidean properties of spike train metric spaces. Physical Review E.
2004; 69(6): 061905. doi.
9.   Stanton JM. Galton, Pearson, and the peas: A brief history of linear regression for statistics
instructors. Journal of Statistics Education. 2001; 9(3). doi.
10.   Schreiber S, Fellous JM, Whitmer D, Tiesinga P, Sejnowski TJ. A new correlation-based measure
of spike timing reliability. Neurocomputing. 2003; 52-54: 925-31. doi: 10.1016/s0925-2312(02)00838-x.
11.   Aimone JB, Wiles J, Gage FH. Computational influence of adult neurogenesis on memory
encoding. Neuron. 2009; 61(2): 187-202. doi: 10.1016/j.neuron.2008.11.026.
12.   Dieni CV, Panichi R, Aimone JB, Kuo CT, Wadiche JI, Overstreet-Wadiche L. Low excitatory
innervation balances high intrinsic excitability of immature dentate neurons. Nat Commun. 2016; 7:
11313. doi: 10.1038/ncomms11313.
13.   Amari S-i. Measure of correlation orthogonal to change in firing rate. Neural computation. 2009;
21(4): 960-72. doi.
14.   de la Rocha J, Doiron B, Shea-Brown E, Josic K, Reyes A. Correlation between neural spike trains
increases with firing rate. Nature. 2007; 448(7155): 802-6. doi: 10.1038/nature06028.
15.   Gourevitch B, Eggermont JJ. A nonparametric approach for detection of bursts in spike trains. J
Neurosci Methods. 2007; 160(2): 349-58. doi: 10.1016/j.jneumeth.2006.09.024.
16.   Cotterill E, Charlesworth P, Thomas CW, Paulsen O, Eglen SJ. A comparison of computational
methods for detecting bursts in neuronal spike trains and their application to human stem cell-derived
neuronal networks. J Neurophysiol. 2016; 116(2): 306-21. doi: 10.1152/jn.00093.2016.
274
275