GenEpi: Gene-based Epistasis Discovery Using Machine Learning

Yu-Chuan Chang[1], June-Tai Wu[2], Ming-Yi Hong[3], Yi-An Tung[4], Ping-Han Hsieh[3], Sook Wah Yee[5], Kathleen M. Giacomini[5,6], Yen-Jen Oyang[1], and Chien-Yu Chen[3,*], for the Alzheimer's Disease Neuroimaging Initiative†

[1]Graduate Institute of Biomedical Electronics and Bioinformatics, National Taiwan University, Taipei 10617, Taiwan. [2]Department of Dermatology, National Taiwan University Hospital, Taipei 10002, Taiwan. [3]Department of Bio-Industrial Mechatronics Engineering, National Taiwan University, Taipei 10617, Taiwan. [4]Genome and Systems biology degree program, Academia Sinica and National Taiwan University, Taipei 10617, Taiwan. [5]Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, San Francisco, California, USA. [6]Institute for Human Genetics, University of California, San Francisco, San Francisco, California, USA.

*To whom correspondence should be addressed.

†Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.ucla.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at:
https://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Chien-Yu Chen
Department of Bio-Industrial Mechatronics Engineering, National Taiwan University, Taipei 10617, Taiwan.
Phone: +886-2-3366-5334, E-mail: chienyuchen@ntu.edu.tw

Abstract

Motivation: Genome-wide association (GWA) studies provide a powerful means to identify

associations between genetic variants and phenotypes. However, GWA techniques for detecting

epistasis, the interactions between genetic variants associated with phenotypes, are still limited.

We believe that developing an efficient and effective GWA method to detect epistasis will be a

key for discovering sophisticated pathogenesis, which is especially important for complex

diseases such as Alzheimer's disease (AD). In this regard, this study presents GenEpi, a

computational package to uncover epistasis associated with phenotypes by the proposed

machine learning approach, and illustrates the application of GenEpi on predicting the diagnosis

and the progression of AD.

Methods: The key concept of GenEpi is a feature extraction process based on gene structures.

Since a gene is the minimal physical and functional unit of heredity, GenEpi considers a gene

as a unit to retrieve genetic variants as features. GenEpi adopts two-element combinatorial

encoding when producing features and constructs the prediction models by randomized

regression. Features are first modeled using these two approaches to detect the epistasis within

a single gene. The selected features for each gene are then considered together in order to

identify cross-gene epistasis. This study uses AD as an example to reveal the capability of

GenEpi in finding disease-related variants and variant interactions that show both biological

meanings and predictive power.

Results: We compared GenEpi with commonly used algorithms for detecting epistasis,

including FastEpistasis, BOOST and ReliefF. The simulation data demonstrated that GenEpi

outperforms the other methods in ranking the true epistasis as the top one. As real data is

concerned, the results suggest that the epistasis selected by GenEpi has the best predictive

power for two major phenotypes in the AD dataset: diagnosis of AD and disease progression.

For diagnosis, the proposed model of predicting AD contains three clinical features (Age,

Gender and Education) and 14 genetic features, including 24 SNPs from 12 genes that contain

the well-known causal gene, APOE. The 2-fold cross validation (CV) and leave-one-out CV

(LOO CV) accuracy of this model are 0.83 and 0.81, respectively. On the other hand, for

predicting progression, the proposed model contains eight clinical features (Age, Gender,

Education, Cognitively Normal (CN), Early Mild Cognitive Impairment (MCI), Late MCI, AD,

and MMSE at baseline) and four genetic features, including seven SNPs from six genes, where

all of the four genetic features are cross-gene epistasis with significant p-values ($< 10^{-11}$). The

average of Pearson and Spearman correlation of 2-fold CV and LOO CV are 0.52 and 0.53,

respectively. The results on AD demonstrated that GenEpi has the ability to detect the epistasis

associated with phenotypes effectively and efficiently. The released package can be generalized

to largely facilitate the studies of many complex diseases in the near future.

Availability: GenEpi is an open-source python package and available free of charge only for

non-commercial users. The package can be downloaded from

https://github.com/Chester75321/GenEpi, and has also been published on The Python Package

Index.

75   Introduction

Genome-wide association studies (GWAS) is a univariate examination of a genome-wide set of

genetic variants to determine if any single variant is associated with the phenotype of interest

[1]. The first GWAS was published in 2002 [2], and three years later, the most remarkable

GWAS regarding age-related macular degeneration (AMD) was published [3]. Their study

investigated the association of 105,980 single nucleotide polymorphisms (SNPs) with AMD on

96 cases and 50 control subjects. This study showed that the SNPs in the complement factor H

(CFH) gene, including a non-synonymous SNP, are significantly associated with AMD. Up to

2017, there have been more than hundreds of thousands individuals being studied in typical

GWAS protocols, and over 72,870 variant-disease associations between 46,589 SNPs and 6,356

phenotypes have been discovered [4]. These studies demonstrated the potential of GWAS to

identify genetic variants associated with many categories of phenotypes, including risk for

diseases such as various cancers, and variation in therapeutic and adverse response to drugs.

However, the success of univariate GWAS is limited to monogenic phenotypes (e.g. Mendelian

diseases). The impact of variant interactions, also known as epistasis on the formation of

diseases [5] is often underestimated in traditional GWAS analysis [6-8].

Researchers have found that GWAS has limitation in identifying the association in complex

diseases [9, 10]. Easton et al. suggested that a number of susceptible loci identified by GWAS

usually have very small effect sizes [11]. A major disadvantage of GWAS is that it considers

only one genetic variant at a time, and ignores underlying epistasis of variants that might have

stronger associations [12]. Studies have also demonstrated that the existence of epistasis is an

important factor contributing to phenotypes, especially in complex diseases such as

hypertension, diabetes and obesity [10]. Therefore, developing analytical methods to identify

epistasis efficiently is critical to understanding the genetic factors that contribute to human traits

[8, 13], and has attracted a wide range of research interests in recent years [7, 14].

Two main challenges to discover epistasis are computational complexity and statistical power

[15]. The first challenge results from the curse of dimensionality. When more genetic variants

are considered, the number of interactions increases exponentially. Based on the specification

of a major commercial technology, Illumina Arrays, a whole-genome array can investigate over

4 million markers per sample simultaneously. In order to evaluate the pairwise interactions from

this microarray, about 8×10^{12} statistical tests need to be processed. Even though Marchini et al.

have demonstrated that pairwise interactions of 3×10^{5} loci is computationally possible with

currently available computational resources, it still remains challenging when the Illumina

Arrays are considered [16]. The second challenge is the issue of statistical power. Since a huge

number of statistical tests are conducted on a limited sample size with high-dimensional

interactions, many false positives arise by random chances. In this regard, new methods have

been developed [10, 17]. Statistical approaches include FastEpistasis [18] and BOOST [19];

the later one has been included in a well-known GWAS software called PLINK [20, 21].

Machine learning approaches such as Multifactor Dimensionality Reduction [22], ReliefF [23],

random forest-like algorithms [24-26] and other methodologies have also been developed for

detecting epistasis [17].

Since the biological experiments used to validate these methodologies are still in demand, there

are no standard analysis methods for epistasis despite the rapid improvement in computational

performance. In 2016, Murk used FastEpistasis and BOOST to search SNP-SNP interactions

on a huge dataset called Genetic Epidemiology Research on Adult Health and Aging (GERA)

that included 78,486 subjects, but still failed to detect a significant and replicable interaction

after exhaustively searching through 45 billion possible interactions for 10 complex diseases of

interest [27]. Alzheimer's disease (AD) is one of these complex diseases and its pathogenesis,

which clearly has a genetic basis, is still ill-defined. In 2016, Sage Bionetworks held a

competition called The Dialogue for Reverse Engineering Assessments and Methods Challenge

(DREAM Challenge) for AD, which tried to use crowdsourcing to assess the capability of

current computational methods to predict the change in cognitive examination based on genetic

data. However, no significant contribution of genetic features except the *APOE* haplotype to

the predictive performance was observed by any competition teams [28]. To address this

problem, this study presents GenEpi, a package to reveal epistasis related to the phenotype

using machine learning and introduces the application of GenEpi on predicting the diagnosis

and the progression of AD.

Methods

The architecture of GenEpi is shown in Fig. 1. GenEpi is designed to group SNPs by gene

boundaries. GenEpi first considers the genetic variants in a gene as features, which have a

higher chance to interact with each other and to influence molecular functions. The idea of

within-gene epistasis analysis followed by cross-gene analysis is not new, which has been

proven successful in previous studies [29-32]. Differently, GenEpi adopts two-element

combinatorial encoding when producing features and models them by randomized regression,

which will be explained in Section 2.2. In the first stage (STAGE 1) of GenEpi, the genotype

features from each single gene will be combinatorically encoded and modeled independently

by randomized regression. In this way, we can estimate the prediction performance of each gene

and detect within-gene epistasis with a low false positive rate. In the second stage (STAGE 2),

both of the individual SNP and the within-gene epistasis features selected by STAGE 1 are

pooled together to generate cross-gene epistasis features. Finally, the genotype features after

feature selection are combined with environmental factors such as clinical features to build the

final prediction models. In addition to the main procedures, two pre-processing steps are also

implemented in GenEpi: retrieving the gene information from public databases and reducing

the gene information from public databases and reducing the dimensionality of the features

using linkage disequilibrium (LD). In the end, we released a Python package that implements

GenEpi (main steps are illustrated in Supplementary Fig. S1). The details of these steps and the

GenEpi method will be described in the following sections.

University of California Santa Cruz (UCSC) database

To obtain the gene information such as official gene symbols and genomic coordinates, we

retrieved kgXref and knownGene data table from the UCSC human genome annotation

database [33, 34]. The version of the database we used is the Feb. 2009 assembly of the human

genome hg19, GRCh37 Genome Reference Consortium Human Reference 37. The two data

tables were merged in order to generate a local database containing the gene symbols as well

as the genomic coordinates of each gene. The in-house script we built could update this local

database automatically. It is noted that there are many different categories of genes in the

RefSeq database. In this study, we only focused on the mRNA and non-coding RNA (22,376

genes in total). The selected transcripts were projected on the genomic coordinates and the

coordinates of corresponding genes were determined based on the leftmost and rightmost

positions of the corresponding transcripts. Moreover, to discover the factors that might affect

the transcription of genes, we also retained the promoter region of each gene. In genetics, the

promoter region is a segment of DNA that initiates the transcription of a particular gene.

Promoters are located near the transcription start sites of genes, on the upstream of the same

DNA strand (towards the 5' region of the sense strand of the transcript). In general, a promoter

region can be 100-1000 base pairs long. In this study, we extracted 1,000 nucleotides on the

upstream of the starting position of each gene as the promoter region.

Estimation of linkage disequilibrium

In GWAS datasets, a SNP often exhibits high dependency with its nearby SNPs because of

linkage disequilibrium (LD). In the practical implantation, we prefer to group these dependent

features to reduce the dimension of features. In other words, we can take the advantages of LD

to reduce the dimensionality of SNP features. In this regard, we adopted the same approach

developed by Lewontin [35] to estimate LD. We used D' >0.9 and r^2 >0.9 as the criteria to group

highly dependent SNP features as blocks. In each block, we chose the features with the largest

minor allele frequency to represent other features in the same block.

Discovery of within-gene epistasis

The main objective of the first stage in GenEpi is to select candidate features from each gene.

In order to extract SNP features for a gene, we used the start and end positions of each gene

from the local UCSC database to split the SNP features after dimension reduction. Since there

are 22,376 genes in the UCSC database, we obtained 22,376 subsets of the SNP features. In

each subset, A SNP feature with the alleles 'A' and 'a' could have three possible genotypes, AA,

Aa and aa, which are used to refer to the pairs of alleles. The pairs of alleles are subsequently

separated into three binary features using one-hot encoding. In order to evaluate epistasis, we

generated interacting features by crossing each pair of genotype features. Considering the false

positive rate and computational complexity, we only focused on pairwise interactions of

epistasis throughout this study. We defined the interaction between two SNPs in following

equation, $\boldsymbol{\alpha_1 SNP_1 + \alpha_2 SNP_2}$ stand for the additive interactions and $\boldsymbol{\alpha_{int(1,2)} SNP_1 \otimes SNP_2}$

represents the synergistic interactions that contain nine terms shown in equation.

$$y \sim \alpha_0 + \sum_{m \in \{AA, Aa, aa\}} \alpha_{1,m} SNP_{1,m} + \sum_{m \in \{AA, Aa, aa\}} \alpha_{2,m} SNP_{2,m} + \sum_{m,n \in \{AA, Aa, aa\}} \alpha_{1,m,2,n} SNP_{1,m} SNP_{2,n}$$

$$= \alpha_0 + \alpha_1 SNP_1 + \alpha_2 SNP_2 + \alpha_{int(1,2)} SNP_1 \otimes SNP_2$$

Before modeling each subset of genotype features, two criteria were adopted to exclude low

quality data. The first criterion is that the genotype frequency of a feature should exceed 5%,

where the genotype frequency means the proportion of genotype among the total samples in the

dataset. The second criterion is regarding the association between the feature and the phenotype.

If the phenotype is dichotomous or categorical, the p-value of χ^2 test should be smaller than

0.01. If the phenotype is a real number, we used F-test to estimate the correlation between the

feature and the phenotype, and the p-value should be smaller than 0.01. In the end, a gene may

have multiple SNPs. The general form of the linear model for a gene with k SNPs is defined as

following equation, which be termed two-element combinatorial encoding.

206
$$y \sim \alpha_0 + \sum_{i=1}^{k} \alpha_i SNP_i + \sum_{i=1}^{k} \sum_{j=1 \cap j \neq i}^{k} \alpha_{int(i,j)} SNP_i \otimes SNP_j$$

We conducted Randomized Lasso and L1-penalized Logistic regression for modeling each gene.

The first advantage is that the L1-penalized model prefers solutions with a smaller number of

features, which effectively reduces the effect of feature dependency. The second advantage is

by using randomization techniques [36], we can largely reduce the false positive rate.

Randomized regression works by resampling and remodeling the training set hundreds or

thousands of times and then picking out the features that are repeatedly selected across

randomization. In this study, we executed this randomization 500 times, and the features

selected by randomization would be retained for the next stage.

Discovery of cross-gene epistasis

In the second stage, we used the features selected by STAGE 1 to generate cross-gene epistasis

features and then applied the same selection procedure described in Section 2.3 to find the

cross-gene epistasis that are associated with the phenotype. The procedures were slightly

modified here. Since we only focused on pairwise interactions, instead of using the entire

features we selected in STAGE 1, we only used single-SNP features to generate cross-gene

epistasis features. Also, we used the genotype frequency and the p-values of χ^2 test or F-test to

control the quality of features and to avoid overfitting. Nevertheless, the p-value of each feature

in this stage should be smaller than 10^{-5}. All of the features from different genes would be

merged for modeling cross-gene epistasis. We conducted Lasso and L1-penalized Logistic

regression for modeling, and the randomization techniques were used once again to select the

final genotype feature set. Since the phenotype may also be affected by environmental factors,

after determining the final set of genotype features, we included the environmental factors such

as clinical assessments for constructing the final model. To obtain the final model, we used

random forests with 1,000 decision trees as the ensemble algorithm. Subsequently, the final

model was evaluated through a process called double cross validation (CV). In the external loop

of double CV, all the instances were divided into two subsets to serve as training and

independent test sets. In this study, we used 2-fold CV and leave-one-out CV (LOO CV) in

external loop for evaluation. In the internal loop, we also used 2-fold CV for model selection.

Materials

This study applied GenEpi on an AD cohort, which was used in Alzheimer's disease Dream

Challenge [28], In total, the cohort consists of 767 participants, who were healthy elderly, mild

cognitive impairment (MCI) and AD patients from the Alzheimer's Disease Neuroimaging

Initiative (ADNI) database. The 767 ADNI participants consist of 241 cognitively normal (CN),

130 Early MCI (EMCI), 273 Late MCI (LMCI) and 123 AD participants. We adopted two types

of features in this study: clinical and genetic data. The clinical assessment includes the 'Mini-

Mental State Examination (MMSE) at baseline' and the '24-month follow-up MMSE'. Other

clinical measures included 'diagnosis at baseline', 'age at baseline' (Age), 'years of education

at baseline' (Education), 'gender' (Gender), and '*APOE* genotype'. All the genetic data has been

pre-processed by the organizers that held the challenge [28]. The genetic data were genotyped

using the Illumina Human610-Quad BeadChip and Illumina HumanOmniExpress BeadChip.

The multidimensional scaling analysis was applied by PLINK using HAPMAP3 to ensure that

samples are within the cluster of European populations. Subsequently, the data were imputed

according to the 1,000 genome haplotypes. After imputation, there were 12,809,667 genotype

features in total. Because GenEpi can be applied to solve both classification and regression

problems, this study illustrated both applications of GenEpi on the ADNI cohort. In predicting

the diagnosis of AD, we used 364 participants, of which the clinical diagnosis are CN or AD,

to predict which samples are control subjects or the AD patients. For predicting the progression

of AD, we used the MMSE difference between '24-month follow-up MMSE' and 'MMSE at

baseline' as the predicting targets.

Experimental Results

This section includes four parts. In the first part, we applied GenEpi and other algorithms for

detecting epistasis, including FastEpistasis [18], BOOST[19] and ReliefF [23, 37] on

simulation data for validation and comparison. In the second part, we applied GenEpi on the

ADNI dataset to categorize each sample as control subjects or AD patients, evaluated by

precision, recall, accuracy and F1 score ($2 \times$ (precision $\times$ recall) / (precision + recall)). In the

third part, we used the same dataset to predict the progression of AD, which is being defined as

the MMSE difference between the baseline and the follow-up after 24 months, and the

performance was assessed using Pearson correlation and Spearman correlation between the

predicting results and the MMSE differences. In the final part, we compared GenEpi with other

algorithms we mentioned in second part on the ADNI dataset in terms of computing time and

prediction performance on real data.

Experiments on simulation data

In this section, we applied different algorithms on simulation data for validation and

comparison. All of the simulation datasets are generated by the simulator GAMETES [38],

which is publicly accessible on the web site

https://popmodels.cancercontrol.cancer.gov/gsr/packages/gametes/. We designed two types of

simulation datasets: basic and complex models. The 'Model 1', 'Model 2' and 'Model 3' are

simulation datasets with the basic model, which means that each dataset only contained one

epistasis consisting of a SNP pair. All of the basic-model datasets are in the same setting as

follows: #individuals = 2000, case/control ratio = 1, #SNPs = 100, #replicates = 100, minor

allele frequency of target SNPs = 0.2, and heritability = 0.2. The complex model means one

dataset contains multiple epistasis from different SNP pairs. In other words, the 'Combined

Model 1+2+3' is a complex model dataset containing three epistasis from the previous three

basic models. Fig. 2 shows the results of these four simulation datasets. Fig. 2(a) shows that the

ranking of the target epistasis reported by GenEpi in 300 replicates of the three basic-model

datasets are always ranked the first. For FastEpistasis and BOOST, the medians of the ranking

of the target epistasis are one but there existed exceptions, the number of the excep-tions of

FastEpistasis and BOOST in 300 replicates of three basic models are 23 and 20, respectively,

which means that the ranked lists contain they will obtain many false positive epistasis when

only one target epistasis should be concerned. For the result of the complex model dataset in

Fig. 2(b), the advantage of GenEpi over other algorithms is more obvious. In the 100 replicates

tested, GenEpi reported the three target epistasis as the top three important features every time.

In contrast, FastEpistasis and Boost did not report that three target epistasis as the top three

important features. On the other hand, since the Python package scikit-rebate [37] that we used

for implementing ReliefF only reports the importance of individual SNPs instead of the scores

for epistasis (SNP pairs), we listed the medians of ReliefF's ranking for each target SNP in

Table 1. Table 1 reveals that ReliefF can detect the SNPs in the target epistasis in the basic

models, but results in some false positives in the complex model dataset.

We believed that the advantage of GenEpi is owing to the proposed two-element combinatorial

encoding of the genotype features and the randomization techniques. In contrast with other

statistical algorithms such as FastEpisasis and BOOST, which only evaluate the epistasis of a

SNP pair one at a time, GenEpi looks for the epistasis among all of the SNP combinations.

Moreover, the false positives among the epistasis can be filtered out by resampling and

remodeling the dataset hundreds or thousands of times. To evaluate the effect of randomization,

we applied both randomized regression and original regression on the complex model dataset

to compare the false positive rates. As shown in Fig. 3. I, randomized regression can reduce the

mean false positive rate effectively and minimized the variance of false positive rate as well.

Classification – Classifying AD patients

In predicting control subjects or AD patients, we applied GenEpi on the 364 samples with CN

(as control) or AD. After dimensionality reduction, 12,102,888 out of the 12,809,667 SNPs in

the ADNI dataset were retained, and 4,916,249 of them are located in 20,206 genes (Table 2).

In the step 4 of selecting epistasis, there are 34,689 genetic features selected and 765 of them

are single SNP features, while the other 33,924 are epistasis features within genes. The final

model contained 3 clinical features (Age, Gender, and Education) and 14 genetic features,

including 24 SNPs from 12 genes. These features contained two single SNP features, 11 within-

gene epistasis features and one cross-gene epistasis feature.

To validate the effectiveness of the genetic features extracted by GenEpi, we compared different

subsets of the features in Table 3. The performance of the model considered only clinical

features outperform that considered only genetic features. When we combined both kind of

features, the performance improved, showing that added genetic features is effective. To reduce

the redundancy, we removed '*APOE* genotype', which can be replaced by the genetic features

selected by GenEpi. We also removed 'MMSE at baseline', which is a clinical assessment to

measure the cognitive impairment. We argued that it does not make sense to use 'MMSE at

baseline' as a feature to predict the status of AD at the moment of diagnosis. The last two

columns of Table 3 showed that the model that used both genetic and clinical features without

APOE genotype and 'MMSE at baseline' is significantly better than the one that considered

only clinical features.

We listed the statistical significance of the selected genetic features in Table 4. The first column

lists each feature by its RSID (Rapid Stain Identification Series) and the genotype (denoted as

RSID_genotype), the pairwise epistasis features are represented using two SNPs. The last

column describes the genes where the SNPs are located according to the genomic coordinates.

We used a star sign to denote the epistasis between genes. Here, only the feature

(rs3130614_BB, rs41276317_AB) is cross-gene epistasis (for MICB and TOB2). The weights

in the second column were extracted from the linear model we defined in Section 2.4. The signs

of the weights indicate if a feature is a causal or protective genotype. On the other hand, the

odds ratio significantly away from 1 also indicates whether the features are potential causal or

protective genotypes. Since low genotype frequency may cause unreliable odds ratios, we also

listed this information in the table. Lastly, the p-value of the χ^2 test showed that these features

are significantly associated with the phenotype. Figure 5 are the bar plots for the features

selected by GenEpi shown in Table 4. We took one single-SNP and the cross-gene epistasis

features as examples. Since each SNP pair has nine SNP combinations, we used the color 'light

blue' to highlight the features selected by GebEpi. We considerd that a good feature can be

judged by two conditions. The first one is that the difference between the amount of case and

control subjects should be large, which suggests that the case and control subjects can be

distinguished by this feature. The second one is that the genotype frequency of this feature

cannot be too small, which suggests that the occurrences of this feature are not only supported

by a small number of samples. Fig. 5(a) reveals that rs429358_AA is a good feature, because

the number of samples in the light blue cell is about 60 percent of the whole dataset and the

proportion of control versus case is 4.425:1. This feature is actually a single SNP feature located

on *APOE*. Moreover, it is a well-known SNP for determining the allele type of *APOE*. Fig. 5(b)

showed all combinations of the SNPs rs3130614 and rs41276317. Even though the number of

samples in the light blue cell is not particularly large, it can still be considered as a good feature

with following explanation. By examining the cells in this figure, two of them are empty cells,

one of them has only one sample, and another five cells have more control subjects than cases

in the cells. Since the proportion of control subjects and cases of this dataset is 2:1, the samples

seem to be randomly distributed into these five cells. In contrast, the proportion of the control

subjects and cases is 1:8 in the light blue cell in the opposite, which indicates the SNP pair with

such combination can be considered as a potential causal genetic variant for AD. These results

show that GenEpi has the ability to identify good features.

Regression – Predicting the progression of AD

In predicting the progression of AD, we applied GenEpi on all of the samples from ADNI. For

this task, the 'diagnosis at baseline' and 'MMSE at baseline' were also adopted as clinical

features. After LD dimension reduction, the original 12,809,667 SNPs decreased to 12,507,343,

and 5,086,639 of them are located on 20,206 genes (Table 2). After selecting epistasis by Step

4, there are 56,427 genetic features left, and 1302 of them are single SNP features, while the

other 55,125 are epistasis features in a single gene. The proposed model contained eight clinical

features (Age, Gender, Education, CN, EMCI, LMCI, AD, and MMSE at baseline) and four

genetic features, which included seven SNPs from six genes, and all of the four genetic features

are cross-gene epistasis. We compared different subsets of features in Table 5. The performance

of the model that only included genetic features is lower than the one that only included the

clinical features. However, when we combined both of them, the performance improved, similar

to the situation observed in the task of classification. It shows that the genetic features are

effective. Again, the *APOE* genotype in the clinical feature set is a genetic feature in essence.

So, we removed it from the list of clinical features. The last two columns showed that even

though we removed the *APOE* genotype, the model with genetic features are still better than

the model without the genetic features. The weights in Table 6 were extracted from the linear

model we described in Section 2.4. The signs of the weights indicate whether the feature is a

causal or protective gene. We applied student's t-test on the samples with the combination of

the selected genetic features (Fig. 6), and all of the p-values of t-test are significant ($< 10^{-11}$).

These results show that GenEpi has the ability to identify good features in the regression task,

too.

Comparison with different algorithms

In this section, we compared GenEpi with other algorithms for detecting epistasis, including

FastEpistasis [18], BOOST [19] and ReliefF [23] in terms of computing time and prediction

performance. We used Microsoft Azure E32 v3 as the computing resource, which contains 32

CPUs and 256 GB RAM. Since the second generation of PLINK has imported FastEpistasis

and BOOST, we used PLINK to test these two algorithms. For ReliefF, we employed a Python

package called scikit-rebate [37] for implementation. Among these algorithms, only

FastEpistasis can afford the computation of the whole set of SNPs. In this regard, 12,809,667

SNPs were used by FastEpistasis (Table 7). However, since FastEpistasis only supports the

classification trial in PLINK, it was skipped in the task of regression. On the other hand, GenEpi

only focuses on the SNPs in the gene regions. In this regard, the number of input SNPs for

estimating epistasis reduced to 4,916,249 and 5,086,639 in classification and regression tasks,

respectively. BOOST took the same subsets of SNPs as GenEpi (Table 7). When taking the

same subset of SNPs as GenEpi and BOOST, ReliefF still caused memory errors. Therefore,

we used the subsets of SNPs that selected by STAGE 1 of GenEpi as the input of ReliefF, which

are 33,868 and 69,842 in classification and regression tasks, respectively. We selected the top

15, 30, 45 and 60 rankings from the results of these algorithms for comparing the prediction

performance. Since different algorithms adopted different strategies to estimate the importance

of the feature, we adopted different formulas defined in Section 2 to generate features for

prediction. FastEpistasis and BOOST used χ^2 statistics to rank the SNP pairs. For these two

algorithms, we put the top SNP pairs into Equation 2 to generate pairwise interactions for each

pair. ReliefF reported significance scores for each SNP, but we cannot find a way to know the

relations between these SNPs. Therefore, we used Equation 3 to generate the interactions for

all the combinations of SNP pairs. Finally, we used logistic and lasso regression to build the

models for classifying AD patients and predicting the progression of AD for comparison. Table

7 show that GenEpi is the most efficient method, which only cost 9.95 and 28.38 CPU-days in

classification and regression tasks, respectively. In both classification or regression tasks,

GenEpi had the best prediction performance despite the fact that GenEpi only uses the subset

of SNPs from the final model. GenEpi shows that the time needed for identifying epistasis can

be drastically reduced, without compromise to the performance. We provided the ROC curves

for the classification task in Fig. 7, and it shows that GenEpi achieved the best performance in

both training and double 2-fold CV procedures, of which the area under the curve (AUC) are

0.95 and 0.85, respectively.

Discussion

The results in the previous section revealed the power of GenEpi to identify phenotype-

associated epistasis efficiently. GenEpi selected 14 features from 12 genes to categorize patients

with AD and four features from six genes to predict the progression of AD. Since AD is a

chronic neurodegenerative disease, our findings would be supported if the gene identified by

GenEpi are expressed in brains. We downloaded the datasets of the GTEx Project [39] to inspect

the gene expression of these genes in different tissues, as shown in Figure. 6. Among the 18

genes selected by GenEpi, 13 have high expression level in the brain tissues. In addition, five

genes, *VSNL1*, *PRKAG2-AS1*, *SYT6*, *CLEC1A* and *LINC00299*, have a similar expression

pattern with *APOE*. These 18 genes are categorized as cross-gene epistasis, single-gene

epistasis and single-SNP features based on the feature types selected by GenEpi. The detail

discussion about these three types of feature will be describe in following sections.

Cross-gene epistasis

GenEpi picked out five cross-gene epistasis in total, which include *CLEC1A * GPR142,*

*CLEC1A * PRKAG2-AS1, IFT20 * RPL32, MFSD6L * PRKAG2-AS1* and *MICB * TOB2*. We

found several direct and indirect evidence to demonstrate that these interactions might have

true association with AD. Liddelow et al. found that neurotoxic reactive astrocytes are induced

by activated microglia [40] and thus are implicated in AD [41]. The expression of *CLEC1A*, a

phagocytic receptor gene, increased in aged microglia, indicative of a pro-inflammatory status

[42]. Based on HaploReg [43], the enhancer of *GPR142* is DNAse sensitive in primary

astrocytes. Together evidence suggests that *CLEC1A* and *GPR142* are expressed in two

inflammatory cells strongly related in AD. Li et al. found that *PRKAG2-AS1* and *PRKAG2* were

differentially regulated [44], suggesting a possible negative regulatory role of *PRKAG2-AS1* on

the *PRKAG2* mRNA level. The *PRKAG2*, AMP-activated protein kinase subunit gamma-2, has

a polymorphism associated with cognitive impairment as well as diabetes in old age [45].

AMPK activation is also related to cognitive impairment in old age, neuroinflammation, and

neurodegeneration [46]. Together, the epistasis between *PRKAG2-AS1* and *CLEC1A* raises an

interesting possibility that *PRKAG2-AS1* may functionally interact with *CLEC1A* through

AMPK regulated neuroinflammation.

We did not find direct link between *RPL32* and AD, but by the data presented in HaploReg

there is another gene called *IQSEC1* that is located in the same LD block with *RPL32*. *IQSEC1*

is expressed in neuron and physically binds to *APOE*, but the implication of this interaction in

AD is not clear. *IFT20* is an essential component of primary cilia that appears to be important

in the survival of dentate granule cell, which plays a key role in memory formation and related

to AD [47]. These studies showed that *IFT20* and *IQSEC1* have been independently implicated

in AD and our results suggest they might together maintain an intact primary cilia that functions

critically in AD. MIC proteins are induced on the cell surface by stress and are ligands of a

common activatory natural killer-cell receptor (*NKG2D*) [48]. They thus have roles in

antimicrobial defense, tumor suppression and autoimmune-like diseases [49]. Quiroga et al. has

found that the human MHC class I chain-related genes (*MICA* and *MICB*) are located within

the HLA region that has stratified association with AD in the cohort of the Oxford Project to

Investigate Memory and Ageing (OPTIMA) even though the association did not survive the

correction for multiple testing [50]. *TOB2* has been reported as a most up-regulated gene [51]

by expression analysis on multiple AD expression datasets from the GEO database to identify

significant genes associated with electrophysiological pathways and attempted determination

of interconnected canonical molecular pathways. The interaction of *MICB* and *TOB2* suggests

a not-yet-explored biological process in AD pathogenesis. However, the significant associations

of these genes in genome-wide association studies associated with immune system diseases and

neurological diseases (see GWAS Catalog, 22561518, 29083406, 28540026), warrant future

investigation. Among all the cross-gene epistasis selected by GenEpi, only one gene *MFSD6L*

cannot find the association with AD so far.

Single-gene epistasis

There are several possible modes of action accounting for intramolecular SNP-SNP interactions

identified in this study. First: a synergistic regulation of transcription. For example, rs12095538

resides in the eQTL of STY6 and rs2774308 resides in a DNAse region of STY6. It is very

possible that variants in both rs12095538 and rs2774308 synergistically affect the expression

of STY6. SYT6 encodes protein control neurotransmitter release. It has been shown that the

expression of SYT6 is significantly down regulated in AD [52], supporting the causative role

of the inherited rs12095538 and rs2774308 haplotype in AD progression. Likewise, rs56233035

and rs3678 reside in the LD blocks containing the eQTL and DNAse region of CACNA1E,

respectively. CACNA1E encodes a subunit of CaV1.2 calcium channel. Its expression level in

reactive astrocytes is associated with amyloid-β plaques formation in an AD mouse model [53].

Another examples are the interactions between rs12926153 and rs12922908, and rs9652600

and rs12922908 that are within DNAse regions controlling the expression of CLEC16A in brain.

CLEC16A encodes a lectin receptor involved in inflammation process and is differentially

expressed in AD brains [54]. Lastly, rs2052573 and rs34580133 reside in DNAse regions within

the LINC00299 and thus may synergistically control the expression of this long non-coding

RNA that has been implicated in brain development [55].

Second: a synergestic interaction between transcriptional and post-transcriptional regulation.

For example, rs9344977, in an intron of BACH2 containing cis-regulatory elements, may affect

the expression of BACH2, whereas rs56148686 in another intron of BACH2 without any cis-

regulatory element. We suspect that rs56148686 may interact with s9344977 through a post-

transcriptional mechanism such as RNA stability, RNA splicing, or microRNA binding.

BACH2 encodes a transcription factor involving cellular responses toward oxidative stress.

Interestingly, BACH2 is upregulated in cultured human neuroblastoma cells upon exposure to

Alzheimer Amyloid β [56]. Similar mechanisms may apply to interaction pairs rs12189429-

rs6881360 and rs12187423-rs6881360 that presumably regulating the abundance of

ADAMTS12 mRNA. The importance of ADAMTS12 in AD is supported by another

independent study using same ANDI dataset [57]. Third: an intramolecular SNP pairs

modulating the expression of two separate neighboring genes. For example, rs11675339 and

rs2710687 of VSNL1 appear to control the expression of GEN1 and SMC6 respectively. We

also found that VSNL (also known as VILIP-1) has strong evidences to AD [58-60]. GEN1 and

SMC6 play pivotal roles in repairing double-strand breaks of genomic DNA. Interestingly,

rDNA instability, a result of failures to repair double-strand breaks of genomic DNA, has been

implicated in AD [61]. Most of the single-gene epistasis selected by GenEpi can be explained

by these three possible modes and only two of the significance SNP-SNP interactions are not

immediately clear at this moment. For example, rs12366151 resides in the eQTL of MICALCL,

an AD associated gene [62] but the interacting rs10831829 resides in the intron of PARVA, a

gene not-yet-reported to be associated with AD. Similarly, rs2421701 and rs200512701 reside

in two eQTLs regulating TNKS2 expression. However, the ADP-ribose polymerase

functionally contributes to AD is not clear.

Single-gene epistasis

There are only two single-SNP features and both of them are located in *APOE*, which is a well-

known causal gene of AD, revealing that GenEpi is an effective tool to identify disease-causing

genes. Moreover, GenEpi successfully selected out the SNP rs429358, which determines the

allele type of *APOE* with rs7412.

Conclusion

While GenEpi has shown its ability to identify epistasis efficiently, it might still has the

following limitations. Firstly, GenEpi can only detect pairwise interactions. Second, GenEpi is

a memory-consuming package, which might cause memory errors when calculating the

epistasis of a gene containing a large number of SNPs. We recommend that the memory for

running GenEpi should be over 256 GB. Finally, a small sample size may lead overfitting,

which forces us to use strict thresholds during feature selection. In this way, GenEpi delivers a

high precision rate, but might suffer having false negatives. This implies different GWAS data

might detect different sets of true positives. In summary, the results of this study demonstrated

that GenEpi is a promising software package to identify causal SNPs and epistasis in GWAS,

and it can be further used to predict the phenotypes. With the demonstrated efficiency, GenEpi

is a powerful tool to explore gene-gene interactions that underlie complex diseases.

Tables

Table 1. The medians of the ranking of the target epistasis for ReliefF

	SNP 1	SNP2	SNP3	SNP4	SNP5	SNP6
Basic-model	1	2	1	2	1	2
Complex-model	7	8.5	9.5	11.5	11	19.5

534 **Table 2.** The summary of the number of features after feature selection in the classification and regression tasks, respectively.

Step	Classification	Regression	Unit
0 (raw data)	12,809,667	12,809,667	SNP
1	22,376	22,376	Gene
2	12,102,888	12,507,343	SNP
3	4,916,249	5,086,639	SNP
	20,206	20,206	Gene
4	44,865	69.843	SNP
	7,933	9,561	Gene
	34,689	56,427	Feature
5	24	7	SNP
	12	6	Gene
	14	4	Feature

536 **Table 3.** Comparison of different subsets of features in predicting control subjects or AD patients.

		Genetic	Clinical	Genetic + Clinical	Clinical - APOE - MMSE	Genetic + Clinical - APOE - MMSE
Training	Precision	0.9633	1.0000	1.0000	0.9658	1.0000
	Recall	0.8537	1.0000	1.0000	0.9187	1.0000
	F1 Score	0.9052	1.0000	1.0000	0.9417	1.0000
	Accuracy	0.9396	1.0000	1.0000	0.9615	1.0000
2-fold CV	Precision	0.7748	0.9280	0.9675	0.3774	0.7699
	Recall	0.6992	0.9431	0.9675	0.3252	0.7073
	F1 Score	0.7350	0.9355	0.9675	0.3493	0.7373
	Accuracy	0.8297	0.9560	0.9780	0.5907	0.8297
LOO CV	Precision	0.8100	0.9286	0.9600	0.3738	0.7596
	Recall	0.6585	0.9512	0.9756	0.3252	0.6423
	F1 Score	0.7265	0.9398	0.9677	0.3478	0.6960
	Accuracy	0.8324	0.9588	0.9780	0.5879	0.8104

537 'Genetic' is short for genetic features; 'Clinical' is short for clinical features. F1 Score = 2 × (Precision × Recall) / (Precision + Recall);
538 'Training' stands for the process of a single-loop CV; '2-fold CV' means that 2-fold CV was used in the external loop of double CV; 'LOO
539 CV' means that LOO CV was used in the external loop of double CV.

541   **Table 4.** The statistical significance of genetic features selected by GenEpi in predicting patients with AD.

Selected SNPs (RSID)	Weight	Odds Ratio	χ^2-test P-value	Genotype Frequency	Gene
rs3130614_BB, rs41276317_AB	3.16	19.23	1.42E-09	0.0742	*MICB*TOB2*
rs12095538_BB, rs2774308_AB	2.41	7.69	6.87E-07	0.0824	*SYT6*
rs12926153_AB, rs12922908_AA	1.18	4.83	6.89E-07	0.1511	*CLEC16A*
rs9652600_AB, rs12922908_AA	0.94	4.83	6.89E-07	0.1511	*CLEC16A*
rs9344977_BB, rs56148686_AB	1.94	4.32	1.14E-06	0.1813	*BACH2*
rs429358_AA	-2.01	0.17	1.73E-06	0.5962	*APOE*
rs56233035_AB, rs3678_AB	2.26	10.16	1.91E-06	0.0604	*CACNA1E*
rs11675339_AA, rs2710687_AA	2.32	3.94	3.55E-06	0.1923	*VSNL1*
rs12189429_BB, rs6881360_AA	1.36	4.34	3.65E-06	0.467	*ADAMTS12*
rs12187423_BB, rs6881360_AA	0.58	4.34	3.65E-06	0.467	*ADAMTS12*
rs10831829_BB, rs12366151_AA	3.48	9.50	4.90E-06	0.0577	*PARVA*
rs2052573_BB, rs34580133_AB	1.80	4.08	5.00E-06	0.1648	*LINC00299*
rs2421701_AB, rs200512701_AB	1.82	4.12	5.29E-06	0.1593	*TNKS2*
rs769449_AA	-1.19	0.16	8.42E-06	0.6648	*APOE*

542   The sign '*' between two gene symbols indicates cross-gene epistasis.

544 **Table 5.** Comparison of different subsets of features in predicting the progression of AD.

		Genetic	Clinical	Genetic + Clinical	Clinical - APOE	Genetic + Clinical - APOE
Training	Pearson	0.4641	0.9582	0.9613	0.9533	0.9600
	Spearman	0.2939	0.9277	0.9375	0.9227	0.9385
	AVG(P+S)	0.3790	0.9430	0.9494	0.9380	0.9493
2-folds CV	Pearson	0.4088	0.5014	0.5771	0.4896	0.5702
	Spearman	0.2286	0.4517	0.4915	0.4289	0.4776
	AVG(P+S)	0.3187	0.4765	0.5343	0.4593	0.5239
LOO CV	Pearson	0.4123	0.4922	0.5806	0.4741	0.5732
	Spearman	-0.1032	0.4441	0.4960	0.4197	0.4854
	AVG(P+S)	0.1545	0.4682	0.5383	0.4469	0.5293

545 'Genetic' is short for genetic features; 'Clinical' is short for clinical features; 'Pearson' means Pearson's Correlation; 'Spearman' means
546 Spearman's Correlation; 'AVG(P+S)' means the average of Pearson and Spearman correlations; 'Training' stands for the process of a single-
547 loop CV; '2-fold CV' means that 2-fold CV was used in the external loop of double CV; 'LOO CV' means that LOO CV was used in the
548 external loop of double CV.

550 **Table 6.** The statistical significance of genetic features selected by GenEpi in predicting the progression of AD.

Selected SNPs (RSID)	Weight	T-test P-value	Genotype Frequency	Gene
rs2306894_BB, rs11653674_BB	-1.2045	5.53E-13	0.3025	*CLEC1A*GPR142*
rs2306894_BB, rs4078431_BB	-0.8244	4.55E-13	0.0574	*CLEC1A*PRKAG2-AS1*
rs7221462_BB, rs11721292_AA	1.3215	3.09E-11	0.8905	*IFT20*RPL32*
rs2242373_AA, rs4078431_BB	-1.2188	3.51E-13	0.0756	*MFSD6L*PRKAG2-AS1*

551 '*' sign between two gene symbols indicates cross-gene epistasis.

553 **Table 7.** The comparison of different algorithms.

Algorithm	# Input SNP	Time Cost	Top 15	Top 30	Top 45	Top 60
Classification – Classifying AD patients						
GenEpi	4,916,249	9.95	0.76	0.72	0.71	0.68
BOOST	4,916,249	2157.6	0.31	0.24	0.30	0.37
ReliefF	33,868	0.11	0.52	0.48	0.45	0.46
FastEpistasis	12,809,667	836.8	0.62	0.61	0.60	0.59
Regression – Predicting the progression of AD						
GenEpi	5,086,639	28.38	0.35	0.48	0.46	0.50
BOOST	5,086,639	327.12	0.08	0.08	0.05	0.04
ReliefF	69,842	1.05	0.16	0.22	0.23	0.26

554 Time is the time spent on identifying the epistasis, which was measured by single CPU time in days. The values in column top 15, top 30,
555 top 45 and top 60 are the 2-fold CV scores. The 2-fold CV scores in classifying AD patients and predicting the progression of AD are the F1
556 scores and the average of Pearson and Spearman correlation, respectively.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

This work has been supported by the Ministry of Science and Technology of Taiwan grant 105-2911-I-002-566, 105-2221-E-002-129-MY3 and 103-2627-M002-015.

Authors' contributions

Y.-C.C. initiated the study, designed the analysis procedures, performed the analysis and wrote the manuscript. J.-T.W. finished the literature survey of the epistasis that selected by GenEpi and wrote the discussion. C.-Y.C. is the main advisor of Y.-C.C., guided this research on right way and provided ideas to optimize the methods. Y.-C.C, M.-Y.H., Y.-A.T, C.-Y.C. and Y.-J.O. were team members of Alzheimer's disease Dream Challenge. C.-Y.C., Y.-J.O., P.-H.H. K.M.G. and S.W.Y. commented on the draft and revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Data collection and sharing for this project was funded by the ADNI (National Institutes of Health Grant U01 AG024904). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Abbott; Alzheimer's Association; Alzheimer's Drug Discovery Foundation; AmorfixLife Sciences Ltd.; AstraZeneca; Bayer HealthCare; BioClinica, Inc.; Biogen Idec Inc.;

Bristol-Myers Squibb Company; Eisai Inc.; Elan Pharmaceuticals Inc.; Eli Lilly and Company;

F.Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; GE Healthcare;

Innogenetics, N.V.; Janssen Alzheimer Immunotherapy Research & Development, LLC.;

Johnson & Johnson Pharmaceutical Research & Development LLC.; Medpace, Inc.; Merck &

Co., Inc.; Meso Scale Diagnostics, LLC.; Novartis Pharmaceuticals Corporation; Pfizer Inc.;

Servier; Synarc Inc. and Takeda Pharmaceutical Company. The Canadian Institutes of Health

Research is providing funds to support ADNI clinical sites in Canada. Private sector

contributions are facilitated by the Foundation for the National Institutes of Health

(www.fnih.org). The grantee organization is the Northern California Institute for Research and

Education and the study is coordinated by the Alzheimer's Disease Cooperative Study at the

University of California, San Diego. ADNI data are disseminated by the Laboratory for Neuro

Imaging at the University of California, Los Angeles. This research was also supported by NIH

grants P30 AG010129, K01 AG030514 and the Dana Foundation.

Reference

1. Kingsmore SF, Lindquist IE, Mudge J, Gessler DD, Beavis WD: **Genome-wide association studies: progress and potential for drug discovery and development**. *Nature Reviews Drug Discovery* 2008, **7**(3):221-230.

2. Ozaki K, Ohnishi Y, Iida A, Sekine A, Yamada R, Tsunoda T, Sato H, Sato H, Hori M, Nakamura Y *et al*: **Functional SNPs in the lymphotoxin-alpha gene that are associated with susceptibility to myocardial infarction**. *Nat Genet* 2002, **32**(4):650-654.

3. Klein RJ, Zeiss C, Chew EY, Tsai JY, Sackler RS, Haynes C, Henning AK, SanGiovanni JP, Mane SM, Mayne ST *et al*: **Complement factor H polymorphism in age-related macular degeneration**. *Science* 2005, **308**(5720):385-389.

4. Pinero J, Bravo A, Queralt-Rosinach N, Gutierrez-Sacristan A, Deu-Pons J, Centeno E, Garcia-Garcia J, Sanz F, Furlong LI: **DisGeNET: a comprehensive platform integrating information on human disease-associated genes and variants**. *Nucleic Acids Res* 2017, **45**(D1):D833-D839.

5. McCarthy MI, Abecasis GR, Cardon LR, Goldstein DB, Little J, Ioannidis JP, Hirschhorn JN: **Genome-wide association studies for complex traits: consensus, uncertainty and challenges**. *Nat Rev Genet* 2008, **9**(5):356-369.

6. Eichler EE, Flint J, Gibson G, Kong A, Leal SM, Moore JH, Nadeau JH: **Missing heritability and strategies for finding the underlying causes of complex disease**. *Nat Rev Genet* 2010, **11**(6):446-450.

7. Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, McCarthy MI, Ramos EM, Cardon LR, Chakravarti A *et al*: **Finding the missing heritability of complex diseases**. *Nature* 2009, **461**(7265):747-753.

8. Frazer KA, Murray SS, Schork NJ, Topol EJ: **Human genetic variation and its contribution to complex traits**. *Nat Rev Genet* 2009, **10**(4):241-251.

9. Carlborg O, Haley CS: **Epistasis: too often neglected in complex trait studies?** *Nat Rev Genet* 2004, **5**(8):618-625.

10. Cordell HJ: **Detecting gene-gene interactions that underlie human diseases**. *Nat Rev Genet* 2009, **10**(6):392-404.

11. Easton DF, Pooley KA, Dunning AM, Pharoah PD, Thompson D, Ballinger DG, Struewing JP, Morrison J, Field H, Luben R *et al*: **Genome-wide association study identifies novel breast cancer susceptibility loci**. *Nature* 2007, **447**(7148):1087-1093.

12. Shriner D, Vaughan LK, Padilla MA, Tiwari HK: **Problems with genome-wide**
association studies. *Science* 2007, **316**(5833):1840-1842.

13. Bodmer W, Bonilla C: **Common and rare variants in multifactorial susceptibility to**
common diseases. *Nat Genet* 2008, **40**(6):695-701.

14. Hindorff LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, Collins FS, Manolio
TA: **Potential etiologic and functional implications of genome-wide association loci**
for human diseases and traits. *Proc Natl Acad Sci U S A* 2009, **106**(23):9362-9367.

15. Moore JH, Asselbergs FW, Williams SM: **Bioinformatics challenges for genome-wide**
association studies. *Bioinformatics* 2010, **26**(4):445-455.

16. Marchini J, Donnelly P, Cardon LR: **Genome-wide strategies for detecting multiple**
loci that influence complex diseases. *Nat Genet* 2005, **37**(4):413-417.

17. Wei WH, Hemani G, Haley CS: **Detecting epistasis in human complex traits**. *Nat Rev*
Genet 2014, **15**(11):722-733.

18. Schupbach T, Xenarios I, Bergmann S, Kapur K: **FastEpistasis: a high performance**
computing solution for quantitative trait epistasis. *Bioinformatics* 2010,
26(11):1468-1469.

19. Wan X, Yang C, Yang Q, Xue H, Fan X, Tang NL, Yu W: **BOOST: A fast approach to**
detecting gene-gene interactions in genome-wide case-control studies. *Am J Hum*
Genet 2010, **87**(3):325-340.

20. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar
P, de Bakker PI, Daly MJ *et al*: **PLINK: a tool set for whole-genome association and**
population-based linkage analyses. *Am J Hum Genet* 2007, **81**(3):559-575.

21. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ: **Second-generation**
PLINK: rising to the challenge of larger and richer datasets. *Gigascience* 2015, **4**:7.

22. Moore JH, Williams SM: **New strategies for identifying gene-gene interactions in**
hypertension. *Ann Med* 2002, **34**(2):88-95.

23. Yang P, Ho JW, Yang YH, Zhou BB: **Gene-gene interaction filtering with ensemble**
of filters. *BMC Bioinformatics* 2011, **12 Suppl 1**:S10.

24. Bureau A, Dupuis J, Falls K, Lunetta KL, Hayward B, Keith TP, Van Eerdewegh P:
Identifying SNPs predictive of phenotype using random forests. *Genet Epidemiol*
2005, **28**(2):171-182.

25. Schwarz DF, Konig IR, Ziegler A: **On safari to Random Jungle: a fast**
implementation of Random Forests for high-dimensional data. *Bioinformatics* 2010,
26(14):1752-1758.

26.   Wan X, Yang C, Yang Q, Xue H, Tang NL, Yu W: **MegaSNPHunter: a learning**
672   **approach to detect disease predisposition SNPs and high level interactions in**
673   **genome wide association study**. *BMC Bioinformatics* 2009, **10**:13.
674   27.   Murk W, DeWan AT: **Exhaustive Genome-Wide Search for SNP-SNP Interactions**
675   **Across 10 Human Diseases**. *G3 (Bethesda)* 2016, **6**(7):2043-2050.
676   28.   Allen GI, Amoroso N, Anghel C, Balagurusamy V, Bare CJ, Beaton D, Bellotti R,
677   Bennett DA, Boehme KL, Boutros PC *et al*: **Crowdsourced estimation of cognitive**
678   **decline and resilience in Alzheimer's disease**. *Alzheimers Dement* 2016, **12**(6):645-
679   653.
680   29.   Ma L, Clark AG, Keinan A: **Gene-based testing of interactions in association studies**
681   **of quantitative traits**. *PLoS Genet* 2013, **9**(2):e1003321.
682   30.   Oh S, Lee J, Kwon MS, Weir B, Ha K, Park T: **A novel method to identify high order**
683   **gene-gene interactions in genome-wide association studies: gene-based MDR**. *BMC*
684   *Bioinformatics* 2012, **13 Suppl 9**:S5.
685   31.   Li S, Cui Y: **Gene-centric gene–gene interaction: A model-based kernel machine**
686   **method**. *The Annals of Applied Statistics* 2012, **6**(3):1134-1161.
687   32.   Wu X, Dong H, Luo L, Zhu Y, Peng G, Reveille JD, Xiong M: **A novel statistic for**
688   **genome-wide interaction analysis**. *PLoS Genet* 2010, **6**(9):e1001131.
689   33.   Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D: **The**
690   **human genome browser at UCSC**. *Genome Res* 2002, **12**(6):996-1006.
691   34.   Rosenbloom KR, Armstrong J, Barber GP, Casper J, Clawson H, Diekhans M, Dreszer
692   TR, Fujita PA, Guruvadoo L, Haeussler M *et al*: **The UCSC Genome Browser**
693   **database: 2015 update**. *Nucleic Acids Res* 2015, **43**(Database issue):D670-681.
694   35.   Lewontin RC: **The Interaction of Selection and Linkage. I. General Considerations;**
695   **Heterotic Models**. *Genetics* 1964, **49**(1):49-67.
696   36.   Meinshausen N, Bühlmann P: **Stability selection**. *Journal of the Royal Statistical*
697   *Society: Series B (Statistical Methodology)* 2010, **72**(4):417-473.
698   37.   Urbanowicz RJ, Meeker M, LaCava W, Olson RS, Moore JH: **Relief-Based Feature**
699   **Selection: Introduction and Review**. *arXiv preprint arXiv:171108421* 2017.
700   38.   Urbanowicz RJ, Kiralis J, Sinnott-Armstrong NA, Heberling T, Fisher JM, Moore JH:
701   **GAMETES: a fast, direct algorithm for generating pure, strict, epistatic models**
702   **with random architectures**. *BioData Min* 2012, **5**(1):16.
703   39.   Consortium GT: **The Genotype-Tissue Expression (GTEx) project**. *Nat Genet* 2013,
704   **45**(6):580-585.

40.   Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, Bennett
ML, Munch AE, Chung WS, Peterson TC *et al*: **Neurotoxic reactive astrocytes are**
induced by activated microglia. *Nature* 2017, **541**(7638):481-487.

41.   von Bernhardi R, Ramirez G: **Microglia-astrocyte interaction in Alzheimer's disease:**
friends or foes for the nervous system? *Biol Res* 2001, **34**(2):123-128.

42.   Raj D, Yin Z, Breur M, Doorduin J, Holtman IR, Olah M, Mantingh-Otter IJ, Van Dam
D, De Deyn PP, den Dunnen W *et al*: **Increased White Matter Inflammation in**
Aging- and Alzheimer's Disease Brain. *Front Mol Neurosci* 2017, **10**:206.

43.   Ward LD, Kellis M: **HaploReg: a resource for exploring chromatin states,**
conservation, and regulatory motif alterations within sets of genetically linked
variants. *Nucleic Acids Res* 2012, **40**(Database issue):D930-934.

44.   Li P, Lahvic JL, Binder V, Pugach EK, Riley EB, Tamplin OJ, Panigrahy D, Bowman
TV, Barrett FG, Heffner GC *et al*: **Epoxyeicosatrienoic acids enhance embryonic**
haematopoiesis and adult marrow engraftment. *Nature* 2015, **523**(7561):468-471.

45.   Kim E, Lee SH, Lee KS, Cheong HK, Namkoong K, Hong CH, Oh BH: **AMPK**
gamma2 subunit gene PRKAG2 polymorphism associated with cognitive
impairment as well as diabetes in old age. *Psychoneuroendocrinology* 2012,
37(3):358-365.

46.   Peixoto CA, Oliveira WH, Araujo S, Nunes AKS: **AMPK activation: Role in the**
signaling pathways of neuroinflammation and neurodegeneration. *Exp Neurol* 2017,
298(Pt A):31-41.

47.   Whitfield JF, Chiarini A, Dal Pra I, Armato U, Chakravarthy B: **The Possible Roles of**
the Dentate Granule Cell's Leptin and Other Ciliary Receptors in Alzheimer's
Neuropathology. *Cells* 2015, **4**(3):253-274.

48.   Vivier E, Tomasello E, Paul P: **Lymphocyte activation via NKG2D: towards a new**
paradigm in immune recognition? *Curr Opin Immunol* 2002, **14**(3):306-311.

49.   Collins RW: **Human MHC class I chain related (MIC) genes: their biological**
function and relevance to disease and transplantation. *Eur J Immunogenet* 2004,
31(3):105-114.

50.   Quiroga I, Lehmann DJ, Barnardo MC, Fuggle S, Cortina-Borja M, Warden DR, Smith
AD: **Association study of MICA and MICB in Alzheimer's disease**. *Tissue Antigens*
2009, **74**(3):241-243.

51.   Mirza Z, Rajeh N: **Identification Of Electrophysiological Changes In Alzheimer's**
Disease: A Microarray Based Transcriptomics And Molecular Pathway Analysis

Study. *CNS Neurol Disord Drug Targets* 2017.

52.   Saura CA, Parra-Damas A, Enriquez-Barreto L: **Gene expression parallels synaptic**
excitability and plasticity changes in Alzheimer's disease. *Front Cell Neurosci* 2015,
9:318.

53.   Daschil N, Obermair GJ, Flucher BE, Stefanova N, Hutter-Paier B, Windisch M,
Humpel C, Marksteiner J: **CaV1.2 calcium channel expression in reactive astrocytes**
is associated with the formation of amyloid-beta plaques in an Alzheimer's disease
mouse model. *J Alzheimers Dis* 2013, **37**(2):439-451.

54.   Porcellini E, Carbone I, Ianni M, Licastro F: **Alzheimer's disease gene signature says:**
beware of brain viral infections. *Immun Ageing* 2010, **7**:16.

55.   Talkowski ME, Maussion G, Crapper L, Rosenfeld JA, Blumenthal I, Hanscom C,
Chiang C, Lindgren A, Pereira S, Ruderfer D *et al*: **Disruption of a large intergenic**
noncoding RNA in subjects with neurodevelopmental disabilities. *Am J Hum Genet*
2012, **91**(6):1128-1134.

56.   Uhrig M, Ittrich C, Wiedmann V, Knyazev Y, Weninger A, Riemenschneider M,
Hartmann T: **New Alzheimer amyloid beta responsive genes identified in human**
neuroblastoma cells by hierarchical clustering. *PLoS One* 2009, **4**(8):e6779.

57.   Wang W, Mandel J, Bouaziz J, Commenges D, Nabirotchkine S, Chumakov I, Cohen
D, Guedj M, Alzheimer's Disease Neuroimaging I: **A Multi-Marker Genetic**
Association Test Based on the Rasch Model Applied to Alzheimer's Disease. *PLoS*
One 2015, **10**(9):e0138223.

58.   Babic Leko M, Borovecki F, Dejanovic N, Hof PR, Simic G: **Predictive Value of**
Cerebrospinal Fluid Visinin-Like Protein-1 Levels for Alzheimer's Disease Early
Detection and Differential Diagnosis in Patients with Mild Cognitive Impairment.
J Alzheimers Dis 2016, **50**(3):765-778.

59.   Kirkwood CM, MacDonald ML, Schempf TA, Vatsavayi AV, Ikonomovic MD, Koppel
JL, Ding Y, Sun M, Kofler JK, Lopez OL *et al*: **Altered Levels of Visinin-Like Protein**
1 Correspond to Regional Neuronal Loss in Alzheimer Disease and
Frontotemporal Lobar Degeneration. *J Neuropathol Exp Neurol* 2016, **75**(2):175-182.

60.   Luo X, Hou L, Shi H, Zhong X, Zhang Y, Zheng D, Tan Y, Hu G, Mu N, Chan J *et al*:
CSF levels of the neuronal injury biomarker visinin-like protein-1 in Alzheimer's
disease and dementia with Lewy bodies. *J Neurochem* 2013, **127**(5):681-690.

61.   Pietrzak M, Rempala G, Nelson PT, Zheng JJ, Hetman M: **Epigenetic silencing of**
nucleolar rRNA genes in Alzheimer's disease. *PLoS One* 2011, **6**(7):e22585.

62.   Rouillard AD, Gundersen GW, Fernandez NF, Wang Z, Monteiro CD, McDermott MG,

Ma'ayan A: **The harmonizome: a collection of processed datasets gathered to serve**

and mine knowledge about genes and proteins. *Database (Oxford)* 2016, **2016**.

Fig. 1. The architecture of GenEpi.

Fig. 2. The modules and data flow in the proposed GenEpi Python package. The first and second modules (STEP 1 and STEP 2) are the two pre-processing steps for downloading the UCSC database and reducing the dimension using LD. The third and fourth modules (STEP 3 and STEP 4) correspond to the STAGE 1 of GenEpi, and the final two modules (STEP 5 and STEP 6) correspond to the STAGE 2 of GenEpi.

Fig. 3. The boxplot for the ranking of the target epistasis in different algorithms. (a) The upper part are the results of three basic-model datasets with one epistasis consisting of a SNP pair. (b) The lower part is the result of the complex-model dataset, which contained three epistasis. The 'S1-S2' means the epistasis between SNP 1 and SNP 2 and so on. The values on the boxplot are the medians of the ranking of the target epistasis.

Fig. 4. The numbers of false positives in randomized and original regression.

Fig. 5. (a) The bar plot for the population of three allele types in rs429358. (b) The bar plot for combinations of SNPs from the cross-gene epistasis feature selected by GenEpi in predicting control subjects or AD patients. The color 'light blue' is used to highlight the features selected by GenEpi.

Fig. 6. The box plots for combinations of SNPs from the genetic features selected by GenEpi in predicting the progression of AD. The y-axes of the plot are MMSE difference between '24-month follow-up MMSE' and 'MMSE at baseline'.

Fig. 7. The ROC curves of different algorithms.

Fig. 8. The heatmap of gene expression in different tissues for the 18 genes selected by GenEpi. The blue box highlights the sub-regions of brain.