GenEpi: Gene-based Epistasis Discovery Using Machine Learning

Yu-Chuan Chang[1], June-Tai Wu[2], Ming-Yi Hong[3], Yi-An Tung[4], Ping-Han Hsieh[1], Sook Wah

Yee[5], Kathleen M. Giacomini[5,6], Yen-Jen Oyang[1], and Chien-Yu Chen[3,*], for the Alzheimer's

Disease Neuroimaging Initiative†

[1]Graduate Institute of Biomedical Electronics and Bioinformatics, National Taiwan University,

Taipei 10617, Taiwan. [2]Department of Dermatology, National Taiwan University Hospital,

Taipei 10002, Taiwan. [3]Department of Bio-Industrial Mechatronics Engineering, National

Taiwan University, Taipei 10617, Taiwan. [4]Genome and Systems biology degree program,

Academia Sinica and National Taiwan University, Taipei 10617, Taiwan. [5]Department of

Bioengineering and Therapeutic Sciences, University of California, San Francisco, San

Francisco, California, USA. [6]Institute for Human Genetics, University of California, San

Francisco, San Francisco, California, USA.

*To whom correspondence should be addressed.

†Data used in preparation of this article were obtained from the Alzheimer's Disease

Neuroimaging Initiative (ADNI) database (adni.loni.ucla.edu). As such, the investigators

within the ADNI contributed to the design and implementation of ADNI and/or provided data

but did not participate in analysis or writing of this report. A complete listing of ADNI

investigators can be found at:

https://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.p

df

Chien-Yu Chen

Department of Bio-Industrial Mechatronics Engineering, National Taiwan University, Taipei

10617, Taiwan.

Phone: +886-2-3366-5334, E-mail: chienyuchen@ntu.edu.tw

Abstract

Genome-wide association studies (GWAS) provide a powerful means to identify associations

between genetic variants and phenotypes. However, GWAS techniques for detecting epistasis,

the interactions between genetic variants associated with phenotypes, are still limited. We

believe that developing an efficient and effective GWAS method to detect epistasis will be a

key for discovering sophisticated pathogenesis, which is especially important for complex

diseases such as Alzheimer's disease (AD). In this regard, this study presents GenEpi, a

computational package to uncover epistasis associated with phenotypes by the proposed

machine learning approach. GenEpi identifies both within-gene and cross-gene epistasis

through a two-stage modeling workflow. In both stages, GenEpi adopts two-element

combinatorial encoding when producing features and constructs the prediction models by

L1-regularized regression with stability selection. The simulated data showed that GenEpi

outperforms other widely-used methods on detecting ground-truth epistasis. As real data is

concerned, this study uses AD as an example to reveal the capability of GenEpi in finding

disease-related variants and variant interactions that show both biological meanings and

predictive power. **Availability:** GenEpi is an open-source python package and available free

of charge only for non-commercial users. The package can be downloaded from

https://github.com/Chester75321/GenEpi, and has also been published on The Python

Package Index.

1. Introduction

Genome-wide association studies (GWAS) is a univariate examination of a genome-wide set of

genetic variants to determine if any single variant is associated with the phenotype of interest

[1]. The first GWAS was published in 2002 [2], and three years later, the most remarkable

GWAS regarding age-related macular degeneration (AMD) was published [3]. Their study

investigated the association of 105,980 single nucleotide polymorphisms (SNPs) with AMD on

96 cases and 50 control subjects. This study showed that the SNPs in the complement factor H

(CFH) gene, including a non-synonymous SNP, are significantly associated with AMD. Up to

2019, there have been more than hundreds of thousands individuals being studied in typical

GWAS protocols, and over 210,498 variant-disease associations between 117,337 SNPs and

10,358 phenotypes have been discovered [4]. These studies demonstrated the potential of

GWAS to identify genetic variants associated with many categories of phenotypes, including

risk for diseases such as various cancers, and variation in therapeutic and adverse response to

drugs. However, the success of univariate GWAS is limited to monogenic phenotypes (e.g.

Mendelian diseases). The impact of variant interactions, also known as epistasis on the

formation of diseases [5] is often underestimated in traditional GWAS analysis [6-8].

A major limitation of traditional GWAS is that it considers only one genetic variant at a time,

and ignores underlying epistasis of variants that might have stronger associations [9].

Researchers have found that GWAS has limitation in identifying the association in complex

diseases [10, 11]. Easton et al. suggested that a number of susceptible loci identified by GWAS

usually have very small effect sizes [12]. Studies have also demonstrated that the existence of

epistasis is an important factor contributing to phenotypes, especially in complex diseases such

as hypertension, diabetes and obesity [11]. Therefore, developing analytical methods to identify

epistasis efficiently is critical to understanding the genetic factors [8, 13], and has attracted a

wide range of research interests in recent years [7, 14].

There are, however, two main challenges to discover epistasis: computational complexity and

statistical power [15]. The first challenge results from the curse of dimensionality. When more

genetic variants are considered, the number of interactions increases exponentially. Based on

the specification of a major commercial technology, Illumina Arrays, a whole-genome array

can investigate over 4 million markers per sample simultaneously. In order to evaluate the

pairwise interactions from this microarray, about 8×10^{12} statistical tests need to be processed.

Even though Marchini et al. have demonstrated that pairwise interactions of 3×10^5 loci is

computationally possible with currently available computational resources, it still remains

challenging when the Illumina Arrays are considered [16]. The second challenge is the issue of

statistical power. Since a huge number of statistical tests are conducted on a limited sample size

with high-dimensional interactions, many false positives arise by random chances. In these

regards, new methods have been developed [11, 17]. Statistical approaches include

FastEpistasis [18] and BOOST [19]; both of them has been included in a well-known GWAS

software called PLINK [20, 21]. Machine learning approaches such as Multifactor

Dimensionality Reduction [22], ReliefF [23], random forest-like algorithms [24-26] and other

methodologies have also been developed for detecting epistasis [17].

Since the biological experiments used to validate these methodologies such as the AMD in

traditional GWAS are still in demand, there are no standard analysis methods for epistasis

despite the rapid improvement in computational performance.In 2016, Murk used FastEpistasis

and BOOST to search SNP-SNP interactions on a huge dataset called Genetic Epidemiology

Research on Adult Health and Aging (GERA) that included 78,486 subjects, but still failed to

detect a significant and replicable interaction after exhaustively searching through 45 billion

possible interactions for 10 complex diseases of interest [27]. Alzheimer's disease (AD) is one

of the most important complex diseases and its pathogenesis, which clearly has a genetic basis,

is still ill-defined. In 2014, Sage Bionetworks held a competition called The Dialogue for

Reverse Engineering Assessments and Methods Challenge (DREAM Challenge) for AD,

which tried to use crowdsourcing to assess the capability of current computational methods to

predict the change in cognitive examination based on genetic data. However, no significant

contribution of genetic features except the *APOE* haplotype to the predictive performance was

observed by any competition teams [28]. To address this problem, this study presents GenEpi, a

package to reveal epistasis related to the phenotype using machine learning and introduces the

application of GenEpi on predicting the diagnosis of AD.

2. Results

This study compared GenEpi with several commonly used algorithms for detecting epistasis,

including FastEpistasis, BOOST and ReliefF. The simulation data demonstrated that GenEpi

outperforms the other methods in ranking the true epistasis as the top one. As real data is

concerned, the results suggest that the epistasis selected by GenEpi has the best predictive

power for diagnosis of AD. The proposed model of predicting AD contains 14 genetic features,

including 24 SNPs from 12 genes that contain the well-known causal gene, APOE. The 2-fold

cross validation (CV) and leave-one-out CV (LOO CV) accuracy of this model are 0.829 and

0.832, respectively. The results on AD demonstrated that GenEpi has the ability to detect the

epistasis associated with the phenotype effectively and efficiently. The released package can be

generalized to largely facilitate the studies of many complex diseases in the near future.

We will demonstrate our experiments in following three parts of this section. In the first part,

we applied GenEpi and other algorithms for detecting epistasis, including FastEpistasis [18],

BOOST [19] and ReliefF [23, 29] on simulation data for validation and comparison. In the

second part, we applied GenEpi on the ADNI dataset to categorize each sample as control

subjects or AD patients, evaluated by precision, recall, accuracy and F1 score ($2 \times$ (precision $\times$

recall) / (precision + recall)). In the final part, we compared GenEpi with other algorithms on

the ADNI dataset in terms of computing time and prediction performance on real data.

Experiments on simulation data

We applied different algorithms on simulation data for validation and comparison. All of the

simulation datasets are generated by the simulator GAMETES [30], which is publicly

accessible on the web site https://popmodels.cancercontrol.cancer.gov/gsr/packages/gametes/.

We designed two types of simulation datasets: basic and complex models. The 'Model 1',

'Model 2' and 'Model 3' are simulation datasets with the basic model, which means that each

dataset contained only one epistasis consisting of a SNP pair. All of the basic-model datasets are

in the same setting as follows: #individuals = 2000, case/control ratio = 1, #SNPs = 100,

#replicates = 100, minor allele frequency of target SNPs = 0.2, and heritability = 0.2. The

complex model means one dataset contains multiple epistasis from different SNP pairs. Here,

the 'Combined Model 1+2+3' is a complex model dataset containing three epistasis from the

previous three basic models.

Fig. 1 provided the results of these four simulation datasets. Fig. 1(a) shows that the ranking of

the target epistasis reported by GenEpi in the 100 replicates of each basic-model dataset are

always ranked as the first. In contrast, for FastEpistasis and BOOST, the medians of the ranking

of the target epistasis among the 100 runs of simulation are one but the averages are not. The

number of failures of FastEpistasis and BOOST in 100 replicates of three basic models are 6, 1,

16 and 5, 1, 14, respectively. For the result of the complex model dataset in Fig. 1(b), the

superiority of GenEpi over other algorithms is more obvious. In the 100 runs of simulation,

GenEpi reported the three target epistasis as the top three important features every time. In

contrast, FastEpistasis and Boost failed to report the three target epistasis as the top three

important features consistently.

When ReliefF was compared, since the Python package scikit-rebate [29] that we used for

implementing ReliefF only reports the importance of individual SNPs instead of the scores for

epistasis (SNP pairs), we listed the medians of ReliefF's ranking for each SNP in the target

epistasis in Table 1. Table 1 reveals that ReliefF can detect the SNPs in the target epistasis in the

basic models, but failed to report the three target epistasis as the top three important features in

the complex model dataset.

The superiority of GenEpi is owing to the proposed two-element combinatorial encoding of the

genotype features and the L1-regularized regression with stability selection. In contrast with

other statistical algorithms such as FastEpisasis and BOOST, which only evaluate the epistasis

of a SNP pair one at a time, GenEpi considers interactions between combinatorial features by

multivariate models. Moreover, the false positives among the epistasis can be filtered out by

resampling and remodeling the dataset hundreds of times. To evaluate the effect of stability

selection, we applied both L1-regularized regression with and without stability selection on the

complex model dataset to compare the number of false positives, which is defined as the

number of non-target epistasis in the final output of GenEpi. As shown in Fig. 2, stability

selection can reduce the mean false positive rate effectively and minimize the variance of false

positive rate as well.

Classifying AD patients

In predicting control subjects or AD patients, we applied GenEpi on the 364 samples with CN

(as control) or AD. After dimensionality reduction, 12,102,888 out of the 12,809,667 SNPs in

the ADNI dataset were retained, and 4,916,249 of them are located in 20,206 genes (Table 2). In

the step 4 of selecting epistasis, there are 34,689 genetic features selected and 765 of them are

single SNP features, while the other 33,924 are epistasis features within genes. The final model

contained 14 genetic features, including 24 SNPs from 12 genes. These features contained two

single SNP features, 11 within-gene epistasis features and one cross-gene epistasis feature. As

shown in Table 3, the 2-fold cross validation (CV) and leave-one-out CV (LOO CV) accuracy

of this model are 0.829 and 0.832, respectively.

We listed the statistical significance of the selected genetic features in Table 4. The first column

lists each feature by its RSID (Reference SNP cluster ID) and the genotype (denoted as

RSID_genotype), the pairwise epistasis features are represented using two SNPs. The last

column describes the genes where the SNPs are located according to the genomic coordinates.

We used a star sign to denote the epistasis between genes. Here, only the feature

(rs3130614_BB, rs41276317_AB) is cross-gene epistasis (for MICB and TOB2). The weights

in the second column were extracted from the linear model we defined in Section 2.4. The signs

of the weights indicate if a feature is a causal or protective genotype, which is consistent with

the corresponding odds ratio. The p-value of the χ^2 test showed that these features are

significantly associated with the phenotype.

Comparison with different algorithms

In this section, we compared GenEpi with other algorithms for detecting epistasis, including

FastEpistasis [18], BOOST [19] and ReliefF [23] in terms of computing time and prediction

performance. We used Microsoft Azure E32 v3 as the computing resource, which contains 32

CPUs and 256 GB RAM. Since the PLINK (version 2.0) has imported FastEpistasis and

BOOST, we used PLINK to test these two algorithms. For ReliefF, we employed a Python

package called scikit-rebate [29] for implementation. Among these algorithms, only

FastEpistasis can afford the computation of the whole set of SNPs. In this regard, 12,809,667

SNPs were used by FastEpistasis (Table 5). On the other hand, GenEpi only focuses on the

SNPs in the gene regions. In this regard, the number of input SNPs for estimating epistasis

reduced to 4,916,249. BOOST took the same subsets of SNPs as GenEpi (Table 5). When

taking the same subset of SNPs as GenEpi and BOOST, ReliefF still caused memory errors.

Therefore, we used the subsets of SNPs that selected by STAGE 1 of GenEpi as the input of

ReliefF, which are 33,868. We selected the top 15, 30, 45 and 60 rankings from the results of

these algorithms for comparing the prediction performance, and used L1-regularized regression

to build the models for classifying AD patients for comparison. Table 5 show that GenEpi is the

most efficient method, which only cost 9.95 CPU-days. Moreover, GenEpi had the best

prediction performance despite the fact that GenEpi only uses the subset of SNPs from the final

model. GenEpi shows that the time needed for identifying epistasis can be drastically reduced,

without compromise to the performance. We provided the ROC curves for the classification

task in Fig. 3, and it shows that GenEpi achieved the best performance in double 2-fold CV

procedures, of which the area under the curve (AUC) is 0.85.

3. Discussion

The results in the previous section revealed the power of GenEpi to identify

phenotype-associated epistasis efficiently. GenEpi selected 14 features from 12 genes to

categorize patients with AD. Since AD is a chronic neurodegenerative disease, our findings

would be supported if the gene identified by GenEpi are expressed in brains. We downloaded

the datasets of the GTEx Project [31] to inspect the gene expression of these genes in different

tissues, as shown in Fig. 4. Among the 12 genes selected by GenEpi, 11 have high expression

level in the brain tissues. In addition, five genes, *CLEC16A, VSNL1*, *SYT6*, *CACNA1E* and

LINC00299, have a similar expression pattern with *APOE*. These 12 genes are categorized as

cross-gene epistasis, single-gene epistasis and single-SNP features based on the feature types

selected by GenEpi. GenEpi detected only one cross-gene epistasis, which is *MICB * TOB2*.

We found several evidence to demonstrate that this interaction might have true association with

AD. MIC proteins are induced on the cell surface by stress and are ligands of a common

activatory natural killer-cell receptor (NKG2D) [32]. They thus have roles in antimicrobial

defense, tumor suppression and autoimmune-like diseases [33]. Quiroga et al. has found that

the human MHC class I chain-related genes (MICA and MICB) are located within the HLA

region that has stratified association with AD in the cohort of the Oxford Project to Investigate

Memory and Ageing (OPTIMA) even though the association did not survive the correction for

multiple testing [34]. TOB2 has been reported as a most up-regulated gene [35] by expression

analysis on multiple AD expression datasets from the GEO database to identify significant

genes associated with electrophysiological pathways and attempted determination of

interconnected canonical molecular pathways. The interaction of MICB and TOB2 suggests a

not-yet-explored biological process in AD pathogenesis. However, the significant associations

of these genes in genome-wide association studies associated with immune system diseases and

neurological diseases (see GWAS Catalog, 22561518, 29083406, 28540026), warrant future

investigation.

About the 11 single-gene epistasis, there are several possible reasons accounting for

intramolecular SNP-SNP interactions identified in this study. The first is a synergistic

regulation of transcription, the second is a synergistic interaction between transcriptional and

post-transcriptional regulation, and the third is an intramolecular SNP pair modulating the

expression of two separate neighboring genes. Most of the single-gene epistasis selected by

GenEpi can be explained by these three possible reasons and only two of the SNP-SNP

interactions are not immediately clear at this moment. First: a synergistic regulation of

transcription. For example, rs12095538 resides in the eQTL of STY6 and rs2774308 resides in

a DNAse region of STY6. It is very possible that variants in both rs12095538 and rs2774308

synergistically affect the expression of STY6. SYT6 encodes protein control neurotransmitter

release. It has been shown that the expression of SYT6 is significantly down regulated in AD

[36], supporting the causative role of the inherited rs12095538 and rs2774308 haplotype in AD

progression. Likewise, rs56233035 and rs3678 reside in the LD blocks containing the eQTL

and DNAse region of CACNA1E, respectively. CACNA1E encodes a subunit of CaV1.2

calcium channel. Its expression level in reactive astrocytes is associated with amyloid-β

plaques formation in an AD mouse model [37]. Another examples are the interactions between

rs12926153 and rs12922908, and rs9652600 and rs12922908 that are within DNAse regions

controlling the expression of CLEC16A in brain. CLEC16A encodes a lectin receptor involved

in inflammation process and is differentially expressed in AD brains [38]. Lastly, rs2052573

and rs34580133 reside in DNAse regions within the LINC00299 and thus may synergistically

control the expression of this long non-coding RNA that has been implicated in brain

development [39].

Second: a synergestic interaction between transcriptional and post-transcriptional regulation.

For example, rs9344977, in an intron of BACH2 containing cis-regulatory elements, may affect

the expression of BACH2, whereas rs56148686 in another intron of BACH2 without any

cis-regulatory element. We suspect that rs56148686 may interact with s9344977 through a

post-transcriptional mechanism such as RNA stability, RNA splicing, or microRNA binding.

BACH2 encodes a transcription factor involving cellular responses toward oxidative stress.

Interestingly, BACH2 is upregulated in cultured human neuroblastoma cells upon exposure to

Alzheimer Amyloid β [40]. Similar mechanisms may apply to interaction pairs

rs12189429-rs6881360 and rs12187423-rs6881360 that presumably regulating the abundance

of ADAMTS12 mRNA. The importance of ADAMTS12 in AD is supported by another

independent study using same ANDI dataset [41]. Third: an intramolecular SNP pairs

modulating the expression of two separate neighboring genes. For example, rs11675339 and

rs2710687 of VSNL1 appear to control the expression of GEN1 and SMC6 respectively. We

also found that VSNL (also known as VILIP-1) has strong evidences to AD [42-44]. GEN1 and

SMC6 play pivotal roles in repairing double-strand breaks of genomic DNA. Interestingly,

rDNA instability, a result of failures to repair double-strand breaks of genomic DNA, has been

implicated in AD [45]. Most of the single-gene epistasis selected by GenEpi can be explained

by these three possible modes and only two of the significance SNP-SNP interactions are not

immediately clear at this moment. For example, rs12366151 resides in the eQTL of MICALCL,

an AD associated gene [46] but the interacting rs10831829 resides in the intron of PARVA, a

gene not-yet-reported to be associated with AD. Similarly, rs2421701 and rs200512701 reside

in two eQTLs regulating TNKS2 expression. However, the ADP-ribose polymerase

functionally contributes to AD is not clear.

Last, there are only two single-SNP features and both of them are located in APOE, which is a

well-known causal gene of AD, revealing that GenEpi is an effective tool to identify

disease-causing genes. Moreover, GenEpi successfully selected out the SNP rs429358, which

determines the allele type of APOE with rs7412. While GenEpi has shown its ability to identify

epistasis efficiently, it might still has the following limitations. Firstly, GenEpi can only detect

pairwise interactions. Second, GenEpi is a memory-consuming package, which might cause

memory errors when calculating the epistasis of a gene containing a large number of SNPs. We

recommend that the memory for running GenEpi should be over 256 GB. Finally, a small

sample size may lead overfitting, which forces us to use strict thresholds during feature

selection. In this way, GenEpi delivers a high precision rate, but might suffer having false

negatives. This implies different GWAS data might detect different sets of true positives. In

summary, the results of this study demonstrated that GenEpi is a promising software package to

identify causal SNPs and epistasis in GWAS, and it can be further used to predict the

phenotypes. With the demonstrated efficiency, GenEpi is a powerful tool to explore gene-gene

interactions that underlie complex diseases.

4. Materials

This study applied GenEpi on an AD cohort, which was used in Alzheimer's disease Dream

Challenge [28], In total, the cohort consists of 767 participants, who were healthy elderly, mild

cognitive impairment (MCI) and AD patients from the Alzheimer's Disease Neuroimaging

Initiative (ADNI) database. The 767 ADNI participants consist of 241 cognitively normal (CN),

130 Early MCI (EMCI), 273 Late MCI (LMCI) and 123 AD participants. We adopted only

genetic features in this study. All the genetic data has been pre-processed by the organizers that

held the challenge [28]. The genetic data were genotyped using the Illumina Human610-Quad

BeadChip and Illumina HumanOmniExpress BeadChip. The multidimensional scaling analysis

was applied by PLINK using HAPMAP3 to ensure that samples are within the cluster of

European populations. Subsequently, the data were imputed according to the 1,000 genome

haplotypes. After imputation, there were 12,809,667 genotype features in total. For predicting

the diagnosis of AD, we used 364 participants, of which the clinical diagnosis are CN or AD, to

predict which samples are control subjects or the AD patients.

5. Methods

The architecture of GenEpi is shown in Fig. 5. GenEpi is designed to group SNPs by gene

boundaries. GenEpi first considers the genetic variants in a gene as features, which have a

higher chance to interact with each other and to influence molecular functions. The idea of

within-gene epistasis analysis followed by cross-gene analysis is not new, which has also been

used in previous studies [47-50]. Differently, GenEpi adopts two-element combinatorial

encoding when producing features and models them by L1-regularized regression with

stability selection, which will be explained in Section 2.2. In the first stage (STAGE 1) of

GenEpi, the genotype features from each single gene will be combinatorically encoded and

modeled independently by L1-regularized regression with stability selection. In this way, we

can estimate the prediction performance of each gene and detect within-gene epistasis with a

low false positive rate. In the second stage (STAGE 2), both of the individual SNP and the

within-gene epistasis features selected by STAGE 1 are pooled together to generate

cross-gene epistasis features, and modeled again by L1-regularized regression with stability

selection as STAGE 1. Finally, the user can combine the selected genetic features with

environmental factors such as clinical features to build the final prediction models. In addition

to the main procedures, two pre-processing steps are also implemented in GenEpi: retrieving

the gene information from public databases and reducing the gene information from public

databases and reducing the dimensionality of the features using linkage disequilibrium (LD).

In the end, we released a Python package that implements GenEpi. The details of these steps

and the GenEpi method will be described in the following sections.

University of California Santa Cruz (UCSC) database

To obtain the gene information such as official gene symbols and genomic coordinates, we

retrieved kgXref and knownGene data table from the UCSC human genome annotation

database [51, 52]. The version of the database we used is the Feb. 2009 assembly of the

human genome hg19, GRCh37 Genome Reference Consortium Human Reference 37. The

two data tables were merged in order to generate a local database containing the gene symbols

as well as the genomic coordinates of each gene. The in-house script we built could update

this local database automatically. It is noted that there are many different categories of genes

in the RefSeq database. In this study, we only focused on the mRNA and non-coding RNA

(22,376 genes in total). The selected transcripts were projected on the genomic coordinates

and the coordinates of corresponding genes were determined based on the leftmost and

rightmost positions of the corresponding transcripts. Moreover, to discover the factors that

might affect the transcription of genes, we also retained the promoter region of each gene. In

genetics, the promoter region is a segment of DNA that initiates the transcription of a

particular gene. Promoters are located near the transcription start sites of genes, on the

upstream of the same DNA strand (towards the 5' region of the sense strand of the transcript).

In general, a promoter region can be 100-1000 base pairs long. In this study, we extracted

1,000 nucleotides on the upstream of the starting position of each gene as the promoter

region.

Estimation of linkage disequilibrium

In GWAS datasets, a SNP often exhibits high dependency with its nearby SNPs because of

linkage disequilibrium (LD). In the practical implantation, we prefer to group these dependent

features to reduce the dimension of features. In other words, we can take the advantages of LD

to reduce the dimensionality of SNP features. In this regard, we adopted the same approach

developed by Lewontin [53] to estimate LD. We used D' >0.9 and r^2>0.9 as the criteria to group

highly dependent SNP features as blocks. In each block, we chose the features with the largest

minor allele frequency to represent other features in the same block.

Discovery of within-gene epistasis

The main objective of the first stage in GenEpi is to select candidate features from each gene. In

order to extract SNP features for a gene, we used the start and end positions of each gene from

the local UCSC database to split the SNP features after dimension reduction. Since there are

22,376 genes in the UCSC database, we obtained 22,376 subsets of the SNP features. In each

subset, A SNP feature with the alleles 'A' and 'a' could have three possible genotypes, AA, Aa

and aa, which are used to refer to the pairs of alleles. The pairs of alleles are subsequently

separated into three binary features using one-hot encoding. In order to evaluate epistasis, we

generated interacting features by crossing each pair of genotype features. Considering the false

positive rate and computational complexity, we only focused on pairwise interactions of

epistasis throughout this study. We defined the interaction between two SNPs in Equation 1. In

Equation 1, $\boldsymbol{\alpha_1 SNP_1}$ + $\boldsymbol{\alpha_2 SNP_2}$ stand for the additive interactions and

$\boldsymbol{\alpha_{int(1,2)} SNP_1 \otimes SNP_2}$ represents the synergistic interactions that contain nine terms.

$$y \sim \alpha_0 + \sum_{m \in \{AA,Aa,aa\}} \alpha_{1,m} SNP_{1,m} + \sum_{m \in \{AA,Aa,aa\}} \alpha_{2,m} SNP_{2,m} + \sum_{m,n \in \{AA,Aa,aa\}} \alpha_{1,m,2,n} SNP_{1,m} SNP_{2,n}$$

$$= \alpha_0 + \alpha_1 SNP_1 + \alpha_2 SNP_2 + \alpha_{int(1,2)} SNP_1 \otimes SNP_2 \qquad Equation\,1$$

Before modeling each subset of genotype features, two criteria were adopted to exclude low

quality data. The first criterion is that the genotype frequency of a feature should exceed 5%,

where the genotype frequency means the proportion of genotype among the total samples in the

dataset. The second criterion is regarding the association between the feature and the phenotype.

We used χ^2 test to estimate the association between the feature and the phenotype, and the

p-value should be smaller than 0.01. In the end, a gene may have multiple SNPs. The general

form of the linear model for a gene with k SNPs is defined as Equation 2, which is termed as

two-element combinatorial encoding.

$$y \sim \alpha_0 + \sum_{i=1}^{k} \alpha_i SNP_i + \sum_{i=1}^{k} \sum_{j=1 \cap j \neq i}^{k} \alpha_{int(i,j)} SNP_i \otimes SNP_j \qquad\qquad Equation 2$$

We conducted L1-regularized regression with stability selection [54] for modeling each gene.

The first advantage is that the sparsity of the L1- regularized model prefers solutions with a

smaller number of features, which effectively reduces the effect of feature dependency. The

second advantage is by using stability selection [54] , we can largely reduce the false positive

rate. Stability selection works by resampling and remodeling the training set hundreds of times,

followed by picking out the features that are repeatedly selected across randomization. In this

study, we executed this randomization 500 times, and the features selected by stability selection

would be retained for the next stage.

Discovery of cross-gene epistasis

In the second stage, we used the features selected by STAGE 1 to generate cross-gene epistasis

features and then applied the same selection procedure described in Section 2.3 to find the

cross-gene epistasis that are associated with the phenotype. The procedures were slightly

modified here. Since we only focused on pairwise interactions, instead of using the entire

features we selected in STAGE 1, we only used single-SNP features to generate cross-gene

epistasis features. Also, we used the genotype frequency and the p-values of χ^2 test to control

the quality of features and to avoid overfitting. Nevertheless, the p-value of each feature in this

stage should be smaller than 10^{-5}. All of the features from different genes would be merged for

modeling cross-gene epistasis. We conducted L1-regularized regression for modeling, and the

stability selection were used once again to select the final genotype feature set. Since the

phenotype may also be affected by environmental factors, after determining the final set of

genotype features, the user can included the environmental factors such as clinical assessments

for constructing the final model. Subsequently, the final model was evaluated through a process

called double cross validation (CV). In the external loop of double CV, all the instances were

divided into two subsets to serve as training and independent test sets. In this study, we used

2-fold CV and leave-one-out CV (LOO CV) in external loop for evaluation. In the internal loop,

we also used 2-fold CV for model selection.

Tables

Table 1. The medians of the rank of the SNPs in the target epistasis for ReliefF

	SNP 1	SNP 2	SNP 3	SNP 4	SNP 5	SNP 6
Basic Model	1	2	1	2	1	2
Complex Model	7	8.5	9.5	11.5	11	19.5

422 **Table 2.** The summary of the number of SNPs, genes and features in each step of GenEpi .

Step	0	1	2	3	4	5
Function	Raw data	Download UCSC Database	Estimate LD Blocks	Split Features By Genes	Select Single-Gene Epistasis	Select Cross-Gene Epistasis
Number of SNP	12,809,667	12,809,667	12,102,888	4,916,249	44,865	24
Number of Gene	-	22,376	22,376	20,206	7,933	12
Number of Feature	-	-	-	-	34,689	14

425 **Table 3.** The score of different models in predicting control subjects or AD patients.

	Precision	Recall	F1 Score	Accuracy
Training	0.9633	0.8537	0.9052	0.9396
2-fold CV	0.7748	0.6992	0.7350	0.8297
LOO CV	0.8100	0.6585	0.7265	0.8324

426 F1 Score = 2 × (Precision × Recall) / (Precision + Recall); 'Training' stands for the process of a single-loop CV; '2-fold CV' means that

427 2-fold CV was used in the external loop of double CV; 'LOO CV' means that LOO CV was used in the external loop of double CV.

429   **Table 4.** The statistical significance of genetic features selected by GenEpi in predicting patients with AD.

Selected SNPs (RSID)	Weight	Odds Ratio	χ^2-test P-value	Genotype Frequency	Gene
rs3130614_BB, rs41276317_AB	3.16	19.23	1.42E-09	0.0742	*MICB*TOB2*
rs12095538_BB, rs2774308_AB	2.41	7.69	6.87E-07	0.0824	*SYT6*
rs12926153_AB, rs12922908_AA	1.18	4.83	6.89E-07	0.1511	*CLEC16A*
rs9652600_AB, rs12922908_AA	0.94	4.83	6.89E-07	0.1511	*CLEC16A*
rs9344977_BB, rs56148686_AB	1.94	4.32	1.14E-06	0.1813	*BACH2*
rs429358_AA	-2.01	0.17	1.73E-06	0.5962	*APOE*
rs56233035_AB, rs3678_AB	2.26	10.16	1.91E-06	0.0604	*CACNA1E*
rs11675339_AA, rs2710687_AA	2.32	3.94	3.55E-06	0.1923	*VSNL1*
rs12189429_BB, rs6881360_AA	1.36	4.34	3.65E-06	0.467	*ADAMTS12*
rs12187423_BB, rs6881360_AA	0.58	4.34	3.65E-06	0.467	*ADAMTS12*
rs10831829_BB, rs12366151_AA	3.48	9.50	4.90E-06	0.0577	*PARVA*
rs2052573_BB, rs34580133_AB	1.80	4.08	5.00E-06	0.1648	*LINC00299*
rs2421701_AB, rs200512701_AB	1.82	4.12	5.29E-06	0.1593	*TNKS2*
rs769449_AA	-1.19	0.16	8.42E-06	0.6648	*APOE*

430   The sign '*' between two gene symbols indicates cross-gene epistasis.

432 **Table 5.** The comparison of different algorithms.

Algorithm	# Input SNP	Time Cost	Top 15	Top 30	Top 45	Top 60
GenEpi	4,916,249	9.95	0.76	0.72	0.71	0.68
BOOST	4,916,249	2157.6	0.31	0.24	0.30	0.37
ReliefF	33,868	0.11	0.52	0.48	0.45	0.46
FastEpistasis	12,809,667	836.8	0.62	0.61	0.60	0.59

433 'Time Cost' is the time spent on identifying the epistasis, which was measured by single CPU time in days. The values in column top 15, top
434 30, top 45 and top 60 are the 2-fold CV scores. The 2-fold CV scores are the F1 scores.

436 **Figures**

Fig. 1. The boxplot for the rank of the target epistasis in different algorithms. (a) The results
of three basic-model datasets with one epistasis consisting of a SNP pair. (b) The result of the
complex-model dataset, which contained three epistasis. The 'S1-S2' means the epistasis
between SNP 1 and SNP 2 and so on. The values on the boxplot are the medians of the rank of
the target epistasis among the 100 runs of simulation.

Fig. 2. The boxplot of false positives in L1-regularized regression with and without stability selection.

447 **Fig. 3.** The ROC curves of different algorithms.

449 **Fig. 4.** The heatmap of gene expression in different tissues for the 12 genes selected by

450 GenEpi. The blue box highlights the sub-regions of brain.

453   **Fig. 5.** The architecture of GenEpi.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

This work has been supported by the Ministry of Science and Technology of Taiwan grant 105-2911-I-002-566, 105-2221-E-002-129-MY3 and 103-2627-M002-015.

Authors' contributions

Y.-C.C. initiated the study, designed the analysis procedures, performed the analysis and wrote the manuscript. J.-T.W. finished the literature survey of the epistasis that selected by GenEpi and wrote the discussion. C.-Y.C. is the main advisor of Y.-C.C., guided this research on right way and provided ideas to optimize the methods. Y.-C.C, M.-Y.H., Y.-A.T, C.-Y.C. and Y.-J.O. were team members of Alzheimer's disease Dream Challenge. C.-Y.C., Y.-J.O., P.-H.H. K.M.G. and S.W.Y. commented on the draft and revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Data collection and sharing for this project was funded by the ADNI (National Institutes of Health Grant U01 AG024904). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Abbott; Alzheimer's Association; Alzheimer's Drug Discovery Foundation; AmorfixLife Sciences Ltd.; AstraZeneca; Bayer HealthCare; BioClinica, Inc.; Biogen Idec Inc.; Bristol-Myers Squibb Company; Eisai Inc.; Elan

Pharmaceuticals Inc.; Eli Lilly and Company; F.Hoffmann-La Roche Ltd and its affiliated
company Genentech, Inc.; GE Healthcare; Innogenetics, N.V.; Janssen Alzheimer
Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical
Research & Development LLC.; Medpace, Inc.; Merck & Co., Inc.; Meso Scale Diagnostics,
LLC.; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Servier; Synarc Inc. and Takeda
Pharmaceutical Company. The Canadian Institutes of Health Research is providing funds to
support ADNI clinical sites in Canada. Private sector contributions are facilitated by the
Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the
Northern California Institute for Research and Education and the study is coordinated by the
Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data
are disseminated by the Laboratory for Neuro Imaging at the University of California, Los
Angeles. This research was also supported by NIH grants P30 AG010129, K01 AG030514 and
the Dana Foundation.

Reference

Reference

1.   Kingsmore SF, Lindquist IE, Mudge J, Gessler DD, Beavis WD: **Genome-wide**
association studies: progress and potential for drug discovery and development.
Nature Reviews Drug Discovery 2008, **7**(3):221-230.

2.   Ozaki K, Ohnishi Y, Iida A, Sekine A, Yamada R, Tsunoda T, Sato H, Sato H, Hori M,
Nakamura Y *et al*: **Functional SNPs in the lymphotoxin-alpha gene that are**
associated with susceptibility to myocardial infarction. *Nat Genet* 2002,
32(4):650-654.

3.   Klein RJ, Zeiss C, Chew EY, Tsai JY, Sackler RS, Haynes C, Henning AK,
SanGiovanni JP, Mane SM, Mayne ST *et al*: **Complement factor H polymorphism**
in age-related macular degeneration. *Science* 2005, **308**(5720):385-389.

4.   Pinero J, Bravo A, Queralt-Rosinach N, Gutierrez-Sacristan A, Deu-Pons J, Centeno E,
Garcia-Garcia J, Sanz F, Furlong LI: **DisGeNET: a comprehensive platform**
integrating information on human disease-associated genes and variants. *Nucleic*
Acids Res 2017, **45**(D1):D833-D839.

5.   McCarthy MI, Abecasis GR, Cardon LR, Goldstein DB, Little J, Ioannidis JP,
Hirschhorn JN: **Genome-wide association studies for complex traits: consensus,**
uncertainty and challenges. *Nat Rev Genet* 2008, **9**(5):356-369.

6.   Eichler EE, Flint J, Gibson G, Kong A, Leal SM, Moore JH, Nadeau JH: **Missing**
heritability and strategies for finding the underlying causes of complex disease.
Nat Rev Genet 2010, **11**(6):446-450.

7.   Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, McCarthy
MI, Ramos EM, Cardon LR, Chakravarti A *et al*: **Finding the missing heritability of**
complex diseases. *Nature* 2009, **461**(7265):747-753.

8.   Frazer KA, Murray SS, Schork NJ, Topol EJ: **Human genetic variation and its**
contribution to complex traits. *Nat Rev Genet* 2009, **10**(4):241-251.

9.   Shriner D, Vaughan LK, Padilla MA, Tiwari HK: **Problems with genome-wide**
association studies. *Science* 2007, **316**(5833):1840-1842.

10. Carlborg O, Haley CS: **Epistasis: too often neglected in complex trait studies?** *Nat*
Rev Genet 2004, **5**(8):618-625.

11. Cordell HJ: **Detecting gene-gene interactions that underlie human diseases**. *Nat*
Rev Genet 2009, **10**(6):392-404.

12. Easton DF, Pooley KA, Dunning AM, Pharoah PD, Thompson D, Ballinger DG,

Struewing JP, Morrison J, Field H, Luben R *et al*: **Genome-wide association study**
identifies   novel   breast   cancer   susceptibility   loci. *Nature*   2007,
447(7148):1087-1093.

13. Bodmer W, Bonilla C: **Common and rare variants in multifactorial susceptibility**
to common diseases. *Nat Genet* 2008, **40**(6):695-701.

14. Hindorff LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, Collins FS, Manolio
TA: **Potential etiologic and functional implications of genome-wide association**
loci for human diseases and traits. *Proc Natl Acad Sci U S A* 2009,
106(23):9362-9367.

15. Moore JH, Asselbergs FW, Williams SM: **Bioinformatics challenges for**
genome-wide association studies. *Bioinformatics* 2010, **26**(4):445-455.

16. Marchini J, Donnelly P, Cardon LR: **Genome-wide strategies for detecting multiple**
loci that influence complex diseases. *Nat Genet* 2005, **37**(4):413-417.

17. Wei WH, Hemani G, Haley CS: **Detecting epistasis in human complex traits**. *Nat*
Rev Genet 2014, **15**(11):722-733.

18. Schupbach T, Xenarios I, Bergmann S, Kapur K: **FastEpistasis: a high performance**
computing solution for quantitative trait epistasis. *Bioinformatics* 2010,
26(11):1468-1469.

19. Wan X, Yang C, Yang Q, Xue H, Fan X, Tang NL, Yu W: **BOOST: A fast approach**
to detecting gene-gene interactions in genome-wide case-control studies. *Am J*
Hum Genet 2010, **87**(3):325-340.

20. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J,
Sklar P, de Bakker PI, Daly MJ *et al*: **PLINK: a tool set for whole-genome**
association and population-based linkage analyses. *Am J Hum Genet* 2007,
81(3):559-575.

21. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ: **Second-generation**
PLINK: rising to the challenge of larger and richer datasets. *Gigascience* 2015,
4:7.

22. Moore JH, Williams SM: **New strategies for identifying gene-gene interactions in**
hypertension. *Ann Med* 2002, **34**(2):88-95.

23. Yang P, Ho JW, Yang YH, Zhou BB: **Gene-gene interaction filtering with ensemble**
of filters. *BMC Bioinformatics* 2011, **12 Suppl 1**:S10.

24. Bureau A, Dupuis J, Falls K, Lunetta KL, Hayward B, Keith TP, Van Eerdewegh P:
Identifying SNPs predictive of phenotype using random forests. *Genet Epidemiol*

568   2005, **28**(2):171-182.

569 25. Schwarz DF, Konig IR, Ziegler A: **On safari to Random Jungle: a fast**
570   **implementation of Random Forests for high-dimensional data**. *Bioinformatics*
571   2010, **26**(14):1752-1758.

572 26. Wan X, Yang C, Yang Q, Xue H, Tang NL, Yu W: **MegaSNPHunter: a learning**
573   **approach to detect disease predisposition SNPs and high level interactions in**
574   **genome wide association study**. *BMC Bioinformatics* 2009, **10**:13.

575 27. Murk W, DeWan AT: **Exhaustive Genome-Wide Search for SNP-SNP Interactions**
576   **Across 10 Human Diseases**. *G3 (Bethesda)* 2016, **6**(7):2043-2050.

577 28. Allen GI, Amoroso N, Anghel C, Balagurusamy V, Bare CJ, Beaton D, Bellotti R,
578   Bennett DA, Boehme KL, Boutros PC *et al*: **Crowdsourced estimation of cognitive**
579   **decline and resilience in Alzheimer's disease**. *Alzheimers Dement* 2016,
580   **12**(6):645-653.

581 29. Urbanowicz RJ, Meeker M, LaCava W, Olson RS, Moore JH: **Relief-Based Feature**
582   **Selection: Introduction and Review**. *arXiv preprint arXiv:171108421* 2017.

583 30. Urbanowicz RJ, Kiralis J, Sinnott-Armstrong NA, Heberling T, Fisher JM, Moore JH:
584   **GAMETES: a fast, direct algorithm for generating pure, strict, epistatic models**
585   **with random architectures**. *BioData Min* 2012, **5**(1):16.

586 31. Consortium GT: **The Genotype-Tissue Expression (GTEx) project**. *Nat Genet* 2013,
587   **45**(6):580-585.

588 32. Vivier E, Tomasello E, Paul P: **Lymphocyte activation via NKG2D: towards a new**
589   **paradigm in immune recognition?** *Curr Opin Immunol* 2002, **14**(3):306-311.

590 33. Collins RW: **Human MHC class I chain related (MIC) genes: their biological**
591   **function and relevance to disease and transplantation**. *Eur J Immunogenet* 2004,
592   **31**(3):105-114.

593 34. Quiroga I, Lehmann DJ, Barnardo MC, Fuggle S, Cortina-Borja M, Warden DR,
594   Smith AD: **Association study of MICA and MICB in Alzheimer's disease**. *Tissue*
595   *Antigens* 2009, **74**(3):241-243.

596 35. Mirza Z, Rajeh N: **Identification Of Electrophysiological Changes In Alzheimer's**
597   **Disease: A Microarray Based Transcriptomics And Molecular Pathway Analysis**
598   **Study**. *CNS Neurol Disord Drug Targets* 2017.

599 36. Saura CA, Parra-Damas A, Enriquez-Barreto L: **Gene expression parallels synaptic**
600   **excitability and plasticity changes in Alzheimer's disease**. *Front Cell Neurosci* 2015,
601   **9**:318.

37. Daschil N, Obermair GJ, Flucher BE, Stefanova N, Hutter-Paier B, Windisch M,
Humpel C, Marksteiner J: **CaV1.2 calcium channel expression in reactive**
astrocytes is associated with the formation of amyloid-beta plaques in an
Alzheimer's disease mouse model. *J Alzheimers Dis* 2013, **37**(2):439-451.

38. Porcellini E, Carbone I, Ianni M, Licastro F: **Alzheimer's disease gene signature**
says: beware of brain viral infections. *Immun Ageing* 2010, **7**:16.

39. Talkowski ME, Maussion G, Crapper L, Rosenfeld JA, Blumenthal I, Hanscom C,
Chiang C, Lindgren A, Pereira S, Ruderfer D *et al*: **Disruption of a large intergenic**
noncoding RNA in subjects with neurodevelopmental disabilities. *Am J Hum*
Genet 2012, **91**(6):1128-1134.

40. Uhrig M, Ittrich C, Wiedmann V, Knyazev Y, Weninger A, Riemenschneider M,
Hartmann T: **New Alzheimer amyloid beta responsive genes identified in human**
neuroblastoma cells by hierarchical clustering. *PLoS One* 2009, **4**(8):e6779.

41. Wang W, Mandel J, Bouaziz J, Commenges D, Nabirotchkine S, Chumakov I, Cohen
D, Guedj M, Alzheimer's Disease Neuroimaging I: **A Multi-Marker Genetic**
Association Test Based on the Rasch Model Applied to Alzheimer's Disease. *PLoS*
One 2015, **10**(9):e0138223.

42. Babic Leko M, Borovecki F, Dejanovic N, Hof PR, Simic G: **Predictive Value of**
Cerebrospinal Fluid Visinin-Like Protein-1 Levels for Alzheimer's Disease Early
Detection and Differential Diagnosis in Patients with Mild Cognitive Impairment.
J Alzheimers Dis 2016, **50**(3):765-778.

43. Kirkwood CM, MacDonald ML, Schempf TA, Vatsavayi AV, Ikonomovic MD, Koppel
JL, Ding Y, Sun M, Kofler JK, Lopez OL *et al*: **Altered Levels of Visinin-Like**
Protein 1 Correspond to Regional Neuronal Loss in Alzheimer Disease and
Frontotemporal Lobar Degeneration. *J Neuropathol Exp Neurol* 2016,
75(2):175-182.

44. Luo X, Hou L, Shi H, Zhong X, Zhang Y, Zheng D, Tan Y, Hu G, Mu N, Chan J *et al*:
CSF levels of the neuronal injury biomarker visinin-like protein-1 in Alzheimer's
disease and dementia with Lewy bodies. *J Neurochem* 2013, **127**(5):681-690.

45. Pietrzak M, Rempala G, Nelson PT, Zheng JJ, Hetman M: **Epigenetic silencing of**
nucleolar rRNA genes in Alzheimer's disease. *PLoS One* 2011, **6**(7):e22585.

46. Rouillard AD, Gundersen GW, Fernandez NF, Wang Z, Monteiro CD, McDermott MG,
Ma'ayan A: **The harmonizome: a collection of processed datasets gathered to**
serve and mine knowledge about genes and proteins. *Database (Oxford)* 2016,

2016.

47.   Ma L, Clark AG, Keinan A: **Gene-based testing of interactions in association**
studies of quantitative traits. *PLoS Genet* 2013, **9**(2):e1003321.

48.   Oh S, Lee J, Kwon MS, Weir B, Ha K, Park T: **A novel method to identify high**
order gene-gene interactions in genome-wide association studies: gene-based
MDR. *BMC Bioinformatics* 2012, **13 Suppl 9**:S5.

49.   Li S, Cui Y: **Gene-centric gene–gene interaction: A model-based kernel machine**
method. *The Annals of Applied Statistics* 2012, **6**(3):1134-1161.

50.   Wu X, Dong H, Luo L, Zhu Y, Peng G, Reveille JD, Xiong M: **A novel statistic for**
genome-wide interaction analysis. *PLoS Genet* 2010, **6**(9):e1001131.

51.   Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D:
The human genome browser at UCSC. *Genome Res* 2002, **12**(6):996-1006.

52.   Rosenbloom KR, Armstrong J, Barber GP, Casper J, Clawson H, Diekhans M, Dreszer
TR, Fujita PA, Guruvadoo L, Haeussler M *et al*: **The UCSC Genome Browser**
database: 2015 update. *Nucleic Acids Res* 2015, **43**(Database issue):D670-681.

53.   Lewontin RC: **The Interaction of Selection and Linkage. I. General**
Considerations; Heterotic Models. *Genetics* 1964, **49**(1):49-67.

54.   Meinshausen N, Bühlmann P: **Stability selection**. *Journal of the Royal Statistical*
Society: Series B (Statistical Methodology) 2010, **72**(4):417-473.