# Identification of differentially methylated cell-types in Epigenome-Wide Association Studies

Shijie C Zheng [1] , Charles E. Breeze [2] , Stephan Beck [3] and Andrew E. Teschendorff [1,3,4]

1. CAS Key Laboratory of Computational Biology, CAS-MPG Partner Institute for Computational Biology, Shanghai Institute of Nutrition and Health, Shanghai Institute for Biological Sciences, University of Chinese Academy of Sciences, Chinese Academy of Sciences, 320 Yue Yang Road, Shanghai 200031, China.

2. Altius Institute for Biomedical Sciences, 2211 Elliott Ave #410, Seattle, WA 98121, USA.

3. UCL Cancer Institute, Paul O'Gorman Building, University College London, 72 Huntley Street, London WC1E 6BT, United Kingdom.

4. Department of Women's Cancer, University College London, 74 Huntley Street, London WC1E 6AU, United Kingdom.

*Corresponding author: Andrew E. Teschendorff- a.teschendorff@ucl.ac.uk

**An outstanding challenge of Epigenome-Wide Association Studies (EWAS) performed in complex tissues is the identification of the specific cell-type(s) responsible for the observed differential DNA methylation. Here, we present a novel statistical algorithm, called CellDMC, which is able to identify not only differentially methylated positions, but also the specific cell-type(s) driving the differential methylation. We provide extensive validation of CellDMC on in-silico mixtures of DNA methylation data generated with different technologies, as well as on real mixtures from epigenome-wide-association and cancer epigenome studies. We demonstrate how CellDMC can achieve over 90% sensitivity and specificity in scenarios where current state-of-the-art methods fail to identify differential methylation. By applying CellDMC to a smoking EWAS performed in buccal swabs, we identify differentially methylated positions occurring in the epithelial compartment, which we validate in smoking-related lung cancer. CellDMC may help towards the identification of causal DNA methylation alterations in disease.**

Somatic DNA methylation (DNAm) alterations have been shown to reflect cumulative exposure to environmental disease risk factors [1], and may contribute to disease risk by modifying cellular phenotypes [2,3]. One major source of DNAm variation which may hamper the identification of DNAm alterations predisposing or driving disease in Epigenome-Wide

Association Studies (EWAS) [4], is cell-type heterogeneity [5,6]. While statistical methods for
identifying differentially methylated cytosines (DMCs) in heterogeneous tissues have been
developed [7-14], none allow the identification of the specific cell-types responsible for the
observed differential methylation [10]. Indeed, the only existing tool that can help pinpoint
differentially methylated cell-types is an enrichment analysis method for cell-type specific
DNase hypersensitive sites that is performed on a relatively large list of DMCs [15], not
allowing for individual CpGs to be ranked according to their likelihood of differential
methylation (DM) in individual cell-types. Here, we present and validate CellDMC, a novel
statistical algorithm that can identify interactions between phenotype and the proportions of
underlying cell-types in the tissue, thus allowing for the detection of differentially methylated
cytosines in individual cell-types (DMCTs).
47

## Results

**Detection of DMCTs with CellDMC: rationale and statistical framework**

We reasoned that identification of DMCTs is possible within the same linear regression
framework normally used to identify DMCs, by further inclusion of statistical interaction
terms between phenotype and estimated cell-type fractions (**Fig.1a, SI fig.S1**): intuitively, if a
DMC is specific to one of the cell-types in the mixture, the observed differential methylation
(DM) should be most prominent when the DM analysis is restricted to samples that contain
the highest fraction of that cell-type (**Fig.1b**). CellDMC analyses the DNAm patterns of
interactions of all cell-types in the mixture to infer DMCTs and their directionality of change
(i.e. hyper or hypomethylation) (**Fig.1**, **Online Methods, SI fig.S1**). Importantly, CellDMC
also works in scenarios where all cell-types are uni-directionally differentially methylated to
a similar degree (**Fig.1c**). CellDMC can also handle more complex scenarios, where a DMC
occurs in two cell-types with opposite directionality (i.e. hypomethylated in one and
hypermethylated in another) (**Fig.1d**), and which may not be identifiable by current
state-of-the-art DMC calling algorithms (see later).

To estimate cell-type fractions, CellDMC applies our previously validated EpiDISH
algorithm [16] in an iterative hierarchical procedure, called HEpiDISH [17], which leads to
improved cell-type fraction estimates in complex tissues by recognizing that cell-types are
naturally arranged along a developmental tree (**Online Methods, SI fig.S2**). In the context of
epithelial tissues, HEpiDISH accomplishes this by using two distinct DNAm reference
matrices, a primary reference matrix for the estimation of total epithelial, total fibroblast and
total immune-cell (IC) fractions, and a separate, secondary, non-overlapping DNAm
reference for the estimation of underlying IC cell subtype fractions. Justification and proof
that this procedure works is given elsewhere [17], while additional proof that HEpiDISH works
is provided here in the context of identifying DMCTs.  In total, HEpiDISH can estimate
cell-type fractions for 9-10 different cell-types commonly found in epithelial tissues,

including epithelial cells, fibroblasts and 7 major types of immune-cells (**Online Methods**) [17].

For tissues like breast, which contain a relatively large fraction of adipocytes [18], the primary

reference also includes a representative DNAm profile for fat cells [17].

**Validation of CellDMC on *in-silico* mixtures**

To test CellDMC's ability to detect DMCTs, we first considered a number of different

simulation scenarios, where we mixed together real DNAm profiles representing epithelial,

fibroblast and immune cell types, in known mixing proportions, introducing DMCs in one,

two or all cell-types, with parallel or opposite directionality, and over a range of different

signal-to-noise ratios (SNRs) (**Online Methods, SI table S1**). Our choice of SNRs are

conservative, corresponding to differences in average DNAm within an affected cell-type of

approximately 0.4 to 0.5 (i.e. 40 to 50%) in the high SNR regime (SNR~3) down to changes

as small as 0.1 (i.e. 10%) for the low SNR regime (SNR<1) (**SI fig.S3**). In effect, the low

SNR regime corresponds to a scenario where only 10% of say single epithelial cells exhibit a

common DNAm change. Considering first scenarios where the underlying cell-type fractions

do not change appreciably between cases and controls, CellDMC obtained a high sensitivity

and specificity to detect DMCTs (**Fig.2a**). For instance, even when SNR<1, CellDMC

obtained a sensitivity above 0.75 for all scenarios where the differential methylation is

unidirectional, regardless of the number of affected cell-types (**Fig.2a**). For scenarios with

bi-directional changes, CellDMC could detect DMCTs with a sensitivity above 0.75 as long

as the SNR was approximately 1.8 or above (**Fig.2a**). In line with these high sensitivity

values, we observed excellent agreement between the predicted DNAm difference in the

underlying DMCTs with the true DNAm difference, irrespective of SNR and whether DNAm

alterations were uni-or-bi-directional (**Fig.2b**). Importantly, CellDMC allows DMCTs to be

ranked in each cell-type according to their statistical significance with the ranking highly

correlated with true effect size, as required (**SI fig.S4**). All of the above results were largely

unchanged if cell-type fractions were allowed to vary between cases and controls (**SI fig.S5**),

or if DMCs only occurred in one of the underlying immune cell subtypes (compared to the

previous case when the DMCs were common to all ICs) (**SI fig.S6-S8**). CellDMC also

attained high sensitivity and specificity if *in-silico* mixtures were generated purely from IC

subtypes (without admixture by epithelial and fibroblast cells), representing the more

common scenario of EWAS performed in whole/peripheral blood (**SI fig.S9**), or if the

phenotype is continuously valued as opposed to binary (**SI fig.S10**). CellDMC was also

robust to binary heterogeneous phenotypes with cases defined by a bi-modal distribution, as

observed for instance in cancer (**SI fig.S11**). We further verified that CellDMC is robust to

the choice of cell-type fraction estimation algorithm (**SI fig.S12**). Underlying this robustness,

we observed that results were also unchanged if cell-type fraction point estimates were

perturbed using either theoretical or empirical confidence intervals (**SI fig.S13-14**).

We also assessed CellDMC's power and specificity to detect DMCTs as a function of sample

size, <u>cell-type complexity and</u> <u>missing cell-types</u> (**SI fig.S15-S18**). At a sample size of 100
controls and 100 cases, power of CellDMC was always greater than 0.75 if the SNR was
larger than 1.8 (corresponding to DNAm changes greater than 0.2 in individual cell-types)
(**Fig.2c**, **SI fig.S15**). <u>Of note, as long as the fraction of the affected cell-type is distributed</u>
<u>uniformly across samples (thus exhibiting a fairly large range), CellDMC's sensitivity was</u>
<u>robust to cell-type complexity</u>, i.e. to the number K of cell-types in the mixture, although due
to limitations on data availability, we could only assess this up to a value of K=7 (**Fig.2d**, **SI**
**fig.S16**). <u>We also assessed performance as a function of the variance of the altered cell-type</u>
<u>fraction, estimating that sensitivity of CellDMC would remain over 50% as long as the range</u>
<u>of fractions exhibited by the affected cell-type is higher than 0.2</u> (**Fig.2e, SI fig.S17**).
<u>CellDMC was also robust if one major unaffected cell-type was missing from the reference</u>
<u>DNAm matrix, while reduction in sensitivity was relatively marginal if an altered cell-type</u>
<u>was missing from the reference</u> (**SI fig.S18**). <u>In the case of blood tissue we simulated a</u>
<u>realistic scenario, where cell-type fractions were modelled as observed in real blood EWAS</u>
<u>but with one cell-type (CD8+ T-cells) missing from our reference: sensitivities to detect</u>
<u>DMCTs in CD4+ T-cells were reduced at most by only 15%</u> (**SI fig.S19**). <u>Thus, these data</u>
<u>support the view that CellDMC can reliably detect DMCTs in a wide range of different</u>
<u>realistic scenarios. However, we also expect CellDMC's power to be strongly influenced by</u>
<u>cell-type complexity, in line with the fact that DNAm references become less reliable as K</u>
<u>increases, and also because specific cell-type proportions may exhibit small variation or fall</u>
<u>below detection levels.</u>

**CellDMC outperforms state-of-the-art reference-based and reference-free DMC calling**
**methods**
Next, we compared CellDMC to state-of-the-art reference-based and reference-free DMC
calling methods (**Fig.3, SI fig.S20**). Although CellDMC outperformed competing methods in
effectively all considered scenarios, differences were especially striking when DNAm
changes were bi-directional (**Fig.3, SI fig.S20-S21**). For instance, when DNAm changes
occurred in two of the underlying cell-types with opposite directionality (denoted "Bi-2C"),
conventional reference-based DMC calling that only includes the estimated cell-type
fractions as covariates [7] would not resolve the underlying DMCs, resulting in sensitivities
well below 0.25 (**Fig.3b, SI fig.S20-S21**). In contrast, CellDMC attained sensitivities that
ranged from over 0.95 for high SNRs to around 0.5 for low SNRs (**Fig.3a**, **SI fig.S20-S21**). If
bi-directional changes occurred in more than 2 cell-types (denoted "Bi-AllC"), CellDMC also
attained significantly higher sensitivity values than conventional non-interaction based DMC
calling (**Fig.3a-b, SI fig.S20-S21**). Similar improvements in sensitivity of CellDMC over
state-of-the-art reference-free methods (SVA & RefFreeEWAS) [8,19,20] were also observed,
which were again particularly prominent in bi-directional scenarios affecting two or more
cell-types (**Fig.3a,c-d**). In line with the sensitivity results, CellDMC's specificity was also

higher and more stable than that of competing methods (**Fig.3e-h, SI fig.S20**). Thus, these
data demonstrate that CellDMC can attain sensitivity and specificity values close to 100% in
scenarios where current state-of-the-art methods would fail to detect DMCs.
155

**Validation of CellDMC on whole-genome bisulfite sequencing data**
All previous simulation models mixed together DNAm profiles generated with Illumina
450k/EPIC technology [21,22], a platform matched to the one used to generate the DNAm
references. Thus, to demonstrate applicability of CellDMC to samples generated with a
different technology, we devised *in-silico* mixture simulation models using whole-genome
bisulfite sequencing (WGBS) profiles of purified cell-types, generated as part of IHEC [23] (**SI**
**fig.S22**). Although CellDMC's performance measures dropped upon application to WGBS
data, the reduction was only relatively marginal, with sensitivity and specificity values
remaining high (sensitivity and specificity was still over 90% for the higher SNR values, **SI**
**fig.S23**). Thus, we conclude that profiling technology is not a major limiting factor for
CellDMC.
167

**Validation of CellDMC in a blood EWAS**
Next, we tested the ability of CellDMC to detect known DMCTs in a real EWAS. A recent
EWAS for Rheumatoid Arthritis (RA) conducted on purified B-cells identified a total of 10
DMCs, which were validated in two independent EWAS cohorts which also profiled purified
B-cells [24]. Thus, we reasoned that application of CellDMC to an independent RA EWAS
performed on 689 blood samples [5], should be able to identify these 10 RA-DMCs as being
differentially methylated specifically in B-cells. CellDMC predicted 8 out of these 10 to be
differentially methylated, and 7 out of 10 to be B-cell DMCTs (FDR<0.05, **Fig.4a**). Of note,
CellDMC predicted the DNAm difference between RA cases and controls to be occurring
primarily in B-cells, and not in the other immune cell subtypes (**Fig.4a**), despite the fact that
on average B-cells only accounted for 2% of blood cell subtypes in the samples, and despite
exhibiting relatively little variation (**Fig.4a**). Although the median absolute difference in
B-cell fractions between the 689 samples was only 1.5%, 5 samples exhibited fractions larger
than 10%, and therefore the ability of CellDMC to detect validated B-cell DMCTs is
consistent with our previous simulation estimates (**Fig.2e**). Importantly, we estimated the
chance that CellDMC would call 7 out of 10 randomly selected CpGs to be B-cell DMCTs at
their observed significance levels, to be less than 0.00001 (**Fig.4a**). We verified that results
were robust to the choice of cell-type fraction estimation algorithm (**SI fig.S24**) or if one of
the other minor blood cell subtypes was removed from the DNAm reference matrix (**SI**
**fig.S25**). Of note, had we applied a standard non-interaction based reference model to
identify DMCs, 6 of the 10 validated RA-DMCTs would have been detected as DMCs using
the same FDR<0.05 threshold but without knowledge of the underlying DMCT being B-cells.

We also note that CellDMC led to a larger than 4-fold reduction of DMCs compared to a
method that did not adjust for cell-type fractions, and a larger than 2-fold reduction of DMCs
compared to a non-interaction based model, with relatively little overlap between DMCs
called by CellDMC and the standard non-interaction model (**SI table.S2**). Thus, like the
standard non-interaction based model, CellDMC is able to remove large numbers of
associations caused by changes in the granulocyte/lymphocyte ratio [5], whilst also identifying
different DMCs to those found using the standard model.
198
199

**Validation of CellDMC in cancer epigenome studies**

To demonstrate the ability of CellDMC to detect DMCTs in solid epithelial tissues, we
applied it to the breast cancer EWAS setting, for which we had previously constructed a
gold-standard set of 19,379 true positive breast cancer epithelial DMCs (bcDMCs) [9,17]. This
set was obtained by intersecting DMCs from a comparison of breast cancer epithelial to
normal mammary epithelial cell-lines (thus representing relatively pure epithelial cell
populations), with a corresponding list of DMCs derived from the TCGA breast cancer study
[25]. We reasoned that applying CellDMC on our independent breast cancer tissue EWAS [18]
should not only validate the breast cancer epithelial DMCs, but should also predict the
epithelial compartment to be the main DMCT. Confirming this, CellDMC's predicted mean
methylation differences between the epithelial cells found in breast cancer and those found in
the normal tissue, exhibited an excellent correlation with the DNAm differences observed in
the actual cell-lines (**Fig.4b**). Overall, the magnitude of DNAm changes were still larger in
the cell-lines possibly owing to cell-culture effects or their longer proliferative history
(**Fig.4b**). CellDMC exhibited approximately 75% sensitivity and 70% specificity to detect
DMCs and over 60% sensitivity to detect the DMCTs as occurring in the epithelial
compartment (**Fig.4b, SI fig.S26a**). Although it is unknown if the 19,379 true positive
bcDMCs are also altered in fibroblasts, fat and immune cells, CellDMC predicted much
smaller fractions of these loci to be altered in fat and ICs (**SI fig.S26b**).
As a third validation on real data, we considered the case of the *HAND2* gene in endometrial
cancer [26,27]. We had previously discovered and validated hypermethylated DMCs in
endometrial cancer, mapping to the 1st exon region of the *HAND2* gene, which is a main
target of the progesterone receptor tumor suppressor pathway [26,27]. *HAND2* is expressed in
normal endometrial tissue including stromal fibroblasts, but lacks expression in endometrial
cancer and the stromal fibroblasts of the cancer tissue [26], suggesting that the observed
hypermethylation occurs in both epithelial and fibroblast cells. Indeed, the relatively large
difference in DNAm ($\Delta\beta \sim 0.5$) observed between endometrial cancer and normal endometrial
tissue [26] is a strong indication that the unidirectional DNAm change is occurring in both
epithelial and fibroblast compartments. Application of CellDMC to the TCGA endometrial
cancer study [28] confirmed that the DNAm change is occurring in the epithelial and fibroblast

cells of the tissue (**Fig.4c-d**), and that the alteration in fibroblasts is local to the 1[st] exon

region (**Fig.4d, SI fig.S27**). Interestingly, CellDMC also predicted the immune-cell

compartment to be differentially methylated, albeit less strongly so (**Fig.4c-d**). This is also

consistent with the observation that these *HAND2* CpG sites become hypermethylated with

age in blood [29], and that the unmatched TCGA endometrial cancers were derived from older

women. Thus, the prediction of CellDMC that these specific *HAND2* CpGs define DMCs in

all 3 major cell-types within the endometrial tissue is entirely consistent with previous

knowledge and data.

**CellDMC identifies smoking-associated DNAm changes in squamous epithelial cells**

Finally, to demonstrate how CellDMC may help gain novel biological insight, we applied it

to the largest EWAS cohort performed in buccal swabs [30], consisting of 790 samples from

women all aged 53 but with a wide range of different lifetime levels of smoking exposure. As

expected, buccal swabs contained mainly epithelial and immune cells (**SI fig.S28**), and thus

we used CellDMC to predict smoking-associated DMCTs in these two cell-types. By

comparison to a gold-standard list of 62 true smoking-associated DMCs derived from whole

blood EWAS (validated in at least three independent whole blood EWAS studies) [27], we

confirmed CellDMC's sensitivity to detect true DMCTs: most of the gold-standard DMCs

were predicted to be hypomethylated specifically in the immune cell compartment of the

buccal tissue (**Fig.5a-d**), in line with their observed hypomethylation in blood EWAS [31].

Confirming CellDMC's specificity, the top ranked DMCTs in immune cells have all been

previously validated as being smoking-associated DMCs in blood (**Fig.5a**). Although most of

the 62 gold-standard smoking-DMCs in blood were not predicted to be altered in the

epithelial cells of the buccal swabs, the great majority (i.e over 90%) of DMCTs were,

however, predicted to occur in the epithelial compartment, with a strong skew towards

hypermethylation (**Fig.5b**). To validate these epithelial DMCTs, we posited that their

variance in DNAm levels would increase with the fraction of epithelial cells in lung

squamous cell carcinoma [32], a cancer strongly associated with smoking, whilst also exhibiting

a concomitant decrease in variance in samples with a higher immune-cell content. We were

able to confirm this pattern for the top-ranked hypomethylated epithelial DMCT (**Fig.5e**), and

for most of the other epithelial DMCTs, at high statistical significance (**Fig.5f, Online**

**Methods**). Moreover, predicted hyper-and-hypo methylated epithelial DMCTs exhibited a

highly significant trend towards positive and negative correlations between their DNAm

levels and the estimated epithelial cell-type fractions, as required (**Fig.5f**). To confirm the

biological and clinical significance of these findings, we posited that the DNAm alterations

happening in the epithelial cells would exhibit increased deviations in lung cancer compared

to the normal adjacent tissue, owing to an increase in the epithelial fraction in the tumors

(**Fig.5g**). Confirming our expectation, hypermethylated and hypomethylated epithelial

DMCTs exhibited increased and decreased levels of DNAm in lung cancer, respectively (**Fig.5h**). Consistent with the total immune cell fraction not being altered between normal and cancer, the top ranked DMCTs specifically hypomethylated in immune cells did not exhibit decreased levels in cancer, but increased levels, probably due to shifts in the specific immune cell subtype proportions (**Fig.5h**) Thus, the epithelial smoking-DMCTs identified here represent epigenetic alterations occurring in the squamous cell of origin of smoking related lung cancer and therefore may mark cells that are being selected for during lung carcinogenesis.

## Conclusions

In summary, we have presented a novel algorithm aimed at identifying DMCTs, and have extensively validated the method on simulated as well as real data, encompassing common EWAS and cancer epigenome scenarios, and different technologies. The ability of CellDMC to rank CpGs according to their probability of being DM in each of the underlying cell-types will be invaluable for EWAS and cancer epigenome studies to help improve biological interpretation, for prioritizing candidates that require functional validation, and to help elucidate causal pathways to disease.

**Software Availability:** CellDMC is freely available as a user-friendly R-script within the EpiDISH Bioconductor package, freely available from github (https://github.com/sjczheng/EpiDISH), and in due course also from Bioconductor (https://www.bioconductor.org).

**Author Contributions:** Study was conceived by AET. SCZ and AET contributed jointly to methods development. SCZ performed the statistical analyses. Manuscript was written by both SCZ and AET. SB and CEB contributed valuable feedback.

**Competing Interests:** The authors declare that they have no competing interests.

**Acknowledgements:** This work was supported by NSFC (National Science Foundation of China) grants 31571359, 31771464 and 31401120, a Wellcome Trust grant (99148) and by a Royal Society Newton Advanced Fellowship (NAF project number: 522438, NAF award number: 164914).

## References

1. Petronis, A. Epigenetics as a unifying principle in the aetiology of complex traits and diseases. *Nature* **465**, 721-7 (2010).
2. Feinberg, A.P., Ohlsson, R. & Henikoff, S. The epigenetic progenitor origin of human cancer. *Nat Rev*

*Genet* **7**, 21-33 (2006).

3.    Lappalainen, T. & Greally, J.M. Associating cellular epigenetic models with human phenotypes. *Nat Rev*
*Genet* **18**, 441-451 (2017).

4.    Rakyan, V.K., Down, T.A., Balding, D.J. & Beck, S. Epigenome-wide association studies for common
human diseases. *Nat Rev Genet* **12**, 529-41 (2011).

5.    Liu, Y. *et al.* Epigenome-wide association data implicate DNA methylation as an intermediary of
genetic risk in rheumatoid arthritis. *Nat Biotechnol* **31**, 142-7 (2013).

6.    Jaffe, A.E. & Irizarry, R.A. Accounting for cellular heterogeneity is critical in epigenome-wide
association studies. *Genome Biol* **15**, R31 (2014).

7.    Houseman, E.A. *et al.* DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC*
*Bioinformatics* **13**, 86 (2012).

8.    Houseman, E.A., Molitor, J. & Marsit, C.J. Reference-free cell mixture adjustments in analysis of DNA
methylation data. *Bioinformatics* **30**, 1431-9 (2014).

9.    Zheng, S.C. *et al.* Correcting for cell-type heterogeneity in epigenome-wide association studies:
revisiting previous analyses. *Nat Methods* **14**, 216-217 (2017).

10.   Teschendorff, A.E. & Relton, C.L. Statistical and integrative system-level analysis of DNA methylation
data. *Nat Rev Genet* **19**, 129-147 (2018).

11.   Zou, J., Lippert, C., Heckerman, D., Aryee, M. & Listgarten, J. Epigenome-wide association studies
without the need for cell-type composition. *Nat Methods* **11**, 309-11 (2014).

12.   Rahmani, E. *et al.* Sparse PCA corrects for cell type heterogeneity in epigenome-wide association
studies. *Nat Methods* **13**, 443-5 (2016).

13.   Rahmani, E. *et al.* Correcting for cell-type heterogeneity in DNA methylation: a comprehensive
evaluation. *Nat Methods* **14**, 218-219 (2017).

14.   Lutsik, P. *et al.* MeDeCom: discovery and quantification of latent components of heterogeneous
methylomes. *Genome Biol* **18**, 55 (2017).

15.   Breeze, C.E. *et al.* eFORGE: A Tool for Identifying Cell Type-Specific Signal in Epigenomic Data. *Cell Rep*
**17**, 2137-2150 (2016).

16.   Teschendorff, A.E., Breeze, C.E., Zheng, S.C. & Beck, S. A comparison of reference-based algorithms for
correcting cell-type heterogeneity in Epigenome-Wide Association Studies. *BMC Bioinformatics* **18**,
105 (2017).

17.   Zheng, S.C. *et al.* A novel cell-type deconvolution algorithm reveals substantial contamination by
immune cells in saliva, buccal and cervix. *Epigenomics* (2018).

18.   Teschendorff, A.E. *et al.* DNA methylation outliers in normal breast tissue identify field defects that are
enriched in cancer. *Nat Commun* **7**, 10478 (2016).

19.   Leek, J.T. & Storey, J.D. Capturing heterogeneity in gene expression studies by surrogate variable
analysis. *PLoS Genet* **3**, 1724-35 (2007).

20.   McGregor, K. *et al.* An evaluation of methods correcting for cell-type heterogeneity in DNA
methylation studies. *Genome Biol* **17**, 84 (2016).

21.   Sandoval, J. *et al.* Validation of a DNA methylation microarray for 450,000 CpG sites in the human
genome. *Epigenetics* **6**, 692-702 (2011).

22.   Moran, S., Arribas, C. & Esteller, M. Validation of a DNA methylation microarray for 850,000 CpG sites
of the human genome enriched in enhancer sequences. *Epigenomics* **8**, 389-99 (2016).

23.   Stunnenberg, H.G., International Human Epigenome, C. & Hirst, M. The International Human
Epigenome Consortium: A Blueprint for Scientific Collaboration and Discovery. *Cell* **167**, 1145-1149

(2016).

24. Julia, A. *et al.* Epigenome-wide association study of rheumatoid arthritis identifies differentially
methylated loci in B cells. *Hum Mol Genet* **26**, 2803-2811 (2017).

25. Koboldt, D.C. *et al.* Comprehensive molecular portraits of human breast tumours. *Nature* **490**, 61-70
(2012).

26. Jones, A. *et al.* Role of DNA Methylation and Epigenetic Silencing of HAND2 in Endometrial Cancer
Development. *PLoS Med* **10**, e1001551 (2013).

27. Jiao, Y., Widschwendter, M. & Teschendorff, A.E. A systems-level integrative framework for
genome-wide DNA methylation and gene expression data identifies differential gene expression
modules under epigenetic control. *Bioinformatics* **30**, 2360-6 (2014).

28. Kandoth, C. *et al.* Integrated genomic characterization of endometrial carcinoma. *Nature* **497**, 67-73
(2013).

29. Yuan, T. *et al.* An integrative multi-scale analysis of the dynamic DNA methylation landscape in aging.
*PLoS Genet* **11**, e1004996 (2015).

30. Hannum, G. *et al.* Genome-wide methylation profiles reveal quantitative views of human aging rates.
*Mol Cell* **49**, 359-67 (2013).

31. Gao, X., Jia, M., Zhang, Y., Breitling, L.P. & Brenner, H. DNA methylation changes of whole blood cells in
response to active smoking exposure in adults: a systematic review of DNA methylation studies. *Clin*
*Epigenetics* **7**, 113 (2015).

32. Cancer Genome Atlas Research, N. Comprehensive genomic characterization of squamous cell lung
cancers. *Nature* **489**, 519-25 (2012).

33. Venables, W.N. & Ripley, B.D. *Modern Applied Statistics with S*, (Springer, New York, 2002).

34. Newman, A.M. *et al.* Robust enumeration of cell subsets from tissue expression profiles. *Nat Methods*
**12**, 453-7 (2015).

35. Nazor, K.L. *et al.* Recurrent variations in DNA methylation in human pluripotent stem cells and their
differentiated derivatives. *Cell Stem Cell* **10**, 620-34 (2012).

36. Absher, D.M. *et al.* Genome-wide DNA methylation analysis of systemic lupus erythematosus reveals
persistent hypomethylation of interferon genes and compositional changes to CD4+ T-cell populations.
*PLoS Genet* **9**, e1003678 (2013).

37. Limbach, M. *et al.* Epigenetic profiling in CD4+ and CD8+ T cells from Graves' disease patients reveals
changes in genes associated with T cell receptor signaling. *J Autoimmun* **67**, 46-56 (2016).

38. Marabita, F. *et al.* An evaluation of analysis pipelines for DNA methylation profiling using the Illumina
HumanMethylation450 BeadChip platform. *Epigenetics* **8**, 333-46 (2013).

39. Nestor, C.E. *et al.* DNA methylation changes separate allergic patients from healthy controls and may
reflect altered CD4+ T-cell population structure. *PLoS Genet* **10**, e1004059 (2014).

40. Reynolds, L.M. *et al.* Age-related variations in the methylome associated with gene expression in
human monocytes and T cells. *Nat Commun* **5**, 5366 (2014).

41. Zilbauer, M. *et al.* Genome-wide methylation analyses of primary human leukocyte subsets identifies
functionally important cell-type-specific hypomethylated regions. *Blood* **122**, e52-60 (2013).

42. Reinius, L.E. *et al.* Differential DNA methylation in purified human blood cells: implications for cell
lineage and studies on disease susceptibility. *PLoS One* **7**, e41361 (2012).

43. Smyth, G.K. Linear models and empirical bayes methods for assessing differential expression in
microarray experiments. *Stat Appl Genet Mol Biol* **3**, Article3 (2004).

**Figure legends**

**Figure-1: Identification of differentially methylated cell-types (DMCTs) using CellDMC.**
**a)** For a given DNAm data matrix, CellDMC uses a reference DNAm matrix encompassing
major cell-types (CTs) in the tissue of interest, to estimate cell-type fractions in each sample,
subsequently adjusting the DNAm data matrix for these estimated fractions. Using the
adjusted beta-values, it then fits statistical models that include interaction terms between the
phenotype and estimated cell-type fractions to identify DMCs in specific cell-types (DMCTs).
These can then be ranked according to statistical significance in each cell-type. **b,c,d)**
Scatterplots of adjusted beta-values against cell-type fraction for 3 different types of DMCTs.
**b)** A DMCT (CpG1) which is hypermethylated in cell-type CT1 but not in cell-types CT2 and
CT3. **c)** A DMCT (CpG2) hypermethylated in all three cell-types. **d)** A DMCT (CpG3)
occurring in two cell-types (CT1 & CT2) but with DNAm changes occurring in opposite
direction (hypermethylated in CT1, hypomethylated in CT2). Given a DNAm data matrix for
an epithelial tissue that includes epithelial, stromal (fibroblast, fat) and all major immune cell
types, CellDMC is able to infer not only DMCs but also the corresponding DMCTs.

**Figure-2: Validation of CellDMC on simulated data. a)** Sensitivity (SE) and specificity
(SP) (y-axis) of CellDMC to detect DMCTs in five different scenarios and for five different
SNRs (x-axis). The scenarios correspond to uni-directional DNAm changes being unique to
one cell-type (Uni-1C), shared by two cell-types (Uni-2C) or occurring in all cell-types
(Uni-AllC), and to bi-directional changes (i.e hypermethylated and hypomethylated) in two
cell-types (Bi-2C) or all cell-types (Bi-AllC). SNR values range from SNR~3 (mean DNAm
difference in affected cell-types of approx. 0.4 to 0.5) to SNR < 1 (mean DNAm difference in
affected cell-types of approx. 0.1). **b)** Scatterplots of the true DNAm difference in affected
cell-types (y-axis) versus the predicted DNAm difference from CellDMC (x-axis) for the
uni-directional (left panel) and bi-directional (right panel) cases. Data points are shown for
100 DMCs, 2 affected cell-types (CT1 and CT2) and for five SNR levels. **c)** Sensitivity to
detect DMCTs as a function of total sample size (numbers of cases and controls are assumed
equal) for five different SNR values in the Uni-1C (left panel) and Uni-AllC (right panel)
scenarios. **d)** Sensitivity to detect DMCTs as a function of the total number of cell-types (K)
in the Uni-1C scenario, with fractions sampled from a uniform Dirichlet distribution. In c-d)
each data point represents the mean over 100 Monte-Carlo runs with 100 "control" and 100

"disease" in-silico mixtures. **e)** As d) but now as a function of the cell-fraction range

exhibited by the affected cell-type in the mixture (K=3). Results shown for 100 Monte-Carlo

runs.

**Figure-3: Benchmarking of CellDMC to state-of-the-art reference based and**

**reference-free DMC calling algorithms. a-d)** From left to right, plots of the sensitivity of

CellDMC, of reference-based DMC calling (which includes cell-type fractions only as

covariates without interaction-terms denoted "NoInt"), of reference-free methods

RefFreeEWAS and SVA, against the SNR for five different differentially methylated cell-type

scenarios, where mean cell-type fractions are not different between cases and controls.

Because the competing methods can only detect DMCs, and not DMCTs, we compare the

methods in terms of their sensitivity to detect DMCs. Results are for 100 Monte-Carlo runs,

where in each run 200 in-silico mixture samples were simulated (100 "control" and 100

"disease"). Note that for the bi-directional scenario, competing state-of-the-art methods

cannot reliably detect DMCs, because the opposite directional DNAm change in two of the

underlying cell-types (assumed here to be of equal magnitude) "cancel-out". **e-h)** As a-d), but

now for the specificity.

**Figure-4: Validation of CellDMC in real EWAS and cancer epigenome data. a)**

Validation of CellDMC in a Rheumatoid Arthritis (RA) EWAS performed on 689 blood

samples (Liu et al [5]). The bar to the left indicates 10 CpG sites that have been validated to be

RA-associated DMCs in 3 independent purified B-cell cohorts, with their directionality of

DNAm change as indicated [24]. The central heatmap displays the predicted DMCTs from

CellDMC. Statistical significance of this result was assessed using 100,000 Monte-Carlo runs,

where in each run 10 CpGs were selected at random and their weighted averaged t-statistic

(grey curve and area) compared to the observed value (vertical red line). Boxplots compare

cell-type fraction estimates in the 689 samples between cell-types. **b)** Validation of CellDMC

to detect true breast cancer epithelial DMCs. Scatterplot of the DNAm difference between

breast cancer cell lines and normal mammary epithelial cell-lines (y-axis) against the

predicted DNAm difference from CellDMC for the epithelial subfraction of breast cancer

tissue compared to the epithelial subfraction of normal breast, for a total of approximately

19,000 true positive breast cancer epithelial DMCs. Fisher-test P-value is given. Barplots to

the right display CellDMC's sensitivity to detect DMCs and DMCTs as well as specificity. **c)**

Validation of CellDMC in the endometrial TCGA cancer study. Scatterplots of the beta value

(adjusted for cell-type fractions) of one of the *HAND2* CpGs against the epithelial, fibroblast

and total immune-cell (IC) fractions, with samples labeled according to endometrial cancer

(red) or normal endometrium (blue). Right panel displayss the predicted DNAm difference

between cancer and normal in each of the 3 cell-types, for all CpGs mapping to *HAND2*.

**Figure-5: Identification of epithelial specific smoking DMCTs and their relevance in lung cancer. a)** Smoothed scatterplot of CellDMC t-statistics in epithelial cells (x-axis) vs immune cells (IC) (y-axis) derived from a 790 buccal swab EWAS with smoking as the phenotype. Blue dashed lines indicate level of statistical significance (FDR<0.05). Red points label CpGs that belong to a gold-standard list of 62 true smoking-DMCs in blood. **b)** Barplots comparing the percentage of DMCTs that occur in epithelial and IC compartments, measured relative to the set of 62 smoking-DMCs in blood, or relative to all called DMCs/DMCTs. **c)** Barplots displaying the number of different types of DMCT, as indicated. Epi=epithelial only, IC=immune cell only, Hyper=hypermethylated in smokers, Hypo=hypomethylated in smokers. All means a DMC that occurs in both cell-types. **d)** Heatmap of CellDMC predictions (significance of signed adjusted P-values) for the 62 true smoking-DMCs. **e)** Scatterplot of the DNAm level (beta-value) of the top-ranked smoking-associated epithelial-specific DMCT (hypomethylated) against the epithelial cell-type fraction (left panel) and total immune cell fraction (right panel) across the 275 lung squamous cell carcinoma samples from the TCGA. Observe the trend towards hypomethylation and increased variance as epithelial fraction increases. **f) Left panel:** Scatterplot of the Pearson Correlation Coefficients (PCC, y-axis) between DNAm and epithelial cell-type fraction, as computed over the lung cancer TCGA samples, against the CellDMC t-statistics for predicted epithelial-specific DMCTs derived in the buccal swab smoking EWAS. Dashed lines indicate level of statistical significance and P-value is from a one-tailed Fisher's exact test on the CpGs passing significance in each quadrant. **Right panel:** Scatterplot of the signed Breusch-Pagan (BP) differential variance statistic of DNAm against immune-cell fraction (y-axis) versus the corresponding statistic for the epithelial fraction, as computed across the TCGA lung cancer samples, for all predicted epithelial-specific smoking-DMCTs derived from buccal swab EWAS. **g)** Boxplots of cell-type fractions estimated with EpiDISH for epithelial, fibroblast and immune cells in the normal and lung squamous cell carcinoma (LSCC) samples from the TCGA. **h)** Average DNAm levels of three categories of DMCTs, derived with CellDMC from the buccal smoking EWAS, in the matched normal-adjacent and LSCC samples from the TCGA. The DMCT categories are the 50 top-ranked hypomethylated and hypermethylated (promoter) epithelial DMCTs, and the top-ranked 50 hypomethylated immune-cell DMCTs. P-values are from a one-tailed Wilcoxon rank sum test.

# Online Methods

**Estimation of cell-type fractions in complex tissues using EpiDISH/HEpiDISH**

Estimation of the fractions of cell-types in a given sample constitutes the first step of the CellDMC algorithm and uses our previously validated EpiDISH [16] and HEpiDISH [17] methods. We use these methods to estimate cell-type fractions for *in-silico* generated mixtures as well as for real samples from EWAS and cancer epigenome studies. Which method we use, i.e. EpiDISH or a hierarchical iterative version of it called HEpiDISH depends on the desired cell-type resolution, as we explain next.

In the case of whole blood, peripheral blood or *in-silico* mixtures of blood cell subtypes, we use the EpiDISH procedure with a DNAm reference matrix defined over 333 IC cell subtype specific DMCs and 7 IC cell subtypes [16]. Briefly, EpiDISH models the DNAm profile of any given sample as a linear mixture over the DNAm profiles for the individual cell-types making up the sample. The associated multivariate linear regression is run using only the 333 IC cell subtype specific DMCs, and is performed using a robust M-estimator, as implemented in the *rlm* function of the *MASS* R-package [33]. We refer to this implementation as Robust Partial Correlations (RPCs) [16]. In this framework, the non-negativity and normalization constraints on the cell-type fractions (i.e. the estimated regression coefficients) are imposed *a-posteriori*, i.e. negative weights are set to zero and all other non-zero weights are normalized so that their sum adds to 1. As shown by us [16], RPCs improves inference, albeit marginally so, over the alternative approach which is to impose these constraints during inference via quadratic programming or constrained projection (CP) optimization [7]. RPCs also performs similarly to another non-constrained approach that uses a penalized support vector regression model called CIBERSORT [16,34]. In this work we use EpiDISH with RPCs throughout and only use CP for comparative purposes.

In the case of solid epithelial tissues and *in-silico* mixtures of epithelial, fibroblast and various IC cell subtypes, and when we only need to estimate cell-type fractions for the total epithelial, total fibroblast and total IC fractions, we once again use EpiDISH with RPCs, but now with a DNAm reference matrix defined over 716 DMCs and 3-cell types (generic Epithelial, generic Fibroblast, and a generic IC, denoted as the EpiFibIC reference matrix), as derived and validated previously by us in Zheng et al [17]. The 716 DMCs are highly discriminative of the 3 main cell-types, exhibiting large differences in mean DNAm between epithelial, fibroblast and ICs [17]. In some of the simulation scenarios, we also estimate fractions for individual IC cell subtypes, in which case we use our validated HEpiDISH algorithm [17]. As explained in Zheng et al, HEpiDISH works by implementing EpiDISH in an iterative hierarchical fashion, first estimating total epithelial, total fibroblast and total IC fractions using the EpiFibIC reference, and then estimating proportions for the 7 individual IC cell subtypes using a 188 DMC subset of the original 333 DMC reference used to estimate fractions in blood. A key point is that the 188 DMCs do not share any overlap with the 716 DMCs of the EpiFibIC reference, and that all 188 DMCs have been selected to ensure

that their median DNAm levels do not vary appreciably (most differences are less than 0.05 and the
maximum difference is 0.3, in absolute terms) between the epithelial, fibroblast and the collection of ICs [17].
This latter criterion helps ensure that the error or noise induced by inferring IC cell subtype fractions in the
mixture using a reference matrix that does not include epithelial and fibroblast references, is less than the
effect size or signal of the 188 DMCs (since any of the original 333 DMCs in the blood reference exhibit
big differences in DNAm, i.e. at least +/-0.5 with the great majority exhibiting over +/-0.8 differences
between at least one IC cell subtype and all others. See Zheng et al [17] for further justification and
validation.

552
553
**The CellDMC algorithm**
CellDMC consists of 3 basic steps: (1) Estimation of cell-type fractions in a given sample using EpiDISH
[16] or Hierarchical EpiDISH (HEpiDISH) [17], depending on the cell-type resolution desired (as described in
previous section), (2) Estimation of differentially methylated cell-types (DMCTs) for a given phenotype of
interest (binary or continuous), (3) Ranking of DMCTs per cell-type.
The key idea behind CellDMC is that a DNAm alteration occurring in a specific cell-type will exhibit a
significant interaction between the phenotype and the corresponding cell-type fraction variable. To
introduce relevant notation, we shall assume that we have a DNA methylation beta-valued matrix $x_{cs}$,
with $c$ ($c = 1,..,C$) labeling the CpGs and $s$ ($s = 1,...,S$) labeling the samples. For each sample $s$, let
$y_s$ denote the phenotype (e.g. a 0 or 1 variable for binary phenotypes, or a continuously-valued number for
continuous phenotypes), and let $\vec{y}$ denote the vector of phenotype values over all samples.
Correspondingly, we define $\vec{x}_c$ to be the vector of beta (DNAm) values for CpG $c$ across the $S$ samples.
We further assume that the fractions for the main cell-types, indexed by $k$ ($k=1...K$ and with $K$ the number
of cell-types), in the given samples have already been estimated using EpiDISH/HEpiDISH (or another
algorithm the user wishes to use). We denote the estimated fractions for cell-type $k$ over the $S$ samples with
the vector $\vec{f}_k$. CellDMC is then based on the following statistical model, which is run separately for each
CpG $c$:
571

$$\vec{x_c} = \mu_c + \sum_{k=1}^{K-1} \mu_{ck}\vec{f_k} + \beta_c\vec{y} + \sum_{k=1}^{K-1} \beta_{ck}^{(I)}\vec{f_k} * \vec{y} + \vec{\varepsilon}_c$$

where $\mu_c, \mu_{ck}, \beta_c, \beta_{ck}^{(I)}$ are regression coefficients to be estimated, * denotes the interaction term and
where we assume that errors are Gaussianly distributed with a variance that may depend on the specific
CpG $c$. We note that the above summations don't run over all $K$ cell-types, because cell-type fractions are
normalized and must add to 1. Because of this normalization constraint, the intercept term and the
phenotype main effects term can be absorbed into the corresponding summations, so that an equivalent
model is
578

$$\vec{x_c} = \sum_{k=1}^{K} \mu_{ck}\vec{f_k} + \sum_{k=1}^{K} \beta_{ck}^{(I)}\vec{f_k} * \vec{y} + \vec{\varepsilon}_c$$

The regression coefficients $\beta_{ck}^{(I)}$ inform us as to whether there is a significant interaction between the

phenotype and the corresponding fraction for cell-type $k$. We note that if differential methylation associated
with the phenotype occurs at a CpG $c$ and in cell-type $k$ that the observed differential methylation should
be larger in samples with high fractions for that cell-type $k$ compared to samples with low content for
cell-type $k$, and should be detectable via a statistically significant interaction term $\beta_{ck}^{(I)}$. We solve the above
model using least squares with the *lm* function in R, which provides estimates for the regression
coefficients and their statistical significance via P-values $P_{ck}^{(I)}$. The P-values $P_{ck}^{(I)}$ for each cell-type $k$ are
adjusted for multiple hypothesis testing using Benjamini-Hochberg (BH) False Discovery Rate (FDR)
estimation. For those CpGs with BH-FDR values less than a predefined significance threshold (i.e.
typically BH FDR<0.05), we call it a DMCT in the given cell-type. Finally, CpGs can be ranked within
each cell-type according to the associated P-value of significance.
Finally, we note that additional covariates representing other biological (e.g. age, gender, ethnicity) or
technical factors (batch) can be included in the above model. For instance, for a set $Q$ of such factors, let
$\vec{z}_q$ $(q=1...Q)$ denote the factor values across the $S$ samples, then the model above becomes

$$\vec{x_c} = \sum_{k=1}^{K} \mu_{ck}\overrightarrow{\widehat{f_k}} + \sum_{k=1}^{K} \beta_{ck}^{(I)}\overrightarrow{\widehat{f_k}} * \vec{y} + \sum_{q=1}^{Q} \rho_{cq}\vec{z}_q + \vec{\varepsilon}_c$$

593
594

**DNA methylation datasets used for *in-silico* mixture simulation experiments**
The following lists the DNAm datasets used in this manuscript to generate the *in-silico* mixtures, with their
GEO (www.ncbi.nlm.nih.gov/geo ) accession numbers or download links. We note that all these samples
were not used in the construction of the DNAm references used in EpiDISH and HEpiDISH and therefore
represent truly independent samples: Illumina 450k data of 4 epithelial cell-lines and 10 fibroblasts
cell-lines from Stem-Cell Matrix Compendium-2 (SCM2) [35] (GSE31848), Illumina 450k data of 8 B-cells,
71 CD4+ T-cells, and 28 monocytes from Absher et al [36] (GSE59250), Illumina 450k data of 31 CD4+
T-cells from Limbach et al [37] (GSE71955), Illumina 450k data of 4 B-cells, 6 CD4+ T-cells, and 5
monocytes from Mamrut et al (GSE71244), Illumina 450k data of 36 monocytes from Marabita et al [38]
(GSE43976), Illumina 450k data of 8 CD4+ T-cells from Nestor et al [39] (GSE50222), Illumina 450k data of
214CD4+ T-cells, and 1202 monocytes from Reynolds et al [40] (GSE56047), Illumina 450k data of 6
B-cells, 6 CD4+ T-cells, and 6 monocytes from Zilbauer et al [41]
(https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-2145/). Two fibroblasts cell-lines were
removed from SCM2 dataset after purity check. All of the above Illumina 450k data together made the
purified cells pool for generating *in-silico* mixtures. In total, we have 4 epithelial cell lines, 8 fibroblast
lines, 66 B-cells, 336 CD4+ T-cells, and 1277 monocytes.
We also downloaded processed WGBS bed files from IHEC (http://epigenomesportal.ca/ihec) samples that
we expected to be of relatively high purity, representing epithelial, fibroblast and immune cell-subtypes.
Samples of low WGBS coverage / sequencing depth were removed. A total of 6 epithelial (4 podocytes and
2 breast epithelial), 2 fibroblast/stromal, 8 B-cells, 11 CD4+ T-cells, and 9 monocyte samples were
included. After merging them together, the number of common CpGs was 220351. Only 381 out of the 716
cell-type CpGs in EpiFibIC reference could be found. Thus, purity estimates were checked using 381
CpGs in the EpiFibIC reference using EpiDISH. One podocyte sample was predicted to have a high
fibroblast fraction and was removed. For the in-silico mixtures, we thus used 5 epithelial, 2 fibroblast, 8
B-cells, 11 CD4+ T-cells, and 9 monocyte WGBS samples.

**Generation of *in-silico* mixtures**

To test performance of CellDMC in terms of sensitivity and specificity, we generated 200 *in-silico* mixtures with half of them representing controls and the other half representing cases (disease). For cases, we altered the DNAm levels of 150 CpGs in specific cell-types (see **SI table S1**), thus defining true DMCTs (and DMCs). For each of the 200 mixtures, we randomly sampled an epithelial cell, fibroblast, B-cell, CD4+ T-cell, and monocyte from the corresponding pools of purified cells, and mixed them together with 5 weights drawn from a Dirichlet distribution. Since for each sample, DNAm changes were made to all immune cells (B-cells, CD4+ T-cells, and monocytes) simultaneously and equally, we could treat this as a three cell-type mixture (Epi, Fib, and IC) problem, where we inferred the total IC fraction alongside the total epithelial and fibroblast fractions. We considered 5 different DMCT scenarios, including unidirectional DNAm changes in all 3 cell-types, unidirectional changes in 2 cell-types, change in 1 cell-type, bidirectional changes in 3 cell-types, and bidirectional changes in 2 cell-types. For each scenario, we also had 5 different signal to noise ratios (SNR) with values of 3, 2.4, 1.8, 1.2, and 0.9, corresponding to approximate absolute DNAm differences in individual cell-types (between cases and controls) of 0.42, 0.32, 0.22, 0.12, and 0.12, respectively. The latter two SNR values differ because we increased the noise/variance in the case where SNR=0.9. For every scenario, we ran 100 Monte-Carlo runs. The specific DMCT scenarios for the three main cell-types are summarized in **SI Table.S1**.

We note that we also considered scenarios, where a DNAm change was only induced in a particular IC cell-subtype (and not in all ICs simultaneously), in which case we used HEpiDISH to estimate the cell-type fractions for all IC cell subtypes, in addition to the total epithelial and fibroblast fractions. We also considered a scenario mimicking a blood EWAS, where we only mixed together purified IC subtypes. The overall strategy used to generate *in-silico* mixtures with a continuous phenotype is similar to that used for a binary phenotype. For each sample, the continuous phenotype $y$ was assumed to take a value between 0 and 1. In the case where $y=1$, the corresponding beta value was sampled as described previously for disease samples in the binary phenotype context, whereas for samples with values $y <1$ , the altered beta-values were correspondingly scaled by $y$. The SNR values were defined at $y=1$.

In the scenario where we allow for a bi-modal distribution within the disease phenotype, we use the same parameters as for the binary phenotype case. The only difference is that for each DMCT, only 70% of the disease samples were randomly selected to be altered, whereas the other 30% disease samples were modelled as control/normal samples.

**Perturbations of cell-type fraction point estimates**

We used two different approaches to assess the robustness of CellDMC to the use of cell-type fraction point estimates. Briefly, in one approach, instead of using the point estimates, we randomly sampled a number from the interval given by point estimate ± standard deviation*3. In the second approach, we bootstrapped  DMCs making up the reference matrix, thus generating a new reference DNAm matrix with the same number of DMC but with several of these appearing multiple times. Inference of cell-type fractions was then performed using this bootstrapped DNAm reference matrix, with the new point estimates subsequently used in CellDMC.

**Simulations involving missing cell-types in DNAm reference matrix**

To study the impact of missing cell-types, we first considered a scenario where there are 3 cell-types with one cell-type being differentially methylated (Uni-1C scenario) and another cell-type missing. In detail, we

only altered epithelial cells introducing 150 true epithelial DMCTs. Cell-type fractions were then estimated
using the EpiFibIC reference matrix but with one cell-type (fibroblasts) missing from the reference DNAm
matrix. Correspondingly, CellDMC was subsequently run with only 2 cell-types. We note that when
generating the *in-silico* mixtures, the cell-type fractions for all 3 cell-types were drawn from a uniform
Dirichlet distribution, so that each cell-type exhibited the same underlying mean and variance for the
fraction.
We also ran a second simulation but now with 2 of the 3 cell-types being altered (Uni-2C scenario), and
with one of the altered cell-types missing from the DNAm reference.
In a third analysis, we focused on blood tissue, considering the Uni-1C scenario. In more detail, we
generated *in-silico* mixtures of 5 blood cell subtypes (B-cells, CD4+ and CD8+ T-cells, Monocytes and
Neutrophils) using our pools of purified blood cell subtypes constructed earlier, drawing cell proportions
from a weighted Dirichlet distribution, with weights determined to yield realistic cell-type fractions in
blood, as inferred from a blood EWAS [5]. As before, 150 DMCTs were generated within one cell-type only
(we chose CD4+ T-cells) with sensitivity and specificity computed by CellDMC using a DNAm reference
matrix that did not include another cell-type (we chose CD8+ T-cells). As in all other simulations, total of
100 Monte-Carlo runs were performed, each run simulating 100 cases and 100 controls. We benchmarked
results against the scenario where none of the 5 cell subtypes was missing from the DNAm reference
matrix.
682
683

**Implementation of CellDMC on *in-silico* mixtures and definition of sensitivity and specificity**
We estimated cell-type fractions of all *in-silico* mixtures using EpiDISH[8] (RPC mode) with a reference [17],
which consists of 716 cell-type specific DMCs, to estimate fractions of 3 major cell-types (generic
epithelial cells, generic fibroblasts, and total immune cells). Next, CellDMC was run with the estimated
fractions, multiple hypothesis correction method set to "FDR", and FDR (adjusted P-value) threshold 0.05,
which resulted in a matrix containing predicted DMCT(s) and a matrix of coefficients for each cell-type
including predicted DNAm change, raw and adjusted P-values, ranked by selected cell-type. As long as a
CpG in one cell-type was predicted to be a DMCT, it was counted as a predicted DMC. Sensitivity to
detect DMCs was defined as the number of predicted true DMCs divided by 150 (true number of DMCs).
Sensitivity to detect DMCT was defined as the mean sensitivity to correctly predict DMCTs (considering
directionality of change) averaged over all relevant cell-types. In the case where we induced DNAm
changes in individual IC cell subtypes, we estimated cell-type fractions for all cell-types using our
HEpiDISH framework, but all subsequent analysis proceeded as for the 3 cell-type scenario above.
697

**Power calculations**
To estimate the appropriate sample size to achieve a certain sensitivity and specificity, we performed
corresponding power calculations for a DNAm change in one cell-type (Uni-1C) and unidirectional
changes in all cell-types (Uni-AllC) scenarios. For each scenario, we did 100 Monte-Carlo runs for 5
SNR(s) as described earlier. We varied sample sizes from 10 to 500 (including 10, 20, 30, 40, 50, 100, 150,
200, 300, and 500), in each case with half of these as controls and the other half as cases. Mean
sensitivities for DMCT detection over 100 Monte-Carlo runs were calculated.
705

**Simulation analysis of cell-type complexity**
For K=3, we mixed epithelial, fibroblast and monocytes and used EpiDISH with centEpiFibIC DNAm

reference to estimate cell-type fractions. For larger K values, we iteratively added one more IC cell subtype

to the mix in the order of neutrophils (K=4), CD4+ T-cells (K=5), B-cells (K=6) and CD8+ T-cells (K=7).

For K>3, we used HEpiDISH. For each choice of K and in each Monte-Carlo run, there were 150 true

DMCTs (75 hyper, and 75 hypo) occurring specifically in monocytes.

**Application of CellDMC to a blood rheumatoid arthritis EWAS to detect B-cell specific rheumatoid**

**arthritis DMCs**

We obtained Illumina 450k data profiling peripheral blood for 335 controls and 354 rheumatoid arthritis

cases from Liu et al[10] (GSE42861). Data was normalized as described elsewhere [29]. We estimated cell-type

fractions of all 7 major immune cell subtypes (B-cells, CD4T-cells, CD8T-cells, NK cells, monocytes,

neutrophils, eosinophils) using EpiDISH with a DNAm reference consisting of 333 IC cell-type specific

DMCs [16]. All parameters of CellDMC were the default choices. We tested CellDMC's predictions against

10 B-cell specific RA CpGs, which had been validated in 3 independent purified B-cell EWAS datasets, as

reported in Julià et al[11]. To assess the statistical significance of CellDMC's DMCT predictions, we

randomly picked 10 CpGs 100,000 times, and compared the weighted t-statistics of the estimated B-cell

specific regression coefficients to the observed one. P value was calculated as the number of runs, where

the statistic was larger than the original observed one, divided by the number of runs.

**Application of CellDMC to endometrial cancer**

We downloaded and processed level 3 Illumina 450k data of uterine corpus endometrial carcinoma(UCEC)

from TCGA[12]. This dataset contains 374 cancer samples and 34 normal adjacent control samples. We

estimated cell-type fractions for the total epithelial, total fibroblast and total immune cell components,

using our previously validated EpiFibIC DNAm reference (716 cell-type specific DMCs) [17].

**Application of CellDMC to a breast cancer EWAS**

To test CellDMC's ability to detect DMCTs occurring in breast cancer epithelial cells, we used a breast

cancer tissue DNAm 450k dataset, which included 92 normal controls and 305 breast cancer cases, from

Teschendorff et al[13] (GSE69914). We estimated cell-type fractions using EpiDISH in conjunction with a

previously validated DNAm reference for breast tissue, consisting of 491 cell-type specific DMCs and 4

cell-types (generic epithelial, generic fibroblast, fat and total IC) [17]. All parameters choices in CellDMC

were the default ones. The true positive breast cancer epithelial DMCTs/DMCs and a corresponding list of

true negatives were constructed as reported in Zheng et al [9,17].

**Application of CellDMC to a buccal smoking-EWAS and validation in lung squamous cell carcinoma**

The smoking buccal Illumina 450k dataset was processed as described in Teschendorff et al [26]. Using

EpiDISH and EpiFibIC DNAm reference [13], we estimated cell-type fractions of total epithelial cells, total

fibroblasts, and total immune cells. Since the estimated fractions of fibroblasts in all samples were quite

small with mean less than 0.05, we only included epithelial cells and immune cells when running

CellDMC, which was run with smoking pack years as the phenotype. Among all 790 samples in the dataset,

we only used 647 samples for which smoking pack year information was available.

DMCTs predicted to occur specifically in the epithelial compartment of buccal swabs were validated in the

Illumina 450k lung squamous cell carcinoma (LSCC) data, generated as part of The Cancer Genome Atlas

(TCGA) [32]. We only validated DMCTs which exhibited similar DNAm levels in epithelial and blood
cell-subtypes in samples not exposed to smoking, since only for these we would expect to see an
association between their DNAm levels and the estimated epithelial fraction in the tumors. To this end, we
collected Illumina 450k data of 11 different epithelial cell-lines (Hipe, Saec, Hre, Hae, Hrpe, Prec, Hee,
Hcpe, Hnpce, Hmec, Hrce) from ENCODE (GSE40699) and a total of 42 purified samples representing all
7 major immune cell types (neutrophils, eosinophils, monocytes, CD4+ and CD8+ T-cells, B-cells and
NK-cells) from Reinius et al [42] (GSE35069) to identify CpGs with the same ground state of DNAm level
in epithelial and immune cells: in detail, we used limma (an empirical Bayes framework) [43], to select
277,801 CpGs with P-values greater than 0.5 or absolute DNAm change between epithelial and immune
cells less than 0.05. Of 17,285 and 6658 DMCTs predicted by CellDMC in buccal swabs to be
hypermethylated and hypomethylated in epithelial cells, respectively, 3629 and 816 of them overlapped
with the 277,801 non-DMCs. We used EpiDISH with EpiFibIC reference to estimate cell-type fractions in
the TCGA LSCC DNAm dataset (275 cancer samples). We computed the Pearson's correlation coefficient
between the DNAm beta values of these predicted epithelial DMCTs and the estimated fraction of
epithelial cells across the 275 cancer samples. To test whether there is an association between DNAm
variance and epithelial cell-type fraction, we used the Breusch-Pagan (BP) test with beta values as
response and cell-type fraction as predictor. We used the signed BP-statistic to test for positive and
negative associations between DNAm variance with epithelial and immune cell-type fractions,
respectively.

**Data availability**
All data analyzed in this manuscript is already publicly available from Gene Expression Omnibus - GEO
(www.ncbi.nlm.nih.gov/geo/), the TCGA data portal (https://gdc.cancer.gov), International Human
Epigenome Consortium - IHEC (http://epigenomesportal.ca/ihec), or from ArrayExpress
(https://www.ebi.ac.uk/arrayexpress/). Accession codes for data from GEO include GSE31848, GSE59250,
GSE71955, GSE71244, GSE43976, GSE50222, GSE56047, GSE42861, GSE69914, GSE40699, and
GSE35069. Accession code for data from ArrayExpress is E-MTAB-2145. All DNA methylation reference
matrices used to estimate cell-type fractions are available on Github
(https://github.com/sjczheng/EpiDISH). Detailed dataset descriptions and how these data were integrated
can be found in Online Methods.

**a** CellDMC flowchart: identification of DMCTs

**b** Examples of DMCTs

**CpG1: Hyper in CT1** | **CpG1: NonDM in CT2** | **CpG1: NonDM in CT3**

**c** **CpG2: Hyper in CT1** | **CpG2: Hyper in CT2** | **CpG2: Hyper in CT3**

**d** **CpG3: Hyper in CT1** | **CpG3: Hypo in CT2** | **CpG3: NonDM in CT3**

**a** Detecting true B-cell specific RA DMCs

True B-cell specific RA DMCs | CellDMC predictions

Estimated cell-type fraction in Liu Data

Hyper
Hypo
Non

Statistical Significance
(100,000 Monte–Carlo runs)
Expected
Observed
$P < 1e{-5}$
Weighted T–stat for B–cell

**b** Detecting true breast cancer epithelial DMCs

*Fisher P* $< 2.2 * 10^{-16}$

Incons.
HypoM
HyperM

Sensitivity (DMC)
Sensitivity (DMCT Epi)
Specificity (DMC)

**c** Patterns of differential methylation vs cell-type fraction
(*HAND2*: cg01580681)

Normal
Cancer(UCEC)

Adj Beta

fEpi        fFib        fIC

**d** All *HAND2* CpG CellDMC predictions
in UCEC(TCGA)

1st Exon

Predicted Mean Methylation Diff

Cell-type
Epi
Fib
IC

Position on chromosome 4

**a** Smoking DMCTs predicted by CellDMC

**b** DMCT percentage

**c** Number of DMCTs

**d** CellDMC predictions among 62 true smoking-DMCs in blood

**e** Epi-DMCT validation in TCGA LSCC

**f** Epi-DMCT validation in TCGA LSCC

**g** Estimated cell-type fraction in TCGA LSCC

**h** DMCTs index in TCGA LSCC