Proximity sensors reveal social information transfer in maternity colonies of Greater noctule bats

Simon Ripperger* [1,2], Linus Günther[1], Hanna Wieser[1], Niklas Duda[3], Martin Hierold[3], Björn Cassens[4], Rüdiger Kapitza[4], Alexander Kölpin[5], Frieder Mayer[1,6]

[1] Museum für Naturkunde, Leibniz-Institute for Evolution and Biodiversity Science, Invalidenstraße 43, 10115 Berlin, Germany

[2] Smithsonian Tropical Research Institute, PO Box 0843-03092, Balboa, Ancón, Republic of Panama

[3] Institute for Electronics Engineering, Friedrich-Alexander University of Erlangen-Nürnberg, Wetterkreuz 15, 91058 Erlangen-Tennenlohe, Germany

[4] Carl-Friedrich-Gauß-Fakultät, Technische Universität Braunschweig, Mühlenpfordtstraße 23, 38106 Braunschweig

[5] Chair for Electronics and Sensor Systems, Brandenburg University of Technology, Siemens-Halske-Ring 14, 03046 Cottbus, Germany

[6] Berlin-Brandenburg Institute of Advanced Biodiversity Research, Altensteinstr. 34, 14195 Berlin, Germany

* Corresponding author: simon.ripperger@gmail.com

Summary

1. Bats are a highly gregarious taxon suggesting that social information should be readily available for making decision. Social information transfer in maternity colonies might be a particularly efficient mechanism for naïve pups to acquire information on resources from informed adults. However, such behaviour is difficult to study in the wild, in particular in elusive and small-bodied animals such as bats.

2. The goal of this study was to investigate the role of social information in acquiring access to two types of resources, which are crucial in the life of a juvenile bat: suitable roosting sites and fruitful feeding grounds. We hypothesized that fledging offspring will make use of social information by following informed members of the social groups to unknown roosts or foraging sites.

3. In the present study we applied for the first time the newly developed miniaturized proximity sensor system 'BATS', a fully automated system for documenting associations among individual bats both while roosting and while on the wing. We quantified associations among juveniles and other group member while switching roosts and during foraging.

4. We found clear evidence for information transfer while switching roosts, mainly among juveniles and their genetically identified mothers. Anecdotal observations suggest intentional guidance behaviour by mothers, indicated by repeated commuting flights among the pup and the target roost. Infrequent, short meetings with colony members other than the mother indicate local enhancement at foraging sites, but no intentional information transfer.

5. Our study illustrates how advances in technology enable researchers to solve long-standing puzzles. Miniaturized proximity sensors facilitate the automated collection of continuous data sets and represent an ideal tool to gain novel insights into the sociobiology of elusive and small-bodied species.

Key-words

Evolved signals, foraging, information centre hypothesis, local enhancement, maternal care, maternal guidance, *Nyctalus noctula*, roost switching

Introduction

The early development is a critical phase for an animal since it paves the way for later life by affecting survival rate and overall fitness. It has been shown for several species that limited access to food resources within the first weeks after birth has a negative impact on reproductive success (Lindström 1999; Lummaa & Clutton-Brock 2002). However, a central question is: how do offspring get access to resources in early life once they start to become independent? Behaviours related to foraging might be genetically predefined, which is the case in many invertebrates but also in vertebrate taxa (van Schaik *et al.* 2017). Over time, 'personal information' which is acquired by direct interaction with the environment (Dall *et al.* 2005) or individual learning by trial and error augment an individual's capabilities (van Schaik *et al.* 2017). If animals are born and raised in presence of their parents or other conspecifics, information may be socially acquired (Galef Jr & Giraldeau 2001). Adopting information from group members may be more efficient and less costly than individual learning and contributes to an individual's behavioural flexibility under changing conditions (Dall *et al.* 2005; van Schaik *et al.* 2017).

Information obtained within colonies or groups of conspecifics enables better decisions in various contexts such as predator avoidance, reduction of parasitism, habitat choice and foraging (Evans, Votier & Dall 2016). The mechanisms of acquiring social information may vary widely in complexity. Individuals may use 'inadvertent social information', which is generated by social cues of conspecifics, i.e., eating may inform about the location of food or

fleeing about the presence of a predator (Dall *et al.* 2005). Such public information is created

non-deliberately and in group-living animals it may be difficult to hide certain information,

e.g., on foraging success. This might particularly apply for breeding colonies, where parents

must return to their young and may inadvertently inform others on foraging success via, e.g.,

time of arrival or fatness (Evans, Votier & Dall 2016). If information is provided deliberately,

'evolved signals' are used to actively exchange information (Dall *et al.* 2005). Black-capped

chickadees, e.g., broadcast alarm calls which contain information on the presence and even

the size of a predator (Templeton, Greene & Davis 2005) and honey bees use the waggle

dance to inform conspecifics on the location of a food source (Leadbeater & Chittka 2007).

The ready availability of information at communal roosts gave rise to the Information Center

Hypothesis (ICH), which states that such assemblages primarily evolved for the efficient

exploitation of unevenly distributed food sources (Ward & Zahavi 1973). According to the

ICH colony members must assess the success of returning foragers, which are later followed

to food patches. However, information can also be generated right at the resource, a

mechanism termed 'local enhancement'. Black-browed albatrosses, e.g., indirectly detect food

patches by approaching aggregations of foraging predators over sea (Grünbaum & Veit 2003).

Both advertent and inadvertent information can be used by juveniles to find food. Juvenile

rats prefer feeding sites where adults are present and scent marks and trails of adults cause

juveniles to explore such sites (reviewed by Galef Jr and Giraldeau (2001)). Juveniles, since

they are naïve, seem to particularly rely on acquiring social information from more

experienced individuals of the group like their parents (termed 'vertical transmission') or

other adults ('oblique transmission') (van Schaik *et al.* 2017).

Bats are an ideal taxon to study the mechanisms of social information use in groups since the

vast majority of species is gregarious and long-lived (Wilkinson & Boughman 1999; Kerth

2008; Smith, Lacey & Hayes 2017). However, there is surprisingly little known on whether

and how juveniles benefit from social information provided by group members. An interesting

case of information transfer across generations is reported for greater sac-winged bats. Here,

vocal development of pups is influenced by imitation of territorial songs of harem males and

leads to a group signature which is independent of relatedness (Knörnschild *et al.* 2010;

Eckenweber & Knörnschild 2013). When it comes to learning where to roost and where to

forage, knowledge on juveniles becomes scarce and existing literature focusses on horizontal

information transfer among adult peers. Empirical studies in the wild demonstrated that

several insectivorous species are attracted to feeding buzzes by conspecifics which may be

used as a signal of foraging success (Gillam 2007; Dechmann *et al.* 2009; Cvikel *et al.* 2015).

Female greater spear-nosed bats coordinate their foraging bouts at the day roost by screech

calls (Wilkinson & Boughman 1998). However, neither local enhancement nor recruitment at

the roost has been demonstrated in foraging juveniles, so far. Spatial association among home

ranges of mothers and offspring in at least three species and simultaneous feeding of mother-

pup pairs in vampire bats suggest that vertical information transfer, possibly in form of

following behaviour, might provide juveniles with insights on where to forage, but this

mechanism has yet has to be demonstrated (Wilkinson 1995; Schnitzler, Moss & Denzinger

2003).

Similarly little is known on how juveniles learn about the location of suitable roosts and the

few existing studies only involved adults. In Greater noctule bats local enhancement by

inadvertent acoustic cues significantly reduces the time required to locate a roost both in

captive and in wild experiments (Ruczyński, Kalko & Siemers 2007; Furmankiewicz *et al.*

2011). Kerth and Reckardt (2003) tracked nightly roost switching behaviour in Bechstein's

bats and assumed that recruitment of naïve by informed individuals already started at the day

roost, but could not unequivocally prove it. An exclusion experiment by Wilkinson (1992)

demonstrated that juvenile bats are able of relocating at a new roost with their mothers and the

author concluded that following behaviour is the only plausible explanation. One of the still

standing mysteries in bat ecology is how juveniles of temperate species locate swarming or

hibernation sites, which are often long distances from where they are born and reared. It has

been hypothesized that mothers guide their offspring (Sachteleben 1991), but so far nobody

was able to track such guidance behaviour. These examples emphasize that there are plenty of

indications on following and guidance behaviour in juvenile bats, but so far technological

limitations prevented the final proof.

While emerging technologies have revolutionized the field of bio-logging and in turn our

understanding of behaviour of wild animals during the past decades, studies on small-bodied

vertebrates still lag behind due to the scarcity of fully automated lightweight devices (Kays *et*

al. 2015). Proximity loggers represent a powerful tool for the study of information transfer

(Rutz *et al.* 2015; St Clair *et al.* 2015), but studies making use of such devices are generally

rare and the loggers used in the aforementioned studies are by far too heavy for tagging

medium-sized bats. Smaller tag versions of acceptable weight, however, show dramatically

reduced runtimes of less than 24 h (Levin *et al.* 2015). In the present study we used the newly

developed miniaturized proximity sensor system 'BATS', a fully automated system for

documenting associations among individuals at a tag weight of one to two gram and runtimes

of at least one to two weeks (Duda, Weigel & Koelpin 2018). Our developments enable us to

study interactions among tagged bats both while roosting and while on the wing. Here we

report on the first extensive study to apply our system and proximity sensors to free-ranging

bats, in general.

The goal of this study was to investigate the use of social information in acquiring access to

two types of resources, which are crucial in the life of a juvenile bat: suitable roosting sites

and fruitful feeding grounds. We hypothesized that fledging offspring will make use of social

information by following either the mother or other informed members of the social groups to

unknown roosts or foraging sites. If juveniles use social information when switching roosts,

we expect that the successfully switching juvenile will be associated with at least one

individual of the group shortly before and shortly after leaving the current roost and shortly

before and shortly after arriving at the new roost. If social information is used for finding

foraging grounds, we would expect juveniles to be associated with at least one roosting

partner shortly before and shortly after starting the bout, during several minutes after starting

while commuting to the foraging ground, and possibly, but not necessarily when returning to

the roost. The BATS-tracking system enabled us to classify and quantify the aforementioned

events.

2. Materials and methods

2.1 Field site and study species

This study was conducted in "Königsheide Forst", a mixed forest in Berlin, Germany, from

June to August 2016 and 2017, respectively. The study site comprises ample of roosting

opportunities for bats such as natural tree holes and roughly 130 bat boxes. During this time

of the year females of the common noctule bat (*Nyctalus noctula*) form temporary groups, so

called maternity colonies, to jointly give birth and rear their young. Mothers give birth to one

or two offspring and individuals of a maternity colony frequently switch roosts, but usually

stay within the area of the nursing colony. Moving among roosts may involve a change in

group composition. However, strong, non-random inter-individual bonds have been observed

in captive studies as well as a certain degree of maternal care such as allogrooming of

offspring (Kleiman 1969; Kozhurina 1993).

The ideal opportunity to observe information transfer in maternity colonies should be the

moment when the offspring start to fledge in order to track their behaviour during the first

nights of independent flight. Therefore, we daily monitored the bat boxes, including checks

after sunset when adults and already flying juvenile had emerged from the roost. We aimed at

tagging the majority of a social group including juveniles, which have started fledging only

recently or which not fledged at all, yet. We therefore prepared to capture on the following

day when around a third of the offspring were still inside the roost while the rest of colony

(including already fledged youngsters) was foraging.

2.2 Sample collection, molecular analysis and identification of mother pup pairs

In 2016 we captured a social group from a single bat box while in 2017 bats were caught from

two different bat boxes which were roughly 300m apart. Bats were kept in cotton cloth bags

until they were weighted, sexed and the forearm was measured using a calliper. If the

epiphyseal gaps were closed and the phalangeal–metacarpal joints were knobby, individuals

were considered adult (Brunet-Rossinni & Wilkinson 2009). We collected tissue samples with

a biopsy punch (Ø 4 mm, Stiefel Laboratorium GmbH, Offenbach, Germany) and preserved

them in 80% ethanol. In the lab we used the salt–chloroform procedure (Miller, Dykes &

Polesky 1988) modified by Heckel *et al.* (1999) for DNA isolation.

We used the DNA Analyser 4300 and the SAGA^{GT} allele scoring software (both: LI–COR

Biosciences, Lincoln, NE, USA) to genotype a total of 75 individuals (n = 33 adult

females, n = 42 juveniles) at 9 polymorphic microsatellite loci. We used the loci P11, P217,

P219 and P223 which were isolated from the focus species *Nyctalus noctula* (Mayer,

Schlötterer & Tautz 2000). Nleis3 and Nleis4 were isolated from the closely related *Nyctalus*

leisleri (Boston, Montgomery & Prodöhl 2009) and G6-Mluc, G31-Mluc, H23-Mluc and

H29-Mluc have originally been isolated from *Myotis myotis* (Castella & Ruedi 2000), but

were subsequently modified for cross-species utility in vespertilionid species (Jan *et al.* 2012).

To calculate allele frequencies all adult individuals from both years (n=33) were used. All

individuals were genotyped at least at eight loci, and genotypes were 99.7% complete. See

Table S1 in supporting information for allele numbers per locus, results of Hardy–Weinberg

tests, null allele frequencies, and non-exclusion probabilities for the nine microsatellite

markers.

Parentage analyses were performed with CERVUS v. 3.0 (Kalinowski, Taper & Marshall

2007) separately for the social groups caught in 2016 and 2017, respectively, since our

objective was to identify mother-pup pairs within year, not across years. The 2016 data set

comprised 20 juveniles and 13 adult females (candidate mothers), while 22 juveniles and 24

candidate mothers were used for 2017. Four of the 24 adult females in 2017 were recaptures

that were already caught in 2016 as juvenile (n=1) and adults (n=3).

Simulations were run with 100 000 cycles, a proportion of 80% sampled candidate mothers,

an estimated genotyping error of 2%, and for two confidence levels (80% and 95%). One

mismatch per mother–offspring dyad was accepted to account for genotyping errors or

mutations. A mother could be assigned to 40 (2016: n=18; 2017: n=22) of the 42 analysed

juveniles. Thirty-three mother-pup pairs were assigned at 95 % confidence with no mismatch,

six at 95 % confidence with one mismatch and only one with 80 % confidence and one

mismatch.

2.3 Automated encounter detection among tagged bats

Our team developed a tracking system for direct encounter detection, which bases on wireless

sensor network technology for field strength related distance estimation between individuals.

The system is fully automated including remote data download and does not require

recapturing tagged animals thus reducing disturbance of the animals to a minimum. The

centrepiece of the tracking hardware is the animal-borne mobile node, in the following

referred to as 'proximity sensor'. Once deployed, a wake-up receiver on the proximity sensor

permanently scans its surroundings for signals of other proximity sensors, which are

constantly broadcasted every two seconds. This operation mode is independent of any further

infrastructure. Whenever one or more tracking sensors are within reception range of ca. 10 m

maximum distance (Ripperger *et al.* 2016), a so called 'meeting' is created. As soon as no

signal has been received by the respective meeting partner for five sending intervals

(corresponding to 10 seconds), the meeting is closed and stored to on-board memory along

with the ID of the meeting partner, a timestamp, total meeting duration, and a maximum

signal strength indicator (RSSI) of the meeting. The signal, which is broadcasted every 2 s, is

simultaneously used as an indicator of presence at a site of interest, e.g. a roost, when the

signal is received by a stationary node, in the following referred to as 'base stations'. We

positioned base stations near potential roosts to detect presence of individual tagged bats

inside a particular bat box or tree hole and we therefore termed a bat signal which are picked

up by base stations 'presence signal'. Base stations also provide remote data download, while

all downloaded data is locally stored and can be accessed by the user. In 2016 the system

could operate a maximum of 30 IDs at a time, while in 2017 the maximum number of

observable individuals has been increased to 60. In the following we give a brief overview of

the hardware components and the functionality of the system. For an elaborate, in-depth

description of the software see Cassens *et al.* (2017) and for hardware see Duda, Weigel and

Koelpin (2018).

2.3.1 Proximity sensors: We used a refined version of miniaturized proximity sensors, which

has been described and tested in free-ranging bats first in Ripperger *et al.* (2016). The

proximity sensor comprises a System-on-Chip (SoC) for communication control and on-board

data processing, a transceiver which enables communication in the 868MHz frequency band

with other proximity sensors or base stations and a wake-up-receiver which activates full

system functionality from an energy-saving low-power mode whenever communication

partners are in range. A lithium-polymer battery powers the mobile node. We built two

versions of the proximity sensor that differ in weight since adult females and offspring of

noctule bats varied considerably in body weight. The low-weight version for tagging offspring

was equipped with a 15 mAh battery and was housed in the fingertip of a nitrile lab glove.

The heavier version for adult females was equipped with either two 15 mAh battery of a

single 24mAh batteries and housed in a 3D-printed plastic case ensuring longer runtime. The

different proximity sensor versions resulted in a total weight of 1.1 to 1.9 g including battery

and housing.

2.3.2 Base stations and data access

The base station contains a receiver for the reception of presence signals and transmitted data.

Presence signals and downloaded data are stored by a Raspberry Pi (Raspberry PI Foundation,

Cambridge, UK) to a SD card along with the ID of the transmitting proximity sensor and the

receiving base station, respectively, and a timestamp which is provided by a GPS unit. At the

same time the Raspberry Pi hosts a WiFi hotspot allowing the user remote data access. The

data is then stored in a MySQL database.

2.4 Tagging and data collection

On July 15[th] 2016 we tagged in total 26 individuals, 10 juveniles and 16 adult females, from a

single bat box. On July 18[th] 2017 we tagged in total 34 individuals, 19 juveniles and 15 adult

females from two bat boxes. This adds up to a total of 60 tracked bats, 29 of which were

juveniles and 31 were adult females. According to individual body weight we used different

versions of the proximity sensors. Bodyweight ranged from 17g to 25g for juveniles and

averaged at 21.26g +/- 2.04, while adult body weight ranged from 23.5g to 35g at an average

of 27.73g +/- 2.27. Individual tag-to-body weight ratios ranged from 4.4% to 7% for juveniles

and from 4.2% to 8% in adults, which is well within the recommendations for short-term

biologging studies in bats (Amelon *et al.* 2009; O'Mara, Wikelski & Dechmann 2014).

Proximity sensors were glued to the fur on the back of the bats using surgical cement (Perma-

Type, Plainville, CT, USA) and drops off when the cement loses its tackiness.

Data collected during the first night after the tagging event was discarded to account for

potential behavioural changes right after tagging and actual data collection started the night

after in order to allow the bats to get used to the tag. In 2016 data collection lasted until July

28th (12 days) and in 2017 until August 8th (20 days). We installed three respectively five base

stations in 2016 and 2017 at day roosts to download data and to receive presence signals for

individual bats. Whenever bats switched to unknown roosts we used a handheld 868 MHz

panel antenna (HSP-868C, WiMo, Herxheim, Germany) connected to a base station to

localize the unknown roost and relocate base station.

2.5 Analysis of tracking data

We used the library RMySQL in R (James & DebRoy 2012) to access the data, which were

managed in HeidiSQL, a Windows client for MariaDB. In a first step we plotted and visually

explored presence signals and meetings received at base stations to define foraging bouts and

roost switching events, respectively, for all individual juveniles (see Fig. 2a-c for examples).

To evaluate potential information transfer we queried the meeting database for events, which

matched the timestamp of foraging bouts or roost switches, respectively. If information

transfer would play a role during foraging or roost switching, we would expect to find

meetings among offspring and other group members associated with these events. In detail,

we proceeded as follows.

2.5.1 Evaluation of information transfer during roost switching

We defined a roost switch as an event during which an individual changes its roost and

potentially its roosting partners without prolonged absence times which may indicate

foraging. A roost switch can be detected if a bat switches between two roosts which are both

equipped with a base station receiving presence signals (Fig. 2b). If at least one roost is

equipped with a base station, presence signals can be used to determine departure time or

arrival, respectively. If the unmonitored roost is occupied by other tagged bats (indicated by

reciprocal, stable meetings) we can at least unequivocally classify this event as a roost switch

(Fig. 2c). However, we cannot determine the time of arrival respectively departure at the unmonitored roost because bats may leave jointly. If juveniles use social information when switching roosts, we expect that the switching juvenile will be associated with at least one individual of the group shortly before and shortly after leaving the current roost or arriving at the new roost. To this end we define the moment of departing from or arriving at a monitored roost, respectively, when the steady reception of signal beacons at a base station gets cut off or starts. We subsequently queried the meeting database for meetings which are active or which started within 60s before and within 60s after the moment of leaving or arriving at a roost.

2.5.2 Evaluation of information transfer during foraging bouts

We defined a foraging bout as an event where an individual starts from a known roost, returns to the same roost and does not visit other monitored roosts or roosts with tagged bats (indicated by stable, lasting meetings) in between (Fig. 2a). We chose these strict rules to ensure that the events we are looking at relate to foraging and do not overlap with roost switching events. If social information would play a role in locating foraging grounds we would expect a juvenile to associate with at least one roosting partner upon starting the bout, during several minutes after departure while commuting to the foraging ground, and possibly, but not necessarily when returning to the roost. As described above we equally defined the start and the end of the foraging bout as the end and the start of the steady reception of the presence signal, respectively. We then queried all meetings which were ongoing or started within 60s before and within 60s after starting a foraging bout and returning, respectively. In addition, we queried all meetings which originated during the entire foraging bout.

2.5.3 Statistical testing

We used a Mantel-Test to test whether social information used by juveniles is obtained by

their mothers in the first place or by any roosting partner. To this end we created a binary

matrix containing "1" for dyads which have been associated while roost switching and "0" for

dyads which have never been observed switching communally. Accordingly, foraging

associations were transformed into a binary martrix. For testing the effect of maternity we

created a second binary correlation matrix which listed the genetically determined identity of

mother-pup pairs as "1", while all other dyads were marked "0". We tested the years 2016 and

2017 separately and ran Mantel tests in the library "ade4" version 1.7-11 in RStudio 1.1.453

using Monte-Carlo permutation tests with 9999 replicates (Dray & Dufour 2007; R

Developing Core Team 2015).

345   3. Results

3.1 Genetic analyses

Mother and juvenile bats were caught in day roots at the time of weaning. In 24 determined

mother-pup pairs, both individuals were tagged with proximity sensors (2016: n=9, all

assigned at 95 % confidence with no mismatch; 2017: n=15, 12 pairs assigned at 95 %

confidence with no and three pairs at 95 % confidence with one mismatch). These 24 mother-

pup pairs generated the data for the following section.

3.2 Tracking results

In 2016 we received a total of 561,795 presence signals and 13,292 meetings from 23

individual bats and in 2017 we received 2,667,409 localization signals and 53,391 meetings

from 33 individual bats. One individual in 2016 and three individuals in 2017 did not get in

contact with base stations. These four individuals may have left the study area between

tagging and the following night.

3.2.1 Evaluation of joint roost switching events

To evaluate information transfer on roosts we screened the data set for joint departures from

and joint arrivals at roosts for all tagged juveniles. In 2016 we observed ten events of seven

individual juveniles being associated with another individual while switching among two

roosts. In all except one event the associated bats arrived together at a new roost, even though

successful switching took several approaches in two cases and temporary roosts may be used

in between (Table 1, Fig. 3). In six cases both roosts have been monitored by a base station, in

two cases the juveniles left a monitored roost and switched to a roost where other tagged bats

have been roosting and in the remaining two cases the juveniles switched from a monitored

roost to an unknown roost where no other tagged bats were present, except the one which

accompanied the juvenile during switching. In all 10 cases the juvenile was in company of its

identified mother and no other tagged bat.

In 2017 we observed six events where 5 individual juveniles switched roosts in company.

Twice, the juvenile switched among two monitored roosts and four times among one

monitored and an unmonitored site. Twice, the juvenile was associated with its identified

mother, in four cases with another adult female.

Some juveniles switched directly among roosts. Such events took only seconds to minutes

(see table xx). During other events stopover sites were used and several attempts of mothers

re-associating with their young were necessary before both arrived at the new roost. Such

unsuccessful tandem flights underline that the offspring was actively flying and not carried by

the mother.

In both years significantly more mother-pup dyad have been observed switching roosts

communally than expected by chance (Mantel tests, 9999 replicates; 2016: r = 0.88, p <

0.001; 2017: r = 0.21, p < 0.01).

3.2.2 Associations during foraging bouts

In total we detected 42 foraging bouts of juveniles, which matched our definition above, conducted by 13 individuals (2016: four juveniles, eight bouts; 2017: nine juveniles, 34 bouts). Foraging bouts lasted on average 1:14:53 h with a standard deviation of 36:19 min. During 6 of these 42 foraging bouts (14 %, n = 7 individual juveniles, all 2017) we detected in total 28 short meetings, which lasted between 1 and 30 seconds (average: 7.4 s +/- 8.6). Two of these meetings occurred within less than 90 s after two co-roosting individuals left a roost; however, no further meetings have been documented during these foraging bouts. All remaining meetings originated at least several minutes after emergence from the roost. Eight times the meeting partner was another juvenile and twice an adult female. Only in one case the meeting partner was the identified mother. Accordingly, meetings among mother-pup dyads have not been observed more often than expected by chance (Mantel tests, 9999 replicates, r = 0.08, p > 0.05).

In 13 out of the 42 foraging bouts of juveniles (2016: n = two individuals; 2017: n = seven individuals) the identified mother was co-roosting before both started a bout. In all 13 cases the mother started its foraging bout considerably earlier than the juvenile (between 4:31 min and 1:26:02 h, average: 36:45 min (+/- 24:57)).

Discussion

The study of information transfer in free-ranging bats is particularly challenging due to their small body size and their elusive, nocturnal life. We tracked bats using novel, miniaturized proximity sensors and demonstrated that juveniles use social information of group members and for finding roosts mothers seem to intentionally guide their young. However, during foraging mothers did not guide their offspring, but meetings with other colony members may reflect local enhancement at feeding grounds.

To the best of our knowledge our study shows for the first time that recruitment to a new roost starts already at the occupied roost. Furthermore, the repeated commuting flights we observed

in at least two cases until the juvenile arrives at the target roost represents anecdotal evidence

that at least in some cases deliberate, evolved signals rather than inadvertent social cues are

used. The existence of evolved signals and the strong bias towards information transfer

among mother-pup pairs suggests that the observed behaviour is best explained by kin

selection. Some studies have reported on the use of social information in bats for finding

suitable roosts, however, studies are scarce and the mechanisms are in parts poorly

understood, in particular when it comes to naïve juveniles. Studies on a range of

vespertilionid species including the focus species *N. noctula* have shown that conspecific calls

enhance roost finding efficiency in captive experiments as well as in the wild (Ruczyński,

Kalko & Siemers 2009; Schöner, Schöner & Kerth 2010; Furmankiewicz *et al.* 2011). These

studies demonstrate that bats may eavesdrop on vocalizations to localize an occupied roost

once within hearing distance. Since playbacks from varying contexts have been used we

conclude that the studied bats relied on inadvertently broadcasted public information. On the

contrary, Spix's disk winged bats deliberately produce signals to facilitate group cohesion, by

a remarkable call-and-response system among flying bats in search of a roost and bats

occupying a roost (Chaverri, Gillam & Vonhof 2010; Chaverri, Gillam & Kunz 2012). A

common theme of all abovementioned studies is that the mechanism of recruitment of

conspecifics is best explained by local enhancement, i.e. the social information is acquired at

the new roost, when searching bats are in hearing distance. Kerth and Reckardt (2003) were

first to present experimental evidence for information transfer about roosts in bats. The

authors presumed that naïve Bechstein's bats are recruited to a novel roost already at the

dayroost by experienced conspecifics, however, they could not unequivocally exclude local

enhancement at the target roost. Our study finally demonstrates that this inferred mechanism

does exist in roost-switching bats.

We classify the advertent information transfer from mothers to their young as a form of

maternal care which has to the best of our knowledge not been observed in free-ranging bats,

so far. Mammalian offspring is usually strongly dependent on maternal care for food,

protection and warmth (Balshine 2012) and maternal investment in young is also wide-spread

in bats (Smith, Lacey & Hayes 2017). Besides weaning maternal care has been demonstrated

in form of post-weaning food provisioning (Wilkinson 1990; Geipel *et al.* 2013), grooming

(Kleiman 1969; Wilkinson 1986; Kozhurina 1993) or pup guarding (Bohn, Moss &

Wilkinson 2009). Carrying young in flight is also commonly observed and Jones (2000)

summarizes some reports where young are possibly carried to temporary roosts or feeding

grounds. However, this is the first study to document maternal guidance to roosts, which has

been hypothesized as a plausible explanation for young to reach swarming and hibernation

sites, but could not be confirmed, possibly due to the lack of appropriate tracking technology

(Sachteleben 1991; Burns & Broders 2015; Stumpf *et al.* 2017).

Previous work on bats indicated that roosts may act as information centres where bats may

obtain information on food by inadvertent cues (Ratcliffe & ter Hofstede 2005; O'Mara,

Dechmann & Page 2014). A considerable part of the diet of Greater noctule bats consists of

insects, which fly in swarms and often over water (Gloor, Stutz & Ziswiler 1995). Such rich

and patchy, but ephemeral foraging sites are required for the establishment of information

centres (Ward & Zahavi 1973) and juveniles in particular might benefit from rich food

patches when collecting experience on where and how to forage. However, we did not

observe recruitment at the roost to feeding grounds in young noctules, which complies with

most foregoing studies that showed that 'ICH' operates in colonial roosts, but is rarely

demonstrated in breeding colonies (summarized by Evans, Votier and Dall (2016)). Our

observation that juveniles start foraging bouts considerably later than their mothers suggests

that juvenile noctules conduct opportunistic, explorative foraging flights. Rare and short

contacts to tagged colony members other than the mother during foraging bouts suggest that

local enhancement by eavesdropping on conspecifics while hunting may play a role as it has

been shown for several insectivorous bat species (Gillam 2007; Dechmann *et al.* 2009; Cvikel

et al. 2015). However, our data cannot unequivocally prove this theory since the exact context

of the meetings remains unknown.

Our observations raise the following question: Why is social information transfer among

mothers and their offspring context dependent? One possible explanation is that group

cohesion is crucial for energy-saving social warming and prolonged lactation periods in bats

require mothers to stay in contact with their young for 3 weeks to 2 months depending on the

species (reviewed by Kerth (2008)). Extended weaning, which was observed in captive

noctules for up to 2 months (Kleiman 1969), and the broad spectrum of insects they feed on

(Gloor, Stutz & Ziswiler 1995) may in turn enable juveniles to forage opportunistically and –

if available – make use of social information by local enhancement. In general, suitable roosts

of high quality may be harder to find opportunistically than insect prey and information on

roosts is likely to accumulate in adults, in particular in philopatric females. This should favour

information transfer on roosts since failing to relocate at an occupied roost might be more

costly than low foraging success, which might subsequently be balanced by extended

weaning. Adverse climatic conditions may have detrimental effects on single bats (Lindström

1999) and might therefore be a strong driver of the evolution of the observed guidance

behaviour, since local enhancement by vocalization at the roost (Furmankiewicz *et al.* 2011)

might not be functional for long-distance localization of roosting partners.

Conclusions

Bats are facing ideal prerequisites for social information transfer, since they are long-lived

and the vast majority of species is living in group. Regarding information use in offspring

Wilkinson and Boughman (1999) speculated already 20 years ago that young bats almost

certainly follow adults in situations other than foraging. However, this is also how long it took

to unequivocally track mother-pup pairs switching among roosting sites. Our study shows that

the current revolution in tracking technology provides powerful tools to investigate

behavioural ecology and sociobiology in free-ranging small bodied animals such as bats.

Authors' contributions

491 SR and FM conceived the ideas and designed the sampling scheme; MH, ND, AK, BC and

492 RK developed and tested all tracking equipment. SR, LG, FM and HW collected the data; SR

493 and LG analysed the data; SR and FM led the writing of the manuscript. All authors

494 contributed critically to the drafts and gave final approval for publication.

Acknowledgements

497 We thank I. Waurick, S. Hayden and E. Jäger for support during molecular lab work. We are

498 grateful to T. Teige for his help and his valuable expertise during field work. All necessary

499 permits have been obtained from SenStadtUm (I E 222/OA-AS/G_1203) and LaGeSo (I C

500 113-G0008/16).

Data accessibility

503 Data will be deposited on GFBio and made available upon acceptance.

Literature

Amelon, S.K., Dalton, D.C., Millspaugh, J.J. & Wolf, S.A. (2009) Radiotelemetry techniques

and analysis. *Ecological and behavioral methods for the study of bats* (eds T.H. Kunz

& S. Parsons), pp. 57-77. The John Hopkins University Press, Baltimore, Maryland.

Balshine, S. (2012) Patterns of parental care in vertebrates. *The evolution of parental care*

(eds N.J. Royle & P.T. Smiseth), pp. 376. Oxford University Press Oxford

Bohn, K.M., Moss, C.F. & Wilkinson, G.S. (2009) Pup guarding by greater spear-nosed bats.

Behavioral Ecology and Sociobiology, **63,** 1693-1703.

Boston, E., Montgomery, I. & Prodöhl, P.A. (2009) Development and characterization of 11

polymorphic compound tetranucleotide microsatellite loci for the Leisler's bat,

Nyctalus leisleri (Vespertilionidae, Chiroptera). *Conservation Genetics,* **10,** 1501.

Brunet-Rossinni, A.K. & Wilkinson, G.S. (2009) Methods for age estimation and the study of

senescence in bats. *Ecological and behavioral methods for the study of bats* (eds T.H.

Kunz & S. Parsons), pp. 315-325. Johns Hopkins University Press, Baltimore.

Burns, L.E. & Broders, H.G. (2015) Who swarms with whom? Group dynamics of *Myotis*

bats during autumn swarming. *Behavioral Ecology,* **26,** 866-876.

Cassens, B., Ripperger, S., Hierold, M., Mayer, F. & Kapitza, R. (2017) Automated

Encounter Detection for Animal-Borne Sensor Nodes. *Proceedings of the 2017*

International Conference on Embedded Wireless Systems and Networks, pp. 120-131.

Junction Publishing.

Castella, V. & Ruedi, M. (2000) Characterization of highly variable microsatellite loci in the

bat Myotis myotis (Chiroptera: Vespertilionidae). *Molecular Ecology,* **9,** 1000-1002.

Chaverri, G., Gillam, E.H. & Kunz, T.H. (2012) A call-and-response system facilitates group

cohesion among disc-winged bats. *Behavioral Ecology,* **24,** 481-487.

Chaverri, G., Gillam, E.H. & Vonhof, M.J. (2010) Social calls used by a leaf-roosting bat to

signal location. *Biology Letters,* **6,** 441-444.

Cvikel, N., Egert Berg, K., Levin, E., Hurme, E., Borissov, I., Boonman, A., Amichai, E. &

Yovel, Y. (2015) Bats Aggregate to Improve Prey Search but Might Be Impaired

when Their Density Becomes Too High. *Current Biology,* **25,** 206-211.

Dall, S.R., Giraldeau, L.-A., Olsson, O., McNamara, J.M. & Stephens, D.W. (2005)

Information and its use by animals in evolutionary ecology. *Trends in Ecology &*

Evolution, **20,** 187-193.

Dechmann, D.K., Heucke, S.L., Giuggioli, L., Safi, K., Voigt, C.C. & Wikelski, M. (2009)
      Experimental evidence for group hunting via eavesdropping in echolocating bats.
      *Proceedings of the Royal Society of London B: Biological Sciences***,** rspb. 2009.0473.

Dray, S. & Dufour, A.-B. (2007) The ade4 package: implementing the duality diagram for
      ecologists. *Journal of statistical software,* **22,** 1-20.

Duda, N., Weigel, R. & Koelpin, A. (2018) Enhanced mobile node design for small size
      animal borne wireless sensor nodes with encounter detection and localization.
      Microwave Conference (GeMiC), 2018 11th German, pp. 123-126. IEEE.

Eckenweber, M. & Knörnschild, M. (2013) Social influences on territorial signaling in male
      greater sac-winged bats. *Behavioral Ecology and Sociobiology,* **67,** 639-648.

Evans, J.C., Votier, S.C. & Dall, S.R. (2016) Information use in colonial living. *Biological
      reviews,* **91,** 658-672.

Furmankiewicz, J., Ruczyński, I., Urban, R. & Jones, G. (2011) Social calls provide
      tree-dwelling bats with information about the location of conspecifics at roosts.
      Ethology, **117,** 480-489.

Galef Jr, B.G. & Giraldeau, L.-A. (2001) Social influences on foraging in vertebrates: causal
      mechanisms and adaptive functions. *Animal Behaviour,* **61,** 3-15.

Geipel, I., Kalko, E.K.V., Wallmeyer, K. & Knörnschild, M. (2013) Postweaning maternal
      food provisioning in a bat with a complex hunting strategy. *Animal Behaviour,* **85,**
      1435-1441.

Gillam, E. (2007) Eavesdropping by bats on the feeding buzzes of conspecifics. *Canadian
      journal of zoology,* **85,** 795-801.

Gloor, S., Stutz, H.-P.B. & Ziswiler, V. (1995) Nutritional habits of the noctule bat Nyctalus
      noctula (Schreber, 1774) in Switzerland. *Myotis,* **32,** 231-242.

Grünbaum, D. & Veit, R.R. (2003) Black-browed albatrosses foraging on Antarctic krill:
      density-dependence through local enhancement? *Ecology,* **84,** 3265-3275.

Heckel, G., Voigt, C.C., Mayer, F. & Von Helversen, O. (1999) Extra-harem paternity in the

white-lined bat Saccopteryx bilineata (Emballonuridae). *Behaviour,* **136,** 1173-1185.

James, D. & DebRoy, S. (2012) RMySQL: R interface to the MySQL database (2011). R

package version 0.8-0.

Jan, C., Dawson, D.A., Altringham, J.D., Burke, T. & Butlin, R.K. (2012) Development of

conserved microsatellite markers of high cross-species utility in bat species

(Vespertilionidae, Chiroptera, Mammalia). *Molecular Ecology Resources,* **12,** 532-

548.

Jones, G. (2000) The ontogeny of behavior in bats: a functional perspective. *Ontogeny,*

functional ecology, and evolution of bats (eds R. ADAMS & S. PEDERSEN), pp.

362-392. Cambridge University Press, Cambridge.

Kalinowski, S.T., Taper, M.L. & Marshall, T.C. (2007) Revising how the computer program

CERVUS accommodates genotyping error increases success in paternity assignment.

Molecular Ecology, **16,** 1099-1106.

Kays, R., Crofoot, M.C., Jetz, W. & Wikelski, M. (2015) Terrestrial animal tracking as an eye

on life and planet. *Science,* **348,** aaa2478.

Kerth, G. (2008) Causes and consequences of sociality in bats. *Bioscience,* **58,** 737-746.

Kerth, G. & Reckardt, K. (2003) Information transfer about roosts in female Bechstein's bats:

an experimental field study. *Proceedings of the Royal Society of London, Series B:*

Biological Sciences, **270,** 511-515.

Kleiman, D.G. (1969) Maternal care, growth rate, and development in the noctule (Nyctalus

noctula), pipistrelle (Pipistrellus pipistrellus), and serotine (Eptesicus serotinus) bats.

Journal of Zoology, **157,** 187-211.

Knörnschild, M., Nagy, M., Metz, M., Mayer, F. & von Helversen, O. (2010) Complex vocal

imitation during ontogeny in a bat. *Biology Letters,* **6,** 156-159.

Kozhurina, E.I. (1993) Social Organization of a Maternity Group in the Noctule Bat, Nyctalus

noctula (Chiroptera: Vespertilionidae). *Ethology,* **93,** 89-104.

Leadbeater, E. & Chittka, L. (2007) Social Learning in Insects — From Miniature Brains to

Consensus Building. *Current Biology,* **17,** R703-R713.

Levin, I.I., Zonana, D.M., Burt, J.M. & Safran, R.J. (2015) Performance of Encounternet

Tags: Field Tests of Miniaturized Proximity Loggers for Use on Small Birds. *PLoS*

ONE, **10,** e0137242.

Lindström, J. (1999) Early development and fitness in birds and mammals. *Trends in Ecology*

& Evolution, **14,** 343-348.

Lummaa, V. & Clutton-Brock, T. (2002) Early development, survival and reproduction in

humans. *Trends in Ecology & Evolution,* **17,** 141-147.

Mayer, F., Schlötterer, C. & Tautz, D. (2000) Polymorphic microsatellite loci in vespertilionid

bats isolated from the noctule bat Nyctalus noctula. *Molecular Ecology,* **9,** 2209-2212.

Miller, S., Dykes, D. & Polesky, H. (1988) A simple salting out procedure for extracting

DNA from human nucleated cells. *Nucleic Acids Research,* **16,** 1215.

O'Mara, M.T., Wikelski, M. & Dechmann, D.K. (2014) 50 years of bat tracking: device

attachment and future directions. *Methods in Ecology and Evolution,* **5,** 311-319.

O'Mara, M.T., Dechmann, D.K. & Page, R.A. (2014) Frugivorous bats evaluate the quality of

social information when choosing novel foods. *Behavioral Ecology,* **25,** 1233-1239.

R Developing Core Team (2015) R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Ratcliffe, J.M. & ter Hofstede, H.M. (2005) Roosts as information centres: social learning of

food preferences in bats. *Biology Letters,* **1,** 72-74.

Ripperger, S., Josic, D., Hierold, M., Koelpin, A., Weigel, R., Hartmann, M., Page, R. &

Mayer, F. (2016) Automated proximity sensing in small vertebrates: design of

miniaturized sensor nodes and first field tests in bats. *Ecology and Evolution,* **6,** 2179-

2189.

Ruczyński, I., Kalko, E.K. & Siemers, B.M. (2007) The sensory basis of roost finding in a

forest bat, Nyctalus noctula. *Journal of Experimental Biology,* **210,** 3607-3615.

Ruczyński, I., Kalko, E.K. & Siemers, B.M. (2009) Calls in the forest: a comparative

approach to how bats find tree cavities. *Ethology,* **115,** 167-177.

Rutz, C., Morrissey, M.B., Burns, Z.T., Burt, J., Otis, B., St Clair, J.J. & James, R. (2015)

Calibrating animal-borne proximity loggers. *Methods in Ecology and Evolution,* **6,**

656–667.

Sachteleben, J. (1991) Zum 'Invasions'—Verhalten der Zwergfledermaus (Pipistrellus

pipistrellus). *Nyctalus (NF),* **4,** 51-66.

Schnitzler, H.-U., Moss, C.F. & Denzinger, A. (2003) From spatial orientation to food

acquisition in echolocating bats. *Trends in Ecology & Evolution,* **18,** 386-394.

Schöner, C.R., Schöner, M.G. & Kerth, G. (2010) Similar is not the same: social calls of

conspecifics are more effective in attracting wild bats to day roosts than those of other

bat species. *Behavioral Ecology and Sociobiology,* **64,** 2053-2063.

Smith, J.E., Lacey, E.A. & Hayes, L.D. (2017) Sociality in Non-Primate Mammals.

Comparative social evolution (eds D.R. Rubenstein & P. Abbot), pp. 284. Camebridge

University Press, Camebridge, UK.

St Clair, J., Burns, Z., Bettaney, E., Morrissey, M., Burt, J., Otis, B., Ryder, B., Fleischer, R.,

James, R. & Rutz, C. (2015) Experimental resource pulses influence social-network

dynamics and the potential for information flow in tool-using crows. *Nature*

Communications, **6,** 7197.

Stumpf, M., Meier, F., Grosche, L., Halczok, T.K., Van Schaik, J. & Kerth, G. (2017) How do

young bats find suitable swarming and hibernation sites? Assessing the plausibility of

the maternal guidance hypothesis using genetic maternity assignment for two

European bat species. *Acta Chiropterologica,* **19,** 319-327.

Templeton, C.N., Greene, E. & Davis, K. (2005) Allometry of alarm calls: black-capped

chickadees encode information about predator size. *Science,* **308,** 1934-1937.

van Schaik, C., Graber, S., Schuppli, C. & Burkart, J. (2017) The Ecology of Social Learning

in Animals and its Link with Intelligence. *The Spanish Journal of Psychology,* **19,**

E99.

Ward, P. & Zahavi, A. (1973) The importance of certain assemblages of birds as

"information-centres" for food-finding. *Ibis,* **115,** 517-534.

Wilkinson, G.S. (1986) Social grooming in the common vampire bat, Desmodus rotundus.

Animal Behaviour, **34,** 1880-1889.

Wilkinson, G.S. (1990) Food sharing in vampire bats. *Scientific American,* **262,** 76-83.

Wilkinson, G.S. (1992) Information transfer at evening bat colonies. *Animal Behaviour,* **44,**

501-518.

Wilkinson, G.S. (1995) Information transfer in bats. *Symposia of the Zoological Society of*

London, pp. 345-360. London: The Society, 1960-1999.

Wilkinson, G.S. & Boughman, J.W. (1998) Social calls coordinate foraging in greater spear-

nosed bats. *Animal Behaviour,* **55,** 337-350.

Wilkinson, G.S. & Boughman, J.W. (1999) Social influences on foraging in bats. *Mammalian*

social learning: Comparative and ecological perspectives (eds H.O. Box & K.R.

Gibson), pp. 188-204. cambridge University Press, Cambridge, UK.

Supporting information

The following Supporting Information is available for this article online.

Table S1: Results from allele frequency calculations with CERVUS v. 3.0 (Kalinovski et al.

2007).

Tables

Table 1: Summary of joint roost switching events of pups and associated partners. Switching durations were only determined when both roosts were known and equipped with base stations, while NA represents unmonitored sites or uncertain switching mode.

pupID	sex	partnerID (mother)	start time [date-time]	end time [date-time]	duration [hh:mm:ss]	Switching mode
9307	f	9338 (m)	2016-07-21 02:59:35	NA	NA	NA
9311	f	9330 (m)	2016-07-16 23:26:22	2016-07-17 01:15:06	01:48:44	3*
9311	f	9330 (m)	2016-07-20 03:18:35	2016-07-20 03:19:31	00:00:56	1
9318	f	9330 (m)	2016-07-16 23:26:33	2016-07-16 23:28:31	00:01:58	1*
9318	f	9330 (m)	2016-07-20 04:14:25	2016-07-20 04:14:54	00:00:29	3
9319	f	9327 (m)	2016-07-16 22:48:01	NA	NA	NA
9312	f	9334 (m)	2016-07-17 01:29:14	NA	NA	1
9323	m	9340 (m)	2016-07-16 23:58:13	2016-07-17 01:37:06	01:38:53	2
9323	m	9340 (m)	2016-07-19 02:05:26	NA	NA	1
9325	m	9336 (m)	2016-07-17 02:44:31	2016-07-17 03:55:19	01:10:48	2
9376	f	9383	2017-07-20 04:23:43	2017-07-20 04:23:49	00:00:06	1
9376	f	9383	2017-07-22 02:25:53	NA	NA	1
9370	f	9368 (m)	2017-07-20 04:35:41	NA	NA	1
9373	m	9412	2017-07-22 02:38:13	2017-07-22 02:38:39	00:00:26	1
9380	m	9413	2017-07-20 04:17:52	NA	NA	1
9391	f	9385 (m)	NA	2017-07-22 02:22:25	NA	1

Switching mode: 1 quick, direct switch among two roosts; 2 switch included a joint stopover at an unknown site indicated by stable meetings; 3 roost switch after commuting flights by mother; * see Fig. 3 for a schematic representation of these events

Figures

Figure 1: Unpackaged proximity sensor (a) and tagged adult Greater noctule bat (*Nyctalus*

noctula) ready for take-off (b).

Figure 2: Visual representation of foraging bouts and roost switches based on presence signals

at bat boxes (base stations) and meeting data. (a) A foraging bout is characterized by an

interrupt of the presence signals of an individual bat which are received by a base station at a

specific roost. Usually, variation of the received signal strength indicator (RSSI) increases

when a bat is leaving a roost compared to when it is roosting (notice the pronounced spike

upon departure and return). (b) A roost switch among two monitored sites is displayed. The

presence signals interrupt at base station two while the strong variation in RSSI indicates that

the bat is flying. Presence signals are then received by base station 1. (c) A roost switch

occurs among an unmonitored to a monitored site. Roosting at the unmonitored site is

indicated by long-lasting stable meetings among three bat individuals. Meetings interrupt

when a bat individual leaves the unmonitored site followed by signal beacons being received

by the base station at the monitored site.

Figure 3: Schematic representation of a mother and its twins switching roosts: repeated

commutes indicate intentional behaviour of the mother. (a) A mother and its twins jointly

leave roost 1 and the mother successfully transfers to roost 2 with pup 1. The meeting to pup 2

aborts and pup 2 is flying back to roost 1. (b) Pup 2 moves solitarily from roost 1 to an

unknown location where it is joined by its mother after a few minutes. Both fly in company

towards roost 2, but pup 2 flies back to roost 1 while a meeting starts among the mother and

pup 1 at roost 2. (c) The mother joins pup 2 in an unknown location and they jointly switch to

roost 1. They jointly leave roost 1, but only the mother arrives at roost 2 starting a meeting

with pup 1, while the pup 2 flies back to roost 1. (d) The mother joins pup 2 in an unknown

location, around 00:50 am the meeting is interrupted for several minutes (possibly because at

least one individual left), before the mother commutes twice between its two pups. Finally,

around 01:15 am the mother successfully switches with pup 2 to roost 2 while a meeting is

ongoing. The triad stays at roost 2 until shortly before 2 am.

(a)

(b)

Legend: ○ Presence signal at base station 1
           △ Presence signal at base station 2

Presence at unknown roost indicated by
stable meetings among three individuals

Roost 1

Unknown location

Roost 2

	Mother (ID 9330)
	Pup 1 (ID 9318)
	Pup 2 (ID 9311)

(a)   (b)   (c)   (d)

23:30   00:00   00:30   01:00

local time Berlin [hh:mm]