Systematic identification and characterization of *Aedes aegypti* long noncoding RNAs

Azali Azlan[1], Sattam M. Obeidat[1], Muhammad Amir Yunus[2] and Ghows Azzam[1]*

[1]*School of Biological Sciences, Universiti Sains Malaysia, 11800 Penang, Malaysia*

[2]*Infectomics Cluster, Advanced Medical & Dental Institute, Universiti Sains Malaysia, Bertam, 13200 Kepala Batas, Pulau Pinang, Malaysia.*

*Corresponding author

Email: ghows@usm.my (G.A)

Abstract

Long noncoding RNAs (lncRNAs) play diverse roles in biological process including developmental regulation and host-pathogen interactions. *Aedes aegypti* (*Ae. aegypti*), a blood-sucking mosquito, is the principal vector responsible for replication and transmission of arboviruses including dengue, zika, and chikungunya virus. Systematic identification and developmental characterisation of *Ae. aegypti* lncRNAs are still limited. We performed genome-wide identification of lncRNAs followed by developmental profiling of lncRNA expression in *Ae. aegypti*. We identified a total of 4,689 novel lncRNA transcripts, of which 2,064, 2,076, and 549 were intergenic, intronic, and antisense respectively. *Ae. aegypti* lncRNAs shared many of the characteristics with other species including low expression, low GC content, short in length, low conservation, and their expression tended to be correlated with neighbouring and antisense protein-coding genes. Subsets of lncRNAs showed evidence that they were maternally inherited, suggesting potential roles in early-stage embryos. Additionally, lncRNAs showed higher tendency to be expressed in developmental and temporal specific manner. Upon infection of *Ae. aegypti* cells with dengue virus serotype 1, we identified 2,335 differentially expressed transcripts, 957 of which were lncRNA transcripts. The systematic annotation, developmental profiling, and transcriptional response upon virus infection provide foundation for future investigation on the function of *Ae. aegypti* lncRNAs.

Introduction

Aedes aegypti (*Ae. aegypti*), a blood-sucking mosquito, is the principal vector responsible for

replication and transmission of arboviruses such as dengue (DENV), Chikugunya (CHIKV), and Zika

(ZIKV) virus. Although virus infections of the vertebrate host are acute and often associated with

disease, the maintenance of these pathogens in nature generally requires the establishment of a

persistent non-lethal infection in the insect host. This is also true in the context of dengue virus

infection. While dengue-infected humans show serious symptoms of illnesses, interestingly, infected

Ae. aegypti mosquitoes are generally asymptomatic, suggesting that these vectors possess intrinsic

antiviral immunity to resist or tolerate the virus infection. A number of studies have shown that, upon

flavivirus infection, major transcriptomic changes occurred within *Ae. aegypti.* Such changes involved

a wide range of host genes including genes related to immunity, metabolic pathways, and trafficking.

Although protein-coding genes have been the central focus, previous report indicated that *Ae. aegypti*

lncRNAs were also involved in virus-host interaction[1,2].

Advances in high-throughput sequencing technologies coupled with bioinformatics analyses

have enabled the identification of a significant number of lncRNAs in various species. Such lncRNAs

exert their functions by a range of mechanisms. For instance, several lncRNAs regulate gene

expression by sequestering miRNAs and transcription factors [3,4]. Other lncRNAs have been shown to

regulate the activity of chromatin modifying proteins; hence, modulating gene expression. In *Ae.*

aegypti, it had been reported that lncRNA was involved in virus-host interaction. Knockdown of

lincRNA_1317 in *Ae. aegypti* cells caused higher replication of DENV[1]. Although lncRNAs have been

systematically identified in many organisms, most lncRNAs have not been functionally characterised.

Recently, the latest version of *Ae. aegypti* genome (AaegL5) was released, and the assembly

was up to chromosome-length scaffolds, which is more contiguous than the previous AaegL3 and

AaegL4 assemblies[5]. This prompted us to perform novel lncRNA identification and characterisation

using the latest genome release as reference. Previous study[1] reported that a total of 3,482 intergenic

lncRNA (lincRNA) was identified in *Ae. aegypti*. However, the identification was performed using

previous version of *Ae. aegypti* genome (AaegL3), and only lncRNAs residing in intergenic region

were annotated.

Here, we report genome-wide identification and characterisation of lncRNAs in *Ae. aegypti*.

Identification of lncRNAs was performed using a combination of RNA-seq based *de novo* transcript

discovery and rigorous filtering of putative protein-coding transcripts. In this study, we defined a high-

confidence set of 4,689 novel lncRNA transcripts -2,064 2,076, and 549 were intergenic, intronic, and

antisense respectively. We then characterised the newly identified lncRNAs by diverse features such as

transcript structures, conservation, and developmental expression. Previous studies have reported

global transcriptional responses of lncRNAs upon infection with dengue virus serotype 2 (DENV2) and

ZIKV. Here, we examined the transcriptome profiles in *Ae. aegypti* cell (Aag2) in response to DENV

serotype 1 (DENV1) infection. Collectively, genome-wide annotation and characterisation of *Ae.*

aegypti lncRNAs provide valuable resources for future genetics and molecular studies in this harmful

mosquito vector.

Identification of lncRNA

To perform lncRNA prediction, we used a total of 117 datasets of Illumina high-throughput

sequencing derived from *Ae. aegypti* mosquito and Aag2 cell, a widely used *Ae. aegypti* derived cell

line[6]. An overview of lncRNA identification pipeline can be found in Fig 1. The pipeline developed in

this study was adapted with few modifications from previous reports [1,7,8]. Briefly, each dataset (both

paired-end and single-end) was individually mapped using HISAT2[9]. The resulting alignment files

were used for transcriptome assembly, and the assemblies were merged into a single unified

transcriptome. Both transcriptome assembly and merging were performed using Stringtie[10]. Then, we

used Gffcompare to annotate and compare the unified transcriptome assembly with reference

annotation (AaegL5.1, VectorBase). We classified lncRNA transcripts based on their position relative to

annotated genes derived from AaegL5.1 assembly (VectorBase). We only selected transcripts with class

code "i", "u", and "x" that denote intronic, intergenic, and antisense to reference genes for downstream

analysis.

To get confident lncRNA transcripts, we performed multiple steps of filtering transcripts having

coding potential or open-reading frame (Fig 1). The steps involved TransDecoder[11], CPAT[12] and finally

BLASTX. We then removed lncRNA candidates that did not have strand information. Detailed

description on the prediction analysis and parameter used can be found in Material and Methods

section. Using this pipeline, we identified a set of 4,689 novel lncRNA transcript isoforms derived from

3,621 loci. Of these 4,689 lncRNA transcripts, 2,064 and 2,076 were intergenic and intronic

respectively, while the remaining 549 transcripts were antisense to reference genes. Currently,

AaegL5.1 annotation catalogs 4,155 lncRNA transcripts. Here, we provided another set of 4,689

lncRNA transcripts, making a total of 8,844 lncRNAs in *Ae. aegypti*.

Characterisation of *Ae. aegypti* lncRNA

To examine whether lncRNAs identified in this study exhibit typical characteristics observed in

other species[8,13–15], we analysed features such as coding potential, sequence length, GC content and

sequence conservation with closely related species. Since lncRNAs are strictly defined by their

inability to code for protein, we determined coding probability of our newly identified lncRNAs and

compared them with known lncRNA, 3'UTR, 5'UTR, and protein-coding mRNA. We discovered that,

similar to other non-coding sequence such as known lncRNA, 3'UTR, and 5'UTR, our novel lncRNA

transcripts have extremely low coding probability when compared to protein-coding mRNA (Fig 2a).

Beside that, we found that both novel and known lncRNAs (provided by AaegL5.1 annotation) were

shorter than protein-coding transcripts (Fig 2b). Mean length of novel and known lncRNAs was 825.4

bp and 745.6 bp respectively, while protein-coding mRNA has an average length of 3330 bp.

Similar to previous reports[1,8], we observed that lncRNAs identified in this study had slightly

lower GC content compared to protein-coding mRNAs (Fig 2c). For instances, mean GC content of

novel lncRNA and mRNA was 40.1% and 46.4% respectively. Known lncRNA, on the other hand, had

relatively similar mean GC content to novel lncRNA (40.8%), while average GC of 5'UTR and 3'UTR

sequence was 43.1% and 34.6% respectively. Overall, GC content of non-coding sequence was

relatively lower than coding sequence.

To determine the conservation level of *Ae. aegypti* lncRNAs, we performed BLASTN against

other insect genomes including *Ae. albopictus, C, quinquifasciatus*, *An, gambiae* and *D. melanogaster,*

Bit score derived from BLAST algorithm was used to determine the level of sequence similarity of *Ae.*

aegypti lncRNAs with previously mentioned insect genomes[1]. Similarly, we also performed BLASTN

of *Ae. aegypti* protein-coding mRNA for comparison. We discovered that both lncRNAs and protein-

coding mRNAs displayed high degree of sequence similarity with *Ae. albopictus,* suggesting that they

were presumably genus specific (Supplementary Fig. 1). In general, compared to protein-coding

mRNA, *Ae. aegypti* lncRNAs exhibited lower sequence conservation.

It was reported that, in contrast to coding gene, lncRNAs harbour higher composition of repeat

elements[16,7]. To test if similar occurrence held true in *Ae. aegypti*, we determined the fraction of repeat

element in the exons of lncRNAs and protein-coding mRNAs. As expected, we discovered that more

than 20% of nt of novel lncRNAs were made up of repeat elements (Fig 2d). Meanwhile, 6% and 1%

of known lncRNAs and protein-coding mRNAs were composed with repeats. Taken together, *Ae.*

aegypti lncRNAs shared many characteristics with other species: relatively short in length, relatively

lower GC content, and higher repeat content.

Developmental expression of *Ae. aegypti* lncRNAs

To examine the developmental expression of *Ae. aegypti* lncRNAs, we analysed public dataset

(SRP026319) which provided RNA-seq data of specific developmental stages, ranging from early

embryo up to adult mosquitoes. These developmental stages include specific time interval of

embryonic development, larval stages, sex-biased expression of male versus female pupae and carcass,

blood-fed versus nonblood-fed ovary and female carcass, and testes-specific expression. Similar to

protein-coding genes, *Ae. aegypti* lncRNA genes exhibited stage-specific (embryo, larva, and pupal

stages), sex-specific, and blood-fed versus nonblood-fed (ovary and female carcass) expression pattern

(Fig 3).In addition, each time point in the development had distinct lncRNA expression pattern. For

instance, there was a subset of lncRNAs that were highly enriched during early embryonic

development (0-8 hour embryo) as compared to later time points (8-76 hour embryo). A total of 1,848

lncRNA genes consisting of 24.7% of the total expressed lncRNAs were specifically highly expressed

at this early embryonic stage. Interestingly, these early embryo-specific lncRNAs were also highly

expressed in blood-fed ovary, suggesting that these lncRNAs may have potentially maternally inherited

and possess essential roles in early development.

Previous report showed that lncRNAs displayed a more temporally specific expression pattern

than protein-coding genes[17]. To test whether or not this is true for *Ae. aegypti* lncRNAs, we computed

specificity score of each lncRNA using Jensen-Shannon (JS) score[18,17]. The JS score ranged from 0 to

1, with 1 indicated perfect specificity. Here, we computed JS score across two stages namely embryo

and larvae, and two conditions which were blood-fed ovary and female carcass, all of which were

sampled in a timely fashion. In all four time-course samples namely embryo, larvae, blood-fed female

carcass and ovary, we discovered that novel and known lncRNAs had higher JS specificity score in all

four samples (Fig. 4). Meanwhile, fraction of protein-coding genes having JS score of 1in all four

samples were lower than lncRNAs (Fig. 4). Although lncRNAs had higher developmental and temporal

specificity, across all four time-course samples being analysed, the overall expression of lncRNAs was

lower than that of protein-coding genes (Supplementary Fig. 2).

Previous studies have shown that lncRNAs are favourably resided in close proximity to genes

with developmental functions[19,20]. In addition, this close physical proximity of lncRNAs and protein-

coding genes resulted in their expression to be strongly correlated. We asked whether *Ae. aegypti*

lncRNAs showed correlation in expression between neighboring protein-coding genes. We also

examined the correlation in expression of lncRNAs that were antisense to protein-coding genes. By

analysing the expression of 42 developmental samples (Fig. 3), we discovered that fraction of

positively correlated (Pearson correlation, p-value < 0.05) antisense lncRNA-coding gene pair was

higher than that of randomly assigned antisense lncRNA-coding gene pair (Fig 5). Analysis of

neighboring genes (within 10 kb) revealed that the expression of lncRNAs and their nearest

neighboring genes showed a slightly higher degree of correlation compared to random gene pairs. In

both cases of random neighbouring pair and random antisense lncRNA-coding gene pair, the majority

of the pairs had correlation near to zero value (Fig 5).

***Ae. aegypti* lncRNAs were differentially expressed following DENV1 infection**

Previous studies reported that lncRNAs in *Ae. aegypti* underwent differential expression upon

dengue virus serotype 2 (DENV2) and ZIKV infection. In this study, we chose another DENV

serotype, namely serotype 1, and observed the transcriptional response of *Ae. aegypti* lncRNAs upon

infection. Therefore, we generated paired-end RNA-seq libraries derived from triplicates of Aag2 cells

infected with DENV1. To avoid bias in library size which affects fold change calculation, we

performed differential expression analysis of protein-coding genes and lncRNAs together in a single

transcriptome reference using Salmon v0.9[21] followed by edgeR[22]. Here, differential expression

analysis was performed at transcript level. A total of 2,335 transcripts were differentially expressed (P-

value < 0.05) upon DENV1 infection, 957 of which were lncRNA transcripts.

In general, we found that across all six samples (3 uninfected and 3 DENV1-infected samples),

expression of lncRNA was lower than protein-coding mRNAs (Fig 6a). Besides, distribution of fold

change (FC) of differentially expressed lncRNAs (P-value < 0.05) were much lower than that of

mRNAs (Fig 6b). This observation suggests that unlike mRNA, lncRNA expression was less

responsive to viral infection. For example, log2 FC of mRNA can extend to more than 10 fold in either

direction (-10 and 10 fold), but in lncRNA, log2 FC ranged from -2 to 5. Besides, most of the

differentially expressed lncRNAs (953 out of 957) transcripts had log2 FC of less than 2 in either

direction.

Discussion

The field of lncRNA has become increasingly important in many areas of biology particularly

infectious disease, immunity, and pathogenesis[1,8,14,23]. High-throughput sequencing combined with

bioinformatics enable scientists to uncover comprehensive repertoire of lncRNA in many species. Here,

we present a comprehensive lncRNA annotation using the latest genome reference of *Ae. aegypti*

(AaegL5). Due to the recent release of *Ae. aegypti* genome (AaegL5) equipped with improved gene set

annotation[5], we decided to perform lncRNA identification using this latest genome reference. Unlike

previous annotation that mainly focused on *Ae. aegypti* intergenic lncRNA, here, we also annotated

lncRNAs residing within the introns and antisense to reference genes. We used large RNA-seq datasets

for lncRNA annotation; hence, enabling identification of *Ae. aegypti* lncRNAs that are lowly expressed.

In addition, we used recently developed and improved softwares including HISAT[9], StringTie[10],

and Salmon[21] for novel lncRNA prediction and expression analysis. HISAT is considered to be faster

with equal or better accuracy than other spliced aligner methods such as Tophat[24] and STAR[25].

Meanwhile, compared to other transcript assembly software such as Cufflinks[26] and Scripture[27],

StringTie has been shown to produce more comprehensive and accurate transcriptome reconstruction

from RNA-seq data[10]. Salmon was used for transcript abundance quantification in this study due to its

rapidness and accuracy since the algorithm is able to correct for fragment GC content bias[21]. Taken

together, the use of large RNA-seq datasets, the most recent genome reference, and improved

algorithms in bioinformatics analyses enable us to provide, to date, the most comprehensive and

confident lncRNA annotation in *Ae. aegypti* transcriptome.

Similar to previous reports[1,8,17,23,28], we discovered that lncRNAs identified in our study

exhibited typical characteristics of lncRNAs found in other species including vertebrates[7]. Such

characteristics are lower GC content, shorter in length, and low sequence conservation even among

closely related species. Analysis of *Ae. aegypti* developmental expression revealed that the expression

of lncRNAs is highly temporally specific relative to that of coding genes. In other words, lncRNAs

have a much narrower time window in expression than coding genes. Such specificity in expression

suggests essential roles of lncRNAs in *Ae. aegypti*, especially in regulating the timing of developmental

transition.

Several studies have pointed out that lncRNAs are essential in development and cell

differentiation[29]. For example, lncRNAs emerge as important regulators of embryogenesis, by means of

directing pluripotency and differentiation[30] In this study, we provided an evidence of maternal

inheritance of *Ae. aegypti* lncRNAs. This further corroborated previous findings that highlighted the

importance of lncRNAs in embryogenesis[20,17,31] We observed a fraction of lncRNAs that was highly

expressed in blood-fed ovary, and the expression persisted up to 8-12 hour embryonic stage. This

narrow time window in early embryo was related to maternal-zygotic transition stage[32]. Since

transcription from the zygotic genome has not been activated during early embryonic stage, these

highly expressed lncRNAs must be maternally provided; suggesting that they might play roles in basic

biosynthesis processes in the early embryo, specification of initial cell fate and pattern formation.

Meanwhile, lncRNAs expressed at later embryonic, larval and pupal stages are putatively responsible

in organogenesis.

Upon DENV1 infection, we observed that both protein-coding and lncRNAs experienced

overall changes in expression level. However, the overall level of fold change displayed by lncRNAs

was not as high as protein-coding genes. This observation however, did not necessarily negate the

possibility of lncRNAs having potential roles in immunity. For example, RNAi-mediated knockdown

of one lncRNA candidate (lincRNA_1317) in *Ae. aegypti* resulted in higher viral replication[1].

Therefore, intensive investigation using loss or gain of function approach on lncRNAs is crucial in

dissecting their functions in mosquito-virus interaction.

In summary, we provided a comprehensive list of lncRNAs in *Ae. aegypti* using the latest

genome version. This study highlights numerous roles of lncRNAs in *Ae. aegypti* especially in

development and virus-host interaction.

Materials and Methods

Cell culture and virus

Ae. aegypti Aag2 cell and *Ae. albopictus* C6/36 cell were cultured in Leibovitz's L-15 medium (Gibco,

41300039), supplemented with 10% Fetal Bovine Serum (FBS, Gibco, 10270) and 10% Tryptose

Phosphate Broth (TPB) solution (Sigma, T9157). Both Aag2 and C6/36 cells were incubated at 25ºC

without CO_2. Vero E6 cells were cultured at 37ºC in Dulbecco's modified Eagles Medium (DMEM,

Gibco, 11995065) supplemented with 10% FBS (Gibco, 10270), and 5% CO_2. DENV1, Hawaiian

strain was propagated in C6/36 cells and titered using Vero E6 cells. Determination of DENV1 titer

was done using 50% tissue culture infectious dose – cytopathic effect (TCID50-CPE) as previously

described[33]. DENV1 used in this study was a gift from Dr. David Perera, University Malaysia Sarawak.

Aag2 cell line was a kind gift from Dr. Ronald P. Van Rij from Radbound Institute for Molecular Life

Sciences, Netherlands.

Virus infection, RNA isolation and sequencing

Aag2 cells were infected with DENV1 at multiplicity of infection (MOI) of 0.5. At 72 hours post

infection (hpi), the culture medium was removed, and RNA isolation was carried out using miRNeasy

Mini Kit 50 (Qiagen, 217004) according to the manufacturer's protocol. Total RNA was then subjected

to next-generation sequencing. The RNA-sequencing libraries were prepared using standard Illumina

protocols and sequenced using HiSeq2500 platform generating paired-end reads of 150 in size.

Verification of DENV1 infection

Total RNA of both uninfected and DENV1-infected samples were subjectred to cDNA synthesis using

Tetro cDNA synthesis kit (Bioline, BIO-65042). Reverse primer for DENV1 was included in the

reaction set-up. This followed with PCR and gel electrophoresis using forward and reverse primers of

DENV1. Primers used for DENV1 verification were taken from previous study[34].

RNA-seq data preparation

RNA-seq datasets generated in this study (Aag2 cell infected with DENV1) are deposited in the Short

Read Archive database under the accession code: PRJNA509977. Publicly available datasets were

downloaded from NCBI Sequence Reads Archive (SRA) and ArrayExpress with accession numbers

SRP041845, SRP047470, SRP046160, SRP115939, E-MTAB-1635, SRP035216, SRP065731,

SRP065119, SRA048559, PRJEB13078, SRP026319[2,35–40]. Adapters were removed using

Trimmomatic version 0.38[41], and reads with average quality score (Phred Score) above 20 were

retained for downstream analysis.

lncRNA identification

Each library (both paired-end and single-end) was individually mapped against *Ae. aegypti* genome

(AaegL5) using HISAT2 version 2.1.0[9]. The resulting alignment files were used as input for transcript

assembly using Stringtie version 1.3.2[10]. We used reference annotation file of AaegL5 (VectorBase) to

guide the assembly only without discarding the assembly of novel transcripts. Here, we set the

minimum assembled transcript length to be 200 bp. Then, the output gtf files were merged into a single

unified transcriptome using Stringtie merge[10]. Only input transcripts of more than 1 FPKM and TPM

were included in the merging. Then, we compared the assembled unified transcript to a reference

annotation of AaegL5 (VectorBase) using Gffcompare (https://github.com/gpertea/gffcompare). For the

purpose of lncRNA prediction, we only retained transcripts with class code "i", "u", and "x". The

transcripts were then subjected to coding potential prediction. We used TransDecoder[11] to identify

transcripts having open-reading frame (ORF), and those having ORF were discarded. The remaining

transcripts were then subjected to a coding potential assessment toll (CPAT)[12]. We set the same cut-off

as previous study in *Ae. aegypti* which is less than 0.3[1]. Transcripts having coding potential more than

0.3 were discarded. To exclude false positive prediction, we used BLASTX against Swissprot database,

and transcripts having E-value of less than 10^{-5} were removed.

Identification of differentially expressed transcripts

307 We used Salmon version 0.10.1[21] to quantify the expression of transcripts. Read count for each

308 transcript was subjected for differential analysis using edgeR[22] in R/Bioconductor environment.

Quantitative real-time PCR (qRT-PCR)

cDNA synthesis was done using QuantiNova Reverse Transcription Kit (Qiagen) followed by qRT-PCR with QuantiNova SYBR Green PCR kit (Qiagen) according to manufacturer's protocol. *RPS17* was used as housekeeping gene[42], and the experiment was performed using Applied Biosystems Step One PlusTM Real-Time PCR System.

Figure Legend

Fig 1 lncRNA identification

Overview of lncRNA identification pipeline.

Fig 2 Characterization of *Ae aegypti* lncRNA

(a) Coding probability of lncRNA, 3'UTR, 5'UTR and mRNA. (b) Sequence length distribution of lncRNA and mRNA. (c) GC content. (d) Repeat content

Fig 3 Hierarchical clustering of protein-coding and lncRNA expression

Hierarchical clustering of protein-coding (a) and lncRNA (b) across developmental stages. Hierarchical clustering analysis was done in Morpheus (https://software.broadinstitute.org/morpheus) based on Pearson correlation of z scores of lncRNA and protein-coding genes. Boxes in (b) indicate a subset of lncRNAs that are presumably maternally inherited.

Fig 4 Maximal JS specificity score

Fig 5 Correlation of expression analysis

(a) Pearson correlation of antisense lncRNA expression with its corresponding genes compared to that

of antisense lncRNA pair with random protein-coding gene (b) Pearson correlation of gene expression

within 10 kb from each other

Fig 6 Overall distribution of expression and fold change of lncRNA and mRNA upon DENV1

infection

List of Supporting Information

S1 Table. Genomic coordinates of lncRNA candidates identified in *Ae. aegypti*

S2 Table. List of differentially expressed lncRNA

S3 Table. List of primers used in this study

S1 Fig. Similarity bit score of lncRNA and mRNA with closely related insect genomes

S2 Fig. Distribution of lncRNA and coding gene expression in embryo, larvae, blood-fed ovary

and female carcass

S3 Fig. Validation of differentially expressed lncRNAs by qRT-PCR

Two-tailed *t*-test was used for all comparisons; **$p < 0.05$, ***$p < 0.01$. Expression levels were

normalised to uninfected samples. Error bars indicate standard error of the mean (SEM) of three

independent experiments.

References

1. Etebari, K., Asad, S., Zhang, G. & Asgari, S. Identification of Aedes aegypti Long Intergenic Non-coding RNAs and Their Association with Wolbachia and Dengue Virus Infection. *PLoS Negl. Trop. Dis.* **10**, e0005069 (2016).

2. Etebari, K. *et al.* Global Transcriptome Analysis of *Aedes aegypti* Mosquitoes in Response to Zika Virus Infection. *mSphere* **2**, e00456-17 (2017).

3. Poliseno, L. *et al.* A coding-independent function of gene and pseudogene mRNAs regulates tumour biology. *Nature* **465**, 1033–1038 (2010).

4. Hung, T. *et al.* Extensive and coordinated transcription of noncoding RNAs within cell-cycle promoters. *Nat. Genet.* **43**, 621–629 (2011).

5. Matthews, B. J. *et al.* Improved reference genome of Aedes aegypti informs arbovirus vector control. *Nature* **563**, 501 (2018).

6. Walker, T., Jeffries, C. L., Mansfield, K. L. & Johnson, N. Mosquito cell lines: history, isolation, availability and application to assess the threat of arboviral transmission in the United Kingdom. *Parasit. Vectors* **7**, 382 (2014).

7. Hezroni, H. *et al.* Principles of Long Noncoding RNA Evolution Derived from Direct Comparison of Transcriptomes in 17 Species. *Cell Rep.* **11**, 1110–1122 (2015).

8. Wu, Y. *et al.* Systematic Identification and Characterization of Long Non-Coding RNAs in the Silkworm, Bombyx mori. *PLOS ONE* **11**, e0147147 (2016).

9. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory requirements. *Nat. Methods* **12**, 357–360 (2015).

10. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat. Biotechnol.* **33**, 290–295 (2015).

11. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-Seq: reference generation and analysis with Trinity. *Nat. Protoc.* **8**, (2013).

12. Wang, L. *et al.* CPAT: Coding-Potential Assessment Tool using an alignment-free logistic regression model. *Nucleic Acids Res.* **41**, e74 (2013).

13. Chodroff, R. A. *et al.* Long noncoding RNA genes: conservation of sequence and brain expression among diverse amniotes. *Genome Biol.* **11**, R72 (2010).

14. Clark, M. B. & Mattick, J. S. Long noncoding RNAs in cell biology. *Semin. Cell Dev. Biol.* **22**, 366–376 (2011).

15. Jenkins, A. M., Waterhouse, R. M. & Muskavitch, M. A. Long non-coding RNA discovery across the genus anopheles reveals conserved secondary structures within and beyond the Gambiae complex. *BMC Genomics* **16**, 337 (2015).

16. Nam, J.-W. & Bartel, D. P. Long noncoding RNAs in C. elegans. *Genome Res.* **22**, 2529–2540 (2012).

17. Chen, B. *et al.* Genome-wide identification and developmental expression profiling of long noncoding RNAs during *Drosophila* metamorphosis. *Sci. Rep.* **6**, 23330 (2016).

18. Cabili, M. N. *et al.* Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes Dev.* **25**, 1915–1927 (2011).

19. Ponjavic, J., Oliver, P. L., Lunter, G. & Ponting, C. P. Genomic and Transcriptional Co-Localization of Protein-Coding and Long Non-Coding RNA Pairs in the Developing Brain. *PLOS Genet.* **5**, e1000617 (2009).

20. Pauli, A. *et al.* Systematic identification of long noncoding RNAs expressed during zebrafish embryogenesis. *Genome Res.* **22**, 577–591 (2012).

21. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon: fast and bias-aware quantification of transcript expression using dual-phase inference. *Nat. Methods* **14**, 417–419 (2017).

22. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

23. Young, R. S. *et al.* Identification and Properties of 1,119 Candidate LincRNA Loci in the Drosophila melanogaster Genome. *Genome Biol. Evol.* **4**, 427–442 (2012).

24. Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* **25**, 1105–1111 (2009).

25. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

26. Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* **28**, 511–515 (2010).

27. Guttman, M. *et al. Ab initio* reconstruction of cell type–specific transcriptomes in mouse reveals the conserved multi-exonic structure of lincRNAs. *Nat. Biotechnol.* **28**, 503–510 (2010).

28. Tang, Z. *et al.* Comprehensive analysis of long non-coding RNAs highlights their spatio-temporal expression patterns and evolutionary conservation in *Sus scrofa*. *Sci. Rep.* **7**, 43166 (2017).

29. Fatica, A. & Bozzoni, I. Long non-coding RNAs: new players in cell differentiation and development. *Nat. Rev. Genet.* **15**, 7–21 (2014).

30. Pauli, A., Rinn, J. L. & Schier, A. F. Non-coding RNAs as regulators of embryogenesis. *Nat. Rev. Genet.* **12**, 136–149 (2011).

31. Karlic, R. *et al.* Long non-coding RNA exchange during the oocyte-to-embryo transition in mice. *DNA Res. Int. J. Rapid Publ. Rep. Genes Genomes* **24**, 129–141 (2017).

32. Akbari, O. S. *et al.* The Developmental Transcriptome of the Mosquito *Aedes aegypti* , an Invasive Species and Major Arbovirus Vector. *G3amp58 GenesGenomesGenetics* **3**, 1493–1509 (2013).

33. Li, J. *et al.* Enzyme-Linked Immunosorbent Assay-Format Tissue Culture Infectious Dose-50 Test for Titrating Dengue Virus. *PLOS ONE* **6**, e22553 (2011).

34. Johnson, B. W., Russell, B. J. & Lanciotti, R. S. Serotype-Specific Detection of Dengue Viruses in a Fourplex Real-Time Reverse Transcriptase PCR Assay. *J. Clin. Microbiol.* **43**, 4977–4983 (2005).

35. Li, Y. *et al.* RNA-Seq Comparison of Larval and Adult Malpighian Tubules of the Yellow Fever Mosquito Aedes aegypti Reveals Life Stage-Specific Changes in Renal Function. *Front. Physiol.* **8**, 283 (2017).

36. Hall, A. B. *et al.* SEX DETERMINATION. A male-determining factor in the mosquito Aedes aegypti. *Science* **348**, 1268–1270 (2015).

37. McBride, C. S. *et al.* Evolution of mosquito preference for humans linked to an odorant receptor. *Nature* **515**, 222–227 (2014).

38. Canton, P. E., Cancino-Rodezno, A., Gill, S. S., Soberón, M. & Bravo, A. Transcriptional cellular responses in midgut tissue of Aedes aegypti larvae following intoxication with Cry11Aa toxin from Bacillus thuringiensis. *BMC Genomics* **16**, 1042 (2015).

39. David, J.-P. *et al.* Comparative analysis of response to selection with three insecticides in the dengue mosquito Aedes aegyptiusing mRNA sequencing. *BMC Genomics* **15**, 174 (2014).

40. Maringer, K. *et al.* Proteomics informed by transcriptomics for characterising active transposable elements and genome annotation in Aedes aegypti. *BMC Genomics* **18**, (2017).

41. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinforma. Oxf. Engl.* **30**, 2114–2120 (2014).

42. Dzaki, N., Ramli, K. N., Azlan, A., Ishak, I. H. & Azzam, G. Evaluation of reference genes at different developmental stages for quantitative real-time PCR in Aedes aegypti. *Sci. Rep.* **7**, 43618 (2017).