Title: Rare genetic variation and balanced polymorphisms are important for survival in global

change conditions

Authors: Reid S. Brennan[1,2], April D. Garrett[1], Kaitlin E. Huber, Heidi Hargarten, Melissa H.

Pespeni[2]

Affiliation: Department of Biology, University of Vermont, Burlington, VT, USA

[1]Authors contributed equally to this work

[2]Corresponding authors

Email: Reid.Brennan@uvm.edu; mpespeni@uvm.edu

Running head: Low pH adaptation

Keywords: standing genetic variation; ocean acidification; adaptation; experimental selection;

select and resequence; extreme events; capture sequencing

Abstract

Standing genetic variation is important for population persistence in extreme

environmental conditions. While some species may have the capacity to adapt to predicted

average future global change conditions, the ability to survive extreme events is largely

unknown. We used single generation selection experiments on hundreds of thousands of

Strongylocentrotus purpuratus sea urchin larvae generated from wild-caught adults to identify

adaptive genetic variation responsive to moderate (pH 8.0) and extreme (pH 7.5) low pH

conditions. Sequencing genomic DNA from pools of larvae, we identified consistent changes in

allele frequencies across replicate cultures of both conditions and observed increased linkage

disequilibrium around selected loci, revealing selection on recombined standing genetic

variation. We found that loci responding uniquely to either selection regime were at low starting

allele frequencies while variants that responded to both pH conditions (11.6% of selected

variants) started at high frequencies. Loci under selection performed functions related to

energetics, pH tolerance, cell growth, and actin/cytoskeleton dynamics. These results highlight

that persistence in future conditions will require two classes of genetic variation: common, pH-

responsive variants maintained by balancing selection in a heterogeneous environment, and rare

variants, particularly for extreme conditions, that must be maintained by large population sizes.

Introduction

Introduction

As temperatures increase and oceans become more acidic, many marine species are at

risk of decline and extinction [1]. In addition to changing global averages, the frequency and

intensity of extreme events are increasing [2]. While organisms may acclimate through

physiological plasticity or migrate to suitable habitats, some amount of genetic adaptation will be

necessary for population persistence, particularly in the context of extreme events [3]. Indeed,

extreme events rather than average conditions play an important role in setting physiological and

biogeographic limits of species [4-6]; however, the genetic mechanisms that allow rapid

adaptation to extreme conditions have rarely been explored [7].

Evolve or select and resequence, a type of experimental evolution, is a powerful and

promising approach for understanding the capacity for adaptation to future conditions and

extreme events. Selection can be artificially induced to directly measure adaptive genetic

responses [8-11]. While these studies typically leverage 15-20 generations of selection [10,12],

empirical studies have found that adaptive genetic variation can be identified with a single

generation of selection by maximizing replication and the number of outbred individuals or

offspring used [13-15], making this approach applicable to a broad diversity of systems.

The purple sea urchin, *Strongylocentrotus purpuratus*, is an ideal model for

understanding the process of adaptation from standing genetic variation. This species is found in

inter- and subtidal rocky reefs and kelp forests from Alaska to Baja California, Mexico in the

California Current Marine Ecosystem (CCME). They experience high heterogeneity in

temperature and pH conditions in both time and space [16-19], increasing the likelihood of the

presence of adaptive standing genetic variation [20-22]. Large census and effective population

sizes contribute to high standing genetic variation [23,24] and there is evidence of adaptation to

local temperature and pH conditions [25]. Previous studies have shown that *S. purpuratus* larvae

have the physiological and genetic capacity to adaptively respond to an acidifying ocean

[15,18,26], but these studies have focused on relatively mild low pH conditions and have not

investigated the capacity to respond to extreme low pH conditions that are predicted to increase

in frequency in the near future [2,17].

Here, we performed single generation selection experiments in moderate (pH 8.0) and

extreme low (pH 7.5) pH conditions using purple sea urchin larvae generated from 25 wild-

caught adults. We used pooled sequencing of genomic DNA to test the hypotheses that (1) larvae

before and after treatments will show consistent changes in allele frequency across replicate

cultures due to selection for genetic variation that impacts survival in low pH conditions; (2)

there will be unique genetic variation responsive to extreme versus moderate low pH conditions;

(3) genomic patterns of variation, such as linkage disequilibrium (LD) and starting allele

frequencies, will provide insight into the evolutionary mechanisms that maintain adaptive

standing genetic variation for survival in future global change conditions.

Methods

Sample collection & experiment:

Purple sea urchin adults (San Diego, CA, USA; n=25 total: 14 females and 11 males)

were induced to spawn using 0.5M KCl in Instant Ocean artificial seawater (ASW) (Instant

Ocean, Blacksburg, VA) at 14°C and salinity of 35ppt. For each female, 200,000 eggs were

placed in each treatment (moderate (pH 8.0) and extreme (pH 7.5) low pH) and fertilized by

evenly pooled sperm from all males. We chose these pH conditions based on empirical data; the

current average open ocean and intertidal pH is 8.1, though conditions in the CCME frequently

drop to pH 8.0 (daily), and rarely drop as low as pH 7.5 (once in a three month upwelling period;

[16,17,19,27]). Fertilized eggs were pooled across all females by pH and seeded into four

replicate culturing vessels per treatment (37,000 eggs per 3.7 L vessel). Developing embryos

were sampled at day 1 and day 7 post fertilization for morphometric, mortality, and genomic

analyses. See supplemental methods for expanded details and Table S1 for water chemistry.

Morphometrics:

Seven-day old 4-armed pluteus larvae were photographed for morphometric analysis

using a Photometrics Scientific CoolSNAP EZ camera (Tuscon, AZ) connected to a Zeiss

Axioscop 2 compound microscope (Jena, Germany). Larval body size data were analyzed in R

[28] with a generalized linear mixed model in the package lme4 [29], with pH as a fixed effect

and culturing vessel as a random effect.

DNA sequencing, Mapping, & SNP-calling:

DNA was extracted from pools of approximately 15,000 to 20,000 surviving larvae from

each replicate vessel using a Zymo ZR-Duet DNA/RNA MiniPrep Plus Kit (Zymo, Irvine, CA).

DNA was shipped to Rapid Genomics (Gainesville, FL) for library prep, capture, and

sequencing. DNA was captured with 46,316 custom probes designed to capture two 120 base

pair regions per gene: one within exon boundaries and one in putative regulatory regions, within

1000 bp upstream of the transcription start site. Barcoded samples were pooled and sequenced

using 150 bp paired-end reads on one lane of an Illumina HiSeqX.

Raw paired-end reads were quality trimmed and mapped to the *S. purpuratus* genome 3.1

(build 7) with bwa mem [30]. Variants were identified using *mpileup* in Samtools and filtered for

minor allele frequency (maf) of 0.01, quality greater than 20, bi-allelic SNPs only, and no

missing data. We used only variants where each pool was sequenced to a depth of $> 40x$ and

with an average minimum depth across all pools $> 50x$, as recommended for pooled sequencing

by Schlotterer et al. [31]. Mean maximum coverage cutoff was 372. Finally, we removed off

target variants, any variant greater than 2kb from a probe. This filtering resulted in 75,368

variant sites. See supplemental methods for more details. Code for data processing and analyses

can be found on GitHub: https://github.com/PespeniLab/urchin_single_gen_selection.

Detecting changes in allele frequency:

A principal components analysis (PCA) was used to visualize genome-wide relationships

between sample allele frequencies using the R package *pcadapt* version 4.0 [32]. We calculated

nucleotide diversity for each treatment using a sliding window approach in Popoolation with the

variance-sliding command [33]. Window size was set to 400 bp with step size of 200 bp.

Loci under selection should increase in frequency consistently among replicates due to

differential survival of larvae with adaptive genotypes. Patterns of genetic variation following a

selective sweep are typically observed after many generations of selection. However, over a

single generation of selection, using wild-caught, outbred parents and 37,000 offspring with

recombined genomes, adaptive alleles can be present on multiple different genomic backgrounds

or haplotypes and selective mortality can result in a signal of elevated LD that decays with

physical distance. In this experiment, selective mortality is analogous to a soft sweep where

multiple individuals or haplotypes (from standing variation) carry the adaptive allele that is

under selection [33]. The consistency of allele frequency changes and increases in LD among

replicates can be used to distinguish selected loci from those undergoing neutral drift.

Cochran-Mantel-Haenszel (CMH) tests, a method for identifying consistent changes in

allele frequency in evolution experiments [10], were conducted in R (*mantelhaen.test*)

comparing allele frequency estimates from the four replicates at time point zero (T_0, day 1, pH

8.0) to the four replicate vessels at day 7, pH 8.0 and to the four replicates at day 7, pH 7.5.

Significance thresholds were determined using a Bonferroni correction. We considered any SNPs

within the same gene to be non-independent due to linkage and calculated the number of tests as

the number of independent gene regions; this resulted in a threshold of 0.05/9828 = 5.09e-6. This

is an alternative to the overly conservative Bonferroni correction that ignores linkage and

considers every site independent [34], and is more stringent than a FDR approach with these data

($P < 0.004$).

We compared LD estimates among pairs of SNPs across the genome for selected (CMH

$P < 5.09e-6$) and neutral (non-selected, CMH $P \geq 5.09e-6$) variants using LDx [35]

(supplemental methods). LDx uses a maximum likelihood approach and leverages haplotype

information from single reads to estimate LD between pairs of variants over short distance. To

estimate the decay in LD, we regressed the log of the physical distance with LD between base

pairs for each treatment group with replicate as a random effect using the R package *nlme* [36].

To assess whether the levels of LD present around selected sites could be due to the low number

of variants relative to neutral sites, we randomly subsampled all variants 500 times to match the

number of selected loci for both pH 7.5 and 8.0. We compared the observed slope of decay and

the intercept to permuted distributions.

We used folded allele frequencies to assess if the distribution of starting frequencies of

adaptive loci were distinct from the distribution of neutral loci. To ensure the patterns observed

were not due to bias of the CMH statistic towards detection of variants at low starting allele

frequencies, the observed distributions were compared to permuted distributions with no true

biological signal as described in the supplemental methods.

Gene ontology (GO) enrichment was performed using the weight algorithm in topGO

version 2.22.0 [37]. GO terms for each gene were retrieved from EchinoBase

(http://www.echinobase.org/). Enrichment tests were conducted for each set of significant

variants and genes that had any SNPs in genic or regulatory regions were considered the target

set. Any gene with multiple significant variants was only considered once.

Figure 1: Morphometric analysis of total larval body length. (a) Boxplot of body size in each selection regime. Inset pictures show representative individuals found in pH 8.0 (b) and 7.5 (c).

Results

Strength of selection

Mean total larval body length was lower in pH 7.5 (mean $\pm$ standard error: 239.1 $\pm$

3.1μm) as compared to pH 8.0 (320.9 $\pm$ 2.9 μm) (Fig. 1, $P < 0.001$). Mortality estimates (the

proportion of larvae dead at day 7, supplemental methods) differed in the same direction as body

length, though not significantly, where pH 7.5 had a higher mean mortality of 51.5% + 11.3%

and pH 8.0 was 50.8% + 5.4% ($P = 0.89$). The median selection coefficient (supplemental

methods), was 0.210 and 0.206 for pH 7.5 and 8.0 selected loci, respectively, and was marginally

different ($P = 0.07$). These results suggest that physiological and selective impact was stronger in

pH 7.5 than pH 8.0.

Consistent allele frequency changes among replicate selection lines

Across the twelve pools of larvae (four replicates of each: T_0 and after 7 days in pH 8.0

and pH 7.5), we identified 75,368 variable sites present in or near 9,828 genes with an average

coverage of 123x per site per sample. PCA showed that samples clustered by day and treatment

(Fig. 2a), indicating consistent allele frequency estimates from each biological replicate and

consistent changes in allele frequency in response to treatment conditions. pH 7.5 samples

showed the largest shift from the starting allele frequencies as expected with increased selective

mortality due to treatment (Fig. 2a). Note that one of the D7 pH 8.0 samples was an outlier. To

Figure 2: (a) Principal component analysis of allele frequencies for all 75,368 SNPs at time point zero (T$_0$) and after seven days in the two pH treatments. (b) Mean change in allele frequency for variants showing significant shifts under pH 7.5 and 8.0, and to both pH regimes. Dashed black line represents the 1:1 expectation.

ensure that this sample did not reduce power, we removed this sample, down sampled all

replicates to n=3, and reran analyses; results were consistent with and without this sample.

We found 787 variants (in or near 606 genes) with consistent changes in allele frequency

in response to selection at pH 7.5 and 572 variants (470 genes) that changed in response to pH

8.0 (CMH, *P*<5.09e-06; Fig. 2b; see Table S2 for allele frequency data and CMH results). The

mean change in allele frequency was 0.115 ± 0.049 (mean ± SD) and 0.110 ± 0.046 for pH 7.5

and pH 8.0 responsive variants, respectively, while the genome-wide mean change was 0.028 ±

0.030. 128 (11.6%) of pH 7.5- and pH 8.0-responsive variants overlapped (100 genes).

Of the total assayed variants, 50,691 (67%) were in genic regions while 24,677 (33%)

were in regulatory regions. Matching expectations of chance sampling, for pH 7.5 selected loci,

537 (68%) were genic and 250 (32%) were regulatory (chi-squared, *P* > 0.05). Similarly, pH 8.0

selected loci consist of 401 (70%) genic and 171 (30%) regulatory loci (chi-squared, *P* > 0.05).

Given the rapid decay in linkage disequilibrium (see below), these results suggest that there are

both important coding and putative regulatory pH-responsive loci segregating in populations.

Signals of shared and pH-specific selection

Shifts in allele frequency showed unique and shared signals between pH selection

regimes (Fig. 2b, S1). Significant loci specific to pH 7.5 show a correlation in average allele

frequency change of 0.67 with their non-significant pH 8.0 counterparts ($P < 0.001$) suggesting

that the same loci were responding to treatment though to a lesser degree in pH 8.0; pH 8.0

significant loci reveal the same pattern with a correlation of 0.71 ($P < 0.001$). As expected, loci

that were identified as significant in both treatments showed the strongest correlation in allele

frequency change (0.89, $P < 0.001$). Interestingly, the loci with the most extreme shifts in allele

frequency were significant in both selection regimes (upper right quadrant of Fig. 2b, Fig. S2). In

comparison, the correlation of allele frequency change in non-significant loci was 0.19.

Elevated LD around selected loci

Selected loci in both pH treatments showed strong linkage disequilibrium with the

slowest decay in LD for SNP pairs involving pH 7.5 selected loci (estimated slope of regression

of log distance and LD ± standard error: -0.069 ± 0.005), followed by pH 8.0 selected loci

(slope: -0.072 ± 0.006); both of these decay rates were slower than the -0.111 mean from the

permutations and fell outside of the 95% distribution (pH 7.5 slope: -0.134 to -0.085; pH 8.0

slope: -0.145 to -0.079; Fig. 3a). We observed rapid decay of LD within 100 base pairs, which is

expected given the high levels of genetic diversity, large effective population size, high

fecundity, and high gene flow of this species. As expected, the observed intercepts, LD at zero

distance, (0.306 ± 0.008 and 0.283 ± 0.009) for pH 7.5 and 8.0 selected loci were not

significantly different from neutral expectations as both sets fall within the range of permutation

95% distributions (pH 7.5: 0.310 (0.298, 0.382); pH 8.0: 0.283 (0.282, 0.393)).

Maintenance of pH-adaptive alleles in natural populations

To understand how variants are segregating and maintained in populations, we explored

the starting allele frequencies (T_0) for loci identified as responsive to pH 7.5 (mean ± standard

error: 0.15 ± 0.01), pH 8.0 (0.16 ± 0.01), and to both treatments (0.19 ± 0.01; Fig. 3b). The

distributions of starting allele frequencies for pH 8.0 and 7.5 selected loci were significantly

lower than neutral loci (0.17 ± 0.0004; two-sided Kolmogorov–Smirnov tests, $P < 0.001$), while

the overlapping selected loci were shifted towards higher frequencies ($P < 0.001$). Loci

Figure 3: (a) Decay in linkage disequilibrium with physical distance between SNP pairs. Dashed lines represent decay around non-significant, putatively neutral loci and solid red and blue lines represent decay around pH 7.5 and 8.0 putatively selected variants, respectively. Solid grey shading is the 95% distribution of 500 permutations of down sampled random variants across the genome (based on pH 8.0 for visualization). Note that each of the four replicates per treatment are plotted though may be difficult to see due to the highly concordant decay. (b) Folded allele frequency at T_0. Colored solid lines are the density plot distributions for each set of selected loci. The solid black line represents the median expectation of neutral loci from 1000 permutations of down sampling to the same number of variants as selected loci; grey shading is the 95% distribution of the permutation. Dashed vertical lines show the mean allele frequencies of each set of variants.

responsive to the most extreme selection regime, pH 7.5, were lower in starting allele frequency
than loci responsive to both pH treatments ($P = 0.03$) and to pH 8.0 alone, though this effect was
not significant for the latter ($P = 0.08$). Conversely, loci responsive to only pH 8.0 did not have
lower starting allele frequency than those responding to both treatments ($P = 0.1$). These patterns
hold when selected loci were polarized by the allele increasing in frequency in response to pH
selection (the putative adaptive allele; Fig. S3). We also compared observed allele frequency
distributions of selected loci to distributions of the same number of randomly sampled loci across
the genome (Fig. 3b). The distributions for pH 7.5, pH 8.0, and overlapping selected variants
were significantly different from all 1000 permutations ($P < 0.001$). Lastly, permutations showed
that the CMH statistic was not biased towards loci that started at low frequencies; randomly
shuffled sample IDs were not enriched for low starting allele frequencies (Fig. S4). Finally, we
found that loci with higher minor allele frequencies showed the greatest changes in frequency in
response to selection (Fig. S5), which matches theoretical and simulation expectations [8]. The

excess of low allele frequency variants among selected loci is consistent across functional classes

of variants (non-synonymous, synonymous, intronic, regulatory; Fig. S6).

Genetic diversity (π) was 0.0158 at T_0 and decreased to 0.0129 and 0.0128 after seven

days at pH 8.0 and 7.5, respectively (one-sided Wilcoxon rank-sum $P < 0.001$). The decrease in

diversity between the pH treatments was not different ($P = 0.12$), demonstrating that, while the

selection regimes were of different magnitude, the relative loss of genetic diversity through time

was similar.

Functional enrichment analysis revealed that genes under selection were enriched for

specific biological functions (Table S3). We observed enrichment for 17, 10, and 14 biological

processes GO terms for pH 7.5, pH 8.0, and overlapping variants, respectively. Of these, only

hexose metabolic process (GO:0019318) was significant across all three variant sets. Other

related terms shared across sets included those involved in cell growth (GO:0030307;

GO:0032467), metabolism and energy production (GO:0001678; GO:0009267; GO:0019318;

GO:0006096; GO:0006869; GO:0009061). pH 7.5 was enriched for unique functional processes,

particularly those involved in cytoskeleton remodeling (GO:0051639; GO:0008154) and

intracellular pH regulation (GO:0007035). pH 8.0 and overlapping variants showed limited

signal of unique enrichment, see Table S3 for all enriched categories.

252
Discussion

We show that genetic response to moderate and extreme low pH relies on shared and

unique mechanisms and that rare variants are important for survival in low pH conditions. We

also identify a class of variants responsive to both pH levels (11.6% of pH-responsive loci) that

are maintained at higher frequencies in natural populations, potentially by balancing selection in

a spatially and temporally variable pH environment. We further demonstrate the utility of single

generation selection experiments to identify the genetic targets of adaptation, which is

particularly useful for testing capacity for response to conditions that will be experienced more

frequently in the future but are rare events in nature today. Using sequence capture of genomic

DNA, we quantified shifts in allele frequency that represent differential survival of genotypes

during low pH selection in sea urchins. This approach has wide potential application for

identifying responsive genetic variants from wild populations. Our results indicate that standing

genetic variation maintained with large population sizes will be critical for survival in future

environmental conditions.

Detecting adaptive loci from standing genetic variation

Recent selective sweeps result in decreased genetic variation and increased linkage

disequilibrium (LD) around adaptive loci [38,39]. Even in a single generation of selection, we

have the potential to detect selective sweeps as increased LD around adaptive loci from standing

variation. In the present experiment, adaptive genetic variation can be present in many different

genomic backgrounds through: 1) the use of 25 wild-caught, outbred parents, 2) the

recombination that occurred between the diploid genomes of each parent during gametogenesis,

and 3) experimental fertilization of eggs from each female with sperm from each male. Seeding

each replicate with 37,000 embryos further amplified the number of uniquely recombined

genomes. We expect that differential survival of larvae with specific combinations of alleles will

yield a pattern consistent with soft selective sweeps where adaptive loci are present on multiple

haplotypes [21,40,41]; we observe increased LD in regions surrounding putatively adaptive loci

to match this expectation. Permutation tests demonstrate that this increase in LD was not a

byproduct of our test statistic or sampling noise (Fig. 3a). In addition, the consistent results

among replicates for each treatment provide strong evidence that our approach identifies

genomic regions that are true targets of selection, rather than due to drift or chance changes in

allele frequency.

Typical evolve and resequence studies rely on multiple generations of selection to

identify adaptive shifts in allele frequency. To increase power to distinguish between drift and

selection, one can increase the selection coefficient, number of generations, population size,

number of replicates, or starting genetic diversity [12,42,43]. Because purple sea urchin take two

years to reach reproductive maturity [44] we can use only a single generation of selection, but

their high fecundity and small offspring size enable us to maximize population size and genetic

diversity. This tradeoff is particularly useful for performing selection experiments in long-lived

organisms. One starts with outbred wild-caught individuals, which will increase starting genetic

diversity relative to inbred lab strains or isofemale lines [12,45,46]. Inference is limited to the

genetic diversity present in the starting individuals and may miss variation present at different

points in space or time and will miss the potential of new beneficial mutations, which may be an

important mode for adaptation [21]. However, starting with outbred parents maximizes the

amount of recombined genetic variation, avoiding the large linkage blocks that can plague

experimental evolution studies from lab lines. These narrow linkage windows also improve the

chances of identifying the specific loci targeted by selection.

Maintenance of adaptive standing genetic variation

Variants that responded to low pH treatments were present in the starting population (T_0)

at low allele frequencies relative to neutral alleles (Fig. 3b). Low starting allele frequencies of

responsive variants could suggest that surviving offspring were from a single parent. However,

this is unlikely due to the decay of LD around selected loci, indicating that many different

genetic backgrounds were targeted by selection. In addition, the large number of responsive loci

and genes suggests that survival in low pH conditions is a polygenic trait, a characteristic of

adaptation to complex environments in other experimental evolution systems [47]. Thus it is

unlikely that surviving offspring inherited all the adaptive alleles from a single parent.

The low starting allele frequency of uniquely response low-pH adaptive variants argues

that these loci are not maintained by balancing selection or overdominance, but by one of three

processes. First, the alleles may be beneficial under low pH but neutral or slightly deleterious at

ambient conditions, known as conditional neutrality [48]. Alternatively, antagonistic pleiotropy

can alter the rank fitness of alleles across different environments, maintaining genetic variation

across space or time [49]. Finally, purple sea urchins inhabit rocky reefs across a broad

geographic range [24,50] where pH conditions vary due to natural processes such as upwelling

[17]. Low pH-responsive loci in the lab have been shown to have allele frequencies correlated

with local pH conditions in the field, suggesting local adaptation [25]. High gene flow across the

geographic range and migration-selection balance can result in the maintenance of low frequency

alleles that are not adaptive in a local environment [51]. Similar selection on low frequency

alleles has been observed in *D. melanogaster* during experimental evolution to high temperature

[52]. Interestingly, this shift was observed for adaptation to high but not low temperature.

Conversely, Barghi et al., find that experimental adaptation to temperature in *Drosophila* recruits

alleles at higher frequency than neutral loci [47], providing additional evidence that the CMH

statistic is not inherently biased towards low starting minor allele frequencies.

In contrast, loci overlapping in response to both moderate and extreme pH had higher

starting allele frequencies than neutral loci and loci responsive to either pH regime alone (Fig.

3b). These "low-pH-essential" alleles may be maintained at higher allele frequency due to

balancing selection in the wild. That is, the spatial and temporal heterogeneity of pH conditions

experienced by purple sea urchins across the species range and across life history stages likely

maintains these alleles at intermediate frequencies through fluctuating selection or spatially

balancing selection [53]. We note that these shared loci may also include variants involved in

selection for the general lab culture conditions. While the design of this study precludes the

isolation of lab-adapted variants, the functional classes of genes we identify here are also

responsive to moderate low pH conditions (pH 7.8) in purple sea urchin larvae generated from

multiple populations [14,23] and are correlated with environmental pH in wild populations [23].

In addition, lower pH results in more extreme patterns of reduced LD decay, allele frequency

change, and morphological impacts than moderate pH conditions, suggesting that, while lab

selection may be present, pH selection is the main driver of the results in the present study.

Mechanisms for response to low pH conditions

Selected loci are enriched for a range of biological processes related to maintaining

cellular homeostasis. Across all sets of selected variants, we observe enrichment in processes

related to metabolism and energy production (GO:0001678, GO:0009267, GO:0019318,

GO:0006096, GO:0006869, GO:0009061). Alterations to metabolic processes, energy demands,

and resource allocations are primary responses to low pH environments [54,55]. Previous

transcriptomic studies in *S. purpuratus* have shown that these classes of genes are differentially

regulated in response to low pH stress [18,56]. The energy required for acid-base regulation is

increased under acidic conditions because the calcification process, production of calcium

carbonate ($CaCO_3$), generates excess protons that must be removed to maintain acid-base balance

[55]. We also see shared enrichment for genes involved in cell proliferation and damage control,

including those implicated in cell growth and replication (GO:0030307, GO:0008361,

GO:0032467) and protein transport (GO:0006606). These results match our morphometric

findings of stunted larval growth and suggest that fine control of cellular growth may be

important for survival in low pH conditions.

Two genes of interest that showed changes in allele frequency greater than 20% in both

pH treatments perform functions related to cytoskeletal and membrane dynamics: ELMOD2 (SP-

ELMOD2/SPU_007564), and Focadhesin (KIAA1797/SPU_015184). ELMOD2 had the

greatest changes in allele frequency in both treatments and plays an important role in regulating

membrane traffic and secretion, phospholipid metabolism, and actin/cytoskeleton dynamics [57-

59]. The Focadhesin protein plays an important role in integrating intra- and extracellular

information as a subcellular structural protein that receives biomechanical and biochemical

signals between the cytoskeleton and the extracellular matrix [60,61]. We identify three SNPs,

two of which change amino acids, which show shifts in allele frequency greater than 20% in this

putatively low-pH adaptive gene.

Selection in pH 7.5, the edge of what *S. purpuratus* urchins experience in nature,

revealed unique adaptive targets to extreme pH selection. We observe enrichment for 'actin

polymerization or depolymerization' (GO:0008154) and actin filament network formation

(GO:0051639). It has been hypothesized that changes in actin abundance are related to

cytoskeleton remodeling due to intracellular stress during acclimation of oysters to climate

change conditions [62,63]. Evans et al., [16] find enrichment for expression of genes involved in

actin folding in *S. purpuratus*, suggesting that the cytoskeleton is a target of pH stress. We also

found enrichment for vacuolar acidification (GO:0007035); regulation of pH in intracellular

compartments can maintain pH homeostasis when extracellular pH is altered [64].

We identify putative adaptive alleles in proteins that play important roles in bicarbonate

uptake and pH regulation. TASK2 is a pH-sensitive K+ transporter [65] and is an important

component of bicarbonate (HCO3-) uptake in higher eukaryotes [66]. We identify pH 7.5-

selected alleles in TASK2 (SPU_003613; Fig. 4). Recent studies in purple sea urchin larvae have

shown that the regulation of intracellular pH is linked to biomineralization and that

compensatory changes in expression of other bicarbonate transporters (Slc4) are necessary to

maintain calcification under acidic conditions [65]. Under extreme low pH conditions, TASK2

function may be under selection to ensure continued HCO3- uptake to regulate intracellular pH

and facilitate skeletogenesis in sea urchin larvae. Empirical work is needed to validate this

hypothesis.

A second protein of interest is a membrane bound and excreted enzyme in the carbonic

anhydrase family (SPU_012518). Carbonic anhydrases catalyzes the hydration of CO_2 to

Figure 4: (a) Mean change in allele frequency of pH treatments relative to T_0 across a ~150 bp region of scaffold 542 in the K$^+$ transporter TASK2. (b) *q*-values corresponding to the allele frequency changes in (a). The dashed horizontal line indicates the significance threshold of 5.09e-06.

bicarbonate [66] and have been shown to be responsive to experimental acidification in many organisms including purple sea urchin [27,56], corals [67], anemones [68], mussels [69], oysters [70], and giant kelp [71]. In the present study, carbonic anhydrase (SPU_012518), appears to be responding to pH 7.5 where 3/12 variants have *P*-values < 0.01. However, these loci do not meet our stringent significance cutoff highlighting that results of any genome-wide scan for selection are sensitive to false negatives (and false positives), missing loci that may truly be under selection. However, given the strong signal of several loci across the gene, the established functional role in skeletogenesis in sea urchin larvae [72], and that the enzyme has been identified as pH-responsive in many ocean acidification studies across a broad range of taxa, carbonic anhydrase is an ideal candidate for future functional studies. Linking the genetic variation identified in the present study to functional phenotypes at the level of this enzyme would improve our understanding of the capacity of organisms to adapt to future conditions from standing genetic variation.

Conclusion

We utilize a single generation selection experiment to reveal that purple sea urchins have

genetic variation responsive to extreme and moderate low pH selection. The low variation among

replicates, increased LD around selected loci, and enrichment for biological functions related to

pH adaptation suggest that this approach accurately identifies adaptive loci. Further, we explore

patterns of starting allele frequency to infer the evolutionary mechanisms underlying the

maintenance of adaptive standing genetic variation. This work provides a framework for future

studies assessing the genetic basis of adaptive responses to environmental stressors. Increasing

the temporal sampling throughout development would reveal how different life stages

differentially respond to selection, while functional studies of top candidates will help to validate

their mechanistic and evolutionary significance. The results presented here show that *S.*

purpuratus possesses genetic variation that is responsive to extreme low pH conditions and

identifies a class of variants essential for adapting to low pH that are likely maintained by

balancing selection. We also identify sets of rare variants that are unique to each pH selection

regime suggesting that the preservation of large population sizes will be important in the

maintenance of this rare adaptive variation.

Data and code availability

Sequence data is available at the National Center for Biotechnology Information (NCBI SRA

BioProject: PRJNA479817). Phenotype data is available as part of the supplementary material.

All code for analysis is available at https://github.com/PespeniLab/urchin_single_gen_selection.

Competing Interests

We have no competing interests.

Author Contributions

RB analyzed data and wrote the manuscript. AG designed and conducted the experiment and

assisted with data analysis and drafting the manuscript. KH helped perform the experiment and

analyzed the morphometric data. HH helped design aquaculture facilities and pilot the

experiment. MP designed the experiment and wrote the manuscript.

Acknowledgements

We thank Pete Halmay and Patrick Leahy for urchin collections. Jeremy Arenos for assistance

with imaging and image analysis, Jason Hodin and Justin McAlister for helpful discussions on

urchin larval culture, Lauren Ashlock for assistance with conducting the experiment, Mike

Austin and the Vermont Advanced Computing Core for server space and maintenance, and all

members of the Pespeni lab and Stephen Keller for helpful discussions.

Funding

This work was supported by the National Science Foundation (NSF) grants OCE-1559075 (to

MHP) and OIA-1736253 (to MHP) and the NSF Graduate Research Fellowship Program DGE-

1451866 (to AG).

References

1. Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N.,
Raven, P. H., Roberts, C. M. & Sexton, J. O. 2014 The biodiversity of species and
their rates of extinction, distribution, and protection. *Science* **344**, 1246752–
1246752. (doi:10.1126/science.1246752)

2. Rahmstorf, S. & Coumou, D. 2011 Increase of extreme events in a warming world.
Proc. Natl. Acad. Sci. **108**, 17905–17909. (doi:10.1073/pnas.1101766108)

3. Hoffmann, A. A. & Sgrò, C. M. 2011 Climate change and evolutionary adaptation.
Nature **470**, 479–485. (doi:10.1038/nature09670)

4. Denny, M. W. & Dowd, W. W. 2012 Biophysics, environmental stochasticity, and
the evolution of thermal safety margins in intertidal limpets. *J. Exp. Biol.* **215**, 934–
947. (doi:10.1242/jeb.058958)

5. Harley, C. D. G. & Paine, R. T. 2009 Contingencies and compounded rare
perturbations dictate sudden distributional shifts during periods of gradual climate
change. *Proc. Natl. Acad. Sci.* **106**, 11172–11176. (doi:10.1073/pnas.0904946106)

6. Lande, R. 2009 Adaptation to an extraordinary environment by evolution of
phenotypic plasticity and genetic assimilation. *J. Evol. Biol.* **22**, 1435–1446.
(doi:10.1111/j.1420-9101.2009.01754.x)

7. Grant, P. R., Grant, B. R., Huey, R. B., Johnson, M. T. J., Knoll, A. H. & Schmitt, J.
2017 Evolution caused by extreme events. *Philos. Trans. Royal Soc. B* **372**,
20160146. (doi:10.1098/rstb.2016.0146)

8. Kofler, R. & Schlotterer, C. 2014 A guide for the design of evolve and resequencing
studies. *Mol. Biol. Evol.* **31**, 474–483. (doi:10.1093/molbev/mst221)

9. Burghardt, L. T., Epstein, B., Guhlin, J., Nelson, M. S., Taylor, M. R., Young, N.
D., Sadowsky, M. J. & Tiffin, P. 2018 Select and resequence reveals relative fitness
of bacteria in symbiotic and free-living environments. *Proc. Natl. Acad. Sci.* **115**,
2425–2430. (doi:10.1073/pnas.1714246115)

10. Schlotterer, C., Kofler, R., Versace, E., Tobler, R. & Franssen, S. U. 2015
Combining experimental evolution with next-generation sequencing: a powerful tool
to study adaptation from standing genetic variation. *Heredity* **114**, 431–440.
(doi:10.1038/hdy.2014.86)

11. Franks, S. J. & Hoffmann, A. A. 2012 Genetics of climate change adaptation. *Annu.*
Rev. Genet. **46**, 185–208. (doi:10.1146/annurev-genet-110711-155511)

12. Kessner, D. & Novembre, J. 2015 Power analysis of artificial selection experiments
using efficient whole genome simulation of quantitative traits. *Genetics* **199**, 991–
1005. (doi:10.1534/genetics.115.175075)

13.   Barrett, R. D. H. et al. 2019 Linking a mutation to survival in wild mice. *Science*
481   **363**, 499–504. (doi:10.1126/science.aav3824)

482   14.   Soria-Carrasco, V. et al. 2014 Stick insect genomes reveal natural selection's role in
483   parallel speciation. *Science* **344**, 738–742. (doi:10.1126/science.1252136)

484   15.   Pespeni, M. H. et al. 2013 Evolutionary change during experimental ocean
485   acidification. *Proc. Natl. Acad. Sci.* **110**, 6937–6942.
486   (doi:10.1073/pnas.1220673110)

487   16.   Chan, F. et al. 2017 Persistent spatial structuring of coastal ocean acidification in the
488   California Current System. *Sci. Rep.* **7**, 2526. (doi:10.1038/s41598-017-02777-y)

489   17.   Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D. & Hales, B. 2008
490   Evidence for Upwelling of Corrosive 'Acidified' Water onto the Continental Shelf.
491   *Science* **320**, 1490–1492. (doi:10.1126/science.1155676)

492   18.   Evans, T. G., Pespeni, M. H., Hofmann, G. E., Palumbi, S. R. & Sanford, E. 2017
493   Transcriptomic responses to seawater acidification among sea urchin populations
494   inhabiting a natural pH mosaic. *Mol. Ecol.* **26**, 2257–2275. (doi:10.1111/mec.14038)

495   19.   Hofmann, G. E. et al. 2011 High-frequency dynamics of acean pH: A multi-
496   ecosystem comparison. *PLoS ONE* **6**. (doi:10.1371/journal.pone.0028983)

497   20.   Chevin, L.-M., Lande, R. & Mace, G. M. 2010 Adaptation, plasticity, and extinction
498   in a changing environment: towards a predictive theory. *PLoS Biol* **8**, e1000357.
499   (doi:10.1371/journal.pbio.1000357.s001)

500   21.   Messer, P. W. & Petrov, D. A. 2013 Population genomics of rapidad aptation by soft
501   selective sweeps. *Trends Ecol. Evol.*, 1–11. (doi:10.1016/j.tree.2013.08.003)

502   22.   Barrett, R. D. & Schluter, D. 2008 Adaptation from standing genetic variation.
503   *Trends Ecol. Evol.* **23**, 38–44. (doi:10.1016/j.tree.2007.09.008)

504   23.   Flowers, J. M., Schroeter, S. C. & Burton, R. S. 2002 The recruitment sweepstakes
505   has many winners: Genetic evidence from the sea urchin *Strongylocentrotus*
506   *purpuratus*. *Evolution* **56**, 1445–1453. (doi:10.1111/j.0014-3820.2002.tb01456.x)

507   24.   Pespeni, M. H. & Palumbi, S. R. 2013 Signals of selection in outlier loci in a widely
508   dispersing species across an environmental mosaic. *Mol. Ecol.* **22**, 3580–3597.
509   (doi:10.1111/mec.12337)

510   25.   Pespeni, M. H., Chan, F., Menge, B. A. & Palumbi, S. R. 2013 Signs of adaptation
511   to local pH conditions across an environmental mosaic in the california current
512   ecosystem. *Integrative and Comparative Biology* **53**, 857–870.
513   (doi:10.1093/icb/ict094)

26.   Kelly, M. W., Padilla-Gamiño, J. L. & Hofmann, G. E. 2013 Natural variation and
the capacity to adapt to ocean acidification in the keystone sea urchin
Strongylocentrotus purpuratus. *Glob Change Biol* **19**, 2536–2546.
(doi:10.1111/gcb.12251)

27.   Evans, T. G., Chan, F., Menge, B. A. & Hofmann, G. E. 2013 Transcriptomic
responses to ocean acidification in larval sea urchins from a naturally variable pH
environment. *Mol. Ecol.* **22**, 1609–1625. (doi:10.1111/mec.12188)

28.   R Core Team 2016 *R: A language and environment for statistical computing.*
Vienna, Austria: R Foundation for Statistical Computing.

29.   Bates, D., Mächler, M., Bolker, B. & Walker, S. 2015 Fitting Linear Mixed-Effects
Models Using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

30.   Li, H. 2013 Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. *arXiv preprint arXiv:1303.3997*

31.   Schlötterer, C., Tobler, R., Kofler, R. & Nolte, V. 2014 Sequencing pools of
individuals — mining genome-wide polymorphism data without big funding. *Nature*
Publishing Group **15**, 749–763. (doi:10.1038/nrg3803)

32.   Luu, K., Bazin, E. & Blum, M. G. B. 2017 pcadapt: an R package to perform
genome scans for selection based on principal component analysis. *Mol. Ecol.*
Resour. **17**, 67–77. (doi:10.1111/1755-0998.12592)

33.   Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik,
A., Kosiol, C. & Schlötterer, C. 2011 PoPoolation: A toolbox for population genetic
analysis of next generation sequencing data from pooled individuals. *PLoS ONE* **6**,
e15925. (doi:10.1371/journal.pone.0015925)

34.   Johnson, R. C., Nelson, G. W., Troyer, J. L., Lautenberger, J. A., Kessing, B. D.,
Winkler, C. A. & O'Brien, S. J. 2010 Accounting for multiple comparisons in a
genome-wide association study (GWAS). *BMC Genomics* **11**, 724.
(doi:10.1186/1471-2164-11-724)

35.   Feder, A. F., Petrov, D. A. & Bergland, A. O. 2012 LDx: Estimation of linkage
disequilibrium from high-throughput pooled resequencing data. *PLoS ONE* **7**,
e48588. (doi:10.1371/journal.pone.0048588)

36.   Pinheiro, J., Bates, D., DebRoy, S.R Core Team 2018 *nlme: Linear and Nonlinear*
Mixed Effects Models. 3.1–137.

37.   Alexa, A. & Rahnenfuhrer, J. 2010 topGO: enrichment analysis for gene ontology.
11, R14. (doi:10.1186/gb-2010-11-2-r14)

38.   Kim, Y. & Nielsen, R. 2004 Linkage disequilibrium as a signature of selective
sweeps. *Genetics* **167**, 1513–1524. (doi:10.1534/genetics.103.025387)

39.   Przeworski, M. 2002 The signature of positive selection at randomly chosen loci. *Genetics* **160**, 1179–1189.

40.   Hermisson, J. & Pennings, P. S. 2017 Soft sweeps and beyond: understanding the patterns and probabilities of selection footprints under rapid adaptation. *Methods Ecol Evol* **8**, 700–716. (doi:10.1111/2041-210X.12808)

41.   Hermisson, J. & Pennings, P. S. 2005 Soft sweeps: Molecular population genetics of adaptation from standing genetic variation. *Genetics* **169**, 2335–2352. (doi:10.1534/genetics.104.036947)

42.   Gompert, Z., Egan, S. P., Barrett, R. D. H., Feder, J. L. & Nosil, P. 2017 Multilocus approaches for the measurement of selection on correlated genetic loci. *Mol. Ecol.* **26**, 365–382. (doi:10.1111/mec.13867)

43.   Baldwin-Brown, J. G., Long, A. D. & Thornton, K. R. 2014 The power to detect quantitative trait loci using resequenced, experimentally evolved populations of diploid, sexual organisms. *Mol. Biol. Evol.* **31**, 1040–1055. (doi:10.1093/molbev/msu048)

44.   Leahy, P. S. 1986 Laboratory Culture of *Strongylocentrotus purpuratus* Adults, Embryos, and Larvae. *Methods Cell Biol.* **27**, 1–13. (doi:10.1016/S0091-679X(08)60339-8)

45.   Turner, T. L. & Miller, P. M. 2012 Investigating natural variation in *Drosophila* courtship song by the evolve and resequence approach. *Genetics* **191**, 633–642. (doi:10.1534/genetics.112.139337)

46.   Orozco-terWengel, P., Kapun, M., Nolte, V., Kofler, R., Flatt, T. & Schlötterer, C. 2012 Adaptation of *Drosophila* to a novel laboratory environment reveals temporally heterogeneous trajectories of selected alleles. *Mol. Ecol.* **21**, 4931–4941. (doi:10.1111/j.1365-294X.2012.05673.x)

47.   Barghi, N. et al. 2019 Genetic redundancy fuels polygenic adaptation in Drosophila. *PLoS Biol* **17**, e3000128. (doi:10.1371/journal.pbio.3000128)

48.   Anderson, J. T., Lee, C.-R., Rushworth, C. A., Colautti, R. I. & Mitchell-Olds, T. 2012 Genetic trade-offs and conditional neutrality contribute to local adaptation. *Mol. Ecol.* **22**, 699–708. (doi:10.1111/j.1365-294X.2012.05522.x)

49.   Mitchell-Olds, T., Willis, J. H. & Goldstein, D. B. 2007 Which evolutionary processes influence natural genetic variation for phenotypic traits? *Nat Rev Genet* **8**, 845–856. (doi:10.1038/nrg2207)

50.   Pespeni, M. H., Oliver, T. A., Manier, M. K. & Palumbi, S. R. 2010 Restriction site tiling analysis: accurate discovery and quantitative genotyping of genome-wide polymorphisms using nucleotide arrays. *Genome Biol* **11**, R44. (doi:10.1186/gb-2010-11-4-r44)

51.   Savolainen, O., Lascoux, M. & Merilä, J. 2013 Ecological genomics of local
adaptation. *Nature Publishing Group* **14**, 807–820. (doi:10.1038/nrg3522)

52.   Tobler, R., Franssen, S. U., Kofler, R., Orozco-terWengel, P., Nolte, V., Hermisson,
J. & Schlötterer, C. 2014 Massive habitat-specific genomic response in *D.*
melanogaster populations during experimental evolution in hot and cold
environments. *Mol. Biol. Evol.* **31**, 364–375. (doi:10.1093/molbev/mst205)

53.   Charlesworth, D. 2006 Balancing selection and Its effects on sequences in nearby
genome regions. *PLoS Genet* **2**, e64–6. (doi:10.1371/journal.pgen.0020064)

54.   Pan, T. C. F., Applebaum, S. L. & Manahan, D. T. 2015 Experimental ocean
acidification alters the allocation of metabolic energy. *Proc. Natl. Acad. Sci. U.S.A.*
112, 4696–4701. (doi:10.1073/pnas.1416967112)

55.   Stumpp, M. et al. 2012 Acidified seawater impacts sea urchin larvae pH regulatory
systems relevant for calcification. *Proc. Natl. Acad. Sci.* **109**, 18192–18197.
(doi:10.1073/pnas.1209174109/)

56.   Todgham, A. E. & Hofmann, G. E. 2009 Transcriptomic response of sea urchin
larvae Strongylocentrotus purpuratus to CO_2-driven seawater acidification. *J. Exp.*
Biol. **212**, 2579–2594. (doi:10.1242/jeb.032540)

57.   Suzuki, M., Murakami, T., Cheng, J., Kano, H., Fukata, M. & Fujimoto, T. 2015
ELMOD2 is anchored to lipid droplets by palmitoylation and regulates adipocyte
triglyceride lipase recruitment. *Mol. Biol. Cell* **26**, 2333–2342.
(doi:10.1091/mbc.E14-11-1504)

58.   East, M. P., Bowzard, J. B., Dacks, J. B. & Kahn, R. A. 2012 ELMO
domains,evolutionary and functional characterization of a novel GTPase-activating
protein (GAP) domain for Arf protein family GTPases. *J. Biol. Chem.* **287**, −39553.
(doi:10.1074/jbc.M112.417477)

59.   Ivanova, A. A., East, M. P., Yi, S. L. & Kahn, R. A. 2014 Characterization of
recombinant ELMOD (cell engulfment and motility domain) proteins as GTPase-
activating proteins (GAPs) for ARF family GTPases. *J. Biol. Chem.* **289**, 11111–
11121. (doi:10.1074/jbc.M114.548529)

60.   Geiger, B., Spatz, J. P. & Bershadsky, A. D. 2009 Environmental sensing through
focal adhesions. *Nat Rev Mol Cell Biol* **10**, 21–33. (doi:10.1038/nrm2593)

61.   Burridge, K. & ChrzanowskaWodnicka, M. 1996 Focal adhesions, contractility, and
signaling. *Annu. Rev. Cell Dev. Biol.* **12**, 463–518.
(doi:10.1146/annurev.cellbio.12.1.463)

62.   Tomanek, L. 2014 Proteomics to study adaptations in marine organisms to
environmental stress. *J Proteomics* **105**, 92–106. (doi:10.1016/j.jprot.2014.04.009)

63. Goncalves, P., Thompson, E. L. & Raftos, D. A. 2017 Contrasting impacts of ocean
acidification and warming on the molecular responses of CO_2-resilient oysters. 1–
15. (doi:10.1186/s12864-017-3818-z)

64. Kane, P. M. 2006 The where, when, and how of organelle acidification by the yeast
vacuolar H^+-ATPase. *Microbiology and Molecular Biology Reviews* **70**, 177–191.
(doi:10.1128/MMBR.70.1.177-191.2006)

65. Hu, M. Y., Yan, J.-J., Petersen, I., Himmerkus, N., Bleich, M. & Stumpp, M. 2018
A SLC4 family bicarbonate transporter is critical for intracellular pH regulation and
biomineralization in sea urchin embryos. *eLife* **7**, 222. (doi:10.7554/eLife.36600)

66. Lindskog, S. 1997 Structure and mechanism of carbonic anhydrase. *Pharmacol.*
Ther. **74**, 1–20. (doi:10.1016/S0163-7258(96)00198-2)

67. Zoccola, D., Innocenti, A., Bertucci, A., Tambutté, E., Supuran, C. & Tambutté, S.
2016 Coral carbonic anhydrases: Regulation by ocean acidification. *Marine Drugs*
14, 109–11. (doi:10.3390/md14060109)

68. Ventura, P., Jarrold, M. D., Merle, P.-L., Barnay-Verdier, S., Zamoum, T., Rodolfo-
Metalpa, R., Calosi, P. & Furla, P. 2016 Resilience to ocean acidification: decreased
carbonic anhydrase activity in sea anemones under high pCO_2 conditions. *Mar.*
Ecol. Prog. Ser. **559**, 257–263. (doi:10.3354/meps11916)

69. Fitzer, S. C., Phoenix, V. R., Cusack, M. & Kamenos, N. A. 2014 Ocean
acidification impacts mussel control on biomineralisation. *Sci. Rep.* **4**, 6218.
(doi:10.1038/srep06218)

70. Wang, X., Wang, M., Jia, Z., Qiu, L., Wang, L., Zhang, A. & Song, L. 2017 A
carbonic anhydrase serves as an important acid-base regulator in pacific oyster
Crassostrea gigas exposed to elevated CO_2: implication for physiological responses
of mollusk to ocean acidification. *Mar Biotechnol* **19**, 22–35. (doi:10.1007/s10126-
017-9734-z)

71. Fernández, P. A., Roleda, M. Y. & Hurd, C. L. 2015 Effects of ocean acidification
on the photosynthetic performance, carbonic anhydrase activity and growth of the
giant kelp *Macrocystis pyrifera*. *Photosynthesis Research*, 1–13.
(doi:10.1007/s11120-015-0138-5)

72. Zito, F., Koop, D., Byrne, M. & Matranga, V. 2015 Carbonic anhydrase inhibition
blocks skeletogenesis and echinochrome production in Paracentrotus lividus and
Heliocidaris tuberculata embryos and larvae. *Development, Growth &*
Differentiation **57**, 507–514.