Title: Rare genetic variation is important for survival in extreme low pH conditions

Authors: Reid S. Brennan[1,2], April D. Garrett[1], Kaitlin E. Huber, Heidi Hargarten, Melissa H.

Pespeni[2]

Affiliation: Department of Biology, University of Vermont, Burlington, VT, USA

[1]Authors contributed equally to this work

[2]Corresponding authors

Email: Reid.Brennan@uvm.edu; mpespeni@uvm.edu

Running head: Low pH adaptation

Keywords: standing genetic variation; ocean acidification; adaptation; experimental evolution;

select and resequence; extreme events

Abstract

Population persistence through increasingly frequent extreme environmental fluctuations

will require selection on standing genetic variation. While some species have shown the capacity

to adapt to mean future conditions, the ability to survive and potentially adapt to extreme events

is unknown. Here we used pooled capture sequencing to test for adaptive capacity and identify

genetic variation responsive to moderate (pH 8.0) and extreme (pH 7.5) low pH conditions using

single generation selection experiments on hundreds of thousands of *Strongylocentrotus*

purpuratus sea urchin larvae generated from wild-caught adults. The single generation of

selection showed consistent shifts in allele frequencies across replicate cultures and increased

linkage disequilibrium around selected loci, revealing selective sweeps from standing variation.

We found extreme pH selection targeted alleles at low frequency in the population while variants

that responded to both moderate and extreme pH selection started at higher allele frequencies,

suggesting maintenance by balancing selection. Variants with the greatest changes in allele

frequencies performed functions related to lipid metabolism, pH tolerance, membrane

trafficking, and regulation of actin/cytoskeleton dynamics. These results highlight that survival

in extreme conditions relies on low frequency standing genetic variation that must be maintained

by large population sizes.

Introduction

As temperatures increase and oceans become more acidic, many marine species are at

high risk of decline and extinction [1]. In addition to changing global averages, the frequency

and intensity of extreme events are also increasing [2]. With small or gradual changes in

conditions, organisms may acclimate through physiological plasticity or migrate to suitable

habitats where possible, but some amount of genetic adaptation will be necessary for continued

population persistence, particularly in the context of extreme events [3]. Indeed, extreme events

rather than average conditions set the physiological and biogeographic limits of individuals and

populations [4-6]. Though limits set by rare extreme events will be critical for population

persistence, the genetic mechanisms that could allow such rapid adaptation have rarely been

explored [7].

Evolve or select & resequence, a type of experimental evolution, is particularly powerful

and promising for understanding capacity to adapt to future conditions and extreme events. In

this approach, one artificially induces the selective treatment and directly measures genetic

response during adaptation [8-11]. While evolve and resequence studies typically leverage 15-20

generations of selection [10,12], we have empirically found that adaptive genetic variation can

be identified with a single generation of selection by using 1000s of small offspring generated

from highly heterozygous, outbred parents collected from the wild [13].

The purple sea urchin, *Strongylocentrotus purpuratus*, is an ideal model for

understanding the process of adaptation from adaptive standing genetic variation. This species is

found on inter- and subtidal rocky reefs and kelp forests across a broad latitudinal range from

Alaska to Baja California, Mexico in the California Current Marine Ecosystem (CCME). They

experience high heterogeneity in environmental conditions in both time and space [14-16],

increasing the likelihood of the presence of adaptive standing genetic variation [17-19]. Further,

this species is a good system to understand the genetics of rapid adaptation due to large census

and effective population sizes that contribute to high standing genetic variation [20,21]. While

gene flow is high due to pelagically dispersing larvae, populations nevertheless show increased

frequencies of variants putatively adaptive to local temperature and pH conditions [22]. Previous

studies have shown that *S. purpuratus* larvae have the physiological and genetic capacity to

adaptively respond to an acidifying ocean [13,15,23], but these studies have focused on relatively

mild conditions and have not investigated the capacity to respond to extreme pH conditions that

are predicted to increase in frequency in the near future [2,24].

Here, we perform single generation selection experiments in moderate (pH 8.0) and

extreme low (pH 7.5) pH conditions using purple sea urchin larvae generated from 25 wild-

caught adults. We test the hypotheses that (1) there will be unique genetic variation responsive

to extreme low pH conditions, and (2) that genomic patterns of variation, such as linkage

disequilibrium and starting allele frequencies, will provide insight into the evolutionary

mechanisms that maintain adaptive standing genetic variation for survival in moderate and

extreme conditions, and (3) pooled sequencing of genomic DNA from larvae before and after

treatments will show consistent changes in allele frequency across replicate cultures due to

selective mortality through development.

Methods

Sample collection & experiment:

Purple sea urchin adults were collected in September 2016 from San Diego, CA, shipped

overnight to the University of Vermont, and the experiment began immediately upon their arrival

(n=25 total: 14 females and 11 males). Spawning was induced with 0.5M KCl in Instant Ocean

artificial seawater (ASW) (Instant Ocean, Blacksburg, VA) at 14°C and salinity of 35ppt. For

each female, 200,000 eggs were put into each treatment, moderate (pH 8.0) and extreme (pH 7.5)

low pH, and fertilized by evenly pooled sperm from all males. We chose these pH conditions

based on empirical data collected in the CCME. The current average open ocean and intertidal

pH is 8.1, though conditions in the CCME frequently drop to pH 8.0 (daily), and only rarely drop

as low as pH 7.5 (once in three month upwelling period; [14,16,24,25]). Fertilized eggs were

pooled across all females by pH and seeded into four replicate culturing vessels per treatment

(37,000 eggs per 3.7 L vessel). Developing embryos were sampled at day 1 and day 7 post

fertilization for morphometric and genomic analyses. See the supplemental methods for

expanded details.

Morphometrics:

7-day old 4-armed pluteus larvae were photographed for morphometric analysis using a

Photometrics Scientific CoolSNAP EZ camera (Tuscon, AZ) connected to a Zeiss Axioscop 2

compound microscope (Jena, Germany). Larval body size data were analyzed in R [26] with a

generalized linear mixed model in the package lme4 [27], with pH as a fixed effect and culturing

vessel as a random effect. See the supplemental methods for expanded details.

DNA sequencing, Mapping, & SNP-calling:

DNA was extracted from pools of larvae for each day and replicate vessel using a Zymo

ZR-Duet DNA/RNA MiniPrep Plus Kit (Zymo, Irvine, CA). High quality DNA was shipped to

Rapid Genomics (Gainesville, FL) for library prep, capture, and sequencing. Following library

generation, DNA libraries were captured with 46,316 custom probes. Probes were designed to

capture two 120 base pair regions per gene: one within exon boundaries and one in putative

regulatory regions. Barcoded samples were then pooled and sequenced using 100 bp paired-end

sequencing on one lane of an Illumina HiSeq 3000.

Raw paired-end reads were quality trimmed and mapped to the *S. purpuratus* genome 3.1

(build 7) with bwa mem [28]. Variants were identified using *mpileup* in Samtools and filtered for

minor allele frequency (maf) of 0.01, quality greater than 20, bi-allelic SNPs only, and no

missing data. Sequencing depth can influence the accuracy of allele frequency estimation, and

we used only variants where each pool was sequenced to a depth of > 40x and with an average

minimum depth across all pools > 50x, as recommended by Schlotterer et al. [29]. Mean

maximum coverage cutoff was 372. Finally, we removed off target variants, which were defined

as any variant greater than 2kb from a probe. This filtering process resulted in 77,449 variant

sites. See supplemental methods for more details. Code for data processing and analyses can be

found on our GitHub page: https://github.com/PespeniLab/urchin_sel_ms_2018.

Detecting changes in allele frequency:

Cochran-Mantel-Haenszel (CMH) tests were conducted in R (*mantelhaen.test*) to identify

significant shifts in allele frequency in response to pH treatment. CMH tests are a standard

method of identifying changes in allele frequency in experimental evolution studies [10]. To

identify consistent changes in allele frequency through developmental time, we compared allele

frequency estimates for the four replicate samples from day 1 at pH 8.0 (T_0) to the four replicate

samples from day 7 at pH 8.0 and to the four replicate samples from day 7 at pH 7.5. P-values

were corrected using the *qvalue* package in R [30]. These data could be affected by long range

linkage disequilibrium [31]; therefore, we take a conservative approach and consider q-values <

0.001 significant.

A principal components analysis (PCA) was used to visualize relationships between

allele frequencies of treatment replicates using the R package *pcadapt* [32]. We calculate

nucleotide diversity for each treatment using a sliding window approach in Popoolation with the

variance-sliding command [33]. Window size was set to 400 bp with step size of 200 bp.

Distributions of nucleotide diversity were compared with a Kolmogorov-Smirnov test using the

ks.test function in R.

Following a selective sweep, sites under selection should show a pattern of higher linkage

disequilibrium (LD) than variants responding to drift alone. To test for this pattern, we compared

LD estimates among pairs of SNPs across the genome for selected (CMH $q < 0.001$) and neutral

(non-selected, CMH $q \geq 0.001$) variants using LDx [34] (supplemental methods). LDx uses a

maximum likelihood approach and leverages haplotype information of SNPs present on single

reads to estimate LD between pairs of variants. An exponential decay model was fit to each

group of LD estimates using the R package *nls* [35]. To assess whether the levels of LD present

in selected sites could be due to the lower number of variants than is present in neutral sites, we

randomly subsampled all variants 500 times to match the number of selected loci and compare

these decay curves to the observed decay in LD using a Kolmogorov-Smirnov test.

To assess the distribution of starting allele frequencies of adaptive loci, selected loci were

polarized by the frequency of the allele increasing in frequency in response to pH selection (i.e.,

the putative adaptive allele). Neutral loci ($q \geq 0.001$) were polarized in the same way, though this

is a random assignment given the lack of a significant shift in frequency. To ensure the patterns

observed were not a byproduct of the CMH statistic having the greatest power to identify

variants at low starting allele frequencies, the observed allele frequency distributions were

compared to permuted distributions with no true biological signal as described in the

supplemental methods.

Gene ontology (GO) enrichment was performed using the weight algorithm in topGO

version 2.22.0 [36]. GO terms for each gene were retrieved from EchinoBase

(http://www.echinobase.org/). Enrichment tests were conducted for each set of significant

variants and genes that had any SNPs in genic or intergenic regions were considered the target

set. Any gene with multiple significant variants was only considered once.

Figure 1: Morphometric analysis of total larval body length. (a) Boxplot of body size in each selection regime. Inset pictures show representative individuals found in pH 8.0 (b) and 7.5 (c).

Results

Morphometrics

Mean total larval body length was significantly lower in pH 7.5 (mean $\pm$ standard error: 239.1 $\pm$ 3.1 micrometers) as compared to pH 8.0 (320.9 $\pm$ 2.9 micrometers) (Fig. 1a, $P < 0.001$). Overall, larvae in the extreme low pH treatment were more stunted, with smaller bodies and spines compared to moderately low pH conditions (Fig. 1b-c). These results suggest that physiological and selective impact was stronger in pH 7.5 than pH 8.0. While we were unable to measure mortality with these data, estimates from other experiments in our lab show 30% survival in ambient (pH 8.1) conditions and 10% survival in extreme low (pH 7.5) conditions after 7 days.

Consistent allele frequency changes among replicate selection lines

We identified 75,368 variable sites present in or near 9,828 genes. 1,078 variants (in 816 genes) showed consistent changes in allele frequency (AF) in response to selection at pH 7.5 and 724 variants (in 579 genes) in response to pH 8.0 (CMH, $q < 0.001$; see Table S1 for a summary of results). 177 (11%) of these significant variants overlapped (in 144 genes, which is 11% of unique genes). However, assessing the overlap of genes that had loci targeted by selection, rather than overlap of specific variants, 205 genes overlapped (17% of unique genes). Overall, 2% of

Figure 2: Principal component analysis of allele frequencies for all 75,368 SNPs at time point zero (T_0) and after seven days in the two the pH treatments.

variants surveyed (1,625/77,449) and 12% of genes surveyed (1,200/9,828) were identified as

responsive to one or both pH treatments at q-value < 0.001.

Of the total assayed variants, 50,686 (67%) were in genic regions while 24,677 (33%)

were in intergenic regions. Matching expectations of chance sampling, for pH 7.5 selected loci,

725 (67%) were genic and 353 (33%) were intergenic (chi square, $P > 0.05$). Similarly, pH 8.0

selected loci consist of 506 (70%) genic and 218 (30%) intergenic loci (chi square, $P > 0.05$).

Given the rapid decay in linkage disequilibrium (see below), these results suggest that there are

both important coding and putative regulatory pH-responsive loci segregating in populations.

Principal component analysis (PCA) showed that the variance in allele frequencies

among larvae sampled from replicate culture vessels clustered by day and treatment (Fig. 2). pH

7.5 samples show the largest shift from the starting allele frequencies as expected with increased

selective mortality due to treatment. Note that one of the D7 pH 8.0 samples was an outlier. To

ensure that this sample did not artificially reduce power, we removed this sample and down

sampled all replicates to n=3, and reran the CMH test. We found the same relative numbers of

significant variants for both pH treatments.

Signals of shared and pH-specific selection

Patterns of LD among loci showed the highest LD between SNP pairs involving pH 7.5

loci, followed by pH 8.0 selected loci. Further, both sets of selected loci had higher LD than the

neutral expectation and putatively neutral sites matched genome wide expectations when

controlling for the number of variants sampled (Fig. 3 and Fig. S1). We observed rapid decay of

LD within 200 base pairs, which is expected given the high levels of genetic diversity, large

effective population size, high fecundity, and high gene flow of this species.

Figure 3: Decay in linkage disequilibrium with physical distance between SNP pairs. Dashed lines are neutral loci and solid red and blue lines represent decay in pH 7.5 and 8.0 selected variants, respectively. Solid grey shading is the 95% distribution of 500 permutations of down sampled random variants across the genome (see text for details).

Shifts in allele frequency showed unique and shared signals between pH selection

regimes (Fig. 4a, S2). Significant loci specific to pH 7.5 show a correlation in average allele

frequency change of 0.68 with their non-significant pH 8.0 counterparts ($P < 0.001$) suggesting

that the same loci were responding to treatment though to a lesser degree in pH 8.0; pH 8.0

significant loci reveal the same pattern with a correlation of 0.69 ($P < 0.001$). As expected, loci

that were identified as significant in both treatments showed the strongest correlation in allele

frequency change ($r^2 = 0.87$, $P < 0.001$). Interestingly, the loci with the most extreme shifts in

allele frequency were significant in both selection regimes (upper right quadrant of Fig. 4a, Fig.

S3).

To understand how variants are segregating and maintained in populations, we explored

the starting allele frequencies (T_0) for loci identified as responsive to pH 7.5, pH 8.0, and in both

treatments (Fig. 4b). All three sets of selected loci had significantly lower starting allele

frequencies than neutral loci (KS test, $P < 0.001$). Loci responsive to the most extreme selection

regime, pH 7.5, had significantly lower starting allele frequency than loci responsive pH 8.0 (KS

test, $P < 0.05$) and than loci responsive to both pH treatments (KS test, $P < 0.1$). Loci responsive

to just pH 8.0 and to both treatments had similar distributions of starting allele frequency (KS

test, $P = 0.38$). We also compared observed allele frequency distributions of selected loci to the

allele frequency distributions of the same number of randomly sampled loci across the genome

(Fig. 4B, S4). The allele frequency distribution for all three groups of selected variants were

significantly different from all 1000 randomly sampled distributions (KS tests, $P < 0.001$).

Lastly, we randomly shuffled sample IDs and compared starting allele frequencies of observed

and permuted "selected" loci. These permutations showed that the CMH statistic was not biased

towards loci that started at low frequencies (Fig. S5). These results suggest that loci underlying

low pH adaptation are rare in the starting population and more severe selection targets less

common variants. However, loci responsive to both pH treatments are more common than pH-

specific loci.

This overall pattern an excess of low allele frequency variants among selected loci is

consistent across functional classes of variants (non-synonymous, synonymous, intronic,

Figure 4: (a) Mean change in allele frequency for variants showing significant shifts under pH 7.5 and 8.0. Dashed black line represents the 1:1 expectation. (b) Allele frequency at T_0 polarized by the allele increasing in frequency during selection. Colored solid lines are the density plot distributions for each set of selected loci. The solid black line represents the median expectation of neutral loci from 1000 permutations of down sampling to the same number of variants as selected loci; grey shading is the 95% distribution of the permutation. Dashed vertical lines are the mean allele frequencies of each set of variants.

intergenic; Fig. S6). Low starting allele frequencies of responsive variants could suggest that

surviving offspring were from a single parent. However, this is unlikely to be the case due to the

rapid decay of linkage disequilibrium for selected loci relative to neutral loci. In addition, the

large number of responsive loci and genes suggests that survival in low pH conditions is a highly

polygenic trait, thus it is unlikely that surviving offspring had all the adaptive alleles from a

single parent.

Genetic diversity at T_0 was 0.0158 and 0.0129 and 0.0128 after seven days at pH 8.0 and

7.5, respectively. Both pH selection regimes resulted in a decrease in genetic diversity compared

to T_0 (KS test, $P < 0.001$). The decrease in diversity between the pH treatments was not different

(KS test, $P = 0.27$), demonstrating that, while the selection regimes were of different magnitude,

the relative loss of genetic diversity through time was similar.

Functional enrichment analysis revealed that genes under selection were enriched for

specific biological functions (Table S2). We observed enrichment for 19, 12, and 10 biological

processes GO terms for pH 7.5, pH 8.0, and overlapping variants, respectively. Of these, only

hexose metabolic process (GO:0019318) was significant across all three variant sets. Other

related terms shared across sets included those involved in mitotic damage control (GO:0044818;

GO:0007095; GO:0044774), cell growth (GO:0007099; GO:0030307; GO:0032467),

metabolism and energy production (GO:0001678; GO:0009267; GO:0019318; GO:0006096).

Discussion

We show that genetic response to moderate and extreme low pH relies on both shared

and unique mechanisms and that rare variants are important for survival in extreme low pH

conditions while common (intermediate frequency) variants are important for survival in both

moderate and extreme low pH regimes. Patterns of linkage disequilibrium and starting allele

frequencies suggest that both neutral and selective processes maintain adaptive variants in

natural populations. We further demonstrate the utility of single generation selection experiments

to identify the genetic basis of adaptation, which is particularly useful for testing capacity for

response to conditions that will be chronic in the future but are not so in nature today. Using

sequence capture of genomic DNA, we quantified shifts in allele frequency that represent

differential survival of genotypes during low pH selection in sea urchins. This approach has wide

potential application for identifying responsive genetic variants from wild populations. Our

results highlight that neutral standing genetic variation maintained with large population sizes

will be critical for survival in extreme environmental conditions.

Detecting adaptive loci from standing genetic variation

In the short term, a selective sweep results in decreased variation and increased LD, both

of which can be leveraged to identify adaptive genomic regions [37,38]. Our results reveal an

increase in LD in regions surrounding putatively adaptive loci, and the permutation shows that

this increase in LD is not a byproduct of our test statistic or sampling noise (Fig. 3, S1). This

provides confidence that our approach identifies genomic regions that are true targets of

selection. In addition, because the experimental design relies on selection on standing genetic

variation where loci will be present on multiple haplotypes, selective sweeps will retain ancestral

variation, soft sweeps, and signals are expected to be weaker than a hard sweep [18,39,40].

Typical evolve and resequence studies rely on multiple generations of selection to

identify adaptive shifts in allele frequency. However, a single generation selection can be

leveraged for organisms with high fecundity and small offspring, which is particularly useful for

long-lived organisms. This approach is unique and has benefits over standard experimental

evolution. First, one starts with outbred wild-caught individuals, which will increase starting

genetic diversity relative to inbred lab strains or isofemale lines [12,41,42]. Inference is limited

to the genetic diversity present in the starting individuals and may miss variation present at

different points in space or time and will miss the potential of new beneficial mutations. For

organisms with large effective population sizes and high mutation rates, new adaptive mutations

may be an important mode for adaptation [18]. However, starting with outbred parents

maximizes the amount of recombined genetic variation, avoids large linkage blocks, which can

plague experimental evolution studies from lab lines, and thereby improves the chances of

identifying selected loci. Moreover, generating a large amount of offspring from all crosses of

many parents creates a vast number of uniquely recombined genotypes upon which selection can

act. Finally, because offspring are small, it is possible to subject replicates of thousands of

individuals to selection, which is not possible with less fecund species with larger offspring.

The lack of genomic resources can limit the utility of a sequence capture approach for

non-model organisms. One solution is to utilize resources from closely related species.

Alternatively, it is possible to use a transcriptome rather than whole genome for probe design or

to utilize expressed exome capture [43]. While this does not allow for the design of probes in

promoter regions, it is typically available at a lower cost than whole genome sequencing.

Previous work in the purple sea urchin has used pooled RNA-sequencing data to identify

potentially adaptive genes and shifts in allele frequencies consistent among replicate pools within

pH treatments [13]. We find in the present study that allele frequencies estimated based on RNA

and genomic DNA from the same pooled larval samples are highly correlated ($r^2 = 0.93$, $P <$

0.001; Fig. S7), demonstrating that RNA-seq is a viable option for generating allele frequency

estimates.

Maintenance of adaptive standing genetic variation

Variants that respond to extreme low pH treatment are present in the starting population

(T_0) at low allele frequencies relative to neutral alleles (Fig. 4B). This pattern suggests that these

adaptive variants are not maintained by balancing selection or overdominance, but rather by one

of three processes. First, these alleles may be beneficial under extreme low pH conditions but

neutral or slightly deleterious at ambient conditions, known as conditional neutrality [44].

Alternatively, antagonistic pleiotropy can alter the rank fitness of alleles across different

environments, thus maintaining genetic variation across space or time [45]. Finally, purple sea

urchins are distributed along the west coast of North America from Alaska to Baja California,

Mexico [21,46] where pH conditions vary due to natural processes such as upwelling, and

populations have putative adaptive genetic variation associated with local pH conditions [22].

High gene flow results in extensive transfer of genetic variation across the geographic range and

migration-selection balance can result in the maintenance of low frequency alleles that are not

adaptive in a local environment [47]. Similar results have been observed in *D. melanogaster*

during experimental evolution to high temperature where allele frequencies of adaptive loci were

shifted towards low starting values [31]. Interestingly, this low shift was only observed for

adaptation to high but not low temperature, providing additional evidence that the CMH statistic

is not inherently biased towards low starting minor allele frequencies.

Interestingly, loci responsive to both moderate and extreme pH treatments had higher

starting allele frequencies than those responsive to extreme low pH alone (Fig. 4B). These "low-

pH-essential" alleles may be maintained at higher allele frequency due to balancing selection in

the wild. That is, the spatial and temporal heterogeneity of pH conditions experienced by purple

sea urchins across the species range and across their life history stages likely maintains these

alleles at intermediate frequencies through fluctuating selection or spatially balancing selection

[48]. We should note that these shared loci may also include variants involved in selection for

the general lab culture conditions. While the experimental design of this study precludes the

isolation of lab-adaptation variants, the concordance of functional classes of genes previously

identified as responsive to moderate low pH conditions (pH 7.8) in purple sea urchin larvae

generated from multiple populations [13,22] suggests that lab selection is unlikely to affect

general results, patterns, or conclusions in the present study.

Mechanisms for response to low pH conditions

Numerous biological processes related to maintaining homeostasis in low pH conditions

are overrepresented among selected loci. Across all sets of selected variants, we observe

enrichment in processes related to metabolism and energy production (GO:0001678,

GO:0009267, GO:0009061, GO:0019318, GO:0006096). Alterations to metabolic processes,

energy demands, and allocation are a primary response to low pH environments [49,50] and

previous *S. purpuratus* transcriptomic work has shown that this class of genes is differentially

regulated in response to low pH stress [15,51]. The production of calcium carbonate ($CaCO_3$)

generates excess protons that must be removed to maintain acid-base balance, and under acidic

conditions and under low pH conditions the energy required for acid-base regulation is increased

[50]. We also see shared enrichment for genes involved in cell proliferation and damage control,

including those implicated in cell growth and replication (GO:0030307, GO:0032467,

GO:0007099, GO:0006272), replication checkpoints (GO:0044818, GO:0044774, GO:0000076,

GO:0007095, and protein transport (GO:0046825, GO:0006606). These results match our

morphometric results of stunted larval growth and suggest that fine control of cellular growth

may be important for survival in low pH conditions. Two genes of interest that showed changes

in allele frequency greater than 20% in both pH treatments included ELMOD2 (SP-ELMOD2,

SPU_007564), the locus with the highest changes in allele frequency in both treatments, and

Focadhesin (KIAA1797, SPU_015184), a gene with three SNPs, two of which change amino

acids (upper right quadrant of Fig. 4a). ELMOD2 plays an important role in regulating

membrane traffic and secretion, phospholipid metabolism, and actin/cytoskeleton dynamics [52-

54] while Focadhesin is an important protein in subcellular structures as it integrates and receives

biomechanical and biochemical signals between the cytoskeleton and the extracellular matrix

[55,56].

In addition to shared responses, pH 7.5 selection, the edge of what these organisms

experience in nature, revealed unique adaptive targets to extreme pH selection. We observe

enrichment for 'actin polymerization or depolymerization' (GO:0008154). It has been

hypothesized that changes in actin abundance are related to cytoskeleton remodeling due to

intracellular stress during acclimation of oysters to climate change conditions [57,58]. Evans et

al., (2017) find enrichment for expression of genes involved in actin folding in *S. purpuratus*,

suggesting that the cytoskeleton is a target of pH stress. There is also enrichment for vacuolar

acidification (GO:0007035) which includes a H^+ transporting V-type ATPase (SPU_016993). V-

type ATPase regulate the pH of cellular compartments and can help maintain pH homeostasis

when extracellular pH is altered [59].

Finally, we identify TASK2 (SPU_003613) as under selection at pH 7.5 (Fig. 5). TASK2

is a pH sensitive K^+ transporter [60] and is an important component of bicarbonate (HCO_3^-)

uptake in mouse kidney [61]. In this mammalian system, HCO_3^- uptake occurs using a Na^+ co-

transporter where Na^+ gradients are maintained by Na^+ excretion through Na^+/K^+-ATPases,

resulting in the buildup of intracellular K^+. Alkalinzation from HCO_3^- activates TASK2 which

exports K^+ and re-establishes membrane polarization. HCO_3^- transport in urchin larvae is

similarly driven by a Na^+ co-transporter and Na^+ gradients are maintained by Na^+ excretion using

Na^+/K^+-ATPases [50]; it is possible that TASK2 is also involved in the urchin uptake process.

Under extreme pH conditions, the pH sensitivity of TASK2 may be under selection to ensure

continued HCO_3^- uptake, however, empirical work is needed to validate this hypothesized

mechanism.

The results of this analysis are sensitive to false negatives as the stringent cutoff of $q <$

0.001 may not identify loci that are truly under selection. For example, carbonic anhydrase

(SPU_012518) appears to be responding to pH 7.5 where 3/12 variants have *P*-values < 0.01,

two of which change amino acid sequence and are in linkage disequilibrium. This gene catalyzes

the hydration of CO_2 to bicarbonate [62] and is responsive to experimental acidification in many

organisms including purple sea urchin [25,51], corals [63], anemones [64], mussels [65], oysters

[66], and giant kelp [67], making it an ideal candidate for adaptation despite not meeting the

significance threshold. This protein, and potentially many others that are not statistically

significant, likely plays an important role in low pH adaptation and is worthy of future functional

investigations.

Figure 5: (a) Mean change in allele frequency of pH treatments relative to T_0 across a ~150 bp region of scaffold 542 that is ~8kb upstream of the K^+ transporter TASK2. (b) q-values corresponding to the allele frequency changes in (a). The dashed horizontal line indicates the significance threshold of 0.001.

Conclusion

We utilize a single generation selection experiment using an outbred, genetically diverse,

highly fecund species to reveal loci responding to extreme and moderate low pH selection. The

low variation among replicates, decreased decay of LD among selected loci, and enrichment for

biological functions related to pH adaptation suggest that this approach accurately identifies

adaptive loci. Further, we demonstrate how patterns of starting allele frequency can be used to

infer the mechanisms underlying maintenance of adaptive standing genetic variation. This work

provides a framework for future studies assessing the genetic basis of adaptive responses to

climate change. Increasing the temporal sampling through development would reveal how

different life stages differentially respond to selection, while functional studies on top candidates

will help to validate their mechanistic and evolutionary significance. Additionally, inter-

population studies will provide insight into how the adaptive potential may differ geographically

and be used to inform models to predict species level responses to climate change. Together, the

results presented here show that *S. purpuratus* possesses genetic variation that is responsive to

extreme low pH conditions and argues that rare genetic variation will be disproportionately

important for surviving future extreme conditions. Taken together, these results suggest an

important warning, given the low starting allele frequency and strength of selection, as evidenced

by the sweep patterns of linkage disequilibrium, selection by extreme events could result in a

major loss of standing genetic variation that could be important for response to other biotic or

abiotic stressors.

Data and code availability

Sequence data is available at the National Center for Biotechnology Information (NCBI SRA

BioProject: PRJNA479817). Phenotype data is available as part of the supplementary material.

All code for analysis is available at https://github.com/PespeniLab/urchin_sel_ms_2018.

Competing Interests

We have no competing interests.

Author Contributions

RB analyzed data and wrote the manuscript. AG designed and conducted the experiment and

assisted with data analysis and drafting the manuscript. KH helped perform the experiment and

analyzed the morphometric data. HH helped design aquaculture facilities and pilot the

experiment. MP designed the experiment and wrote the manuscript.

Acknowledgements

We thank Pete Halmay and Patrick Leahy for urchin collections. Jeremy Arenos for assistance

with imaging and image analysis, Jason Hodin and Justin McAlister for helpful discussions on

urchin larval culture, Lauren Ashlock for assistance with conducting the experiment, Mike

Austin and the Vermont Advanced Computing Core for server maintenance, and all members of

the Pespeni lab and Stephen Keller for helpful discussions.

Funding

This work was supported in part by the National Science Foundation (NSF) grant OCE-1559075

(to M.H.P.). AG was supported by the NSF Graduate Research Fellowship Program DGE-

1451866.

References

1.   Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N.,
Raven, P. H., Roberts, C. M. & Sexton, J. O. 2014 The biodiversity of species and
their rates of extinction, distribution, and protection. *Science* **344**, 1246752–
1246752. (doi:10.1126/science.1246752)

2.   Rahmstorf, S. & Coumou, D. 2011 Increase of extreme events in a warming world.
Proc. Natl. Acad. Sci. **108**, 17905–17909. (doi:10.1073/pnas.1101766108)

3.   Hoffmann, A. A. & Sgrò, C. M. 2011 Climate change and evolutionary adaptation.
Nature **470**, 479–485. (doi:10.1038/nature09670)

4.   Denny, M. W. & Dowd, W. W. 2012 Biophysics, environmental stochasticity, and
the evolution of thermal safety margins in intertidal limpets. *J. Exp. Biol.* **215**, 934–
947. (doi:10.1242/jeb.058958)

5.   Harley, C. D. G. & Paine, R. T. 2009 Contingencies and compounded rare
perturbations dictate sudden distributional shifts during periods of gradual climate
change. *Proc. Natl. Acad. Sci.* **106**, 11172–11176. (doi:10.1073/pnas.0904946106)

6.   Lande, R. 2009 Adaptation to an extraordinary environment by evolution of
phenotypic plasticity and genetic assimilation. *J. Evol. Biol.* **22**, 1435–1446.
(doi:10.1111/j.1420-9101.2009.01754.x)

7.   Grant, P. R., Grant, B. R., Huey, R. B., Johnson, M. T. J., Knoll, A. H. & Schmitt, J.
2017 Evolution caused by extreme events. *Philos. Trans. Royal Soc. B* **372**,
20160146. (doi:10.1098/rstb.2016.0146)

8.   Kofler, R. & Schlotterer, C. 2014 A guide for the design of evolve and resequencing
studies. *Mol. Biol. Evol.* **31**, 474–483. (doi:10.1093/molbev/mst221)

9.   Burghardt, L. T., Epstein, B., Guhlin, J., Nelson, M. S., Taylor, M. R., Young, N.
D., Sadowsky, M. J. & Tiffin, P. 2018 Select and resequence reveals relative fitness
of bacteria in symbiotic and free-living environments. *Proc. Natl. Acad. Sci.* **115**,
2425–2430. (doi:10.1073/pnas.1714246115)

10.   Schlotterer, C., Kofler, R., Versace, E., Tobler, R. & Franssen, S. U. 2015
Combining experimental evolution with next-generation sequencing: a powerful tool

to study adaptation from standing genetic variation. *Heredity* **114**, 431–440.
(doi:10.1038/hdy.2014.86)

11.   Franks, S. J. & Hoffmann, A. A. 2012 Genetics of climate change adaptation. *Annu.*
Rev. Genet. **46**, 185–208. (doi:10.1146/annurev-genet-110711-155511)

12.   Kessner, D. & Novembre, J. 2015 Power analysis of artificial selection experiments
using efficient whole genome simulation of quantitative traits. *Genetics* **199**, 991–
1005. (doi:10.1534/genetics.115.175075)

13.   Pespeni, M. H. et al. 2013 Evolutionary change during experimental ocean
acidification. *Proc. Natl. Acad. Sci.* **110**, 6937–6942.
(doi:10.1073/pnas.1220673110)

14.   Chan, F. et al. 2017 Persistent spatial structuring of coastal ocean acidification in the
California Current System. *Sci. Rep.* **7**, 2526. (doi:10.1038/s41598-017-02777-y)

15.   Evans, T. G., Pespeni, M. H., Hofmann, G. E., Palumbi, S. R. & Sanford, E. 2017
Transcriptomic responses to seawater acidification among sea urchin populations
inhabiting a natural pH mosaic. *Mol. Ecol.* **26**, 2257–2275. (doi:10.1111/mec.14038)

16.   Hofmann, G. E. et al. 2011 High-frequency dynamics of acean pH: A multi-
ecosystem comparison. *PLoS ONE* **6**. (doi:10.1371/journal.pone.0028983)

17.   Chevin, L.-M., Lande, R. & Mace, G. M. 2010 Adaptation, plasticity, and extinction
in a changing environment: towards a predictive theory. *PLoS Biol* **8**, e1000357.
(doi:10.1371/journal.pbio.1000357.s001)

18.   Messer, P. W. & Petrov, D. A. 2013 Population genomics of rapidad aptation by soft
selective sweeps. *Trends Ecol. Evol.*, 1–11. (doi:10.1016/j.tree.2013.08.003)

19.   Barrett, R. D. & Schluter, D. 2008 Adaptation from standing genetic variation.
Trends Ecol. Evol. **23**, 38–44. (doi:10.1016/j.tree.2007.09.008)

20.   Flowers, J. M., Schroeter, S. C. & Burton, R. S. 2002 The recruitment sweepstakes
has many winners: Genetic evidence from the sea urchin *Strongylocentrotus*
purpuratus. *Evolution* **56**, 1445–1453. (doi:10.1111/j.0014-3820.2002.tb01456.x)

21.   Pespeni, M. H. & Palumbi, S. R. 2013 Signals of selection in outlier loci in a widely
dispersing species across an environmental mosaic. *Mol. Ecol.* **22**, 3580–3597.
(doi:10.1111/mec.12337)

22.   Pespeni, M. H., Chan, F., Menge, B. A. & Palumbi, S. R. 2013 Signs of adaptation
to local pH conditions across an environmental mosaic in the california current
ecosystem. *Integrative and Comparative Biology* **53**, 857–870.
(doi:10.1093/icb/ict094)

23. Kelly, M. W., Padilla-Gamiño, J. L. & Hofmann, G. E. 2013 Natural variation and
the capacity to adapt to ocean acidification in the keystone sea urchin
Strongylocentrotus purpuratus. *Glob Change Biol* **19**, 2536–2546.
(doi:10.1111/gcb.12251)

24. Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D. & Hales, B. 2008
Evidence for Upwelling of Corrosive 'Acidified' Water onto the Continental Shelf.
Science **320**, 1490–1492. (doi:10.1126/science.1155676)

25. Evans, T. G., Chan, F., Menge, B. A. & Hofmann, G. E. 2013 Transcriptomic
responses to ocean acidification in larval sea urchins from a naturally variable pH
environment. *Mol. Ecol.* **22**, 1609–1625. (doi:10.1111/mec.12188)

26. R Core Team 2016 *R: A language and environment for statistical computing.*
Vienna, Austria: R Foundation for Statistical Computing.

27. Bates, D., Mächler, M., Bolker, B. & Walker, S. 2015 Fitting Linear Mixed-Effects
Models Using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

28. Li, H. 2013 Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. *arXiv preprint arXiv:1303.3997*

29. Schlötterer, C., Tobler, R., Kofler, R. & Nolte, V. 2014 Sequencing pools of
individuals — mining genome-wide polymorphism data without big funding. *Nature*
Publishing Group **15**, 749–763. (doi:10.1038/nrg3803)

30. Storey, J. D. 2002 A direct approach to false discovery rates. *Journal of the Royal*
Statistical Society: Series B (Statistical Methodology) **64**, 479–498.
(doi:10.1111/1467-9868.00346)

31. Tobler, R., Franssen, S. U., Kofler, R., Orozco-terWengel, P., Nolte, V., Hermisson,
J. & Schlötterer, C. 2014 Massive habitat-specific genomic response in *D.*
melanogaster populations during experimental evolution in hot and cold
environments. *Mol. Biol. Evol.* **31**, 364–375. (doi:10.1093/molbev/mst205)

32. Luu, K., Bazin, E. & Blum, M. G. B. 2017 pcadapt: an R package to perform
genome scans for selection based on principal component analysis. *Mol. Ecol.*
Resour. **17**, 67–77. (doi:10.1111/1755-0998.12592)

33. Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik,
A., Kosiol, C. & Schlötterer, C. 2011 PoPoolation: A toolbox for population genetic
analysis of next generation sequencing data from pooled individuals. *PLoS ONE* **6**,
e15925. (doi:10.1371/journal.pone.0015925)

34. Feder, A. F., Petrov, D. A. & Bergland, A. O. 2012 LDx: Estimation of linkage
disequilibrium from high-throughput pooled resequencing data. *PLoS ONE* **7**,
e48588. (doi:10.1371/journal.pone.0048588)

35.   Baty, F., Ritz, C., Charles, S., Brutsche, M., Flandrois, J.-P. & Delignette-Muller,
533   M.-L. 2015 A toolbox for nonlinear regression in R: the package nlstools. *J. Stat.*
534   *Softw.* **66**, 1–21. (doi:10.18637/jss.v066.i05)

535   36.   Alexa, A. & Rahnenfuhrer, J. 2010 topGO: enrichment analysis for gene ontology.
536   **11**, R14. (doi:10.1186/gb-2010-11-2-r14)

537   37.   Kim, Y. & Nielsen, R. 2004 Linkage disequilibrium as a signature of selective
538   sweeps. *Genetics* **167**, 1513–1524. (doi:10.1534/genetics.103.025387)

539   38.   Przeworski, M. 2002 The signature of positive selection at randomly chosen loci.
540   *Genetics* **160**, 1179–1189.

541   39.   Hermisson, J. & Pennings, P. S. 2017 Soft sweeps and beyond: understanding the
542   patterns and probabilities of selection footprints under rapid adaptation. *Methods*
543   *Ecol Evol* **8**, 700–716. (doi:10.1111/2041-210X.12808)

544   40.   Hermisson, J. & Pennings, P. S. 2005 Soft sweeps: Molecular population genetics of
545   adaptation from standing genetic variation. *Genetics* **169**, 2335–2352.
546   (doi:10.1534/genetics.104.036947)

547   41.   Turner, T. L. & Miller, P. M. 2012 Investigating natural variation in *Drosophila*
548   courtship song by the evolve and resequence approach. *Genetics* **191**, 633–642.
549   (doi:10.1534/genetics.112.139337)

550   42.   Orozco-terWengel, P., Kapun, M., Nolte, V., Kofler, R., Flatt, T. & Schlötterer, C.
551   2012 Adaptation of *Drosophila* to a novel laboratory environment reveals
552   temporally heterogeneous trajectories of selected alleles. *Mol. Ecol.* **21**, 4931–4941.
553   (doi:10.1111/j.1365-294X.2012.05673.x)

554   43.   Puritz, J. B. & Lotterhos, K. E. 2018 Expressed exome capture sequencing: A
555   method for cost-effective exome sequencing for all organisms. *Mol. Ecol. Resour.*
556   **166**, 1724. (doi:10.1111/1755-0998.12905)

557   44.   Anderson, J. T., Lee, C.-R., Rushworth, C. A., Colautti, R. I. & Mitchell-Olds, T.
558   2012 Genetic trade-offs and conditional neutrality contribute to local adaptation.
559   *Mol. Ecol.* **22**, 699–708. (doi:10.1111/j.1365-294X.2012.05522.x)

560   45.   Mitchell-Olds, T., Willis, J. H. & Goldstein, D. B. 2007 Which evolutionary
561   processes influence natural genetic variation for phenotypic traits? *Nat Rev Genet* **8**,
562   845–856. (doi:10.1038/nrg2207)

563   46.   Pespeni, M. H., Oliver, T. A., Manier, M. K. & Palumbi, S. R. 2010 Restriction site
564   tiling analysis: accurate discovery and quantitative genotyping of genome-wide
565   polymorphisms using nucleotide arrays. *Genome Biol* **11**, R44. (doi:10.1186/gb-
566   2010-11-4-r44)

47. Savolainen, O., Lascoux, M. & Merilä, J. 2013 Ecological genomics of local
adaptation. *Nature Publishing Group* **14**, 807–820. (doi:10.1038/nrg3522)

48. Charlesworth, D. 2006 Balancing selection and Its effects on sequences in nearby
genome regions. *PLoS Genet* **2**, e64–6. (doi:10.1371/journal.pgen.0020064)

49. Pan, T. C. F., Applebaum, S. L. & Manahan, D. T. 2015 Experimental ocean
acidification alters the allocation of metabolic energy. *Proc. Natl. Acad. Sci. U.S.A.*
112, 4696–4701. (doi:10.1073/pnas.1416967112)

50. Stumpp, M. et al. 2012 Acidified seawater impacts sea urchin larvae pH regulatory
systems relevant for calcification. *Proc. Natl. Acad. Sci.* **109**, 18192–18197.
(doi:10.1073/pnas.1209174109/)

51. Todgham, A. E. & Hofmann, G. E. 2009 Transcriptomic response of sea urchin
larvae Strongylocentrotus purpuratus to CO_2-driven seawater acidification. *J. Exp.*
Biol. **212**, 2579–2594. (doi:10.1242/jeb.032540)

52. Suzuki, M., Murakami, T., Cheng, J., Kano, H., Fukata, M. & Fujimoto, T. 2015
ELMOD2 is anchored to lipid droplets by palmitoylation and regulates adipocyte
triglyceride lipase recruitment. *Mol. Biol. Cell* **26**, 2333–2342.
(doi:10.1091/mbc.E14-11-1504)

53. East, M. P., Bowzard, J. B., Dacks, J. B. & Kahn, R. A. 2012 ELMO
domains,evolutionary and functional characterization of a novel GTPase-activating
protein (GAP) domain for Arf protein family GTPases. *J. Biol. Chem.* **287**, −39553.
(doi:10.1074/jbc.M112.417477)

54. Ivanova, A. A., East, M. P., Yi, S. L. & Kahn, R. A. 2014 Characterization of
recombinant ELMOD (cell engulfment and motility domain) proteins as GTPase-
activating proteins (GAPs) for ARF family GTPases. *J. Biol. Chem.* **289**, 11111–
11121. (doi:10.1074/jbc.M114.548529)

55. Geiger, B., Spatz, J. P. & Bershadsky, A. D. 2009 Environmental sensing through
focal adhesions. *Nat Rev Mol Cell Biol* **10**, 21–33. (doi:10.1038/nrm2593)

56. Burridge, K. & ChrzanowskaWodnicka, M. 1996 Focal adhesions, contractility, and
signaling. *Annu. Rev. Cell Dev. Biol.* **12**, 463–518.
(doi:10.1146/annurev.cellbio.12.1.463)

57. Tomanek, L. 2014 Proteomics to study adaptations in marine organisms to
environmental stress. *J Proteomics* **105**, 92–106. (doi:10.1016/j.jprot.2014.04.009)

58. Goncalves, P., Thompson, E. L. & Raftos, D. A. 2017 Contrasting impacts of ocean
acidification and warming on the molecular responses of CO_2-resilient oysters. 1–
15. (doi:10.1186/s12864-017-3818-z)

59.   Kane, P. M. 2006 The where, when, and how of organelle acidification by the yeast
vacuolar H^+-ATPase. *Microbiology and Molecular Biology Reviews* **70**, 177–191.
(doi:10.1128/MMBR.70.1.177-191.2006)

60.   Reyes, R., Duprat, F., Lesage, F., Fink, M., Salinas, M., Farman, N. & Lazdunski,
M. 1998 Cloning and expression of a novel pH-sensitive two pore domain K^+
channel from human kidney. *J. Biol. Chem.* **273**, 30863–30869.
(doi:10.1074/jbc.273.47.30863)

61.   Warth, R. et al. 2004 Proximal renal tubular acidosis in TASK2 K^+ channel-
deficient mice reveals a mechanism for stabilizing bicarbonate transport. *Proc. Natl.*
Acad. Sci. **101**, 8215–8220. (doi:10.1073/pnas.0400081101)

62.   Lindskog, S. 1997 Structure and mechanism of carbonic anhydrase. *Pharmacol.*
Ther. **74**, 1–20. (doi:10.1016/S0163-7258(96)00198-2)

63.   Zoccola, D., Innocenti, A., Bertucci, A., Tambutté, E., Supuran, C. & Tambutté, S.
2016 Coral carbonic anhydrases: Regulation by ocean acidification. *Marine Drugs*
14, 109–11. (doi:10.3390/md14060109)

64.   Ventura, P., Jarrold, M. D., Merle, P.-L., Barnay-Verdier, S., Zamoum, T., Rodolfo-
Metalpa, R., Calosi, P. & Furla, P. 2016 Resilience to ocean acidification: decreased
carbonic anhydrase activity in sea anemones under high pCO_2 conditions. *Mar.*
Ecol. Prog. Ser. **559**, 257–263. (doi:10.3354/meps11916)

65.   Fitzer, S. C., Phoenix, V. R., Cusack, M. & Kamenos, N. A. 2014 Ocean
acidification impacts mussel control on biomineralisation. *Sci. Rep.* **4**, 6218.
(doi:10.1038/srep06218)

66.   Wang, X., Wang, M., Jia, Z., Qiu, L., Wang, L., Zhang, A. & Song, L. 2017 A
carbonic anhydrase serves as an important acid-base regulator in pacific oyster
Crassostrea gigas exposed to elevated CO_2: implication for physiological responses
of mollusk to ocean acidification. *Mar Biotechnol* **19**, 22–35. (doi:10.1007/s10126-
017-9734-z)

67.   Fernández, P. A., Roleda, M. Y. & Hurd, C. L. 2015 Effects of ocean acidification
on the photosynthetic performance, carbonic anhydrase activity and growth of the
giant kelp *Macrocystis pyrifera*. *Photosynthesis Research*, 1–13.
(doi:10.1007/s11120-015-0138-5)

Figure legends:

Figure 1: Morphometric analysis of total larval body length. (a) Boxplot of body size in each

selection regime. Inset pictures show representative individuals found in pH 8.0 (b) and 7.5 (c).

Figure 2: Principal component analysis of allele frequencies for all 75,368 SNPs at time point

zero (T_0) and after seven days in the two the pH treatments.

Figure 3: Decay in linkage disequilibrium with physical distance between SNP pairs. Dashed

lines are neutral loci and solid red and blue lines represent decay in pH 7.5 and 8.0 selected

variants, respectively. Solid grey shading is the 95% distribution of 500 permutations of down

sampled random variants across the genome (see text for details).

Figure 4: (a) Mean change in allele frequency for variants showing significant shifts under pH

7.5 and 8.0. Dashed black line represents the 1:1 expectation. (b) Allele frequency at T_0

polarized by the allele increasing in frequency during selection. Colored solid lines are the

density plot distributions for each set of selected loci. The solid black line represents the median

expectation of neutral loci from 1000 permutations of down sampling to the same number of

variants as selected loci; grey shading is the 95% distribution of the permutation. Dashed vertical

lines are the mean allele frequencies of each set of variants.

Figure 5: (a) Mean change in allele frequency of pH treatments relative to T_0 across a ~150 bp

region of scaffold 542 that is ~8kb upstream of the K^+ transporter TASK2. (b) *q*-values

corresponding to the allele frequency changes in (a). The dashed horizontal line indicates the

significance threshold of 0.001.