Eco-evolutionary   processes   underlying   early   warning   signals   of

population declines

Gaurav Baruah[1], Christopher F. Clements[1,2], Arpat Ozgul[1]

[1] Department of Evolutionary Biology and Environmental studies, University of Zurich,

Winterthurerstrasse 30, 4055 Zurich

[2] School of Biosciences, University of Melbourne, Parkville VIC 3052, Australia

Melbourne

Corresponding author: Gaurav Baruah

Email: gaurav.baruah@ieu.uzh.ch

Abstract

Environmental change can impact the stability of a population and cause a rapid decline

in abundance. Abundance-based early warning signals have been proposed to predict

such declines, but these have been shown to have limited success, leading to the

development of warning signals based on the distribution of fitness-related phenotypic

traits such as body size. The dynamics of such traits in response to external

environmental perturbations are controlled by a range of underlying factors such as

reproductive rate, genetic variation, and plasticity. However, it remains unknown how

such ecological and evolutionary factors affect the stability landscape of populations and

the detectability of abundance and trait-based warning signals of population decline.

Here, we apply a trait-based demographic approach and investigate both trait and

population dynamics in response to gradual changes in the environment. We explore a

range of ecological and evolutionary constraints under which the stability of a

population may be affected. We show both analytically and with model-based

simulations that strength of abundance-based early warning signals is significantly

affected by ecological and evolutionary factors. Finally, we show that a unified approach,

combining trait- and abundance-based information, significantly improves our ability to

predict population declines.

1. Introduction

Projected anthropogenic influences on earth systems will continue to increase over the next 100 years and will significantly change the earth's biota (Mora et al. 2013). The impact of anthropogenic forcing can often bring natural populations and complex ecosystems into the region of tipping points (Wissel 1984; Scheffer et al. 2001; Scheffer et al. 2012), This point represents the threshold at which any slight increase in pressure or stochastic fluctuation can cause a transition from a stable state to an alternative state – exemplified by the collapse of populations (Frank et al. 2011) or change in ecosystem functions (Jackson et al. 2001; Frank et al. 2011). Predicting such transitions are thus critical if we are to mitigate the effects of human pressures and minimize the loss of biodiversity (Cardillo et al. 2005; Thomas et al. 2004). These transitions are however difficult to predict (Clements et al. 2015; Boettiger & Hastings 2013; Clements & Ozgul 2018).

One suite of methods, which may achieve this, are early warning signals derived from dynamic systems theory (van Nes & Scheffer 2007; M. Scheffer et al. 2009). Such methods assume that a dynamical system, for example a biological population, that is under increasing continuous external pressure, will show a property called critical slowing down (CSD) as it approaches a tipping point, whereby the system takes longer to return to its original state after a perturbation (Wissel 1984; Dakos et al. 2008; Scheffer et al. 2009; Strogatz 1994). The occurrence of CSD is a direct implication of the dominant eigenvalue of the system approaching zero (for continuous dynamical system) (van Nes & Scheffer 2007) or 1 (for a discrete dynamical system) (Krko ek & Drake 2014). As a direct consequence of CSD, statistical moments embedded within abundance time-series data, particularly variance and autocorrelation will show marked increase over time. These two indicators are the prominent abundance-based statistical metrics, and, along with various other metrics (Dakos et al. 2012; C. Boettiger & Hastings 2012;

74   Clements & Ozgul 2016; Clements & Ozgul 2018; Drake & Griffen 2010), are commonly

75   known as early warning signals (EWS) or leading indicators.

76   Changes in autocorrelation and variance are measures of the stability of a

system, derived from theory of alternative stable states (Wissel 1984; Clements & Ozgul

2018). Holling's (1973) fundamental work on stability and resilience laid the foundation

for future work on deriving stability criteria for systems with alternative equilibrium

states. Stability of a dynamical system can be characterized by the classic ball-in-cup

diagram (Holling 1973; Nolting & Abbott 2015; Clements & Ozgul 2018), where the 'ball'

represents the state of the system and the 'cup' represents the potential energy

landscape. The stability of the system is thus quantified by the depth of the energy

landscape, and resilience by the size of the cup (Villa Martín et al. 2015; Dai et al. 2015;

Clements & Ozgul 2018). In response to continuous exogenous pressure, the system

loses resilience (for example, decline in population numbers) and the energy landscape

shrinks. Concurrent with the shrinkage of the landscape, there is also a decrease in the

depth of the 'ball-in-cup' landscape signaling the loss in stability. It is this loss in

stability, which statistical EWSs such as variance and autocorrelation measure as an

alternative for loss of global resilience of the system (Clements and Ozgul 2018). The

shape of the ball-in-cup landscape thus determines the dynamics of the state variable in

response to an exogenous environmental change (Beisner et al. 2003).

However, the shape of the 'ball-in-cup' landscape can be changed not only by exogenous

pressures, but can also be altered by intrinsic factors associated with the system's

dynamics (Beisner et al. 2003). Changes in these intrinsic factors might alter the location

of an equilibrium state, or destabilize the current state, causing the system to arrive at a

point closer to the other alternative state (see Fig. 1). For a biological population, the

state of the population will be characterized by its current abundance. Responses of this

state variable (abundance) to an exogenous environmental pressure will be a

combination of ecological and evolutionary responses controlled by factors such as

genetic variation or plasticity (Price et al. 2003; Gienapp et al. 2008; Ozgul et al. 2010).

For example, plastic responses to fluctuating changes in the exogenous environment are

usually fast which consequently stabilizes population fluctuations (Reed et al. 2010).

However, if the environment keeps on changing, the population might not only deplete

its plastic capacity but also its standing genetic variation, which in turn leads to its

eventual decline. Further, such a decline would also be conditional on the population's

reproductive rate, with higher reproductive rate leading to a slower decline (Juan-Jordá

et al. 2015). In addition, due to continuous changes in the exogenous environment,

selection pressures can subsequently increase leading to contemporary evolution

(Yoshida et al. 2003). Adaptive evolution would then depend on the amount of genetic

variation present in the population with higher variation leading to faster evolution,

which can either stabilize or destabilize population dynamics (Sanchez & Gore 2013;

Cortez 2016). Since the dynamics of a population in response to an exogenous pressure

is determined by the shape of its potential energy landscape (Beisner et al. 2003), it is

thus highly probable that changes in these factors namely plasticity, genetic variation,

and reproductive rate have the capacity to influence the shape of the ball-in-cup

landscape. In turn, given that EWS are stability measures derived from alternative stable

states theory (Beisner et al. 2003; Scheffer et al. 2009; Clements & Ozgul 2018), it is very

likely that strength in EWS and hence predictability of population decline might also be

affected by these factors.

The EWSs described above are phenomenological signals which require very few

assumptions about the mechanistic processes driving system dynamics, which makes

them applicable to a wide variety of systems (Wissel 1984; Scheffer et al. 2009;

Clements et al. 2017; Clements and Ozgul 2018; Jarvis et al. 2016; Dutta, Sharma, and

Abbott 2018). However, such statistical tools have a multitude of issues associated with

it, for example- low signal to noise ratio and the requirement of high-quality time-series

data (Clements et al. 2015). This had led to the development of alternatives like

maximum likelihood-based (Boettiger & Hastings 2012; Brock & Carpenter 2012) and

generalized system-specific modelling approaches (Lade & Gross 2012). However, for

many natural and ecological systems specific models might not be easily available, and a

generic EWS approach then becomes necessary.

Abundance-based measures of the stability of an ecological system might not be the only

indicators of a population being forced by an external environment (Clements & Ozgul

2016) . Phenotypic traits, such as body size, is one of the important traits that

determines the fate of individuals as well as dynamics of population, and thus can

provide crucial information regarding the current state of the population (Clements and

Ozgul 2016b; Ozgul et al. 2014; Spanbauer et al. 2016). Recently, Clements & Ozgul

(2016) showed, using data from experimental microcosms, that including body size

information with EWS could significantly improve the predictive accuracy of population

collapse. Such trait-based signals, have been further shown to be present in the historic

collapses of whale populations (Clements et al. 2017). However, extensive research on

the efficacy of such signals in comparison to abundance-based EWS is still in its infancy

(Clements & Ozgul 2018).

In this paper, we used a quantitative genetics framework to simulate the dynamics of a

population in response to gradual environmental change. We track both the population

dynamics and trait distribution until the simulated population collapses. We show that

this particular model exhibits critical slowing down behavior and hence is expected to

show EWS (van Nes and Scheffer 2007). We then assess whether genetic variation,

plasticity in the trait, and net reproductive rate alter the ball-in-cup landscape (stability

landscape) and consequently the strength in EWS. In addition, we assess the efficacy of

trait-based EWS in comparison with abundance-based EWS and reiterate the ability of

trait-based signals in accurately informing population collapses.

2. Materials and Methods

2.1 Mathematical model:

We model the joint dynamics of quantitative trait evolution and population size under

density dependent regulation via the effects of selection (Gomulkiewicz & Holt 1995;

Chevin et al. 2010; Chevin & Lande 2010) . We consider a population that has discrete

generations where individuals have fitness that is determined by a single quantitative

trait z under stabilizing selection with linear reaction norms (Gavrilets & Scheiner

1993). The phenotype of an individual *i* in this finite population before selection is given

by:

$$z_{i,t} = a_{i,t} + bu_t + \epsilon \qquad (1)$$

where a is the breeding value of the individual i. The breeding value is normally

distributed with mean $\overline{a}$ and additive genetic variance σ_a^2, and ϵ is the residual

component with mean 0 and variance σ_ϵ^2. The slope b in our model is constant meaning

that plasticity in the trait cannot evolve. We model an environment that determines the

optimal phenotypic value θ_t for the primary phenotype z_t and also cues the plastic

response. The optimal phenotype. θ_t, is assumed to be linearly dependent on the

environment E that selects for a particular phenotypic value such that, $\theta_t = BE_t$.

We included stochasticity in the environment E_t as a Gaussian white noise process. The

optimal phenotype hence fluctuates as E_t fluctuates over a mean value. As in Reed et al.

(2010), the environmental cue u and the optimal phenotype θ are drawn from a

bivariate normal distribution at each generation t with equal variances. The covariance

of u and θ is given by $\sigma_{\theta,u} = \xi \sigma_\theta \sigma_u$ where ξ is the correlation between θ and the cue *u.*

The environmental cue u is drawn from the distribution with its mean value U_t over

time given by:

$$U_t = \begin{cases} 0 & , t < t_{500} \\ 1 - c\frac{t}{500} & , \ t \geq t_{500} \end{cases} \qquad (2)$$

And for optimal phenotype θ is drawn from the bivariate distribution with mean value

given by:

$$\theta_t = BE_t = \begin{cases} 0 & , t < t_{500} \\ B - B\dfrac{t}{500} & , \ t \geq t_{500} \end{cases} \qquad (3)$$

We assumed that variances in the cue and the optimum are equal. This mimics the

situation, for instance, when variation in food availability is correlated with inter annual

variation in snow melt (Vuren & Armitage 1991). ξ determines predictability of the

environment such that $\xi = 1$ means the environmental cue is correlated with the

optimum phenotype in a way that the plastic response tracks the optimum environment

perfectly. In such a case, the strength in plastic response of the trait will be determined

by b.

The mean phenotype at any generation t is then: $\bar{z} = \bar{a} + bu_t$ and has variance:

$\sigma_z^2 = \sigma_a^2 + \sigma_\epsilon$ (We consider $\sigma_\epsilon \approx 0, \sigma_z^2 = \sigma_a^2$). Without loss of generality, at the

reference environment, i.e., $u_t = 0$, the phenotype is assumed to be at its optimum

value. Each individual in this population has a net reproductive fitness of R_0 that is under

density-dependent regulation. The primary trait z determines its reproductive fitness in

the sense that:

$$R(z) = R_0^{\left(1 - \frac{f(N_t)}{f(K)}\right)} W(z) \qquad (4)$$

where, $\dfrac{f(N_t)}{f(K)}$ is a density-dependent function and $W(z)$ is the Gaussian stabilizing

fitness function :

$$W(z, \theta) = e^{-(z-\theta)^2/2\sigma_w^2} \qquad (5)$$

If the primary trait z is far from its optimum phenotype θ then its fitness will decline

depending on the width σ_w^2 of the fitness function $W(z, \theta)$, which signifies the strength

of selection. Assuming that the phenotypic distribution to be normal *p(z)*, the average

reproductive fitness of the population will be

204
$$\bar{R} = R_0^{\left(1-\frac{f(N_t)}{f(K)}\right)} \int W(z)p(z)dz \qquad (6)$$

205   If time *t* is in generations, then population size at *t+1* is

206
$$N_{t+1} = N_t \bar{R} \qquad (7)$$

207   And the dynamics of mean phenotypic value is:

208
$$\bar{z}_{t+1} = \bar{a}_t + \sigma_a^2 \frac{\partial ln \overline{W}}{\partial \bar{a}} + bu_t \qquad (8)$$

where, $\overline{W} = \int W(z)p(z)dz = \sqrt{\frac{\sigma_w^2}{\sigma_z^2+\sigma_w^2}}\ e^{-(\bar{z}_t-\theta_t)^2/2(\sigma_w^2+\sigma_z^2)}$.

The above model is based on the infinitesimal quantitative genetic model (Falconer &

Mackay 1996).

2.2 Stability landscape and effective potential:

One approach to quantifying stability which is also fundamental to dynamical systems

with alternative stable states is the 'ball-in-cup' analogy (Haydon 2003; Villa Martín et

al. 2015). The different states of the dynamical system can be represented by a

landscape and the state of the population at a particular moment by a ball in that

landscape. The landscape of the dynamical system can be represented by an effective

potential $V(N)$ (Villa Martín et al. 2015; Beisner et al. 2003). If the dynamical system

can be represented by $\frac{dn}{dt} = g(n)$, then $V(n) = -\int g(n)dn$. For our dynamical system,

$g(n) = \log\left(R_0^{\left(1-\frac{n}{k}+\beta-\gamma\right)}\right)$. Here *n* is the state of the system, which in our case is the

population size. The effective potential of our dynamical system is then (supplementary

4):

$$V(n) = -\left(rn - \frac{\frac{r}{k}n^2}{2} + \beta rn - \gamma rn \right)$$

$$= \left(rn - \frac{\frac{r}{k}n^2}{2} + \frac{\log\left(\frac{\sqrt{(\sigma_W^2)}}{\sqrt{(\sigma_W^2 + \sigma_Z^2)}} \right)}{\log(R_0)} rn - \frac{(\bar{a} + bu_t - \theta_t)^2}{(R_0(\sigma_W^2 + \sigma_Z^2))} rn \right) \qquad (9)$$

It is evident from the above equation (9) that the effective potential function is

dependent on three parameters: r, β, γ, where $r = \log(R_0)$; $\beta = \dfrac{\log\left(\frac{\sqrt{(\sigma_W^2)}}{\sqrt{(\sigma_W^2 + \sigma_Z^2)}} \right)}{\log(R_0)}$;

$\gamma = \dfrac{(\bar{z} - \theta_t)^2}{(R_0(\sigma_W^2 + \sigma_Z^2))}$; $n = \log(N)$; $k = \log(K)$.

2.3 Equilibria and eigenvalues of the model:

The extinction equilibrium for the model is at $N^* = 0$ and the positive equilibrium for the

model is at $N^* = K + \beta K - \gamma K$, where $\gamma = \dfrac{(\bar{z} - \theta_t)^2}{(R_0(\sigma_W^2 + \sigma_Z^2))}$; $\beta = \log\left(\frac{\sqrt{(\sigma_W^2)}}{\sqrt{(\sigma_W^2 + \sigma_Z^2)}} \right) / \log(R_0)$

(see supplementary 1). Stability of the two equilibria are based on their corresponding

eigenvalues, $\lambda = \frac{\partial}{\partial N} f(N)|_{N=N^*}$. Equilibria will be stable if magnitude of $\lambda < 1$ and

unstable if magnitude of $\lambda > 1$ (Supplementary 1). Local stability analysis gives for the

extinction equilibrium N* = 0, $\lambda_1 = R_0^{1+\beta-\gamma}$ and for positive equilibrium $N^* = K + \beta K -$

γK, $\lambda_2 = 1 - (1 + \beta - \gamma)\log R_0$. Thus, the extinction equilibrium N* = 0, is always

unstable since $\lambda_1 = R_0^{1+\beta-\gamma} > 1$ whenever $R_0 > 1$. On the other hand, the positive

equilibrium $N^* = K + \beta K - \gamma K$ is stable conditional on the forcing parameter $\gamma =$

$\dfrac{(\bar{z} - \theta_t)^2}{(R_0(\sigma_W^2 + \sigma_Z^2))}$ and $R_0 > 1$. Assuming that the phenotype is at its optimum $(\bar{z} = \theta)$ i.e.,

$\gamma = 0$, then for $R_0 > 1$ the positive equilibrium is stable due to $\lambda_2 = 1 - (1 + \beta -$

$\gamma)\log R_0 < 1$. But as the forcing parameter $\gamma = \dfrac{(\bar{z} - \theta_t)^2}{(R_0(\sigma_W^2 + \sigma_Z^2))}$ increases over time (due to

θ_t shifting and the phenotype lagging behind), λ_2 slowly approaches 1 and the positive

equilibrium which is $N^* = K + \beta K - \gamma K$ is initially stable but becomes unstable once λ_2

reaches 1. This happens when increases in the forcing parameter γ or increases in

maladaptation causes average population growth rate $\bar{R}$ (4) also to reach 1. Once this

happens, the extinction equilibrium then becomes stable at $\bar{R} = 1$ and the two

equilibrium exchange stability at $\bar{R} = 1$. This is the definition of transcritical bifurcation.

Moreover, as $\bar{R}$ approaches 1, the magnitude of λ_2 becomes larger and the recovery rate

from external perturbation becomes slower (see supplementary 1 and 2). This indicates

that there is CSD. λ_2 is directly related to return time, from which indicators like

autocorrelation and standard deviation can be calculated (supplementary 2).

2.4 Simulations of population collapse

To complement our theoretical findings we did simulations and forced the populations

to collapse by increasing the rate of change in the environment E_t. We chose a rate of

change in the environment θ_t in a way that populations collapsed regardless of any

combination of parameter values. This was done by setting B =60 in (3) in all of our

simulations. Environmental change was introduced at time point t = 500, after

populations have reached their carrying capacity. In all of our simulations,

environmental predictability was high at $\xi = 1$. For each parameter value of a factor

(genetic variation, plasticity, net reproductive rate) we did 200 independent simulations

of population collapse. For each population time series, we first estimated the

bifurcation point by fitting GAMS to $\frac{N_{t+1}}{N_t}$ over time t. $\frac{N_{t+1}}{N_t} < 1$ is then our estimated

bifurcation point in our simulations.

2.4.1 Abundance-based EWS

For abundance-based EWS analyses, the length of the time-series used was of the same

length for each factor in question. For each parameter value of each factor in question,

we used time-series data to the estimated bifurcation point. We discarded time-series

data after the bifurcation point. The bifurcation point in the time-series however varied

due to different levels of parameter values for a factor in question, resulting in a range of

abundance time-series lengths (from as low as 30 to 110 time points from t = 500 where

the optimum environments shifts). Hence due to the possible effect of length of time-

series on strength of EWS of population collapse, we standardized the time-series length

such that each population time-series analyzed for EWS had the same length. We then

applied generic EWS of population collapse by using the *earlywarnings* package in R. We

focused our analysis on two specific indicators namely autocorrelation at the first-lag

(AR1) and standard deviation (SD) as most of the other indicators – such as first-order

autoregressive coefficient or coefficient of variation – could be theoretically derived

from these parameters (Dakos et al. 2012). The two indicators AR1 and SD were

calculated using a predefined sliding window, typically 50% of the abundance time-

series being analyzed. We normalized the leading indicators so that it would be easier

to incorporate information from phenotypic traits later in our analyses. We used

Gaussian detrending to remove any trend in the abundance time-series and used

Kendall's Tau correlation coefficients of the indicators prior to the estimated bifurcation

point as an indicator of an approaching collapse. Strong positive correlations of the

statistical indicators (SD and AR1) with time would indicate an approaching population

collapse which would be evident in a higher value of Kendall's Tau coefficient (Dakos et

al. 2012).

2.4.2 Trait-based EWS

To evaluate the efficacy of trait-based EWS (Clements & Ozgul 2016), information from

mean phenotypic trait was incorporated with the leading indicators in an additive

manner. To include information from average phenotypic value, we first normalized the

phenotypic time-series. The length of the phenotypic time-series that was used was the

same as the corresponding abundance time-series. Standardized EWS such as standard

deviation and auto-correlation (SD, AR1) were added additively with standardized mean

phenotypic time series to quantitatively create trait-based signals. We evaluated three

trait-based EWS metrics namely *AR1+trait, AR1+SD+trait, SD+trait* and compared how

the metrics that included information from the mean phenotype improved predictability

of population collapse when compared with the standardized abundance-based EWS

like *AR1, SD, AR1+SD*. We quantified Kendall's Tau correlation coefficient as an indicator

for strength of predictability of population collapse.

2.4.3 Adaptive plasticity: To assess the role of adaptive plasticity b, we simulated 200

replicate population collapses for each parameter value of b, which ranged from low to

high: 0.02, 0.06, 0.1, 0.2, 0.4, and 0.6. While we varied plasticity, we fixed $\sigma_a^2 = 0.2$, and

R_0 at 1.2. The width of the fitness function determines the strength of selection in the

trait. We fixed the width of the fitness function at $\sigma_w^2 = 25$, which would indicate

medium strength in selection. Usually, $\sigma_w^2 = 50$ would indicate weak strength in

selection and $\sigma_w^2 = 5$ would indicate strong selection (Chevin and Lande 2010).

2.4.4 Genetic variation: The levels of genetic variation σ_a^2 used for the simulations of

population collapse are 0.02, 0.06, 0.1, 0.2,0.3,0.5, 0.6. Adaptive plasticity b, net

reproductive rate R_0 and σ_w^2 were kept at 0.2, 1.2 and 25 respectively.

We also did simulations where we relaxed the assumption of genetic variation (σ_a^2)

remaining constant during population collapses. For details see supplementary 5.

2.4.5 Net reproductive rate: To assess the role R_0, we simulated 200 replicate population

collapses. The levels of R_0 used for the simulations of population collapse were 1.1, 1.2,

1.3, 1.4, and 1.5. Adaptive plasticity b, genetic variation σ_a^2 and σ_w^2, were kept at 0.2, 0.2

and 25 respectively.

3. Results

3.1 Effective potential

3.1.1 Strength in plasticity

The term $\gamma = \frac{(\bar{z}-\theta_t)^2}{(R_0(\sigma_w^2+\sigma_z^2))}$ in equation (9) is the evolutionary lag due to maladaptation of

the mean phenotype with the optimum (Lande 2009). The effective potential becomes

narrower with increasing strength in plasticity and consequentially increases stability,

i.e., the potential function becomes more negative (Fig. 2). Since return time and

stability are inversely related (Dai et al. 2015), return time decreases with increasing

strength in plasticity (Supplementary 2, Fig. S1). In other words, the population quickly

recovers with every perturbation if strength in plasticity is high.

3.1.2 Genetic variation

The mean phenotype has variance of $\sigma_z^2 = \sigma_a^2 + \sigma_\epsilon$. Assuming residual component σ_ϵ

to be zero, phenotypic variance is then equal to additive genetic variance, $\sigma_z^2 = \sigma_a^2$.

Inspection of the equation (9) reveals that high additive genetic variance causes the

effective potential to be more positive when compared with low σ_a^2 and consequently

decreases stability, i.e., the effective potential becomes less negative (Fig. 2; the more

negative the effective potential, more stable the population is). Loss of stability and

effective potential energy is higher for populations with low genetic variance whenever

they are forced to collapse compared to populations with high genetic variance (Fig. 2,

supplementary 3). Further, return time also increases with increasing additive genetic

variance (supplementary 2, Fig. S1).

3.1.3 Net reproductive rate

From equation (9), we also see that higher values of R_0 lead to higher stability and

narrow potential curve for a given state of the population (Fig. 2). Loss in stability is

more when R_0 is higher for the same rate of change in the environmental change when

compared with a lower R_0. Further, return time also decreases with increasing R_0

(supplementary 2, Fig. S1).

3.2 Simulations of population collapse: abundance-based EWS

3.2.1 Strength in plasticity

The strength of EWSs of population collapse (Kendall's Tau value) was much higher in

populations which had high adaptive plasticity *b* = 0.6 (Fig. 3a, d) when compared with

populations with low adaptive plasticity.

3.2.2 Genetic variation

Our model simulations showed that genetic variation altered the detectability of

population collapse for the same rate of environmental change (Fig. 3b, 3e). Higher

genetic variation in the population led to slight decrease in detectability of population

collapse than in populations with lower genetic variation.

3.2.3 Net reproductive rate

Strength in EWS of populations collapse was significantly altered by net reproductive

rate. Strength in EWS was lower in populations with lower R_0 when compared with

higher R_0 (Fig. 3c, 3f). Both the leading indicators SD and AR1 showed similar behavior

in this aspect (Fig. 3c, 3f), but Kendall's tau value of SD showed significant increases in

strength as net reproductive rate increased.

3.3 Simulations of population collapse: trait-based EWS

Including information from trait value alongside abundance-based EWS significantly

improved the strength of EWS of population collapse (Fig. 4, Fig S8-S9). In principle, *AR1*

was the least efficient of all the signals compared. However, when trait information was

included with *AR1*, strength in trait-based *AR1 (trait+AR1)* improved significantly

(Fig.4). Overall, the Kendall's Tau value of *SD+trait* was relatively higher than all the

other trait-based and abundance-based EWS. The extent of the distribution of Kendall's

tau values of *SD+trait* signal was much lower than that of just *SD* (Fig. 4). This was

consistent across all the factors analyzed such as genetic variation, plasticity (Fig.4, Fig.

S8-S9).

4. Discussion

The early detection of population collapse with the help of EWS provides a unique tool to highlight conservation efforts to a population approaching an impending transition. These tools are relatively simple, easy to use, as they only require state-based data like abundance and hence are applicable to a variety of ecological systems. However, we show here that factors intrinsic to a population in question such as genetic variation, plasticity or net reproductive rate could significantly alter stability of a population as well as strength in EWS of population decline.

The detectability of EWSs is dependent not only on the relation between stability and resilience but is also determined by a variety of factors including sampling frequency, and the strength and type of external forcing (Dai, Korolev, and Gore 2015; Clements et al. 2015a; Clements and Ozgul 2016). Here we add to this body of knowledge by demonstrating that the effective potential of a system can also be altered by intrinsic factors such as genetic variation, plasticity and net reproductive rate. These changes to the effective potential in turn led to stronger EWS being observed, when changes in the external optimum caused a population to lose significant amount of potential energy. In other words, stronger EWSs would be observed whenever a population is forced to move from a deeper point in the potential energy curve (more stable) to its tipping point compared to another population, which is lying on a shallower potential (less stable). Our findings also suggested that the utility of EWS was strongly determined by factors that are intrinsic to a population such as genetic variation, net reproductive rate and plasticity.

Higher strength in plasticity in the phenotype significantly affected the detectability of population collapse (Fig. 2). Strength in plasticity altered the response and the dynamics of a population to changes in the external environment. When plasticity was high the response of the phenotype tracked the environment perfectly causing the population to

stabilize in response to fluctuating changes in the environment (Fig. S3) (Reed et al.

2011; Charmantier et al. 2008). The stability of a population is also reflected in short

return time to equilibrium (van Nes & Scheffer 2007). This was true for the case where

the strength in plasticity was high (Fig. S1). Moreover, the basin of attraction for a highly

plastic population was deeper when compared to a less plastic population, which was

shallower. To push a highly plastic population from the deepest point in its basin of

attraction to the tipping point, the energy loss would be more, compared to a less plastic

population which was lying in a much shallower basin. The loss in energy from driving

the highly plastic population from the deepest point to its tipping point was reflected in

stronger EWS when one compared it with another less plastic population. Consequently,

for the same rate of change in the environment, differences in Kendall's Tau values for

different values of plasticity was observed, with higher values of plasticity leading to

stronger EWS of population collapse.

Similarly, the effective potential of our dynamical system was also altered by genetic

variation (Fig. 2), with higher genetic variation leading to a shallower stability landscape

than that of a population with lower genetic variation. High additive genetic variance

created what was called a 'genetic load' that lowered the overall population growth rate

(Lande 1976; Lande and Shannon 1996; Chevin, Lande, and Mace 2010). This was due to

stabilizing selection acting on the mean phenotype. Due to this genetic load caused by

high additive genetic variance, population growth rate at equilibrium was lower causing

the effective potential to be closer towards the tipping point compared to another

population having a lower genetic variation. Hence for the same rate of external

environmental change, the loss of potential was different when one compared a

population with high genetic variation to that with low genetic variation. As a result, the

performance of EWS was affected with lower detectability of population collapse when

there was higher genetic variation (Fig. 3). This result however did not mean that

genetic variation was detrimental for long-term population persistence. Time to

extinction was however longer for populations with higher genetic variation when

compared with a population which had a lower genetic variation in the trait (Fig

S2)(Robinson et al. 2013; Yvonne & Hoffmann 2009). High genetic variation is necessary

for rapid adaptation via genetic evolution, which then aids in evolutionary rescue

(Chevin & Lande 2010). However our model simulations and analyses of EWS were

mainly focused on short transient dynamics after the start of the external optimum

change. Further, due to continuous and fast rate of environmental deterioration,

adaptation or evolutionary rescue by genetic evolution (Lande 1976; Lande 2009;

Chevin and Lande 2010) was not possible in our model and the populations eventually

collapsed.

Decreasing population size in response to environmental deterioration and directional

selection acting towards a mean phenotype could both lead to depletion of additive

genetic variance (Barton & Keightley 2002). We took this account in our simulations by

following a framework developed in (Bürger 2000). Using this framework to simulate

population collapse we compared strength of EWS between two cases: 1) when there

was decrease in genetic variation over time due to environmental deterioration and, 2)

when genetic variation remained constant over time, however the results were largely

similar (See Fig S6). Depletion of genetic variation over time did not affect the strength

of EWS compared with when genetic variation remained constant.

Net reproductive rate also had a significant affect on the stability of populations and

detectability of EWS of population collapse. Populations with a lower net reproductive

rate declined steeply and earlier than populations with higher net reproductive rate for

the same rate of external environmental change (Fig. S5). Higher net reproductive rate

buffered the population from declining immediately after the environment shifted when

compared to a population with lower reproductive rate (Juan-Jordá et al. 2015).

Consequently, for the same rate of change in the external optimum environment,

populations with lower reproductive rate recovered slowly due to a wide basin of

attraction than populations with higher net reproductive rate (Fig. 2, Fig. S1) and hence

strength in EWS was more pronounced in the latter.

Our modeling results suggested that including fitness-related trait information could

significantly improve the strength in EWS of population collapse. Trait-based EWS

(*SD+trait*) produced the strongest signal. Inclusion of phenotypic trait information

alongside the leading indicators produced less distributed, consistent and stronger

Kendall's tau values compared to the generic abundance-based EWS (Fig. 4, Fig S8-S9).

In all the different biological factors examined in our study the *SD+trait* metric

performed the best overall. In response to a directional change in the external optimum

environment, shift in the mean phenotypic trait occurred over time even if there were

not significant increases in either AR1 or SD. Furthermore, including information from

the mean phenotype alongside EWS statistics would depend not only on whether the

mean phenotype responded to the external forcing but also on the nature of the external

environmental forcing itself (Clements & Ozgul 2016). Nevertheless, body size is a

candidate to be included with the leading indicators (Clements & Ozgul 2016). Shifts in

body size have previously been shown to occur before population collapse in whales,

with true positive EWS being detectable earlier when information from the body size

was included in the EWS (Clements et al. 2017).

In conclusion, we showed that predictability of population collapse was affected by

factors intrinsic to a population like genetic variation, strength in adaptive plasticity and

net reproductive rate. Changes in these factors can not only alter the underlying

phenotypic and population dynamics, but also the potential energy landscape.

Consequently, due to the alteration of the potential energy landscape, the detectability

of EWS was severely hampered. However, if information from fitness-related trait was

incorporated with EWS, the strength in predicting population collapses increased

significantly. Our results add to the body of evidence suggesting that the inclusion of

trait dynamic information alongside the generic EWS should provide more accurate

forecasts of the future state of biological systems.

Author's Contribution: GB designed the study, analyzed the model and did the

simulations with feedback from CC and AO. All the authors contributed to writing the

paper.

REFERENCES

Barton, N.H. & Keightley, P.D., 2002. Understanding quantitative genetic variation.

Nature reviews. Genetics, 3(1), pp.11–21.

Beisner, B., Haydon, D. & Cuddington, K., 2003. Alternative stable states in ecology.

Frontiers in Ecology and the Environment, 1(7), pp.376–382. Available at:

https://esajournals.onlinelibrary.wiley.com/doi/10.1890/1540-

9295%282003%29001%5B0376%3AASSIE%5D2.0.CO%3B2 [Accessed May 22,

2018].

Boettiger, C. & Hastings, A., 2013. No early warning signals for stochastic transitions:

insights from large deviation theory. *Proceedings of the Royal Society, Series B*,

280(1766), p.20131372. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/23843397.

Boettiger, C. & Hastings, A., 2012. Quantifying limits to detection of early warning for

critical transitions. *Journal of the Royal Society, Interface / the Royal Society*.

Boettiger & Hastings, 2012. Quantifying limits to detection of early warning for critical

transitions. *Journal of The Royal Society Interface*, 9(75), pp.2527–2539.

Brock, W.A. & Carpenter, S.R., 2012. Early Warnings of Regime Shift When the

Ecosystem Structure Is Unknown. *PLoS ONE*, 7(9).

Bürger, R. (Reinhard), 2000. *The mathematical theory of selection, recombination, and*

mutation, Wiley.

Cardillo, M. et al., 2005. Multiple Causes of High Extinction Risk in Large Mammal

Species. *Science*, 309(5738), pp.1239–1241. Available at:

http://www.sciencemag.org/content/309/5738/1239%5Cnhttp://www.ncbi.nlm.

nih.gov/pubmed/16037416%5Cnhttp://www.sciencemag.org/content/309/5738

/1239.full%5Cnhttp://www.sciencemag.org/content/309/5738/1239.full.pdf.

Charmantier, A. et al., 2008. Adaptive phenotypic plasticity in response to climate

change in a wild bird population. *Science (New York, N.Y.)*, 320(5877), pp.800–803.

Chevin, L.M., Lande, R. & Mace, G.M., 2010. Adaptation, plasticity and extinction in a

changing enviroment: towards a predictive theory J. G. Kingsolver, ed. *PLoS biology*,

8(4), pp.1–8.

Chevin & Lande, 2010. When do adaptive plasticity and genetic evolution prevent

extinction of a density-regulated population? *Evolution*, 64(4), pp.1143–1150.

Clements et al., 2017. Body size shifts and early warning signals precede the historic

collapse of whale stocks. *Nature Ecology & Evolution*.

Clements et al., 2015. Factors Influencing the Detectability of Early Warning Signals of

Population Collapse. *The American Naturalist*, 186(1), pp.50–58. Available at:

http://www.jstor.org/stable/10.1086/681573%5Cnhttp://www.jstor.org/stable/

pdf/10.1086/681573.pdf?acceptTC=true.

Clements, C.F. & Ozgul, A., 2016. Rate of forcing and the forecastability of critical

transitions. *Ecology and Evolution*, In press(April), pp.7787–7793.

Clements & Ozgul, A., 2016. Including trait-based early warning signals helps predict

population collapse. *Nature Communications*, 7, p.10984. Available at:

http://www.nature.com/doifinder/10.1038/ncomms10984.

Clements & Ozgul, A., 2018. Indicators of transitions in biological systems J. Metcalf, ed.

Ecology Letters. Available at: http://doi.wiley.com/10.1111/ele.12948 [Accessed

May 3, 2018].

Cortez, M.H., 2016. How the Magnitude of Prey Genetic Variation Alters Predator-Prey

Eco-Evolutionary Dynamics. *The American naturalist*, 188(3), pp.329–41. Available

at: https://www.journals.uchicago.edu/doi/10.1086/687393 [Accessed May 14,

2018].

Dai, L., Korolev, K.S. & Gore, J., 2015. Relation between stability and resilience

determines the performance of early warning signals under different

environmental drivers. *Proceedings of the National Academy of Sciences of the*

United States of America, 112(32), pp.10056–61. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/26216946 [Accessed April 4, 2018].

Dakos, V. et al., 2012. Methods for detecting early warnings of critical transitions in time

series illustrated using simulated ecological data. *PLoS ONE*, 7(7).

Dakos, V. et al., 2008. Slowing down as an early warning signal for abrupt climate

change. *Proceedings of the National Academy of Sciences of the United States of*

America, 105(38), pp.14308–12. Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2567225&tool=pmce

ntrez&rendertype=abstract.

Dakos, V. & Bascompte, J., 2014. Critical slowing down as early warning for the onset of

collapse in mutualistic communities. *Proceedings of the National Academy of*

Sciences, 111(49), pp.17546–17551. Available at:

http://www.pnas.org/content/111/49/17546%5Cnhttp://www.ncbi.nlm.nih.gov

/pubmed/25422412%5Cnhttp://www.pnas.org/content/111/49/17546.full.pdf.

Drake, J.M. & Griffen, B.D., 2010. Early warning signals of extinction in deteriorating

environments. *Nature*, 467(7314), pp.456–9. Available at:

http://dx.doi.org/10.1038/nature09389 [Accessed September 3, 2015].

Dutta, P.S., Sharma, Y. & Abbott, K.C., 2018. Robustness of early warning signals for

catastrophic and non-catastrophic transitions. *Oikos*. Available at:

http://doi.wiley.com/10.1111/oik.05172 [Accessed August 30, 2018].

Falconer, D.S. & Mackay, T.F.C., 1996. *Introduction to Quantitative Genetics (4th Edition)*,

Available at: http://www.amazon.com/Introduction-Quantitative-Genetics-

Douglas-Falconer/dp/0582243025.

Frank, K.T. et al., 2011. Transient dynamics of an altered large marine ecosystem.

Nature, 477(7362), pp.86–9. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/21796120.

Gavrilets, S. & Scheiner, S.M., 1993. The genetics of phenotypic of reaction norm shape V

. Evolution. *Journal of Evolutionary Biology*, 6(1), pp.31–48. Available at:

http://onlinelibrary.wiley.com/doi/10.1046/j.1420-

9101.1993.6010031.x/abstract.

Gienapp, P. et al., 2008. Climate change and evolution: Disentangling environmental and

genetic responses. *Molecular Ecology*, 17(1), pp.167–178.

Gomulkiewicz, R. & Holt, R.D., 1995. When does evolution by natural selection prevent

extinction? *Evolution*, 49(1), pp.201–207. Available at:

http://www.jstor.org/stable/10.2307/2410305%5Cnpapers2://publication/uuid

/AF75FB13-CA45-471F-9595-604002DDAB20.

Haydon, D., 2003. Alternative stable states in ecology. *Frontiers in Ecology*.

Holling, C.S., 1973. Resilience and Stability of Ecological Systems. *Annual Review of*

Ecology and Systematics, 4(1), pp.1–23. Available at:

http://www.annualreviews.org/doi/10.1146/annurev.es.04.110173.000245

[Accessed May 22, 2018].

Jackson, J.B. et al., 2001. Historical overfishing and the recent collapse of coastal

ecosystems. *Science (New York, N.Y.)*.

Jarvis, L. et al., 2016. Early warning signals detect critical impacts of experimental

warming. *Ecology and Evolution*, 6(17), pp.6097–6106. Available at:

http://doi.wiley.com/10.1002/ece3.2339 [Accessed August 30, 2018].

Juan-Jordá, M.J. et al., 2015. Population declines of tuna and relatives depend on their

speed of life. *Proceedings of the Royal Society B: Biological Sciences*, 282(1811),

p.20150322. Available at:

http://rspb.royalsocietypublishing.org/lookup/doi/10.1098/rspb.2015.0322.

Krko ek, M. & Drake, J.M., 2014. On signals of phase transitions in salmon population

dynamics. *Proceedings of the Royal Society B: Biological Sciences*, 281(1784),

pp.20133221–20133221. Available at:

http://rspb.royalsocietypublishing.org/cgi/doi/10.1098/rspb.2013.3221.

Lade, S.J. & Gross, T., 2012. Early warning signals for critical transitions: A generalized

modeling approach. *PLoS Computational Biology*.

Lande, 2009. Adaptation to an extraordinary environment by evolution of phenotypic

plasticity and genetic assimilation. *Journal of Evolutionary Biology*, 22(7), pp.1435–

1446.

Lande, R., 1976. NATURAL SELECTION AND RANDOM GENETIC DRIFT IN PHENOTYPIC

EVOLUTION. *Evolution*, 30(2), pp.314–334. Available at:

http://doi.wiley.com/10.1111/j.1558-5646.1976.tb00911.x [Accessed April 18,

2018].

Lande & Shannon, 1996. THE ROLE OF GENETIC VARIATION IN ADAPTATION AND

POPULATION PERSISTENCE IN A CHANGING ENVIRONMENT. *Evolution*, 50(1),

pp.434–437. Available at: http://doi.wiley.com/10.1111/j.1558-

5646.1996.tb04504.x [Accessed April 18, 2018].

Mora, C. et al., 2013. Biotic and Human Vulnerability to Projected Changes in Ocean

Biogeochemistry over the 21st Century. *PLoS Biology*.

van Nes, E.H. & Scheffer, M., 2007. Slow recovery from perturbations as a generic

indicator of a nearby catastrophic shift. *The American naturalist*, 169(6), pp.738–

747.

Nolting, B.C. & Abbott, K.C., 2015. Balls, cups, and quasi-potentials: quantifying stability

in stochastic systems. *Ecology*, 97(4), p.15–1047.1. Available at:

http://doi.wiley.com/10.1890/15-1047.1 [Accessed May 21, 2018].

Ozgul, A. et al., 2010. Coupled dynamics of body mass and population growth in

response to environmental change. *Nature*, 466(7305), pp.482–485. Available at:

http://dx.doi.org/10.1038/nature09210.

Ozgul, A. et al., 2014. Linking body mass and group dynamics in an obligate cooperative

breeder A. Mysterud, ed. *Journal of Animal Ecology*, 83(6), pp.1357–1366. Available

at: http://doi.wiley.com/10.1111/1365-2656.12239 [Accessed May 13, 2018].

Price, T.D., Qvarnström, A. & Irwin, D.E., 2003. The role of phenotypic plasticity in

driving genetic evolution. *Proceedings of the Royal Society - Biological sciences*.

Reed, T.E. et al., 2010. Phenotypic plasticity and population viability: the importance of

environmental predictability. *Proceedings. Biological sciences / The Royal Society*,

277(1699), pp.3391–400. Available at:

http://rspb.royalsocietypublishing.org/content/277/1699/3391.short.

Reed, T.E., Schindler, D.E. & Waples, R.S., 2011. Efectos Interactivos de la Plasticidad

Fenot??pica y Evoluc??on sobre la Persistencia Poblacional en un Clima Cambiante.

Conservation Biology, 25(1), pp.56–63.

Robinson, J.D., Wares, J.P. & Drake, J.M., 2013. Extinction hazards in experimental

Daphnia magna populations: Effects of genotype diversity and environmental

variation. *Ecology and Evolution*.

Sanchez, A. & Gore, J., 2013. Feedback between Population and Evolutionary Dynamics

Determines the Fate of Social Microbial Populations S. P. Ellner, ed. *PLoS Biology*,

11(4), p.e1001547. Available at: http://www.ncbi.nlm.nih.gov/pubmed/23637571

[Accessed April 24, 2018].

Scheffer et al., 2009. Early-warning signals for critical transitions. *Nature*, 461, pp.53–59.

Available at: http://dx.doi.org/10.1038/nature08227%5Cn%3CGo to

ISI%3E://WOS:000269478800026%5Cnhttp://www.nature.com/nature/journal/

v461/n7260/pdf/nature08227.pdf.

Scheffer, M. et al., 2012. Anticipating Critical Transitions. *Science*, 338(6105), pp.344–

348.

Scheffer, M. et al., 2001. Catastrophic shifts in ecosystems. *Nature*, 413(6856), pp.591–6.

Available at: http://www.ncbi.nlm.nih.gov/pubmed/15448261.

Scheffer, M. et al., 2009. Early-warning signals for critical transitions. *Nature*, 461(7260),

pp.53–59. Available at: http://dx.doi.org/10.1038/nature08227%5Cn%3CGo to

ISI%3E://WOS:000269478800026%5Cnhttp://www.nature.com/nature/journal/

v461/n7260/pdf/nature08227.pdf.

Spanbauer, T.L. et al., 2016. Body size distributions signal a regime shift in a lake

ecosystem. *Proceedings of the Royal Society B*, 283, p.20160249.

Strogatz, S.H., 1994. Nonlinear Dynamics and Chaos. *Book*, pp.1–505.

Thomas, C.D. et al., 2004. Extinction risk from climate change. *Nature*, 427(6970),

pp.145–8. Available at: http://www.ncbi.nlm.nih.gov/pubmed/18653892.

Villa Martín, P. et al., 2015. Eluding catastrophic shifts. *Proceedings of the National

Academy of Sciences*.

Vuren, D. Van & Armitage, K.B., 1991. Duration of snow cover and its influence on life-

history variation in yellow-bellied marmots. *Canadian Journal of Zoology*.

Wissel, C., 1984. A universal law of the characteristic return time near thresholds.

Oecologia, 65(1), pp.101–107.

Yoshida, T. et al., 2003. Rapid evolution drives ecological dynamics in a predator–prey

system. *Nature*.

Yvonne, W. & Hoffmann, A.A., 2009. Demographic factors and genetic variation influence

population persistence under environmental change. *Journal of Evolutionary

Biology*.

System state (Population size)

Figure 1. Illustration of the 'ball-in-cup' potential energy curve in relation to stability and

resilience of a one-dimensional dynamical system. The stable state of the dynamical system in

this illustration is indicated by the blue ball, and the unstable state (or the tipping point) is

indicated by the red ball in both A and B. (1) shows that the slope of the potential curve (i.e.,

stability) is steep and the basin where the ball lies is also deep signifying that both resilience as

well as stability is high. This means that the amount of potential energy (2) to push the ball from

its stable state (blue) to the unstable state (red) or in other words the loss in stability will be

large. However, due to changes in ecological or evolutionary factors intrinsic to the dynamical

system (3), the 'ball-in-cup' potential energy curve can shift from (6) to (4) where the slope is

now less steep, signifying a loss in stability. Along with this, there is also a loss in resilience. Now,

the amount of potential energy (5) to move the blue ball at (4) to the unstable state for (B) is less.

This could reflect in the strength of EWS of population collapse.

Figure 2. Effective potential $V(N)$ plotted as a function of the state variable abundance, N. Note

the effective potential is altered by factors like genetic variation, adaptive plasticity and net

reproductive rate. The lower the value of the effective potential, more stable the system is.

Figure 3. Box plots showing the distribution of Kendall's Tau correlation coefficient (high values

around 0.7 are usually considered an EWS) for 200 simulation of population collapse for

different levels of adaptive plasticity *b*, genetic variation σ_a^2, reproductive rate R_0. The leading

indicators that are shown here are autocorrelation at lag-1 (AR1) and standard deviation (SD). X-

axis is the different levels of the above factors.

Figure 4. Box plots showing the distribution of Kendall's Tau correlation coefficient for 200

simulation of population collapse for different levels of net reproductive rate R_0. On the X-axis are

the different early warning metrics shown : autocorrelation at lag-1 (AR1), standard deviation

(SD), (*AR1+SD*),(*AR1+Tr*), (*SD+Tr*) ,(*AR1+SD+Tr*). *Tr* corresponds to the mean value of the

phenotypic trait. Colored boxplots denote the trait-based EWS while white boxplots denote

abundance-based EWS.