*Eco-evolutionary processes underlying early warning signals of population declines*

Gaurav Baruah[1], Christopher F. Clements[1,2], Arpat Ozgul[1]

[1] Department of Evolutionary Biology and Environmental studies, University of

Zurich

[2] School of Biological Sciences, University of Bristol

Keywords: stability, early warning signals, population declines, eco-evolutionary

factors, trait dynamics, quantitative genetics

**Corresponding author**: Gaurav Baruah; gaurav.baruah@ieu.uzh.ch

**Abstract**

1. Environmental change can impact the stability of populations and can cause rapid

declines in abundance.  Abundance-based warning signals have been proposed to

predict such declines, but these have been shown to have limited success, leading to

the development of warning signals based on the distribution of fitness-related traits

such as body size.

2. The dynamics of such traits in response to external environmental perturbations are

controlled by a range of underlying factors such as reproductive rate, genetic

variation, and plasticity. However, it remains unknown how such ecological and

evolutionary factors affect the stability landscape of populations and the detectability

of abundance and trait-based warning signals of population decline.

3. Here, we apply a trait-based demographic approach and investigate both trait and

population dynamics in response to gradual changes in the environment.  We explore

a range of ecological and evolutionary constraints under which the stability of a

population may be affected.

4.  We show both analytically and with model-based simulations that strength of

abundance-based early warning signals is significantly affected by ecological and

evolutionary factors.

5. Finally, we show that a unified approach, combining trait- and abundance-based

information, significantly improves our ability to predict population declines. Our

study suggests that the inclusion of trait dynamic information alongside generic

warning signals should provide more accurate forecasts of the future state of

biological systems.

**1. Introduction**

Predicting catastrophic and non-catastrophic collapses of populations and ecosystems

in response to novel environmental pressures are critical if we are to mitigate the

effects of global change and minimize the loss of biodiversity (Cardillo et al., 2005;

Thomas et al., 2004). These collapses are however difficult to predict (Boettiger &

Hastings, 2012b; Clements, Drake, Jason, & Ozgul, 2015; Clements & Ozgul, 2018).

One suite of methods, which may achieve this, are early warning signals derived from

dynamic systems theory (Scheffer et al., 2009b; van Nes & Scheffer, 2007). Such

methods assume that a dynamical system, for example a biological population, that is

under increasing continuous external pressure, will show a property called critical

slowing down (CSD) as it approaches a tipping point- the point at which the system's

state can substantially change in response to a slight perturbation. Around the vicinity

of this tipping point, the system takes longer to return to its original state after every

perturbation (Wissel, 1984; Strogatz, 1994; Scheffer *et al.*, 2009).

The occurrence of CSD is a direct implication of the dominant eigenvalue of the

system approaching zero (for continuous dynamical system) (van Nes & Scheffer,

2007) or 1 (for a discrete dynamical system) (Krkoek & Drake, 2014). As a direct

consequence of CSD, statistical moments embedded within abundance time-series

data, particularly variance and autocorrelation will show marked increase over time.

These two indicators are the prominent abundance-based statistical metrics, and,

along with various other temporal (Boettiger & Hastings, 2012a; Chevalier &

Grenouillet, 2018; Vasilis Dakos et al., 2012; Drake & Griffen, 2010; Jarvis,

McCann, Tunney, Gellner, & Fryxell, 2016) and spatial (Butitta, Carpenter, Loken,

Pace, & Stanley, 2017; Dai, Korolev, & Gore, 2013) metrics, are commonly known as

early warning signals (EWS).

Changes in autocorrelation and variance are measures of the stability of a system,

derived from the theory of alternative stable states (Clements & Ozgul, 2018; Wissel,

1984). Holling's (1973) fundamental work on stability and resilience laid the

foundation for future work on deriving stability criteria for systems with alternative

equilibrium states. Stability of a dynamical system can be characterized by the classic

ball-in-cup diagram (Clements & Ozgul, 2018; Holling, 1973; Nolting & Abbott,

2015), where the 'ball' represents the state of the system and the 'cup' represents the

potential energy landscape (Fig.1). The stability of the system is thus quantified by

the depth of the energy landscape, and resilience by the size of the cup (Clements &

Ozgul, 2018; Villa Martín, Bonachela, Levin, & Muñoz, 2015). In response to

continuous exogenous pressure, the system loses resilience (for example, decline in

population numbers) and the energy landscape shrinks (Fig. 1). Concurrent with the

shrinkage of the landscape, there is also a decrease in the depth of the 'ball-in-cup'

landscape signaling a loss in stability (Fig. 1). It is this loss in stability, which

statistical EWSs such as variance and autocorrelation measure as an alternative for

loss of global resilience of the system (Clements and Ozgul 2018). The shape of the

ball-in-cup landscape thus determines the dynamics of the state variable in response

to an exogenous environmental change (Beisner, Haydon, & Cuddington, 2003).

However, the shape of the 'ball-in-cup' landscape can be changed not only by

exogenous pressures, but can also be altered by intrinsic factors associated with the

system's dynamics (Beisner et al., 2003). Changes in these intrinsic factors might

alter the location of an equilibrium state, or destabilize the current state, causing the

system to arrive at a point closer to the other alternative state (Fig. 1). For a biological

population, the state of the population will be characterized by its current abundance.

Responses of this state variable (abundance) to an exogenous environmental pressure

will be a combination of ecological and evolutionary responses controlled by factors

such as genetic variation or plasticity (Ozgul et al., 2010; Price, Qvarnström, & Irwin,

2003). For example, plastic responses to fluctuating changes in the exogenous

environment are usually fast which consequently stabilizes population fluctuations

(Reed, Waples, Schindler, Hard, & Kinnison, 2010). However, if the environment

keeps on changing, the population might not only deplete its plastic capacity but also

its standing genetic variation, which in turn leads to its eventual decline. Further, such

a decline would also be conditional on the population's reproductive rate, with higher

reproductive rate leading to a slower decline (Juan-Jordá, Mosqueira, Freire, &

Dulvy, 2015). In addition, due to continuous changes in the exogenous environment,

selection pressures can subsequently increase leading to contemporary evolution

(Yoshida, Jones, Ellner, Fussmann, & Hairston, 2003). Adaptive evolution would

then depend on the amount of genetic variation present in the population with higher

variation leading to faster evolution, which can either stabilize or destabilize

population dynamics (Cortez, 2016; Sanchez & Gore, 2013). Since the dynamics of a

population in response to an exogenous pressure is determined by the shape of its

potential energy landscape (Beisner et al., 2003), it is thus expected that these factors

namely plasticity, genetic variation, and reproductive rate have the capacity to

influence the shape of the ball-in-cup landscape. In turn, given that EWS are stability

measures derived from alternative stable states theory (Beisner et al., 2003; Clements

& Ozgul, 2018; Scheffer et al., 2009a), it is very likely that strength in EWS and

hence predictability of population decline might also be affected by these factors.

Such theoretical expectations raise important practical questions: can predictability of

population decline with the help of EWS be affected by such ecological and

evolutionary factors?

Abundance-based measures of the stability of an ecological system might not be the

only indicators of a population being forced by an external environment (Clements &

Ozgul, 2016b) . Phenotypic traits, such as body size, is one of the important traits that

determines the fate of individuals as well as dynamics of population, and thus can

provide crucial information regarding the current state of the population (Clements

and Ozgul 2016b; Ozgul et al. 2014; Spanbauer et al. 2016). Recently, Clements &

Ozgul (2016) showed, using data from experimental microcosms, that including body

size information with EWS could significantly improve the predictive accuracy of

population declines. Such trait-based signals, have been further shown to be present in

the historic collapses of whale populations (Clements et al., 2017). However,

extensive research on the efficacy of such signals in comparison to abundance-based

EWS is still in its infancy (Clements & Ozgul, 2018).

Here in this paper, we used a quantitative genetics framework to simulate the

dynamics of a population in response to gradual environmental change. We track both

the population dynamics and trait distribution until the simulated population declines

significantly. We show that this particular model exhibits critical slowing down

behavior and hence is expected to show EWS. We then assess whether genetic

variation, plasticity in the trait, and net reproductive rate alter the ball-in-cup

landscape (stability landscape) and consequently the strength in EWS. In addition, we

assess the efficacy of trait-based EWS in comparison with abundance-based EWS and

reiterate the ability of trait-based signals in accurately informing population declines.

**2. Materials and Methods**

*2.1 Mathematical model:*

We model the joint dynamics of quantitative trait evolution and population size under

density dependent regulation via the effects of selection (Chevin, Lande, & Mace,

2010; Chevin & Lande, 2010; Gomulkiewicz & Holt, 1995) . We consider a

population that has discrete generations where individuals have fitness that is

determined by a single quantitative trait z under stabilizing selection with linear

reaction norms (Gavrilets & Scheiner, 1993). Under these assumptions the dynamics

of the population and the mean value of the trait can be written as (Lande, 2009) :

$$N_{t+1} = N_t \, \bar{R}_t = \ R_0^{(1-\frac{N}{K})} \, \overline{W} \, , \tag{1}$$

$$\bar{z}_{t+1} = \bar{a}_t + \sigma_a^2 \frac{\partial ln \overline{W}}{\partial \bar{a}} + b u_t. \tag{2}$$

$\overline{R_t}$ is the growth rate of the population at generation $t$, $R_0$ is the net reproductive rate

which is under density dependent selection given by the exponent $\left(1 - \frac{N}{K}\right)$ (Chevin &

Lande, 2010). The average fitness of the population is given by the equation $\overline{W}$ which

is due to the quantitative trait $z$. $\overline{z}_{t+1}$ is the average value of the trait at generation $t+1$,

$\overline{a}_t$ is the mean breeding value or genotypic value of the population at generation $t$,

$\frac{\partial ln \overline{W}}{\partial \overline{a}}$ is the gradient of selection on the mean trait $\overline{z}$, $\sigma_a^2$ is the additive genetic

variance, $bu_t$ quantifies the average plastic response of the trait at time $t$ , and $u$ is the

environmental cue (table 1).

Expanding $\overline{W}$ in equation (1) as follows:

$\overline{W} = \int W(z, \theta) p(z) dz = \sqrt{\frac{\sigma_w^2}{\sigma_z^2 + \sigma_w^2}} \, e^{-(\overline{z_t} - \theta_t)^2 / 2(\sigma_w^2 + \sigma_z^2)}.$        (3)

The integral in (3) over all trait values in the population gives the average fitness of

the population. $p(z)$ is the distribution of the trait in the population, $\sigma_z^2$ is the variance

in the phenotypic distribution $p(z)$ and $W(z, \theta)$ is the gaussian stabilizing fitness

function given as:

$W(z) = e^{-(z - \theta)^2 / 2(\sigma_w^2)}$                          ,        (4)

with width $w_z^2$ and optimum phenotype of $\theta$. Individuals with trait values far from the

optimum will have lower fitness compared to individuals whose trait value matches

the optimum phenotype $\theta$. Hence an individual's fitness will be determined by how

far its trait value $z$ is from the optimum phenotype $\theta$ (Chevin and Lande, 2010).

Finally, the response of the primary phenotype $z$ of an individual in this finite

population to the external environment is modeled using linear reaction norms

(Gavrilets and Scheiner, 1993; Lande and Shannon, 1996) :

$z_{i,t} = a_{i,t} + bu_t + \epsilon$                                    (5)

where $a$ is the breeding value of the individual $i$. The breeding value is normally

distributed with mean $\bar{a}$ and additive genetic variance $\sigma_a^2$, and $\epsilon$ is the residual

component with mean 0 and variance $\sigma_\epsilon^2$. The slope $b$ quantifies how plastic the trait

$z$ is in response to the environment. In our model $b$ is constant meaning that plasticity

in the trait cannot evolve (Chevin & Lande, 2010). We model an environment and the

environmental cue (see below) that determines the optimal phenotypic value $\theta_t$ for the

primary phenotype $z$. The optimal phenotype $\theta_t$, is assumed to be linearly dependent

on the external environment $E$ that selects for a particular phenotypic value such that,

$\theta_t = BE_t.$                               (6)

Expanding equation 6, the dynamics of the optimal phenotype is:

$\theta_t = BE_t = \begin{cases} 0 & , t < 500 \\ B - B\frac{t}{500} & , t \geq 500 \end{cases}$        (6a)

And the environmental cue $u$ in equation (5) is modeled as:

$u_t = \begin{cases} 0 & , t < 500 \\ c - c\frac{t}{500} & , t \geq 500 \end{cases}$        (7)

Environmental change is modeled at t = 500, where a shift in the external

environmental value $E_t$ also triggers a shift in the optimum phenotypic value. The

rate of the shift in the optimum phenotypic value is controlled by the parameter B

(equation 6a, see section 2.4 below). As in Reed et al. (2010), the dynamics of the

optimal phenotype $\theta$ in equation (6a) and the environmental cue $u$ in equation (7) are

modelled to be correlated over time. This correlation is modeled by taking the help of

a bivariate normal distribution: at each generation $t$, a value for the optimal phenotype

$\theta$ and environmental cue $u$ are drawn randomly from a bivariate normal distribution

with means given by equation (6a) and (7). The covariance of the bivariate normal

distribution is given by $\sigma_{\theta,u} = \xi \sigma_\theta \sigma_u$ where $\xi$ is the correlation between $\theta$ and the

cue $u$. If $\xi = 1$, then the values of the optimum phenotype and the environmental cue

drawn from the bivariate normal distribution will be perfectly correlated over time $t$

and when $\xi = 0$, they will not be correlated. Thus, $\xi$ determines predictability of the

external environment such that $\xi = 1$ means that an individual is able to perceive

changes in the external environmental perfectly with the help of an environmental cue

$u$ that is correlated with the changes in the external environment. In such a case, the

plastic response will track the changes in the environment and the strength of the

plastic response of the trait will be determined by $b$. We assumed that variances in the

cue $\sigma_u$ and the optimum $\sigma_\theta$ are equal (see table 1). This mimics the situation, for

instance, when variation in food availability is correlated with inter annual variation

in snow melt (Vuren & Armitage, 1991) - where in this case, inter annual variation of

snow melt is the environmental cue $u$ that the organism perceives, and variation in

food availability is the external environment $E$. Depending on the changes in the

external environment (for example, changes in food availability), the optimum

phenotype $\theta$ will change which is given by equation 6.

The above model is based on the infinitesimal quantitative genetic model (Chevin &

Lande, 2010; Falconer & Mackay, 1996).

*2.2 Stability landscape and effective potential:* One approach to quantifying stability

which is also fundamental to dynamical systems with alternative stable states is the

'ball-in-cup' analogy (Haydon, 2003; Villa Martín et al., 2015). The different states of

the dynamical system can be represented by a landscape and the state of the

population at a particular moment by a ball in that landscape. The landscape of the

dynamical system can be represented by an effective potential $V(n)$ (Beisner et al.,

2003; Villa Martín et al., 2015). If a dynamical system (example a population) can be

represented by a continuous differential equation in the form of: $\frac{dn}{dt} = g(n)$, then the

effective potential of that dynamical system can be written as : $V(n) = -\int g(n)dn$

(Beisner et al., 2003; Nolting & Abbott, 2015). For our dynamical system, $g(n) =$

$\log\left(R_0^{\left(1-\frac{n}{k}+\beta-\gamma\right)}\right)$ (see appendix 4). Here $n$ is the state of the system, which in our

case is the log of the population size $N$. The effective potential of our dynamical

system is then (appendix 4):

$$V(n) = -\left( rn - \frac{\frac{r}{k}n^2}{2} + \beta rn - \gamma rn \right)$$

$$= -\left( rn - \frac{\frac{r}{k}n^2}{2} + \frac{\log\left(\frac{\sqrt{(\sigma_w^2)}}{\sqrt{(\sigma_w^2 + \sigma_z^2)}}\right)}{\log(R_0)} rn - \frac{(\bar{a} + bu_t - \theta)^2}{(R_0(\sigma_w^2 + \sigma_z^2))} rn \right) \tag{8}$$

It is evident from the above equation (8) that the effective potential function is

dependent on three parameters: $r, \beta, \gamma$, where $r = \log(R_0)$; $\beta = \dfrac{\log\left(\frac{\sqrt{(\sigma_w^2)}}{\sqrt{(\sigma_w^2 + \sigma_z^2)}}\right)}{\log(R_0)}$ ;

$\gamma = \dfrac{(\bar{z}-\theta)^2}{(R_0(\sigma_w^2 + \sigma_z^2))}$ ; $n = \log(N)$; $k = \log(K)$. $\beta$ represents the standing load of the

population and $\gamma$ represents the lag-load.

*2.3 Equilibria and eigenvalues of the model:* The extinction equilibrium for the

model is at $N^* = 0$ and the positive equilibrium for the model is at $N^* = K + \beta K -$

$\gamma K$, where $\gamma$ is the lag-load and $\beta$ represents the standing load of the population (see

appendix 1). Stability of the two equilibria are based on their corresponding

eigenvalues, $\lambda = \frac{\partial}{\partial N} f(N)|_{N=N^*}$. Equilibria will be stable if magnitude of $\lambda < 1$ and

unstable if magnitude of $\lambda > 1$. For details of linear stability analysis and occurrence

of CSD see appendix 1 and appendix 2.

*2.4 Simulations of population declines:* We performed stochastic simulations of the

model described in section 2.1. Dynamics of the trait, population and the external

environment were iteratively updated using equations (1- 7). Without loss of

generality, the mean phenotypic value $\bar{z}_0 = \theta = 0$, at the start of the simulation, i.e.,

at t=0 (equation 6a). This meant that initially before a shift in the external

environment, average phenotypic value of the population is at its optimum. External

environmental change in our model was enforced at t=500 after the populations had

reached the carrying capacity. Stochasticity in the model was introduced at each

generation *t* by randomly drawing both optimum phenotypic value and the

environmental cue from a bivariate normal distribution with means given by equation

6a and 7 respectively, and covariance given by $\sigma_{\theta,u} = \xi \sigma_\theta \sigma_u$, where $\sigma_\theta$ and $\sigma_u$

quantifies the variance in the fluctuation of optimum phenotype and the

environmental cue over time *t*. Environmental predictability in our model was set

at $\xi = 1$, which meant that the external environmental change and the environmental

cue drawn at each generation *t* were highly correlated (see Reed et al. 2010). We

forced the populations to decline by changing the rate of change in the environment

$E_t$, at t= 500.  We chose a rate of change in the environment in a way that populations

declined regardless of any combination of parameter values. This was done by setting

B =20 in (6a) in all of our simulations, after $t \geq 500$. In all of our simulations, for

each parameter value of a factor (genetic variation, plasticity, net reproductive rate)

we did 100 independent simulations of population declines.

*2.4.1 Abundance-based EWS:* For abundance-based EWS analyses, the length of the

time-series used was of the same length for each factor (genetic variation, plasticity,

net reproductive rate) in question. For each parameter value of each factor in question,

we used 30 time-series data after the time point of environmental change i.e., after t=

500 to t=530. We discarded the rest of time-series data. We chose a rate of

environmental change in a way (B=20) that population decline was not immediate.

Significant population declines were observed across replicates for different factors

after t = 530 time point (see timeseries data, Fig. S3-S5). Next, we analyzed 30

abundance data points after t=500, when the environmental change sets in. This

ensured normalization of the length of each time series and minimized the effect of

varying length of timeseries. We then applied generic EWS of population collapse by

using the *earlywarnings* package in R. We focused our analysis on two specific

indicators namely autocorrelation at the first-lag (AR1) and standard deviation (SD)

as most of the other indicators – such as first-order autoregressive coefficient or

coefficient of variation – could be theoretically derived from these parameters (Dakos

et al. 2012). These two EWSs are theorized to increase over time before a significant

population decline (Dakos, Carpenter, van Nes, & Scheffer, 2014; Scheffer et al.,

2009a). Thus, the important question is whether these two EWSs increases before

significant population declines that was observed after time point t = 530 (see Fig. S3-

5). If the two EWSs increased over time before significant population declines, were

their performances affected by the different ecological and evolutionary factors?

The two indicators AR1 and SD were calculated using a predefined sliding window

which was 50% of the abundance time-series being analyzed.  We z-standardized the

leading indicators so that it would be easier to incorporate information from

phenotypic trait later in our analyses. We used Gaussian detrending to remove any

trend in the abundance time-series and used Kendall's Tau correlation coefficient of

the indicators prior to time point t=530 as an indicator of an approaching decline.

Kendall's tau is a non-parametric measure of rank correlation used to identify an

increasing trend. Kendall's tau takes values in the range of [-1,1].  Strong positive

Kendall's Tau correlations of the statistical indicators (SD and AR1) with time would

indicate an approaching population decline (Dakos et al. 2012). If Kendall's tau

correlation coefficient for an EWS was less than 0.5, we considered that EWS as a

false negative. Hence, we quantified reliability of EWS as the rate of false negatives:

the number of times in replicate simulations the Kendall's tau correlation coefficient

was less than 0.50. Higher proportion of false negatives would indicate less reliability

of EWS and vice-versa.

*2.4.2 Trait-based EWS:* To evaluate the efficacy of trait-based EWS (Clements &

Ozgul, 2016b), information from mean phenotype $\bar{z}$ was incorporated with the leading

indicators in an additive manner (Clements *et al.*, 2017; Clements and Ozgul, 2018).

To include information from average phenotypic value, we first z-standardized the

trait time-series.  The length of the phenotypic time-series that was used was the same

as the corresponding abundance time-series. Before a population decline, SD and

AR1 are expected to increase over time, while in our model, because of the direction

of the optimum environmental change, the trait is expected to decline (it is also

possible for trait to increase over time, but that would depend on the direction of the

optimum phenotypic change). Such a phenotypic trait, for example, could be body

size. Body size is a quantitative trait fundamental to life-history theory and population

dynamics (Cameron, O'Sullivan, Reynolds, Piertney, & Benton, 2013; Gibert, Allen,

Hruska, & DeLong, 2017; Pigeon, Ezard, Festa-Bianchet, Coltman, & Pelletier,

2017),  which has been shown to decline in response to decreases in resource

availability (Clements & Ozgul, 2016b), increases in temperature (Forster, Hirst, &

Atkinson, 2011, 2012; Gardner, Peters, Kearney, Joseph, & Heinsohn, 2011; Tseng et

al., 2018), and increases in harvest rate (Frank *et al.*, 2011; Clements *et al.*, 2017).

Whether including trait information improves predictability of population decline will

be dependent on whether the trait responds to an external environmental change.

Here, in our model, the optimum environmental change will affect the dynamics of

the phenotype. However, whether the change in the dynamics of the phenotype will

occur within the time series length being analyzed here and hence be informative of

population decline is unknown. It has been shown that it is possible for shifts in

phenotypic dynamics to precede population declines, but only under certain

environmental circumstances (Baruah *et al.*, 2018 *(in press)*). As of yet, extensive

analysis of the efficacy of EWS and trait-based EWSs has not been done.

Since the phenotype is expected to decrease in response to the change in the optimum

environment, the z-standardized values of phenotypic timeseries were multiplied by -

1 so that they could be included with the other two standardized statistical metrics

such as SD and AR1. The values of standardized EWS such as SD and AR1 were

added additively with standardized mean phenotypic time series to quantitatively

create trait-based signals or trait-based EWS.  We evaluated three trait-based EWS

metrics namely *AR1+trait, AR1+SD+trait, SD+trait* and compared how the metrics

that included information from the mean phenotype improved predictability of

population declines when compared with the abundance-based EWS like *AR1, SD.*

We quantified Kendall's Tau correlation coefficient as an indicator for strength of

predictability of population declines and reliability of the trait-based EWS as the rate

of false negatives.

*2.4.3 Adaptive plasticity:* To assess the role of adaptive plasticity *b*, we simulated 100

replicate population declines for each parameter value of *b,* which ranged from low to

high: 0.05, 0.1, 0.3, 0.4, and 0.8. While we varied plasticity, we fixed $\sigma_a^2 = 0.25$, and

$R_0$ at 1.2. The width of the fitness function determines the strength of selection in the

trait. We fixed the width of the fitness function at $\sigma_w^2 = 20$, which would indicate

medium strength in selection. Usually, $\sigma_w^2 = 50$ would indicate weak strength in

selection and $\sigma_w^2 = 5$ would indicate strong selection (Chevin and Lande 2010).

*2.4.4 Genetic variation:* The levels of genetic variation $\sigma_a^2$ used for the simulations of

population declines are 0.05, 0.1, 0.2, 0.3, 0.4, 0.5. Adaptive plasticity *b*, net

reproductive rate $R_0$ and $\sigma_w^2$ were kept at 0.2, 1.2 and 20 respectively.

We also did simulations where we relaxed the assumption of genetic variation ($\sigma_a^2$)

remaining constant during population collapses. For details see appendix 5.

*2.4.5 Net reproductive rate:* To assess the role $R_0$, we simulated 100 replicate

population declines. The levels of $R_0$ used for the simulations of population collapse

were 1.1, 1.2, 1.3, 1.4, and 1.5. Adaptive plasticity *b*, genetic variation $\sigma_a^2$ and $\sigma_w^2$,

were kept at 0.2, 0.25 and 20 respectively.

**3. Results**

*3.1 Effective potential*

*3.1.1 Strength of plasticity :* The term $\gamma = \frac{(\bar{z}-\theta)^2}{(R_0(\sigma_w^2+\sigma_z^2))}$ in equation (8) is the

evolutionary lag due to maladaptation of the mean phenotype with the optimum

(Lande, 2009). The effective potential becomes narrower with increasing strength of

plasticity and consequentially increases stability, i.e., the potential function becomes

more negative (Fig. 2). Since return time and stability are inversely related (Dai et al.,

2015), return time decreases with increasing strength of plasticity (Appendix 2, Fig.

S1). In other words, the population quickly recovers with every perturbation if

strength of plasticity is high.

*3.1.2 Genetic variation:* The mean phenotype has variance of $\sigma_z^2 = \sigma_a^2 + \sigma_\epsilon$.

Assuming residual component $\sigma_\epsilon$ to be zero, phenotypic variance is then equal to

additive genetic variance, $\sigma_z^2 = \sigma_a^2$. Inspection of the equation (8) reveals that high

additive genetic variance causes the effective potential to be more positive when

compared with low $\sigma_a^2$ and consequently decreases stability, i.e., the effective

potential becomes less negative (Fig. 2; the more negative the effective potential,

more stable the population is). Reduction of stability and effective potential energy is

higher for populations with low genetic variance whenever they are forced to collapse

compared to populations with high genetic variance (Fig. 2). Further, return time also

increases with increasing additive genetic variance (Appendix 2, Fig. S1).

*3.1.3 Net reproductive rate:* From equation (8), we also see that higher values of $R_0$

lead to higher stability and narrow potential curve for a given state of the population

(Fig. 2). Loss in stability is more when $R_0$ is higher for the same rate of change in the

environmental change when compared with a lower $R_0$. Further, return time also

decreases with increasing $R_0$ (appendix 2, Fig. S1).

*3.2 Simulations of population declines: abundance-based EWS*

*3.2.1 Strength of plasticity:* The strength of EWSs of population collapse (Kendall's

Tau value) was much higher in populations which had high adaptive plasticity $\boldsymbol{b} = 0.8$

(Fig. 3A, B) when compared with populations with low adaptive plasticity. Moreover,

the rate of false negatives decreased for all the abundance and trait-based metrics as

strength of plasticity increased (Fig. 5A). Rate of false negatives however remained

more or less same for AR1 across different plasticity levels (Fig. 5A).

*3.2.2 Genetic variation:* Higher genetic variation in the population led to slight

decrease in detectability of population decline than in populations with lower genetic

variation, particularly for SD (Fig. 3C, D). Moreover, the rate of false negatives for

SD, AR1 remained more or less same as genetic variation decreased (Fig. 5B). For

trait-based EWSs, particularly for *SD+trait*, rate of false negatives was the lowest

(Fig. 5B).

*3.2.3 Net reproductive rate:* Strength in EWS was lower in populations with lower $R_0$

when compared with higher $R_0$, particularly for SD (Fig. 3E). AR1 however showed a

slightly different result (Fig. 3F). In case of reliability of abundance-based EWS, the

rate of false negatives decreased for SD as net reproductive rate increased (Fig. 5C).

However, for AR1 rate of false negatives remained unchanged as $R_0$ increased (Fig.

5C).

*3.3 Simulations of population declines: trait-based EWS:* Trait-based EWS improved

the strength of EWS of population declines (Fig. 4, Fig S8-S9). In principle, *AR1* was

the least efficient of all the signals compared. However, when trait information was

included with *AR1*, strength in trait-based *AR1 (trait+AR1)* improved significantly

(Fig.4, Fig. 5). Overall, the Kendall's Tau value of SD and *SD+trait* was relatively

higher than all the other trait-based and abundance-based EWS.  This was consistent

across all the factors analyzed such as genetic variation, plasticity (Fig.4, Fig. S8-S9).

The rate of false negatives was highest for AR1 across all factors analyzed (Fig. 5A-

C). For plasticity, the rate of false negatives decreased as strength of plasticity

increased, but this was however opposite for genetic variation where the false

negatives slightly increased as genetic variance increased (Fig. 5B). Furthermore,

even though AR1 had the highest number of false negatives, false negatives of

*AR1+trait* significantly decreased (Fig. 5).

**4. Discussion**

The early detection of population declines with the help of EWS provides a unique

tool to highlight conservation efforts to a population approaching an impending

transition. We show in this paper that factors intrinsic to a population in question such

as genetic variation, plasticity or net reproductive rate could significantly alter

stability of a population as well as strength and reliability of EWS of population

decline.

The detectability of EWSs is dependent not only on the relation between stability and

resilience but is also determined by a variety of factors including sampling frequency,

and the strength and type of external forcing (Dai, Korolev, and Gore 2015; Clements

et al. 2015a; Clements and Ozgul 2016). Here we add to this body of knowledge by

demonstrating that the stability of a system can also be altered by intrinsic factors

such as genetic variation, plasticity and net reproductive rate. These changes to the

effective potential in turn led to stronger EWS being observed, whenever changes in

the external optimum caused a population to lose a significant amount of effective

potential energy (Fig. 1). In other words, stronger EWSs would be observed whenever

a population is forced to move from a deeper point in the potential energy curve

(more stable) to its tipping point compared to another population, which is lying on a

shallower potential (less stable) (Fig. 2). Our findings also suggested that the utility of

EWS was strongly determined by factors that are intrinsic to a population such as

genetic variation, net reproductive rate and plasticity.

Higher strength of plasticity in the phenotype significantly affected the detectability

of population collapse (Fig. 2). Strength of plasticity altered the response and the

dynamics of a population to changes in the external environment. When plasticity was

high the response of the phenotype tracked the environment perfectly causing the

population to stabilize in response to fluctuating changes in the environment (Fig. S3)

(Charmantier et al., 2008; Reed, Schindler, & Waples, 2011). The stability of a

population is also reflected in short return time to equilibrium (van Nes & Scheffer,

2007). This was true for the case where the strength of plasticity was high (Fig. S1).

Moreover, the basin of attraction for a highly plastic population was deeper when

compared to a less plastic population, which was shallower. To push a highly plastic

population from the deepest point in its effective potential energy curve to the tipping

point, the energy loss would be more, compared to a less plastic population which

was lying in a much shallower basin. The loss in energy from driving the highly

plastic population from the deepest point to its tipping point was reflected in stronger

EWS when one compared it with another less plastic population. Consequently, for

the same rate of change in the environment, higher values of plasticity lead to stronger

and more reliable EWS of population collapse.

Similarly, the effective potential of our dynamical system was also altered by genetic

variation (Fig. 2), with higher genetic variation leading to a shallower stability

landscape than that of a population with lower genetic variation. High additive genetic

variance created what was called a 'genetic load' that lowered the overall population

growth rate (Lande 1976; Lande and Shannon 1996; Chevin, Lande, and Mace 2010).

This was due to stabilizing selection acting on the mean phenotype. Due to this

genetic load caused by high additive genetic variance, population growth rate at

equilibrium was lower causing the effective potential to be closer towards the tipping

point compared to another population having a lower genetic variation. Hence for the

same rate of external environmental change, the loss of potential energy was different

when one compared a population with high genetic variation to that with low genetic

variation. As a result, the performance of EWS was affected with slightly lower

detectability and lower reliability of population collapse when there was higher

genetic variation (Fig. 3, Fig. 5B).  This result however did not mean that genetic

variation was detrimental for long-term population persistence. Time to extinction

was however longer for populations with higher genetic variation when compared

with a population which had a lower genetic variation in the trait (Fig S2)(Robinson,

Wares, & Drake, 2013; Willi & Hoffmann, 2009). High genetic variation is necessary

for rapid adaptation via genetic evolution, which then aids in evolutionary rescue (

Ashander, Chevin and Baskett, 2016).  However, our model simulations and analyses

of EWS were mainly focused on short transient dynamics after the start of the

external optimum change. Further, due to continuous and fast rate of environmental

deterioration, adaptation or evolutionary rescue by genetic evolution (Lande 2009;

Chevin and Lande 2010) was not possible in our model and the populations

eventually declined.

Decreasing population size in response to environmental deterioration and directional

selection acting towards a mean phenotype could both lead to depletion of additive

genetic variance (Barton & Keightley, 2002). We took this account in our simulations

by following a framework developed in (Bu☐rger, 2000) (see appendix 5). Using this

framework, we find that depletion of genetic variation over time did not affect the

strength of EWS compared with when genetic variation remained constant (See Fig

S6).

Net reproductive rate also had a significant effect on the stability of populations,

detectability and reliability of EWS of population collapse.  Higher net reproductive

rate buffered the population from declining immediately after the environment shifted

when compared to a population with lower reproductive rate (Juan-Jordá et al., 2015).

Consequently, for the same rate of change in the external optimum environment,

populations with lower reproductive rate recovered slowly due to a wide basin of

attraction than populations with higher net reproductive rate (Fig. 2, Fig. S1) and

hence strength in EWS was more pronounced in the latter.

Our modeling results suggested that including fitness-related trait information could

significantly improve the strength and reliability in EWS of population collapse.

Trait-based EWS (*SD+trait*) produced strongest and reliable signals. Inclusion of

phenotypic trait information alongside the leading indicators produced less

distributed, consistent and stronger Kendall's tau values compared to the generic

abundance-based EWS (Fig. 4, Fig S7-S8). This is because in response to a

directional change in the external optimum environment, shift in the mean phenotypic

trait occurred over time even if there were no significant increases in either AR1 or

SD. Furthermore, including information from the mean phenotype alongside EWS

statistics would depend not only on whether the mean phenotype responded to the

external forcing but also on the nature of the external environmental forcing itself

(Clements & Ozgul, 2016b). Body  size is suggested to be an ideal candidate to be

included with the leading indicators (Clements & Ozgul, 2016b). Shifts in body size

have previously been shown to occur before population collapse in whales, with true

positive EWS being detectable earlier when information from the body size was

included in the EWS (Clements et al., 2017). Furthermore, it has been suggested that

in certain environmental change scenarios, it is possible for body size to shift before a

population decline and thus act as a warning signal (Baruah *et al.*, 2018 *(in press)*).

In conclusion, we showed that predictability of population decline was affected by

factors intrinsic to a population like genetic variation, strength in adaptive plasticity

and net reproductive rate. Changes in these factors can not only alter the underlying

phenotypic and population dynamics, but also the effective potential landscape.

Consequently, due to the alteration of the effective potential landscape, the

detectability of EWS was hampered. However, if information from fitness-related

trait was incorporated with EWS, the strength in predicting population collapses

increased significantly. Our results suggests that the inclusion of trait dynamic

information alongside the generic EWS should provide more accurate forecasts of the

future state of biological systems.

**5. Code availability**: R version 3.5.1. R script for the code is made available:

https://github.com/GauravKBaruah/Eco-Evo-EWS

**6. Data availability:** will be made available at Dryad repository.

**7. Author contributions:**

GB conceived the ideas and designed the methodology; GB analyzed the data with

feedback from CFC and AO; All authors contributed to the drafts and final approval

for submission.

**8. References**

Ashander, J., Chevin, L.-M., & Baskett, M. L. (2016). Predicting evolutionary rescue

via evolving plasticity in stochastic environments. *Proceedings of the Royal*

521   *Society B: Biological Sciences*, *283*(1839), 20161690.

522   doi:10.1098/rspb.2016.1690

523  Barton, N. H., & Keightley, P. D. (2002). Understanding quantitative genetic

524   variation. *Nature Reviews. Genetics*, *3*(1), 11–21. doi:10.1038/nrg700

525  Baruah, G., Clements, C. F., Guillaume, F., & Ozgul, A. (2018). When do shifts in

526   trait dynamics precede population declines? *BioRxiv*, 424671.

527   doi:10.1101/424671

528  Beisner, B., Haydon, D., & Cuddington, K. (2003). Alternative stable states in

529   ecology. *Frontiers in Ecology and the Environment*, *1*(7), 376–382.

530   doi:10.1890/1540-9295(2003)001[0376:ASSIE]2.0.CO;2

531  Boettiger, C., & Hastings,  a. (2012a). Early warning signals and the prosecutor's

532   fallacy. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1748),

533   4734–4739.

534  Boettiger, & Hastings. (2012b). Quantifying limits to detection of early warning for

535   critical transitions. *Journal of The Royal Society Interface*.

536   doi:10.1098/rsif.2012.0125

537  Bu⬜rger, R. (Reinhard). (2000). *The mathematical theory of selection,*

538   *recombination, and mutation*. Wiley.

539  Butitta, V. L., Carpenter, S. R., Loken, L. C., Pace, M. L., & Stanley, E. H. (2017).

540   Spatial early warning signals in a lake manipulation. *Ecosphere*, *8*(10), e01941.

541   doi:10.1002/ecs2.1941

542  Cameron, T. C., O'Sullivan, D., Reynolds, A., Piertney, S. B., & Benton, T. G.

(2013). Eco-evolutionary dynamics in response to selection on life-history.

*Ecology Letters*, *16*(6), 754–63. doi:10.1111/ele.12107

Cardillo, M., Mace, G. M., Jones, K. E., Bielby, J., Bininda-Emonds, O. R. P.,

Sechrest, W., … Purvis, A. (2005). Multiple causes of high extinction risk in

large mammal species. *Science (New York, N.Y.)*, *309*(5738), 1239–41.

doi:10.1126/science.1116030

Charmantier, A., McCleery, R. H., Cole, L. R., Perrins, C., Kruuk, L. E. B., &

Sheldon, B. C. (2008). Adaptive phenotypic plasticity in response to climate

change in a wild bird population. *Science (New York, N.Y.)*, *320*(5877), 800–803.

doi:10.1126/science.1157174

Chevalier, M., & Grenouillet, G. (2018). Global assessment of early warning signs

that temperature could undergo regime shifts. *Scientific Reports*, *8*(1), 10058.

doi:10.1038/s41598-018-28386-x

Chevin, L. M., Lande, R., & Mace, G. M. (2010). Adaptation, plasticity and extinction

in a changing enviroment: towards a predictive theory. *PLoS Biology*, *8*(4), 1–8.

Chevin, & Lande. (2010). When do adaptive plasticity and genetic evolution prevent

extinction of a density-regulated population? *Evolution*, *64*(4), 1143–1150.

doi:10.1111/j.1558-5646.2009.00875.x

Clements, Blanchard, J. L., Nash, K. L., Hindell, M. A., & Ozgul, A. (2017). Body

size shifts and early warning signals precede the historic collapse of whale

stocks. *Nature Ecology & Evolution*, *1*(7), 0188. doi:10.1038/s41559-017-0188

Clements, C. F., & Ozgul, A. (2016a). Rate of forcing and the forecastability of

critical transitions. *Ecology and Evolution*, *In press*(April), 7787–7793.

doi:10.1002/ece3.2531

Clements, Drake, J. M., Jason, I. G., & Ozgul, A. (2015). Factors Influencing the

Detectability of Early Warning Signals of Population Collapse. *The American*

*Naturalist*, *186*(1), 50–58. doi:10.1086/681573

Clements, & Ozgul, A. (2016b). Including trait-based early warning signals helps

predict population collapse. *Nature Communications*, *7*, 10984. Retrieved from

http://www.nature.com/doifinder/10.1038/ncomms10984

Clements, & Ozgul, A. (2018). Indicators of transitions in biological systems.

*Ecology Letters*, *21*(6), 905–919. doi:10.1111/ele.12948

Cortez, M. H. (2016). How the Magnitude of Prey Genetic Variation Alters Predator-

Prey Eco-Evolutionary Dynamics. *The American Naturalist*, *188*(3), 329–41.

doi:10.1086/687393

Dai, L., Korolev, K. S., & Gore, J. (2013). Slower recovery in space before collapse

of connected populations. *Nature*, *496*(7445), 355–358. doi:10.1038/nature12071

Dai, L., Korolev, K. S., & Gore, J. (2015). Relation between stability and resilience

determines the performance of early warning signals under different

environmental drivers. *Proceedings of the National Academy of Sciences of the*

*United States of America*, *112*(32), 10056–61. doi:10.1073/pnas.1418415112

Dakos, V., Carpenter, S. R., Brock, W. A., Ellison, A. M., Guttal, V., Ives, A. R., …

Scheffer, M. (2012). Methods for detecting early warnings of critical transitions

in time series illustrated using simulated ecological data. *PLoS ONE*, *7*(7).

doi:10.1371/journal.pone.0041010

Dakos, V., Carpenter, S. R., van Nes, E. H., & Scheffer, M. (2014). Resilience

indicators: prospects and limitations for early warnings of regime shifts.

*Philosophical Transactions of the Royal Society B: Biological Sciences*,

*370*(1659), 20130263–20130263. doi:10.1098/rstb.2013.0263

Dakos, V., Scheffer, M., van Nes, E. H., Brovkin, V., Petoukhov, V., & Held, H.

(2008). Slowing down as an early warning signal for abrupt climate change.

*Proceedings of the National Academy of Sciences of the United States of*

*America*, *105*(38), 14308–12. doi:10.1073/pnas.0802430105

Drake, J. M., & Griffen, B. D. (2010). Early warning signals of extinction in

deteriorating environments. *Nature*, *467*(7314), 456–9. doi:10.1038/nature09389

Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to Quantitative Genetics*

*(4th Edition). Trends in Genetics* (Vol. 12). Retrieved from

http://www.amazon.com/Introduction-Quantitative-Genetics-Douglas-

Falconer/dp/0582243025

Forster, J., Hirst, A. G., & Atkinson, D. (2011). How do organisms change size with

changing temperature? The importance of reproductive method and ontogenetic

timing. *Functional Ecology*, *25*(5), 1024–1031. doi:10.1111/j.1365-

2435.2011.01852.x

Forster, J., Hirst, A. G., & Atkinson, D. (2012). Warming-induced reductions in body

size are greater in aquatic than terrestrial species. *Proceedings of the National*

*Academy of Sciences of the United States of America*, *109*(47), 19310–4.

doi:10.1073/pnas.1210460109

Frank, K. T., Petrie, B., Fisher, J. A., & Leggett, W. C. (2011). Transient dynamics of

an altered large marine ecosystem. *Nature*. doi:10.1038/nature10285

Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011).

Declining body size: A third universal response to warming? *Trends in Ecology*

*and Evolution*. doi:10.1016/j.tree.2011.03.005

Gavrilets, S., & Scheiner, S. M. (1993). The genetics of phenotypic of reaction norm

shape V . Evolution. *Journal of Evolutionary Biology*, *6*(1), 31–48.

doi:10.1046/j.1420-9101.1993.6010031.x

Gibert, J. P., Allen, R. L., Hruska, R. J., & DeLong, J. P. (2017). The ecological

consequences of environmentally induced phenotypic changes. *Ecology Letters*,

*20*(8), 997–1003. doi:10.1111/ele.12797

Gomulkiewicz, R., & Holt, R. D. (1995). When does evolution by natural selection

prevent extinction? *Evolution*, *49*(1), 201–207. Retrieved from

http://www.jstor.org/stable/10.2307/2410305%5Cnpapers2://publication/uuid/A

F75FB13-CA45-471F-9595-604002DDAB20

Haydon, D. (2003). Alternative stable states in ecology. *Frontiers in Ecology*.

doi:10.1890/1540-9295(2003)001[0376:ASSIE]2.0.CO;2

Holling, C. S. (1973). Resilience and Stability of Ecological Systems. *Annual Review*

*of Ecology and Systematics*, *4*(1), 1–23.

doi:10.1146/annurev.es.04.110173.000245

Jarvis, L., McCann, K., Tunney, T., Gellner, G., & Fryxell, J. M. (2016). Early

warning signals detect critical impacts of experimental warming. *Ecology and*

*Evolution*. doi:10.1002/ece3.2339

Juan-Jordá, M. J., Mosqueira, I., Freire, J., & Dulvy, N. K. (2015). Population

declines of tuna and relatives depend on their speed of life. *Proceedings of the*

*Royal Society B: Biological Sciences*, *282*(1811), 20150322.

doi:10.1098/rspb.2015.0322

Krko ek, M., & Drake, J. M. (2014). On signals of phase transitions in salmon

population dynamics. *Proceedings of the Royal Society B: Biological Sciences*,

*281*(1784), 20133221–20133221. doi:10.1098/rspb.2013.3221

Lande. (2009). Adaptation to an extraordinary environment by evolution of

phenotypic plasticity and genetic assimilation. *Journal of Evolutionary Biology*,

*22*(7), 1435–1446. doi:10.1111/j.1420-9101.2009.01754.x

Lande, R. (1976). NATURAL SELECTION AND RANDOM GENETIC DRIFT IN

PHENOTYPIC EVOLUTION. *Evolution*, *30*(2), 314–334. doi:10.1111/j.1558-

5646.1976.tb00911.x

Lande, & Shannon. (1996). THE ROLE OF GENETIC VARIATION IN

ADAPTATION AND POPULATION PERSISTENCE IN A CHANGING

ENVIRONMENT. *Evolution*, *50*(1), 434–437. doi:10.1111/j.1558-

5646.1996.tb04504.x

Nolting, B. C., & Abbott, K. C. (2015). Balls, cups, and quasi-potentials: quantifying

stability in stochastic systems. *Ecology*, *97*(4), 15–1047.1. doi:10.1890/15-

1047.1

Ozgul, A., Bateman, A. W., English, S., Coulson, T., & Clutton-Brock, T. H. (2014).

Linking body mass and group dynamics in an obligate cooperative breeder.

*Journal of Animal Ecology*, *83*(6), 1357–1366. doi:10.1111/1365-2656.12239

Ozgul, A., Childs, D. Z., Oli, M. K., Armitage, K. B., Blumstein, D. T., Olson, L. E.,

… Coulson, T. (2010). Coupled dynamics of body mass and population growth

in response to environmental change. *Nature*, *466*(7305), 482–485. Retrieved

from http://dx.doi.org/10.1038/nature09210

Pigeon, G., Ezard, T. H. G., Festa-Bianchet, M., Coltman, D. W., & Pelletier, F.

(2017). Fluctuating effects of genetic and plastic changes in body mass on

population dynamics in a large herbivore. *Ecology*, *98*(9), 2456–2467.

doi:10.1002/ecy.1940

Price, T. D., Qvarnström, A., & Irwin, D. E. (2003). The role of phenotypic plasticity

in driving genetic evolution. *Proceedings of the Royal Society - Biological*

*Sciences*. doi:10.1098/rspb.2003.2372

Reed, T. E., Schindler, D. E., & Waples, R. S. (2011). Efectos Interactivos de la

Plasticidad Fenot??pica y Evoluc??on sobre la Persistencia Poblacional en un

Clima Cambiante. *Conservation Biology*, *25*(1), 56–63. doi:10.1111/j.1523-

1739.2010.01552.x

Reed, T. E., Waples, R. S., Schindler, D. E., Hard, J. J., & Kinnison, M. T. (2010).

Phenotypic plasticity and population viability: the importance of environmental

predictability. *Proceedings. Biological Sciences / The Royal Society*, *277*(1699),

3391–400. doi:10.1098/rspb.2010.0771

Robinson, J. D., Wares, J. P., & Drake, J. M. (2013). Extinction hazards in

experimental Daphnia magna populations: Effects of genotype diversity and

environmental variation. *Ecology and Evolution*. doi:10.1002/ece3.449

Sanchez, A., & Gore, J. (2013). Feedback between Population and Evolutionary

Dynamics Determines the Fate of Social Microbial Populations. *PLoS Biology*,

*11*(4), e1001547. doi:10.1371/journal.pbio.1001547

Scheffer, Bascompte, Brock, W. A., Brovkin, V., Carpenter, S. R., Dakos, V., …

Sugihara, G. (2009a). Early-warning signals for critical transitions. *Nature*, *461*,

53–59. Retrieved from http://dx.doi.org/10.1038/nature08227%5Cn%3CGo to

ISI%3E://WOS:000269478800026%5Cnhttp://www.nature.com/nature/journal/v

461/n7260/pdf/nature08227.pdf

Scheffer, M., Bascompte, J., Brock, W. A., Brovkin, V., Carpenter, S. R., Dakos, V.,

… Sugihara, G. (2009b). Early-warning signals for critical transitions. *Nature*,

*461*(7260), 53–59. doi:10.1038/nature08227

Spanbauer, T. L., Allen, C. R., Angeler, D. G., Eason, T., Fritz, S. C., Garmestani, A.

S., … Sundstrom, S. M. (2016). Body size distributions signal a regime shift in a

lake ecosystem. *Proceedings of the Royal Society B*, *283*, 20160249.

doi:http://dx.doi.org/10.1098/rspb.2016.0249

Strogatz, S. H. (1994). Nonlinear Dynamics and Chaos. *Book*, 1–505.

doi:9780738204536

Thomas, C. D., Thomas, C. D., Cameron, A., Cameron, A., Green, R. E., Green, R.

E., … Williams, S. E. (2004). Extinction risk from climate change. *Nature*,

*427*(6970), 145–8. doi:10.1038/nature02121

Tseng, M., Kaur, K. M., Soleimani Pari, S., Sarai, K., Chan, D., Yao, C. H., …

Fograscher, K. (2018). Decreases in beetle body size linked to climate change

and warming temperatures. *Journal of Animal Ecology*, *87*(3), 647–659.

doi:10.1111/1365-2656.12789

van Nes, E. H., & Scheffer, M. (2007). Slow recovery from perturbations as a generic

indicator of a nearby catastrophic shift. *The American Naturalist*, *169*(6), 738–

747. doi:10.1086/516845

Villa Martín, P., Bonachela, J. A., Levin, S. A., & Muñoz, M. A. (2015). Eluding

catastrophic shifts. *Proceedings of the National Academy of Sciences*.

doi:10.1073/pnas.1414708112

Vuren, D. Van, & Armitage, K. B. (1991). Duration of snow cover and its influence

on life-history variation in yellow-bellied marmots. *Canadian Journal of*

*Zoology*, *69*(7), 1755–1758. doi:10.1139/z91-244

Willi, Y., & Hoffmann, A. A. (2009). Demographic factors and genetic variation

influence population persistence under environmental change. *Journal of*

*Evolutionary Biology*, *22*(1), 124–133. doi:10.1111/J.1420-9101.2008.01631.X

Wissel, C. (1984). A universal law of the characteristic return time near thresholds.

*Oecologia*, *65*(1), 101–107. doi:10.1007/BF00384470

Yoshida, T., Jones, L. E., Ellner, S. P., Fussmann, G. F., & Hairston, N. G. (2003).

Rapid evolution drives ecological dynamics in a predator–prey system. *Nature*.

doi:10.1038/nature01767

**Table 1:** List of parameter values and variables used in the model simulations. The

variables are not given any value but factors for which the model simulations are

tested are given values.

| Parameters/variables | Description | Value |
|---|---|---|
| $N_t$ | Abundance of the population at time t | .... |
| $\bar{z}_t$ ; $\bar{a}_t$ | Mean trait value at time t; mean breeding value at time t | .... |
| $\sigma_z^2$; $\sigma_a^2$ | Variance of the mean trait; additive genetic variance or variance of the mean breeding value | [0.05, 0.1, 0.2, 0.3, 0.4,0.5] ; [0.05,0.1, 0.2, 0.3, 0.4,0.5] |
| $b$ | Strength in phenotypic plasticity | [0.05, 0.1, 0.3,0.4, 0.8] |
| $R_0$ | Net reproductive rate | [1.1, 1.2,1.3,1.4, 1.5] |
| $u_t$ | Environmental cue at time $t$ | $$U_t = \begin{cases} 0 & , t < 500 \\ 1 - c\dfrac{t}{500} & , t \geq 500 \end{cases}$$ |
| $\theta_t$ | Optimum phenotype at time $t$ | $$\theta_t = BE_t = \begin{cases} 0 & , t < 500 \\ B - B\dfrac{t}{500} & , t \geq 500 \end{cases}$$ (B = 20) |
| $w_z^2$ | Width of the Gaussian stabilizing fitness function. A measure of the strength in selection | 20 |
| $z_i, a_i$ | Trait value for an individual $i$ ; breeding value for an individual $i$ | .... |
| $\epsilon$ | Residual component or unexplained component of a trait z | 0 |
| $\sigma_\epsilon^2$ | Variance in that residual component $\epsilon$ | 0 |
| $E_t$ | External environmental value at time $t$ | …. |
| $B$ | Coefficient that captures the strength in the external environmental change | 20 |
| $c$ | Coefficient that controls the rate of change in the environmental cue | 5 |
| $\xi$ | Environmental predictability; | 1 |

| | quantifies how much the external environment is correlated with the environmental cue | |
|---|---|---|
| $\sigma_\theta, \sigma_u$ | Variance in optimum phenotype, variance in environmental cue | $1.25^2$, $1.25^2$ |

*Figure Captions:*

**Figure 1.** Illustration of the 'ball-in-cup' potential energy curve in relation to

stability and resilience of a one-dimensional dynamical system. The stable state of

the dynamical system (A) which is an isolated population, can be indicated by the

blue ball, and the unstable state (or the tipping point) can be indicated by the red ball

in both A and B. (1)- shows that the slope of the potential curve (i.e., stability) is steep

and the basin where the ball lies is also wide signifying that both resilience as well as

stability is high. This means that the amount of effective potential energy needed (2)

to push the ball from its stable state (blue) to the unstable state (red) (or in other

words the reduction in stability) will be large. However, for another isolated

population, there might be intrinsic differences in ecological or evolutionary factors

controlling the dynamics of that isolated population and this might cause (3) a shift in

the 'ball-in-cup' potential energy curve from (6) to (4), for that population. In (4) the

slope is now less steep, signifying a reduction in intrinsic stability. Along with this,

there is also a reduction in intrinsic resilience. Now, the amount of energy (5) to move

the blue ball at (4) to the unstable state for (B) is less. This could reflect in the

performances of EWS of population declines.

**Figure 2.** Effective potential $V(N)$ plotted as a function of the state variable

abundance, N. Note the effective potential is altered by factors like genetic variation,

adaptive plasticity and net reproductive rate. The lower the value of the effective

potential, more stable the system is.

**Figure 3.** Box plots showing the distribution of Kendall's Tau correlation coefficient

(high values around 0.5 are usually considered an EWS) for 100 simulation of

population declines for different levels of adaptive plasticity $b$, genetic variation $\sigma_a^2$,

reproductive rate $R_0$. The leading indicators that are shown here are autocorrelation at

lag-1 (AR1) and standard deviation (SD). X-axis is the different levels of the above

factors.

**Figure 4.** Box plots showing the distribution of Kendall's Tau correlation coefficient

for 100 simulation of population declines for different levels of adaptive plasticity $b$.

On the X-axis are the different early warning metrics shown: autocorrelation at lag-1

(AR1), standard deviation (SD), *(AR1+SD), (AR1+Tr), (SD+Tr ),(AR1+SD+Tr). Tr*

corresponds to the mean value of the phenotypic trait. Colored boxplots denote the

trait-based EWS while white boxplots denote abundance-based EWS.

**Figure 5.** Proportion of false negatives shown here for a threshold of 0.5 (which

means Kendall's tau value below 0.5 are considered not an EWS) on the Y-axis for

abundance and trait-based EWS metrics for different levels of (A) plasticity strength ,

(B) genetic variation, (C) net reproductive rate.

770                                                    System state (Population size)

771    Figure 1.

781    Figure 2.

784    Figure 3.

787    Figure. 4

790    Figure 5.