Title Page

No clear monogenic links between left-handedness and *situs inversus*

Merel C. Postema[1], Amaia Carrion-Castillo[1], Simon E. Fisher[1,2], Guy Vingerhoets[3], Clyde Francks[1,2] *

[1] Department of Language & Genetics, Max Planck Institute for Psycholinguistics, Nijmegen, Netherlands

[2] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, Netherlands

[3] Department of Experimental Psychology, Ghent University, Belgium

*Correspondence to:

Clyde Francks D.Phil.

Max Planck Institute for Psycholinguistics

Wundtlaan 1, Nijmegen 6525XD

The Netherlands

Tel +31 24 35 21929

clyde.francks@mpi.nl

Conflicts of interest

The authors declare no conflicts of interest.

Abstract

Left-handedness is a complex trait which might sometimes involve rare, monogenic contributions. *Situs inversus* (SI) of the visceral organs can occur with Primary Ciliary Dyskinesia (PCD), due to mutations which affect left-right axis formation. Roughly 10% of people with SI and PCD are left-handed, similar to the general population. However, in non-PCD SI, the rate of left-handedness may be elevated. We sequenced the genomes of nine non-PCD SI people who show an elevated rate of left-handedness (five out of nine). We also sequenced six SI people with PCD as positive controls, and fifteen unaffected people as technical controls. Recessive mutations in known PCD genes were found in all positive controls with PCD. Of the nine non-PCD SI cases, two had recessive mutations in known PCD genes, suggesting reduced penetrance for PCD, and one had a recessive mutation in the non-PCD laterality gene *PKD1L1*. However, six of the nine non-PCD SI cases, including most of the left-handers, had no mutations in likely candidate genes, nor any significant biological pathways affected by their mutations. Therefore we did not identify a molecular link between visceral and brain laterality. While we cannot exclude a monogenic basis for non-PCD SI with left-handedness, multifactorial and non-genetic models must also be considered.

Keywords: *Situs inversus*, primary ciliary dyskinesia, left-handedness, whole genome sequencing

Introduction

A fundamental question in human neurobiology is how the brain becomes functionally left-right asymmetrical. For example, approximately 90% of people are right-handed and have left-hemisphere dominance for language, among other lateralized functions, but the developmental basis for this asymmetry remains unknown[1]. One possibility is that early embryonic mechanisms which give rise to asymmetries of the visceral organs also impact on brain asymmetries. However, this has not been previously addressed by genetic mutation screening in people who are both left-handed and have altered forms of visceral laterality.

Roughly 1:6,000-8,000 people have *situs inversus* (SI), a mirror reversal of the normal asymmetrical arrangement of the viscera [2,3]. SI can occur alone or in combination with Primary Ciliary Dyskinesia (PCD), a recessive genetic disorder which involves mutations that disrupt motile cilia [2]. Cilia are hair-like organelles that protrude from the cell surface into the extracellular space [4]. They are expressed in various tissues including the respiratory epithelium [5], so that a disruption of ciliary motility can cause symptoms such as chronic bronchitis, inflamed or infected sinuses [6], which are often present in PCD.

Motile cilia are also expressed early in development, within an embryonic structure called the 'node', where they generate a leftward fluid flow that helps to create the left-right body axis [5]. The leftward direction of the nodal flow may be explained by a posterior tilt of the cilia together with their clockwise rotation, arising ultimately from molecular chirality of their component proteins [7-9]. When leftward nodal flow is absent due to recessive PCD-causing mutations, many of which affect component proteins of the ciliary cytoskeleton, it becomes a matter of chance whether the viscera will take up the normal or mirror-reversed positioning [10].

However, roughly three quarters of people with SI do not have PCD [2], and the causes of their SI remain largely unknown. A few genes have been reported to be involved in SI without PCD, and some of these can also cause partial disruptions of visceral laterality, known as heterotaxy or *situs ambiguus* (SA), including *ZIC3* [MIM:300265] [11], *CCDC11* [MIM:614759] [12], *WDR16*

[MIM:609804] [13], *NME7* [MIM:613465] [14], and *PKD1L1* [MIM:609721] [15]. The mechanisms
by which these non-PCD, SI-causing genes influence visceral laterality are not well understood,
but most code for cilia-associated proteins, rather than coding directly for cytoskeletal
components of cilia.

Intriguingly, the general population rate of 85-90% right-handedness is not altered in SI people
with PCD [16,17]. This implies a developmental dissociation between brain laterality for hand
motor control and nodal-ciliary visceral patterning. However, in the only study of handedness
in SI to include non-PCD cases, Vingerhoets *et al*. reported that five of nine SI cases without
PCD were left-handed [18]. Although based on a small sample, this suggests developmental
mechanisms which might indeed link handedness and visceral laterality, but independently of
genes involved in PCD.

One study reported a possible genetic link between a continuous measure of left-versus-right
hand motor skill and genes involved in visceral laterality [19], based on analyzing genetic variants
which are common in the population. However, the sample size of under 3000 subjects was low
for complex-trait genome-wide association analysis using common genetic variants. A much
larger study of over 300,000 subjects from the UK Biobank found no support for a link of left-
handedness to genes involved in visceral asymmetry [20]. The same large study identified an
association of a common variant at the *MAP2* gene with left-handedness, with a very small
effect [20]. Left-handedness has a heritability of roughly 25% based on twin and family data [21],
but only around 2% based on genome-wide genotype data for common polymorphisms, within
the UK Biobank dataset [22].

It has been proposed that left-handedness may sometimes occur due to genetic mutations which
are rare in the population, but might have substantial effects on brain laterality when present
[23,24]. Rare genetic effects are not well captured in genome-wide association studies, which are
focused on relatively common variation [20]. Here we performed an exploratory whole genome
sequencing study in the same set of 15 SI subjects studied by Vingerhoets *et al*., as well as 15
healthy controls matched for age, sex, education, and handedness (**Table 1**). The goal was to
identify rare, highly penetrant mutations in the nine people with non-PCD SI, as they show an
elevated rate of left-handedness. This approach had the potential to yield novel insights into the
developmental origins of left-right patterning of both brain and body.

Methods

Participants

Fifteen people with radiologically documented SI, and 15 controls with normal *situs* matched
for age, sex, education and handedness, were recruited from Ghent University Hospital and
Middelheim Hospital Antwerp (approved by Research Ethics Committee). Table 1 gives an
overview of the participants and their characteristics. All participants gave informed consent
for DNA sample collection and genomic analysis in relation to body, brain and behavioural
laterality. All methods were performed in accordance with the relevant guidelines and
regulations.

Details about the recruitment, diagnosis and selection/exclusion procedure have been described
previously [18]. Radiological information (RX or CT) of thorax and complete abdomen was
available in eight SI participants, and of thorax and upper abdomen in seven SI participants.
The medical reports confirmed that all 15 participants presented with radiologically
documented SI.

In five participants with SI, a formal diagnosis of PCD or Kartagener syndrome was found in their medical records. Kartagener syndrome refers to the clinical triad of situs inversus, chronic sinusitis, and bronchiectasis[10]. A sixth SI-participant was identified on account of a radiological consultation regarding infertility. The participant also presented with chronic sinusitis, mild chronic bronchitis. Although no formal diagnosis of PCD was obtained in this case, the presence of chronic upper and lower respiratory infection and infertility in an individual with SI warrants suspicion of Kartagener syndrome. Consequently, we included the participant within the PCD group. In addition, all six members of the PCD group reported having a daily wet cough, and had PICADAR scores of between 8 and 12 [25], and thus predictive probabilities of having PCD of between 66% and 99%, based on this recently-developed, questionnaire-based tool (note that five of these subjects anyway had formal medical diagnoses).

A seventh SI subject (SI03) had no medical record pertaining to PCD but did report daily wet cough. This subject had a PICADAR score of 8. In the study of Vingerhoets *et al.* this SI subject was classified as non-PCD, before the PICADAR assessment was available. For the purposes of the present study, we also treated this subject as non-PCD given the lack of formal medical history of PCD and the ambiguous PICADAR score, but we repeated some genetic analyses having excluded the person from the non-PCD group, in order to account for this uncertainty (see below).

Eight other SI subjects had no medical record of PCD or PCD-like symptoms, and were classified as non-PCD SI cases. Six of these reported no daily wet cough, and two did not answer in this regard. PICADAR scores can only be calculated in the presence of daily wet cough, so that none of these eight cases received PICADAR scores. Three of these cases had been previously diagnosed with congenital heart disease that required surgical treatment, and their radiological files all referred to their cardiac condition. Congenital heart disease is a frequent comorbidity of SI, as the cardiac circulation appears particularly sensitive to perturbation in normal left-right positional information [26].

Handedness was assessed using the Edinburgh Handedness Inventory (EHI) [27]. Note that one non-PCD SI subject reported being forced to switch from left- to right-handedness in childhood, in which case five out of nine non-PCD SI cases were naturally left-handed. One of the six cases with PCD also reported forced left-to-right switching, otherwise the rest were right-handed (**Table 1**).

Whole Genome Sequencing (WGS) and Pre-processing

DNA was extracted from saliva samples using the Oragene kit (Oragene). WGS was performed by Novogene (Hong Kong) using Illumina high throughput sequencing (HiSeq-PE150), creating paired end reads with a length of 150 base pairs (bp). Raw reads, stored in BAM files, were aligned to the human reference genome (the extended "decoy" version of b37) using Burrows-Wheeler Aligner (BWA) software [28], and sorted and reordered using SAMtools (v1.3.1) [29]. PCR duplicates, which could arise during cluster amplification, were marked using Picard (v2.9.0). Genome Analysis Toolkit (GATK v3.7) [30] software was used to realign reads around insertions/deletions (indels) and to recalibrate base quality scores per sample.

Variant Calling and Quality Control

Indels (insertion/deletions) and single nucleotide polymorphisms (SNPs) were called as recommended by the GATK Best Practices workflow. Variant Quality Score Recalibration (VQSR) was performed, and variants with a high probability of being false positive were flagged according to their sensitivity tranche (90, 99, 99.9 and 100). All SNPs and indels within a VQSR tranche of 99.9% or higher were discarded. Variants with a quality depth ≤ 9 or a call rate ≤ 0.8 were also removed. Vcftools v1.13 was used to create a summary table in the Variant

139 Call Format (vcf). A total of 13,989,941 SNPs and indels were identified across the 30 subjects,
140 with an average number of 5,186,055 (min= 5,053,188, max = 5,272,561) alternative alleles
141 per subject (i.e. different to the reference genome, build37 decoy version).

Genomic-level evaluation

For overall descriptive analysis of the participant genomes, a subset of 40,387 variants
distributed genome-wide was used that had known Minor Allele Frequencies (MAFs) > 0.1,
and were the result of pruning to be in low linkage disequilibrium with one another. For this,
the flag --indep-pairwise function in Plink (v.1.90b3w) [31] was applied with a pairwise linkage
disequilibrium r^2 greater than 0.2, based on a SNP-SNP correlation matrix of 1500 by 150 in
window size. The resulting data were then used as the basis for inferring pairwise identity by
descent (IBD) sharing between subjects (i.e. genetic relatedness), inbreeding, and
inconsistencies with reported sex, using the Plink operations --genome, --ibc/het, and --check-
sex.

A different subset of 61,467 independent variants distributed genome-wide was used for
visualizing population structure through Multi-Dimensional Scaling (MDS) analysis with
respect to the 1000 genomes reference dataset (v37) [32] (downloaded from:
ftp://climb.genomics.cn/pub/10.5524/100001_101000/100116/1kg_phase1_all.tar.gz) using
the Plink operation –mds-plot.

Annotation and filtering of single nucleotide variants affecting protein sequences

Variants were normalized using the software tool *vt normalize* (v0.5772-60f436c3) [33], which
ensures consistent representation of variants across the genome. Normalized SNVs were
annotated using Annovar [34] and Variant Effect Predictor (v88) [35]. Gemini (v 0.20.0) was used
to select protein coding variants with 'MEDIUM' or 'HIGH' impact severity annotations, as
well as non-coding variants with 'HIGH' impact severity annotations (in practice those altering
splice donor or acceptor sites). Additional filtering was done in R and comprised the removal
of synonymous variants, and of 'MEDIUM' variants with a PolyPhen [36] or Sift [37] prediction
score of "benign" or "tolerant" respectively. Data were then processed and analyzed separately
under recessive and dominant models:

Recessive model

For the recessive model we further excluded variants with a known population frequency >
0.005 in any of the following databases: GNOMAD [38], ESP [39], 1000 Genomes [32] and ExAC
databases [38]. The remaining low-frequency variants were considered as putative mutations.
Gene-level mutation counts per subject were then made, with a given gene being assigned as
recessively mutated when it carried two copies of the same mutation (homozygous) or two
different mutations (possible compound heterozygous). Integrative Genome Viewer (IGV
v2.3.97) [40] was used to visualize the possible compound heterozygous mutations, and genes
carrying these were discarded when both mutations were definitely present on the same allele
(i.e. "in phase") on a given sequence read. Finally, genes recessively mutated according to these
criteria in any of the fifteen unaffected control subjects were excluded as being potentially
causative in cases, and also removed for the purposes of Gene Set Enrichment Analysis (GSEA;
see below): this step had the advantage of removing false variants from potential technical
artifacts, or variants which are common in the population but not annotated as such in on-line
databases. In total, 17 genes were excluded based on overlap with the unaffected control
subjects. Genes on chromosome X were processed as part of the recessive pipeline, such that
females would need to carry two mutations in a given gene, and males one mutation.

Dominant model

185 A maximum population frequency threshold of 5×10^{-5} was applied in this case, and genes
186 carrying at least one rare variant according to this criterion were considered as potentially
187 causative. Again, genes mutated according to this criterion in any of the fifteen unaffected
188 control subjects (N = 47 genes) were excluded as being potentially causative in cases, and
189 removed for the purposes of GSEA analysis (below).

Gene Set Enrichment Analysis

To test whether a list of mutated genes in a given set of subjects contained functionally related
genes across subjects, we performed Gene Set Enrichment Analysis (GSEA) using the
g:Profiler R package (version 0.6.1) [41]. A gene classification scheme derived from the Gene
Ontology (GO) database [42,43] was used. This specified a total of 6380 functionally defined gene-
sets, based on a background of 19,327 genes with functional annotations. Many genes are not
represented in the GO, particularly mRNA transcripts of no known function or homology, so
that the numbers of mutated genes in a given set of subjects was always higher than the subset
used as input for GSEA.

Mutated genes on chromosome X were combined with recessively mutated autosomal genes
for the purposes of GSEA. GSEAs were performed separately for the mutated gene lists in SI
subjects with PCD (54 genes, of which 40 are in the GO), non-PCD SI subjects (60 genes, 38
in the GO), left handed non-PCD SI subjects (42 genes, 26 in the GO), and cases that were
unsolved under the recessive model (42 genes, 22 in the GO). These analyses were also repeated
after excluding subject SI03 (see above and **Table 1**), in the non-PCD SI group (55 genes, 35
in the GO), the left handed non-PCD SI group (37 genes, 23 in the GO), and in the unsolved
group (36 genes, 19 in the GO).

For the dominant model, GSEA was performed separately for mutated genes in the non-PCD
SI subjects (330 genes, 271 in the GO), the subset of non-PCD SI subjects that remained
unsolved under the recessive model (217 genes, 175 in the GO), and the left handed subset of
non-PCD subjects (201 genes, 163 in the GO). PCD subjects were not tested under a dominant
model, as PCD is known to be a recessive phenotype. Dominant analyses were repeated after
excluding subject SI03, for non-PCD SI subjects (285 genes, 235 in the GO), unsolved non-
PCD SI subjects (170 genes, 139 in the GO) and non-PCD left handed SI subjects (156 genes,
127 in the GO).

In order to confirm that unaffected controls showed no significant pathway enrichment among
their mutated genes, a mirrored exclusion was performed whereby any genes mutated in cases
were excluded from the control gene lists. This resulted in 56 genes exclusively mutated in
controls for the recessive model (of which 34 genes are in the GO), and 533 genes exclusively
mutated in controls under the dominant model (440 in the GO).

The identities of genes were first converted using the g:Convert tool [41] to ensure recognition by
the GO schema. The following settings were then used for GSEA: minimum set size = 15,
maximum set size = 500, minimum intersect number = 2, hierarchical filtering = moderate. P-
values were corrected for multiple testing across gene sets, based on the gSCS correction
method in g:Profiler, separately for each input list of mutated genes corresponding to a given
set of subjects. This method of multiple testing correction takes into account the hierarchical
structure of the sets [41]. We applied a cut-off P value of adjusted 0.01.

Candidate Gene Lists

We queried the genetic data with respect to candidate gene lists for some purposes (see Results).
An initial list of candidate genes was created in R (v3.3), based on searching the Online
Mendelian Inheritance in Man (OMIM) database [44] for the terms: "situs inversus", "heterotaxy
| heterotaxia | situs ambiguus", "congenital heart disease", "PCD | ciliary dyskinesia |

Kartagener", "left-right", and "asymmetry | laterality". Additionally, the Clinvar column of our
annotated variant data, which contains information based on the Clinvar database [45], was
searched for these terms, and genes that were not yet in the initial list of candidate genes were
accordingly included.

A broader candidate gene list was also created which included genes from the literature that
were suggested to be associated with human laterality phenotypes and/or ciliopathies, but not
(yet) present in our initial list of candidate genes (**Supplementary Table S1**). Specifically, a
list of ciliopathy genes from a review of this topic [6] was searched for the terms: "PCD",
"heterotaxy", "situs", "left-right", "asymmetry" or "laterality", yielding 18 additional candidate
genes. Additionally, a list of genes potentially associated with human laterality disorders, as
compiled in a 2015 review [2], resulted in the addition of 25 candidate genes. Eleven additional
genes were included, of which three were reported as potentially associated with PCD or
heterotaxy [46], three had potential associations with non-syndromic heterotaxy [47], and five were
considered as possibly causal in a recent exome sequencing study of various laterality defects
[48]. Finally, 39 more genes – arising from a search for the 'situs inversus' phenotype - were
retrieved from the Mouse Genome Database (MGD) [49] at the Mouse Genome Informatics
website, the Jackson laboratory, Bar Harbor, Maine. (URL: http://www.informatics.jax.org)
[Oct, 2017].

Structural Variants

For all participant genomes, structural Variants (SVs) (>50 kilobases in length) were called
using a combination of two SV calling algorithms: CNVnator (v0.3.3) [50] and Lumpy (v 0.2.13)
[51]. These algorithms complement each other regarding the kinds of signals in WGS data that
they can detect, as CNVnator is based on read-depth, whereas Lumpy is based on paired-end
mapping [52]. For CNVnator the bin size was set to 100 base pairs for all genomes except for two,
where it was 150 base pairs (we determined a roughly optimal bin size for each subject's
genome, such that that the average read depth for that genome would be at least 4 standard
deviations from zero). Lumpy was run using default parameters in "lumpyexpress".

SVs were then annotated using SV2 [53]. Variants that were present in any of the 15 healthy
controls were removed from consideration as potentially causative for SI. Only variants that
passed the default SV2 filtering criteria for quality were included [53].

Results

Protein-altering single nucleotide variants

Recessive mutations

Our variant calling, filtering and annotation pipeline produced between 5 and 15 recessively
mutated genes per SI subject. We included six SI subjects with PCD as positive controls, in
order to ensure that the variant calling and mutation definition criteria were well calibrated. As
PCD is known to be a recessive phenotype for which at least 30 different genes have already
been identified [2], we expected most, or all, of these six subjects to have identifiable mutations
in known PCD-causing genes. As expected, each of these six cases had just one recessively
mutated gene which was annotated 'Kartagener' or 'PCD' in the Clinvar database [45], and was
therefore the most likely monogenic cause for their condition (**Table 1**). These genes were
LRRC6 [MIM:614930], *DNAH11* [MIM:603339], *DNAAF1* [MIM:613190], *CCDC114*
[MIM:615038], and *DNAH5* [MIM: 603335] (the latter mutated in two SI cases with PCD,
consistent with *DNAH5* being the most common cause of PCD in European-ancestry
populations [54]) (**Table 1**). The PCD subject with a homozygous *LRRC6* mutation (subject SI06)

was the only individual to show an elevated inbreeding coefficient and non-European ancestry
(**Supplementary Table S2, Supplementary Figure S1**).

None of the fifteen unaffected control subjects had any recessively mutated genes annotated
'Kartagener', 'PCD', 'SA' or 'SI' in Clinvar.

Surprisingly, two of the nine non-PCD SI cases had recessive mutations in genes annotated
'Kartagener' in the Clinvar database [45]. These were subject SI12 (again involving mutations in
DNAH5), and subject SI16 (in *CCDC151*) (**Table 1**). As these subjects had no medical records
pertaining to PCD-like symptoms, and no daily wet cough, then the findings suggest reduced
penetrance for PCD.

One of the nine non-PCD subjects – i.e., SI02 - had a recessive mutation in a gene that is
annotated in Clinvar as '*situs ambiguus*' and '*situs inversus totalis*', but not annotated as PCD-
causing (**Table 1**). This gene is *PKD1L1* [MIM: 617205]. A homozygous missense mutation in
PKD1L1 was previously reported in an individual with SI and congenital heart disease but no
PCD, as well as recessive splicing mutations in two individuals with heterotaxy [15]. Our subject
SI02 had no diagnosis of congenital heart disease (CHD) (**Table 1**). As *PKD1L1* is a known
recessive cause of non-PCD SI, we consider this gene to be the most likely cause of the non-
PCD SI in subject SI02.

Subject SI03, who had no formal medical history of PCD, but had reported an intermediate
PICADAR score, showed no recessive mutations in known PCD genes, which tends to support
non-PCD status for this subject. This meant that six non-PCD SI cases did not have recessive
mutations in genes known to cause human laterality disorders, as annotated in Clinvar, and
therefore remained 'unsolved' under a recessive model (**Table 1**). Among these six non-PCD
SI cases, four were left-handed, and were therefore of primary interest for the present study.
Three of these same subjects also had CHD (**Table 1**).

We constructed an extended list of known or suspected laterality genes with reference to the
literature and mouse laterality phenotypes (Methods; **Supplementary Table S1**), but none of
these genes had recessive mutations in the six unsolved non-PCD SIT subjects.

We note possible compound heterozygous mutations in *PKD1* [MIM:601313], as a paralogue
of *PKD1L1* [MIM:609721], in subject SI14 (**Supplementary Table S3**). However, mutations
in this gene would be expected to cause autosomal dominant Polycystic Kidney Disease [55,56],
and since there was no such diagnostic record for this subject, one or both of these specific
mutations probably has limited functional impact and is therefore unlikely to be a monogenic
cause for SI either.

KIF13B [MIM:607350] was putatively recessively mutated in subjects SI14 (unsolved) and
SI12 (solved) (**Supplementary Table S3**). Although no literature has linked *KIF13B* to PCD
or laterality phenotypes, *KIF3A* [MIM:604683], another kinesin encoding gene, is known to be
involved in LR determination [57]. Moreover, *KIF3B* [MIM: 603754] is known to affect motility
of nodal cilia, accordingly affecting LR determination [58]. Together with dyneins, kinesins allow
ciliary proteins to enter the organelle via the transition zone by transporting them as cargo [59,60],
and accordingly play an important role in ciliary construction and maintenance [59,60]. The
mutations in *KIF13B* might therefore potentially cause SI without PCD in subject SI14, and
perhaps also affect the phenotypic outcome in subject SI12 who has likely causal mutations in
DNAH5, but we cannot confidently assign a role to *KIF13B* on the basis of this evidence.

Dominant mutations

For the six non-PCD SI cases who remained unsolved under a recessive model, we also
considered a dominant model using a maximum known mutation frequency of 5×10^{-5}, and again

cross-referenced the mutated genes against Clinvar and our extended candidate gene list (**Supplementary Table S1**), but no likely causative genes emerged (**Supplementary Table S3**) (see Discussion).

Subject SI05 showed a heterozygous mutation in *LRRC6* [MIM:614930] (**Supplementary Table 3**), which was included among our candidate genes. However, since recessive – but not dominant - mutations in this gene have been associated with PCD [61], it is unlikely that this mutation is causal for non-PCD SI in this subject.

We also note a heterozygous mutation in *WDR62* [MIM:613583] in the unsolved case SI09 (**Supplementary Table 3**). Although mice with mutations in this gene have shown dextrocardia and right pulmonary isomerism (MGI:5437081) [49], humans with recessive *WDR62* mutations do not show altered laterality. Instead, they have shown infantile spasm, microcephaly and intellectual disability [62]. It is therefore unlikely that *WDR62* is a dominant cause of altered laterality in humans.

The gene *CEP290* [MIM:610142], mutated in subject SI14 (**Supplementary Table 3**), is linked to left sided isomerism in mice (MGI:5438068)[49]. In humans, recessive mutations have been linked to a variety of ciliopathies, ranging from nephronophthisis, retinal degeneration and Joubert syndrome, to Bardet-Biedl syndrome and Meckel-Grüber syndrome [63,64]. However, similar to the aforementioned genes, mutations in *CEP290* have not been associated with laterality phenotypes in humans, and we therefore consider this to be an unlikely cause of non-PCD SI.

Subject SI09 had a heterozygous mutation in PLXND1 (Supplementary Table 3), a gene which appears among search results for the phenotype 'situs inversus' within the Mouse Genome Database [49]. However, while PLXND1 is annotated as a cause of aortic arch and atrial abnormalities in this database, it is not annotated with situs inversus or heterotaxia, so that the basis for the search result is unclear. We did not find evidence for this gene's involvement in visceral laterality in a further literature search.

Gene set enrichment analysis

We first performed gene set enrichment analysis using the positive control set of six SI subjects with PCD. As noted above, the six PCD subjects had likely recessive monogenic causes in five different genes (two subjects had mutations in the same gene, *DNAH5*). As expected, from the total combined list of recessively or X-linked mutated genes in these subjects, of which 40 genes were included in the GO schema, we obtained significant results for various cilia-related pathways, such as 'axoneme' (p = 0.004), 'outer dynein arm assembly' (p = 3.85×10^{-5}), 'dynein complex' (p = 4.89×10^{-5}), and 'microtubule based movement' (p = 1.41×10^{-5}) (**Table 2**) (all P values adjusted for multiple testing across gene sets, see Methods). As expected, when the single most likely causative gene for each PCD subject was removed from the gene list, which left 36 recessively or X-linked mutated genes in these subjects that are present in the GO schema, there were no longer any significant enrichment terms, which further supports that the monogenic causes had been correctly identified in these subjects. The gene set enrichment analysis in the PCD subjects confirmed that, despite a relatively small number of subjects (i.e. six), the analysis was well powered to identify affected biological processes, even when most individual subjects had monogenic causes in different genes, and each subject had other, non-causative mutated genes.

With this in mind, we then performed gene set enrichment analysis in the non-PCD SI cases, who were of primary interest for the present study due to a potential link with left-handedness. However, no significant enrichments were found when testing the list of recessively or X-linked mutated genes in the nine subjects with non-PCD SI, of which 38 genes were included in the

GO schema. There was also no significant functional enrichment when testing the recessively
mutated or X-linked genes in the five left-handed subjects with non-PCD SI, of which 26 genes
were in the GO schema, and neither when testing the list of genes in the six unsolved non-PCD
SI subjects, of which 22 genes were present in the GO schema (**Table 2**). Repeating the analysis
after excluding subject SI03 from these subsets made no difference, there were still no
significant gene sets. In addition, gene-set enrichment analysis with dominantly mutated genes
as input did not produce significant results in the non-PCD SI cases, nor the left-handed or
unsolved subsets.

As expected, the lists of recessively/X-linked and dominantly mutated genes in the fifteen
unaffected control subjects did not produce any significant gene set enrichment terms (**Table
2**).

Structural variant analysis

We additionally screened the genomes of all subjects for structural variants (such as larger-
scale deletions and duplications), using two complementary algorithms (see Methods). SI
subjects had SVs affecting an average of 9.7 genes per subject (min = 2 SVs, max = 16 SVs),
and controls had SVs affecting an average of 9.6 genes per subject (min = 5 SVs, max = 16
SVs). None of the SI subjects, regardless of PCD status, had SVs affecting genes that were
annotated SI, PCD, Kartagener, *situs ambiguus* (SA), or Heterotaxy (HTX) in Clinvar, nor
affecting genes from our broader list of candidate laterality genes.

Discussion

In this study we aimed to identify rare, highly penetrant genetic mutations that might link
visceral body asymmetry with handedness, by analysing whole genome sequence data from
nine SI subjects without medical histories of PCD, five of whom were left-handed. We
additionally included six SI subjects with PCD as positive technical controls, and fifteen
unaffected subjects as negative controls.

Likely monogenic causes were identified in all positive controls, i.e. each of the six PCD
subjects had one recessively mutated gene (usually a different gene in each subject) that is
already known to cause PCD when mutated. The six PCD subjects also confirmed that
significant pathway enrichment could be detected on the basis of their mutated gene lists, as
multiple gene sets related to ciliary functions were detected. The fifteen unaffected control
subjects showed no recessive mutations in genes known to cause PCD or laterality phenotypes,
as expected.

Two of the nine non-PCD SI subjects had likely recessive monogenic causes in known PCD
genes. This may indicate reduced penetrance of these mutations for PCD-like symptoms such
as lung symptoms, although they can apparently affect *situs* in early development. One non-
PCD SI subject, who was right-handed, had a homozygous mutation in a gene already known
to cause SI without PCD, i.e. *PKD1L1* [15]. This gene encodes a component of a calcium channel
which is associated with non-motile cilia [65].

However, the six remaining non-PCD SI subjects had no obvious monogenic basis for their
condition, i.e. they did not have likely causative mutations in genes known to cause human
laterality disorders as annotated in the Clinvar database, nor within an extended list of known
or suspected laterality genes based on literature searching and mouse phenotypes. Among the
six non-PCD SI subjects, four were left-handed, and therefore comprised the bulk of left-
handers in the dataset. Furthermore, gene set enrichment analysis of their mutated genes did not
identify significant biological pathways, in either the whole set, or the left-handed subset, or

the subset that was 'unsolved' by recessive monogenic causes. In other words, the biology of their non-PCD SI could not be linked via the mutations that they carried. Finally, we also considered larger genomic rearrangements known as Copy Number Variants (CNVs), but no obvious candidate genes emerged.

A monogenic model is still possible for the majority of the non-PCD SI cases, and/or for the left-handed subset specifically, but would have to involve genetic heterogeneity across a set of genes which are not currently linked in terms of their known biology, at least to an extent which would have been discernible in this dataset, as it was for the PCD subjects. We did not therefore identify a genetic-developmental pathway that links handedness and visceral asymmetry in this study. The question of whether, and how, functional brain laterality is linked developmentally to visceral laterality in humans remains unanswered [20].

Genetic contributions to non-PCD SI and left-handedness might also involve non-coding variation, or rare combinations of multiple common variants. The noncoding genome comprises 98% of the genome, but interpreting the variation within these regions is challenging. Several attempts have been made to rank potentially causative variants across the genome based on scores that integrate different types of information, including conservation of DNA sequence, regulatory information [66], and population genomic data [67-72]. However, these ranking approaches are currently not very concordant with each other [73]. For the present study we did not address these possibilities, which must await larger sample sizes and an improved understanding of the role of rare, non-coding mutations in phenotypic variation.

In utero environmental effects such as prenatal drug exposure might also affect left-right determination of body and brain [74]. Handedness itself is known to associate with various early life factors including birthweight and breastfeeding, although not to a degree which is remotely predictive at the individual level [75]. As noted in the introduction, left-handedness has a modest heritability of roughly 25% in family and twin studies, and lower in SNP-based heritability studies. Regardless, the primary causes of this trait remain unknown.

In this study there was a degree of diagnostic uncertainty as regards the PCD status of some SI subjects. However, it was not the purpose of the present study to achieve a clinical diagnosis of the presence or absence of PCD, nor to confirm already-known PCD genes. In this context we did not, for example, sequence the mutations in the PCD subjects by another technique for validation, nor confirm allelic phase in the compound heterozygotes. Rather, we were concerned to identify potentially causative mutations in the nine SI subjects without medical histories of PCD who show an elevated rate of left-handedness, with the potential to yield important basic insights into body and brain laterality. If we had found clear candidate mutations in left-handed members of the non-PCD SI group, in genes not previously linked to PCD, then further validation and functional characterisation of those mutations would have been appropriate, but this did not arise.

Regardless of the PCD status of any individual SI subject in this study, the overall pattern of results was clearly different between the PCD and non-PCD groups, where all six positive control subjects in the PCD group had mutations in known PCD-causing genes, while six out of nine in the non-PCD group had no obvious monogenic mutations, among whom were most of the left-handers. Also, the pathway analyses in various different subsets of the non-PCD group showed a consistent lack of significant enrichment, whereas clear signals emerged from the PCD group, which further supports an overall distinction of the groups. Nonetheless, further detailed investigation of subjects SI03, SI12 and SI16 with PCD diagnostic tools might reveal lung and other ciliary symptoms to an extent [76].

Although ciliary-induced nodal flow plays a crucial role in symmetry breaking, some organisms, such as chicks and fruit flies (*Drosophilia melanogaster*), do not have motile nodal

cilia, yet they do show asymmetrical organs [77]. For example, left-right asymmetry in chicks involves rearranging the relative orientations of cells expressing critical genes at the node, in a non-ciliary-dependent manner [78]. Furthermore, left-right asymmetry of the *Drosophilia* gut is established by intracellular cytoskeletal organization that may give rise to cellular shape chirality, by means of unidirectional tilting of radial fibers, and anti-clockwise swirling of transverse fibers [79]. Whether such mechanisms also influence left-right organ asymmetry in mammals is unclear. In the present study we saw no mutations in the homologues of two genes implicated in cellular chirality in *Drosophilia*, *MYO1D* or *MYO1C* [80], nor in the homologues of two genes thought to affect laterality in chicks and frogs, *SLC6A4* and *SLC18A2* [81]

Future studies in larger human cohorts may help to identify genetic contributions to non-PCD SI and left-handedness in some cases. Candidate biological pathways which emerge from research on non-mammalian mechanisms of asymmetry development should be considered in future studies.

Data availability

Requests to access the genomic datasets generated for the current study will be considered in relation to the consents, relevant rules and regulations, and can be made via the corresponding author.

References

Francks, C. Exploring human brain lateralization with molecular genetics and genomics. *Ann N Y Acad Sci* **1359**, 1-13, doi:10.1111/nyas.12770 (2015).

Deng, H., Xia, H. & Deng, S. Genetic basis of human left-right asymmetry disorders. *Expert Rev Mol Med* **16**, e19, doi:10.1017/erm.2014.22 (2015).

Arnold, G. L., Bixler, D. & Girod, D. Probable autosomal recessive inheritance of polysplenia, situs inversus and cardiac defects in an Amish family. *American journal of medical genetics* **16**, 35-42, doi:10.1002/ajmg.1320160107 (1983).

Mirvis, M., Stearns, T. & James Nelson, W. Cilium structure, assembly, and disassembly regulated by the cytoskeleton. *The Biochemical journal* **475**, 2329-2353, doi:10.1042/bcj20170453 (2018).

Fliegauf, M., Benzing, T. & Omran, H. When cilia go bad: cilia defects and ciliopathies. *Nature reviews. Molecular cell biology* **8**, 880-893, doi:10.1038/nrm2278 (2007).

Reiter, J. F. & Leroux, M. R. Genes and molecular pathways underpinning ciliopathies. *Nature reviews. Molecular cell biology* **18**, 533-547, doi:10.1038/nrm.2017.60 (2017).

Okada, Y., Takeda, S., Tanaka, Y., Belmonte, J. I. & Hirokawa, N. Mechanism of nodal flow: a conserved symmetry breaking event in left-right axis determination. *Cell* **121**, 633-644, doi:10.1016/j.cell.2005.04.008 (2005).

Cartwright, J. H., Piro, O. & Tuval, I. Fluid-dynamical basis of the embryonic development of left-right asymmetry in vertebrates. *Proc Natl Acad Sci U S A* **101**, 7234-7239, doi:10.1073/pnas.0402001101 (2004).

Hilfinger, A. & Julicher, F. The chirality of ciliary beats. *Physical biology* **5**, 016003, doi:10.1088/1478-3975/5/1/016003 (2008).

Leigh, M. W. *et al.* Clinical and genetic aspects of primary ciliary dyskinesia/Kartagener syndrome. *Genet Med* **11**, 473-487, doi:10.1097/GIM.0b013e3181a53562 (2009).

Ware, S. M. *et al.* Identification and functional analysis of ZIC3 mutations in heterotaxy and related congenital heart defects. *Am J Hum Genet* **74**, 93-105, doi:10.1086/380998 (2004).

Gur, M. *et al.* Roles of the cilium-associated gene CCDC11 in left-right patterning and in laterality disorders in humans. *The International journal of developmental biology* **61**, 267-276, doi:10.1387/ijdb.160442yc (2017).

Ta-Shma, A. *et al.* A human laterality disorder associated with a homozygous WDR16 deletion. *European journal of human genetics : EJHG* **23**, 1262-1265, doi:10.1038/ejhg.2014.265 (2015).

| 515 | 14 | Reish, O. *et al.* A Homozygous Nme7 Mutation Is Associated with Situs Inversus Totalis. *Human mutation* **37**, 727-731, doi:10.1002/humu.22998 (2016). |

14 Reish, O. *et al.* A Homozygous Nme7 Mutation Is Associated with Situs Inversus Totalis.
Human mutation **37**, 727-731, doi:10.1002/humu.22998 (2016).
15 Vetrini, F. *et al.* Bi-allelic Mutations in PKD1L1 Are Associated with Laterality Defects in
Humans. *Am J Hum Genet* **99**, 886-893, doi:10.1016/j.ajhg.2016.07.011 (2016).
16 McManus, I. C., Martin, N., Stubbings, G. F., Chung, E. M. & Mitchison, H. M. Handedness
and situs inversus in primary ciliary dyskinesia. *Proc Biol Sci* **271**, 2579-2582,
doi:10.1098/rspb.2004.2881 (2004).
17 Afzelius, B. A. & Stenram, U. Prevalence and genetics of immotile-cilia syndrome and left-
handedness. *The International journal of developmental biology* **50**, 571-573,
doi:10.1387/ijdb.052132ba (2006).
18 Vingerhoets, G. *et al.* Brain structural and functional asymmetry in human situs inversus totalis.
Brain Struct Funct, doi:10.1007/s00429-017-1598-5 (2018).
19 Brandler, W. M. *et al.* Common variants in left/right asymmetry genes and pathways are
associated with relative hand skill. *PLoS genetics* **9**, e1003751,
doi:10.1371/journal.pgen.1003751 (2013).
20 de Kovel, C. G. F. & Francks, C. The molecular genetics of hand preference revisited. *bioRxiv*,
447177, doi:10.1101/447177 (2018).
21 Medland, S. E. *et al.* Genetic influences on handedness: data from 25,732 Australian and Dutch
twin families. *Neuropsychologia* **47**, 330-337, doi:10.1016/j.neuropsychologia.2008.09.005
(2009).
22 Neale, B. Heritability of >2,000 traits and disorders in the UK Biobank. (2017).
23 Armour, J. A., Davison, A. & McManus, I. C. Genome-wide association study of handedness
excludes simple genetic models. *Heredity* **112**, 221-225, doi:10.1038/hdy.2013.93 (2014).
24 Kavaklioglu, T., Ajmal, M., Hameed, A. & Francks, C. Whole exome sequencing for
handedness in a large and highly consanguineous family. *Neuropsychologia* **93**, 342-349,
doi:10.1016/j.neuropsychologia.2015.11.010 (2016).
25 Behan, L. *et al.* PICADAR: a diagnostic predictive tool for primary ciliary dyskinesia. *The
European respiratory journal* **47**, 1103-1112, doi:10.1183/13993003.01551-2015 (2016).
26 Kosaki, K. & Casey, B. Genetics of human left-right axis malformations. *Seminars in cell &
developmental biology* **9**, 89-99, doi:10.1006/scdb.1997.0187 (1998).
27 Oldfield, R. C. The assessment and analysis of handedness: the Edinburgh inventory.
Neuropsychologia **9**, 97-113 (1971).
28 Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).
29 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
2079, doi:10.1093/bioinformatics/btp352 (2009).
30 McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome research* **20**, 1297-1303, doi:10.1101/gr.107524.110
(2010).
31 Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage
analyses. *Am J Hum Genet* **81**, 559-575, doi:10.1086/519795 (2007).
32 Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68-74,
doi:10.1038/nature15393 (2015).
33 Tan, A., Abecasis, G. R. & Kang, H. M. Unified representation of genetic variants.
Bioinformatics **31**, 2202-2204, doi:10.1093/bioinformatics/btv112 (2015).
34 Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants from
high-throughput sequencing data. *Nucleic acids research* **38**, e164, doi:10.1093/nar/gkq603
(2010).
35 McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome biology* **17**, 122,
doi:10.1186/s13059-016-0974-4 (2016).
36 Adzhubei, I. A. *et al.* A method and server for predicting damaging missense mutations. *Nature
methods* **7**, 248-249, doi:10.1038/nmeth0410-248 (2010).
37 Kumar, P., Henikoff, S. & Ng, P. C. Predicting the effects of coding non-synonymous variants
on protein function using the SIFT algorithm. *Nature protocols* **4**, 1073-1081,
doi:10.1038/nprot.2009.86 (2009).

38 Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285-
291, doi:10.1038/nature19057 (2016).
39 Fu, W. *et al.* Analysis of 6,515 exomes reveals the recent origin of most human protein-coding
variants. *Nature* **493**, 216-220, doi:10.1038/nature11690 (2013).
40 Robinson, J. T. *et al.* Integrative genomics viewer. *Nature biotechnology* **29**, 24-26,
doi:10.1038/nbt.1754 (2011).
41 Reimand, J. *et al.* g:Profiler-a web server for functional interpretation of gene lists (2016
update). *Nucleic acids research* **44**, W83-89, doi:10.1093/nar/gkw199 (2016).
42 Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. *Nature genetics* **25**, 25-29, doi:10.1038/75556 (2000).
43 Ashburner, M. *et al.* Expansion of the Gene Ontology knowledgebase and resources. *Nucleic
acids research* **45**, D331-d338, doi:10.1093/nar/gkw1108 (2017).
44 Amberger, J. S., Bocchini, C. A., Schiettecatte, F., Scott, A. F. & Hamosh, A. OMIM.org:
Online Mendelian Inheritance in Man (OMIM(R)), an online catalog of human genes and
genetic disorders. *Nucleic acids research* **43**, D789-798, doi:10.1093/nar/gku1205 (2015).
45 Landrum, M. J. *et al.* ClinVar: improving access to variant interpretations and supporting
evidence. *Nucleic acids research* **46**, D1062-d1067, doi:10.1093/nar/gkx1153 (2018).
46 Hagen, E. M. *et al.* Copy-number variant analysis of classic heterotaxy highlights the
importance of body patterning pathways. *Human genetics* **135**, 1355-1364, doi:10.1007/s00439-
016-1727-x (2016).
47 Peeters, H. & Devriendt, K. Human laterality disorders. *European journal of medical genetics*
49, 349-362, doi:10.1016/j.ejmg.2005.12.003 (2006).
48 Li, A. H. *et al.* Genetic architecture of laterality defects revealed by whole exome sequencing.
European journal of human genetics : EJHG, doi:10.1038/s41431-018-0307-z (2019).
49 Smith, C. L., Blake, J. A., Kadin, J. A., Richardson, J. E. & Bult, C. J. Mouse Genome Database
(MGD)-2018: knowledgebase for the laboratory mouse. *Nucleic acids research* **46**, D836-d842,
doi:10.1093/nar/gkx1006 (2018).
50 Abyzov, A., Urban, A. E., Snyder, M. & Gerstein, M. CNVnator: an approach to discover,
genotype, and characterize typical and atypical CNVs from family and population genome
sequencing. *Genome research* **21**, 974-984, doi:10.1101/gr.114876.110 (2011).
51 Layer, R. M., Chiang, C., Quinlan, A. R. & Hall, I. M. LUMPY: a probabilistic framework for
structural variant discovery. *Genome biology* **15**, R84, doi:10.1186/gb-2014-15-6-r84 (2014).
52 Pirooznia, M., Goes, F. S. & Zandi, P. P. Whole-genome CNV analysis: advances in
computational approaches. *Front Genet* **6**, 138, doi:10.3389/fgene.2015.00138 (2015).
53 Antaki, D., Brandler, W. M. & Sebat, J. SV2: Accurate Structural Variation Genotyping and De
Novo Mutation Detection from Whole Genomes. *Bioinformatics*,
doi:10.1093/bioinformatics/btx813 (2017).
54 Hornef, N. *et al.* DNAH5 mutations are a common cause of primary ciliary dyskinesia with
outer dynein arm defects. *American journal of respiratory and critical care medicine* **174**, 120-
126, doi:10.1164/rccm.200601-084OC (2006).
55 Ranjzad, F. *et al.* Identification of Three Novel Frameshift Mutations in the PKD1 Gene in
Iranian Families with Autosomal Dominant Polycystic Kidney Disease Using Efficient
Targeted Next-Generation Sequencing. *Kidney & blood pressure research* **43**, 471-478,
doi:10.1159/000488471 (2018).
56 Xu, D. *et al.* Novel Mutations in the PKD1 and PKD2 Genes of Chinese Patients with
Autosomal Dominant Polycystic Kidney Disease. *Kidney & blood pressure research* **43**, 297-
309, doi:10.1159/000487899 (2018).
57 Hirokawa, N., Tanaka, Y. & Okada, Y. Left-right determination: involvement of molecular
motor KIF3, cilia, and nodal flow. *Cold Spring Harbor perspectives in biology* **1**, a000802,
doi:10.1101/cshperspect.a000802 (2009).
58 Nonaka, S. *et al.* Randomization of left-right asymmetry due to loss of nodal cilia generating
leftward flow of extraembryonic fluid in mice lacking KIF3B motor protein. *Cell* **95**, 829-837
(1998).
59 Goetz, S. C. & Anderson, K. V. The primary cilium: a signalling centre during vertebrate
development. *Nat Rev Genet* **11**, 331-344, doi:10.1038/nrg2774 (2010).

Verhey, K. J., Dishinger, J. & Kee, H. L. Kinesin motors and primary cilia. *Biochemical Society transactions* **39**, 1120-1125, doi:10.1042/bst0391120 (2011).

Horani, A. *et al.* LRRC6 mutation causes primary ciliary dyskinesia with dynein arm defects. *PLoS One* **8**, e59436, doi:10.1371/journal.pone.0059436 (2013).

Nardello, R. *et al.* A novel mutation of WDR62 gene associated with severe phenotype including infantile spasm, microcephaly, and intellectual disability. *Brain & development* **40**, 58-64, doi:10.1016/j.braindev.2017.07.003 (2018).

Zhang, Y. *et al.* BBS mutations modify phenotypic expression of CEP290-related ciliopathies. *Human molecular genetics* **23**, 40-51, doi:10.1093/hmg/ddt394 (2014).

Hildebrandt, F., Benzing, T. & Katsanis, N. Ciliopathies. *The New England journal of medicine* **364**, 1533-1543, doi:10.1056/NEJMra1010172 (2011).

Masyuk, A. I., Gradilone, S. A. & LaRusso, N. F. Calcium signaling in cilia and ciliary-mediated intracellular calcium signaling: are they independent or coordinated molecular events? *Hepatology (Baltimore, Md.)* **60**, 1783-1785, doi:10.1002/hep.27331 (2014).

Dunham, I. *et al.* An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57-74 (2012).

Kircher, M. *et al.* A general framework for estimating the relative pathogenicity of human genetic variants. *Nat. Genet.* **46**, 310-315 (2014).

Quang, D., Chen, Y. & Xie, X. DANN: a deep learning approach for annotating the pathogenicity of genetic variants. *Bioinformatics* **31**, 761-763 (2015).

Ritchie, G. R., Dunham, I., Zeggini, E. & Flicek, P. Functional annotation of noncoding sequence variants. *Nat. Methods* **11**, 294-296 (2014).

Jagadeesh, K. A. *et al.* M-CAP eliminates a majority of variants of uncertain significance in clinical exomes at high sensitivity. *Nat. Genet.* **48**, 1581-1586 (2016).

Dong, C. *et al.* Comparison and integration of deleteriousness prediction methods for nonsynonymous SNVs in whole exome sequencing studies. *Hum. Mol. Genet.* **24**, 2125-2137 (2015).

Ioannidis, N. M. *et al.* REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am. J. Hum. Genet.* **99**, 877-885 (2016).

Eilbeck, K., Quinlan, A. & Yandell, M. Settling the score: variant prioritization and Mendelian disease. *Nat. Rev. Genet.* (2017).

van Veenendaal, N. R., Kusters, C. D., Oostra, R. J., Bergman, J. E. & Cobben, J. M. When the right (Drug) should be left: Prenatal drug exposure and heterotaxy syndrome. *Birth defects research. Part A, Clinical and molecular teratology* **106**, 573-579, doi:10.1002/bdra.23497 (2016).

de Kovel, C. G. F., Carrion-Castillo, A. & Francks, C. A large-scale population study of early life factors influencing left-handedness. *bioRxiv*, 305425, doi:10.1101/305425 (2018).

Shapiro, A. J. *et al.* Diagnosis of Primary Ciliary Dyskinesia. An Official American Thoracic Society Clinical Practice Guideline. *American journal of respiratory and critical care medicine* **197**, e24-e39, doi:10.1164/rccm.201805-0819ST (2018).

Vandenberg, L. N. & Levin, M. A unified model for left-right asymmetry? Comparison and synthesis of molecular models of embryonic laterality. *Developmental biology* **379**, 1-15, doi:10.1016/j.ydbio.2013.03.021 (2013).

Gros, J., Feistel, K., Viebahn, C., Blum, M. & Tabin, C. J. Cell movements at Hensen's node establish left/right asymmetric gene expression in the chick. *Science* **324**, 941-944, doi:10.1126/science.1172478 (2009).

Tee, Y. H. *et al.* Cellular chirality arising from the self-organization of the actin cytoskeleton. *Nature cell biology* **17**, 445-457, doi:10.1038/ncb3137 (2015).

Lebreton, G. *et al.* Molecular to organismal chirality is induced by the conserved myosin 1D. *Science* **362**, 949-952, doi:10.1126/science.aat8642 (2018).

Fukumoto, T., Blakely, R. & Levin, M. Serotonin transporter function is an early step in left-right patterning in chick and frog embryos. *Developmental neuroscience* **27**, 349-363, doi:10.1159/000088451 (2005).

Acknowledgements

Thanks to all of the study participants. This research was funded by the Max Planck Society (Germany) and the Netherlands Organization for Scientific Research (NWO, grant No. 054-15-101) as part of the FLAG-ERA consortium project 'MULTI-LATERAL', a Partner Project to the European Union's Flagship Human Brain Project. The study was also funded by the Fonds Wetenschappelijk Onderzoek-Vlaanderen (FWO-grant n⁰ G.0114.16N) assigned to GV.

Author contributions

GV recruited and characterized the subjects. GV, SEF & CF conceived the study. MCP performed the genetic data analysis with support from ACC. MCP and CF wrote the manuscript. GV, ACC & SEF edited the manuscript. CF directed the study.

Additional information

The authors declare no competing interests.

Tables (see next page)

Table 1. The most likely causal recessive mutations in the fifteen SI subjects

Subj	SI group	Sex/Age	EHI	NH	CHD	Daily wet cough	Type	Gene	Clinvar	rs ID	Start position	ref	alt	MAF	AAC	impact
SI02	non-PCD	M/50	0.9	R	0	?	hom	*PKD1L1*	SA | SI	rs528302390	47870810	T	TCA	0.00109	-	splice donor
SI03	non-PCD	F/26	-0.8	L	0	yes		unsolved								
SI04	non-PCD	M/23	-1	L	1	no		unsolved								
SI05	non-PCD	M/27	0.9	R	0	no		unsolved								
SI07	non-PCD	F/35	0.9	R	1	?		unsolved								
SI09	non-PCD	F/36	0.7	L§	0	no		unsolved								
SI12	non-PCD	F/40	0.9	R	0	no	chet	*DNAH5*	Kartagener	None	13900415	A	C	-	E/*	stop gained
							chet	*DNAH5*	Kartagener	None	13769691	G	GC	0.00002	A/X	frameshift
SI14	non-PCD	M/18	-0.8	L	1	no		unsolved								
SI16	non-PCD	M/21	-1	L	0	no	hom	*CCDC151*	Kartagener	rs765121016	11533429	T	TCTC	0.00011	E/-	inframe deletion
SI06	PCD	M/46	1	R	0	yes	hom	*LRRC6*	Kartagener	rs767624733	133687728	T	C	0.00017	-	splice donor
SI08	PCD	F/23	0.9	R	0	yes	hom	*DNAH11*	PCD	rs373706559	21659620	A	C	0.00006	Y/*	stop gained
SI11	PCD	F/32	0.9	R	0	yes	chet	*DNAAF1*	Kartagener	rs569633512	84203963	C	T	-	-	splice donor
							chet	*DNAAF1*	Kartagener	rs373103805	84193302	G	C	0.00023	N/K	missense
SI13	PCD	M/48	0.6	L§	0	yes	hom	*CCDC114*	Kartagener	rs779459076	48814907	CACG	C	0.00093	-/R	Inframe insert
SI15	PCD	F/31	0.7	R	0	yes	chet	*DNAH5*	Kartagener	None	13786289	A	T	-	K/*	stop gained
							chet	*DNAH5*	Kartagener	rs397515540	13753397	G	GA	0.00034	D/X	frameshift
SI17	PCD	M/39	0.5	R	0	yes	chet	*DNAH5*	Kartagener	rs548521732	13839638	C	T	0.00021	-	splice acceptor
							chet	*DNAH5*	Kartagener	rs769458738	13753597	T	C	0.00004	R/H	missense

696 Subjects shown with two mutations have possible compound heterozygous mutations, otherwise they have homozygous mutations. The Genome Reference Consorium (GRC) build 37 decoy
697 version was used as reference sequence. EHI: Edinburgh Handedness Inventory score; NH: natural handedness; CHD: Congenital Heart Disease. Type: type of genetic mutation, i.e.,
698 homozygous (hom) or compound heterozygous (chet); MAF: minor allele frequency in population databases, if known; AAC: amino acid change. §Self-identified natural lefthander made to
699 convert to right-handedness.

701 **Table 2.** Gene set enrichment analysis under a recessive mutation model

Group	p-value[1]	term size	query size[2]	overlap size	term ID	term name	intersection	samples	Pval Fisher[3]
Non-PCD SI (n=9)[4]	NS		38						
Non-PCD SI LH (n=5)[4]	NS		26						
Unsolved cases (n=6)[4]	NS		22						
SI with PCD (n=6)	3.85×10^{-5}	17	40	4	GO:0036158	outer dynein arm assembly	DNAH5, CCDC114, LRRC6, DNAAF1	SI06, SI11, SI12, SI13, SI15, SI17	7.39×10^{-13}
SI with PCD (n=6)	1.41×10^{-4}	296	40	8	GO:0007018	microtubule-based movement	DNAH5, CCDC114, DNAH11, WDR60, LRRC6, DNAAF1, FMN2, DNAH12	SI06, SI08, SI11, SI12, SI13, SI15, SI17	8.94×10^{-15}
SI with PCD (n=6)	4.89×10^{-5}	48	40	5	GO:0030286	dynein complex	DNAH5, CCDC114, DNAH11, WDR60, DNAH12	SI08, SI11, SI12, SI13, SI15, SI17	1.59×10^{-13}
SI with PCD (n=6)	0.004	113	40	5	GO:0005930	axoneme	DNAH5, CCDC114, DNAH11, DNAAF1, RP1L1	SI03, SI08, SI11, SI12, SI13, SI15, SI17	1.31×10^{-11}
SI with PCD (n=6)	0.002	427	40	8	GO:0099568	cytoplasmic region	DNAH5, CCDC114, DNAH11, SHROOM2, DNAAF1, FMN2, RP1L1, PCLO	SI03, SI06, SI08, SI11, SI12, SI13, SI15, SI17	1.73×10^{-13}
Unaffected controls (n=15)	NS		34						

702 LH: left-handed.[1] P-values are corrected for multiple testing using the gSCS method. NS: no significant gene sets. [2] The number of mutated genes present in the GO schema.
703 [3] Uncorrected P-values were calculated using Fisher's exact test. [4] Results were the same after excluding subject SI03.
704