1 **Epigenetic signatures associated with imprinted paternally-expressed genes in**

2 **the *Arabidopsis* endosperm**

Jordi Moreno-Romero[1], Gerardo Del Toro-De León, Vikash Kumar Yadav, Juan Santos-

González and Claudia Köhler*

Department of Plant Biology, Uppsala BioCenter, Swedish University of Agricultural

Sciences and Linnean Center for Plant Biology, Uppsala, Sweden

[1]Present address: Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-

UAB-UB, Campus UAB, Bellaterra, Barcelona, Spain.

*Corresponding Author (claudia.kohler@slu.se)

14 **Running title: Epigenetic signatures of imprinted genes**

ABSTRACT

Imprinted genes are epigenetically modified during gametogenesis and maintain the established epigenetic signatures after fertilization, causing parental-specific gene expression. In this study, we show that imprinted paternally-expressed genes (PEGs) in the *Arabidopsis* endosperm are marked by an epigenetic signature of Polycomb-mediated H3K27me3 together with heterochromatic H3K9me2 and CHG methylation, which specifically mark the silenced maternal alleles of PEGs. Based on the presence of this epigenetic signature on maternal alleles we were able to predict novel PEGs and could confirm several of those using published and novel datasets generated in this study. The combined activity of the otherwise largely exclusive H3K27me3 together with the heterochromatic H3K9me3 has also been reported to silence the maternal *Xist* locus in mammalian preimplantation embryos, suggesting an evolutionary conserved employment of both pathways during reproduction.

AUTHOR SUMMARY

Genomic imprinting is an epigenetic phenomenon causing parental alleles to be active

depending on their parent-of-origin. In plants, imprinted genes are mainly confined to

the endosperm, an ephemeral tissue supporting embryo development. Differential

methylation of histone H3 on lysine 27 (H3K27me3) established by the Polycomb

Repressive Complex 2 (PRC2) is a major regulatory mechanism determining activity of

paternally expressed imprinted genes (PEGs) in animals and plants. Here, we show

that the coding region of many PEGs is marked by an epigenetic signature of

H3K27me3, H3K9me2 and CHG methylation and that the combination of these three

modifications correlates with paternally-biased gene expression in the endosperm. The

maternal alleles of PEGs are marked by CHG methylation in the central cell, indicating

that the repressive epigenetic signature of PEGs is established before fertilization. We

use the presence of the three modifications to predict novel PEGs and propose that

genomic imprinting is substantially more common than previously estimated based on

expression data.

INTRODUCTION

Genomic imprinting is an epigenetic phenomenon causing maternal and paternal alleles to be differentially expressed after fertilization. In plants, genomic imprinting is mainly confined to the endosperm, an ephemeral nutritive tissue supporting embryo growth, similar to the placenta in mammals [1]. The endosperm is the product of a double fertilization event, where one of the haploid sperm cells fertilizes the haploid egg cell giving rise to the diploid embryo, while the second sperm cell fertilizes the diploid central cell to give rise to the triploid endosperm [2]. Imprinted genes are epigenetically modified during gamete formation and the established epigenetic asymmetry is maintained after fertilization. Differential DNA methylation is established by the DNA glycosylase DEMETER (DME) that removes methylated cytosine residues in the central cell of the female gametophyte [3]. DME is not active in sperm, leading to differential DNA methylation between male and female genomes in the endosperm. DME acts on small transposable elements (TEs) in the vicinity of genes [4] and its activity has been connected to the expression of maternally-expressed imprinted genes (MEGs). Hypomethylation can furthermore expose binding sites for the Fertilization Independent Seed (FIS)-Polycomb Repressive Complex 2 (PRC2) [5, 6], an evolutionary conserved chromatin modifying complex that applies a trimethylation mark on histone H3 at lysine 27 (H3K27me3) [7]. The Arabidopsis FIS-PRC2 consists of the subunits MEDEA (MEA), FIS2, FERTILIZATION INDEPENDENT ENDOSPERM (FIE), and MULTICOPY SUPPRESSOR OF IRA1 (MSI1) [8] and is specifically active in the central cell of the female gametophyte and in the endosperm [9]. Repression of the maternal alleles of PEGs is mediated by the activity of the FIS-PRC2 [10], consistent with maternal PEG

alleles being marked by H3K27me3 [6]. In this manuscript, we addressed the

mechanism of maternal allele repression in PEGs. We surprisingly found that PEGs are

regulated by two otherwise largely exclusive epigenetic repressive pathways, the FIS2-

PRC2 and the pathway establishing the heterochromatin-localized H3K9me2

modification [11]. We demonstrate that both modifications, H3K27me3 and H3K9me2,

overlap on the maternal alleles of the majority of PEGs. Our data suggest that most

likely FIS-PRC2 acts first and is required to establish H3K9me2. Furthermore, we find

maternal alleles of PEGs to be marked by CHG methylation in the central cell, indicating

that repressive pathways establishing H3K27me3, H3K9me2 and CHG methylation act

in the central cell of the female gametophyte. Finally, we use the presence of the three

modifications to predict novel PEGs and propose that the number of PEGs predicted

based on expression data strongly underestimates the real number of PEGs.

RESULTS

The maternal alleles of PEGs are marked by H3K27me3, H3K9me2, and CHG
methylation

In a previous study, we revealed that H3K27me3 and H3K9me2 overlapped at paternal

heterochromatic regions in the endosperm, suggesting a partial functional redundancy

of both modifications [6]. We now addressed the question whether this redundancy

extends to other regions of the genome. We found that about one-third of genes marked

by H3K27me3 on the maternal alleles were also marked by H3K9me2 (Fig 1A,

hypergeometric test, P=0; S1 Fig, P=0). Genes containing both modifications on their

maternal alleles had significantly higher levels of H3K27me3 compared to those only

marked by H3K27me3 (Fig 1B, S1 Fig). The majority of double-marked genes contained

both modifications specifically on the maternal but not the paternal alleles (Fig 1A, S1

Fig) and increasing levels of H3K27me3 on maternal alleles correlated with increasing

levels of H3K9me2 on the maternal but not the paternal alleles (S2 Fig). We thus

proposed that the presence of both modifications on the maternal alleles correlates with

paternally-biased expression. Consistent with this notion we found that genes previously

identified as PEGs [12] were substantially enriched for both modifications on their

maternal alleles, while MEGs did not show enrichment of both marks on either maternal

or paternal alleles (Fig 1C and 1D, S1 Fig). This trend was independent of the direction

of the cross and similarly observed in Col x L*er* and L*er* x Col crosses (Fig 1 and S1

Fig). The presence of H3K27me3 is generally confined to gene bodies [13], consistent

with the observed enrichment of H3K27me3 in gene bodies of PEG maternal alleles

(Fig 1E and S1 Fig). Interestingly, H3K9me2 was similarly restricted to gene bodies of

PEGs (Fig 1E and S1 Fig), contrasting the exclusion of H3K9me2 from genic regions in

sporophytic tissues [14]. The maternal alleles of PEGs were also strongly enriched for

CHG methylation (Fig 1F, S1 and S2 Figs), consistent with the chromomethylase 3

(CMT3) acting in a positive feedback loop with the histone methyltransferase proteins

KYP/SUVH4, SUVH5, and SUVH6 [11]. Levels of CHH methylation were generally

higher on maternal than on paternal alleles (Fig 1F, S1 Fig), consistent with depletion of

CHH methylation in sperm [4, 15].

Maternal-specific CHG methylation is established in the central cell and depends

on FIS-PRC2

PEGs had increased levels of CHG methylation in the central cell, but low CHG

methylation in sperm and vegetative cells of pollen (Fig 2A), revealing that differences in

CHG methylation are established before fertilization. The FIS-PRC2 had been proposed

to promote non-CG methylation [4], raising the hypothesis that increased CHG

methylation on the maternal alleles of PEGs depends on the activity of FIS-PRC2 in the

central cell. Consistently, we observed strongly decreased CHG methylation on the

maternal alleles of PEGs in seeds lacking maternal FIE activity (Fig 2B), suggesting that

FIS-PRC2 activity is required to recruit CHG methylation. The DNA glycosylase DME is

required for the activation of *MEA* and *FIS2*, both encoding subunits of the FIS-PRC2

[16, 17]. Maternal PEG alleles had reduced levels of CHG methylation in seeds lacking

maternal DME activity (Fig 2B), supporting the notion that FIS-PRC2 activity is required

for CHG methylation establishment on maternal PEG alleles.

Previous work revealed that loss of FIS-PRC2 function caused activation of maternal

PEG alleles [10]. We addressed the question whether loss of CMT3 and the

redundantly acting SUVH4,5,6 [18] similarly affect silencing of maternal PEG alleles.

Based on published expression studies, *CMT3*, *SUVH4*, and *SUVH6* are expressed in

the central cell of the female gametophyte [19]. The maternal alleles of seven tested

PEGs remained silenced in reciprocal crosses of wild type with *cmt3* and *suvh4,5,6*

triple mutants (S3 Fig), indicating that stable silencing of the maternal alleles of PEGs

does not depend on CMT3 and SUVH4,5,6 activity before fertilization. We tested the

requirement of CMT3 and SUVH4,5,6 for PEG regulation after fertilization by monitoring

PEG expression in homozygous mutant *cmt3* and *suvh4,5,6* seeds. None of the seven

tested genes was significantly upregulated in seeds of *cmt3* or *suvh4,5,6* mutants (S3

Fig), indicating that CMT3 and SUVH4,5,6 are not required for repression of maternal

PEG alleles after fertilization. In the *suvh4,5,6* triple mutant, H3K9me2 is genomewide

eliminated in vegetative tissues [20]; however, whether H3K9me2 is similarly eliminated

in the central cell and endosperm remains to be shown. Therefore, deciphering the role

of H3K9me2 in repressing the maternal alleles of PEGs remains to be subject of future

investigations.

Paternally-biased expression coincides with the combination of H3K27me3,

H3K9me2, and CHG methylation

We addressed the question whether the presence of H3K9me2 and CHG methylation is

functionally relevant for maternal allele repression by testing whether the presence of

both modifications either alone or in combination with H3K27me3 correlates with

maternal allele repression. Strikingly, while the presence of either CHG methylation,

H3K9me2, or H3K27me3 on maternal alleles did not shift the allelic balance, the

combination of more than one modification shifted the balance towards preferential

paternal allele expression (Fig 3A, S5 Fig), suggesting that the combined presence of

H3K27me3, H3K9me2, and CHG methylation is a hallmark for maternal allele

repression. Consistently, increased bias towards paternal allele expression correlated

with increased levels of H3K27me3, H3K9me2, and CHG methylation on maternal

alleles (Fig 3B, S5 Fig).

We tested which combination of marks would most reliably allow predicting paternally-

biased genes. We assigned scores for the allele-specific presence of the three

modifications (see S1 Table) and tested whether highly scoring genes were enriched for

previously predicted PEGs [12]. The combination of CHG methylation in the central cell

together with maternal-specific H3K27me3 and H3K9me2 allowed to predict the highest

number of previously described PEGs in Col and L*er* accessions (24 out of 42 (57%),

[12]) in relation to the number of genes in the category with the highest score (Fig 4A,

S1 Table) and was chosen for further analysis. The category with the highest score was

significantly enriched for PEGs (hypergeometric test, P=1.0e-32); while categories with

lower scores contained only few PEGs (Fig 4A). Similarly, out of 64 PEGs that had been

predicted by a recent study re-evaluating previously published imprintome datasets [21],

40 (62%) PEGs were present in the highest score category (Fig 4A). Nearly half (96

genes, 46.4%) of those genes in the highest score category were significantly paternally

biased (Chi-square <0.05, Bonferroni corrected, S2 Table), which was significantly more

than the 8% paternally-biased genes identified among all genes tested (Hypergeometric

test, P= 8.9e-52). We thus conclude that the presence of the three modifications, CHG

in the central cell, H3K27me3 and H3K9me2 on maternal alleles in the endosperm

allows to predict genes with paternally-biased expression. Paternally biased genes were

particularly involved in transcriptional regulation (P=7.54e-5) and chromatin organization

(P=1.57e-3), consistent with previous reports on the functional role of PEGs [12, 22].

Maternal seed coat contamination restricts the identification of PEGs

Published endosperm transcriptome data contain a substantial fraction of transcripts

from the maternal seed coat, which may limit the correct prediction of paternally-biased

genes [21]. We hypothesized that there are several genes that based on their

epigenetic modifications (score 12, Fig 4A and S2 Table) are likely to be PEGs but

based on available expression data are not correctly classified. A prediction of this

hypothesis is that genes in the highest score category that have been classified as

biallelically expressed or maternally-biased are more highly expressed in the seed coat

than those with paternally-biased expression. We tested this hypothesis using available

expression data of seed coat tissue [23] at a similar developmental stage as published

endosperm expression data [12]. Indeed, genes with paternally-biased expression were

significantly less expressed in the seed coat compared to genes that based on available

expression data are either predicted to be biallelically expressed or maternally-biased

(Wilcoxon rank sum test, *P*<0.005; Fig 4B), strongly suggesting that maternal tissue

contamination in available expression data limits the prediction of PEGs. To further test

this hypothesis, we generated transgenic reporter lines containing the promoter ($\simeq$2kb)

and downstream genic regions fused to the green fluorescent protein (GFP) of seven

genes belonging to the highest score category but predicted to be maternally

(*AT2G33620*, *AT1G43580*, *AT1G47530*, *AT1G64660*, *AT2G30590*, *AT4G15390*) or

biallelically expressed (*AT5G53160*). We detected a GFP signal in the endosperm only

for construct *AT1G64660* (Fig 5, S3Table). For construct *AT1G47530* a signal was

detected in the seed coat, while for the other constructs no GFP signal was detected in

seeds, indicating that the regulatory elements required for the expression of those

genes are located outside the promoter and genic regions used to generate the reporter

lines. Reciprocal crosses using the *AT1G64660* reporter lines revealed that this gene is

indeed a PEG and strongly expressed in the endosperm when paternally, but not when

maternally inherited (Fig 5). This data support the hypothesis that seed coat

contamination limits the transcriptome-based identification of PEGs.

To test whether we could confirm additional PEGs that have been predicted based on

their epigenetic signatures, we applied INTACT (isolation of nuclei tagged in specific cell

types) to purify endosperm nuclei at 4 days after pollination (DAP) from Col × L*er* and

L*er* × Col reciprocal crosses. Isolated RNA was sequenced and profiled for allele-

specific gene expression. By analyzing the maternal to total reads ratio in each

epigenetic category, we confirmed that the genes in the highest score category (group

with score 12) indeed showed a clear trend towards paternally-biased expression (Fig

6A, S4 Table). Following previously established criteria [12], we predicted 107 PEGs

that were reciprocally imprinted in both directions of the crosses. There was a

significantly higher number of PEGs present in the highest score category compared to

a representative random sample of genes with informative reads (Fig 6B). Furthermore,

the highest score category had the highest frequency of PEGs, with other categories

having significantly fewer PEGs (Fig 6C). Of the 107 genes that we predicted as PEGs

based on our RNA sequencing data, 38 were present in the highest score category,

which is significantly more than expected by chance (p=3.6 e-59, Fig 6D). Out of those,

20 were previously predicted based on published data [10, 12, 24, 25], while 18 genes

are likely new high-confidence PEGs, revealing that PEGs are more common than

previously estimated.

DISCUSSION

In this study, we identified the concomitant presence of maternal-specific CHG

methylation, H3K27me3, and H3K9me2 as an epigenetic signature for paternally-biased

expression in the endosperm. We furthermore predict that there are substantially more

PEGs than previously reported in Arabidopsis [10, 12, 24], suggesting that in

Arabidopsis the number of PEGs exceeds the number of MEGs. Recent re-evaluation of

published imprintome data of Arabidopsis revealed that a large number of previously

predicted MEGs were seed coat expressed genes, while the number of PEGs was

underestimated [21]. Our study supports and extends this notion by showing that there

is a large number of paternally-biased genes that likely failed to be identified in previous

studies because of maternal seed coat contamination or early stage-specific

expression.

A previous study reported that the maternal alleles of PEGs in *A. lyrata* are marked by

CHG methylation and implicated that closely related species use different mechanisms

to regulate imprinted gene expression [26]. Our study reveals that similar to *A. lyrata* the

maternal alleles of many PEGs are also marked by CHG methylation in *A. thaliana*,

highlighting that epigenetic mechanisms employed to maintain monoallelic expression

are rather conserved between related species. Interestingly, while the maternal alleles

of PEGs in maize are marked by H3K27me3, they are not marked by CHG methylation

[27], indicating that diverged species may use different mechanism in maintaining

maternal allele repression.

How H3K9me2 and CHG methylation are established at PRC2 target genes remains to

be studied; however, the strong activation of maternal PEG alleles upon loss of FIS-

PRC2 function [10] suggests that H3K9me2 and CHG methylation require FIS-PRC2

function. The PRC2 is generally targeting genes with specific roles during development

[28], while complexes establishing H3K9me2 are mainly targeting TEs localized in

heterochromatic regions of the genome [29]. This functional division of PRC2 and

machineries establishing heterochromatic marks is conserved in plants as well as in

mammals [30]; however, there are notable exceptions to this rule in both groups of

organisms. In rice seedlings, about one-third of H3K27me3 marked genes are also

marked by CHG and CHH methylation [31]. Similar to the findings reported in our study,

higher levels of H3K27me3 correlate with higher CHG and CHH methylation in gene

bodies of rice [31]. The rice H3K27me3 methyltransferase SDG711 physically interacts

with the CHH methyltransferase OsDMR2 and the SRA-domain containing SUVH

protein SDG703, uncovering a mechanistic connection between PRC2 and non-CG

methylation. In mammalian cells, H3K9 methyltransferases colocalize with PRC2 [32-

34], revealing crosstalk between these two major epigenetic silencing pathways that

likely is required for stable gene silencing. The imprinted maternal *Xist* locus encoding

an X-linked long-noncoding RNA is covered by H3K27me3 and H3K9me3 in

preimplantation embryos [35, 36]. Importantly, loss of maternal H3K27me3 induces *Xist*

activation, indicating that maternal H3K27me3 is the major imprinting mark of *Xist* [36].

This is strikingly similar to findings made in this study revealing H3K27me3 as the major

repressive mark for PEGs. Recent work revealed that maternal H3K27me3 controls

DNA methylation-independent imprinting in mammalian preimplantation embryos [37].

While imprinted expression of most genes is lost in the embryonic cell lineage, few

genes maintain their imprinted expression in the extra-embryonic cell lineage [37]. The

Xist locus that is marked by H3K27me3 and H3K9me3 is among those loci that remain

imprinted in extra-embryonic tissues [36]. Whether the presence of both marks

distinguishes those genes that maintain their imprinted expression from those that

become biallelically expressed remains to be tested, but we consider this a very

attractive hypothesis. We speculate that the presence of both marks in certain tissue

types of mammals and flowering plants is a conserved epigenetic signature marking

stably repressed genes.

MATERIAL AND METHODS

Plant Material and Growth Conditions

All seeds were surface sterilized (5% sodium hypochlorite and 0.01% Triton X-100),

stratified for 2 days at 4°C and germinated on half-strength Murashige and Skoog

medium containing 1% sucrose under long-day conditions (16 h light/8 h darkness;

21°C). Plants were transferred to soil after 10 to 12 days and grown under long day

conditions. The *cmt3-11* (SALK_148381; [38]) and *suvh456* mutants (kindly provided by

Judith Bender) used in this study are in the Col-0 background.

Imprinting Assays and Expression Analysis

To generate siliques of indicated crosses, three to five flowers were emasculated, hand-

pollinated, and harvested at 4 DAP. RNA extraction was performed using the MagJET

Plant RNA Purification Kit (Thermo Scientific) following the manufacturer´s instructions.

Residual DNA was removed using Invitrogen DNase I (Amplification Grade), and cDNA

was synthesized using the Fermentas first strand cDNA synthesis kit according to the

manufacturer's instructions. Quantitative PCR was performed using a MyiQ5 real-time

PCR detection system (Bio-Rad) and Solis BioDyne-5x Hot FIREPol EvaGreen qPCR

Mix Plus (ROX, Solis BioDyne). For the imprinting-by-sequencing assay, the PCR

products were purified and analyzed by Sanger sequencing. For the imprinting-by-

restriction enzyme digestion assay, the PCR products were purified and digested.

Restriction enzymes and primers used are listed in the S5 Table.

Data analysis

We made use of endosperm-specific ChIP-seq data that have been previously

generated in our group [6]. Data correspond to pooled biological triplicates with ChIP

signals being normalized with H3 ChIP data by calculating the log2 ratio in 150-bp bins

across the genome. Data were standardized and normalized with a z-score

transformation [39]. Metagene plots over genes were constructed between -2 kb and

+2 kb by calculating mean levels of methylation signals in 100-bp bins in the flanks of

the genes and in 40 equally long bins between the transcriptional start and stop. Gene

z-scores were calculated as an average of z-scores over the gene body. DNA

methylation data of *fie* and *dme* and their corresponding wild types are from [4]. DNA

methylation data of the central cell, sperm, and vegetative cells are from [3] and [15].

Endosperm expression data are from [12] and seed coat expression data from [23].

Endosperm nuclei isolation and RNA sequencing

We performed Col × L*er* reciprocal crosses using *Arabidopsis* lines expressing

PHE1::NTF and *PHE1::BirA* (lines referred as INT hereafter) [40]. To facilitate the

crosses, we used the male sterile mutants *pistillata* (*pi-1*, in L*er* accession) and *dde2* (in

Col accession containing INT) as female parents and pollinated them with the INT line

(Col accession) and L*er* wild type, respectively. A total of 250mg of siliques per sample

was collected at 4DAP. Tissue homogenization and nuclei purification were performed

from three biological replicates as previously described [40]. Total RNA was extracted

from purified nuclei using the mirVana Isolation Kit Protocol (Ambion). mRNA extraction

was performed using NEBNext Poly(A) mRNA Magnetic Isolation and the Libraries were

prepared with the NEBNext Ultra II RNA Library Prep Kit from Illumina. Samples were

sequenced at the National Genomic Infrastructure (NGI) from SciLife Laboratory

(Uppsala, Sweden) on an Illumina HiSeq2500 in paired-end 125bp read length. Reads

were trimmed and mapped in single-end mode to the Arabidopsis (TAIR10) genome

previously masked for *rRNA* genes and for SNP positions between Col and

L*er*accessions, using TopHat v2.1. [41] (parameters adjusted as -g 1 -a 10 -i 40 -I 5000

-F 0 -r 130). Gene expression was normalized to reads per kilobase per million mapped

reads (RPKM) using GFOLD40 [42]. For discrimination of maternal and paternal

transcripts, reads were assigned to the Col or L*er* genomes using single nucleotide

polymorphisms (SNPs) between the accessions using SNPsplit [43]. We calculated

contamination levels based on the deviation of the read counts from the expected 2:1

maternal/paternal genome ratio in the endosperm [40] (S6 Table). Based on this

analysis, one of the triplicates of the cross direction Col × L*er* was not included in the

downstream analyses. To increase the statistical power to detect parentally-biased

genes, we merged libraries from two replicates of Col × L*er* and three replicates of L*er* ×

Col for downstream analysis.

Allele-specific expression analysis

We defined a minimum threshold of 20 and 30 informative reads for Col × L*er* (2

replicates) and L*er* × Col (3 replicates) crosses, respectively. Statistical differences

between maternal and paternal read counts for each gene were calculated using a Chi-

square test, considering genes with a false discovery rate adjusted *P*-value of less than

0.01. Additionally, MEGs required to have at least 85% maternal informative reads in

both directions of the reciprocal cross and PEGs to have at least 50% paternal

informative reads in both directions of the reciprocal cross, following previously defined

conditions [12].

Generation of reporter constructs and transgenic lines

For the generation of reporter constructs, we used the ClonExpress® MultiS One Step

Cloning kit. Promoters ($\simeq$2kb) and genic sequences of *AT2G33620*, *AT1G43580*,

AT1G47530, *AT1G64660*, *AT2G30590*, *AT4G15390*, and *AT5G53160* were amplified

from WT Col-0 genomic DNA using primers specified in S5 Table and cloned into vector

pB7FWG.0. Constructs were transformed into *Agrobacterium tumefaciens* strain

GV3101 and *Arabidopsis* plants were transformed using the floral dip method [44]. Ten

transgenic lines per construct were generated and analyzed.

Imprinting analyses

For reciprocal crosses, designated female partners were emasculated at 1–2 days prior

to anthesis. Two days after emasculation pistils were hand-pollinated with respective

pollen donors. Seeds were dissected from the siliques and mounted on a microscope

slide for imaging and counting at 2 and 4 DAP. For fluorescence analyses, seeds were

stained with 0.1 mg/mL propidium iodide (PI) solution in 7% glucose. Seeds of

reciprocal crosses of reporter lines were analyzed under confocal microscopy on a

Zeiss 800 Inverted Axio Observer with a supersensitive GaASp detector. Images were

acquired, analyzed and exported using Zeiss ZEN software. For each reporter construct

50-60 seeds per line were analyzed.

Acknowledgements

This research was supported by a European Research Council Starting Independent

Researcher grant (to C.K.), a grant from the Swedish Science Foundation (to C.K.), a

grant from the Knut and Alice Wallenberg Foundation (to C.K.), and an EMBO

fellowship (to G.D.T.D.L). Sequencing was performed by the SNP&SEQ Technology

Platform, Science for Life Laboratory at Uppsala University, a national infrastructure

supported by the Swedish Research Council (VRRFI) and the Knut and Alice

Wallenberg Foundation.

References:

1. Rodrigues JA, Zilberman D: Evolution and function of genomic imprinting in
plants. Genes Dev 2015;29: 2517-2531.

2. Li J, Berger F: Endosperm: food for humankind and fodder for scientific
discoveries. New Phytol 2012;195: 290-305.

3. Park K, Kim MY, Vickers M, Park JS, Hyun Y, Okamoto T, Zilberman D, Fischer
RL, Feng X, Choi Y et al: DNA demethylation is initiated in the central cells of
Arabidopsis and rice. Proc Natl Acad Sci USA 2016;113: 15138-15143.

4. Ibarra CA, Feng X, Schoft VK, Hsieh TF, Uzawa R, Rodrigues JA, Zemach A,
Chumak N, Machlicova A, Nishimura T et al: Active DNA demethylation in plant
companion cells reinforces transposon methylation in gametes. Science
2012;337: 1360-1364.

5. Weinhofer I, Hehenberger E, Roszak P, Hennig L, Köhler C: H3K27me3 profiling
of the endosperm implies exclusion of Polycomb group protein targeting by DNA
methylation. PLoS Genet 2010;6: e1001152.

6.   Moreno-Romero J, Jiang H, Santos-Gonzalez J, Kohler C: Parental epigenetic
asymmetry of PRC2-mediated histone modifications in the Arabidopsis
endosperm. EMBO J 2016;35: 1298-1311.

7.   Simon JA, Kingston RE: Occupying chromatin: Polycomb mechanisms for getting
to genomic targets, stopping transcriptional traffic, and staying put. Mol Cell
2013;49: 808-824.

8.   Mozgova I, Hennig L: The Polycomb Group protein regulatory network. Annu Rev
Plant Biol 2015;66: 269-296.

9.   Luo M, Bilodeau P, Dennis ES, Peacock WJ, Chaudhury A: Expression and
parent-of-origin effects for FIS2, MEA, and FIE in the endosperm and embryo of
developing Arabidopsis seeds. Proc Natl Acad Sci USA 2000;97: 10637-10642.

10.   Hsieh TF, Shin J, Uzawa R, Silva P, Cohen S, Bauer MJ, Hashimoto M, Kirkbride
RC, Harada JJ, Zilberman D et al: Regulation of imprinted gene expression in
Arabidopsis endosperm. Proc Natl Acad Sci USA 2011;108: 1755-1762.

11.   Law JA, Jacobsen SE: Establishing, maintaining and modifying DNA methylation
patterns in plants and animals. Nat Rev Genet 2010;11: 204-220.

12.   Pignatta D, Erdmann RM, Scheer E, Picard CL, Bell GW, Gehring M: Natural
epigenetic polymorphisms lead to intraspecific variation in Arabidopsis gene
imprinting. Elife 2014;3: e03198.

13.   Shu H, Nakamura M, Siretskiy A, Borghi L, Moraes I, Wildhaber T, Gruissem W,
Hennig L: Arabidopsis replacement histone variant H3.3 occupies promoters of
regulated genes. Genome Biol 2014;15: R62.

14.   Roudier F, Ahmed I, Berard C, Sarazin A, Mary-Huard T, Cortijo S, Bouyer D,
Caillieux E, Duvernois-Berthet E, Al-Shikhley L et al: Integrative epigenomic
mapping defines four main chromatin states in Arabidopsis. EMBO J 2011;30:
1928-1938.

15.   Calarco JP, Borges F, Donoghue MT, Van Ex F, Jullien PE, Lopes T, Gardner R,
Berger F, Feijo JA, Becker JD et al: Reprogramming of DNA methylation in
pollen guides epigenetic inheritance via small RNA. Cell 2012;151: 194-205.

16.   Choi Y, Gehring M, Johnson L, Hannon M, Harada JJ, Goldberg RB, Jacobsen
SE, Fischer RL: DEMETER, a DNA glycosylase domain protein, is required for
endosperm gene imprinting and seed viability in Arabidopsis. Cell 2002;110: 33-
42.

17.   Jullien PE, Kinoshita T, Ohad N, Berger F: Maintenance of DNA methylation
during the Arabidopsis life cycle is essential for parental imprinting. Plant Cell
2006;18: 1360-1372.

18. Ebbs ML, Bender J: Locus-specific control of DNA methylation by the
Arabidopsis SUVH5 histone methyltransferase. Plant Cell 2006;18 1166-1176.

19. Wuest SE, Vijverberg K, Schmidt A, Weiss M, Gheyselinck J, Lohr M, Wellmer F,
Rahnenfuhrer J, von Mering C, Grossniklaus U et al: Arabidopsis female
gametophyte gene expression map reveals similarities between plant and animal
gametes. Curr Biol 2010;20: 506-512.

20. Stroud H, Greenberg MV, Feng S, Bernatavichute YV, Jacobsen SE:
Comprehensive analysis of silencing mutants reveals complex regulation of the
Arabidopsis methylome. Cell 2013;152: 352-64.

21. Schon MA, Nodine MD: Widespread Contamination of Arabidopsis Embryo and
Endosperm Transcriptome Data Sets. Plant Cell 2017;29: 608-617.

22. Waters AJ, Bilinski P, Eichten SR, Vaughn MW, Ross-Ibarra J, Gehring M,
Springer NM: Comprehensive analysis of imprinted genes in maize reveals allelic
variation for imprinting and limited conservation with other species. Proc Natl
Acad Sci USA 2013;110: 19639-19644.

23. Belmonte MF, Kirkbride RC, Stone SL, Pelletier JM, Bui AQ, Yeung EC,
Hashimoto M, Fei J, Harada CM, Munoz MD et al: Comprehensive
developmental profiles of gene activity in regions and subregions of the
Arabidopsis seed. Proc Natl Acad Sci USA 2013;110: 435-444.

24. Wolff P, Weinhofer I, Seguin J, Roszak P, Beisel C, Donoghue MT, Spillane C,
Nordborg M, Rehmsmeier M, Köhler C et al: High-Resolution Analysis of Parent-
of-Origin Allelic Expression in the Arabidopsis Endosperm. PLoS Genet 2011;7:
e1002126.

25. Gehring M, Missirian V, Henikoff S: Genomic Analysis of Parent-of-Origin Allelic
Expression in Arabidopsis thaliana Seeds. PLoS One 2011;6: e23687.

26. Klosinska M, Picard CL, Gehring M: Conserved imprinting associated with unique
epigenetic signatures in the Arabidopsis genus. Nat Plants 2016;2: 16145.

27. Zhang M, Xie S, Dong X, Zhao X, Zeng B, Chen J, Li H, Yang W, Zhao H, Wang
G et al: Genome-wide high resolution parental-specific DNA and histone
methylation maps uncover patterns of imprinting regulation in maize. Genome
Res 2014;24: 167-176.

28. Mozgova I, Kohler C, Hennig L: Keeping the gate closed: functions of the
polycomb repressive complex PRC2 in development. Plant J 2015;83: 121-132.

29. Du J, Johnson LM, Jacobsen SE, Patel DJ: DNA methylation pathways and their
crosstalk with histone methylation. Nat Rev Mol Cell Biol 2015;16: 519-532.

30. Alabert C, Groth A: Chromatin replication and epigenome maintenance. Nat Rev Mol Cell Biol 2012;13: 153-167.

31. Zhou S, AUID- ORCID: ht, Liu X, Zhou C, Zhou Q, Zhao Y, Li G, Zhou DX: Cooperation between the H3K27me3 Chromatin Mark and Non-CG Methylation in Epigenetic Regulation. Plant Physiol 2016;172: 1131-1141.

32. Bilodeau S, Kagey MH, Frampton GM, Rahl PB, Young RA: SetDB1 contributes to repression of genes encoding developmental regulators and maintenance of ES cell state. Genes Dev 2009;23: 2484-2489.

33. Yuan P, Han J, Guo G, Orlov YL, Huss M, Loh YH, Yaw LP, Robson P, Lim B, Ng HH et al: Eset partners with Oct4 to restrict extraembryonic trophoblast lineage potential in embryonic stem cells. Genes Dev 2009;23: 2507-2520.

34. Mozzetta C, Pontis J, Fritsch L, Robin P, Portoso M, Proux C, Margueron R, Ait-Si-Ali S: The histone H3 lysine 9 methyltransferases G9a and GLP regulate polycomb repressive complex 2-mediated gene silencing. Mol Cell 2014;53: 277-289.

35. Fukuda A, Tomikawa J, Miura T, Hata K, Nakabayashi K, Eggan K, Akutsu H, Umezawa A: The role of maternal-specific H3K9me3 modification in establishing imprinted X-chromosome inactivation and embryogenesis in mice. Nat Commun 2014;5: 5464.

36. Inoue A, Jiang L, Lu F, Zhang Y: Genomic imprinting of Xist by maternal H3K27me3. Genes Dev 2017;31: 1927-1932.

37. Inoue A, Jiang L, Lu F, Suzuki T, Zhang Y: Maternal H3K27me3 controls DNA methylation-independent imprinting. Nature 2017;547: 419-424.

38. Zhang X, Yazaki J, Sundaresan A, Cokus S, Chan SW, Chen H, Henderson IR, Shinn P, Pellegrini M, Jacobsen SE et al: Genome-wide high-resolution mapping and functional analysis of DNA methylation in arabidopsis. Cell 2006;126: 1189-1201.

39. Cheadle C, Vawter MP, Freed WJ, Becker KG: Analysis of microarray data using Z score transformation. J Mol Diagn 2003;5: 73-81.

40. Moreno-Romero J, Santos-Gonzalez J, Hennig L, Köhler C: Applying the INTACT method to purify endosperm nuclei and to generate parental-specific epigenome profiles. Nat Protoc 2017;12: 238-254.

41. Trapnell C, Pachter L, Salzberg SL: TopHat: discovering splice junctions with RNA-Seq. Bioinformatics 2009;25: 1105-1111.

42. Feng J, Meyer CA, Wang Q, Liu JS, Shirley Liu X, Zhang Y: GFOLD: a
generalized fold change for ranking differentially expressed genes from RNA-seq
data. Bioinformatics 2012;28: 2782-2788.

43. Krueger F, Andrews SR: SNPsplit: Allele-specific splitting of alignments between
genomes with known SNP genotypes. F1000Res 2016;5: 1479.

44. Clough SJ, Bent AF: Floral dip: a simplified method for agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J 1998;16: 735-743.

Figure Legends

Fig 1. H3K27me3, H3K9me2, and CHG methylation overlap on maternal alleles of

PEGs.

(A) Overlap of genes marked by H3K27me3 and H3K9me2 on the maternal alleles.

Dark red inset marks genes that have H3K27me3 and H3K9me2 exclusively on their

maternal alleles (excluding genes also marked on the paternal alleles).

(B) Genes marked by H3K27me3 and H3K9me2 have higher levels of H3K27me3. Box

plots show mean values of z-scored H3K27me3 and H3K9me2 of genes where the

maternal alleles are either marked by H3K27me3 alone or by a combination of

H3K27me3 and H3K9me2. Genes containing maternal-specific H3K27me3 and

H3K9me2 are particularly enriched for H3K27me3. Boxes show medians and the

interquartile range, error bars show the full range excluding outliers. Notches show the

95% confidence interval for the median. Asterisks mark significance (Wilcoxon rank sum

test, P <1.0E-3). Presence of the mark corresponds to the mark highest score defined in

S1 Table.

(C) The majority of PEGs are marked by H3K27me3 and H3K9me2 on their maternal

alleles. Overlap of PEGs [12] with maternal and paternal H3K27me3 and H3K9me2.

(D) Levels of H3K27me3 and H3K9me2 on PEGs, MEGs, and all genes. Box plots (as

specified in panel B) show mean values of z-scored H3K27me3 and H3K9me2.

Asterisks mark significance (Wilcoxon rank sum test, P <1.0E-3).

(E) Metagene plots showing average distribution of z-score normalized H3K27me3 and

H3K9me2 levels on maternal and paternal alleles of PEGs [12].

(F) Levels of DNA methylation in each sequence context (mCG, mCHG and mCHH) on

PEGs, MEGs, and all genes. Box plots (as specified in panel B) show mean values of

relative DNA methylation. Asterisks mark significance (Wilcoxon rank sum test, P<5.0E-

3). Data shown correspond to cross Col × L*er* [6], for reciprocal cross direction see S1

Fig.

Fig 2. Maternal CHG methylation is established in the central cell and depends on

FIS-PRC2.

(A) Box plots showing mean values of relative DNA methylation in the central cell,

sperm cell, and vegetative cell for PEGs, MEGs, and all genes. Boxes show medians

and the interquartile range, error bars show the full range excluding outliers. Notches

show the 95% confidence interval for the median. Asterisk marks significance (Wilcoxon

rank sum test, P <5.0E-3).

(B) Box plots showing mean values of CHG methylation for PEGs and MEGs on

maternal and paternal alleles in wild-type, *fie* (left panels) and *dme* (right panels)

endosperm. Asterisk marks significance (Wilcoxon rank sum test, *P* <5.0E-2).

Fig 3. Paternally-biased expression coincides with the combination of H3K27me3,

H3K9me2, and CHG methylation.

(A) Box plots show mean values of maternal to total reads for genes marked by one or

several of the following modifications: CHG methylation in the central cell, difference of

CHG methylation, H3K27me3, or H3K9me2 levels on maternal and paternal alleles in

the endosperm. For each modification only genes present in the highest scoring

category were included in the analysis (see S1 Table). Boxes show medians and the

interquartile range, error bars show the full range excluding outliers. Widths of boxes

are proportional to square-root of the number of genes in each category. Blue line

indicates the expected ratio for biallelically expressed genes.

(B) Paternal expression bias correlates with high levels of CHG methylation in the

central cell and H3K27me3 and H3K9me2 on maternal alleles in the endosperm.

Plotted are mean values of CHG methylation, H3K27me3, and H3K9me2 on maternal

and paternal alleles for genes with deviating parental expression levels from the

expected two maternal to one paternal ratio.

Data shown correspond to cross Col × L*er*, for reciprocal cross direction see S4 Fig.

Fig 4. Epigenetic signatures allow predicting paternally-expressed imprinted

genes (PEGs).

(A) The combination of CHG methylation in the central cell together with maternal-specific H3K27me3 and H3K9me2 allows predicting previously described PEGs [12, 21]. Scores are calculated as described in the S1 Table. High and low scores correspond to high or low levels of CHG methylation, H3K27me3, and H3K9me2, respectively. Number of PEGs corresponds to PEGs previously predicted [12] (i) and [21] (ii) in Col and L*er* accessions. Total genes correspond to all genes present in the corresponding categories. Box plots show mean values of maternal to total reads in the indicated categories. Boxes show medians and the interquartile range, error bars show the full range excluding outliers. Blue line indicates the expected ratio for biallelically expressed genes.

(B) Box plots show relative expression level of genes in the seed coat of torpedo stage embryos [23]. Genes present in the highest score category (S2 Table) were grouped in paternally biased genes (Chi-square test, *P*<0.05 Bonferroni corrected) and not-paternally biased genes. Asterisks mark significance (Wilcoxon rank sum test, *P* <5.0E-3).

Fig 5. Reporter lines for *AT1G64660* confirm imprinted expression.

Representative images of seeds derived after reciprocal crosses of the *AT1G64660* reporter line (fusion with green fluorescent protein (GFP)) with wild-type (WT) plants. GFP fluorescence was detected in the seed coat when the *AT1G64660* reporter was maternally inherited, but endosperm-specific expression was only detected when the *AT1G64660* reporter was paternally inherited. Seeds at 2 DAP were used for imaging. Red stain is propidium iodide. Scale bars correspond to 50 μm.

Fig 6. Confirmation of predicted PEGs in RNA samples of INTACT-purified endosperm nuclei.

(A) Distribution of maternal to total read values for genes in each epigenetic category. The distribution for the total population of genes with informative reads (U) is included. Genes with the highest levels of CHG methylation, H3K27me3, and H3K9me2 show a clear trend towards paternally-biased expression (around 0.5). Values correspond to combined values of Col × L*er* and L*er* × Col crosses. Boxes show medians and the interquartile range, lines show the full range of outliers. Blue line indicates the expected ratio for biallelically expressed genes (0.66).

(B) Parent-of-origin distribution of the group of genes with score 12, compared with a random group of the same size. Significantly more PEGs are detected in the group of genes with score 12, compared with the random sample (Wilcoxon test, *P*-values correspond to the comparison with the total population of PEGs in the whole RNAseq data). The scatter plots show maternal to total read values for Col × L*er* and L*er* × Col reciprocal crosses. Colors indicate genes with significant parent-of-origin biased expressed: MEGs, 85% or more of maternal reads in each reciprocal cross; PEGs, 50% or more of paternal reads in each reciprocal cross. Maternally and paternally biased genes show significantly biased expression ($P <= 0.01$) in both reciprocal crosses.

(C) Percentage of PEGs, MEGs and biallelically expressed genes in each epigenetic category. Color code indicates the percentage of genes within each category. Categories with genes having lower levels of CHG methylation, H3K27me3, and H3K9me2 marks have significantly fewer PEGs than the highest scoring category (Chi-

square test, $P<1e-8$ in categories 1-10; $P=0.001$ in category 11). Only PEGs, MEGs and Biallelic categories were considered.

(D) Overlap of PEGs that were detected in our INTACT-based RNAseq samples (blue) with genes predicted as PEGs based on their epigenetic score 12 (green) and previously reported PEGs (yellow) by [10, 12, 24, 25]. New predicted and confirmed PEGs in bold. Statistical significance was calculated using a hypergeometric test.

Supporting information

S1 Table. Scores assigned to genes based on the presence of CHG methylation in the central cell as well as H3K9me2 and H3K27me3 on maternal alleles in the endosperm.

S2 Table. Seed coat expression of genes with maximum score (12) based on S1Table.

S3 Table: Fluorescence analysis of 4DAP seeds derived after indicated reciprocal crosses.

S4 Table. Parent-of-origin RNAseq dataset of 4DAP INTACT-purified endosperm of Col × L*er* reciprocal crosses.

S5 Table. List of primers used in this study.

S6 Table. Quality of sequencing samples.

S1 Fig. H3K27me3, H3K9me2, and CHG methylation overlap on maternal alleles of PEGs.

S2 Fig. Total and parental-specific H3K27me3, H3K9me2, and CHG methylation levels in the endosperm of previously described PEGs [12].

S3 Fig. Maternal alleles remain silenced in *suvh4,5,6* and *cmt3* mutants.

640　**S4 Fig.** Increasing levels of H3K27me3 on maternal alleles correlates with increasing

641　levels of H3K9me2.

642　**S5 Fig.** Paternally-biased expression coincides with the combination of H3K27me3,

643　H3K9me2, and CHG methylation.

A

score	PEGs (i)	PEGs (ii)	PEGs (shared)	Total genes
12	24	40	17	207
11	3	3	2	80
10	3	3	2	114
9	1	0	0	108
8	0	1	0	103
7	1	0	0	206
6	1	3	1	405
5	0	4	0	787
4	2	1	0	4516
3	2	4	2	2490
2	1	1	1	351
1	1	1	1	972
0	3	2	2	1995

Maternal/Total Ratio

1   0.8   0.66   0.5   0

♀   ♂

B

Relative expression in seed coat

Pat.-biased   Not Pat.-biased

*

AT1G64660-GFP × WT   WT × *AT1G64660 -GFP*